A natural single nucleotide mutation in the small regulatory RNA ArcZ of

***Dickeya solani* switches off the antimicrobial activities against yeast and**

bacteria.

Running title: ArcZ role in *Dickeya solani* anti-microbial activities

Authors: Brual T[1$], Effantin G[1$], Baltenneck J[1], Attaiech L[2], Grosbois C[1], Royer M[4], Cigna J[5],

Faure D[6], Hugouvieux-Cotte-Pattat N[1] & Gueguen E[1]*.

Affiliation:

[1] Univ Lyon, Université Claude Bernard Lyon1, CNRS, INSA Lyon, UMR5240 MAP, F-

69622, LYON, France

[2] CIRI, Centre International de Recherche en Infectiologie, Team "Horizontal gene transfer in

bacterial pathogens" (Horigene), Inserm, U1111, Université Claude Bernard Lyon 1, CNRS,

UMR5308, Ecole Normale Supérieure de Lyon, Univ Lyon, 69100, Villeurbanne, France.

[4] CIRAD, UMR PHIM, Montpellier, F-34398, France

[5] French Federation of Seed Potato Growers (FN3PT-inov3PT), 43-45 rue de Naples, 75008

Paris, France

[6] Paris-Saclay University, CNRS, CEA, Institute for Integrative Biology of the Cell (I2BC), Gif-

sur-Yvette, France

[$] These authors contributed equally to this work.

*corresponding author: erwan.gueguen@univ-lyon1.fr

Orcid I.D.

G.E.   0000-0001-8784-1491

H-C-P.N   0000-0002-4322-160X

F. D.   0000-0002-5379-8867

R. M.   0000-0003-0421-982X

A. L.   0000-0002-6188-1839

E.G.   0000-0002-7296-0831

B.J.   0000-0001-8456-4126

C. J.   0000-0002-1219-7096

B.T   0000-0001-5179-8882

ABSTRACT

The necrotrophic plant pathogenic bacterium *Dickeya solani* emerged in the potato

agrosystem in Europe. All isolated strains of *D. solani* contain several large polyketide

synthase/non-ribosomal peptide synthetase gene clusters. Analogy with genes described in

other bacteria suggests that the clusters *ooc* and *zms* are involved in the production of

secondary metabolites of the oocydin and zeamine families, respectively. A third cluster that

we named *ssm* for solani secondary metabolite had an unknown function. In this study, we

constructed mutants impaired in each of the three secondary metabolite clusters *ssm*, *ooc*,

and *zms* to compare first the phenotype of the *D. solani* wild-type strain D s0432-1 with its

associated mutants. We demonstrated the antimicrobial functions of these three PKS/NRPS

clusters against bacteria, yeasts or fungi. The secondary metabolite cluster *ssm*, conserved

in several other *Dickeya* species, produces a secondary metabolite inhibiting yeasts.

Phenotyping and comparative genomics of different *D. solani* wild-type isolates revealed that

the small regulatory RNA ArcZ plays a major role in the control of the clusters *ssm* and *zms*.

A single-point mutation, conserved in some *Dickeya* wild-type strains, including the type strain

IPO 2222, impairs the ArcZ function by affecting its processing into an active form.

197 words

AUTHOR SUMMARY

The development of new antibacterial molecules is critical in tackling the emergence of new

pathogens or bacterial strains resistant to already available antibiotics. Bacterial

phytopathogens can potentially synthesize novel compounds capable of targeting a specific

type of microorganism. *Dickeya solani* is the only of the twelve described *Dickeya* species that

has the three secondary metabolic pathways *ssm*, *ooc* and *zms*. An investigation of the

functions of these three clusters allowed us to identify the anti-yeast activity of the *ssm* cluster,

a potential new molecule of clinical importance. By comparing the antimicrobial activity of

several *Dickeya solani* strains, we identified the small RNA regulator ArcZ as a critical

regulator in the activation of the *ssm* and *zms* clusters. Our study showed that single-

nucleotide polymorphisms of sRNA encoding genes can have huge impacts on bacterial

phenotypes. It is thus critical to pay attention to the allele diversity of sRNA genes.

148 words

INTRODUCTION

Bacterial phytopathogens of the genus *Dickeya* and *Pectobacterium* are pectinolytic necrotrophic bacteria with a broad host plant spectrum [1,2]. These members of the family *Pectobacteriaceae* [3] cause substantial agricultural losses worldwide by affecting many vegetables, ornamentals and crops, of which the potato is the most important economically. These bacteria are able to invade and degrade the plant tissues through the coordinated expression of genes encoding virulence factors, with a major role of pectate lyases that dissociate the plant cell wall constituents [4].

The *Dickeya* genus was established in 2005 [5], resulting from the reclassification of *Pectobacterium chrysanthemi* (formerly *Erwinia chrysanthemi*). To date, twelve species of *Dickeya* have been described, *D. aquatica*, *D. chrysanthemi*, *D. dadantii, D. dianthicola* , *D. fangzhongdai*, *D. lacustris*, *D. oryzae*, *D. parazeae, D. poaceiphila*, *D. solani*, *D. undicola*, and *D. zeae* [5–14].

The species *D. solani* was officially established in 2014 [9] but *D. solani* isolates have attracted attention since its emergence on the potato agrosystem in Europe in the early 2000s. It causes symptoms in both subtropical and temperate climates. Many scientific efforts have been made to provide information on this phytopathogen, resulting in 76 *D. solani* genomes available in May 2021 [15]. Comparative genomics was performed to identify the genetic basis for the different levels of virulence between *D. solani* strains [15–20]. Most *D. solani* strains isolated from different regions show a low level of genetic variation, suggesting a clonal origin [20]. The *D. solani* genomes share a high similarity and synteny with those of the model strain *D. dadantii* 3937, prompting comparison between the two species. Only a few hundred genes were specific to each species, including a few dozen distinctive genomic regions [16,17]. Three of these regions encode polyketide synthases (PKS), non-ribosomal peptide synthetases (NRPS) and amino acid adenylation domain proteins, which are typically involved in the production of secondary metabolites [16].

PKSs and NRPSs are able to synthesize molecules by sequential condensation of carboxylic

acids and amino acids, respectively. PKS and NRPS modules can combine together to form

hybrid PKS/NRPS systems capable of producing compounds of great structural diversity [21].

The molecules synthesized may have siderophore, antibiotic or phytotoxic properties that

promote the virulence of a plant pathogen. Three PKS/NRPS clusters are present in all

sequenced *D. solani* strains and found in a few other *Dickeya* species and related genera:

ooc, *zms*, and a cluster of unknown function [22]. In *Serratia plymuthica,* the cluster *ooc* is

involved in the synthesis of oocydin A, a halogenated macrolide with antifungal and anti-

oomycete activity [23]. The cluster *zms*, previously found in the genomes of *S. plymuthica* and

Dickeya oryzae, leads to the biosynthesis of a polyamino-amide antibiotic, zeamine [24,25].

The third cluster *ssm* (for *solani* secondary metabolite) is found in a few *Dickeya* species but

the nature and function of the synthesized molecule are unknown. To elucidate the

contribution of the *ssm*, *ooc* and *zms* clusters of *D. solani* in competition with other living

organisms, we constructed mutants of each of the three loci involved in the biosynthesis of

secondary metabolites in the highly virulent *D. solani* strain D s0432-1. We showed that the

D. solani D s0432-1 clusters *zms* and *ooc,* encoding zeamine and oocydin biosynthesis, are

involved in growth inhibition of bacteria and fungi, respectively. Recently we, and others

identified the third cluster as implicated in *Dickeya solani*'s ability to inhibit the growth of

different yeast species [26,27]. We named this cluster *ssm* (for *solani* secondary metabolite)

while Matilla and colleagues named it sol (for solanimycin, the molecule produced by this

cluster). Finally, the comparative analysis of the genomes and phenotypes of several wild-

type *D. solani* strains revealed the fundamental role of the small RNA ArcZ. This sRNA plays

a major role in the regulation of the *ssm* and *zms* clusters, which can be disrupted by a

mutation in the *arcZ* sequence, conserved in some natural isolates.

RESULTS

Description and distribution of the three selected PKS/NRPS secondary metabolite

clusters of *D. solani*.

The gene clusters encoding complex NRPS and PKS involved in the production of secondary

metabolites were named *ssm*, *ooc* and *zms,* respectively (Fig. 1A). Their distribution in

Dickeya species is summarized in figure 1B (see also Table S1 for a complete list of the

Dickeya strains with the indication of the presence or absence of each cluster).

The ~42-kbp cluster *ssm* contains the 12 genes *ssmABCDEFGHIJKL* (Fig. 1A). It is widely

conserved in the genus *Dickeya*, *i.e.*, in all sequenced *D. solani, D. aquatica, D. fangzhongdai,*

D. poaceiphila and *D. zeae* genomes, in some *D. dadantii* strains (such as NCPPB 898,

NCPPB 3537, DSM18020, but not the model strain 3937), in some *D. undicola* strains (FVG1,

FVG10), and in some *D. oryzae* strains (such as S20 and FVG03). The structure of the

metabolite produced from this cluster has not been elucidated yet but the function of this

cluster has been described by us in a 2021 preprint article [27] and others in 2022 [26]. The

ssm cluster produces solanimycin that targets yeast.

The ~80-kbp cluster *ooc* (Fig. 1A) is highly similar to the *oocBCDEFGJKLMNOPQRSTUVW*

cluster of *S. plymuthica* A153. It is present in all the sequenced genomes of *D. solani* and *D.*

dianthicola, in some *D. oryzae* strains (such as ZYY5, EC1, DZ2Q and ZJU1202) and in a few

other species. In *S. plymuthica* A153, disruption of this gene cluster abolished bioactivity

against the fungi *Verticillium dahliae* and the oomycetes *Pythium ultimum*. This cluster

produces oocydin A [23], a chlorinated macrolide, powerfully active against plant pathogenic

oomycetes [28]. Since various *D. solani* strains inhibit *V. dahliae* and *P. ultimum* growth [29],

it was suggested, on the basis of gene sequence homologies and similar cluster organization,

that *D. solani* also produces oocydin A.

The ~55-kbp cluster *zms* encodes mixed fatty acid synthase FAS/PKS and hybrid NRPS/PKS

enzymes (Fig.1A). Its genomic organization is identical to the *D. oryzae* EC1 zeamine cluster

and related to the *S. plymuthica* AS12 zeamine cluster [24,30]. While the

zmsABCDEFGIJKLMNPQRS cluster directing zeamine biosynthesis is present only in a few

D. oryzae strains, it is conserved in all the sequenced *D. solani* and *D. fangzhongdai* genomes,

which suggest secondary acquisition by horizontal gene transfer in *D. oryzae* [22,30]. After

reclassification of several *D. zeae* strains in the novel species *D. oryzae* [14], the *zms* cluster

appeared to be absent in the genomes of true *D. zeae* strains. It is present in some, but not

all, *D. oryzae* rice strains (ZYY5, EC1, DZ2Q and ZJU1202) and in only one *D. dadantii* strain

(A622-S1-A17). The zeamine biosynthetic clusters from *D. oryzae* EC1 and *D. solani* Ds0432-

1 share from 59 to 94% identity at individual protein level [30]. Zeamine-related antibiotics are

polyamino-amide molecules toxic to a wide range of pro- and eukaryotic organisms such as

bacteria, fungi, oomycetes, plants, and nematodes [31]. Mutation of the zeamine synthase

gene *zmsA* in *D. oryzae* EC1 attenuates the inhibition of rice seed germination [24] and

suppresses antibacterial activity against *E. coli* [24]. Zeamine produced by *S. plymuthica* kills

nematodes and yeast [32]. *D. solani* IPO 2222 can also kill the nematode *Caenorhabditis*

elegans but not as quickly as *S. plymuthica* [32].

To interrupt the synthesis of the secondary molecules produced by these three clusters, we

constructed from a Nal^R Gm^R derivative of WT *D. solani* D s0432-1 in-frame deletion mutants

inactivating a key gene of each cluster. The inhibitory effects of the mutants Δ*ssmG*, Δ*oocL*

and Δ*zmsA* against fungi, bacteria and yeasts were compared with that of the parental strain.

To further analyze possible synergistic effects, double mutants Δ*ssmG*Δ*oocL*, Δ*ssmG*Δ*zmsA*

and Δ*oocL*Δ*zmsA* as well as the triple mutant (hereafter called Δ3) were also constructed.

The *D. solani* oocydin cluster ooc inhibits *Ascomycota* growth, and the *ssm* cluster is

also involved in the inhibition of some fungi.

We compared the ability of the WT *D. solani* strain D s0432-1 and its derived mutants to inhibit

the growth of *Botrytis cinerea*, *Magnaporthe oryzae* and *Sclerotinia sclerotiorum*, three fungi-

like eukaryotes of the phylum *Ascomycota*. The center of a potato dextrose agar (PDA) plate

was inoculated with fungal mycelium, and 5 µl of overnight bacterial culture of each *D. solani*

strain grown in M63 minimal medium with sucrose was deposited at the periphery of the plate

(Fig. 2). The different *D. solani* strains showed similar overall growth in liquid culture (Fig. S1).

After incubation at 25°C for several days, we observed a growth inhibition of the three fungi

by the *D. solani* WT and the Δ*ssmG* and Δ*zmsA* mutants. In contrast, the Δ*oocL* mutant

seemed less effective: *B. cinerea* and *M. oryzae* mycelium growth was completely unaffected,

while a small zone of inhibition was observed for *S. sclerotiorum* (Fig. 2). This suggests that

the Δ*oocL* mutant may also produce another molecule that inhibits this fungus. The

phenotypes were fully restored in the reverted strain Δ*oocL/oocL*$^+$. Double and triple mutants

of the secondary metabolite clusters containing the *oocL* deletion showed no differences from

the Δ*oocL* mutant with *B. cinerea* and *M. oryzae*. However, the Δ*oocL*Δ*ssmG* double mutant

and the Δ3 triple mutant were completely unable to inhibit *S. sclerotiorum* growth (Fig. 2).

These data suggest that the *ooc* cluster is mainly responsible for the majority of *Ascomycota*

inhibition and that the *ssm* cluster is secondary involved in the inhibition of some fungi. In

contrast, the *zms* cluster does not seem to be involved in fungal inhibition.

The *D. solani* zeamine cluster inhibits bacterial growth.

The zeamine cluster of *D. oryzae* EC1 is responsible for the a bactericidal activity against

Escherichia coli DH5α [24]. *D. oryzae* EC1 and *D. solani* zeamine biosynthetic genes share a

high degree of similarity [30]. We thus evaluated the capacity of the *D. solani* strain D s0432-

1 and its derived mutants to inhibit the growth of gram-positive and gram-negative bacteria

(Fig. 3). Only the Δ*zmsA* mutant was unable to inhibit *Bacillus subtilis* growth, indicating that

WT *D. solani* produces an active zeamine antibiotic (Fig. 3). The WT phenotype was restored

in the reverted Δ*zmsA/zmsA*+ strain. We also tested the ability of *D. solani* to inhibit growth of

the other gram-positive bacteria *Streptomyces scabiei*, a plant pathogen causing the potato

disease common scab [33]. An inhibition of *S. scabiei* growth was observed, except with the

Δ*zmsA* mutant. No phenotype difference was observed between the single Δ*zmsA* mutant and

the double and triple mutants containing the *zmsA* deletion, suggesting that only the cluster

zms is required for Gram-positive bacterial growth inhibition. We also tested the ability of *D.*

solani to inhibit growth of the gram-negative bacteria *E. coli*, *D. dadantii* and *Pectobacterium*

atrosepticum (Fig. 3). *E. coli* DH5α was, like *B. subtilis* and *S. scabiei*, inhibited by the WT

strain through *zms* activity, albeit much less potently (Fig. 3 and Fig. S2). No inhibition of the

two other pectinolytic bacteria *D. dadantii* and *P. atrosepticum* growth was observed. In

conclusion, out of the three clusters analyzed, only the zeamine-like molecule potentially

produced by the *zms* cluster has an anti-bacterial activity against some gram-positive and

gram-negative bacteria.

The *D. solani* clusters *ssm* and *zms* inhibits yeast growth.

Since zeamine produced by *S. plymuthica* A153 has previously been shown to be toxic to the

ascomycete yeast *Saccharomyces cerevisiae* [32], we tested the capacity of *D. solani* D

s0432-1 and its derivative mutant to inhibit the growth of the yeasts *S. cerevisiae*,

Kluyveromyces lactis, a predominant eukaryote during cheese productions [34], and *Candida*

albicans, an opportunistic human pathogen. In a growth inhibition assay performed in Yeast

Peptone Dextrose (YPD) solid agar medium, we observed a strong inhibition of *K. lactis* by

WT *D. solani*, a lower inhibition of *S. cerevisiae*, and a slight inhibition of *C. albicans* (Fig. 4).

The $\Delta ssmG$ mutant was almost completely deficient in its ability to inhibit yeast growth, while

the $\Delta ssmG$/*ssmG*$^+$ revertant strain showed a restored WT phenotype. The cluster *ssm* is thus

primarily responsible for the anti-eukaryotic action against these yeasts (Fig. 4). Nevertheless,

a weak halo of inhibition of the yeasts *K. lactis* and *S. cerevisiae* was observable around the

mutant $\Delta ssmG.$ This halo disappeared around the double mutant $\Delta ssmG \Delta zmsA$ and the triple

mutant $\Delta 3,$ suggesting that the *zms* cluster is also implicated in yeast inhibition, but at a lower

level than that observed for the *ssm* cluster. In conclusion, although the zeamine cluster of *D.*

solani D s0432-1 is involved in the inhibition of yeast growth, as shown for *S. plymuthica* A153

[32], *D. solani* D s0432-1 possesses a second cluster *ssm* that produces another anti-yeast

molecule.

The WT *D. solani* strains IPO 2222 and IFB0223 are unable to inhibit yeast and bacterial growth.

We showed that *D. solani* D s0432-1 has the capacity to inhibit the growth of a wide range of microorganisms, including filamentous fungi, bacteria, and yeasts. We therefore tested whether other WT *D. solani* strains isolated from the environment in Europe, either from lesions of potato tubers or plants, or from potato rhizosphere, possess this ability. We focused our work on *D. solani* strains available in national strain collections and whose genome sequence is available. These strains are D s0432-1, IFB0099, IFB0158, IFB0223, IFB0484, IPO 3337, IPO 3494, IPO 3793 and the *D. solani* type strain IPO 2222. We tested their capacity to inhibit the bacterium *B. subtilis*, the yeast *K. lactis* and the fungus *S. sclerotiorum* (Fig. 5A). All strains were able to inhibit the growth of *S. sclerotiorum*. The inhibition of *B. subtilis* and *K. lactis* was clearly observed for all the strains except IPO 2222 and IFB0223. These strong phenotypic differences between IPO 2222 and IFB0223 and the other WT *D. solani* strains could not be explained by any nucleotide variabilities in the *ssm*, *zms*, and *ooc* clusters. All the selected strains carry the *ssm*, *zms*, and *ooc* clusters, with 100% conservation at the nucleotide level (including in the promoter regions). Nonetheless, there are certain changes at the genome level, most notably some nucleotide variability with the presence of SNPs (single-nucleotide polymorphism) and InDel (insertion or deletion), outside the clusters *ssm, zms,* and *ooc*. Hence, we searched for variations in the whole genomes by comparing two groups: strains IFB0223 and IPO 2222 that are defective for the growth inhibition of yeast and bacteria, and strains D s0432-1, IPO 3337 and IFB0099 that are efficient for these inhibitions (Table S2).

A SNP was observed with an identical nucleotide in the strains D s0432-1, IPO3337 and IFB0099 (a G at position 90 of *arcZ*) but different nucleotide in the strain IPO2222 (a A at position 90 of *arcZ*) (position 2530087 according to IPO 2222 genome). This variation is positioned in the 3' region of the small RNA *arcZ*, known to be a key area interacting with targeted mRNA for post-translational regulation in other bacterial species (Fig. 5C). The ArcZ variants of D s0432-1 and IPO 2222 will be referred to as $ArcZ_1$ and $ArcZ_2$ respectively in the

remainder of this article. In the strain IFB0223, we observed a 3-kbp chromosomal deletion

covering the gene *arcZ*, and extending between the genes *arcB* and *osmY* (Fig. 5B). In other

D. solani strains, this region includes *arcZ*, *elbB*, *mtgA*, and a gene encoding an alginate lyase

domain (pfam05426). These discoveries led us to hypothesize that the inability of strains

IFB0223 and IPO 2222 to inhibit yeast and bacterial growth could be due to an ArcZ deficiency.

Before going further, we ordered the *D. solani* IPO 2222 type strain stored in the BCCM

collection (LMG25993) to analyse its genotype and phenotype. Using PCR amplification and

sanger sequencing of *arcZ,* we confirmed that the *arcZ* allele of LMG 25993 is *arcZ$_2$*, similarly

to the IPO 2222 strain used in our experiments Similarly, the strain IPO 2222 used in our

experiments and LMG 25993 have the same inhibition phenotype, i.e., an incapacity to inhibit

yeast and bacterial growth (Fig. S3).

The *arcZ* gene is required for *zms* and *ssm* anti-microbial activities.

sRNAs are posttranscriptional regulators that most commonly influence gene

translation by base-pairing to target mRNAs. In *E. coli*, this regulation often requires the RNA

chaperones Hfq or ProQ [35–37]. ArcZ belongs to what is called the "core sRNAs" because it

is highly conserved among enterobacterial species [38]. In order to determine if ArcZ is

required for secondary metabolite production, the Δ*arcZ* mutants of *D. solani* strains D s0432-

1 and IPO 2222 were constructed, and their phenotypes compared to those of the WT strains

(Fig. 6A). In M63 sucrose, the mutants grew as well as the parental WT strains (Fig. S4). In

our inhibition assay against *B. subtilis* and *K. lactis,* no inhibition zones were observed around

the Δ*arcZ$_1$* mutant of D s0432-1 (Fig. 6A), demonstrating that ArcZ$_1$ is required for the

synthesis of the metabolites produced from the *ssm* and *zms* clusters in D s0432-1. However,

the Δ*arcZ$_1$* mutant of *D. solani* D s0432-1 was still able to inhibit the growth of the fungus *S.*

sclerotiorum, although a slight decrease in the anti-fungal effect was observed. Thus, ArcZ

does not appear to be fully required for *ooc* cluster expression, which is consistent with the

results obtained with *D. solani* IFB0223, the naturally *arcZ* deletion mutant capable of inhibiting

S. sclerotiorum growth (Fig. 5A and Fig. 6A). *D. solani* IPO 2222 WT inhibited neither bacteria

nor yeast but inhibited the growth of fungi, and the phenotype of the IPO 2222 $\Delta arcZ_2$ mutant

was similar to that of the IPO 2222 WT or the D s0432-1 $\Delta arcZ_1$ mutant. This suggests that

ArcZ$_2$ might be inactive.

To verify the effect of the G90A mutation in *arcZ*, we performed a heterologous

complementation experiment by transferring the allele $arcZ_1$ from D s0432-1 or $arcZ_2$ from IPO

2222 in *D. solani* WT strains D s0432-1 or IPO 2222 or their $\Delta arcZ$ derivatives. The alleles

$arcZ_1$ and $arcZ_2$ were also transferred into *D. solani* IFB0223 naturally deleted for *arcZ*.

The G90A mutation in arcZ$_2$ of *D. solani* IPO 2222 causes a loss of function of *zms* and

***ssm* clusters.**

The two allelic forms $arcZ_1$ and $arcZ_2$ were cloned with their own promoter in the plasmid

pWSK29-oriT, a mobilizable low copy plasmid with pSC101 origin and ampicillin resistance

gene. DNA sequencing confirmed that the only difference between the two plasmids is the

G90A mutation in pWSK29-oriT-$arcZ_2$. The recombinant plasmids and the empty vector were

transferred by conjugation into WT or $\Delta arcZ$ *D. solani* D s0432-1 or IPO 2222 strains, as well

as WT *D. solani* IFB0223 (naturally deleted for *arcZ*). The phenotypes of the conjugants were

determined by testing their anti-microbial activity against *B. subtilis*, *K. lactis* and *S.*

sclerotiorum (Fig. 6B).

The transfer of $arcZ_1$ in $\Delta arcZ_1$ Ds0432-1 restored the WT phenotypes, proving that the allele

$arcZ_1$ activates the anti-microbial activities of the *ssm* and *zms* clusters, while the anti-fungal

activity was unaffected. In contrast, the transfer of $arcZ_2$ in the $\Delta arcZ_1$ DS0432-1 strain did not

restore the WT phenotypes. Thus, the difference in complementation between $arcZ_1$ and $arcZ_2$

is due to the presence of the G90A mutation that impairs ArcZ$_2$ function.

An identical result was obtained with the $\Delta arcZ_2$ mutant of IPO 2222 or the IFB0223 strain

carrying the pWSK29-oriT, pWSK29-oriT-$arcZ_1$ and pWSK29-oriT-$arcZ_2$. Indeed, the transfer

of $arcZ_1$ in strains WT IPO 2222, $\Delta arcZ_2$ IPO 2222 or IFB0223 activates the anti-microbial

activities of the *ssm* and *zms* clusters. In contrast, the transfer of $arcZ_2$ in these strains did not

activate the *ssm* and *zms* anti-microbial activities. These data demonstrated that the presence

of *arcZ* and the nature of the *arcZ* allele are the main factors governing the difference in

phenotypes for antimicrobial activities between the *D. solani* strains D s0432-1 and IPO 2222

or IFB0223.

Finally, we questioned if an *arcZ* mutation could be found in additional *D. solani* natural

isolates. We extended this genomic analysis to 57 *D. solani* genomes, including IPO 2222. It

revealed that only four strains have an *arcZ* single-nucleotide mutation, all at different positions

in the 3' region of *arcZ*. It is not known whether these other mutations affect the antimicrobial

activity of these strains. The *arcZ* DNA sequence of *D. solani* IPO 2222 was also used as a

query in a BlastN search against the NCBI nucleotide database. Three *D. fangzhongdai strains*

DSM 101947 (CP025003.1), QZH3 (CP031507.1), and LN1 (CP031505.1) had the G90A

mutation in *arcZ*. A G90T mutation was also discovered in *D. parazeae* Ech586 (CP001836.1)

(Fig. S5). Therefore, the G90 mutation to A or T of *arcZ* can be found in other *Dickeya* species.

The G90A mutation in *arcZ$_2$* of *D. solani* IPO 2222 alters ArcZ$_2$ processing.

To gain further insight into the consequence of the G90A mutation in *arcZ*, Northern-blot

experiments were performed to detect the sRNAs ArcZ$_1$ in WT D s0432-1 and ArcZ$_2$ in WT

IPO 2222 (Fig. 7). A probe specific to the 3' part of ArcZ$_1$ and ArcZ$_2$ was used (annealing from

nucleotide 91 to 132). An abundant smaller transcript (~60nt) representing the processed 3′

fragment of ArcZ$_1$ was detected in WT D s0432-1 but not in the mutant D s0432-1 Δ*arcZ$_1$* (Fig.

7, compared lanes 1 and 2). The full-length form corresponding to the ArcZ$_1$ precursor was

barely detected. This confirms that the processed form of ArcZ$_1$ is predominant, like observed

in *E. coli*, *Salmonella*, and *Photorhabdus* [38–40]. In IPO 2222, only the full-length ArcZ$_2$ was

detected in the WT strain whereas no ArcZ$_2$ transcript was detected in the IPO 2222 Δ*arcZ$_2$*

mutant (Fig. 7, compared lanes 5 and 6). This result suggests that the G90A mutation present

in the allele *arcZ$_2$* of IPO 2222 prevents ArcZ$_2$ processing in a shorter functional transcript.

Next, we wondered if we could relate the importance of ArcZ processing to the activation of

the *ssm* and *zms* clusters. To do that, Northern-blot analyses were realized with the *D. solani*

$\Delta arcZ$ strains expressing either the $arcZ_1$ or $arcZ_2$ allele from the plasmid pWSK29-oriT-$arcZ$.

On one hand, Northern blot showed that D s0432-1$\Delta arcZ_1$, IPO 2222 $\Delta arcZ_2$ and IFB0223

(natural $\Delta arcZ$) expressing $arcZ_1$ from D s0432-1 accumulate the processed short transcript

of ArcZ$_1$ (Fig. 7, lanes 3, 7 and 10) and have the capacity to inhibit yeast and bacteria (Fig. 6).

On the other hand, D s0432-1$\Delta arcZ_1$, IPO 2222 $\Delta arcZ_2$ and IFB0223 expressing $arcZ_2$ from

IPO 2222 showed the accumulation of the non-processed transcript of ArcZ$_2$ (Fig. 7, lanes 3,

7 and 10) and an inability to inhibit yeast and bacteria (Fig. 6). These results supported the

hypothesis that the G90A mutation in ArcZ alters the processing of the ArcZ precursor in a

shorter form, leading to its inability to play a correct regulatory function on the ssm and zms

clusters.

DISCUSSION

PKS/NRPS secondary metabolite pathways are a great source of molecules with anti-

eukaryotic or anti-prokaryotic activity, giving the bacteria that synthesize them a competitive

advantage over other organisms. Targeted mutagenesis of the *D. solani* D s0432-1

chromosome has allowed us to specifically study the involvement of three secondary

metabolite biosynthesis clusters encoded by all the *D. solani* strains for which genome

sequence is available. We focused our work on strain D s0432-1, one of the most virulent *D.*

solani strains [18]. We have demonstrated that *D. solani* D s0432-1 is able to inhibit the growth

of a variety of living microorganisms. We tested bio-activities against Gram-negative bacteria

(*E. coli*, *P. atrosepticum*, *D. dadantii*), Gram-positive bacteria (*B. subtilis*, *S. scabiei*), yeasts

(*S. cerevisiae*, *K. lactis, C. albicans*), and fungi (*B. cinerea*, *M. oryzae* and *S. sclerotiorum*).

The cluster *ooc* encodes a biosynthesis pathway that produces an oocydin-like molecule, a

chlorinated macrocyclic lactone molecule having antifungal, antioomycete and antitumor

activities in *S. plymuthica* [23,29]. It was previously shown that the *D. solani* strains MK10,

MK16 and IPO 2222, all encoding the oocydin cluster, inhibit the growth of the fungus *V.*

dahliae and the oomycete *P. ultimum* [29]. Our study showed that *D. solani* D s0432-1

prevents the growth of the fungi *B. cinerea*, *M. oryzae* and *S. sclerotiorum*, and that inactivation

of the *oocL* gene suppressed the inhibition of the three fungi without any visible effect on the

bacteria or yeasts tested.

By testing other mutants of the other secondary metabolite clusters, we showed that the

mutant Δ*ssmG* was nearly completely unable of preventing the growth of *S. cerevisiae* and *K.*

lactis. Therefore, the *ssm* cluster appears to be responsible for yeast inhibition. *D. solani* D

s0432-1 also inhibits the human pathogen yeast *C. albicans*, but its inhibitory activity appears

to be weaker than that observed with *S. cerevisiae* and *K. lactis* in our inhibition assay. This

observation on the *ssm* cluster paves the way for the identification of a new molecule with

highly specific inhibitory activity against yeasts. The *ssm* cluster is conserved in several

Dickeya genomes but rarely in other bacterial genera, except in some *Rouxiella* species. The

structure of the molecule and its target remains to be elucidated.

Our study also showed that *D. solani* D s0432-1 has an anti-bacterial activity linked to the

cluster *zms*, which encodes a zeamine biosynthetic pathway. Zeamine produced by *S.*

plymuthica kills *B. subtilis,* the yeasts *S. cerevisiae* and *Schizosaccharomyces pombe,* and

the nematode *Caenorhabditis elegans* [32]. We showed that *D. solani* D s0432-1, but not the

Δ*zmsA* mutant, can clearly inhibit the growth of two Gram-positive bacteria, *B. subtilis* and *S.*

scabies. Since *D. solani* and *S. scabiei* are two potato pathogens, they might be in competition

for the same ecological niche. It is thus interesting to note that *D. solani* can inhibit the growth

of another potato pathogen. We also observed a slight inhibition exerted by *D. solani* D s0432-

1 against *E. coli*, but no inhibition towards *D. dadantii* and *P. atrosepticum*. Zeamine resistance

of *D. dadantii* could be explained by the presence in its genome of the genes *desAB* encoding

an RND pump identified in *D. oryzae* EC1 and involved in zeamine efflux [41]. In-frame

deletion of *desA* or *desB* in *D. oryzae* EC1 leads to a zeamine sensitive phenotype [41]. In *in*

planta Tn-seq experiments with *D. dadantii* 3937 [42], mutants of the genes *desA* and *desB*

(Dda3937_00787 and Dda3937_00786 respectively) did not display a significant negative or

positive variation (Log2 fold-changes -0.09 and +0.44, respectively), suggesting that this RND

efflux pump does not play a significant role during *D. dadantii* chicory infection, contrary to the

D. dadantii RND efflux pump AcrAB that appeared to be essential for virulence [42]. While *P.*

atrosepticum is not inhibited by *D. solani* D s0432-1, it does not have the genes *desAB*.

Zeamine resistance could be provided by another efflux pump or by a different mechanism in

P. atrosepticum. It's worth remembering that the *D. solani* D s0432-1 Δ*zmsA* mutant grows

slightly faster than the WT strain after reaching OD_{600nm} 0.4 (Fig. S1). This small difference in

growth might be explained by the lack of need of the mutant to adapt to the presence of

zeamine in the medium. We also observed that *D. solani* D s0432-1 Δ*ssmG* produced a very

slight halo of inhibition against *S. cerevisiae* and *K. lactis*, barely visible in our tests. This weak

halo disappears in the double mutant Δ*ssmG*Δ*zmsA* and the triple mutant

Δ*ssmG*Δ*oocL*Δ*zmsA,* showing that the zeamine produced by the *D. solani zms* cluster has

also an anti-yeast action. It is worthwhile to remind that zeamine produced by *S. plymuthica*

A153 is bioactive against *S. cerevisiae* and *S. pombe* [32].

Thus, the presence of three clusters encoding secondary metabolites allows *D. solani* D

s0432-1 to produce an arsenal of bioactive secondary metabolites against a variety of living

microorganisms. The cluster *ssm* produces an unkown molecule active against yeasts; the

cluster *ooc* produces an oocydin-like molecule active against fungi and the cluster *zms*

produces a zeamine-like molecule active against bacteria and yeast. In future studies, It would

be instructive to investigate *D. solani* behavior toward other organisms found in the

environment, such as amoebas, nematodes, or even insects.

We then expanded our studies to a larger set of *D. solani* strains isolated in different European

countries and years. Through the combination of phenotypic analysis and comparative

genomics, confirmed by mutant construction, we demonstrated that the ArcZ sRNA plays a

major role in activating the antimicrobial activities of the clusters *ssm* and *zms*. We first

observed that some *D. solani* WT strains are defective for yeast inhibition and possess

different *arcZ* alleles, a deletion in the case of strain IFB0223, or the presence of a SNP G90A

in the 3' part of *arcZ* in the type strain IPO 2222. We demonstrated that these arcZ alterations

significantly impair the *D. solani* ability to inhibit other bacteria and yeasts. These *arcZ*-related

genomic changes represent a minority of *D. solani* isolates, only two of the nine strains

phenotypically tested . Hence, our results show that despite a probable clonal origin,

environmental strains of *D. solani* can present very different characteristics due to a single

mutation into a regulatory sRNA. The *arcZ* polymorphism may be one of the explanations for

the variability of phenotypes and virulence observed between genetically close *D. solani*

strains. A global investigation of the polymorphism of all the short RNAs within the genomes

of the same species could give surprising results. This study shows that it is critical to pay

attention to the allelic diversity due to SNPs present not only in protein coding genes but also

in sRNA genes.

ArcZ is a well-known regulatory sRNA that is associated with the chaperone Hfq. The ArcZ

primary transcript (121 to 134-nt) is converted by the essential endoribonuclease RNAse E

[43] to a shorter stable approximative 60-nt RNA retaining the seed region. The precise length

of this short processed RNAs depends on the bacterial species [38,40,44,45]. In *E. coli*, the

processed short form of ArcZ promotes *rpoS* translation while suppressing the expression of

several other genes. In anaerobic conditions, *arcZ* is repressed and translation of RpoS is low

[44]. RpoS is the RNA polymerase alternative sigma factor that governs the general stress

response, which is activated by a wide range of conditions. It controls 10% of the *E. coli*

genome [46]. RpoS can control positively or negatively the production of metabolites such as

antibiotics [29,47]. *rpoS* mRNA is not the only ArcZ direct target. In *Photorhabdus* and

Xenorhabdus, ArcZ pairs with the mRNA encoding HexA, a transcriptional repressor of the

expression of specific metabolite gene clusters [40]. In *Dickeya* species, the regulator gene

pecT is the orthologous gene to *hexA*. In *D. dadantii* 3937, the *pecT* mRNA is also a direct

target of ArcZ. Short processed ArcZ binds to Hfq and directly to the 5'UTR region of *pecT*

mRNA, repressing its translation [45]. PecT is directly and indirectly involved in the expression

of genes encoding plant cell wall degrading enzymes and virulence factors [45]. In the

phytopathogen *Erwinia amylovora,* ArcZ participates also in the positive control of T3SS,

exopolysaccharide production, biofilm formation, and motility [48]. Finally, another known

direct target of ArcZ is the *flhD* mRNA [49,50]. FlhD forms with FlhC the master regulator

FlhDC that controls motility in bacteria [51]. Although we demonstrated the critical role of ArcZ

in the regulation of secondary metabolite production in *D. solani*, it remains to understand the

precise mechanism of ArcZ action on the ssm or zms clusters. We can only speculate that this

could occur through the control of *rpoS*, *flhD*, or *pecT*, three regulators known to be ArcZ

targets. However, other actors could be involved. Another point to consider is a more global

role of *D. solani* ArcZ exerted on other cellular functions such as motility and virulence. ArcZ

targets in *D. solani* need to be investigated in the future. It is worth pointing out that the G90

nucleotide is in the 3' region of ArcZ that interacts with the target mRNAs *rpoS*, *flhD*, or *pecT*

in other bacteria (Fig. S6). Northern-blot analysis revealed that ArcZ$_2$ of *D. solani* IPO 2222 is

not processed in the stable shorter sRNA as ArcZ$_1$ of *D. solani* D s0432-1. The G90A mutation

in IPO 2222 ArcZ$_2$ is predicted to modify the secondary structure of the ArcZ precursor form

(Fig. 7B). Since maturation by RNase E is essential for target regulation by the ArcZ sRNA in

Salmonella, we conclude that the absence of a short form of ArcZ in IPO 2222 is probably the

main cause of the loss of regulatory function of the IPO 2222 ArcZ$_2$.

The type strain *D. solani* IPO 2222 is largely used in academic laboratories as a model to

study the *D. solani* species. In light of our findings, it could be important to compare data

obtained with the strain IPO 2222 with those obtained with other *D. solani* strains. The *arcZ*

mutations may be responsible, at least partially, for the lower virulence of IPO 2222 and

IFB0223 observed previously in comparison to D s0432-1 [18]. Even strains with the same

arcZ, *ssm*, *ooc*, and *zms* genomic sequences as D s0432-1, showed heterogeneity in their

ability to inhibit other micro-organisms. For example, IPO 3494 and IFB0484 inhibited *B.*

subtilis better than other *D. solani* strains (Fig. 5A). Other specific mutations in the genomes

of these strains, which could also modify the activity of secondary metabolite clusters, could

explain these phenotypic variations.

We first described the inhibitory function of the *ssm* cluster of the *D. solani* strain D s0432-1

against the yeasts *S. cerevisiae* and *K. lactis* in a 2021 bioRxiv preprint [27]. Very recently, in

November 2022, Matilla *et al.* published results on the *ssm* cluster of the *D. solani* strain MK10

[26]. The *ssm* cluster was named *sol* in this study and the authors described the antifungal

activity of the uncharacterized molecule that they called solanimycin [26]. Matilla *et al.* also

report the inhibition observed with the *D. solani* strain IPO 2222 against *Verticillium dahliae*

and *Schizosaccharomyces pombe* ([26], Fig S1). In our conditions, strain IPO 2222 was able to inhibit the fungus *Sclerotinia sclerotiorum* but not the yeast *Kluyveromyces lactis* (Fig. 5). In comparison to our study, Matilla *et al* used different fungal species but also different growth conditions (medium, temperature, exposure time, etc). By random mutagenesis of strain MK10, they identified RsmA and the two *Dickeya* quorum-sensing systems ExpI/R and Vfm as regulators of the *ssm/sol* cluster such [26]. Interestingly, the Rsm regulatory pathway of *D. dadantii* 3937 is regulated by PecT, which is under the ArcZ control [45]. Further studies are now necessary to specify the biological targets of solanimycin and to clarify the role in *ssm/sol* expression of the sRNA ArcZ, quorum-sensing, and transcriptional regulators PecT or RsmA, in order to better understand the regulatory network controlling the production of secondary metabolites in *D. solani*. Our lab is currently investigating the regulatory pathway of secondary metabolite clusters from ArcZ.

In conclusion, our results provide the basis for further investigation of a novel metabolite capable of limiting yeast growth and, at the same time, open doors to understand how the ArcZ sRNA acts as a global regulator to regulate two different clusters of secondary metabolites in *D. solani*. It also shows how a single mutation in a small RNA can greatly modify the phenotype of bacterial strains that are very close phylogenetically.

MATERIAL AND METHODS

Bacterial and fungal strains, plasmids and growth conditions.

The *E. coli* and *Dickeya* bacterial strains, plasmids and oligonucleotides used in this study are described in Table S3 and S4. The genome accession number of *D. solani* D s0432-1 is NZ_CP017453. The following strains have been used in the study: *Sclerotinia sclerotiorum* S5, *Botrytis cinerea* B05.10, *Magnaporthe oryzae* Guy11, *Saccharomyces cerevisiae* BY4743 (*MATa/α his3Δ1/his3Δ1 leu2Δ0/leu2Δ0 LYS2/lys2Δ0 met15Δ0/MET15 ura3Δ0/ura3Δ0*), *Kluyveromyces lactis* MWL9S1 [52], *Candida albicans* SC5314, *Dickeya dadantii* 3937, *Pectobacterium atrosepticum* SCRI1043, *Streptomyces scabiei* CFBP4517, *Bacillus subtilis* PY79. *E. coli* was grown routinely at 37°C in LB. Fungus strains were grown at 25°C onto

Potato Dextrose Agar (PDA). *S. scabiei* was grown in tryptic soy broth (TSB) medium at 28°C.

B. subtilis, *P. atrosepticum* and the *Dickeya* strains were cultivated in LB unless specified.

Yeast cells were grown at 30°C in rich medium consisting of complete yeast extract-peptone

(YP) medium containing 1% Bacto yeast extract, 1% Bacto peptone (Difco) supplemented with

2% glucose (yeast extract-peptone-dextrose [YPD] medium). For the bacteria, yeast and fungi

inhibition assay, M63 medium supplemented with sucrose (2 g $(NH_4)_2SO_4$, 13.6 g KH_2PO_4, 2.5

mg $FeSO_47H_2O$, 0.2 g $MgSO_47H_2O$, 10 g sucrose, per liter) were employed to grow overnight

the *D. solani* strains before performing the assay.

When required, antibiotics were added at the following concentrations: ampicillin (Amp), 100

µg/L; nalidixic acid (Nal), 10 µg/mL. Diaminopimelic acid (DAP) (57 µg/mL) was added for the

growth of the *E. coli* MFDpir strain. Media were solidified with 12 g/L agar.

Growth inhibition assay of bacteria and yeast.

D. solani strains were grown for 24 h at 30°C with shaking in M63 medium supplemented with

1% sucrose. *B. subtilis*, *E. coli*, *D. dadantii* and *P. atrosepticum* were grown in LB at 30°C with

shaking overnight. The next day, the OD_{600} of the cultures of *D. solani* were adjusted to 2. The

temperature of melted LB agar was lowered to around 40°C (just before the agar re-solidified).

100 mL of the LB agar in surfusion were mixed with 100 µl of the OD_{600} 1 culture of *B. subtilis*,

E. coli, *D. dadantii* or *P. atrosepticum*. 30 mL of inoculated LB agar were poured in 12- by 12-

cm square plates. Then 5 µl of the OD_{600} 2 cultures of *D. solani* were spotted onto the

inoculated square plates which were incubated at 30°C for 24-48 h before visualization of the

inhibition zone.

The same experiment was conducted with *S. scabiei*, except that it was grown for 3 days in

TSB at 28°C and TSB agar was poured in the square plates. With the yeasts *S. cerevisiae*, *K.*

lactis, and *C. albicans*, a similar protocol was used with YPD medium and with OD_{600} 2 culture

of yeast per 30mL of medium.

Growth inhibition of fungal strains.

S. sclerotiorum, *B. cinerea* and *M. oryzae* were grown onto PDA plates at 25°C for 5, 7 and

10 days respectively. *D. solani* strains were grown for 24 h at 30°C with shaking in M63

medium supplemented with 1% sucrose. The OD_{600} of these overnight cultures were adjusted

to 2. Then, 5 µl of the bacterial suspensions were spotted onto PDA plates with 5 mm agar

plugs of fungus at the center of the Petri dish. The radius of fungus inhibition zone was

measured.

Construction of the MLSA tree positioning strains within the *Dickeya* genus.

Details on the methods used for the construction of MLSA tree positioning strains are given in

Supplementary Material and Methods.

Constructions of the mutant strains and plasmids

All the material and methods for the construction of the strains and plasmids used in this study

are given in Supplementary Material and Methods.

RNA isolation and Northern detection

All the material and methods for RNA isolation and Northern detection performed in this study

are given in Supplementary Material and Methods.

ACKNOWLEDGMENTS

B.T. was supported by a PhD grant from the Ministère de l'Enseignement Supérieur, de la

Recherche et de l'Innovation. We thank Veronique Utzinger for technical assistance, the

members of the Membrane Trafficking and Signaling in Bacteria (MTSB) team for discussions,

Amélie De Vallée, Nathalie Poussereau and Christophe Bruel for advices on fungi growth and

providing the strains, Alexandre Soulard, Marc Lemaire and Jade Ravent for advices on yeast

growth and providing the strains and medium. We would also like to thank the past reviewers

whose comments allowed us to greatly improve this study.

AUTHOR CONTRIBUTION

B.T., E.G. B.J. A.L., G.C., C.J., G.E carried out the experiments. G.E wrote the manuscript with support from B.T, E.G., A.L., R.M., F.D. and H-C.P.N.

F.D., C.J. H-C-P.N. and R.M. performed the phylogenetic and bioinformatic analyses. G.E conceived the original idea with advises from H-C-P.N. G.E supervised the project. All authors provided critical feedback and helped shape the research, analysis and manuscript.

CONFLICT OF INTEREST

The authors state that the research was conducted in the absence of any commercial or financial relationship that could be interpreted as a potential conflict of interest.

FUNDING

This work was supported by a grant from Agence Nationale de la Recherche to L.A. (Project RNAchap, ANR-17-CE11-0009-01) and to E.G. (Project Tn-Phyto, ANR- 19-CE35-0016). E.G was also supported by the FR BioEEnVis, annual credits from the University Lyon I and the CNRS at regular basis.

REFERENCES

1. Hugouvieux-Cotte-Pattat N, Condemine G, Gueguen E, Shevchik VE. Dickeya Plant Pathogens. eLS. American Cancer Society; 2020. pp. 1–10. doi:10.1002/9780470015902.a0028932

2. Charkowski A, Blanco C, Condemine G, Expert D, Franza T, Hayes C, et al. The role of secretion systems and small molecules in soft-rot Enterobacteriaceae pathogenicity. Annu Rev Phytopathol. 2012;50: 425–449. doi:10.1146/annurev-phyto-081211-173013

3. Van Gijsegem F, Toth IK, van der Wolf JM. Soft Rot Pectobacteriaceae: A Brief Overview. In: Van Gijsegem F, van der Wolf JM, Toth IK, editors. Plant Diseases Caused by Dickeya and Pectobacterium Species. Cham: Springer International Publishing; 2021. pp. 1–11. doi:10.1007/978-3-030-61459-1_1

4. Hugouvieux-Cotte-Pattat N, Condemine G, Shevchik VE. Bacterial pectate lyases, structural and functional diversity. Environ Microbiol Rep. 2014;6: 427–440.

5. Samson R, Legendre JB, Christen R, Saux MF-L, Achouak W, Gardan L. Transfer of Pectobacterium chrysanthemi (Burkholder et al. 1953) Brenner et al. 1973 and Brenneria paradisiaca to the genus Dickeya gen. nov. as Dickeya chrysanthemi comb.

nov. and Dickeya paradisiaca comb. nov. and delineation of four novel species,
Dickeya dadantii sp. nov., Dickeya dianthicola sp. nov., Dickeya dieffenbachiae sp. nov.
and Dickeya zeae sp. nov. Int J Syst Evol Microbiol. 2005;55: 1415–1427.
doi:10.1099/ijs.0.02791-0

6. Hugouvieux-Cotte-Pattat N, Van Gijsegem F. Diversity within the Dickeya zeae
complex, identification of Dickeya zeae and Dickeya oryzae members, proposal of the
novel species Dickeya parazeae sp. nov. Int J Syst Evol Microbiol. 2021;71.
doi:10.1099/ijsem.0.005059

7. Brady CL, Cleenwerck I, Denman S, Venter SN, Rodríguez-Palenzuela P, Coutinho TA,
et al. Proposal to reclassify Brenneria quercina (Hildebrand and Schroth 1967) Hauben
et al. 1999 into a new genus, Lonsdalea gen. nov., as Lonsdalea quercina comb. nov.,
descriptions of Lonsdalea quercina subsp. quercina comb. nov., Lonsdalea quercina
subsp. iberica subsp. nov. and Lonsdalea quercina subsp. britannica subsp. nov.,
emendation of the description of the genus Brenneria, reclassification of Dickeya
dieffenbachiae as Dickeya dadantii subsp. dieffenbachiae comb. nov., and emendation
of the description of Dickeya dadantii. Int J Syst Evol Microbiol. 2012;62: 1592–1602.
doi:10.1099/ijs.0.035055-0

8. Parkinson N, DeVos P, Pirhonen M, Elphinstone J. Dickeya aquatica sp. nov., isolated
from waterways. Int J Syst Evol Microbiol. 2014;64: 2264–2266.
doi:10.1099/ijs.0.058693-0

9. van der Wolf JM, Nijhuis EH, Kowalewska MJ, Saddler GS, Parkinson N, Elphinstone
JG, et al. Dickeya solani sp. nov., a pectinolytic plant-pathogenic bacterium isolated
from potato (Solanum tuberosum). Int J Syst Evol Microbiol. 2014;64: 768–774.
doi:10.1099/ijs.0.052944-0

10. Tian Y, Zhao Y, Yuan X, Yi J, Fan J, Xu Z, et al. Dickeyafangzhongdai sp. nov., a plant-
pathogenic bacterium isolated from pear trees (Pyrus pyrifolia). Int J Syst Evol
Microbiol. 2016;66: 2831–2835. doi:10.1099/ijsem.0.001060

11. Hugouvieux-Cotte-Pattat N, Jacot-des-Combes C, Briolay J. Dickeya lacustris sp. nov.,
a water-living pectinolytic bacterium isolated from lakes in France. Int J Syst Evol
Microbiol. 2019;69: 721–726. doi:10.1099/ijsem.0.003208

12. Hugouvieux-Cotte-Pattat N, Brochier-Armanet C, Flandrois J-P, Reverchon S. Dickeya
poaceiphila sp. nov., a plant-pathogenic bacterium isolated from sugar cane
(Saccharum officinarum). Int J Syst Evol Microbiol. 2020;70: 4508–4514.
doi:10.1099/ijsem.0.004306

13. Oulghazi S, Pédron J, Cigna J, Lau YY, Moumni M, Van Gijsegem F, et al. Dickeya
undicola sp. nov., a novel species for pectinolytic isolates from surface waters in
Europe and Asia. Int J Syst Evol Microbiol. 2019;69: 2440–2444.
doi:10.1099/ijsem.0.003497

14. Wang X, He S-W, Guo H-B, Han J-G, Thin KK, Gao J-S, et al. Dickeya oryzae sp. nov.,
isolated from the roots of rice. Int J Syst Evol Microbiol. 2020;70: 4171–4178.
doi:10.1099/ijsem.0.004265

15. Blin P, Robic K, Khayi S, Cigna J, Munier E, Dewaegeneire P, et al. Pattern and causes
of the establishment of the invasive bacterial potato pathogen Dickeya solani and of the
maintenance of the resident pathogen D. dianthicola. Mol Ecol. 2021;30: 608–624.
doi:https://doi.org/10.1111/mec.15751

16. Garlant L, Koskinen P, Rouhiainen L, Laine P, Paulin L, Auvinen P, et al. Genome
Sequence of Dickeya solani, a New soft Rot Pathogen of Potato, Suggests its
Emergence May Be Related to a Novel Combination of Non-Ribosomal
Peptide/Polyketide Synthetase Clusters. Diversity. 2013;5: 824–842.
doi:10.3390/d5040824

17. Pédron J, Mondy S, des Essarts YR, Van Gijsegem F, Faure D. Genomic and
metabolic comparison with Dickeya dadantii 3937 reveals the emerging Dickeya solani
potato pathogen to display distinctive metabolic activities and T5SS/T6SS-related toxin
repertoire. BMC Genomics. 2014;15: 283. doi:10.1186/1471-2164-15-283

18. Golanowska M, Potrykus M, Motyka-Pomagruk A, Kabza M, Bacci G, Galardini M, et al.
Comparison of Highly and Weakly Virulent Dickeya solani Strains, With a View on the
Pangenome and Panregulon of This Species. Front Microbiol. 2018;9: 1940.
doi:10.3389/fmicb.2018.01940

19. Motyka-Pomagruk A, Zoledowska S, Misztak AE, Sledz W, Mengoni A, Lojkowska E.
Comparative genomics and pangenome-oriented studies reveal high homogeneity of
the agronomically relevant enterobacterial plant pathogen Dickeya solani. BMC
Genomics. 2020;21: 449. doi:10.1186/s12864-020-06863-w

20. Khayi S, Blin P, Pédron J, Chong T-M, Chan K-G, Moumni M, et al. Population
genomics reveals additive and replacing horizontal gene transfers in the emerging
pathogen Dickeya solani. BMC Genomics. 2015;16: 788. doi:10.1186/s12864-015-
1997-z

21. Cane DE, Walsh CT. The parallel and convergent universes of polyketide synthases
and nonribosomal peptide synthetases. Chem Biol. 1999;6: R319–R325.
doi:10.1016/S1074-5521(00)80001-0

22. Duprey A, Taib N, Leonard S, Garin T, Flandrois JP, Nasser W, et al. The
phytopathogenic nature of Dickeya aquatica 174/2 and the dynamic early evolution of
Dickeya pathogenicity. Environ Microbiol. 2019;21: 2809–2835. doi:10.1111/1462-
2920.14627

23. Matilla MA, Stöckmann H, Leeper FJ, Salmond GPC. Bacterial Biosynthetic Gene
Clusters Encoding the Anti-cancer Haterumalide Class of Molecules. J Biol Chem.
2012;287: 39125–39138. doi:10.1074/jbc.M112.401026

24. Zhou J, Zhang H, Wu J, Liu Q, Xi P, Lee J, et al. A Novel Multidomain Polyketide
Synthase Is Essential for Zeamine Production and the Virulence of Dickeya zeae. Mol
Plant-Microbe Interactions®. 2011;24: 1156–1164. doi:10.1094/MPMI-04-11-0087

25. Masschelein J, Mattheus W, Gao L-J, Moons P, Houdt RV, Uytterhoeven B, et al. A
PKS/NRPS/FAS Hybrid Gene Cluster from Serratia plymuthica RVH1 Encoding the
Biosynthesis of Three Broad Spectrum, Zeamine-Related Antibiotics. PLOS ONE.
2013;8: e54143. doi:10.1371/journal.pone.0054143

26. Matilla MA, Monson RE, Murphy A, Schicketanz M, Rawlinson A, Duncan C, et al.
Solanimycin: Biosynthesis and Distribution of a New Antifungal Antibiotic Regulated by
Two Quorum-Sensing Systems. Sperandio V, editor. mBio. 2022; e02472-22.
doi:10.1128/mbio.02472-22

27. Effantin G, Brual T, Rahbé Y, Hugouvieux-Cotte-Pattat N, Gueguen E. Dickeya solani
D s0432-1 produces an arsenal of secondary metabolites with anti-prokaryotic and anti-

eukaryotic activities against bacteria, yeasts, fungi, and aphids. bioRxiv; 2021. p.
2021.07.19.452942. doi:10.1101/2021.07.19.452942

28. Strobel G, Li J-Y, Sugawara F, Koshino H, Harper J, Hess WM. Oocydin A, a
chlorinated macrocyclic lactone with potent anti- oomycete activity from Serratia
marcescens. Microbiology. 1999;145: 3557–3564. doi:10.1099/00221287-145-12-3557

29. Matilla MA, Leeper FJ, Salmond GPC. Biosynthesis of the antifungal haterumalide,
oocydin A, in Serratia, and its regulation by quorum sensing, RpoS and Hfq. Environ
Microbiol. 2015;17: 2993–3008. doi:10.1111/1462-2920.12839

30. Zhou J, Cheng Y, Lv M, Liao L, Chen Y, Gu Y, et al. The complete genome sequence
of Dickeya zeae EC1 reveals substantial divergence from other Dickeya strains and
species. BMC Genomics. 2015;16: 571. doi:10.1186/s12864-015-1545-x

31. Masschelein J, Jenner M, Challis GL. Antibiotics from Gram-negative bacteria: a
comprehensive overview and selected biosynthetic highlights. Nat Prod Rep. 2017;34:
712–783. doi:10.1039/C7NP00010C

32. Hellberg JEEU, Matilla MA, Salmond GPC. The broad-spectrum antibiotic, zeamine,
kills the nematode worm Caenorhabditis elegans. Front Microbiol. 2015;6.
doi:10.3389/fmicb.2015.00137

33. Loria R, Kers J, Joshi M. Evolution of Plant Pathogenicity in Streptomyces. Annu Rev
Phytopathol. 2006;44: 469–487. doi:10.1146/annurev.phyto.44.032905.091147

34. Rodicio R, Heinisch JJ. Yeast on the milky way: genetics, physiology and biotechnology
of Kluyveromyces lactis. Yeast. 2013;30: 165–177. doi:https://doi.org/10.1002/yea.2954

35. Valentin-Hansen P, Eriksen M, Udesen C. MicroReview: The bacterial Sm-like protein
Hfq: a key player in RNA transactions. Mol Microbiol. 2004;51: 1525–1533.
doi:10.1111/j.1365-2958.2003.03935.x

36. Vogel J, Luisi BF. Hfq and its constellation of RNA. Nat Rev Microbiol. 2011;9: 578–
589. doi:10.1038/nrmicro2615

37. Dutta T, Srivastava S. Small RNA-mediated regulation in bacteria: A growing palette of
diverse mechanisms. Gene. 2018;656: 60–72. doi:10.1016/j.gene.2018.02.068

38. Papenfort K, Said N, Welsink T, Lucchini S, Hinton JCD, Vogel J. Specific and
pleiotropic patterns of mRNA regulation by ArcZ, a conserved, Hfq-dependent small
RNA. Mol Microbiol. 2009;74: 139–158. doi:10.1111/j.1365-2958.2009.06857.x

39. Argaman L, Hershberg R, Vogel J, Bejerano G, Wagner EGH, Margalit H, et al. Novel
small RNA-encoding genes in the intergenic regions of Escherichia coli. Curr Biol.
2001;11: 941–950. doi:10.1016/S0960-9822(01)00270-6

40. Neubacher N, Tobias NJ, Huber M, Cai X, Glatter T, Pidot SJ, et al. Symbiosis,
virulence and natural-product biosynthesis in entomopathogenic bacteria are regulated
by a small RNA. Nat Microbiol. 2020;5: 1481–1489. doi:10.1038/s41564-020-00797-5

41. Liang Z, Huang L, He F, Zhou X, Shi Z, Zhou J, et al. A Substrate-Activated Efflux
Pump, DesABC, Confers Zeamine Resistance to Dickeya zeae. mBio. 2019;10.
doi:10.1128/mBio.00713-19

42. Royet K, Parisot N, Rodrigue A, Gueguen E, Condemine G. Identification by Tn-seq of
Dickeya dadantii genes required for survival in chicory plants. Mol Plant Pathol.
2019;20: 287–306. doi:10.1111/mpp.12754

43. Chao Y, Li L, Girodat D, Förstner KU, Said N, Corcoran C, et al. In Vivo Cleavage Map
Illuminates the Central Role of RNase E in Coding and Non-coding RNA Pathways. Mol
Cell. 2017;65: 39–51. doi:10.1016/j.molcel.2016.11.002

44. Mandin P, Gottesman S. Integrating anaerobic/aerobic sensing and the general stress
response through the ArcZ small RNA. EMBO J. 2010;29: 3094–3107.
doi:10.1038/emboj.2010.179

45. Yuan X, Zeng Q, Khokhani D, Tian F, Severin GB, Waters CM, et al. A feed-forward
signalling circuit controls bacterial virulence through linking cyclic di-GMP and two
mechanistically distinct sRNAs, ArcZ and RsmB. Environ Microbiol. 2019;21: 2755–
2771. doi:10.1111/1462-2920.14603

46. Battesti A, Majdalani N, Gottesman S. The RpoS-mediated general stress response in
Escherichia coli. Annu Rev Microbiol. 2011;65: 189–213. doi:10.1146/annurev-micro-
090110-102946

47. Wilf NM, Salmond GPC. The stationary phase sigma factor, RpoS, regulates the
production of a carbapenem antibiotic, a bioactive prodigiosin and virulence in the
enterobacterial pathogen Serratia sp. ATCC 39006. Microbiology. 2012;158: 648–658.
doi:10.1099/mic.0.055780-0

48. Zeng Q, McNally RR, Sundin GW. Global Small RNA Chaperone Hfq and Regulatory
Small RNAs Are Important Virulence Regulators in Erwinia amylovora. J Bacteriol.
2013;195: 1706–1717. doi:10.1128/JB.02056-12

49. De Lay N, Gottesman S. A complex network of small non-coding RNAs regulate motility
in Escherichia coli. Mol Microbiol. 2012;86: 524–538. doi:10.1111/j.1365-
2958.2012.08209.x

50. Schachterle JK, Zeng Q, Sundin GW. Three Hfq-dependent small RNAs regulate
flagellar motility in the fire blight pathogen Erwinia amylovora. Mol Microbiol. 2019;111:
1476–1492. doi:10.1111/mmi.14232

51. Fraser GM, Hughes C. Swarming motility. Curr Opin Microbiol. 1999;2: 630–635.
doi:10.1016/S1369-5274(99)00033-8

52. Wésolowski-Louvel M. An efficient method to optimize Kluyveromyces lactis gene
targeting. FEMS Yeast Res. 2011;11: 509–513. doi:10.1111/j.1567-1364.2011.00741.x

53. Clamp M, Cuff J, Searle SM, Barton GJ. The Jalview Java alignment editor. Bioinforma
Oxf Engl. 2004;20: 426–427. doi:10.1093/bioinformatics/btg430

54. Edgar RC. MUSCLE: a multiple sequence alignment method with reduced time and
space complexity. BMC Bioinformatics. 2004;5: 113. doi:10.1186/1471-2105-5-113

FIGURE LEGENDS

<u>**Figure 1**</u>. **Organization and distribution of the secondary metabolite clusters *ssm*, *ooc*,**

and *zms* of *D. solani* D s0432-1.

(A) Organization of the clusters *ssm*, *ooc*, and *zms*. Genes are indicated using the NCBI

nomenclature of the NCBI reference genome sequence NZ_CP017453.1 of *D. solani* D s0432-

1. XXXXX are the digital number in BJD21_RSXXXXX, which corresponds to the locus tag.

Arrowheads show gene orientations. Color code indicates gene function. The red-framed

arrows indicate genes targeted for in-frame deletion performed in this study. (B) MLSA tree

positioning strains within the *Dickeya* genus. The evolutionary history was inferred by using

the Maximum Likelihood method. The presence (green frame) or absence (red frame) of the

cluster *ssm*, *ooc*, and *zms* are indicated in front of each strain. The distribution of these clusters

in 155 *Dickeya* strains whose genome has been sequenced is given in Table S1.

<u>**Figure 2**</u>. **Inhibition of fungal growth by *D. solani* Ds0432-1 and mutant derivatives.**

5 µL of bacteria culture at $OD_{600nm}= 2$ from *D. solani* Ds0432-1 (WT) or derivatives were

spotted onto PDA plates inoculated with plugs of *S. sclerotiorum, B. cinerea or M. oryzae*

mycelium. Plates were incubated at 25°C until the mycelium covers the plate. Lengths of fungi

inhibition zone were measured in 4 independent experiments. A statistical difference was

significant only between the WT and Δ*oocL* mutants (Mann-Whitney test; * p-value < 0.05).

<u>**Figure 3**</u>. **Inhibition of bacteria by *D. solani* Ds0432-1 and mutant derivatives.**

Bioassay plates were prepared by mixing bacterial culture (*B. subtilis, S. scabiei, E. coli, D.*

dadantii and P. atrosepticum) with warm LB agar. 5 µL of bacterial cultures at $OD_{600nm}= 2$ from

D. solani Ds0432-1 (WT) or derivatives were spotted onto the plate and incubation was

performed at 30°C during 48 h. A slight inhibition zone was observed with *B. subtilis* and *E.*

coli, except with the *Δzms* mutants. All experiments were carried in 4 replicates.

<u>**Figure 4**</u>. **Inhibition of yeasts by *D. solani* Ds0432-1 and mutant derivatives.**

Bioassay plates were prepared by mixing yeast culture (*S. cerevisiae, K. lactis, C. albicans*)

with warm YPD agar. (A) 5 μL of bacterial culture at $OD_{600nm}= 2$ from *D. solani* Ds0432-1 (WT)

or derivatives were spotted onto the plates and incubation was performed at 30°C during 48

h. (B) The radius of each inhibition zone was measured. A statistical difference was significant

only between the WT and the *Δssm* mutants (Mann-Whitney test; * p-value < 0.05). For *C.*

albicans, the inhibition zone was too small to be quantified. All experiments were carried in 4

replicates.

<u>**Figure 5**</u>. ***D. solani* WT strains deficient for anti-microbial activity are mutated for the**

sRNA *arcZ*.

(A) Inhibition of *B. subtilis*, *K. lactis* and S. *sclerotiorum* by diverse WT *D. solani* strains. The

tests were performed as previously described using the nine strains IFB 0099

(NZ_CP024711), IFB 0158 (NZ_PENA00000000), IFB 0223 (NZ_CP024710), D s0432-1

(NZ_CP017453), IPO 3494 (NZ_CM001842), IFB 0484 (NZ_CM001860), IPO 3337

(NZ_CP016928), IPO 3793 (NZ_CP017454) or the type strain IPO 2222 (NZ_CP015137).

Only IFB 0223 and IPO 2222 were unable to inhibit *B. subtilis* and *K. lactis.* (B) Organization

and conservation of the *arcZ* locus in the nine *D. solani* strains. The single mutation G90A in

IPO 2222 *arcZ* is symbolized by a dark spot in the red *arcZ* arrow. (C) Zoom on the nucleotide

sequence around *arcZ* of *D. solani* D s0432-1. This allele is named *arcZ₁*. Positions and

orientations of the genes *elbB*, *arcZ* and *arcB* are indicated. The location of the G90A mutation

present in IPO 2222 is indicated. This allele is named *arcZ₂*.

<u>**Figure 6**</u>: **Inhibition of *K. lactis* and *B. subtilis* by *D. solani* is regulated by ArcZ.**

Yeast bioassay plates were prepared by mixing culture of *K. lactis* with warm YPD agar.

Bacteria bioassay plates were prepared by mixing culture of *B. subtilis* with warm LB agar.

PDA plates were inoculated with plugs of *S. sclerotiorum.* For each experiment, 5 µL of

bacterial culture at OD_{600nm}=2 from the different *D. solani* strains were spotted onto the plate.

(A) Inhibition zones observed with Ds0432-1, IPO 2222, their respective Δ*arcZ* mutants and

IFBO223. $ArcZ_1$ is required for the synthesis of metabolites produced by the clusters *ssm* and

zms (inhibition of *K. lactis* and *B. subtilis*, respectively) but not for the metabolite produced by

the cluster *ooc* (inhibition of *S. sclerotiorum*). (B) Complementation tests and heterologous

expression of the alleles *arcZ*1 or *arcZ*2 in the three *D. solani* strains. $ArcZ_1$ activates the anti-

microbial activities of *ssm* and *zms* clusters in IPO 2222 and IFBO223. All experiments were

carried in 4 replicates.

<u>Figure 7</u>. Expression of different ArcZ alleles: the G90A mutation in $ArcZ_2$ does not

prevent its production but its processing

(A) Detection of the sRNA $ArcZ_1$ and $ArcZ_2$ by Northern Blot in *D. solani* strains. A small

transcript (~60 nt) corresponding to the processed 3′ fragment of $ArcZ_1$ was detected in D

s0432-1 and in all strains expressiong $arcZ_1$ on plasmid. In IPO 2222, only the full-length form

(~132nt) $ArcZ_2$ was detected, as well as in strains expressing the allele $arcZ_2$ on plasmid. The

5S RNA is used as a loading control. This experiment was performed twice with biological

replicates and gave identical results. (B) Prediction of secondary structure of the precursor

forms of $ArcZ_1$ and $ArcZ_2$ by RNA fold. Bases are colored according to base-pairing

probabilities (see color scale). The G90A mutation present in $ArcZ_2$ seems to modify the

structure of the ArcZ sRNA precursor.

<u>Figure S1</u>. Growth curves of *D. solani* strains and mutant derivatives.

A 96-well plate containing M63 medium with 1% sucrose was inoculated with the strains used

in this study at an OD of 0.06. The growth of each strain was determined by measuring OD_{600}

every 20 minutes during 2 days in TECAN. The different strains showed similar overall growth, except for the light grey points where a low but significant fitness gain was observed for the mutants Δ*zms,* Δ*ssm* Δ*zms,* Δ*ooc* Δ*zms* and Δ*ssm* Δ*ooc* Δ*zms* (Mann-Whitney test, p-value<0.05).

Figure S2. **Inhibition of *E. coli* growth by *D. solani* D s0432-1 and mutant derivatives (zoom on drop borders).**

Bioassay plates were prepared by mixing *E. coli* culture with melted LB agar as described in experimental procedures. 5 µL of bacterial culture at OD_{600nm} = 2 of *D. solani* D s0432-1 (WT) or derivatives were spotted onto the plate and incubated at 30°C during 48 h. A slight inhibition zone was observed except with the Δ*zms* mutant. All experiments were carried in 4 replicates.

Figure S3. **Phenotypes and *arcZ* sequence of WT *D. solani* IPO 2222 from the BCCM collection (LMG 25993).**

(A) Inhibition assay of *K. lactis, B. subtilis* and *S. sclerotiorum* by WT *D. solani* IPO 2222 LMG 25993 from BCCM collection, compared to the strains used in this study. (B) Sanger sequencing results of *D. solani* IPO 2222 *arcZ* (LMG 25993). The mutation G90A is highlighted by a red frame.

Figure S4. Growth curves of WT and Δ*arcZ D. solani* strains used in this study.

(A) Growth curves of D s0432-1, IPO 2222, their respective Δ*arcZ* mutants and IFB0223. The latter (in grey) has a slight growth delay (Mann-Whitney test, p-value<0.05). (B) Growth curves of strains used in complementation and heterologous expression tests. The plasmids do not cause any growth defect in D s0432-1, IPO 2222 and Δ*arcZ* mutants. IFB0223 strains (in grey) have growth delay already observed in the experiment without the plasmid (Mann-Whitney test, p-value<0.05). All experiments were carried in 4 biological replicates.

Figure S5. **Alignment of *arcZ* sequences.** The *arcZ* DNA sequences were retrieved by

running a BlastN on the NCBI database using the *arcZ* sequence of *D. solani* D s0432-1 as

query. Search was limited to the *Dickeya* genomes. Then, the *Dickeya arcZ* sequences were

aligned with *E. coli* MG1655 *arcZ* by using Jalview [53] and Muscle [54]. Known regions of

interactions with the *pecT*, *flhD* and *rpoS* 5'UTR mRNA in *D. dadantii* and *E. coli* are indicated

Supplementary Material and Methods. DOCX file with supplementary material and

methods.

Table S1. Distribution of the clusters *ssm*, *ooc*, and *zms* in 155 *Dickeya* strains whose

genome is sequenced. For each *Dickeya* species, the strains were classified on the basis of

the presence or absence of the cluster *ssm*, *ooc*, and *zms*.

Table S2. Lists of SNPs in *D. solani* IFB0099, D S0432.1, IFB0223, IPO3337 and IPO 2222.

Table S3. Strains and plasmids used in the study.

Table S4. Oligonucleotides used in the study.

Figure 1: Organization and distribution of the secondary metabolite clusters *ssm*, *ooc* and *zms* of *D. solani* D s0432-1.

(A) Organization of the clusters *ssm, ooc,* and *zms.* Genes are indicated using the NCBI nomenclature of the NCBI reference genome sequence NZ_CP017453.1 of *D. solani* D s0432-1. XXXXX are the digital number in BJD21_RSXXXXX, which corresponds to the locus tag. Arrowheads show gene orientations. Color code indicates gene function. The red-framed arrows indicate genes targeted for in-frame deletion performed in this study. (B) MLSA tree positioning strains within the *Dickeya* genus. The evolutionary history was inferred by using the Maximum Likelihood method. The presence (green frame) or absence (red frame) of the cluster *ssm, ooc,* and *zms* are indicated in front of each strain. The distribution of these clusters in 155 *Dickeya* strains whose genome has been sequenced is given in Table S1.

Figure 2: Inhibition of fungal growth by *D. solani* D s0432-1 and mutant derivatives.

5 µL of bacteria culture at OD_{600nm} = 2 from *D. solani* D s0432-1 (WT) or derivatives were spotted onto PDA plates inoculated with plugs of *S. sclerotiorum*, *B. cinerea* or *M. oryzae* mycelium. Plates were incubated at 25°C until the mycelium covers the plate. Lengths of fungi inhibition zone were measured in 4 independent experiments. A statistical difference was significant only between the WT and *ΔoocL* mutants (Mann-Whitney test; * p-value < 0.05).

Figure 3: Inhibition of bacteria by *D. solani* D s0432-1 and mutant derivatives.

Bioassay plates were prepared by mixing bacterial culture (*B. subtilis, S. scabiei, E. coli, D. dadantii* and *P. atrosepticum*) with warm LB agar. 5 µL of bacterial cultures at OD_{600nm} = 2 from *D. solani* D s0432-1 (WT) or derivatives were spotted onto the plate and incubated at 30°C during 48h. A slight inhibition zone was observed with *B. subtilis* and *E. coli* plates, except with the *Δzms* mutants. All experiments were carried in 4 replicates.

Figure 4: Inhibition of yeasts by *D. solani* and mutant derivatives.
Bioassay plates were prepared by mixing yeast culture (*S. cerevisiae, K. lactis, C. albicans*) with warm YPD agar. (A) 5 µL of bacterial cultures at OD_{600nm} = 2 from *D. solani* D s0432-1 (WT) or derivatives were spotted onto the plates and incubation was performed at 30°C during 48 h. (B) The radius of each inhibition zone was measured. A statistical difference was significant only between the WT and the *Δssm* mutants (Mann-Whitney test; * p-value < 0.05). For *C. albicans*, the inhibition zone was too small to be quantified. All experiments were carried in 4 replicates.

Figure 5: *D. solani* WT strains deficient for anti-microbial activity are mutated for the sRNA *arcZ*.

(A) Inhibition of *B. subtilis, K. lactis* and *S. sclerotiorum* by diverse WT *D. solani* strains. The tests were performed as previously described using the nine strains IFB0099 (NZ_CP024711), IFB 0158 (NZ_PENA00000000), IFB0223 (NZ_CP024710), D s0432-1 (NZ_CP017453), IPO 3494 (NZ_CM001842), IFB0484 (NZ_CM001860), IPO 3337 (NZ_CP016928), IPO 3793 (NZ_CP017454) or the type strain IPO 2222 (NZ_CP015137). Only IFB0223 and IPO 2222 were unable to inhibit *B. subtilis* and *K. lactis.* (B) Organization and conservation of the *arcZ* locus in the nine *D. solani* strains. The single mutation G90A in IPO 2222 *arcZ* is symbolized by a dark spot in the red *arcZ* arrow. (C) Zoom on the nucleotide sequence around *arcZ* of *D. solani* D s0432-1. This allele is named *arcZ₁*. Positions and orientations of the genes *elbB, arcZ* and *arcB* are indicated. The location of the G90A mutation present in IPO 2222 is indicated. This allele is named *arcZ₂*.

A

B

Figure 6: Inhibition of *K. lactis* and *B. subtilis* by *D. solani* is regulated by ArcZ.
Yeast bioassay plates were prepared by mixing culture of *K. lactis* with warm YPD agar. Bacteria bioassay plates were prepared by mixing culture of *B. subtilis* with warm LB agar. PDA plates were inoculated with plugs of *S. sclerotiorum*. For each experiment, 5 µL of bacteria culture at OD_{600nm} = 2 from the different *D. solani* strains were spotted onto the plate. (A) Inhibition zones observed with D s0432-1, IPO 2222, their respective Δ*arcZ* mutants and IFBO223. ArcZ$_1$ is required for the synthesis of metabolites produced by the clusters *ssm* and *zms* clusters (inhibition of *K. lactis* and *B. subtilis*, respectively) but not for the metabolite produced by the cluster *ooc* (inhibition of *S. sclerotiorum*).
(B) Complementation tests and heterologous expression of the alleles *arcZ$_1$* or *arcZ$_2$* in the three *D. solani* strains. ArcZ$_1$ activates the anti-microbial activities of *ssm* and *zms* clusters in IPO 2222 and IFBO223. All experiments were carried in 4 replicates.

Figure 7. Expression of different ArcZ alleles: the G90A mutation in ArcZ$_2$ does not prevent its production but its processing.

(A) Detection of the sRNA ArcZ$_1$ and ArcZ$_2$ by Northern Blot in *D. solani* strains. A small transcript (~60 nt) corresponding to the processed 3' fragment of ArcZ$_1$ was detected in D s0432-1 and in all strains expressing the allele *arcZ$_1$* on plasmid. In IPO 2222, only the full-length form (~132nt) ArcZ$_2$ was detected, as well as in strains expressing the allele *arcZ$_2$* on plasmid. The 5S RNA is used as a loading control. This experiment was performed twice with biological replicates and gave identical results. (B) Prediction of secondary of the precursor forms of ArcZ$_1$ and ArcZ$_2$ by RNA fold. Bases are colored according to base-pairing probabilities (see color scale). The G90A mutation present in ArcZ$_2$ seems to modify the structure of the ArcZ sRNA precursor.

Figure S1. Growth curves of *D. solani* strains and mutant derivatives.
A 96-well plate containing M63 medium with 1% sucrose was inoculated with the strains used in this study at an OD of 0.06. The growth of each strain was determined by measuring OD_{600} every 20 minutes during 2 days in TECAN. The different strains showed similar overall growth, except for the light grey points where a low but significant fitness gain was observed for the mutants Δ*zms*, Δ*ssm* Δ*zms*, Δ*ooc* Δ*zms and* Δ*ssm* Δ*ooc* Δ*zms* (Mann-Whitney test, p-value<0.05).

Figure S2. Inhibition of *E. coli* growth by *D. solani* D s0432-1 and mutant derivatives (zoom on drop borders).

Bioassay plates were prepared by mixing *E. coli* culture with melted LB agar as described in experimental procedures. 5 µL of bacterial culture at OD$_{600nm}$ = 2 of *D. solani* D s0432-1 (WT) or derivatives were spotted onto the plate and incubated at 30°C during 48 h. A slight inhibition zone was observed except with the Δ*zms* mutant. All experiments were carried in 4 replicates.

Figure S3. Phenotypes and *arcZ* sequence of WT *D. solani* IPO 2222 from the BCCM collection (LMG25993).

(A) Inhibition assay of *K. lactis*, *B. subtilis* and *S. sclerotiorum* by WT *D. solani* IPO 2222 LMG25993 from BCCM collection, compared to the strains used in this study.

(B) Sanger sequencing results of *D. solani* IPO 2222 *arcZ* (LMG25993)**.** The mutation G90A is highlighted by a red frame.

Figure S4. Growth curves of WT and Δ*arcZ D. solani* strains used in this study.
(A) Growth curves of D s0432-1, IPO 2222, their respective Δ*arcZ* mutants and IFB0223. The latter (in grey) has a slight growth delay (Mann-Whitney test, p-value<0.05). (B) Growth curves of strains used in complementation and heterologous expression tests. The plasmids do not cause any growth defect in D s0432-1, IPO 2222 and Δ*arcZ* mutants. IFB0223 strains (in grey) have growth delay already observed in the experiment without the plasmid (Mann-Whitney test, p-value<0.05). All experiments were carried in 4 biological replicates.

Figure S5. Alignment of *arcZ* sequences.

The *arcZ* DNA sequences were retrieved by running a BlastN on the NCBI database using the *arcZ* sequence of *D. solani* D s0432-1 as query. Search was limited to the *Dickeya* genomes. Then, the *Dickeya arcZ* sequences were aligned with *E. coli* MG1655 arcZ by using Jalview (56) and Muscle (57). Known regions of interactions with the *pecT*, *flhD* and *rpoS* 5'UTR mRNA in *D. dadantii* and *E. coli* are indicated