1

Dickeya solani D s0432-1 produces an arsenal of secondary metabolites with anti-prokaryotic and anti-eukaryotic activities against bacteria, yeasts, fungi, and aphids.

Running title: *Dickeya solani* secondary metabolites

Authors: Effantin G[1], Brual T[1], Rahbé Y[1,2], Hugouvieux-Cotte-Pattat N[1] & Gueguen
E[1]*.

Affiliation:
[1] Univ Lyon, Université Claude Bernard Lyon1, CNRS, INSA Lyon, UMR5240 MAP,
F-69622, LYON, France
[2] INRAE, Univ. Lyon, UMR5240 MAP, F-69622, Lyon, France

*corresponding author: erwan.gueguen@univ-lyon1.fr

Orcid I.D.
G.E.   0000-0001-8784-1491
H-C-P.N   0000-0002-4322-160X
R. Y.   0000-0002-0074-4443

Key words: *Dickeya solani*, deletion mutagenesis, secondary metabolites, oocydin,
zeamine, inhibition, bacteria, yeasts, fungi, streptomyces, aphid

SUMMARY

The necrotrophic plant pathogenic bacterium *Dickeya solani* is a new invader of potato agrosystem in Europe. All isolated strains of *D. solani* contain several large polyketide/fatty acid/non-ribosomal peptide synthetase clusters. Analogy with genes described in other bacteria, suggests that two clusters are involved in the production of secondary metabolites of the oocydin and zeamine family. In this study, we constructed by an approach of reverse genetics mutants affected in the three secondary metabolite clusters *ssm*, *ooc* and *zms* in order to compare the phenotype of the *D. solani* strain D s0432-1 with its derived mutants. We demonstrated that the zeamine cluster *zms* inhibits growth of gram-positive and gram-negative bacteria. It is also implicated in a toxicity against aphids. The oocydin cluster *ooc* inhibits growth of fungi of the phylum *Ascomycota*. Finally, we unveiled the function of a new secondary metabolite cluster *ssm* (for *solani* secondary metabolite), only conserved in some *Dickeya* species. This cluster produces a secondary metabolite inhibiting yeasts. *D. solani* therefore produces several molecules that are toxic to a wide range of living and potentially interacting organisms, from bacteria to insects. The expression of these secondary metabolite pathways could contribute to the rapid spread of *D. solani* in Europe.

INTRODUCTION

Bacterial phytopathogens of the genus *Dickeya* and *Pectobacterium* are pectinolytic necrotrophic bacteria with a broad host plant spectrum (Charkowski *et al.*, 2012; Hugouvieux-Cotte-Pattat *et al.*, 2020). These members of the family *Pectobacteriaceae* (Van Gijsegem, Toth, *et al.*, 2021) cause substantial agricultural losses worldwide by affecting many vegetables, ornamentals and crops, of which the potato is the most important economically. These bacteria are able to invade and degrade the plant tissues through the coordinated expression of genes encoding virulence factors, with a major role of pectate lyases that dissociate the plant cell wall constituents (Hugouvieux-Cotte-Pattat *et al.*, 2014).

The *Dickeya* genus was established in 2005 (Samson *et al.*, 2005), resulting from the reclassification of *Pectobacterium chrysanthemi* (formerly *Erwinia chrysanthemi*). To date, twelve species of *Dickeya* have been described, *D. aquatica*, *D. chrysanthemi*, *D. dadantii D. dianthicola* , *D. fangzhongdai*, *D. lacustris*, *D. oryzae*, *D. paradisiaca*, *D. poaceiphila*, *D. solani*, *D. undicola*, and *D. zeae* (Samson *et al.*, 2005; Brady *et al.*, 2012; Parkinson *et al.*, 2014; van der Wolf *et al.*, 2014; Tian *et al.*, 2016; Hugouvieux-Cotte-Pattat *et al.*, 2019, 2020; Oulghazi *et al.*, 2019; Wang *et al.*, 2020).

The species *D. solani* was officially established in 2014 (van der Wolf *et al.*, 2014) but *D. solani* isolates have attracted attention since its emergence on the potato agrosystem in Europe in

the early 2000s. It appeared to be highly aggressive in both subtropical and temperate
climates. Recently, seed companies have adopted a zero-tolerance policy to *D. solani* due to
its invasive and aggressive nature (Van Gijsegem, van der Wolf, *et al.*, 2021). Many scientific
efforts have been made to provide information on this phytopathogen, resulting in 76 *D. solani*
genomes available in May 2021 (Blin *et al.*, 2021). Comparative genomics was performed to
identify the genetic basis for the high virulence level of *D. solani* (Garlant *et al.*, 2013; Pédron
et al., 2014; Khayi *et al.*, 2015; Golanowska *et al.*, 2018; Motyka-Pomagruk *et al.*, 2020; Blin
et al., 2021). Most *D. solani* strains Isolated from different regions show a low level of genetic
variation, suggesting a clonal origin (Khayi *et al.*, 2015). The genomes of *D. solani* share a
high similarity and synteny with those of the model strain *D. dadantii* 3937, prompting
comparison between the two species. Only a few hundred genes were specific to each
species, including a few dozen distinctive genomic regions (Garlant *et al.*, 2013; Pédron *et al.*,
2014). Three of these regions have been shown to encode polyketide synthases (PKS), non-
ribosomal peptide synthetases (NRPS) and amino acid adenylation domain proteins,
suggesting that they encode proteins involved in the production of secondary metabolites
(Garlant *et al.*, 2013).

PKSs and NRPSs are able to synthesize molecules by sequential condensation of carboxylic
acids and amino acids, respectively. PKS and NRPS modules can combine together to form
hybrid PKS/NRPS systems capable of producing compounds of great structural diversity
(Cane and Walsh, 1999). The molecules synthesized may have siderophore, antibiotic or
phytotoxic properties that promote the virulence of a plant pathogen. Three PKS/NRPS
clusters are present in all sequenced *D. solani* strains and found in a few other *Dickeya*
species (Duprey *et al.*, 2019). In *Serratia plymuthica,* the cluster *ooc* is involved in the
synthesis of oocydin A, a halogenated macrolide with antifungal and anti-oomycete activity
(Matilla *et al.*, 2012). The cluster *zms*, previously found in the genomes of *S. plymuthica* and
Dickeya oryzae, leads to the biosynthesis of a polyamino-amide antibiotic, zeamine (Zhou *et*
al., 2011; Masschelein *et al.*, 2013). A third cluster is found in a few *Dickeya* species but the
nature and function of the molecule synthesized are unknown.

Mutagenesis is one of the most powerful genetic tools for analyzing the function of a gene and
its involvement in bacterial virulence. A random mutagenesis approach was used in the *D.*
solani type strain IPO2222 with a transposon Tn5 harboring a promoterless *gusA* reporter
gene to obtain mutants affected under given conditions (Fikowicz-Krosko and Czajkowski,
2017; Lisicka *et al.*, 2018; Czajkowski *et al.*, 2020). Some mutants affected for a selected gene
have also been constructed in various strains of *D. solani* using a generalized transduction
method with phage, to transfer a mutation previously constructed in *D. dadantii* 3937 (Potrykus
et al., 2014, 2018). However, this method is strictly limited to genes common to *D. dadantii*

and *D. solani*. Given the need for in-depth studies on this economically important species, we

have tested a general method to precisely inactivate a selected gene using the SacB-based

reverse genetics method. We used this method to construct mutants affected for the three loci

involved in the biosynthesis of secondary metabolites in the highly virulent *D. solani* strain

Ds0432-1. This method is potentially applicable to all genes and in all strains of *D. solani*. We

showed that *D. solani* Ds0432-1 clusters *zms* and *ooc,* encoding zeamine and oocydin

biosynthesis, are involved in growth inhibition of bacteria and fungi, respectively. We also

showed evidence that the *D. solani* Ds0432-1 cluster *ssm* (for <u>s</u>olani <u>s</u>econdary <u>m</u>etabolite)

produces a third secondary metabolite that prevents yeast growth.

RESULTS

Use of the SacB-based reverse genetics method to delete genes of *D. solani*

A technique commonly used in bacterial genetics to delete a gene from a Gram-negative

bacterium requires a suicide plasmid with the R6K origin of replication and the counter-

selection gene *sacB1*, which confers sucrose sensitivity (Edwards *et al.*, 1998). The vector

pRE112 had previously been successfully used for allelic exchange in *D. dadantii* 3937

(Koskiniemi *et al.*, 2013; Royet *et al.*, 2019). We tried to use it with *D. solani*.

To evaluate the effectiveness of the reverse genetic approach with *D. solani*, we first tried to

delete the *araBA* operon of the *D. solani* strain Ds0432-1. *araA* encodes the L-arabinose

isomerase and *araB* a ribulokinase. A Δ*araBA* mutant, deleted for this operon, is unable to

catabolize arabinose and it forms white colonies on MacConkey-arabinose agar plates. This

mutant is thus easily distinguishable from the wild-type Ara$^+$ strain forming red colonies. Two

1-kb fragments upstream and downstream *araBA* were assembled and cloned directly into

linearized pRE112 vector (Fig. 1A). The plasmid pRE112-Δ*araBA* was transferred from *E. coli*

MFDpir to *D. solani* Ds0432-1 by conjugation (Fig. 1B). Plasmid integrants were selected by

their chloramphenicol resistance (Cm^R) on LB agar plates supplemented with

chloramphenicol. On this selective medium, the *E. coli* donor strain cannot survive because

LB agar does not contain diaminopimelic acid (DAP). The plasmid is integrated either

upstream or downstream of the *araBA* locus (Fig. 1A). The integrated plasmid can be lost by

homologous recombination between the tandemly duplicated *ara* sequences (Fig. 1A).

Plasmid loss is stimulated in the presence of sucrose, *i.e.* by spreading the integrants onto LB

agar without NaCl containing 5% sucrose. The sucrose-resistant colonies were then replica-

plated onto MacConkey-arabinose agar, allowing the identification of Ara$^-$ segregants, and on

LB agar-Cm, to confirm the plasmid loss by their Cm^S phenotype (Fig. 1B). Finally, the correct

structure of the mutants was confirmed by colony PCR analysis (Fig. 1C). By this first
experiment, we were able to verify that this method works perfectly and is easy to implement
in *D. solani*. In addition, this method was shown to work by cloning only 500 bp DNA fragments
upstream and downstream of *araBA*, instead of 1000 bp. We also used this protocol to
construct a Δ*araBA* mutant of the *D. solani* type strain IPO2222 with comparable efficiency
(data not shown).

Figure 1

Figure 1. Summary of the method used to obtain in-frame deletion mutants. (A) Genetic
steps to obtain a ∆*araAB* deletion mutant of *Dickeya solani*. The 1 kb upstream and
downstream DNA regions of the *araBA* genes are cloned between the KpnI and SacI
restriction sites of the suicide plasmid pRE112 by Gibson assembly. The resulting plasmid
pRE112-∆*araBA* is then transferred into *D. solani* by conjugation. The plasmid is integrated
into the chromosome by homologous recombination either upstream or downstream the *araBA*
locus. The structures of both possible integrants are shown as structures 1 and 2. Next,
sucrose-resistant segregants lose the integrated plasmid pRE112-∆araBA by homologous
recombination between the tandemly duplicated upstream or downstream *ara* sequences.
This either regenerates a WT *araBA* locus or an *araBA* deletion on the chromosome, shown
as structure 3.
(B) Microbiological steps to obtain a ∆*araBA D. solani* mutant. The *E. coli* donor strain
MFDpir(pRE112-∆*araBA*) was mated with *D. solani* (step I). *D. solani* recombinants were
selected onto LB agar medium supplemented with chloramphenicol (step II). *D. solani*
recombinant were then spread onto LB agar without NaCl and supplemented with 5% sucrose.
Sucrose resistant colonies (step III) were transferred onto MacConkey agar plate with 1 %
arabinose (step IV). White colonies are formed by bacteria unable to catabolize arabinose
because they acquired the ∆*araBA* mutation. (C) Their genotype was confirmed by PCR on
colonies using oligo pairs L1127/L1128. The desired mutants give a PCR product of 2 kb,
whereas PCR done on WT *D. solani* gives a PCR product > 5 kb. Detail procedure is indicated
in the material and methods section.
165

Using this reverse genetic approach, we decided to study the three secondary metabolite

clusters of *D. solani* which are absent in *D. dadantii*. We focused our work on *D. solani* D

s0432-1 since it is one of the most aggressive *D. solani* strain (Khayi *et al.*, 2015; Golanowska

et al., 2018). This strategy should allow us to assess the contribution of these three clusters

on the competitiveness of *D. solani* against eukaryotic and prokaryotic organisms.

Description of the three selected PKS/NRPS secondary metabolite clusters of *D. solani*.

The clusters A, B and C encoding complex NRPS and PKS involved in the production of

secondary metabolites were named *ssm*, *ooc* and *zms,* respectively (Fig.2).

The ~42-kbp cluster A contains the 12 genes *ssmABCDEFGHIJKL* (Fig.2). It is widely

conserved in the genus *Dickeya*, *i.e.*, in all sequenced *D. solani, D. aquatica, D. fangzhondai,*

D. poaceiphila and *D. zeae* genomes, in some *D. dadantii* strains (NCPPB 898, NCPPB 3537,

but not the model strain 3937), in some *D. undicola* strains (FVG1, FVG10), and in some *D.*

oryzae strains (EC2, NCPPB 3531 and CSLRW192). The role and the structure of the

metabolite produced from this cluster have not been elucidated yet.

The ~80-kbp cluster B is highly similar to the *oocBCDEFGJKLMNOPQRSTUVW* cluster of *S.*

plymuthica A153. It is present in all the sequenced genomes of *D. solani* and *D. dianthicola*,

in four *D. oryzae* strains (ZYY5, EC1, DZ2Q and ZJU1202), and in two *D. paradisiaca* strains,

Ech703 and NCPPB2511. In *S. plymuthica* A153, disruption of this gene cluster abolished

bioactivity against the fungi *Verticillium dahliae* and the oomycetes *Pythium ultimum*. This

cluster produces oocydin A (Matilla *et al.*, 2012), a chlorinated macrolide, powerfully active
against plant pathogenic oomycetes (Strobel *et al.*, 1999). Since various *D. solani* strains
inhibit *V. dahliae* and *P. ultimum* growth (Matilla *et al.*, 2015), it was suggested, on the basis
of gene sequence homologies and similar cluster organization, that *D. solani* also produces
oocydin A.
The ~55-kbp cluster C encodes mixed fatty acid synthase (FAS)/PKS and hybrid NRPS/PKS
enzymes. Its genomic organization is identical to the *D. oryzae* EC1 zeamine cluster and
related to the *S. plymuthica* AS12 zeamine cluster (Zhou *et al.*, 2011, 2015). The *zms* cluster
is conserved in all the sequenced *D. solani* and *D. fangzhongdai* genomes. After
reclassification of several *D.zeae* strains in the novel species *D. oryzae* (Wang *et al.*, 2020),
this cluster appeared to be absent in the genomes of true *D. zeae* strains. It is present in some,
but not all, *D. oryzae* rice strains (ZYY5, EC1, DZ2Q and ZJU1202). The zeamine biosynthetic
clusters from *D. oryzae* EC1 and *D. solani* Ds0432-1 share from 59 to 94% identity at individual
protein level (Zhou *et al.*, 2015).
202

Figure 2

Figure 2. **Organization of the secondary metabolite clusters of *D. solani* D s0432-1.**
Genes are indicated using the NCBI nomenclature of the NCBI reference genome sequence
NZ_CP017453.1 of *D. solani* D s0432-1. XXXXX are the digital number in BJD21_RSXXXXX,
which corresponds to the locus tag. Arrowheads show gene orientations. Color code indicates
gene function. The red-framed arrows indicate genes targeted for in-frame deletion performed
in this study. Cluster function is noted on the right.

While the *zmsABCDEFGIJKLMNPQRS* cluster directing zeamine biosynthesis is present only
in some *D. oryzae* strains isolated from rice, it is conserved in all *D. fangzhongdai* and *D.*
solani strains, which suggest secondary acquisition by horizontal gene transfer in *D. oryzae*
(Zhou *et al.*, 2015; Duprey *et al.*, 2019). Zeamine-related antibiotics are polyamino-amide
molecules toxic to a wide range of pro- and eukaryotic organisms such as bacteria, fungi,
oomycetes, plants, and nematodes (Masschelein *et al.*, 2017). Mutation of the zeamine
synthase gene *zmsA* in *D. oryzae* EC1 attenuates the inhibition of rice seed germination (Zhou
et al., 2011) and suppresses antibacterial activity against *E. coli* (Zhou *et al.*, 2011). Zeamine
produced by *S. plymuthica* kills nematodes and yeast (Hellberg *et al.*, 2015). *D. solani*
IPO2222 can also kill *C. elegans* but not as quickly as *S. plymuthica* (Hellberg *et al.*, 2015).
To interrupt the synthesis of the secondary molecules produced by these three clusters, we
constructed in-frame deletion mutants inactivating a key gene of each cluster. The mutants
ΔssmG (cluster A), ΔoocL (oocydin cluster B) and ΔzmnA (zeamine cluster C) were
constructed using the described reverse genetic approach with the pRE112 suicide plasmid.
Inhibitory effects of these mutants against fungi, bacteria, yeasts, aphids and plants were
compared with that of the wild-type strain.
228
229 **The *D. solani* oocydin cluster inhibits *Ascomycota* growth**
230
We compared the ability of the *D. solani* Ds0432-1 DS58 strain (a Nal[R] Gm[R] derivative of the
WT strain) and its derived mutants to inhibit the growth of *Botrytis cinerea*, *Magnaporthe*
oryzae and *Sclerotinia sclerotiorum*, three fungi-like eukaryotes of the phylum *Ascomycota*. A
potato dextrose agar (PDA) plate was inoculated at the periphery of the Petri dish with fungal
mycelium and 10 μl of overnight bacterial culture of each *D. solani* strain was deposited at the
center of the plate (Fig. 3). After incubation at 25°C for several days, we observed a growth
inhibition of the three fungi by the *D. solani* WT and the ΔssmG and ΔzmsA mutants. In
contrast, the ΔoocL mutant did not inhibit mycelium growth (Fig. 3). Thus, we conclude that
the gene *oocL* is involved in the production of an oocydin-like secondary metabolite that has
an anti-fungal activity.

Figure 3. Radial diffusion assays to assess fungal growth inhibition on PDA plates with *D. solani* D s0432-1 or derived mutants. 5 µl of *D. solani* D s0432-1 at OD_{600nm} of 0.1 was deposited in the center of a 90 mm Petri dish inoculated with plugs of *Sclerotinia sclerotiorum*, *Botrytis cinerea* or *Magnaporthe oryzae* mycelium. Plates were incubated at 25 °C until the mycelium covers the plate. Lengths of fungi inhibition zone were measured in 3 independent experiments. A statistical difference was significant only between the WT and Δ*oocL* mutant (Mann-Whitney test; * P value < 0.05).

The *D. solani* zeamine cluster inhibits bacterial growth.

The zeamine cluster of *D. oryzae* EC1 is responsible of the bactericidal activity against *E. coli* DH5α (Zhou *et al.*, 2011). Because *D. oryzae* EC1 and *D. solani* zeamine biosynthetic genes share a high degree of similarity (Zhou *et al.*, 2015), we evaluated the capacity of *D. solani* WT DS58 and mutants to inhibit the growth of gram-positive and gram-negative bacteria. It was previously shown that D. *oryzae* EC1 produced 128 µg.mL^{-1} of zeamine in the minimum medium LS4+yeast extract, while zeamine production was not detected in LB (Liao *et al.*, 2014). Based on this work, we first used LS4+yeast extract medium for production of zeamine

by *D. solani*. The *D. solani* strains were grown in LS4+yeast extract minimal medium before addition of 5 μl of the overnight culture onto a LB agar plate pre-inoculated with *B. subtilis*. Only a very slight zone of *B. subtilis* inhibition was observed around the drop of *D. solani* in our bioassay (Fig. S1). We then tested *D. solani* growth in minimal medium M63 supplemented with sucrose and yeast extract. A large diameter of inhibition was observed, allowing differentiation between strains capable or not of inhibiting *B. subtilis* growth (Fig. S1). Thus, we used this medium for overnight growth of *D. solani* in all the following experiments. The *D. solani* WT strain DS58 and the three mutants were tested (Fig. 4A). Only the Δ*zmnA* mutant was unable to inhibit *B. subtilis* growth, indicating that *D. solani* produces an active zeamine antibiotic (Fig 4A). We also tested the ability of *D. solani* to inhibit growth of other bacteria (Fig 4A) such as *E. coli* DH5α, *Streptomyces scabiei* CFBP 4517, *D. dadantii* 3937 and *P. atrosepticum* SCRI 1043. The *D. solani* zeamine cluster was also the sole responsible for the growth inhibition of *S. scabiei*, a plant pathogen causing the potato disease common scab (Loria *et al.*, 2006) (Fig 4A). The *zms* cluster also inhibits the growth of *E. coli*, but with less efficiency than that observed for the two Gram[+] bacteria tested. No inhibition of the two other pectinolytic bacteria *D. dadantii* and *P. atrosepticum* growth was observed. In conclusion, out of the three clusters analyzed, only the zeamine-like molecule has an anti-bacterial activity.

The *D. solani* cluster *sms* but not the zeamine cluster inhibits yeast growth.

Since zeamine produced by *S. plymuthica* A153 has previously been shown to be toxic to the ascomycete yeast *Saccharomyces cerevisiae* (Hellberg *et al.*, 2015), we tested the capacity of *D. solani* D s0432-1 and its mutants to inhibit the growth of the yeasts *S. cerevisiae* and *Kluyveromyces lactis*. *K. lactis* has been isolated from milk and constitutes the predominant eukaryote during cheese productions (Rodicio and Heinisch, 2013)*. S. cerevisiae* and *K. lactis* were grown in the rich medium YPD. We observed the inhibition of *S. cerevisiae* and *K. lactis* growths by the *D. solani* WT strain DS58 (Fig. 4B), grown either in M63+sucrose+yeast extract or in YPD medium (Fig S1). The anti-eukaryotic activity against these yeasts was caused by the cluster *sms* since only the Δ*smsG* mutant did not inhibit yeast growth (Fig 4B). Therefore, it appears that zeamine is not the primary factor responsible for yeast inhibition in *D. solani* D s0432-1. Conversely, we identified a novel secondary metabolite cluster that has never been studied previously and produces a molecule with anti-eukaryotic activity.

Figure 4

A

B

Figure 4. Plate-based bioassays for metabolite production by *D. solani* and mutant derivatives. Bioassay plates were prepared by mixing different bacteria culture (*B. subtilis, S. scabiei, E. coli, D. dadantii* or *P. atrosepticum*) with warmed LB agar. (A) 5 µl of overnight bacteria culture from *D. solani* D s0432-1 or derivative mutants were spotted onto the bioassay plate. (B) Bioassay with the yeasts *Saccharomyces cerevisiae* and *Kluyveromyces lactis* in YPD agar plates. The experiment was repeated at least three times. The photograph was taken 24 h after incubation at 30°C.

The *D. solani* zeamine cluster contributes to the killing of aphid.

Since *D. dadantii* 3937 was previously shown to be able to kill aphids (Grenier *et al.*, 2006), we tested the effect of *D. solani* D s0432-1 on these insects. Aphid survival was heavily affected by infection with *D.solani*, similarly to what was observed with *D.dadantii*. Among the three mutants analyzed, the mutant ΔzmsA was clearly altered in its virulence towards the pea aphid, with a significant survival outcome over the WT strain (p=0.0008, Log-rank test) (Fig. 5). Mean lethal time was increased from 1.68 days ± [1.41-1.99] (mean ± *conf. interv.*) to 2.66 days ± [2.16-3.28]. The mutant ΔoocL seemed slightly affected for aphid survival, (Fig. 5), but the results were not significant with the non-parametric tests used throughout (p=0.23), only showing significance for a global Weibull model (p=0.027), with a mean survival time increased to 1.90 days ± [1.50-2.41]. The mutant ΔssmG was not affected in its capacity to kill aphids. Globally however, and as this occurred for the *cyt* proteins toxin cluster in *D. dadantii* 3937 (Costechareyre *et al.*, 2010), the capacity of both mutants ΔzmsA and ΔoocL to kill aphids remained high as all aphids were dead at the end of the 7 days period. Thus, the analyzed clusters only supported partial causality for the aphid-killing phenotype harbored by *D. solani* D s0432-1. In conclusion, these results showed that the zeamine cluster is partially implicated in aphid killing by *D. solani* D s0432-1.

Figure 5

Figure 5. Bioactivity assay on pea aphids. Survival curves of pea aphid individuals (n=30) after 24h exposure to *D. solani* WT and mutants. Control survival with no infection, in green (top). Curves fitted with Weibull distributions (dashed lines), showing ΔzmsA mutant with significantly altered infection capacity (see text for statistical data).

Plant cell wall maceration capacity of the *D. solani* mutants affected for secondary metabolite production .

In *D. oryzae* EC1, the mutation of *zmsA* attenuates the inhibitory activity observed on rice seed germination (Zhou *et al.*, 2011). Thus, zeamine was suggested to be a phytotoxin that could affect virulence against plants.

We examined the role of clusters *ssm*, *ooc*, and *zms* on the ability of *D. solani* to infect chicory leaves. The *D. solani* WT strain and its derived mutants were inoculated on a small wound made on a chicory leaf. After overnight incubation, no difference in maceration was observed (Fig. 6), suggesting that the wild-type strain of *D. solani* does not require any of the three PKS/NRPS secondary metabolites to efficiently infect plants.

Figure 6

A

B

Figure 6. Maceration of chicory leaves by *D. solani* and mutant derivatives. (A) The photograph was taken 24 h after incubation at 30°C. **(B)** The length of rotten tissue was measured from 9 infected leaves. No statistical difference was measured between the different strains (Mann–Whitney U-test).

DISCUSSION

Secondary metabolite pathways are a great source of molecules with anti-eukaryotic or anti-

prokaryotic activity giving the bacteria that synthesize them a competitive advantage over

other organisms. The development of targeted mutagenesis of the *D. solani* chromosome has

allowed us to specifically study the involvement of three secondary metabolite synthesis

clusters encoded by all the *D. solani* strains for which genome sequence is available. We

focused our work on D s0432-1, one of the most virulent *D. solani* strains (Golanowska *et al.*,

2018). We have demonstrated that this *D. solani* strain is able to inhibit the growth of a variety

of living organisms. We tested bio-activities against Gram-negative bacteria (*E. coli*, *P.*

atrosepticum, *D. dadantii*), Gram-positive bacteria (*B. subtilis*, *S. scabiei*), yeasts (*S.*

cerevisiae, *K. lactis*), fungi (*B. cinerea*, *M. oryzae* and *S. sclerotiorum*), and pea aphids (*A.*

pisum). The targeted mutagenesis approach revealed that each of the three secondary

metabolite synthesis pathways studied has the capacity of inhibiting a particular type of

organism. Cluster B , or *ooc*, encodes an enzymatic pathway that produces an oocydin-like

molecule, a chlorinated macrocyclic lactone molecule having antifungal, anti-oomycete and

antitumor activities in *S. plymuthica* (Matilla *et al.*, 2012, 2015). In contact with fungi *B. cinerea*,

M. oryzae and *S. sclerotiorum*, *D. solani* D s0432-1 prevents their growth. It was previously

shown that the *D. solani* strains MK10, MK16, IPO 2222, 3337, D-s0432-1 and GBBC 2040,

all encoding the oocydin cluster, inhibit the growth of the fungus *V. dahliae* and the oomycete

P. ultimum (Matilla *et al.*, 2015). Our study showed that inactivation of the *oocL* gene of this

cluster in *D. solani* D-s0432-1 suppressed the inhibition of the three fungi but it had no

apparent effect on the other organisms tested.

Our data also demonstrated that *D. solani* D s0432-1 has an anti-bacterial activity linked to

cluster C, or *zsm*, which encodes a zeamine biosynthetic pathway. Zeamine produced by *S.*

plymuthica kills *B. subtilis,* the yeasts *S. cerevisiae* and *Schizosaccharomyces pombe,* and

the nematode *C. elegans* (Hellberg *et al.*, 2015). In our study, we showed that *D. solani* D

s0432-1, but not the *zmsA* mutant, can clearly inhibit the growth of two Gram-positive bacteria,

B. subtilis and *S. scabies*. We tested *S. scabiei* because it is the principal causal agent of the

common scab disease of potato (Loria *et al.*, 2006). Since *D. solani*, is also a potato pathogen,

D. solani and *S. scabiei* might be in competition for the same ecological niche. It is thus

interesting to note that *D. solani* can inhibit the growth of another potato pathogen. We also

observed a slight inhibition exerted by *D. solani* D s0432-1 against *E. coli*, but no inhibition

towards *D. dadantii* and *P. atrosepticum*. Zeamine resistance of *D. dadantii* could be explained

by the presence in its genome of the genes *desAB* encoding the RDN pump DesAB involved

in zeamine efflux (Liang *et al.*, 2019). In-frame deletion of *desA* or *desB* in *D. oryzae* EC1 leads to a zeamine sensitive phenotype (Liang *et al.*, 2019). In *in planta* Tn-seq experiments with *D. dadantii* 3937 (Royet *et al.*, 2019), mutants of the genes *desA* and *desB* (Dda3937_00787 and Dda3937_00786 respectively) did not display a significant negative or positive variation (Log2 fold-changes -0.09 and +0.44, respectively), suggesting that this RND efflux pump does not play a significant role during *D. dadantii* plant infection, contrary to the *D. dadanti* RND efflux pump AcrAB that appeared to be essential for virulence (Royet *et al.*, 2019). *P. atrosepticum* does not have the genes *desAB* but it is not inhibited by *D. solani* D s0432-1. Another RND efflux pump could allow zeamine efflux, or zeamine resistance could be provided by a different mechanism.

We also showed that zeamine is involved in insect killing. We tested pea aphids because they are susceptible to infection by *D. dadantii* (Grenier *et al.*, 2006). The *D. solani* strain D s0432-1 is also able to kill pea aphids and the zeamine synthesis pathway (absent in *D. dadantii*) is partially implicated in pea aphid killing. We did not test inhibition of other multicellular organisms. Hellberg *et al.* tested nematodes with *S. plymuthica* A153 and a few *D. solani* strains (Hellberg *et al.*, 2015), they showed that *S. plymuthica* A153 but not its zeamine-deficient mutants can kill *C. elegans* in a few hours. In contrast, *D. solani* strains MK10, MK16, and IPO2222 that also carry the zeamine cluster behaves like the zeamine-deficient mutants of *S. plymuthica*. The authors suggested the *D. solani* zeamine cluster might be cryptic under their conditions or that it produces a zeamine molecule with somewhat different biological properties (Hellberg *et al.*, 2015).

Zeamine produced by *S. plymuthica* A153 is bioactive against *S. cerevisiae* and *S. pombe* (Hellberg *et al.*, 2015). We observed that *D. solani* D s0432-1 is bioactive against *S. cerevisiae* and *K. lactis*. However, the zeamine-deficient mutant of *D. solani* behaves exactly like the WT strain, ruling out the involvement of zeamine in yeast inhibition. By testing other mutants of the secondary metabolites clusters, we observed that the mutant s*smG* was totally unable to prevent the growth of *S. cerevisiae* and *K. lactis*. Therefore, the *ssm* cluster appears to be responsible for yeast inhibition. This cluster was not required for bioactivities against bacteria, aphids, and fungi. This result opens the way to the discovery of a new molecule with very specific activity against yeasts. The *ssm* cluster is only conserved in a few *Dickeya* genomes but not in other bacterial genera. The structure of the molecule and its target remains to be elucidated.

Finally, we assessed the virulence of the *D. solani* mutants Δ*ssmG*, Δ*oocL* and Δ*zsmA* by testing the maceration of chicory leaves. No difference was observed between the WT strain D s0432-1 DS58 and the mutants affected for the production of a secondary metabolite, indicating that the three secondary metabolites tested play no significant role in the degradation of the plant cell wall. Zeamine was described as a major virulence determinant of

D. oryzae since it is involved in inhibition of rice seed germination (Zhou *et al.*, 2011). Zeamine

could have a phytotoxic activity mostly efficient at the level of seed germination.

In conclusion, *D. solani* D s0432-1 produces an arsenal of bioactive secondary metabolites

against a variety of living organisms. The cluster *ssm* produces an unkown molecule active

against yeasts; the cluster *ooc* produces an oocidin-like molecule active against fungi and the

cluster *zms* produces a zeamine-like molecule active against bacteria and insects. The

complementary activities of this set of molecules may have favored the rapid spread of *D.*

solani in Europe by giving the capacity to compete with other microorganisms.

EXPERIMENTAL PROCEDURES

Bacterial and fungal strains, plasmids and growth conditions.

The *E. coli* and *Dickeya* bacterial strains, plasmids and oligonucleotides used in this study are

described in Table S1 and S2. The genome accession number of *D. solani* D S0432-1 is

NZ_CP017453. The following strains have been also used in the study: *Sclerotinia*

sclerotiorum S5, *Botitris cinerea* B05.10, *Magnaporthe oryzae* strain Guy11, *Saccharomyces*

cerevisiae BY4743 (*MATa/α his3Δ1/his3Δ1 leu2Δ0/leu2Δ0 LYS2/lys2Δ0 met15Δ0/MET15*

ura3Δ0/ura3Δ0), *Kluyveromyces lactis* MWL9S1 (Wésolowski-Louvel, 2011), *Dickeya dadantii*

3937, *Pectobacterium atrosepticum* SCRI1043*, Streptomyces scabiei* CFBP4517, *Bacillus*

subtilis PY79. *E. coli* was grown routinely at 37°C in LB. Fungus strains were grown at 25°C

onto Potato Dextrose Agar (PDA). *S. scabiei* was cultivated in tryptic soy broth (TSB) medium

at 28°C. *B. subtilis*, *P. atrosepticum* and the *Dickeya* strains were cultivated in LB unless

specified. Yeast cells were grown at 30°C in rich medium consisting of complete yeast extract-

peptone (YP) medium containing 1% Bacto yeast extract, 1% Bacto peptone (Difco)

supplemented with 2% glucose (yeast extract-peptone-dextrose [YPD] medium). For the

bacteria and yeast inhibition assay, M63 medium supplemented with sucrose and yeast

extract (2 g $(NH_4)_2SO_4$, 13.6 g KH_2PO_4, 2.5 mg $FeSO_4 7H_2O$, 0.2 g $MgSO_4 7H_2O$, 10 g sucrose,

0.1 g yeast extract, per liter) or LS4 medium supplemented with yeast extract (9.25 g K_2HPO_4,

3.3 g KH_2PO_4, 1.4 g NH_4NO_3, 12.7 g sucrose, 1 g KCl, 0.1 g yeast extract and 0.25 g $MgSO_4$,

pH 7.0, per liter) (Liao *et al.*, 2014) were employed to grow the *D. solani* strains.

When required, antibiotics were added at the following concentrations: ampicillin (Amp), 100

µg/L; kanamycin (Kan), 50 µg/mL; nalidixic acid (Nal), 10 µg/mL; chloramphenicol (Cm), 20

µg/mL or 4 µg/mL for *E. coli* or *D. solani,* respectively. Diaminopimelic acid (DAP) (57 µg/mL)

was added for the growth of the *E. coli* MFDpir strain. Media were solidified with 12 g/L agar.

PCR conditions

To amplify DNA for cloning, a bacterial suspension of the *D. solani* was prepared by boiling
an isolated colony resuspended into 40 µl of sterilized water for 10 minutes at 95°C. The PCR
was carried out in a 50 µl reaction mix containing 1 µl of bacterial suspension, 1.5 µl of 10 µM
of each primer (Eurofins), 25 µL of primestar master mix 2x (Takara) and demineralized water.
Thermocycling for amplification of a 1 kb upstream and downstream fragment of operon *araBA*
consisted of 34 cycles of 98°C for 15 seconds, 55°C for 15 seconds and 72°C for 15 seconds.
To check the correct deletion of a gene of interest in *D. solani*, a bacterial suspension was
made by boiling an isolated colony resuspended into 40 µl of sterilized water for 10 minutes
at 95°C. The suspension was centifuged 1min at 12000 g to remove cellular debris. The
supernatant was used as a template for PCR. To amplify up to 3-kb DNA fragments, colony
PCR was performed with DreamTaq Green PCR Master mix (2X) (Thermofisher, Waltham,
MA, USA). However, the amplification from colonies with the dreamtaq kit did not work when
the amplified fragments was larger than 3-kb. In this case the colony PCR was carried out with
the Q5 High-Fidelity DNA Polymerase (Biolabs) by following manufacturer recommendations.
469
470 **Construction of the Δ*araBA D. solani* mutants.**
471
To construct the in-frame deletion Δ*araBA* of *D. solani*, the *sacB* counter-selection method
was used (Edwards *et al.*, 1998). pRE112 is an R6K-based suicide plasmid carrying the *sacB*
gene and the cat gene (Cm^R). The plasmid pRE112-Δ*araBA* was constructed by cloning
simultaneously two PCR fragments corresponding to the upstream and downstream 1-kbp
DNA of the *araBA* genes into SacI/KpnI digested pRE112 using the Gibson's assembly
method. Chemical ultracompetent DH5α λpir cells were prepared with the Mix & Go! *E.coli*
Transformation Kit using standard procedures (Zymo Research). They were transformed with
5 µl of the Gibson reaction. Transformants were selected onto LB plate supplemented with
chloramphenicol. Transformants with the correct plasmid were found by colony PCR with oligo
pairs L762/L763 and Dreamtaq DNA polymerase. Plasmids were extracted with the
NucleoSpin Plasmid kit (Macherey-Nagel) and checked by restriction digestion (NEB) and
sequencing (Eurofins).
Then, plasmids were transferred into competent *E. coli* strain MFDpir (Ferrières *et al.*, 2010)
prepared with the TSS method (Chung *et al.*, 1989). *E. coli MFDpir* produces the RP4
conjugation machinery, which allows the transfer of the suicide plasmid into *D. solani* by
conjugation. To do that, a few colonies of *D. solani* and MFDpir were mixed in the same
proportion in 500 µl LB and centrifuged for 2 min at 8000 rpm. The pellet was resuspended in
90 µl LB with 5 µl DAP at 57 mg/mL, and deposited onto a LB agar plate incubated at 30°C.
After 18h, the bacteria were resuspended in 1 ml LB, diluted in 10-fold series from 10^{-1} to 10^{-7}
and spread onto LB agar supplemented with chloramphenicol at 4 µg/l to select the first event

of recombination. Transconjugants re-isolated on this medium were then spread onto LB agar
without NaCl supplemented with 5% sucrose and incubated at 19°C for 2-3 days to allow the
second event of recombination. Sucrose-resistant colonies were then replicated onto Mac
Conkey medium supplemented with 1% arabinose incubated for 18h at 30°C to check the
arabinose catabolism, and LB-Cm plate to check plasmid loss.
497
498 **Construction of the secondary metabolite in-frame deletion mutants of *D. solani*.**
499
A spontaneous mutant of *D. solani* D s0432-1 resistant to nalidixic acid (Nal^R) was obtained
by growing the WT strain at 30°C in LB medium supplemented with Nal at 5 µg/mL for 18 h
before spreading the liquid culture onto LB agar plates with Nal at 10 µg/mL. The Nal^R strain
was named DS50 (Table S1). Then to discriminate in-frame deletion mutants from the wild-
type DS50 strain, a Gm^R derivative of strain DS50 was constructed by insertion of a mini-Tn7-
Gm cassette into the attTn7 site of *D. solani,* as previously performed with *D. dadantii* 3937
(Royet *et al.*, 2019) by using the plasmids pTn7-M (Zobel *et al.*, 2015) and pTnS3 (Choi *et al.*,
2008). The correct integration of the Gm^R cassette was then checked by colony PCR using
oligo pairs L365/L848 (amplification of 350 bp for the correct integration) (Table S2). This gave
the Nal^R Gm^R strain DS58. Subsequently, the in-frame deletion mutations Δ*ssmG* (NCBI
Reference Sequence: WP_022634121.1; cluster A), Δ*oocL* (NCBI Reference Sequence:
WP_023638021.1; cluster B) and Δ*zsmA* (NCBI Reference Sequence: WP_022632849.1;
cluster C) were constructed in strain DS58. Derivatives of the pRE112 suicide plasmids
containing upstream and downstream 500-bp DNA of the gene to delete were also constructed
(Table S2). In-frame deletions were checked by PCR as previously described (Table S2).
515
Growth inhibition assay of bacteria and yeast.
First, *B. subtilis*, *E. coli*, *D. dadantii* and *P. atrosepticum* were grown overnight in LB at 30°C
with shaking. LB agar was cooled until the temperature reached about 50°C (just before the
agar re-solidified). The OD$_{600}$ of the overnight culture of *B. subtilis*, *E. coli*, *D. dadantii* or *P.*
atrosepticum were adjusted to 1 with fresh LB. 100 mL of the LB agar in surfusion were mixed
with 100 µl of the OD$_{600}$ 1 culture of *B. subtilis*, *E. coli*, *D. dadantii* or *P. atrosepticum* . 25 mL
of inoculated LB agar was poured in 10- by 10-cm square plates. The plates were dried for a
few hours. *D. solani* strains were grown for 24h at 30°C with shaking in M63 medium
supplemented with sucrose and yeast extract or in LS4 medium supplemented with yeast
extract. The OD$_{600}$ of the *D. solani* cultures were adjusted to 1 and 5 µl were spotted onto the
inoculated square plates which were incubated at 30°C for 24-48 h before visualization of the
inhibition zone.

The same experiment was conducted with *Streptomyces scabiei*, except that it was grown for
3 days in TSB at 28°C and TSB agar was poured in the square plates. With the yeasts *S.*
cerevisiae and *K. lactis*, YPD medium was used.

Growth inhibition of fungal strains.
S. sclerotiorum, *B. cinerea* and *M. oryzae* were grown onto PDA plate at 25°C for 5, 7 and 10
days respectively. At the start of the experiments, Colonies of *D. solani* D s0432-1 strains were
resuspended in M63 medium and OD_{600} was adjusted to 0.1. Then, 10 µl of the bacterial
suspensions were deposited at the center of a 90-mm Petri dish with two 5 mm agar plugs of
fungus at 4.5-cm around. The radius of fungus inhibition zone was measured.

Pea aphid survival assay
Aphid survival assays were performed at 20-25°C as previously described (Grenier *et al.*,
2006). Pea aphid (*Acyrthosiphon pisum*, LL01 clone, alfalfa strain reared on faba bean)
individuals aged 0-24 h were transferred on Ap3 diet (Febvay *et al.*, 1988) loaded with
10^7/mL cells of *D. solani* D S0432-1 or its corresponding mutants. After a full day on diet,
aphids were transferred on single *V. faba* plantlets and scored daily for survival up to the 7[th]
day. Survival data were analysed with JMP software (SAS Inc) and its survival platform, and
data adjusted through Kaplan-Meier graphing and analysis, followed by a Weibull fit (best of
tested log-normal or exponential fits). Data are fully compatible with previously published
work on *D. dadantii* (Costechareyre *et al.*, 2010), except that temperature was less
stringently controlled.

Chicory inoculation experiments
The *D. solani* WT strain and the different mutants were grown overnight in LB medium.
Bacteria were washed in M63 medium and OD_{600} was adjusted to 1. One microliter of the
bacterial suspension was inoculated into a small hole made into the leaves by a yellow pipette
tip. The wound was covered with mineral oil and the leaves were incubated at 30°C at high
humidity for 24 h. The length of rotten tissue was measured.

ACKNOWLEDGMENTS
This work was supported by the Centre National de la Recherche Scientifique (CNRS), Institut
National des Sciences Appliquées (INSA) and by the University Lyon I (UCBL). We thank them
for their continued financial support. T.B. was supported by a PhD grant from the Ministère de
l'Enseignement Supérieur, de la Recherche et de l'Innovation. We thank Veronique Utzinger
for technical assistance, the members of the Membrane Trafficking and Signaling in Bacteria
(MTSB) team for discussions, Nathalie Poussereau and Mathias Choquer for advice on fungi

growth and providing the strains, Alexandre Soulard and Jade Ravent for advice on yeast
growth and providing strains and medium. We do not thank the Sars-CoV-2 virus that forced
us to close the lab for several months in 2020, delaying experiments.
568

569   **AUTHOR CONTRIBUTION**
570

571   E.G., B.T, R.Y and G.E carried out the experiments. G.E and H-C-P.N wrote the manuscript
572   with support from E.G, B.T and R.Y. H-C-P.N and G.E conceived the original idea. G.E
573   supervised the project. All authors provided critical feedback and helped shape the research,
574   analysis and manuscript.
575

580   **FUNDING**
581   This work was supported by annual credits from the University Lyon I and the CNRS, given to
582   the laboratory without writing a funding application.
583

584   **REFERENCES**
585

Blin, P., Robic, K., Khayi, S., Cigna, J., Munier, E., Dewaegeneire, P., et al. (2021) Pattern
and causes of the establishment of the invasive bacterial potato pathogen Dickeya solani and
of the maintenance of the resident pathogen D. dianthicola. *Mol Ecol* **30**: 608–624.
Brady, C.L., Cleenwerck, I., Denman, S., Venter, S.N., Rodríguez-Palenzuela, P., Coutinho,
T.A., and De Vos, P. (2012) Proposal to reclassify Brenneria quercina (Hildebrand and Schroth
1967) Hauben et al. 1999 into a new genus, Lonsdalea gen. nov., as Lonsdalea quercina
comb. nov., descriptions of Lonsdalea quercina subsp. quercina comb. nov., Lonsdalea
quercina subsp. iberica subsp. nov. and Lonsdalea quercina subsp. britannica subsp. nov.,
emendation of the description of the genus Brenneria, reclassification of Dickeya
dieffenbachiae as Dickeya dadantii subsp. dieffenbachiae comb. nov., and emendation of the
description of Dickeya dadantii. *Int J Syst Evol Microbiol* **62**: 1592–1602.
Cane, D.E. and Walsh, C.T. (1999) The parallel and convergent universes of polyketide
synthases and nonribosomal peptide synthetases. *Chem Biol* **6**: R319–R325.
Charkowski, A., Blanco, C., Condemine, G., Expert, D., Franza, T., Hayes, C., et al. (2012)
The role of secretion systems and small molecules in soft-rot Enterobacteriaceae
pathogenicity. *Annu Rev Phytopathol* **50**: 425–449.

Choi, K.-H., Mima, T., Casart, Y., Rholl, D., Kumar, A., Beacham, I.R., and Schweizer, H.P.
(2008) Genetic tools for select-agent-compliant manipulation of Burkholderia pseudomallei.
Appl Environ Microbiol **74**: 1064–1075.

Chung, C.T., Niemela, S.L., and Miller, R.H. (1989) One-step preparation of competent
Escherichia coli: transformation and storage of bacterial cells in the same solution. *Proc Natl*
Acad Sci U S A **86**: 2172–2175.

Costechareyre, D., Dridi, B., Rahbe, Y., and Condemine, G. (2010) Cyt toxin expression
reveals an inverse regulation of insect and plant virulence factors of Dickeya dadantii. *Env*
Microbiol **12**: 3290–301.

Czajkowski, R., Fikowicz-Krosko, J., Maciag, T., Rabalski, L., Czaplewska, P., Jafra, S., et al.
(2020) Genome-Wide Identification of Dickeya solani Transcriptional Units Up-Regulated in
Response to Plant Tissues From a Crop-Host Solanum tuberosum and a Weed-Host Solanum
dulcamara. *Front Plant Sci* **11**: 580330.

Duprey, A., Taib, N., Leonard, S., Garin, T., Flandrois, J.P., Nasser, W., et al. (2019) The
phytopathogenic nature of Dickeya aquatica 174/2 and the dynamic early evolution of Dickeya
pathogenicity. *Environ Microbiol* **21**: 2809–2835.

Edwards, R.A., Keller, L.H., and Schifferli, D.M. (1998) Improved allelic exchange vectors and
their use to analyze 987P fimbria gene expression. *Gene* **207**: 149–157.

Febvay, G., Delobel, B., and Rahbé, Y. (1988) Influence of the amino acid balance on the
improvement of an artificial diet for a biotype of Acyrthosiphon pisum (Homoptera: Aphididae).
Can J Zool **66**: 2449–2453.

Ferrières, L., Hémery, G., Nham, T., Guérout, A.-M., Mazel, D., Beloin, C., and Ghigo, J.-M.
(2010) Silent Mischief: Bacteriophage Mu Insertions Contaminate Products of Escherichia coli
Random Mutagenesis Performed Using Suicidal Transposon Delivery Plasmids Mobilized by
Broad-Host-Range RP4 Conjugative Machinery. **192**: 6418–6427.

Fikowicz-Krosko, J. and Czajkowski, R. (2017) Fast and reliable screening system to preselect
candidate Dickeya solani Tn5 mutants in plant tissue-induced genes. *Eur J Plant Pathol* **149**:
1023–1027.

Garlant, L., Koskinen, P., Rouhiainen, L., Laine, P., Paulin, L., Auvinen, P., et al. (2013)
Genome Sequence of Dickeya solani, a New soft Rot Pathogen of Potato, Suggests its
Emergence May Be Related to a Novel Combination of Non-Ribosomal Peptide/Polyketide
Synthetase Clusters. *Diversity* **5**: 824–842.

Golanowska, M., Potrykus, M., Motyka-Pomagruk, A., Kabza, M., Bacci, G., Galardini, M., et
al. (2018) Comparison of Highly and Weakly Virulent Dickeya solani Strains, With a View on
the Pangenome and Panregulon of This Species. *Front Microbiol* **9**: 1940.

Grenier, A.-M., Duport, G., Pagès, S., Condemine, G., and Rahbé, Y. (2006) The
Phytopathogen Dickeya dadantii (Erwinia chrysanthemi 3937) Is a Pathogen of the Pea Aphid.

639 *Appl Environ Microbiol* **72**: 1956–1965.

640 Hellberg, J.E.E.U., Matilla, M.A., and Salmond, G.P.C. (2015) The broad-spectrum antibiotic,

641 zeamine, kills the nematode worm Caenorhabditis elegans. *Front Microbiol* **6**:.

642 Hugouvieux-Cotte-Pattat, N., Brochier-Armanet, C., Flandrois, J.-P., and Reverchon, S.

643 (2020) Dickeya poaceiphila sp. nov., a plant-pathogenic bacterium isolated from sugar cane

644 (Saccharum officinarum). *Int J Syst Evol Microbiol* **70**: 4508–4514.

645 Hugouvieux-Cotte-Pattat, N., Condemine, G., Gueguen, E., and Shevchik, V.E. (2020)

646 Dickeya Plant Pathogens. In *eLS*. American Cancer Society, pp. 1–10.

647 Hugouvieux-Cotte-Pattat, N., Condemine, G., and Shevchik, V.E. (2014) Bacterial pectate

648 lyases, structural and functional diversity. *Environ Microbiol Rep* **6**: 427–440.

649 Hugouvieux-Cotte-Pattat, N., Jacot-des-Combes, C., and Briolay, J. (2019) Dickeya lacustris

650 sp. nov., a water-living pectinolytic bacterium isolated from lakes in France. *Int J Syst Evol*

651 *Microbiol* **69**: 721–726.

652 Khayi, S., Blin, P., Pédron, J., Chong, T.-M., Chan, K.-G., Moumni, M., et al. (2015) Population

653 genomics reveals additive and replacing horizontal gene transfers in the emerging pathogen

654 Dickeya solani. *BMC Genomics* **16**: 788.

655 Koskiniemi, S., Lamoureux, J.G., Nikolakakis, K.C., t'Kint de Roodenbeke, C., Kaplan, M.D.,

656 Low, D.A., and Hayes, C.S. (2013) Rhs proteins from diverse bacteria mediate intercellular

657 competition. *Proc Natl Acad Sci U S A* **110**: 7032–7037.

658 Liang, Z., Huang, L., He, F., Zhou, X., Shi, Z., Zhou, J., et al. (2019) A Substrate-Activated

659 Efflux Pump, DesABC, Confers Zeamine Resistance to Dickeya zeae. *mBio* **10**:.

660 Liao, L., Cheng, Y., Liu, S., Zhou, J., An, S., Lv, M., et al. (2014) Production of Novel Antibiotics

661 Zeamines through Optimizing Dickeya zeae Fermentation Conditions. *PLoS ONE* **9**: e116047.

662 Lisicka, W., Fikowicz-Krosko, J., Jafra, S., Narajczyk, M., Czaplewska, P., and Czajkowski, R.

663 (2018) Oxygen Availability Influences Expression of Dickeya solani Genes Associated With

664 Virulence in Potato (Solanum tuberosum L.) and Chicory (Cichorium intybus L.). *Front Plant*

665 *Sci* **9**: 374.

666 Loria, R., Kers, J., and Joshi, M. (2006) Evolution of Plant Pathogenicity in Streptomyces.

667 *Annu Rev Phytopathol* **44**: 469–487.

668 Masschelein, J., Jenner, M., and Challis, G.L. (2017) Antibiotics from Gram-negative bacteria:

669 a comprehensive overview and selected biosynthetic highlights. *Nat Prod Rep* **34**: 712–783.

670 Masschelein, J., Mattheus, W., Gao, L.-J., Moons, P., Houdt, R.V., Uytterhoeven, B., et al.

671 (2013) A PKS/NRPS/FAS Hybrid Gene Cluster from Serratia plymuthica RVH1 Encoding the

672 Biosynthesis of Three Broad Spectrum, Zeamine-Related Antibiotics. *PLOS ONE* **8**: e54143.

673 Matilla, M.A., Leeper, F.J., and Salmond, G.P.C. (2015) Biosynthesis of the antifungal

674 haterumalide, oocydin A, in Serratia, and its regulation by quorum sensing, RpoS and Hfq.

675 *Environ Microbiol* **17**: 2993–3008.

Matilla, M.A., Stöckmann, H., Leeper, F.J., and Salmond, G.P.C. (2012) Bacterial Biosynthetic
Gene Clusters Encoding the Anti-cancer Haterumalide Class of Molecules. *J Biol Chem* **287**:
39125–39138.
Motyka-Pomagruk, A., Zoledowska, S., Misztak, A.E., Sledz, W., Mengoni, A., and Lojkowska,
E. (2020) Comparative genomics and pangenome-oriented studies reveal high homogeneity
of the agronomically relevant enterobacterial plant pathogen Dickeya solani. *BMC Genomics*
21: 449.
Oulghazi, S., Pédron, J., Cigna, J., Lau, Y.Y., Moumni, M., Van Gijsegem, F., et al. (2019)
Dickeya undicola sp. nov., a novel species for pectinolytic isolates from surface waters in
Europe and Asia. *Int J Syst Evol Microbiol* **69**: 2440–2444.
Parkinson, N., DeVos, P., Pirhonen, M., and Elphinstone, J. (2014) Dickeya aquatica sp. nov.,
isolated from waterways. *Int J Syst Evol Microbiol* **64**: 2264–2266.
Pédron, J., Mondy, S., des Essarts, Y.R., Van Gijsegem, F., and Faure, D. (2014) Genomic
and metabolic comparison with Dickeya dadantii 3937 reveals the emerging Dickeya solani
potato pathogen to display distinctive metabolic activities and T5SS/T6SS-related toxin
repertoire. *BMC Genomics* **15**: 283.
Potrykus, M., Golanowska, M., Hugouvieux-Cotte-Pattat, N., and Lojkowska, E. (2014)
Regulators involved in Dickeya solani virulence, genetic conservation, and functional
variability. *Mol Plant-Microbe Interact MPMI* **27**: 700–711.
Potrykus, M., Hugouvieux-Cotte-Pattat, N., and Lojkowska, E. (2018) Interplay of classic Exp
and specific Vfm quorum sensing systems on the phenotypic features of Dickeya solani strains
exhibiting different virulence levels. *Mol Plant Pathol* **19**: 1238–1251.
Rodicio, R. and Heinisch, J.J. (2013) Yeast on the milky way: genetics, physiology and
biotechnology of Kluyveromyces lactis. *Yeast* **30**: 165–177.
Royet, K., Parisot, N., Rodrigue, A., Gueguen, E., and Condemine, G. (2019) Identification by
Tn-seq of Dickeya dadantii genes required for survival in chicory plants. *Mol Plant Pathol* **20**:
287–306.
Samson, R., Legendre, J.B., Christen, R., Saux, M.F.-L., Achouak, W., and Gardan, L. (2005)
Transfer of Pectobacterium chrysanthemi (Burkholder et al. 1953) Brenner et al. 1973 and
Brenneria paradisiaca to the genus Dickeya gen. nov. as Dickeya chrysanthemi comb. nov.
and Dickeya paradisiaca comb. nov. and delineation of four novel species, Dickeya dadantii
sp. nov., Dickeya dianthicola sp. nov., Dickeya dieffenbachiae sp. nov. and Dickeya zeae sp.
nov. *Int J Syst Evol Microbiol* **55**: 1415–1427.
Strobel, G., Li, J.-Y., Sugawara, F., Koshino, H., Harper, J., and Hess, W.M. (1999) Oocydin
A, a chlorinated macrocyclic lactone with potent anti- oomycete activity from Serratia
marcescens. *Microbiology* **145**: 3557–3564.
Tian, Y., Zhao, Y., Yuan, X., Yi, J., Fan, J., Xu, Z., et al. (2016) Dickeyafangzhongdai sp. nov.,

a plant-pathogenic bacterium isolated from pear trees (Pyrus pyrifolia). *Int J Syst Evol*

Microbiol **66**: 2831–2835.

Van Gijsegem, F., Toth, I.K., and van der Wolf, J.M. (2021) Soft Rot Pectobacteriaceae: A

Brief Overview. In *Plant Diseases Caused by Dickeya and Pectobacterium Species*. Van

Gijsegem, F., van der Wolf, J.M., and Toth, I.K. (eds). Cham: Springer International Publishing,

pp. 1–11.

Van Gijsegem, F., van der Wolf, J.M., and Toth, I.K. eds. (2021) Plant Diseases Caused by

Dickeya and Pectobacterium Species, Cham: Springer International Publishing.

Wang, X., He, S.-W., Guo, H.-B., Han, J.-G., Thin, K.K., Gao, J.-S., et al. (2020) Dickeya

oryzae sp. nov., isolated from the roots of rice. *Int J Syst Evol Microbiol* **70**: 4171–4178.

Wésolowski-Louvel, M. (2011) An efficient method to optimize Kluyveromyces lactis gene

targeting. *FEMS Yeast Res* **11**: 509–513.

van der Wolf, J.M., Nijhuis, E.H., Kowalewska, M.J., Saddler, G.S., Parkinson, N., Elphinstone,

J.G., et al. (2014) Dickeya solani sp. nov., a pectinolytic plant-pathogenic bacterium isolated

from potato (Solanum tuberosum). *Int J Syst Evol Microbiol* **64**: 768–774.

Zhou, J., Cheng, Y., Lv, M., Liao, L., Chen, Y., Gu, Y., et al. (2015) The complete genome

sequence of Dickeya zeae EC1 reveals substantial divergence from other Dickeya strains and

species. *BMC Genomics* **16**: 571.

Zhou, J., Zhang, H., Wu, J., Liu, Q., Xi, P., Lee, J., et al. (2011) A Novel Multidomain Polyketide

Synthase Is Essential for Zeamine Production and the Virulence of Dickeya zeae. *Mol Plant-*

Microbe Interactions® **24**: 1156–1164.

Zobel, S., Benedetti, I., Eisenbach, L., de Lorenzo, V., Wierckx, N., and Blank, L.M. (2015)

Tn7-Based Device for Calibrated Heterologous Gene Expression in Pseudomonas putida.

ACS Synth Biol **4**: 1341–1351.

SUPPLEMENTAL DATA

Figure S1. Plate-based bioassays against *B. subtilis* and *K. lactis* for metabolite

production by *Dickeya solani* WT cultivated in various medium.

Table S1. Strains and plasmids used in the study.

Table S2. oligonucleotides used in the study.