The genetic basis of interspecies cooperation and conflict in naturally-occurring isolates of a model plant symbiont

Rebecca T. Batstone[a,1], **Liana T. Burghardt**[b], **and Katy D. Heath**[a,c]

[a]Carl R. Woese Institute for Genomic Biology, University of Illinois at Urbana-Champaign, 1206 West Gregory Drive, Urbana, IL 61801 USA; [b]Department of Plant Science, The Pennsylvania State University, 103 Tyson Building, University Park, PA, 16802 USA; [c]Department of Plant Biology, University of Illinois at Urbana-Champaign, 286 Morrill Hall, 505 South Goodwin Avenue, Urbana, IL 61801 USA

Given the need to predict the outcomes of evolution in host-associated microbiomes, whether microbial and host fitnesses tend to trade off,
generating conflict, remains a pressing question. Examining the relationships between host and microbe fitness proxies at both the phenotypic
and genomic levels can illuminate the mechanisms underlying interspecies cooperation and conflict. We conducted genome-wide association
studies (GWAS) to examined naturally-occurring genetic variation in 191 strains of the model microbial symbiont, *Ensifer meliloti*, paired with
each of two host *Medicago truncatula* genotypes in single- or multi-strain experiments. Using this data, we determine how multiple proxies
of microbial and host fitness were related to one another and test key predictions about mutualism evolution at the genomic scale, while
also addressing the challenge of measuring microbial fitness. We found little evidence for interspecies fitness conflict; loci tended to have
concordant effects on both microbe and host fitnesses, even in environments with multiple co-occurring microbes. Our results emphasize the
importance of quantifying microbial relative fitness for understanding microbiome evolution and thus harnessing microbiomes to improve host
fitness. Additionally, we find that mutualistic coevolution between hosts and microbes potentially acts to maintain, rather than erode, genetic
diversity, thus, explaining why variation in mutualism traits persists in nature.

conflict | cooperation | GWAS | mutualism | pleiotropy

A central goal of genetics research is to build a reliable map of genotype-to-phenotype relationships (G→P map) in order to uncover the genetic basis of important organismal phenotypes, including plant yield in agriculture or disease-related traits in humans (1–3). Recent advances in sequencing and microbiome research have revealed the ubiquity of microbial symbioses, meaning that many of these host phenotypes (yield, disease resistance, et c.) are actually **symbiotic extended phenotypes**, their variation being influenced by loci present within interacting microbial symbionts in addition to the host (4–9). Studies of the G→P map in single organisms have highlighted the importance of **pleiotropy**, i.e., loci that influence more than one trait. Pleiotropy can constrain adaptation when it underlies trade-offs among fitness-related traits of the same organism (10–12), and thus, has important consequences for development, aging, and disease (reviewed in 13). In the G→P map of symbiotic extended phenotypes, pleiotropic loci can influence fitness-related traits of both host and symbiont, which we henceforth refer to as **symbiotic pleiotropy**. Importantly, symbiotic pleiotropy determines the degree to which partners' fitnesses are aligned (i.e., same-sign or concordant effects) or in conflict (i.e., opposite-sign or discordant effects). Identifying the loci underlying symbiotic pleiotropy is therefore critical not only for illuminating the genetic basis of symbiotic extended phenotypes, but also for predicting how hosts and symbionts coevolve in nature.

In symbiotic mutualisms, wherein partners trade fitness benefits (14), whether fitness conflict or alignment drives the evolution of these interactions is hotly debated (15–20). Individuals that

benefit from the interaction without paying the costs, also known as **cheaters**, are predicted to be selectively favoured, given that they should outcompete cooperative counterparts (21–24). At the phenotypic level, cheaters would be seen as symbiont genotypes that gain fitness at the host's expense (i.e., points in the bottom right quadrant of **Fig. 1A**), while fitness conflict would be seen as an overall negative correlation (grey line in **Fig. 1A**). In contrast, cooperators would be seen as symbiont genotypes whose increase in fitness is associated with an increase in host fitness (i.e., points in the top right quadrant of **Fig. 1A**), while fitness alignment would be seen as an overall positive correlation (orange line in **Fig. 1A**).

Empirical evidence supporting conflict or alignment in symbiotic mutualisms is mixed: when examining relationships between host and symbiont fitness proxies, some studies have observed an overall negative (4, 25, 26) while others have found a positive (15, 27) correlation. Recent experimental evolution studies (28–30) have failed to detect the evolution of cheaters and instead found that microbial adaptation to particular host genotypes is associated with an increase, rather than a decrease, in **partner quality**, i.e., the fitness benefit a symbiont genotype confers to its host. Yet, fitness alignment at the phenotypic level does not necessarily preclude fitness conflict at the genomic level: rather than dichotomous categories of "cooperator" or "cheater", mutualist genomes are better viewed as mosaics of loci (8), some underlying cooperation while others underlie conflict. Whether coevolution resulting from symbiosis leads to more beneficial interactions and greater mutualism stability, or alternatively, more antagonism and less stable interactions, therefore requires examining the relationships between host and symbiont fitness proxies at both the phenotypic and genomic levels.

Genome-wide association studies (i.e., GWAS) can be used to reveal the genes, as well as specific segregating mutations (i.e., variants), that underlie variation in symbiotic extended phenotypes in natural populations (28, 31–35). Because they provide an estimate of both the strength and direction of effects of particular alleles on the trait, henceforth referred to as **allelic effect sizes**, GWAS are especially useful for identifying loci underlying symbiotic pleiotropy, and thus, defining the mutational landscape of mutualism evolution in nature. For example, if symbiotic pleiotropy is extensive and its effects on fitness-related traits in the interacting partners tend to be discordant (grey quadrants in **Fig. 1B**), then conflict should underlie the evolution of mutualism, allowing for the possibility of cheating individuals that are competitively superior, as mutualism theory

R.T.B., K.D.H, and L.T.B conceived of the work. R.T.B led the data analysis and drafted the article with guidance from K.D.H. and L.T.B., who also provided critical revisions and approved the final version for submission.

The authors declare no competing interests.

[1]To whom correspondence should be addressed. E-mail: rtbatstone@gmail.com

predicts ([21–24](#)). In contrast, if pleiotropic effects are overwhelmingly concordant (orange quadrants in **Fig. 1B**), fitness alignment rather than conflict should be the null hypothesis in mutualism, and models relying on cheating genotypes as the main driver of mutualism evolution may not be suitable. GWAS can also facilitate the study of how historical selection has acted on loci underlying symbiotic pleiotropy in the past ([8, 36](#)). For example, loci underlying fitness conflict might resemble those underlying antagonistic arms race dynamics, showing signatures of rapid turnover and higher rates of evolutionary change as rare alleles are selectively favoured ([37](#)). In contrast, loci underlying fitness alignment may exhibit signatures of evolutionary stability and lower rates of evolutionary change as more common alleles are selectively favoured ([37](#)). Because GWAS has the power to reveal individual allelic effects, it can be used to identify loci underlying fundamental sources of conflict at the genomic-level (e.g., grey quadrants, **Fig. 1B**) even if fitness alignment is realized at the phenotypic-level (e.g., orange line, **Fig. 1A**).

Over the past ∼25 years, legume-rhizobium symbioses have been developed as models for understanding mutualism evolution ([38–41](#)). This interaction is one of the most ecologically and economically important symbioses, contributing upwards of 20 million tonnes of nitrogen (N) to the global N-cycle, and saving billions of dollars that would have otherwise been spent on synthetic N fertilizer production ([42](#)). Legumes house rhizobia within specialized root nodules and supply them with photosynthate, while rhizobia within the nodules convert atmospheric N to a plant-usable form, resulting in a beneficial exchange of resources. Key traits of this symbiosis, such as plant biomass and nodule number, are known to be influenced by variants in the genomes of both host and microbes, as well as the epistatic interactions between them (Genotype-by-Genotype or G x G interactions; e.g., [4, 34, 38, 43–46](#)), making this symbiosis an excellent model for understanding mutualistic coevolution.

A quantitative comparison of single-strain and multi-strain proxies of rhizobium fitness, as well as their correlations with plant fitness, is needed to reveal interspecies conflict. Seed number, or close correlates such as aboveground (shoot) biomass and leaf chlorophyll A content, are well-established proxies for reproductive fitness in annual plants ([25, 31, 47](#)) (**Fig. 1C**). In contrast, estimating rhizobium fitness has been an empirical and conceptual challenge ([15–17, 48, 49](#)). Early attempts to estimate rhizobium fitness relied on single-strain experiments, whereby plants were inoculated with a single rhizobium strain, and fitness proxies including nodule number and nodule size were measured (**Fig. 1D**). Both measures reflect rhizobium fitness because a rhizobium that establishes a nodule in symbiosis can gain a fitness benefit on the order of $10^5 - 10^7$ ([50](#)), larger nodules release more rhizobia ([26, 47, 49](#)), and intuitively, strains that produce more nodules will release more rhizobia ([4, 26, 41](#)). While rhizobium fitness proxies measured in single-strain experiments can be directly correlated with

partner quality (e.g., shoot biomass), these measures have been criticized for producing spurious positive correlations due to stronger fitness feedbacks between host plants and rhizobia (16). In contrast, multi-strain experiments decouple individual strain fitness from host growth and better reflect rhizobia fitness in natural and agricultural soils where many strains coexist. Recent advances merging population genomics and metagenomics have enabled measuring **rhizobium relative fitness**, i.e., a strain's ability to compete for nodulation opportunities and extract host resources once in nodules when other strains are present (33, 51). Multi-strain experiments that have estimated rhizobium relative fitness have so far hinted at a surprising lack of correlation between single-strain and multi-strain measures of rhizobium fitness (33), motivating our comprehensive analysis here.

Here we combine two datasets: first, GWAS that test for the association between rhizobium variants and both plant and rhizobium fitness proxies measured in single-strain experiments (**Fig. 1C**), whereby 191 strains of *Ensifer meliloti* collected from natural populations were inoculated individually onto one of two genotypes of the host plant *Medicago truncatula*. Second, a GWAS based on a new dataset that measured rhizobium relative fitness (**Fig. 1D**) in a multi-strain experiment, whereby the same two host genotypes were inoculated with 89 of the 191 strains used in the single-strain experiments. We combine these datasets to first ask: What are the relationships among rhizobium fitness proxies measured in both single-strain (nodule number, nodule size) and multi-strain (rhizobium relative fitness) experiments? Are these proxies genetically distinct and thus likely to evolve independently, or are they linked through pleiotropy, and thus, evolve together? We use this information to next address the potential for genomic conflict in this symbiosis, asking: Do variants tend to have aligned or conflicting effects on host and symbiont fitnesses? Finally we ask whether there is any evidence that historical selection has differentially shaped loci underlying fitness alignment versus conflict, as we might predict under different models of mutualism evolution. To our knowledge, this is the first study to interrogate the genetic basis of cooperation and conflict across the entire rhizobium genome using naturally occurring populations of strains from their native range.

Materials and Methods

Study system. Full details are provided in Riley et al. (52) and **SI Methods**. *Ensifer* (formerly *Sinorhizobium*) *meliloti* is an Alphaproteobacteria that forms symbiosis with various *Medicago* spp., fixing atmospheric N in return for plant photosynthate. All 191 strains used here were collected from the native range of the host *Medicago truncatula*, spanning Spain to France (as detailed in 52). *E. meliloti* has a multipartite genome, including a large (∼3.7 Mbp) chromosome and two symbiosis plasmids (pSymA and pSymB, ∼1.4 Mbp and ∼1.7 Mbp, respectively); pSymA contains many of

the canonical genes for nodulation and N-fixation ([53], [54]). We used two lines of *M. truncatula* DZA
315.16 (hereafter DZA) and A17 in separate single-strain experiments and a multi-strain experiment
detailed below.

Single-strain experiments. Full details are provided in Batstone et al. ([46]), and **SI Methods**.
Briefly, we conducted two separate experiments in Sept and Nov 2018, one for each *M. truncatula*
line, DZA and A17, respectively. At least three replicate seeds of each plant line were planted
individually into pots, and were singly inoculated with one of 191 strains of *E. meliloti* described
above. Within each experiment, we measured proxies for both plant fitness (shoot biomass, leaf
chlorophyll A content) and rhizobium fitness (nodule number, per nodule weight; **Fig. 1C,D**).

Multi-strain experiment. We grew A17 and DZA hosts with a multi-strain inoculum composed of 89
of the 191 strains used in the single-strain experiments (described in more detail in **SI Methods**).
Using the "select-and-resequence" approach ([33]), this experiment allowed us to generate new data
on **rhizobium relative fitness**, which represents a strain's ability to both compete for nodulation
opportunities and extract host resources once in nodules when 88 other strains were present (**Fig. 1D**). This fitness proxy was obtained by sequencing pooled nodule samples from each plant at the
end of the experiment and estimating each strain's frequency using a haplotype reconstruction
method ([55]). We then obtained each strain's relative fitness by calculating the fold change in the
frequency of a strain at the end of the experiment relative to its mean frequency at the beginning
(see **SI Methods**).

Phenotypic analyses. As described in Batstone et al. ([46]), we calculated the estimated marginal
means for each fitness proxy in each experiment, correcting for the effect of rack, using the emmeans
package (v1.4.1, [56]) in the R environment ([57]). We then conducted linear pairwise regressions (lm
option in base R) for each fitness proxy against the other within each plant line.

DNA isolation, whole-genome sequencing, and variant calling. Detailed methods are provided in
Riley et al. ([52]), and **SI Methods**. We obtained DNA from each of the 191 rhizobium isolates,
sequenced their full genomes including the chromosome and two symbiosis plasmids, used a common
reference-based assembly pipeline to align sequences and call single nucleotide polymorphisms
(SNPs), and filtered the resulting SNPs based on sequence quality, depth, minor allele frequency,
and missingness, resulting in a total of 36,526 filtered SNPs.

Genome-wide association tests. Detailed methods are provided in **SI Methods**. We conducted
linear mixed models (LMMs) implemented in GEMMA (v0.98.1, [58]) to quantify **allelic effect sizes**

that represent both the strength and direction of the association between variants and fitness proxies after correcting for rhizobium population structure. We ran ten separate association analyses in GEMMA, one for each of the four fitness proxies measured in single-strain experiments (**Fig. 1C**), plus rhizobium relative fitness measured in the multi-strain experiment, and for both host plant lines (DZA, A17; five proxies x two hosts = ten runs).

Genomic analyses. We first identified pleiotropic variants as those that were significantly associated with more than one trait on the same host line, significance determined using a permutation method (described in more detail in the **SI Methods**). When variants were significantly associated with two or more rhizobium fitness proxies, they were designated as underlying **rhizobium fitness pleiotropy**, whereas when they were associated with at least one rhizobium AND plant fitness proxy, they were designated as underlying **symbiotic pleiotropy**. Next, we categorized whether pleiotropic variants had discordant (opposite-sign; +,- or -,+) or concordant (same-sign; +,+ or -,-) effects on pairwise fitness proxy combinations (**Fig. 1B**). Finally, to test whether selection acted on genes underlying rhizobium fitness and symbiotic pleiotropy, we used the R package PopGenome (v2.7.5, 59) to compute several commonly used statistics that test for signatures of historical selection and/or departures of neutrality, namely, nucleotide diversity (i.e., π), Tajima's D, as well as Fu and Li's F and D (60, 61). Additional details appear in **SI Methods**.

Data availability. Strains and plant lines are available upon request. All raw data and analysis code are available on GitHub (see "Genetics_conflict_cooperation" folder). The Supplementary Information doc contains additional methods, results, figures and tables, as well as descriptions of datasets that have been uploaded along with this manuscript, and can be additionally accessed from bioRχiv and GitHub. Once raw sequence reads and assemblies are archived and made accessible on NCBI, accession numbers will be added to this manuscript.

Results

Relationships among rhizobium fitness proxies: nodule number, nodule size, and rhizobium relative fitness. In single-strain experiments, when we regressed strain means for nodule number and nodule weight, we found significant negative correlations for both hosts (**Fig. 2A**, left), indicating that strains creating larger nodules tended to form fewer total nodules on both host genotypes. At the genomic-level, most variants had discordant effects (**Fig. 2A**, right), similarly indicating a trade-off whereby variants that were positively associated with nodule number tended to be negatively associated with nodule weight, or vice-versa.

Comparing single- and multi-strain experiments, when we regressed strain means for rhizobium

relative fitness and nodule number, we found a weak but significant negative relationship for host line DZA only (**Fig. 2B**, left), indicating that strains that were more common in the multi-strain experiment formed fewer nodules in the single-strain experiment. At the genomic-level on DZA, most variants had discordant effects (**Fig. 2B**, right), again indicating a trade-off between nodule number and rhizobium relative fitness. For host line A17, most pleiotropic variants (20/28) had discordant effects (**Fig. 2B**, right), despite no relationship between these fitness proxies at the phenotypic level (**Fig. 2B**, left).

We found a significant, positive relationship between strain means for rhizobium relative fitness and nodule weight, again for DZA only (**Fig. 2C**, left). This result indicates that strains that were more commonly found in nodules in the multi-strain experiment formed larger nodules in the single-strain experiment. At the genomic-level, all pleiotropic variants were concordant (i.e., appearing in the top right or bottom left quadrants of **Fig. 2C**, right). For A17, while we found a lack of significant phenotypic correlation between nodule weight and rhizobium relative fitness (**Fig. 2C**, left), at the genomic level, all pleiotropic variants had discordant effects on these fitness proxies (**Fig. 2C**, right), indicating a trade-off at this scale. However, we found fewer total pleiotropic variants underlying the relationship between rhizobium relative fitness and nodule weight compared to the other two fitness proxy combinations (**SI Fig. S2A**, right), suggesting that nodule weight and rhizobium relative fitness are largely governed by different molecular mechanisms, and thus, likely to evolve independently.

These results overall support trade-offs at the phenotypic level, plus underlying discordant pleiotropy at the genomic level, suggesting that the genetic mechanisms rhizobia use to form more nodules in single-strain experiments make them less able to proliferate within and compete for nodules in multi-strain experiments. We additionally found host-dependent relationships among rhizobium fitness proxies at both the phenotypic and genomic levels, especially when comparing fitness proxies measured in single- versus multi-strain experiments.

Relationships between plant and rhizobium fitness proxies. We found no correlation, in either host, between strain means for nodule number and shoot biomass (**Fig. 3A** left), or nodule number and leaf chlorophyll A content on A17 (**SI Fig. 3A**, left). However we found a significantly negative correlation between nodule number and chlorophyll A for DZA (**SI Fig. 3A**, left). Assuming chlorophyll A is related to the N-fixation efficiency of each strain (62–64), this result suggests that strains forming more numerous (and smaller) nodules tended to fix less N. At the genomic level, most variants had concordant effects for nodule number and shoot biomass on DZA, whereas the opposite was true for the same proxy pair on A17 (**Fig. 3A**, right). Most variants had discordant

effects on nodule number and chlorophyll A on both hosts (**SI Fig. 3A**, right).

Strain means for nodule weight and shoot biomass were significantly positively correlated for both hosts (**Fig. 3B**, left). Nodule weight and chlorophyll A were significantly positively correlated on DZA, but not on A17 (**SI Fig. 3B**, left). At the genomic level for both hosts, all pleiotropic variants had concordant effects for nodule weight and shoot biomass (**Fig. 3B**, right), as well as nodule weight and chlorophyll A (**SI Fig. 3B**, right).

Finally, rhizobium relative fitness and shoot biomass were significantly positively correlated for DZA but not for A17 (**Fig. 3C**, left), whereas rhizobium relative fitness and leaf chlorophyll A content were significantly positively correlated on both hosts (**SI Fig. 3C**, left). At the genomic level, most pleiotropic variants had concordant effects on rhizobium relative fitness and shoot biomass (**Fig. 3C**, right), and this pattern was even stronger between rhizobium relative fitness and chlorophyll A on DZA with all but one pleiotropic variant having concordant effects (**SI Fig. 3C** right). Variant effects were mixed on A17 for rhizobium relative fitness and leaf chlorophyll A content (**SI Fig. 3C**, right).

Overall, we found mostly concordant relationships between rhizobium and plant fitnesses at both the phenotypic and genomic levels, suggesting a strong signal of fitness alignment in natural rhizobium populations. However, we note a lack of pleiotropic variants underlying the relationship between leaf chlorophyll A content and both rhizobium fitness proxies measured in single-strain experiments (i.e., nodule number and nodule weight; **SI Fig. S2C**, left & middle), suggesting these proxies are governed by different molecular mechanisms, and thus, should evolve independently.

Selection acts differently on genes underlying alignment versus conflict. Using multiple diversity and neutrality metrics, we found that rhizobium genes underlying fitness alignment exhibit higher nucleotide diversity and stronger signatures of balancing selection compared to any other gene category analyzed. Specifically, genes underlying concordant symbiotic pleiotropy (i.e., solid orange lines in **Fig. 4**) had significantly elevated statistics relative to the null (i.e., distributions in **Fig. 4**, based on all genes containing significant variants identified by GWAS) for three of the four test statistics (**SI Table S1**). We did not see any significant deviations from the null for genes underlying discordant symbiotic pleiotropy (i.e., dotted grey lines in **Fig. 4**), or for genes underlying both concordant and discordant rhizobium fitness pleiotropy (**SI Table S1, Fig. S4, Dataset S1**).

The genetic basis of symbiotic pleiotropy is host-dependent. Comparing the identities and putative functions of variants underlying symbiotic pleiotropy on both host lines revealed little overlap – concordant variants giving rise to both high host and symbiont fitness largely differed between the two host genotypes. Specifically, we identified a total of 168 variants underlying

symbiotic pleiotropy, corresponding to 128 coding-genes (see **SI Dataset S2** for variant-level and **SI Table S2 & Dataset S3** for gene-level summaries). 60 and 93 of these variants were uniquely associated with fitness proxies measured on plant lines A17 and DZA, respectively, while only 15 variants were shared between hosts. We highlight some of the noteworthy genes identified in our analysis for each pleiotropic category in **SI Results**.

Discussion

Leveraging genomics to quantify the genetic architecture underlying symbiotic extended phenotypes gives us the power to address long-standing issues in mutualism evolution with genome-scale resolution. Our results overall suggest that: 1) fitness alignment between hosts and rhizobia is common at both the phenotypic and genomic levels, with genes underlying alignment showing elevated nucleotide diversity and signatures of balancing selection; and 2) the lack of a relationship or even trade-offs between rhizobium nodule number and rhizobium relative fitness mean that measures of rhizobium fitness in multi-strain experiments should be prioritized when we want to predict rhizobium evolution. We discuss these main points in turn below.

Alignment of host and symbiont fitnesses. Many models of mutualism rely on the key assumption that cooperation is inherently costly and thus that selection should favour less-cooperative, potentially 'cheating', partners that forgo paying the costs whilst continuing to receive benefits – particularly in one-to-many symbioses where a 'tragedy of the commons' (65) should favour less-beneficial symbionts (21–23). Using genetic correlations, GWAS, and analyses of nucleotide variation to study the genetic basis of mutualism variation in natural populations of rhizobia, we find little evidence for trade-offs between rhizobium and plant fitnesses at the phenotypic and genomic levels. Instead, we find strong alignment between host and symbiont fitnesses in nature. Indeed, the lack of points in the bottom right quadrants of figures correlating plant and rhizobium fitness proxies, particularly for DZA, suggests little evidence of competitively superior 'cheating' strains that benefit at the expense of their host (19).

Fitness alignment is ultimately governed by the degree to which mutualistic trait optima are shared among partners, as well as the degree of fitness feedbacks that enforce alignment between hosts and symbionts (8, 18, 21, 66). For example, legume host plants have autoregulation of nodulation to limit the formation of costly nodules once sufficient N levels are achieved (67). No such constraint exists for rhizobia; every nodule formed is expected to lead to greater potential fitness benefits (68). Thus, the optimum number of nodules for a plant is likely to be much lower than that for rhizobia. Indeed, the strongest evidence for host-symbiont conflict in our experiment comes from

regressing plant fitness proxies on nodule number, suggesting conflict over a host's total investment in nodulation along with previous studies (e.g., 4, 25, 26).

In one-to-many mutualisms, such as a single plant associating with a diverse population of rhizobia, fitness feedbacks operating on a per partner basis are predicted to align host and symbiont interests (21–24). Empirical data supports the presence of such feedbacks via **host sanctions** wherein nodules fixing more nitrogen receive more plant carbon (39, 69, 70), although the mechanisms are likely not perfectly accurate and have rarely been measured in the wild. Our observations of abundant alignment between host fitness proxies, nodule size, and rhizobium relative fitness are particularly striking given that neither host line was collected from the same sites as the rhizobia strains, and thus, are unlikely to naturally co-occur. Overall, our results suggest that fitness feedbacks operating to align host and symbiont fitness are present even in diverse communities irrespective of coevolutionary history.

Genomic resolution of conflict and alignment in symbiosis. Using genomics to dig below the phenotypic level can illuminate the genetic mechanisms that favour cooperation over conflict by quantifying the extent of symbiotic pleiotropy and suggesting patterns of historical selection acting on pleiotropic loci. We found little evidence for discordant pleiotropic variants underlying host and symbiont fitness, alongside evidence that selection has acted to maintain genetic variation within rhizobium genomes through time and/or space. An abundance of concordant pleiotropic variants, like we have found, may have resulted from physiological constraints that make discordant variants impossible (i.e., alleles associated with larger nodules and small host biomass are rare or non-existent), or because of correlational selection that removes discordant variants from the population altogether (71, 72). In addition to direction, the extent of symbiotic pleiotropy (i.e., the number of pleiotropic variants) can inform whether traits are likely to evolve together or independently. Fitness proxy pairs with abundant pleiotropy (e.g., nodule size vs. shoot biomass) suggest a highly polygenic basis governed by many small-effect pleiotropic variants; other proxies with little pleiotropy (e.g., nodule number vs. chlorophyll) are largely governed by different sets of genetic mechanisms, and are thus likely to evolve independently.

One challenge in mutualism research has been identifying patterns of selection acting on traits that determine fitness outcomes in natural populations, particularly for microbial symbioses (73, 74). Key studies examining patterns of molecular variation have found stabilizing or purifying selection acting on candidate genes associated with partner recognition or quality, rather than patterns suggesting rapid turnover of alleles underlying conflict (75–78). In a recent in silico GWAS (8), conflict over mutualistic trait optima between hosts and microbes tended to increase genetic variance due to

repeated sweeps, while alignment scenarios resulted in stabilizing selection and decreased genetic variance as trait optima were reached (8). In our empirical dataset, we found abundant fitness alignment combined with evidence that the genes underlying this alignment harbour particularly high nucleotide diversity and experience balancing selection.

Although the selective agents responsible for maintaining diversity remain to be determined, host genotypes potentially act as distinct selective environments for rhizobia, meaning that rhizobium alleles associated with higher rhizobium fitness and host benefits largely differ across host lines. Indeed, we found that concordant variants for both host and symbiont fitnesses were largely non-overlapping between the two host genotypes, a pattern also found in a recent experimental evolution study (28). In general, balancing selection acting on genes underlying fitness alignment provides solution to the **market paradox**, whereby host-driven selection for the 'best' symbiont genotype is expected to reduce overall symbiont diversity, yet quantitative variation in symbiotic outcomes remains (reviewed by 73, 74). If the genotype of the 'best' symbiont differs among host genotypes, then host selection should lead to greater, and not reduced, symbiont diversity.

Despite abundant concordance in our data at both levels, our paired approach did reveal several instances of discordance at the genomic level despite alignment at the phenotypic level (e.g., shoot biomass vs rhizobium relative fitness on DZA). Linkage disequilibrium (LD) could lead to discordant alleles being "packaged" into multilocus genotypes that show fitness alignment. Such LD results from multiple, non-mutually exclusive factors including epistatic interactions among individual variants that render discordant variants effectively neutral, and/or past selection favouring allelic combinations that increase both host and symbiont fitness (and disfavours discordant combinations; 79, 80). Overall, these findings highlight the polygenic nature of partner quality variation in nature — where the collective action of individual mutations, their additive and nonadditive effects (46), and a history of selection shapes the trait variation currently present in natural populations (81, 82).

Trade-offs among rhizobium fitness proxies and the rhizobium competition problem. Understanding how microbial symbioses, which are ubiquitous and important in nature, evolve (or coevolve) requires accurate estimates of symbiont fitness in ecologically realistic settings. Given our evidence at both the phenotypic and genomic levels for trade-offs among proxies, and because diverse strains of rhizobia co-occur in nature, relative fitness proxies should be used whenever possible (33). Nevertheless these proxies are not replacements for those measured in single-strain experiments because they cannot be used to assign individual genetic means for whole-plant measures of partner quality (plant biomass and seed number) to individual strains (e.g., 4, 34, 43, 46), necessitating that partner quality and rhizobium fitness be measured on separate individuals.

Comparing even the same fitness proxy across single- and multi-strain experiments (e.g., nodule number) may be more like comparing apples to oranges if the physiological or signalling processes governing the proxy differ between the two types of experiments. In single-strain experiments, plant autoregulation of nodulation (67) means that plants singly inoculated with a less-effective N-fixing strain will tend to form more total nodules than if they were inoculated with a more effective N-fixing strain (e.g., 4). In multi-strain experiments, however, nodule number for a particular strain depends on several factors: direct strain-strain competition (83), host signaling interactions (84), as well as any **partner choice**, i.e., overrepresentation of more beneficial strains in nodules (15, 41, 85). Therefore, the predominant processes that determine nodule number in single-strain experiments likely differ from those governing nodule number in multi-strain experiments, the latter being more representative of ecological conditions.

Together our results suggest that the genetic architectures underlying rhizobium fitness proxies, and their relationships, are host-dependent, and thus that their evolutionary trajectories are influenced not only by variation in the rhizobium's genome, but also by variation in host traits. For example, host genotypes could differ in their ability to exert sanctions or partner choice, quantitative traits known to vary in host populations (25, 27). At the genomic-level, distinct variants underlying rhizobium fitness pleiotropy on each host genotype suggests that the genetic mechanisms (i.e., genes, pathways, metabolic processes) governing the relationship between fitness proxies are largely non-overlapping when rhizobia interact with different hosts. Such host genotype-dependent shifts in the rhizobium genetic architecture of these symbiotic extended phenotypes is supported by transcriptomics studies of G x G interactions (33, 44) and GWAS revealing distinct sets of candidate genes in different host backgrounds (34, 46). Similar genetic variation exists in hosts (86) and undoubtedly interacts with the variants we identify here, thus, should be accounted for if we are to build a reliable G→P map.

ACKNOWLEDGMENTS. For their insightful feedback on the manuscript, we thank two anonymous reviewers and an associate editor. For greenhouse help, we thank Alex Riley, Hanna Lindgren, Cassandra Allsup, Laura Goralka, Mannix Burns, Brendan Epstein, and Amy-Marshall Colon, as well as the greenhouse staff, particularly Debbie Black. For feedback on the results and analyses, we thank Peter Tiffin, Brendan Epstein, and Robert Williamson. We acknowledge funding from the National Science Foundation (K.D. Heath: IOS-1645875 and NPGI-1401864; L.T. Burghardt: IOS-1724993 and IOS-1856744), a Carl R. Woese Institute for Genomic Biology postdoctoral fellowship to R.T. Batstone, and strain sequencing by Joint Genome Institute (CSP-1223795). This work was additionally supported by the USDA National Institute of Food and Agriculture Federal Appropriations under Project PEN04760 and Accession no. 1025611.

1. TF Mackay, EA Stone, JF Ayroles, The genetics of quantitative traits: challenges and prospects. *Nat. Rev. Genet.* **10**, 565–577 (2009).

2. M Pigliucci, Genotype–phenotype mapping and the end of the 'genes as blueprint' metaphor. *Philos. Transactions Royal Soc. B* **365**, 557–566 (2010).

3. B Lehner, Genotype to phenotype: lessons from model organisms for human genetics. *Nat. Rev. Genet.* **14**, 168–178 (2013).

4. KD Heath, Intergenomic epistasis and coevolutionary constraint in plants and rhizobia. *Evolution* **64**, 1446–1458 (2010).

5. ML Friesen, et al., Microbially mediated plant functional traits. *Annu. Rev. Ecol. Evol. Syst.* **42**, 23–46 (2011).

6. AL Gould, et al., Microbiome interactions shape host fitness. *Proc. Natl. Acad. Sci.* **115**, E11951–E11960 (2018).

7. J Lynch, E Hsiao, Microbiomes as sources of emergent host phenotypes. *Science* **365**, 1405–1409 (2019).
8. AM O'Brien, CN Jack, ML Friesen, ME Frederickson, Whose trait is it anyways? Coevolution of joint phenotypes and genetic architecture in mutualisms. *Proc. Royal Soc. B* **288**, 20202483 (2021).
9. RT Batstone, Genomes within genomes: nested symbiosis and its implications for plant evolution (*New Phytologist.* https://doi.org/10.1111/nph.17847) (2021).
10. SC Stearns, Trade-offs in life-history evolution. *Funct. Ecol.* **3**, 259–268 (1989).
11. N Barton, Pleiotropic models of quantitative variation. *Genetics* **124**, 773–782 (1990).
12. SP Otto, Two steps forward, one step back: the pleiotropic effects of favoured alleles. *Proc. Royal Soc. B* **271**, 705–714 (2004).
13. GP Wagner, J Zhang, The pleiotropic structure of the genotype–phenotype map: the evolvability of complex organisms. *Nat. Rev. Genet.* **12**, 204–213 (2011).
14. JL Bronstein, *Mutualism.* (Oxford University Press, USA), (2015).
15. ML Friesen, Widespread fitness alignment in the legume–rhizobium symbiosis. *New Phytol.* **194**, 1096–1111 (2012).
16. ET Kiers, WC Ratcliff, RF Denison, Single-strain inoculation may create spurious correlations between legume fitness and rhizobial fitness. *New Phytol.* **198**, 4–6 (2013).
17. ML Friesen, KD Heath, One hundred years of solitude: integrating single-strain inoculations with community perspectives in the legume–rhizobium symbiosis. *New Phytol.* **198**, 7–9 (2013).
18. ME Frederickson, Rethinking mutualism stability: cheaters and the evolution of sanctions. *The Q. Rev. Biol.* **88**, 269–295 (2013).
19. EI Jones, et al., Cheaters must prosper: reconciling theoretical and empirical perspectives on cheating in mutualism. *Ecol. Lett.* **18**, 1270–1284 (2015).
20. ME Frederickson, Mutualisms are not on the verge of breakdown. *Trends Ecol. & Evol.* **32**, 727–734 (2017).
21. JJ Bull, W Rice, Distinguishing mechanisms for the evolution of co-operation. *J. Theor. Biol.* **149**, 63–74 (1991).
22. S West, ET Kiers, I Pen, R Denison, Sanctions and mutualism stability: when should less beneficial mutualists be tolerated? *J. Evol. Biol.* **15**, 830–837 (2002).
23. SA West, ET Kiers, EL Simms, RF Denison, Sanctions and mutualism stability: why do rhizobia fix nitrogen? *Proc. Royal Soc. B* **269**, 685–694 (2002).
24. KR Foster, H Kokko, Cheating can stabilize cooperation in mutualisms. *Proc. Royal Soc. B* **273**, 2233–2239 (2006).
25. AK Simonsen, JR Stinchcombe, Standing genetic variation in host preference for mutualist microbial symbionts. *Proc. Royal Soc. B* **281**, 20142036 (2014).
26. SS Porter, EL Simms, Selection for cheating across disparate environments in the legume-rhizobium mutualism. *Ecol. Lett.* **17**, 1121–1129 (2014).
27. RT Batstone, EM Dutton, D Wang, M Yang, ME Frederickson, The evolution of symbiont preference traits in the model legume *Medicago truncatula*. *New Phytol.* **213**, 1850–1861 (2017).
28. RT Batstone, AM O'Brien, TL Harrison, ME Frederickson, Experimental evolution makes microbes more cooperative with their local host genotype. *Science* **370**, 476–478 (2020).
29. KW Quides, et al., Experimental evolution can enhance benefits of rhizobia to novel legume hosts. *Proc. Royal Soc. B* **288**, 20210812 (2021).
30. E Li, et al., Rapid evolution of bacterial mutualism in the plant rhizosphere. *Nat. Commun.* **12**, 1–13 (2021).
31. ML Friesen, et al., The ecological genomic basis of salinity adaptation in tunisian *Medicago truncatula*. *BMC Genomics* **15**, 1160 (2014).
32. CR Klinger, JA Lau, KD Heath, Ecological genomics of mutualism decline in nitrogen-fixing bacteria. *Proc. Royal Soc. B* **283**, 20152563 (2016).
33. LT Burghardt, et al., Select and resequence reveals relative fitness of bacteria in symbiotic and free-living environments. *Proc. Natl. Acad. Sci.* **115**, 2425–2430 (2018).
34. B Epstein, et al., Genome-wide association analyses in the model rhizobium *Ensifer meliloti*. *MSphere* **3** (2018).
35. SS Porter, J Faber-Hammond, AP Montoya, ML Friesen, C Sackos, Dynamic genomic architecture of mutualistic cooperation in a wild population of *Mesorhizobium*. *The ISME J.* **13**, 301–315 (2019).
36. TF Mackay, W Huang, Charting the genotype–phenotype map: lessons from the *Drosophila melanogaster* genetic reference panel. *Wiley Interdiscip. Rev. Dev. Biol.* **7**, e289 (2018).
37. JL Sachs, CJ Essenberg, MM Turcotte, New paradigms for the evolution of beneficial infections. *Trends Ecol. & Evol.* **26**, 202–209 (2011).
38. MA Parker, Plant fitness variation caused by different mutualist genotypes. *Ecology* **76**, 1525–1535 (1995).
39. ET Kiers, RA Rousseau, SA West, RF Denison, Host sanctions and the legume–rhizobium mutualism. *Nature* **425**, 78–81 (2003).
40. EL Simms, et al., An empirical test of partner choice mechanisms in a wild legume–rhizobium interaction. *Proc. Royal Soc. B* **273**, 77–81 (2006).
41. KD Heath, P Tiffin, Stabilizing mechanisms in a legume–rhizobium mutualism. *Evolution* **63**, 652–662 (2009).
42. CH Foyer, et al., Neglecting legumes has compromised human health and sustainable food production. *Nat. Plants* **2**, 1–10 (2016).
43. J Burdon, P Thrall, Spatial and temporal patterns in coevolving plant and pathogen associations. *The Am. Nat.* **153**, S15–S33 (1999).
44. KD Heath, PV Burke, JR Stinchcombe, Coevolutionary genetic variation in the legume-rhizobium transcriptome. *Mol. Ecol.* **21**, 4735–4747 (2012).
45. LT Burghardt, et al., Transcriptomic basis of genome by genome variation in a legume-rhizobia mutualism. *Mol. Ecol.* **26**, 6122–6135 (2017).
46. R Batstone, et al., The complex genetics of symbiotic extended phenotypes across multiple environments in a model mutualism (bioRχiv: https://www.biorxiv.org/content/10.1101/2021.08.03.454976) (2021).
47. KD Heath, P Tiffin, Context dependence in the coevolution of plant and rhizobial mutualists. *Proc. Royal Soc. B* **274**, 1905–1912 (2007).
48. RF Denison, Legume sanctions and the evolution of symbiotic cooperation by rhizobia. *The Am. Nat.* **156**, 567–576 (2000).
49. WC Ratcliff, K Underbakke, RF Denison, Measuring the fitness of symbiotic rhizobia. *Symbiosis* **55**, 85–90 (2011).
50. RF Denison, ET Kiers, Life histories of symbiotic rhizobia and mycorrhizal fungi. *Curr. Biol.* **21**, R775–R785 (2011).
51. LT Burghardt, B Epstein, P Tiffin, Legacy of prior host and soil selection on rhizobial fitness in planta. *Evolution* **73**, 2013–2023 (2019).
52. A Riley, M Grillo, B Epstein, P Tiffin, Partners in space: Discordant population structure between legume hosts and rhizobium symbionts in their native range (bioRχiv: https://www.biorxiv.org/content/10.1101/2021.06.30.449460) (2021).
53. TM Finan, et al., The complete sequence of the 1,683-kb psymb megaplasmid from the n2-fixing endosymbiont *Sinorhizobium meliloti*. *Proc. Natl. Acad. Sci.* **98**, 9889–9894 (2001).
54. F Galibert, et al., The composite genome of the legume symbiont *Sinorhizobium meliloti*. *Science* **293**, 668–672 (2001).
55. D Kessner, TL Turner, J Novembre, Maximum likelihood estimation of frequencies of known haplotypes from pooled sequence data. *Mol. Biol. Evol.* **30**, 1145–1158 (2013).
56. SR Searle, FM Speed, GA Milliken, Population marginal means in the linear model: an alternative to least squares means. *The Am. Stat.* **34**, 216–221 (1980).
57. RC Team, R foundation for statistical computing. *R: a language environment for statistical computing.* (2016).
58. X Zhou, M Stephens, Efficient multivariate linear mixed model algorithms for genome-wide association studies. *Nat. Methods* **11**, 407–409 (2014).
59. B Pfeifer, U Wittelsbürger, SE Ramos-Onsins, MJ Lercher, Popgenome: an efficient swiss army knife for population genomic analyses in r. *Mol. Biol. Evol.* **31**, 1929–1936 (2014).
60. F Tajima, Statistical method for testing the neutral mutation hypothesis by dna polymorphism. *Genetics* **123**, 585–595 (1989).
61. YX Fu, WH Li, Statistical tests of neutrality of mutations. *Genetics* **133**, 693–709 (1993).
62. D Bullock, D Anderson, Evaluation of the minolta spad-502 chlorophyll meter for nitrogen management in corn. *J. Plant Nutr.* **21**, 741–755 (1998).
63. JM Swiader, A Moore, Spad-chlorophyll response to nitrogen fertilization and evaluation of nitrogen status in dryland and irrigated pumpkins. *J. Plant Nutr.* **25**, 1089–1100 (2002).
64. DJ Weese, KD Heath, BT Dentinger, JA Lau, Long-term nitrogen addition causes the evolution of less-cooperative mutualists. *Evolution* **69**, 631–642 (2015).
65. G Hardin, The tragedy of the commons. *Science* **162**, 1243–1248 (1968).
66. JL Sachs, UG Mueller, TP Wilcox, JJ Bull, The evolution of cooperation. *The Q. Rev. Biol.* **79**, 135–160 (2004).
67. DE Reid, BJ Ferguson, S Hayashi, YH Lin, PM Gresshoff, Molecular mechanisms controlling legume autoregulation of nodulation. *Annals Bot.* **108**, 789–795 (2011).
68. JL Sachs, KW Quides, CE Wendlandt, Legumes versus rhizobia: a model for ongoing conflict in symbiosis. *New Phytol.* **219**, 1199–1206 (2018).
69. JU Regus, et al., Cell autonomous sanctions in legumes target ineffective rhizobia in nodules with mixed infections. *Am. J. Bot.* **104**, 1299–1312 (2017).
70. A Westhoek, et al., Conditional sanctioning in a legume–rhizobium mutualism. *Proc. Natl. Acad. Sci.* **118** (2021).
71. F Guillaume, MC Whitlock, Effects of migration on the genetic covariance matrix. *Evolution* **61**, 2398–2409 (2007).
72. JB Saltz, FC Hessel, MW Kelly, Trait correlations in the genomics era. *Trends Ecol. & Evol.* **32**, 279–290 (2017).
73. KD Heath, JR Stinchcombe, Explaining mutualism variation: a new evolutionary paradox? *Evolution* **68**, 309–317 (2014).
74. KS Stoy, AK Gibson, NM Gerardo, LT Morran, A need to consider the evolutionary genetics of host–symbiont mutualisms. *J. Evol. Biol.* **33**, 1656–1668 (2020).
75. JA Kimbrel, et al., Mutualistic co-evolution of typeIII effector genes in *Sinorhizobium fredii* and *Bradyrhizobium japonicum*. *PLoS Pathog.* **9**, e1003204 (2013).
76. MA Grillo, S De Mita, PV Burke, KL Solórzano-Lowell, KD Heath, Intrapopulation genomics in a model mutualist: Population structure and candidate symbiosis genes under selection in *Medicago truncatula*. *Evolution* **70**, 2704–2717 (2016).
77. JB Yoder, Understanding the coevolutionary dynamics of mutualism with population genomics. *Am. J. Bot.* **103**, 1742–1752 (2016).
78. B Epstein, P Tiffin, Comparative genomics reveals high rates of horizontal transfer and strong purifying selection on rhizobial symbiosis genes. *Proc. Royal Soc. B* **288**, 20201804 (2021).
79. B Sinervo, E Svensson, Correlational selection and the evolution of genomic architecture. *Heredity* **89**, 329–338 (2002).
80. AG Jones, SJ Arnold, R Bürger, Stability of the G-matrix in a population experiencing pleiotropic mutation, stabilizing selection, and genetic drift. *Evolution* **57**, 1747–1760 (2003).
81. JK Conner, DL Hartl, , et al., *A Primer of Ecological Genetics.* (Sinauer Associates Incorporated), (2004).
82. M Lynch, B Walsh, *Genetics and Analysis of Quantitative Traits.* (Sinauer, Sunderland, MA) Vol. 1, (1998).
83. D Dowling, W Broughton, Competition for nodulation of legumes. *Annu. Rev. Microbiol.* **40**, 131–157 (1986).
84. Q Wang, J Liu, H Zhu, Genetic and molecular mechanisms underlying symbiotic specificity in legume-rhizobium interactions. *Front. Plant Sci.* **9**, 313 (2018).
85. BS Younginger, ML Friesen, Connecting signals and benefits through partner choice in plant–microbe interactions. *FEMS Microbiol. Lett.* **366**, fnz217 (2019).
86. J Stanton-Geddes, et al., Candidate genes and genetic architecture of symbiotic and agronomic traits revealed by whole-genome, sequence-based association genetics in *Medicago truncatula*. *PloS*

One **8**, e65688 (2013).

Fig. 1. Interpreting fitness alignment and conflict at both the phenotypic and genomic levels. A) Phenotypic-level correlations of genotypic (strain) means for rhizobium fitness proxies (e.g., nodule number, nodule size, rhizobium relative fitness) or plant fitness proxies (shoot biomass or leaf chlorophyll A content) on the x- and y-axes, respectively. The positive (orange) or negative (grey) correlations represent fitness alignment or conflict, respectively. **B)** Genomic-level correlations of allelic effect sizes determined in GWAS for rhizobium fitness proxies or plant fitness proxies on the x- and y-axes, respectively. Points appearing in orange or grey quadrants represent pleiotropic variants with concordant (same-sign) or discordant (opposite sign) effects, respectively, on both host and symbiont. **C)** Plant and rhizobium fitness proxies measured, parentheses indicating the experiment type for which proxies were measured (SS = single-strain; MS = multi-strain). **D)** Rhizobium fitness proxies corresponding to the pink rhizobium strain. In single-strain experiments, nodule number indicates the number of reproductive cells released per plant, while nodule size indicates the number of reproductive cells released per nodule. In the multi-strain experiment, rhizobium relative fitness is a composite metric that combines competition among strains for nodule occupancy and for host resources once in nodules. Pink versus blue nodules are used here to illustrate two different rhizobium strains competing, however, a total of 89 strains were inoculated together onto plants in the multi-strain experiment.

Fig. 2. Trade-offs among rhizobia (*Ensifer meliloti*) fitness proxies prevail at both the phenotypic (left panels) and genomic-levels (right panels).
Phenotypic: Genetic correlations between pairwise fitness proxies measured on plant lines DZA (green, top rows) or A17 (pink, bottom rows), based on 89 *Ensifer meliloti* strains. Dots represent strain estimated marginal means for nodule number and nodule weight, both being measured in single-strain experiments, or medians for rhizobium relative fitness measured in multi-strain experiments. Numbers at top right of each correlation represent Pearson correlation coefficients, while asterisks represent significance: * = p < 0.05; ** = p < 0.01; *** = p < 0.001. **Genomic**: Dots represent allelic effect sizes (i.e., beta scores calculated in GEMMA), whereby effects were regressed for each pairwise fitness proxy combination. Dots falling along the diagonal (orange quadrants) represent variants with concordant (i.e., same-sign) effects, while those falling on the off-diagonal (grey quadrants) represent variants with discordant (i.e., opposite-sign) effects. Coloured dots represent variants that were significantly associated with one of the two fitness proxies, while black dots represent pleiotropic variants, i.e., significantly associated with both fitness proxies. Numbers outside of and percentages within parentheses at the top left and right of each plot represent the pleiotropic variant counts and the proportion of total significant variants, respectively, that are discordant (left, in grey) or concordant (right, in orange).

Fig. 3. Fitness alignment between rhizobia (*Ensifer meliloti*) and host (*Medicago truncatula*) prevails at both the phenotypic (left panels) and genomic (right panels) levels. Phenotypic: Genetic correlations between pairwise fitness proxies measured on plant lines DZA (green, top rows) or A17 (pink, bottom rows), based on 89 *Ensifer meliloti* strains. Dots represent strain estimated marginal means for shoot biomass as well as nodule number and nodule weight, all being measured in single-strain experiments, or medians for rhizobium relative fitness measured in multi-strain experiments. Numbers at top right of each correlation represent Pearson correlation coefficients, while asterisks represent significance: * = P < 0.05; ** = P < 0.01; *** = P < 0.001. **Genomic**: Dots represent allelic effect sizes (i.e., beta scores calculated in GEMMA), whereby effects were regressed for each pairwise fitness proxy combination. Dots falling along the diagonal (orange quadrants) represent variants with concordant (i.e., same-sign) effects, while those falling on the off-diagonal (grey quadrants) represent variants with discordant (i.e., opposite-sign) effects. Coloured dots represent variants that were significantly associated with one of the two fitness proxies, while black dots represent pleiotropic variants, i.e., significantly associated with both fitness proxies. Numbers outside of and percentages within parentheses at the top left and right of each plot represent the pleiotropic variant counts and the proportion of total significant variants, respectively, that are discordant (left, in grey) or concordant (right, in orange).

Fig. 4. Neutrality statistics show elevated values for genes underlying fitness alignment. Vertical lines represent the average values calculated for four separate statistics, grey for discordant and orange for concordant genes. Distributions represent the same statistics calculated for all genes containing significant variants based on GWAS. Dashed and solid lines represent non-significant (p > 0.1) and significant (p < 0.1) differences, respectively, between each focal gene category and all significant genes (i.e., distributions).