Strategies of tolerance reflected in two North American maple genomes

Susan L. McEvoy[1], U. Uzay Sezen[2], Alexander Trouern-Trend[1], Sean M. McMahon[2], Paul

G. Schaberg[3], Jie Yang[4], Jill L. Wegrzyn[1], Nathan G. Swenson[5]

[1] Department of Ecology and Evolutionary Biology, University of Connecticut, Storrs,

Connecticut, 06269

[2] Smithsonian Environmental Research Center, Edgewater, Maryland, 21037

[3] Forest Service, U.S. Department of Agriculture, Northern Research Station, Burlington, VT

05405

[4] CAS Key Laboratory of Tropical Forest Ecology, Xishuangbanna Tropical Botanical

Garden, Chinese Academy of Sciences, Mengla, 666303, Yunnan, China

[5] Department of Biological Sciences, University of Notre Dame, Notre Dame, IN, 46556

[#]Corresponding authors: Jill Wegryzn, email: jill.wegrzyn@uconn.edu, Nate Swenson, email

nswenson@nd.edu

Keywords: Acer saccharum, Acer negundo, genome, differential expression, nutrient stress,

calcium, aluminum, plasticity, abiotic stress, tolerance

Significance statement: The first chromosome-scale assemblies for North American members

of the *Acer* genus, sugar maple (*Acer saccharum*) and boxelder (*Acer negundo*), as well as

transcriptomic evaluation of abiotic stress response in *A. saccharum*. This integrated study

describes in-depth aspects contributing to each species' approach to tolerance and applies

current knowledge in many areas of plant genome biology with *Acer* physiology to help

convey the genomic complexities underlying tolerance in broadleaf tree species.

Abstract

Maples (the genus *Acer*) represent important and beloved forest, urban, and ornamental trees distributed throughout the Northern hemisphere. They exist in a diverse array of native ranges and distributions, across spectrums of tolerance or decline, and have varying levels of susceptibility to biotic and abiotic stress. Among *Acer* species, several stand out in their importance to economic interest. Here we report the first two chromosome-scale genomes for North American species, *Acer negundo* and *Acer saccharum*. Both assembled genomes contain scaffolds corresponding to 13 chromosomes, with *A. negundo* at a length of 442 Mb, N50 of 32 Mb and 30,491 genes, and *A. saccharum* at 626 Mb, N50 of 46 Mb, and 40,074 genes. No recent whole genome duplications were detected, though *A. saccharum* has local gene duplication and more recent bursts of transposable elements, as well as a large-scale translocation between two chromosomes. Genomic comparison revealed that *A. negundo* has a smaller genome with recent gene family evolution that is predominantly contracted and expansions that are potentially related to invasive tendencies and tolerance to abiotic stress. Examination of expression from RNA-Seq obtained from *A. saccharum* grown in long-term aluminum and calcium soil treatments at the Hubbard Brook Experimental Forest, provided insights into genes involved in aluminum stress response at the systemic level, as well as signs of compromised processes upon calcium deficiency, a condition contributing to maple decline.

Introduction

Acer saccharum (sugar maple) is a long-lived and dominant species in New England forests with a native range representing Eastern Canada, and the Northcentral and Northeastern United States. Widely known for its vibrant autumn hues, high quality timber, and as the preferred species for the production of maple syrup, *A. saccharum* also plays a key role in its native ecosystems, altering soil mineral content (Lucash et al., 2012), moisture levels (Emerman & Dawson, 1996), and mycorrhizae communities (Cong et al., 2015). *A. saccharum* provides food and shelter to many mammals (Godman et al., 1990), resident and migratory birds (Darveau et al., 1992), and over 300 species of caterpillar. Although seeds from this predominantly monoecious species are wind dispersed, the early flowers are an important pollen source for bees in late winter (Blitzer et al., 2016).

One of the most phylogenetically (~50 MY) and morphologically distinctive *Acer* species from *A.*

saccharum is *Acer negundo* (box elder). *A. negundo* is a short-lived tree that has the largest range

of all North American *Acer*. It is native to predominantly lower elevation regions of Canada, the

United States, and Mexico (Figure 1; Enquist et al., 2016; Maitner et al., 2018). This adaptable

pioneer species is often seen in disturbed sites and urban settings. *A. negundo* has soft wood, less

concentrated sugars for syrup, grows rapidly, and can tolerate low nutrient soils, moderate

salinity, and drought conditions. Its invasive status in large portions of Europe, South America,

and Asia are indicative of greater phenotypic plasticity (Lamarque et al., 2013). *A. negundo* has a

number of distinctive attributes including dioecy (Renner et al., 2007) and pinnately compound

leaves, mostly seen only in close relatives. Within such a significant genus, these two species

reflect a breadth of social and ecological diversity and importance that recommends better

understanding of their genetic distinctions.

Forest ecosystems of the Northeastern U.S. are facing significant changes in composition driven

by climate change (Rogers et al., 2017). "Maple decline" is a term referring to the loss of maple

populations, originally referring to *A. saccharum*, but now applicable to *A. platanoides* and *A.*

rubrum in the Northeast, and most recently, *A. macrophyllum* in the Northwest. Loss of *A.*

saccharum has been documented over the last century, beginning in the late 1950s, leading to the

first comprehensive, multidisciplinary study of this condition (Giese & Benjamin, 1964; Horsley

et al., 2002). Maple decline is characterized by crown dieback, reduction in overall health and

vigor, and a decrease in regeneration (Bishop et al., 2015). Episodic decline has increased in

recent decades (Oswald et al., 2018). Decline and crown dieback of dominant *A. saccharum*

provides a release for sympatric species such as *Fagus grandifolia* (American beech) which

displays a higher level of tolerance to soil conditions and foliar aluminum ratios leading to

shifting forest composition (Halman et al., 2015). Studies examining potential factors of maple

decline have largely agreed that modified soil conditions, largely due to acid deposition, are the

leading cause, compounded by additional climatic, pathogenic, and anthropogenic stressors (Bal

et al., 2015). Acidic soils rapidly leach the essential cations calcium, magnesium and potassium,

while mobilizing aluminum within the soil and contributing to more phytotoxic forms (Likens et

al., 1998; Likens & Lambert, 1998). Competition between aluminum and calcium at the roots

further decreases levels of available calcium within tissues, while increasing aluminum damages

plasma membranes, cell walls, DNA, and increases the burden of oxidative stress. Such nutrient

interactions and their broader consequences on physiology and ecology are studied at the Hubbard

Brook Experimental Forest (HBEF), a Long Term Ecological Research (LTER) site. It was here

that acid deposition was first discovered in North America (Likens & Bormann, 1974) and

continues to be studied through the Nutrient Perturbation (NuPert) program (Berger et al., 2001).

It provides a replicated high elevation natural ecosystem to examine current, future, and past soil

conditions, and has been the site of several key studies on native species, including *A. saccharum*,

A. balsamea, *F. grandifolia*, and *P. rubens*. At HBEF, no studies to-date on *A. saccharum* have

focused at the genomic level, where variation in gene expression or signs of adaptation among

gene families may be more immediately informative in these slow-growing organisms. For such

analysis, a high-quality, chromosomal-length genome is necessary to more accurately detect these

forms of variation.

Genomic resources necessary to guide *Acer* conservation are very limited. Only three genomes

exist to-date: *A. truncatum* (purpleblow maple), widely distributed across East Asia (Ma et al.,

2020), and the endangered *A. yangbiense* and *A. catalpifolium*, native to the Yuhan Province (J.

Yang et al., 2019) and West China (Yu et al., 2021), respectively. Here, were present the first two

North American *Acer* genomes, *A. saccharum* and *A. negundo*. With these chromosome-scale

references, we describe differences in genomic characteristics that may reflect their alternative

tolerance strategies. We conducted a differential expression study with stem tissue from *A.*

saccharum individuals from HBEF in order to identify genes that may be involved in aluminum

response and calcium deficiency. Identification of key processes in the expression study helped

to provide focus to the following analysis of comparative gene family dynamics. Together, these

approaches highlighted families associated with various abiotic stress responses, including those

that also have significant dynamics or novel isoforms in *A. saccharum* or *A. negundo* relative to

other broadleaf tree species. And it allowed investigation of the effects of calcium availability at

the molecular level, a significant factor associated with maple decline.

Figure 1. a) Native distributions of *A. saccharum* (blue) and *A. negundo* (orange) in North America. Leaves indicate location of individuals selected for the reference genomes; *A. saccharum* from the University of Maryland campus, and *A. negundo* from the Smithsonian Environmental Research Center. HBEF (Hubbard Brook Experimental Forest) is the location of the 9 individuals used for RNA-seq. b) All records of occurrence, including native, introduced, and unknown, per BIEN 4.2. Non-native occurrences are predominantly *A. negundo*.

Results

Genome size estimation and quality control

A. saccharum DNA sequencing resulted in 63 Gb of PacBio data with an N50 of 21 Kb (max

read length 94 Kb) and Illumina PE data totalling 225 Gb. *A. negundo* was similar with 61 Gb of

PacBio data, an N50 of 17 Kb (max read length 95 Kb), and 223 Gb of Illumina PE data.

Genome size estimation using short reads resulted in smaller than expected estimates (Contreras

& Shearer, 2018; Leitch et al., 2019) at 636 Mb and 319 Mb for *A. saccharum* and *A. negundo*,

respectively Figure S1. Using the short-read estimations of genome length, DNA sequence read

coverage was high, with long reads at 111x and 141x and short reads at 180x and 208x for *A.*

saccharum and *A. negundo*, respectively (Table S1). RNA sequencing of the reference

individuals resulted in 61 M reads for *A. saccharum* (92% mapped) and 62 M for *A. negundo*

(93% mapped). RNA sequencing of samples for the differential expression study resulted in 207

M reads with mapping rates that ranged from 82 to 92%.

Genome assembly

Testing of multiple assembly approaches found the FALCON/Purge Haplotigs assembly to be

the most contiguous and closest to *A. saccharum's* estimated size (Figure 2). Statistics for the set

of primary contigs from FALCON did not change much between the assembly, unzip, and

polishing stages (File S1). The final total length for *A. saccharum* settled at 970 Mb, with an N50

of 691 Kb and 2549 contigs. These last two statistics indicated better contiguity compared to the

other assemblers tested. The associated haplotype contigs rose from 109 Mb after assembly, to

320 Mb after unzipping, and dropped to 264 Mb after polishing. Removal of under-collapsed

haplotypes reduced the genome size to 668 Mb across 1210 contigs with an N50 of 951 Kb.

BUSCO (embryophyte) reported 94.8% complete, but with a somewhat high percentage of

duplication (11.7%).

The FALCON primary assembly for *A. negundo* was also consistent in size across pipeline

stages, with a total length matching estimates at 481 Mb across 1481 contigs with an N50 of 625

Kb. Removal of haplotype duplication from the primary assembly decreased the overall length to

442 Mb, number of contigs to 1063, and increased the N50 to 700Kb. The BUSCO score was

94.1% with only 5.8% duplication.

Figure 2. Results of assembly testing with *A. saccharum*, comparing fragmentation in terms of

total contigs versus assembly length. The dashed line represents the estimated genome size. Gray

dots are short-read assemblers, shown as highly fragmented. Blue dots are long-read tests of

assembly workflows. Canu refers to the use of reads error-corrected by the Canu pipeline. The

red dot is the selected draft assembly, and the green dot shows scaffolding results following Hi-

C. Detailed assembly statistics are available in File S1.

Hi-C scaffolding

The FALCON assembly was selected over Flye due to the substantially more contiguous

assembly it produced for *A. saccharum*, though it should be noted that the statistics for both

assemblers were comparable for *A. negundo*. Hi-C reads provided 65x coverage of the *A.*

saccharum genome. The final assembly was 626.33 Mb in 388 scaffolds with an N50 of 45.72

Mb and GC% of 35.7%. The 13 pseudo-chromosomes represented 97% of the genome length,

and BUSCO (embryophyte) scores were 97.7% complete with 3.0% duplicate, 0.7% fragmented,

and 1.6% missing.

A. negundo Hi-C reads provided 100x coverage and the FALCON assembly was used for

scaffolding due to potential mis-assembly in the Flye version (Figure S2). The final assembly

was 442.39 Mb in 108 scaffolds with an N50 of 32.30 Mb and GC% of 34.1%. The thirteen

pseudo-chromosomes represented 99.74% of the total length. BUSCO embryophyta scores were

97.4% complete with 5.6% duplicate, 0.9% fragmented, and 1.7% missing. (File S1). Earlier

work based on low coverage sequencing postulated higher GC content for both *Acer* (Staton et

al., 2015), and *A. negundo* in particular (Contreras & Shearer, 2018), but whole genome

sequencing supports their place at the lower range among angiosperm plants (Trávníček et al.,

2019).

Genome annotation

Annotations for *A. saccharum* resulted in 40,074 gene models of which 8,765 were monoexonics

verified by the presence of a protein domain and start and stop codons. Transcriptome

comparison based on 15,234 transcript loci supported 13,997 gene models. Functional annotation

was applied via similarity search or gene family assignment for 35,304 models. *A. negundo* had

30,491 genes, 5,558 of which were monoexonic, and 16,168 transcript loci supported 14,682 of

the *de novo* models. Functional annotations were determined for 27,077 of the models. (File S2).

A. saccharum repeat content was at 64.4% while *A. negundo* was 58.6%.

Whole genome duplication and Acer synteny

Categorization of putative paralogs revealed a higher percentage of each type in *A. saccharum*

relative to *A. negundo*. Plots of Ks distribution for WGD genes in syntenic regions show a single

clear peak at a Ks range consistent with the core eudicot WGT reported in other species using the

same pipeline (Figure 3a). *A. yangbiense* does not have a recent WGD, and when compared to *A.*

saccharum which had an additional small recent peak, further investigation identified small

blocks of collinearity, a minimum of five genes in palindromic or tandem arrangements. These

blocks are predominantly located on a few scaffolds and are not reflective of the general

distribution typical of WGD (File S3). Macrosynteny analysis found that *A. negundo* and *A.*

yangbiense are syntenic, while comparisons between each of these and *A. saccharum* revealed a large-scale translocation where two chromosomes from *A. negundo*, including the largest, are split with sections exchanged to form two different chromosomes in *A. saccharum* (Figure 3b, Figure S4).

Figure 3. Ks distribution for WGD synteny blocks with a summary of duplication types in (a) *A. negundo* and (b) *A. saccharum*. Abbreviations for categories of duplication: WGD, whole genome duplication; TD, tandem duplication; PD, proximal duplication; TRD, transposed duplication; DSD, dispersed duplication. (c) Circos plot of the thirteen chromosomes ordered largest to smallest for *A. negundo* (orange bars) and *A. saccharum* (blue bars) with distributions of gene density (green) and transposable element frequency (purple). Syntenic regions are linked in gray with darker shades to visually highlight larger recombinations.

Expression analysis of A. saccharum aluminum and calcium treatments

The final annotated genome for *A. saccharum* served as a reference for the expression study. In

total, there were 245 unique differentially expressed genes with 181 informatively described by

sequence similarity descriptors. Of those with no similarity match, four were completely novel

with no identifiable protein domain. Initial analysis produced six up and nine downregulated

genes comparing the aluminum to calcium treatments, and the other pairwise comparisons had

similarly small totals. Clustering of the expression results showed season had a strong effect

(Figure S3), so the analysis was repeated for each season individually to remove this variable

from treatment comparisons. For brevity, abbreviations are used according to the following

definitions: All, across seasons; Fa, fall; Sp, spring; Al, aluminium; Ca, calcium; and Un,

unamended. FaAl to FaCa had 26 upregulated and 17 down, FaAl to FaUn had 7 up and 12

down, and FaUn to FaCa had 28 up and 27 down. SpAl to SpCa had 39 up and 19 down, SpAl to

SpUn had the greatest number with 33 up and 41 down, and SpUn to SpCa had 28 up and 11

down (Figure 4, File S4).

There were only two instances where the same gene was found in both separate seasonal

analyses. Transcription factor-like E2FE was upregulated in both unamended and aluminum

treatments in fall, spring, and all. It had the most isoforms expressed, and was also the most

highly upregulated for fall (8-fold). E2FE represses endoreduplication, reducing this type of

growth in response to stress (Hendrix et al., 2018). The second instance was disease resistance

protein At4g27190-like, expressed primarily in SpAl, but also FaUn and FaCa. Another

At4g27190 isoform was upregulated in SpCa and SpAl, but downregulated in SpUn.

GO enrichment results for fall showed sugar and carbohydrate transmembrane transporter

activity upregulated in the FaUn compared to FaCa (Table S3). Several sugar transporters,

SWEETs and ERD6-like, were seen in both seasons, though more often in fall. There were two

SWEET15 (tandem duplicates), one upregulated in FaUn to FaCa, and SWEET15-like

upregulated in SpUn relative to both SpCa and SpAl (7.6-fold). SWEET2a was upregulated in

FaAl to FaCa. Three different ERD6 were upregulated in FaUn with a fourth in SpAl.

Triterpenoid biosynthetic process was also enriched FaUn to FaCa, supported by two

downregulated beta-amyrin synthase genes, one of which was significantly downregulated even

further in FaAl to FaUn (5.4-fold). Other strongly differentiated genes included accelerated cell

death 6-like (ACD6) upregulated in FaCa (5-fold) relative to FaAl. It is associated with flavin-

containing monooxygenase 1 (FMO1), the most highly upregulated DEG in FaUn (22-fold) and

FaCa, both relative to FaAl.

Figure 4. a) Differential expression study design showing number of samples collected in fall and spring from treatment plots at the Hubbard Brook Experimental Forest, Nutrient Perturbation study. b) Differentially expressed genes (up and downregulated) for each treatment and season comparison. Charts display both significance and relative expression denoted as log-fold change. Dotted lines indicate thresholds of significance (0.1 p-adjusted, 1.5 $\log_2$ fold change).

In spring, DEGs upregulated in SpUn compared to SpAl were enriched in processes related to biotic and abiotic stress, including gene ontology terms for defense response, and acid, oxygen-containing, and antimicrobial response. There were six disease resistance genes, all largely in SpUn and SpCa compared to two in fall. Also present in large numbers, were serine/threonine kinases, including LRR receptor-like, with seven out of twelve of these in the FaUn. Heat shock proteins were also common, though more equally split between the two seasons and shifted toward unamended or aluminum. There were two copies of ITN1, involved in salicylic acid signaling, one of which was the most upregulated DEG in SpCa to SpAl (23.6-fold). In addition to direct stress response, there was an interesting increase of expression in three Holliday junction resolvases, and a lamin-like gene, which was the second highest DEG in these comparisons (19.7-fold). It can play a role in nuclear membrane integrity, chromatin organization, and gene expression (Hu et al., 2019).

Metal tolerance via ligation, sequestration, and transport
In SpAl, upregulation of both metallothionein-like 3 (6.6-fold), and aluminum-activated malate transporter (ALMT) 10 (4.5-fold) was observed. An EH domain-containing gene (4.6-fold) is involved in endocytosis, vesicle transport, and signal transduction (Naslavsky & Caplan, 2005), and several cytoskeletal related genes were seen such as kinesin in fall (6.8-fold), and myosin-6 in spring (8-fold).

In addition to the sugar transporters, ABC transporters were present as multiple isoforms in FaUn, and one was the most upregulated DEG in FaUn compared to FaCa (13-fold). Two were downregulated in FaAl compared to FaCa, but still elevated in FaUn, and were from family C which contains pumps for glutathione S-conjugates, which have been shown to remove cadmium in *Arabidopsis* (Tommasini et al., 1998). Additional transport included upregulation of ATPases

across seasons, cyclic and mechanosensitive ion channels in spring and fall, respectively, and a

K+/H antiporter upregulated in FaAl.

Calcium dependent proteins

Three calcium-transporting ATPases were upregulated in either FaCa or FaUn along with an

unspecified plasma membrane ATPase. Ca-based external signal relay mechanisms upregulated

in SpCa included a cyclic nucleotide-gated ion channel, G-type lectin S-receptor like

serine/threonine-protein kinases, and a glutamate receptor (5.5-fold) (X.-L. Sun et al., 2013;

Toyota et al., 2018). A calcium-dependent protein kinase (4.5-fold) was upregulated in FaUn.

Calmodulin-binding proteins (60d at 5-fold) were up-regulated in both SpAl and SpCa, and are

associated with salicylic acid synthesis and immunity in *Arabidopsis* (L.-S. Li et al., 2021).

Reactive oxygen species response

Antioxidant and redox genes included glutathione peroxidase, upregulated in FaUn, and

glutathione S-transferase, upregulated in both FaCa and SpAl. Two cytosolic sulfotransferases

and six thioredoxin genes were upregulated in SpAl. Other redox related genes include a highly

expressed epoxide hydrolase (9-fold) and a carotenoid cleavage dioxygenase (4.6-fold) in FaAl;

and peroxidase, aldehyde dehydrogenase (6.7-fold), and germin in SpAl. Ascorbate-dependant

oxidoreductase SRG1 was upregulated in SpUn, cytochrome P450 71D11 (a monooxygenase) in

SpAl, and zinc finger (C2H2 type) in SpUn.

Cell wall and membrane integrity

MDIS1-interacting receptor kinase, upregulated in FaUn relative to FaAl, and expansin, FaUn to

FaCa, are associated with the cell wall. Two ADP-ribosylation factor GTPase-activating, one of

which is the most highly upregulated gene in spring (19.9 fold-change SpAl to SpCa), assists

with cell signaling by recruiting cargo-sorting coat proteins at the membrane and regulating lipid

composition in support of development and defense (Donaldson & Jackson, 2011).

Hormone crosstalk

Auxin was indicated by three WAT genes (seen in each season, greatest at 7-fold in SpAl to

SpCa) and indole-3-acetic acid (highest of AllAl to AllUn at 9.7-fold). Two ethylene synthesis

genes were present: 1-aminocyclopropane-1-carboxylate oxidase, FaAl to FaCa, and

methylthioribose kinase, SpAl to SpCa. Ent-kaurenoic acid oxidase is related to brassinosteroid

homeostasis and gibberellin biosynthesis (Helliwell et al., 2001) and was present in SpAl to

SpUn (7-fold), and obtusifoliol 14α-demethylase, which mediates brassinosteroid synthesis, was

upregulated SpUn to SpCa (Xia et al., 2015). Jasmonic acid activity was upregulated in FaCa,

negative regulation of cytokinin was up in AllAl to AllUn, and ABA-induced HVA22 which

inhibits gibberellin and is possibly involved in vesicular traffic, was in SpAl to SpCa (8-fold) as

well as AllAl to AllCa.

Gene family evolution with expression study integration

By leveraging 22 high-quality plant genomes, gene family dynamics between *A. negundo* and *A.*

saccharum revealed distinct characteristics. Comparisons among the plant proteomes resulted in

20,234 orthogroups with a mean size of 26.4 genes. Of these, 4,262 were shared by all species,

and 79 were single-copy. 88.5% of 603,640 genes were contained in orthogroups, and 0.7% were

species-specific. All species had at least 80% of their genes contained in orthogroups with the

exception of *Ginkgo biloba, Nymphaea colorata*, and *Oryza sativa* (Table S4).

All three *Acer* shared 11,156 orthogroups. *A. negundo* and *A. saccharum* had the largest overlap,

A. saccharum and *A. yangbiense* had the second largest overlap, and *A. saccharum* had the most

unshared groups. (Figure 5b). Comparing *Acer* against the other woody angiosperms (*B.*

pendula, C. papaya, C. clementina, C. sinensis, E. grandis, J. hindsii, J. regia, P. vera, P.

trichocarpa, P. persica, Q. lobata, Q. robur, T. grandis), 728 *Acer* orthogroups were expanded,

14 were contracted, 1992 were novel, and *Acer* was estimated to be missing from 202

orthogroups. To clarify, these may not be fully absent, but didn't have representation in

orthogroups including those that were more lineage specific. Comparing the Sapindales to the

other trees resulted in 340 expanded, 4 contracted, 2788 novel, and 160 missing. Dynamics in

common between the two *Acer* included 270 expanded groups, 1 contracted, 0 novel, and 247

missing (File S5).

Acer families expanded among the woody angiosperms were enriched for RNA modification,

DNA replication strand elongation, and processes of organic cyclic compound, cellular aromatic

compound, and heterocycle metabolisms. Novel genes were highly enriched for cell periphery

localization and marginally for sesquiterpenoid and triterpenoid biosynthesis (Figure 5a, Table

S5). When focusing on absent gene families, none were found to be missing exclusively in

Sapindales, which includes the two *Citrus* species and *Pistacia vera*. A total of two families were

absent in all *Acer*, phosphatidylcholine transfer protein-like and cellulose synthase interactive 3,

but were present in all other species.

Several interesting gene families more novel to *Acer* overlap with the HBEF DEGs. There are

twenty orthogroups associated with disease resistance At4g27190, two seen as DEGs, and the

specific families containing these DEGs are larger for *A. saccharum* with one novel to the *Acer*.

Two additional non-DEG families are rapidly expanding in *A. saccharum*, with one of these also

expanding in *A. negundo*, but both contracting in *A. yangbiense*. In fall, another more novel DEG

ACD6-like belongs to a family with limited species membership (Sapindales and *V. vinifera*).

Compared to *Arabidopsis* ACD6, which has two 3-repeating ankyrin domains, the *A. saccharum*

ACD6-like had varying ankyrin positions, as do other members of this family. A third example,

seen only in spring, acetyl-coenzyme A synthetase (ACS) is a member of a novel *Acer* family

consisting of 1 *A. negundo*, 6 *A. saccharum*, and 18 *A. yangbiense*. Comparison with *A. negundo*

and ACS isoforms reveals the *A. saccharum* gene contains a longer ACL domain, with only

~67% query coverage and ~43% percent identity to *A. negundo* which is much more similar to

other ACS (87-90% length; ~88% identity).

Within the broader set of species, *A. saccharum* gene families were characterized by more

expansion, with 1827 expanded, 18 contracted, 127 novel, and 511 absent, and more rapidly

expanding, with 99 compared to 18 contracting. *A. negundo* had 1068 expanded, 23 contracted,

89 novel, and 558 absent orthogroups. Rapidly contracting families were greater in this species

with 52 compared to 26 rapidly expanding (Figure 5, File S5, File S6).

A. saccharum gene families

Functional enrichment of *A. saccharum* in the full species comparison revealed that expansions

were processes of ncRNA metabolism, RNA modification, organic cyclic compound

metabolism, heterocycle metabolism, and intracellular membrane-bound organelle localization

(Table S5). Almost half of the *A. saccharum* families had limited annotation information, due to

either missing descriptors or uncharacterized protein matches. Relative to *Acer*, *A. saccharum's*

expanded families are enriched in a larger list of stress response associated functions that are

fairly specific, including water deprivation, hypoxia, salinity, heat, cold, xenobiotic, nematode,

karrikin, acid chemical, and hormone (File S13). Other significant processes include regulation

of indolebutyric acid stimulus (auxin family), RNA splicing, chloroplast RNA processing,

phospholipid translocation, brassinosteroid homeostasis, lignin synthesis, microsporogenesis,

phenylpropanoid biosynthesis, cadmium ion transmembrane transport, cyclin-dependent

serine/threonine kinase, and calcium-transporting ATPase. Rapidly expanding families are

associated with various biotic and abiotic responses, such as fungal, salt stress, and xenobiotic

response (File S6). Interesting genes include patatin-like 2, involved in membrane repair via

removal of lipids modified by oxidation (W.-Y. Yang et al., 2012) and ALP1 negative regulation

of polychrome group chromatin silencing (Liang et al., 2015). Those that overlap with HBEF

DEGs include disease resistance At4g27190 and DSC1, FMO1, rapidly expanding SRG1, and

rapidly contracting disease resistance At1g50180.

Compared to other *Acer*, contracted families are enriched for pollen wall assembly, extracellular

matrix assembly and organization, chlorophyll binding, NADH dehydrogenase (ubiquinone)

activity, DNA-directed 5'-3' RNA polymerase activity, programmed cell death, and myb-like

transcription factors (File S11, File S12). Genes absent in *A. saccharum* but present in all other

species total 28 (File S15), including red chlorophyll catabolite reductase (ACD2) and S-

adenosyl-L-homocysteine hydrolase (HOG1), which is necessary to hydrolyze the by-product of

the activity of S-adenosyl-L-methionine-dependent methyltransferase, and one of these was also

absent. The apparent absence of HOG1 requires further investigation as mutants display a

number of problematic phenotypes and variants display association with fiber length in *P.*

tomentosa (Du et al., 2014).

A. negundo gene families

In the broad comparison of species, *A. negundo* expanded families were enriched in RNA

modification, microgametogenesis, and metabolic processes of nucleobase-containing

compound, organic cyclic compound, heterocycle, and cellular aromatic compound (Table S5).

The small number of rapidly expanding families were by far mostly uncharacterized proteins or

missing sequence similarity descriptions with only four out of 26 genes having a description,

including glutathione-S-transferase, two disease-resistance proteins, At4g27190-like and

At5g66900, a receptor-like 12, and additional functional descriptors such as E3 ubiquitin-protein

ligase, LRR receptor-like ser/thr kinase, and more (File S6). Relative to other *Acer*, *A. negundo's*

expanded families are enriched in a short list of specific stress response including UV, UV-B,

radiation, bacterium, cadmium, metal ion, drug, chemical, and osmotic stress (File S9, File S10).

Other processes include proanthocyanidin biosynthesis, lignin synthesis via cinnamyl-alcohol

and sinapyl-alcohol dehydrogenase, starch metabolism and glucan catabolism, error-prone

translesion synthesis, and other DNA damage response and repair. Processes related to

reproduction were present, especially pollen development. For example, decreased size exclusion

limit 1 (DSE1, aka aluminum tolerant 2 (ALT2)) is a transcription factor that regulates the size

of molecules that can travel through plasmodesmata, as channel aperture is not static, changing

in response to stress and decreasing during embryo development (Xu et al., 2012). DSE1 is

single copy in all species with expansions only in *A. negundo*, *Q. rober*, and *T. grandis*.

Contractions relative to other *Acer* include transcription by RNA polymerase III, chloroplast

RNA processing, and lignin biosynthetic processes (File S7, File S8). Rapidly contracted

families were enriched in quercetin 3-O- and 7-O-glucosyltransferase activity (Table S6).

Examples of contracted families include 7-ethoxycoumarin O-deethylase, which metabolizes a

wide range of xenobiotics (Robineau et al., 1998), and disease resistance-like protein DSC1, both

rapidly expanding families in *A. saccharum* (File S6). There are 23 genes absent in *A. negundo*

that are present in all other species (File S14) . Several of these are curious, as they appear to be

required components of important processes, such as AUGMIN subunits 2 and 7 that help form a

complex that plays a role in spindle microtubule generation (Tian & Kong, 2019), and cell

division cycle 45-like, which is required for meiosis in *Arabidopsis* (Stevens et al., 2004). The

absence of formamidopyrimidine-DNA glycosylase is also interesting as it is involved in base

excision repair of DNA damage, a notable area of specific enrichments described below.

Figure 5. a) Gene ontology enrichments for *Acer* (all three species combined), *A. negundo*, and *A. saccharum*. Abbreviations for gene family dynamics: E, expanded; N, novel; RC, rapidly contracting. b) Total gene families, shared and unique, among the *Acer*. c) Reconstructed gene tree showing contracted gene families in red and expanded in green.

Discussion

Acer saccharum and *A. negundo* differ in species distribution and tolerance to abiotic stressors, and their genomes reveal support for their contrasting life histories. *A. saccharum* has slow growth until release of canopy coverage, doesn't achieve reproductive maturity until 40 years of age, and is able to live 400 years. It requires high nutrient soils, prefers mesic environments, and is tolerant of cold, but not salinity. Its range crosses a more narrow latitudinal gradient, but a

stronger elevational one, where *A. negundo* tends to be more limited. While both genomes are

small in size and diploid, *A. saccharum* is 42% larger, contains twice as many transposable

elements, and 38% more gene duplications, many of which are very recent. Its gene families tend

to be larger, more diverged, and undergoing rapid expansion compared to *A. yangbiense* or *A.*

negundo.

In contrast, *A. negundo* grows quickly, is able to reproduce after only five years, and has a

relatively short lifespan of 60 years. It is tolerant of a range of conditions and is widespread

throughout North America. *A. negundo* is considered an aggressive invasive in Europe, South

Africa, parts of Asia and North America, and it rapidly dominates disturbed habitats (CABI,

2021). It is sexually dimorphic and both native and invasive conspecifics are highly plastic in

growth, leaf phenology, and physiological traits such as maximum assimilation rate, nitrogen

content and photosynthetic efficiency (Dawson & Ehleringer, 1993; Lamarque et al., 2015).

Relative to native heterospecifics, invasive populations of *A. negundo* maximized growth in high

light and nutrient conditions, with dramatically reduced growth and survivorship in deficient

environments (Porté et al., 2011).

The *A. negundo* reference genome is a small diploid with high heterozygosity from the native

species range. Macrosynteny between *A. yangbiense* and *A. negundo* is largely collinear and

without rearrangement as shown in the two *A. saccharum* chromosomes (Figure S4). Invasive

plant species are often associated with smaller genomes and *A. negundo* is characterized by

contracting gene families. Traits such as fast growth rate, germination time, stomatal

responsiveness, and dispersal ability are cell size or division rate dependent (Pyšek et al., 2018).

The greater surface area to volume ratio of small cells, derived from small genomes, reduces the

metabolic and signaling requirements, but does not preclude additional growth or activation, thus

extending the range of expression, or plasticity, for a wider set of traits (Roddy et al., 2019; Suda

et al., 2015). The adaptive potential conferred by polyploidism can also be leveraged for success

of invasives, but polyploidism has not been documented in the native or introduced range.

If larger genome size presupposes increased metabolic, transport, and nutrient demands (Pellicer

et al., 2018) perhaps *A. saccharum's* susceptibility to nutrient deficiencies, may be in part due to

the extra burden of a larger genome and its particular enrichments. The decrease in soil nutrient

availability in its native range over the past decades is at odds with resources necessary to

tolerate stressors brought on by a changing environment, and previous adaptations are now

maladaptive. From the foundational differences between *A. negundo* and *A. saccharum* in

morphological, physiological, and genomic characteristics, we turn to the integration of gene

family dynamics and expression data from a study on nutrient stress to further elucidate the

contrasting strategies of tolerance seen in these species.

Integration of Acer saccharum expression studies

Integration of gene expression sampling with gene family dynamics highlighted the genetic

factors that may influence maple decline and adaptation. HBEF NuPert *A. saccharum* saplings

grown in native ecological and environmental conditions within replicated long-term soil

treatments provided an opportunity to examine differences in gene expression related to soil

acidity, calcium deficiency, and increased aluminum availability (Table S7). Calcium

amendment of NuPert plots was designed to represent levels from 50 years ago, before extensive

acid deposition in the HBEF range led to the deficient conditions seen in unamended control

plots (Berger et al., 2001). Aluminum amended plots further decrease calcium availability via

competition for uptake between the two cations. They represent more extreme calcium depletion

and an increase in aluminum availability that is observed with soil acidity. Examination of trees

on the landscape in these conditions allowed us to study the effects of long-term nutrient stress in

conjunction with other life-history processes. Seasonal differences between the differentially

expressed genes were striking, revealing the effect of calcium levels on processes occurring at

the time of sampling.

Signaling and metabolic processes are sensitive to lower calcium levels

Calcium is the common limiting element underlying maple decline, and trees improve long-term

with the addition of calcium (Moore & Ouimet, 2021). Calcium transport and binding is well

studied for its ability to activate and regulate primary components of stress response systems,

including internalization of external signals, modulation of cytosolic levels for transient

signaling, and initiation and perpetuation of ROS responses, hormone modulation, and pathways

of secondary metabolism (Estravis-Barcala et al., 2020). In physiological studies conducted at

HBEF, calcium amendments resulted in a 50% increase of calcium in foliar tissues of *A.*

saccharum, while non-significantly decreasing aluminum concentrations. Trees in calcium plots

devoted more carbon to growth than storage, were better able to flush after a late spring frost,

produced more flowers, and increased seed germination (Halman et al., 2013).

Calcium amendment facilitated upregulated gene expression relative to unamended and

aluminum treatments in areas related to calcium signaling and senescence activities. In fall, a

gene involved in leaf senescence was highly expressed in calcium treatments, and the relative

paucity of other differential expression in this season made it all the more interesting. ACD6 is

important to both natural and stress response senescence, shown to regulate rosette biomass,

chlorophyll degradation, and leaf nitrogen and carbon levels in *A. thaliana* (Jasinski et al., 2021).

It is also a factor in nitrogen translocation from leaves to seeds before cell death. ACD6

stimulates calcium signaling in *A. thaliana* and could potentially be an ion channel itself (Zhu et

al., 2021), and several calcium dependent signaling and calmodulin binding genes were

upregulated in the fall calcium and unamended plots, including the critical calcium-dependent

protein kinase 1. The ACD6 isoform expressed here is limited to Sapindales and quite diverged

from the *A. thaliana* form. Interestingly, a second senescence gene that is expressed in *Acer*

rubrum (Z. Chen et al., 2019) is absent from the *A. saccharum* genome. Red chlorophyll

catabolite reductase (RCCR aka ACD2) is a non-green chlorophyll degradation, and while the

exact role of RCCR is unclear (Jockusch & Kräutler, 2020), it appears possible that like many

senescence related genes, it provides additional fitness advantages (Mach et al., 2001). Timing,

duration, and intensity of gene expression are all important during senescence activities leading

up to cell death, including adjustment of signaling, growth, metabolism, and nutrient

remobilization. The susceptibility of ACD6 to calcium deficiency could potentially create

complications in fall that could affect spring growth, reproduction, and ability to respond to

stress.

Throughout *A. saccharum's* native range, when calcium is deficient, additional nutrient

imbalances, such as magnesium, phosphorus, potassium, and nitrogen deficiency, as well as high

aluminum and manganese concentrations, climatic effects, and biotic stressors such as

defoliation, create compounding stress (Bal et al., 2015; Long et al., 2019). Samples from

unamended plots show an increase in stress response perhaps reflective of calcium deficiency

combined with external stressors. Signs include enrichments of sugar and other transporters in

fall samples (Julius et al., 2017; Miao et al., 2017). Spring samples reported upregulation of

several putative disease resistance genes, heat shock proteins, leucine-rich repeats proteins and

kinases, and signs of oxidative stress.

Oxidative stress response is high in aluminum amendment conditions

Reactive oxygen species (ROS) are generated as a result of both stress and normal cellular

processes, and reactive oxygen can take different forms, with varying toxicities. ROS is reduced

by a variety of potential "scavengers" to prevent accumulation of levels that are damaging to

lipids, proteins, DNA, and RNA, leading to cell death. ROS production and redox activity vary

according to the primary processes of each cellular compartment (i.e., metabolism in

mitochondria), and also in response to different stressors or stress combinations (i.e., drought and

heat), so it is thought that cells develop complex ROS signatures or hotspots that influence signal

transduction and metabolic regulation in nuanced ways (Castro et al., 2021; Choudhury et al.,

2017). Aluminum increases peroxidation of lipids in membranes and produces H_2O_2 which can

participate in retrograde signaling, regulating expression of additional genes (Castro et al., 2021).

Aluminum derived oxidative stress causes inhibition of cell growth and is an early signal of

aluminum toxicity (Yamamoto et al., 2003).

Antioxidants and redox genes are commonly observed as a response to ROS stress and a

combination of these were observed in different seasons and treatments, though more were found

in spring, as fits with the trend of increased overall activity in that season. Antioxidants,

including thioredoxins, peroxidases, and a germin-like, were all upregulated in the aluminum

treatment, implying significant stress response activation. Glutathione system members, which

can also act as chelators, are more broadly seen in both unamended and aluminum. A number of

antioxidant DEGs were also from expanded families, in particular the significantly expanded

thioredoxins H3 and YLS8 and SRG1 ([Figure 6](#)). The antioxidant DEGs from this study can be

compared with a number of other transcriptomic studies on metal stress in trees where

glutathione S-transferase, peroxidases, and thioredoxins were in common. For example,

cadmium accumulators *Salix integra* and *Populus x canadensis* 'Neva' both expressed

superoxide dismutase with glutathione pathway genes and peroxidases, respectively (X. Li et al.,

2021; Shi et al., 2016). Nickel stress in resistant versus susceptible genotypes of *Betula*

papyrifera found glutathione S-transferase and thioredoxins upregulated in resistant trees

(Theriault et al., 2016). *A. rubrum*, a nickel avoider, had root-level expression of superoxide

dismutase, but downregulated glutathione S-transferase (Nkongolo et al., 2018) similar to one of

the *A. saccharum* DEGs. In comparisons of aluminum treated *Citrus* root tissue expression,

peroxidases and germin-like proteins were upregulated. It is interesting that superoxide

dismutase was not significantly expressed in *A. saccharum* given that it is from an expanded

gene family and seen in several other species. In physiological studies conducted at HBEF,

Halman et al. (2013) found *A. saccharum* were under increasing oxidative stress in unamended

and aluminum plots. Aluminum treated trees had elevated levels of glutathione reductase activity

as measured by spectrophotometric assay. Ascorbate peroxidase activity was also elevated, but

this particular peroxidase was not observed in our gene expression study.

Figure 6. Orthogroup sizes for aluminum tolerance gene families are presented by species. Families were selected for inclusion based on documented aluminum tolerance and/or presence in the HBEF RNA-Seq differential expression results. Color represents the proportion of gene membership per species, with darker purple equating to more contracted families relative to the

median, and dark green indicating expansion. (H) Family contains HBEF differentially expressed

gene; (E) Expanded in *A. saccharum*; (C) Contracting; (M) Missing; (N) Novel; (*) Rapidly

expanding; Categorization of tolerance is according to literature describing aluminum stress

phenotypes. The undetermined category contains species where tolerance to aluminum or acidic

soils has not been reported. [1]*B. pendula* is undetermined due to high variability in tolerance by

genotype.

Aluminum and metal response is minimal in aluminum amendment conditions

Deciphering aluminum response mechanisms can be complicated as aluminum affects several

targets within cells, and resistance and tolerance mechanisms can vary widely, even within

species. Accumulation of aluminum occurs primarily in the root, binding to pectin in the cell

wall, increasing its rigidity and decreasing permeability. Throughout the plant, aluminum can

disrupt plasma membrane potential and membrane-bound solute transporters (Kar et al., 2021).

Damage to proteins and DNA reduces enzymatic activity, and oxidative stress and hormone

signaling from DNA damage can lead to growth inhibition (Sade et al., 2016).

Studies from locations such as Vermont, the Adirondack Mountains, and the Allegheny Plateau,

have linked foliar aluminum and calcium levels (Long et al., 2019; Oswald et al., 2018; Sullivan

et al., 2013) and stem aluminum levels to branch dieback as seen in maple decline (Mohamed et

al., 1997). At HBEF, previous studies examined physiological effects of aluminum with a multi-

tissue study on mature *A. saccharum.* Using inductively coupled plasma atomic emission

spectroscopy, Halman et al. (2015) found foliar levels of aluminum did not vary much by

treatment and all were well below thresholds of toxicity. Calcium treatments caused

concentrations of aluminum to drop slightly, while aluminum treatments caused calcium to drop

more strongly in dominant than non-dominant trees. The main effects noted were moderate root

damage and foliar antioxidant activity (Halman et al., 2013). Within our stem tissue expression

analysis, signs of metal response were not clear, but we can detect levels of general stress

response expression between the unamended and aluminum treatments.

Genes currently known to be associated with aluminum resistance or tolerance were not

expressed in the stem tissue samples, but upregulated genes related to metal transport and

sequestration are potentially involved in aluminum remediation (Hasan et al., 2017;

Hasanuzzaman et al., 2017). Metallothionein 3 was highly expressed in the combined season

aluminum comparisons, though unevenly distributed throughout the replicates. Similar to other

metallothioneins, it works by ROS scavenging and metal ion homeostasis via chelation. (Hasan

et al., 2017). In salt-tolerant *Oryza sativa*, this gene responds to cadmium, salinity, and oxidative

stress (Mekawy et al., 2018).

Most of *A. saccharum's* aluminum resistance gene families are modestly sized relative to other

species examined, with the exception of PEP carboxylase and a large family of mixed MATEs,

with members likely similar to homologs studied in *P. trichocarpa* and *Arabidopsis* (N. Li et al.,

2017). Overexpression of either of these increases efflux of organic acids at the root (Begum et

al., 2009). *A.saccharum* had slightly more expansion in resistance families than the other *Acer*,

while *A. negundo* had the least. Given the limited number of DEGs indicating metal toxicity, and

the non-toxic aluminum tissue concentrations observed in physiological studies, it seems likely

resistance mechanisms exist at the root in *A. saccharum*. Comparing aluminum/metal tolerance

gene families, *A. yangbiense* had the fewest expanded and there are no reports on aluminum

tolerance for this species. *A. negundo* and *A. saccharum* were not notably different with the

exception of the metallothionein 3 family, which was rapidly expanding, much larger in *A.

saccharum* relative to other broadleaf tree species, and observed in the expression results for

both seasons. Expanded resistance and tolerance families were seen more frequently in high-

tolerance species such as *P. vera*, *P. trichocarpa*, *T. grandis*, *Q. lobata*, and *E. grandis*, though

the mix of families varies ([Figure 6](#); Sork et al., 2016; Tuskan et al., 2006; Zeng et al., 2019;

Zhao et al., 2019). Gene family size is only a potential factor in response and it does not

necessarily serve as a proxy for expression. The redox families selected for comparison are

primarily limited to those containing DEGs, and in this case there is good overlap between gene

family size and expressed redox genes. Many were larger in *A. saccharum*, but *A. negundo* was

also enriched for a different set of redox families.

Indications of growth reduction and DNA damage in response to aluminum

Strong upregulation of transcription factor E2FE was notable as it is one of only two genes seen

in both seasons. E2FE represses endoreduplication (Liu et al., 2019), and while

endoreduplication can be induced by certain types of stress such as drought and salinity (Lang &

Schnittger, 2020), cadmium toxicity was observed to reduce endoreduplication and cell division

in *A. thaliana* leaves (Hendrix et al., 2018). With cadmium, pathways that delay cell cycle

progression due to oxidative DNA damage were upregulated along with E2FE/DEL1. Aluminum

is known to cause double-stranded breaks which requires homology-dependent recombination

repair (P. Chen et al., 2019). In *A. saccharum,* DNA damage might be indicated by upregulation

of several Holliday junction resolvases which help repair double stranded breaks using

homologous recombination (Bauknecht & Kobbe, 2014). Patterns of expression point toward

oxidative DNA damage from aluminum or another source, inducing repression of

endoreduplication similar to cadmium.

The hormone Indole-3-acetic acid (IAA)-amido synthetase was the most highly upregulated gene

in the combined season comparison of aluminum treatment to unamended. The gene family for

this DEG is novel to *Acer* and significantly expanded in *A. saccharum*. IAA is the most abundant

auxin in plants, and IAA-amido synthetase helps with its homeostasis by leading to IAA-amino

acid conjugates (Ludwig-Müller, 2011). Increased expression of IAA-amido synthetase produces

highly reduced growth forms, altered architecture, and shortened roots (C. Ding et al., 2021; M.

Sun et al., 2020). In *O. sativa*, over-expression of IAA-amido synthetase resulted in improved

drought tolerance (Zhang et al., 2009), fungal pathogen response (X. Ding et al., 2008; Domingo

et al., 2009) and altered growth. Park et al (2007) observed reduced growth in *A. thaliana* from

overexpression of an IAA-conjugating gene (GH3) that was highly linked to biotic and abiotic

stress response. They proposed that modulation of auxin is an adaptive strategy that trades

reduced growth for enhanced stress resistance.

Stress response in Acer saccharum and calcium deficiency versus aluminum toxicity

Replicated samples of *A. saccharum* reveal complex expression changes indicating multiple

forms of stress response in aluminum treatments and to a lesser degree in unamended plots.

Differentially expressed genes, often related to signaling, redox, hormones, and growth

reduction, are seen in aluminum or unamended treatments, but not as extensively in calcium,

where there is enhancement of other processes, such as senescence and calcium signaling.

Ecosystem and environmental conditions at the time of sampling likely had interacting effects

and contributed to complexity of expression, but trends by season and treatment were still

evident. With this data, we provide preliminary molecular support to the many studies

correlating maple decline with calcium-poor, acidic soils. Given the minimal expression of metal

tolerance genes and lower level of foliar aluminum seen in dominant sugar maple at HBEF,

along with large amount of variability in aluminum response sometimes seen between

conspecific genotypes in other species, further transcriptomic studies of root tissue would be

beneficial for a better understanding of aluminum resistance in *A. saccharum*.

Herbivory, DNA damage, and light harvesting enrichment in Acer negundo

Gene family expansion in *A. negundo* was specifically limited to UV, bacteria, metal, cadmium,

chemical, and osmotic response rather than the more extensive set seen in *A. saccharum*. A high

proportion of proanthocyanidin synthesis related gene families were expanded, which are

precursors to condensed tannins that protect against herbivory, bacteria and fungal pathogens,

and encroaching of neighboring plants (He et al., 2008). All four families were relatively novel

to *A. negundo* with absence in all but one or two other species. Proanthocyanidins also have

antioxidant and radical scavenging functions, and have been compared with lignins in terms of

pathogen defense mechanisms (Stafford, 1988).

Both *Acer* are enriched in gene families with DNA damage recognition and repair functionality,

but *A. negundo* has a greater number, and they are largely categorized as either translesion

synthesis or meiosis-related repair. Translesion synthesis is often activated by UV damage. It is

an error-prone repair mechanism designed to quickly eliminate lesions at the replication fork that

might lead to stalling and double-stranded breaks (Sakamoto, 2019). *A. negundo* has three

families in this category expanded relative to *Acer,* two of which are significantly expanded

among the full comparison of species. (Nisa et al., 2019). Among *A. negundo's* DNA damage

related enrichment are secondary enrichments that include regulation of leaf, seed, flower

development, and post-embryonic structures, response to abiotic and biotic stress, and growth

such as cell division and endoreduplication ([Figure 7](#)). Too much DNA damage can inhibit

growth and development or initiate programmed cell death. Perhaps a more robust response to

DNA damage is a factor in *A. negundo's* plasticity of growth and tolerance.

In addition to UV-related DNA repair, there is another potential adaptation related to

photooxidative stress. The early light induced protein 1 (ELIP) gene family stood out as

significantly expanded in *A. negundo* relative to all other species, with twice as many genes

compared to other *Acer*. Research has focused on ELIP's protection against high light (Huang et

al., 2012), but it also regulates seed germination in response to environmental factors (Rizza et

al., 2011), and is also expressed in roots, and in response to some abiotic stressors including

nitrogen in *Populus* (Luo et al., 2015), indicating additional protective functions. Capacity for

regulation of photosynthesis and DNA repair mechanisms, within a largely streamlined genome,

could represent investments with large-scale effects, altering growth according to resources or

stress as described for *A. negundo* in various nutrient and light conditions.

Figure 7. *A. negundo* gene families with ontology related to DNA damage and repair, and

secondary enrichments categorized by color. Circles with multiple colors indicate multiple

ontology assignments. Lines indicated known or predicted interactions, or other association via

text-mining, co-expression, or protein homology.

Conclusion

In this study we present new chromosomal-length genomes for *A. saccharum* and *A. negundo* and demonstrate the greater repeat, paralog, GC content, and expanding character of *A. saccharum* versus *A. negundo* which was reduced and rapidly contracting. We conducted a stem tissue expression study of *A. saccharum* subjected to long-term aluminum and calcium treatments. Larger trends related to the abiotic stress response and calcium deficiency were revealed against a complex background of expression. Gene families related to aluminum stress resistance and tolerance were compared showing moderate expansion of families in *A. saccharum*. Gene family dynamics combined with physiological and expression results, point toward potential aluminum resistance mechanisms in effect at the root. Broader gene family comparisons revealed expansion in traits associated with invasiveness and stress tolerance in *A. negundo*. Release of these genomes and a complementary expression analysis of trees in the HBEF NuPert study shed further light on mechanisms of tolerance and potential adaptations of increasing importance due to climate change.

Methods

Sequencing

Leaf material for *A. saccharum* was collected from the University of Maryland College Park Campus (GPS location 38°59'16.0"N 76°56'32.8"W). *A. negundo* was sourced from the ForestGEO Forest Dynamics plot (tag # 91603) located in Smithsonian Environmental Research Center, Edgewater, MD. Leaves were dark treated for at least one day prior to collection and shipped to Amplicon Express Inc. (Pullman, WA, USA) for DNA extraction. Both species were sequenced with Pacific BioSciences Sequel v2.1 using a total of fourteen SMRTcells, with the SMRTbell® Express Template Prep Kit v1.0, insert size 40 Kb, library size selection of 66 Kb for *A. saccharum* and 55 Kb for *A. negundo*. The resulting short read sequencing consisted of two lanes of Illumina HiSeq 2500 150PE with insert sizes ranging from 500 to 600 bp (Genomics Resource Center, University of Maryland School of Medicine Institute for Genome Sciences, Baltimore, MD, USA). For Hi-C, DNA leaf material was collected from the same individuals and extracted with the Phase Genomics Proximo Hi-C Plant kit (Phase Genomics,

Seattle, WA, USA). Sequencing consisted of two lanes of mid output Illumina NextSeq 500

75PE with an average of 150 M reads per species (Center for Genome Innovation (CGI) at the

University of Connecticut, Storrs, CT, USA). For the RNA-Seq used in annotation, leaf samples

were collected from one individual per species at two years of age grown at the Michigan State

University greenhouse. Sequencing consisted of Illumina HiSeq at 100PE.

Stem tissue from *A. saccharum* was sampled from nine trees across two seasons from the

Nutrient Perturbation study plots at HBEF (North Woodstock, NH, USA). A single sapling (dbh

< 21 cm) was selected from three plots in each of the three treatments (calcium, control,

aluminum). A total of 18 libraries (nine in Oct 2017 (Fall); nine in May 2018 (Spring)) were

sampled (Table S2). Tissue was sampled from five years of growth measured by spring wood

internodes. Sections were frozen in liquid nitrogen in the field and RNA were extracted after

grinding tissues in liquid nitrogen. Extractions were run on Agilent Bioanalyzer Tapestation for

quantification and RNA integrity. Two samples were unsuccessful (one aluminum and one

calcium) from the spring sampling. Libraries were prepared using Illumina TruSeq Stranded

mRNA and sequenced with Illumina NextSeq 500, 150PE (CGI).

Draft genome assembly

In advance of assembly, genome size was estimated with Jellyfish v2.2.6 (21-mers) and

GenomeScope (Vurture et al., 2017) using the trimmed Illumina short-reads. The *A. saccharum*

reads were test assembled using Illumina short-read, raw and corrected long-reads, and a hybrid

of both. Draft assemblies were evaluated in relation to expected genome size, contiguity (N50

and number of contigs), conserved seed plant orthologs, and genomic/transcriptomic read

alignments.

The best draft assemblies leveraged the deep PacBio sequencing (*A. saccharum* 103x, *A.*

negundo 141x) and prioritized assembling repetitive regions of the genome and resolving the

heterozygosity, found in both species. During this phase of the process, two draft assemblies of

comparable results were used to investigate scaffolding potential. One was created with

FALCON (pb-assembly v0.0.6) a *de novo,* diploid-aware assembler for PacBio (Chin et al.,

2016), and the other was done using Canu (v1.6) error-corrected PacBio reads as input (Koren et

al., 2017) to Flye v2.3.7, an efficient haploid assembler that leverages repeat graphs with read

alignment techniques to resolve areas of repetition (Kolmogorov, 2019).

Alternate heterozygous contigs (haplotigs) were separated from the primary assemblies using

Purge Haplotigs v1.0.4 (Roach et al., 2018). To determine coverage, PacBio reads were aligned

to the primary assemblies with Minimap2 v2.15-r911-dirty (H. Li, 2018). RepeatModeler v1.0.8

was used to create the repeat libraries for this analysis.

Hi-C scaffolding

Long-range scaffolding of the FALCON/Purge Haplotigs assembly with Hi-C reads followed

processes recommended for the following suite of tools (*Genome Assembly Cookbook*, 2019).

HiC reads were aligned with BWA mem -5SP and PCR duplicates removed with samblaster

v0.1.24 (Faust & Hall, 2014). Scripts from the Juicer v1.5.6 (Durand, Shamim, et al., 2016)

pipeline were modified to identify the Sau3AI restriction enzyme. The resulting Hi-C alignment

file was provided to 3D-DNA v180419 (Dudchenko et al., 2017) for scaffolding, leveraging

different parameters for each species according to the differing draft assembly characteristics. In

particular, --diploid was added to *A. saccharum* to address remaining under-collapsed

heterozygosity. JuiceBox was used to visualize Hi-C mapping against each scaffold created by

the different parameter tests to visually detect which incorporated the majority of the contigs into

the expected 13 pseudo-chromosomes(Dudchenko et al., 2018; Durand, Robinson, et al., 2016).

Genome annotation

RepeatModeler v2.01 (Flynn et al., 2020) and RepeatMasker v4.0.6 (Smit et al., 2013) were used

to softmask the assembly. Trimmed RNA reads were aligned to the assembly with Hisat2 v2.1.0

(Kim et al., 2015). GenomeThreader v1.7.1 (Gremme et al., 2005) was used to align protein

sequences (derived from *de novo* transcriptome assembly; -gcmincoverage 80, -dpminexonlen 20

-startcodon -finalstopcodon -introncutout). Structural gene prediction was executed with

BRAKER2 v2.0.5 (Hoff et al., 2019). The process converted RNA-Seq alignments to exon

support in GeneMark-ET v4.32 (Lomsadze et al., 2014), and combined this output with protein

alignments for two rounds of training with AUGUSTUS v3.2.3 (Camacho et al., 2009; Stanke et

al., 2006, 2008) to predict coding regions in the genome. Extensive filtering was performed on

the predicted gene space using gFACS v1.1 (Caballero & Wegrzyn, 2019). Evaluation of

structural annotations were conducted with BUSCO and the PLAZA CoreGF rosids database

v2.5 (Van Bel et al., 2019; Veeckman et al., 2016). Transcriptomic alignments were used to

identify where they fully overlapped BRAKER-based models or provided additional support to

those that did not pass previous filtering criteria. Transcriptome assemblies were conducted *de*

novo with Trinity v2.20 (Grabherr et al., 2011). EnTAP v0.8.0 (Hart et al., 2020) was used to

frame-select, functionally annotate, and identify potential contaminants for filtering, including

bacteria, archaea, opisthokonta, rhizaria, and fungi. The resulting translated protein sequences

were clustered with USEARCH v9.0.2132 (Edgar, 2010) at an alignment identity of 0.90.

Transcriptomic alignments created with GMAP v2019-06-10 (Wu & Watanabe, 2005) and

gFACs were compared against the genome annotation using GffCompare v0.11.5 (Pertea, 2018).

Functional annotation was performed using EnTAP v0.9.1, a pipeline that integrates both

similarity search and other annotation sources including gene family (eggNOG), protein domains

(Pfam), gene ontology, and pathway assignment (KEGG). The following public databases were

included: NCBI RefSeq Complete, EMBL-EBI UniProt, and *Arabidopsis* (TAIR11) .

Comparative genomics

The translated gene space of 22 plant species were used for gene family analysis. *Acer* included

A. saccharum, *A. negundo*, and *A. yangbiense*. The remaining species were selected from high

quality public annotations. Fourteen broadleaf trees (*Betula pendula, Carica papaya*, *Citrus*

clementina and *sinensis*, *Eucalyptus grandis*, *Juglans hindsii* and *regia*, *Pistacia vera*, *Populus*

trichocarpa, *Prunus persica*, *Quercus lobata* and *robur*, and *Tectona grandis*), plus one woody

angiosperm, *Vitis vinifera*, were included, along with one gymnosperm, *Ginkgo biloba*. *Oryza*

sativa Kitaake was the representative monocot, along with *Amborella trichopoda* and *Nymphaea*

colorata, representing other more basal lineages, and *Arabidopsis thaliana*, being the primary

plant model system. OrthoFinder v2.3.7 (Emms & Kelly, 2018) was used to generate

orthogroups. Resulting gene counts per orthogroup for *A. negundo*, *A. saccharum*, and the mean

of the combined three *Acer* were each compared to the mean of other species to identify

potentially expanded, contracted, missing, and novel gene families. The initial delineation of

expansion and contraction was set at 2-fold above the standard deviation. Full absence was

verified with alignment of the *Arabidopsis* protein sequence against the assembly. CAFE v5

(Mendes et al., 2020) was used to identify rapidly evolving gene families. Values from the

newick species tree produced by OrthoFinder were multiplied by 100 to prevent issues with

rounding in CAFE, and the tree was made ultrametric using OrthoFinder. The poisson root

frequency distribution was run three times on gene families filtered by size to ensure

convergence of a lambda value representing birth and death rate. The selected lambda value was

then used to run the large family set. Functional enrichment of the resulting families was

obtained from gProfiler v:e99_eg46_p14_f929183 (Raudvere et al., 2019) using the annotation

of the representative (longest) gene when aligned to *Vitis vinifera.* This well annotated woody

angiosperm represents an ideal source having no recent WGD (Tang et al., 2008).

Whole genome duplication and synteny analysis

Characterization of putative paralogs, including whole genome duplication was done as

previously described (Qiao et al., 2019) using DupGen_finder to separate whole genome,

tandem, proximal, transposed, and dispersed duplicates. Categorization can be helpful in

speculating on the origin of duplication such as transposable element activity, localized

replication error, larger segmental translocations, or ploidy events. The whole genome duplicate

frequency distribution was plotted by Ks value for analysis of peaks. Microsynteny of the small

peak of recent supposed whole genome duplication seen in *A. saccharum* was further analyzed

with MCScanX-based collinearity scripts (Nowell et al., 2018), as well as overall macrosynteny

between the three *Acer.*

Differential expression analysis

HBEF NuPert plots were used as a basis for this study as they were designed to reflect acidity

and calcium levels over time as described by Berger et al. 2001. They consist of 12 *A. saccharum*

dominant plots near reference watershed 6, with four receiving annual $AlCl_2$ treatments 12 times

from 1995 to 2015, and $CaCl_2$ treatments were applied to 4 other plots for 4 years, followed by

applications of slow-release wollastonite in 1999 and 2015 (Table S7). Three samples were

collected from aluminum, calcium, and unamended control plots as described in the Sequencing

section. Trimmed reads for each of the sixteen successful HBEF-sourced libraries were aligned

to the *A. saccharum* reference genome with Hisat2 v2.1.0, and read counts were extracted with

htseq-count v0.11.2. The R Bioconductor package, DESeq2 v1.26.0 (Love et al., 2014), was used

for the expression analysis with the calcium as the control in pairwise comparisons of

unamended to calcium and aluminum to unamended, representing increasing levels of aluminum,

and then aluminum to calcium, contrasting the extremes. P-adjusted values greater than 0.1 were

filtered. Pairwise comparisons, specific to each season (fall and spring) as well as combined

resulted in a total of nine sets. Gene Ontology (GO) enrichment was conducted with g:Profiler

(database version e99_eg46_p14_f929183) using alignments of differentially expressed protein

models to both *Vitis vinifera* (Phytozome v12.1) and *Arabidopsis thaliana* (TAIR11) as baseline

annotations.

A. thaliana was used as a representative model for pathway analysis in the Genemania

application for Cytoscape v3.8.0. and *V. vinifera* (NCBI taxon ID:29760) (Smoot et al., 2011)

was used similarly with STRINGDB v.11 in Cytoscape v2.7.1. Differentially expressed proteins

for each pairwise comparison were used to visualize the fold-change values in context of the

supported pathways. Networks were constructed with a confidence score of 0.4 and 20 maximal

additional interactions and additional networks for protein models reported to be responsive to

calcium deficiency and aluminum biotoxicity were imported and merged.

Data Availability

Acknowledgements

NGS was funded by the National Science Foundation (NSF DEB-2029997; NSF EF-1638488).

Hi-C library preparation and sequencing was funded by the Ronald Bamford Fund Endowment

for Ecology and Evolutionary Biology to the Department of Ecology and Evolutionary Biology,

University of Connecticut. Support for the HBEF RNA-Seq was provided by the University of

Connecticut Center for Biological Risk.

The authors would like to thank the Institute for Systems Genomics (ISG) at UConn for

sequencing support, the Microbial Analysis Resources and Services (MARS) within the Center

for Open Research Resources & Equipment (COR^{2}E) at UConn for RNA extraction support, the

Computational Biology Core for HPC services, and the Hubbard Brook Experimental Forest for

access to the NuPert plots. Thanks to Laura Figueroa Corona for assistance with the range map,

and Sumaira Zaman, Alison Scott, and Nasim Rahmatpour for sharing computational approaches

and scripts. Thank you to Dr. Yaowu Yuan for advice on both data analysis and the manuscript.

The *A. negundo* individual sequenced is a part of the Smithsonian Environment Research Center

Forest Dynamics Plot, which is part of the Smithsonian's ForestGEO network. We thank

ForestGEO and Director Stuart Davies for encouraging tree genome research in the network and

for facilitating and funding (NSF DEB-1046113; NSF DEB-1545761) as well as previous

working group discussions on the topic. The authors declare no conflict of interest.

Author Contributions

SM, NS, JW, PS, and UU conceived and designed various aspects of the study; JW and NS

coordinated and managed the study; SM, UU, NS, PS, AT performed the sampling and

experiments; AT performed transcriptomic analysis; SM performed the assembly, annotation,

and related comparative genomic and expression analysis under the advisement of JW; SM, JW

wrote the manuscript; UU, NS, and AT contributed sections. All authors read and approved the

final manuscript.

References

944 Bal, T. L., Storer, A. J., Jurgensen, M. F., Doskey, P. V., & Amacher, M. C. (2015). Nutrient

946   stress predisposes and contributes to sugar maple dieback across its northern range: A

947   review. *Forestry: An International Journal of Forest Research*, *88*(1), 64–83.

948   https://doi.org/10.1093/forestry/cpu051

949 Bauknecht, M., & Kobbe, D. (2014). AtGEN1 and AtSEND1, Two Paralogs in Arabidopsis,

950   Possess Holliday Junction Resolvase Activity. *Plant Physiology*, *166*(1), 202–216.

951   https://doi.org/10.1104/pp.114.237834

952 Berger, T. W., Eagar, C., Likens, G. E., & Stingeder, G. (2001). Effects of calcium and

953   aluminum chloride additions on foliar and throughfall chemistry in sugar maples. *Forest*

954   *Ecology and Management*, *149*(1), 75–90. https://doi.org/10.1016/S0378-1127(00)00546-

955   6

956 Bishop, D. A., Beier, C. M., Pederson, N., Lawrence, G. B., Stella, J. C., & Sullivan, T. J.

957   (2015). Regional growth decline of sugar maple (Acer saccharum) and its potential

958   causes. *Ecosphere*, *6*(10), art179. https://doi.org/10.1890/ES15-00260.1

959 Blitzer, E. J., Gibbs, J., Park, M. G., & Danforth, B. N. (2016). Pollination services for apple are

960   dependent on diverse wild bee communities. *Agriculture, Ecosystems & Environment*,

961   *221*, 1–7. https://doi.org/10.1016/j.agee.2016.01.004

962 Caballero, M., & Wegrzyn, J. (2019). gFACs: Gene Filtering, Analysis, and Conversion to Unify

963   Genome Annotations Across Alignment and Gene Prediction Frameworks. *Genomics,*

964   *Proteomics & Bioinformatics*, *17*(3), 305–310. https://doi.org/10.1016/j.gpb.2019.04.002

965 CABI. (2021). *Acer Negundo*. Invasive Species Compendium. https://www.cabi.org/ISC

966 Camacho, C., Coulouris, G., Avagyan, V., Ma, N., Papadopoulos, J., Bealer, K., & Madden, T.

967   L. (2009). BLAST+: Architecture and applications. *BMC Bioinformatics*, *10*(1), 421.

https://doi.org/10.1186/1471-2105-10-421

Castro, B., Citterico, M., Kimura, S., Stevens, D. M., Wrzaczek, M., & Coaker, G. (2021).
Stress-induced reactive oxygen species compartmentalization, perception and signalling.
Nature Plants, *7*(4), 403–412. https://doi.org/10.1038/s41477-021-00887-0

Chen, P., Sjogren, C. A., Larsen, P. B., & Schnittger, A. (2019). A multi-level response to DNA
damage induced by aluminium. *The Plant Journal*, *98*(3), 479–491.
https://doi.org/10.1111/tpj.14231

Chen, Z., Lu, X., Xuan, Y., Tang, F., Wang, J., Shi, D., Fu, S., & Ren, J. (2019). Transcriptome
analysis based on a combination of sequencing platforms provides insights into leaf
pigmentation in Acer rubrum. *BMC Plant Biology*, *19*(1), 240.
https://doi.org/10.1186/s12870-019-1850-7

Chin, C.-S., Peluso, P., Sedlazeck, F. J., Nattestad, M., Concepcion, G. T., Clum, A., Dunn, C.,
O'Malley, R., Figueroa-Balderas, R., Morales-Cruz, A., Cramer, G. R., Delledonne, M.,
Luo, C., Ecker, J. R., Cantu, D., Rank, D. R., & Schatz, M. C. (2016). Phased diploid
genome assembly with single-molecule real-time sequencing. *Nature Methods*, *13*(12),
1050–1054. https://doi.org/10.1038/nmeth.4035

Choudhury, F. K., Rivero, R. M., Blumwald, E., & Mittler, R. (2017). Reactive oxygen species,
abiotic stress and stress combination. *The Plant Journal*, *90*(5), 856–867.
https://doi.org/10.1111/tpj.13299

Cong, J., Yang, Y., Liu, X., Lu, H., Liu, X., Zhou, J., Li, D., Yin, H., Ding, J., & Zhang, Y.
(2015). Analyses of soil microbial community compositions and functional genes reveal
potential consequences of natural forest succession. *Scientific Reports*, *5*(1), 1–11.
https://doi.org/10.1038/srep10007

Contreras, R. N., & Shearer, K. (2018). Genome Size, Ploidy, and Base Composition of Wild

and Cultivated Acer. *Journal of the American Society for Horticultural Science*, *143*(6),

470–485. https://doi.org/10.21273/JASHS04541-18

Darveau, M., DesGranges, J. L., & Gauthier, G. (1992). Habitat Use by Three Breeding

Insectivorous Birds in Declining Maple Forests. *The Condor*, *94*(1), 72–82.

https://doi.org/10.2307/1368797

Dawson, T. E., & Ehleringer, J. R. (1993). Gender-Specific Physiology, Carbon Isotope

Discrimination, and Habitat Distribution in Boxelder, Acer Negundo. *Ecology*, *74*(3),

798–815. https://doi.org/10.2307/1940807

Ding, C., Lin, X., Zuo, Y., Yu, Z., Baerson, S. R., Pan, Z., Zeng, R., & Song, Y. (2021).

Transcription factor OsbZIP49 controls tiller angle and plant architecture through the

induction of indole-3-acetic acid-amido synthetases in rice. *The Plant Journal*, *108*(5),

1346–1364. https://doi.org/10.1111/tpj.15515

Ding, X., Cao, Y., Huang, L., Zhao, J., Xu, C., Li, X., & Wang, S. (2008). Activation of the

Indole-3-Acetic Acid–Amido Synthetase GH3-8 Suppresses Expansin Expression and

Promotes Salicylate- and Jasmonate-Independent Basal Immunity in Rice. *The Plant*

Cell, *20*(1), 228–240. https://doi.org/10.1105/tpc.107.055657

Domingo, C., Andrés, F., Tharreau, D., Iglesias, D. J., & Talón, M. (2009). Constitutive

Expression of OsGH3.1 Reduces Auxin Content and Enhances Defense Response and

Resistance to a Fungal Pathogen in Rice. *Molecular Plant-Microbe Interactions®*, *22*(2),

201–210. https://doi.org/10.1094/MPMI-22-2-0201

Donaldson, J. G., & Jackson, C. L. (2011). ARF family G proteins and their regulators: Roles in

membrane transport, development and disease. *Nature Reviews. Molecular Cell Biology*,

12(6), 362–375. https://doi.org/10.1038/nrm3117

Du, Q., Wang, L., Zhou, D., Yang, H., Gong, C., Pan, W., & Zhang, D. (2014). Allelic variation

within the S-adenosyl-L-homocysteine hydrolase gene family is associated with wood

properties in Chinese white poplar (Populus tomentosa). *BMC Genetics*, *15*(1), S4.

https://doi.org/10.1186/1471-2156-15-S1-S4

Dudchenko, O., Batra, S. S., Omer, A. D., Nyquist, S. K., Hoeger, M., Durand, N. C., Shamim,

M. S., Machol, I., Lander, E. S., Aiden, A. P., & Aiden, E. L. (2017). De novo assembly

of the Aedes aegypti genome using Hi-C yields chromosome-length scaffolds. *Science*,

356(6333), 92–95. https://doi.org/10.1126/science.aal3327

Dudchenko, O., Shamim, M. S., Batra, S. S., Durand, N. C., Musial, N. T., Mostofa, R., Pham,

M., Hilaire, B. G. S., Yao, W., Stamenova, E., Hoeger, M., Nyquist, S. K., Korchina, V.,

Pletch, K., Flanagan, J. P., Tomaszewicz, A., McAloose, D., Estrada, C. P., Novak, B. J.,

… Aiden, E. L. (2018). The Juicebox Assembly Tools module facilitates de novo

assembly of mammalian genomes with chromosome-length scaffolds for under $1000.

BioRxiv, 254797. https://doi.org/10.1101/254797

Durand, N. C., Robinson, J. T., Shamim, M. S., Machol, I., Mesirov, J. P., Lander, E. S., &

Aiden, E. L. (2016). Juicebox Provides a Visualization System for Hi-C Contact Maps

with Unlimited Zoom. *Cell Systems*, *3*(1), 99–101.

https://doi.org/10.1016/j.cels.2015.07.012

Durand, N. C., Shamim, M. S., Machol, I., Rao, S. S. P., Huntley, M. H., Lander, E. S., & Aiden,

E. L. (2016). Juicer Provides a One-Click System for Analyzing Loop-Resolution Hi-C

Experiments. *Cell Systems*, *3*(1), 95–98. https://doi.org/10.1016/j.cels.2016.07.002

Edgar, R. C. (2010). Search and clustering orders of magnitude faster than BLAST.

Bioinformatics, *26*(19), 2460–2461. https://doi.org/10.1093/bioinformatics/btq461

Emerman, S. H., & Dawson, T. E. (1996). Hydraulic Lift and Its Influence on the Water Content

of the Rhizosphere: An Example from Sugar Maple, Acer saccharum. *Oecologia*, *108*(2),

273–278. JSTOR.

Emms, D. M., & Kelly, S. (2018). OrthoFinder2: Fast and accurate phylogenomic orthology

analysis from gene sequences. *BioRxiv*. https://doi.org/10.1101/466201

Enquist, B., Condit, R., Peet, R., Schildhauer, M., & Thiers, B. (2016). Cyberinfrastructure for

an integrated botanical information network to investigate the ecological impacts of

global climate change on plant biodiversity. PeerJ Preprints 4: E2615v2. *Int. J. Climatol*,

37, 4302–4315.

Estravis-Barcala, M., Mattera, M. G., Soliani, C., Bellora, N., Opgenoorth, L., Heer, K., &

Arana, M. V. (2020). Molecular bases of responses to abiotic stress in trees. *Journal of*

Experimental Botany, *71*(13), 3765–3779. https://doi.org/10.1093/jxb/erz532

Faust, G. G., & Hall, I. M. (2014). SAMBLASTER: Fast duplicate marking and structural

variant read extraction. *Bioinformatics*, *30*(17), 2503–2505.

https://doi.org/10.1093/bioinformatics/btu314

Flynn, J. M., Hubley, R., Goubert, C., Rosen, J., Clark, A. G., Feschotte, C., & Smit, A. F.

(2020). RepeatModeler2 for automated genomic discovery of transposable element

families. *Proceedings of the National Academy of Sciences*, *117*(17), 9451–9457.

https://doi.org/10.1073/pnas.1921046117

Genome Assembly Cookbook. (2019). DNA Zoo. https://www.dnazoo.org/methods

Giese, R. L., & Benjamin, D. M. (1964). The insect complex associated with maple blight.

University of Wisconsin, 21–57.

Godman, R. M., Yawney, H. W., & Tubbs, C. H. (1990). Acer saccharum Marsh. Sugar maple.

Silvics of North America, *2*(654), 78.

Grabherr, M. G., Haas, B. J., Yassour, M., Levin, J. Z., Thompson, D. A., Amit, I., Adiconis, X.,

Fan, L., Raychowdhury, R., Zeng, Q., Chen, Z., Mauceli, E., Hacohen, N., Gnirke, A.,

Rhind, N., di Palma, F., Birren, B. W., Nusbaum, C., Lindblad-Toh, K., … Regev, A.

(2011). Full-length transcriptome assembly from RNA-Seq data without a reference

genome. *Nature Biotechnology*, *29*(7), 644–652. https://doi.org/10.1038/nbt.1883

Gremme, G., Brendel, V., Sparks, M. E., & Kurtz, S. (2005). Engineering a software tool for

gene structure prediction in higher organisms. *Information and Software Technology*,

47(15), 965–978. https://doi.org/10.1016/j.infsof.2005.09.005

Halman, J. M., Schaberg, P. G., Hawley, G. J., Hansen, C. F., & Fahey, T. J. (2015). Differential

impacts of calcium and aluminum treatments on sugar maple and American beech growth

dynamics. *Canadian Journal of Forest Research*, *45*(1), 52–59.

https://doi.org/10.1139/cjfr-2014-0250

Halman, J. M., Schaberg, P. G., Hawley, G. J., Pardo, L. H., & Fahey, T. J. (2013). Calcium and

aluminum impacts on sugar maple physiology in a northern hardwood forest. *Tree*

Physiology, *33*(11), 1242–1251. https://doi.org/10.1093/treephys/tpt099

Hart, A. J., Ginzburg, S., Xu, M. (Sam), Fisher, C. R., Rahmatpour, N., Mitton, J. B., Paul, R., &

Wegrzyn, J. L. (2020). EnTAP: Bringing faster and smarter functional annotation to non-

model eukaryotic transcriptomes. *Molecular Ecology Resources*, *20*(2), 591–604.

https://doi.org/10.1111/1755-0998.13106

Hasan, M. K., Cheng, Y., Kanwar, M. K., Chu, X.-Y., Ahammed, G. J., & Qi, Z.-Y. (2017).

Responses of Plant Proteins to Heavy Metal Stress—A Review. *Frontiers in Plant*

Science, *8*. https://doi.org/10.3389/fpls.2017.01492

Hasanuzzaman, M., Nahar, K., Anee, T. I., & Fujita, M. (2017). Glutathione in plants:

Biosynthesis and physiological role in environmental stress tolerance. *Physiology and*

Molecular Biology of Plants, *23*(2), 249–268. https://doi.org/10.1007/s12298-017-0422-2

He, F., Pan, Q.-H., Shi, Y., & Duan, C.-Q. (2008). Biosynthesis and Genetic Regulation of

Proanthocyanidins in Plants. *Molecules*, *13*(10), 2674–2703.

https://doi.org/10.3390/molecules13102674

Helliwell, C. A., Chandler, P. M., Poole, A., Dennis, E. S., & Peacock, W. J. (2001). The

CYP88A cytochrome P450, ent-kaurenoic acid oxidase, catalyzes three steps of the

gibberellin biosynthesis pathway. *Proceedings of the National Academy of Sciences*,

98(4), 2065–2070. https://doi.org/10.1073/pnas.98.4.2065

Hendrix, S., Keunen, E., Mertens, A. I. G., Beemster, G. T. S., Vangronsveld, J., & Cuypers, A.

(2018). Cell cycle regulation in different leaves of Arabidopsis thaliana plants grown

under control and cadmium-exposed conditions. *Environmental and Experimental*

Botany, *155*, 441–452. https://doi.org/10.1016/j.envexpbot.2018.06.026

Hoff, K., Lomsadze, A., Borodovsky, M., & Stanke, M. (2019). Whole-Genome Annotation with

BRAKER. *Methods in Molecular Biology (Clifton, N.J.)*, *1962*, 65–95.

https://doi.org/10.1007/978-1-4939-9173-0_5

Horsley, S. B., Long, R. P., Bailey, S. W., Hallett, R. A., & Wargo, P. M. (2002). Health of

Eastern North American Sugar Maple Forests and Factors Affecting Decline. *Northern*

Journal of Applied Forestry, *19*(1), 34–44. https://doi.org/10.1093/njaf/19.1.34

Hu, B., Wang, N., Bi, X., Karaaslan, E. S., Weber, A.-L., Zhu, W., Berendzen, K. W., & Liu, C.

(2019). Plant lamin-like proteins mediate chromatin tethering at the nuclear periphery.

Genome Biology, *20*(1), 87. https://doi.org/10.1186/s13059-019-1694-3

Huang, S., Chen, Z., Huang, G., Yu, T., Yang, P., Li, J., Fu, Y., Yuan, S., Chen, S., & Xu, A.

(2012). HaploMerger: Reconstructing allelic relationships for polymorphic diploid

genome assemblies. *Genome Research*, *22*(8), 1581–1588.

https://doi.org/10.1101/gr.133652.111

Jasinski, S., Fabrissin, I., Masson, A., Marmagne, A., Lécureuil, A., Bill, L., & Chardon, F.

(2021). ACCELERATED CELL DEATH 6 Acts on Natural Leaf Senescence and

Nitrogen Fluxes in Arabidopsis. *Frontiers in Plant Science*, *11*.

https://doi.org/10.3389/fpls.2020.611170

Jockusch, S., & Kräutler, B. (2020). The red chlorophyll catabolite (RCC) is an inefficient

sensitizer of singlet oxygen – photochemical studies of the methyl ester of RCC.

Photochemical & Photobiological Sciences, *19*(5), 668–673.

https://doi.org/10.1039/D0PP00071J

Julius, B. T., Leach, K. A., Tran, T. M., Mertz, R. A., & Braun, D. M. (2017). Sugar

Transporters in Plants: New Insights and Discoveries. *Plant and Cell Physiology*, *58*(9),

1442–1460. https://doi.org/10.1093/pcp/pcx090

Kar, D., Pradhan, A. A., & Datta, S. (2021). The role of solute transporters in aluminum toxicity

and tolerance. *Physiologia Plantarum*, *171*(4), 638–652.

https://doi.org/10.1111/ppl.13214

Kim, D., Langmead, B., & Salzberg, S. L. (2015). HISAT: A fast spliced aligner with low

memory requirements. *Nature Methods*, *12*(4), 357–360.

https://doi.org/10.1038/nmeth.3317

Kolmogorov, M. (2019). *Fast and accurate de novo assembler for single molecule sequencing*

reads: Fenderglass/Flye [C++]. https://github.com/fenderglass/Flye (Original work

published 2016)

Koren, S., Walenz, B. P., Berlin, K., Miller, J. R., Bergman, N. H., & Phillippy, A. M. (2017).

Canu: Scalable and accurate long-read assembly via adaptive k-mer weighting and repeat

separation. *Genome Research*, *27*(5), 722–736. https://doi.org/10.1101/gr.215087.116

Lamarque, L. J., Lortie, C. J., Porté, A. J., & Delzon, S. (2015). Genetic differentiation and

phenotypic plasticity in life-history traits between native and introduced populations of

invasive maple trees. *Biological Invasions*, *17*(4), 1109–1122.

https://doi.org/10.1007/s10530-014-0781-3

Lamarque, L. J., Porté, A. J., Eymeric, C., Lasnier, J.-B., Lortie, C. J., & Delzon, S. (2013). A

Test for Pre-Adapted Phenotypic Plasticity in the Invasive Tree Acer negundo L. *PLOS*

ONE, *8*(9), e74239. https://doi.org/10.1371/journal.pone.0074239

Lang, L., & Schnittger, A. (2020). Endoreplication—A means to an end in cell growth and stress

response. *Current Opinion in Plant Biology*, *54*, 85–92.

https://doi.org/10.1016/j.pbi.2020.02.006

Leitch, I. J., Johnston, E., Pellicer, J., Hidalgo, O., & Bennett, M. (2019). *Angiosperm DNA C-*

values Database (release 9.0, Apr 2019). https://cvalues.science.kew.org/search

Li, H. (2018). Minimap2: Pairwise alignment for nucleotide sequences. *Bioinformatics*, *34*(18),

3094–3100. https://doi.org/10.1093/bioinformatics/bty191

Li, L.-S., Ying, J., Li, E., Ma, T., Li, M., Gong, L.-M., Wei, G., Zhang, Y., & Li, S. (2021).

Arabidopsis CBP60b is a central transcriptional activator of immunity. *Plant Physiology*,

kiab164. https://doi.org/10.1093/plphys/kiab164

Li, N., Meng, H., Xing, H., Liang, L., Zhao, X., & Luo, K. (2017). Genome-wide analysis of

MATE transporters and molecular characterization of aluminum resistance in Populus. *Journal of Experimental Botany*, *68*(20), 5669–5683. https://doi.org/10.1093/jxb/erx370

Li, X., Mao, X., Xu, Y., Li, Y., Zhao, N., Yao, J., Dong, Y., Tigabu, M., Zhao, X., & Li, S. (2021). Comparative transcriptomic analysis reveals the coordinated mechanisms of Populus × canadensis 'Neva' leaves in response to cadmium stress. *Ecotoxicology and Environmental Safety*, *216*, 112179. https://doi.org/10.1016/j.ecoenv.2021.112179

Liang, S. C., Hartwig, B., Perera, P., Mora-García, S., Leau, E. de, Thornton, H., Alves, F. L. de, Rapsilber, J., Yang, S., James, G. V., Schneeberger, K., Finnegan, E. J., Turck, F., & Goodrich, J. (2015). Kicking against the PRCs – A Domesticated Transposase Antagonises Silencing Mediated by Polycomb Group Proteins and Is an Accessory Component of Polycomb Repressive Complex 2. *PLOS Genetics*, *11*(12), e1005660. https://doi.org/10.1371/journal.pgen.1005660

Likens, G. E., & Bormann, F. H. (1974). Acid Rain: A Serious Regional Environmental Problem. *Science*, *184*(4142), 1176–1179. https://doi.org/10.1126/science.184.4142.1176

Likens, G. E., Driscoll, C. T., Buso, D. C., Siccama, T. G., Johnson, C. E., Lovett, G. M., Fahey, T. J., Reiners, W. A., Ryan, D. F., Martin, C. W., & Bailey, S. W. (1998). The biogeochemistry of calcium at Hubbard Brook. *Biogeochemistry*, *41*(2), 89–173. https://doi.org/10.1023/A:1005984620681

Likens, G. E., & Lambert, K. F. (1998). The Importance of Long-Term Data in Addressing Regional Environmental Issues. *Northeastern Naturalist*, *5*(2), 127–136. https://doi.org/10.2307/3858583

Liu, Z., Chen, G., Gao, F., Xu, R., Li, N., Zhang, Y., & Li, Y. (2019). Transcriptional Repression of the APC/C Activator Genes CCS52A1/A2 by the Mediator Complex Subunit MED16

Controls Endoreduplication and Cell Growth in Arabidopsis. *The Plant Cell*, *31*(8), 1899–1912. https://doi.org/10.1105/tpc.18.00811

Lomsadze, A., Burns, P. D., & Borodovsky, M. (2014). Integration of mapped RNA-Seq reads into automatic training of eukaryotic gene finding algorithm. *Nucleic Acids Research*, *42*(15), e119–e119. https://doi.org/10.1093/nar/gku557

Long, R. P., Horsley, S. B., Bailey, S. W., Hallett, R. A., & Hall, T. J. (2019). Sugar maple decline and lessons learned about Allegheny Plateau soils and landscapes. *In: Stout, Susan L., Ed. SILVAH: 50 Years of Science-Management Cooperation. Proceedings of the Allegheny Society of American Foresters Training Session; 2017 Sept. 20-22; Clarion, PA. Gen. Tech. Rep. NRS-P-186. Newtown Square, PA: U.S. Department of Agriculture, Forest Service, Northern Research Station: 80-97.*, 80–97. https://doi.org/10.2737/NRS-GTR-P-186-Paper8

Love, M., Anders, S., & Huber, W. (2014). Differential analysis of count data–the DESeq2 package. *Genome Biol*, *15*(550), 10–1186.

Lucash, M. S., Yanai, R. D., Blum, J. D., & Park, B. B. (2012). Foliar Nutrient Concentrations Related to Soil Sources across a Range of Sites in the Northeastern United States. *Soil Science Society of America Journal*, *76*(2), 674–683. https://doi.org/10.2136/sssaj2011.0160

Ludwig-Müller, J. (2011). Auxin conjugates: Their role for plant development and in the evolution of land plants. *Journal of Experimental Botany*, *62*(6), 1757–1773. https://doi.org/10.1093/jxb/erq412

Luo, J., Zhou, J., Li, H., Shi, W., Polle, A., Lu, M., Sun, X., & Luo, Z.-B. (2015). Global poplar root and leaf transcriptomes reveal links between growth and stress responses under

nitrogen starvation and excess. *Tree Physiology*, *35*(12), 1283–1302.

https://doi.org/10.1093/treephys/tpv091

Ma, Q., Sun, T., Li, S., Wen, J., Zhu, L., Yin, T., Yan, K., Xu, X., Li, S., Mao, J., Wang, Y., Jin,

S., Zhao, X., & Li, Q. (2020). The Acer truncatum genome provides insights into

nervonic acid biosynthesis. *The Plant Journal*, *104*(3), 662–678.

https://doi.org/10.1111/tpj.14954

Mach, J. M., Castillo, A. R., Hoogstraten, R., & Greenberg, J. T. (2001). The Arabidopsis-

accelerated cell death gene ACD2 encodes red chlorophyll catabolite reductase and

suppresses the spread of disease symptoms. *Proceedings of the National Academy of*

Sciences, *98*(2), 771–776. https://doi.org/10.1073/pnas.98.2.771

Maitner, B. S., Boyle, B., Casler, N., Condit, R., Donoghue, J., Durán, S. M., Guaderrama, D.,

Hinchliff, C. E., Jørgensen, P. M., Kraft, N. J. B., McGill, B., Merow, C., Morueta-

Holme, N., Peet, R. K., Sandel, B., Schildhauer, M., Smith, S. A., Svenning, J.-C., Thiers,

B., … Enquist, B. J. (2018). The bien r package: A tool to access the Botanical

Information and Ecology Network (BIEN) database. *Methods in Ecology and Evolution*,

9(2), 373–379. https://doi.org/10.1111/2041-210X.12861

Mekawy, A. M. M., Assaha, D. V. M., Munehiro, R., Kohnishi, E., Nagaoka, T., Ueda, A., &

Saneoka, H. (2018). Characterization of type 3 metallothionein-like gene (OsMT-3a)

from rice, revealed its ability to confer tolerance to salinity and heavy metal stresses.

Environmental and Experimental Botany, *147*, 157–166.

https://doi.org/10.1016/j.envexpbot.2017.12.002

Mendes, F. K., Vanderpool, D., Fulton, B., & Hahn, M. W. (2020). CAFE 5 models variation in

evolutionary rates among gene families. *Bioinformatics*, *btaa1022*.

https://doi.org/10.1093/bioinformatics/btaa1022

Miao, H., Sun, P., Liu, Q., Miao, Y., Liu, J., Zhang, K., Hu, W., Zhang, J., Wang, J., Wang, Z.,

    Jia, C., Xu, B., & Jin, Z. (2017). Genome-wide analyses of SWEET family proteins

    reveal involvement in fruit development and abiotic/biotic stress responses in banana.

    Scientific Reports, *7*(1), 3536. https://doi.org/10.1038/s41598-017-03872-w

Mohamed, H. K., Pathak, S., Roy, D. N., Hutchinson, T. C., McLaughlin, D. L., & Kinch, J. C.

    (1997). RELATIONSHIP BETWEEN SUGAR MAPLE DECLINE AND

    CORRESPONDING CHEMICAL CHANGES IN THE STEM TISSUE. *Water, Air, and*

    Soil Pollution, *96*(1), 321–327. https://doi.org/10.1023/A:1026429122530

Moore, J.-D., & Ouimet, R. (2021). Liming still positively influences sugar maple nutrition,

    vigor and growth, 20 years after a single application. *Forest Ecology and Management*,

    490, 119103. https://doi.org/10.1016/j.foreco.2021.119103

Naslavsky, N., & Caplan, S. (2005). C-terminal EH-domain-containing proteins: Consensus for a

    role in endocytic trafficking, EH? *Journal of Cell Science*, *118*(18), 4093–4101.

    https://doi.org/10.1242/jcs.02595

Nisa, M.-U., Huang, Y., Benhamed, M., & Raynaud, C. (2019). The Plant DNA Damage

    Response: Signaling Pathways Leading to Growth Inhibition and Putative Role in

    Response to Stress Conditions. *Frontiers in Plant Science*, *10*.

    https://doi.org/10.3389/fpls.2019.00653

Nkongolo, K., Theriault, G., & Michael, P. (2018). Nickel-induced global gene expressions in

    red maple (Acer rubrum): Effect of nickel concentrations. *Plant Gene*, *14*, 29–36.

    https://doi.org/10.1016/j.plgene.2018.04.003

Nowell, R. W., Almeida, P., Wilson, C. G., Smith, T. P., Fontaneto, D., Crisp, A., Micklem, G.,

Tunnacliffe, A., Boschetti, C., & Barraclough, T. G. (2018). Comparative genomics of

bdelloid rotifers: Insights from desiccating and nondesiccating species. *PLOS Biology*,

16(4), e2004830. https://doi.org/10.1371/journal.pbio.2004830

Oswald, E. M., Pontius, J., Rayback, S. A., Schaberg, P. G., Wilmot, S. H., & Dupigny-Giroux,

L.-A. (2018). The complex relationship between climate and sugar maple health: Climate

change implications in Vermont for a key northern hardwood species. *Forest Ecology*

and Management, *422*, 303–312. https://doi.org/10.1016/j.foreco.2018.04.014

Park, J.-E., Park, J.-Y., Kim, Y.-S., Staswick, P. E., Jeon, J., Yun, J., Kim, S.-Y., Kim, J., Lee,

Y.-H., & Park, C.-M. (2007). GH3-mediated Auxin Homeostasis Links Growth

Regulation with Stress Adaptation Response in Arabidopsis*. *Journal of Biological*

Chemistry, *282*(13), 10036–10046. https://doi.org/10.1074/jbc.M610524200

Pellicer, J., Hidalgo, O., Dodsworth, S., & Leitch, I. J. (2018). Genome Size Diversity and Its

Impact on the Evolution of Land Plants. *Genes*, *9*(2), 88.

https://doi.org/10.3390/genes9020088

Pereira, J. F., & Ryan, P. R. (2019). The role of transposable elements in the evolution of

aluminium resistance in plants. *Journal of Experimental Botany*, *70*(1), 41–54.

https://doi.org/10.1093/jxb/ery357

Pertea, G. (2018). *GffCompare*. http://ccb.jhu.edu/software/stringtie/gffcompare.shtml

Porté, A. J., Lamarque, L. J., Lortie, C. J., Michalet, R., & Delzon, S. (2011). Invasive Acer

negundo outperforms native species in non-limiting resource environments due to its

higher phenotypic plasticity. *BMC Ecology*, *11*(1), 28. https://doi.org/10.1186/1472-

6785-11-28

Pyšek, P., Skálová, H., Čuda, J., Guo, W.-Y., Suda, J., Doležal, J., Kauzál, O., Lambertini, C.,

Lučanová, M., Mandáková, T., Moravcová, L., Pyšková, K., Brix, H., & Meyerson, L. A.

(2018). Small genome separates native and invasive populations in an ecologically

important cosmopolitan grass. *Ecology*, *99*(1), 79–90. https://doi.org/10.1002/ecy.2068

Qiao, X., Li, Q., Yin, H., Qi, K., Li, L., Wang, R., Zhang, S., & Paterson, A. H. (2019). Gene

duplication and evolution in recurring polyploidization–diploidization cycles in plants.

Genome Biology, *20*(1), 38. https://doi.org/10.1186/s13059-019-1650-2

Raudvere, U., Kolberg, L., Kuzmin, I., Arak, T., Adler, P., Peterson, H., & Vilo, J. (2019).

g:Profiler: A web server for functional enrichment analysis and conversions of gene lists

(2019 update). *Nucleic Acids Research*, *47*(W1), W191–W198.

https://doi.org/10.1093/nar/gkz369

Renner, S. S., Beenken, L., Grimm, G. W., Kocyan, A., & Ricklefs, R. E. (2007). The Evolution

of Dioecy, Heterodichogamy, and Labile Sex Expression in Acer. *Evolution*, *61*(11),

2701–2719. https://doi.org/10.1111/j.1558-5646.2007.00221.x

Rizza, A., Boccaccini, A., Lopez-Vidriero, I., Costantino, P., & Vittorioso, P. (2011).

Inactivation of the ELIP1 and ELIP2 genes affects Arabidopsis seed germination. *New*

Phytologist, *190*(4), 896–905. https://doi.org/10.1111/j.1469-8137.2010.03637.x

Roach, M. J., Schmidt, S. A., & Borneman, A. R. (2018). Purge Haplotigs: Allelic contig

reassignment for third-gen diploid genome assemblies. *BMC Bioinformatics*, *19*(1), 460.

https://doi.org/10.1186/s12859-018-2485-7

Roddy, A. B., Théroux-Rancourt, G., Abbo, T., Benedetti, J. W., Brodersen, C. R., Castro, M.,

Castro, S., Gilbride, A. B., Jensen, B., Jiang, G.-F., Perkins, J. A., Perkins, S. D.,

Loureiro, J., Syed, Z., Thompson, R. A., Kuebbing, S. E., & Simonin, K. A. (2019). The

Scaling of Genome Size and Cell Size Limits Maximum Rates of Photosynthesis with

Implications for Ecological Strategies. *International Journal of Plant Sciences*, *181*(1),

75–87. https://doi.org/10.1086/706186

Rogers, B. M., Jantz, P., & Goetz, S. J. (2017). Vulnerability of eastern US tree species to

climate change. *Global Change Biology*, *23*(8), 3302–3320.

https://doi.org/10.1111/gcb.13585

Sade, H., Meriga, B., Surapu, V., Gadi, J., Sunita, M. S. L., Suravajhala, P., & Kavi Kishor, P. B.

(2016). Toxicity and tolerance of aluminum in plants: Tailoring plants to suit to acid

soils. *BioMetals*, *29*(2), 187–210. https://doi.org/10.1007/s10534-016-9910-z

Sakamoto, A. N. (2019). Translesion Synthesis in Plants: Ultraviolet Resistance and Beyond.

Frontiers in Plant Science, *10*, 1208. https://doi.org/10.3389/fpls.2019.01208

Shi, X., Sun, H., Chen, Y., Pan, H., & Wang, S. (2016). Transcriptome Sequencing and

Expression Analysis of Cadmium (Cd) Transport and Detoxification Related Genes in

Cd-Accumulating Salix integra. *Frontiers in Plant Science*, *7*.

https://doi.org/10.3389/fpls.2016.01577

Smit, A., Hubley, R., & Green, P. (2013). *RepeatMasker Open-4.0*. http://www.repeatmasker.org

Smoot, M. E., Ono, K., Ruscheinski, J., Wang, P.-L., & Ideker, T. (2011). Cytoscape 2.8: New

features for data integration and network visualization. *Bioinformatics*, *27*(3), 431–432.

https://doi.org/10.1093/bioinformatics/btq675

Sork, V. L., Fitz-Gibbon, S. T., Puiu, D., Crepeau, M., Gugger, P. F., Sherman, R., Stevens, K.,

Langley, C. H., Pellegrini, M., & Salzberg, S. L. (2016). First Draft Assembly and

Annotation of the Genome of a California Endemic Oak Quercus lobata Née (Fagaceae).

G3: Genes, Genomes, Genetics, *6*(11), 3485–3495.

https://doi.org/10.1534/g3.116.030411

Stafford, H. A. (1988). Proanthocyanidins and the lignin connection. *Phytochemistry*, *27*(1), 1–6.

https://doi.org/10.1016/0031-9422(88)80583-1

Stanke, M., Diekhans, M., Baertsch, R., & Haussler, D. (2008). Using native and syntenically

mapped cDNA alignments to improve de novo gene finding. *Bioinformatics*, *24*(5), 637–

644. https://doi.org/10.1093/bioinformatics/btn013

Stanke, M., Schöffmann, O., Morgenstern, B., & Waack, S. (2006). Gene prediction in

eukaryotes with a generalized hidden Markov model that uses hints from external

sources. *BMC Bioinformatics*, *7*(1), 62. https://doi.org/10.1186/1471-2105-7-62

Staton, M., Best, T., Khodwekar, S., Owusu, S., Xu, T., Xu, Y., Jennings, T., Cronn, R.,

Arumuganathan, A. K., Coggeshall, M., Gailing, O., Liang, H., Romero-Severson, J.,

Schlarbaum, S., & Carlson, J. E. (2015). Preliminary Genomic Characterization of Ten

Hardwood Tree Species from Multiplexed Low Coverage Whole Genome Sequencing.

PLOS ONE, *10*(12), e0145031. https://doi.org/10.1371/journal.pone.0145031

Stevens, R., Grelon, M., Vezon, D., Oh, J., Meyer, P., Perennes, C., Domenichini, S., &

Bergounioux, C. (2004). A CDC45 Homolog in Arabidopsis Is Essential for Meiosis, as

Shown by RNA Interference–Induced Gene Silencing. *The Plant Cell*, *16*(1), 99–113.

https://doi.org/10.1105/tpc.016865

Suda, J., Meyerson, L. A., Leitch, I. J., & Pyšek, P. (2015). The hidden side of plant invasions:

The role of genome size. *New Phytologist*, *205*(3), 994–1007.

https://doi.org/10.1111/nph.13107

Sullivan, T. J., Lawrence, G. B., Bailey, S. W., McDonnell, T. C., Beier, C. M., Weathers, K. C.,

McPherson, G. T., & Bishop, D. A. (2013). Effects of Acidic Deposition and Soil

Acidification on Sugar Maple Trees in the Adirondack Mountains, New York.

Environmental Science & Technology, *47*(22), 12687–12694.

https://doi.org/10.1021/es401864w

Sun, M., Li, H., Li, Y., Xiang, H., Liu, Y., He, Y., Qi, M., & Li, T. (2020). Tomato YABBY2b

controls plant height through regulating indole-3-acetic acid-amido synthetase (GH3.8)

expression. *Plant Science*, *297*, 110530. https://doi.org/10.1016/j.plantsci.2020.110530

Sun, X.-L., Yu, Q.-Y., Tang, L.-L., Ji, W., Bai, X., Cai, H., Liu, X.-F., Ding, X.-D., & Zhu, Y.-

M. (2013). GsSRK, a G-type lectin S-receptor-like serine/threonine protein kinase, is a

positive regulator of plant tolerance to salt stress. *Journal of Plant Physiology*, *170*(5),

505–515. https://doi.org/10.1016/j.jplph.2012.11.017

Tang, H., Bowers, J. E., Wang, X., Ming, R., Alam, M., & Paterson, A. H. (2008). Synteny and

Collinearity in Plant Genomes. *Science*, *320*(5875), 486–488.

https://doi.org/10.1126/science.1153917

Theriault, G., Michael, P., & Nkongolo, K. (2016). Comprehensive Transcriptome Analysis of

Response to Nickel Stress in White Birch (Betula papyrifera). *PLOS ONE*, *11*(4),

e0153762. https://doi.org/10.1371/journal.pone.0153762

Tian, J., & Kong, Z. (2019). The role of the augmin complex in establishing microtubule arrays.

Journal of Experimental Botany, *70*(12), 3035–3041. https://doi.org/10.1093/jxb/erz123

Tommasini, R., Vogt, E., Fromenteau, M., Hörtensteiner, S., Matile, P., Amrhein, N., &

Martinoia, E. (1998). An ABC-transporter of Arabidopsis thaliana has both glutathione-

conjugate and chlorophyll catabolite transport activity. *The Plant Journal*, *13*(6), 773–

780. https://doi.org/10.1046/j.1365-313X.1998.00076.x

Toyota, M., Spencer, D., Sawai-Toyota, S., Jiaqi, W., Zhang, T., Koo, A. J., Howe, G. A., &

Gilroy, S. (2018). Glutamate triggers long-distance, calcium-based plant defense

signaling. *Science*, *361*(6407), 1112–1115. https://doi.org/10.1126/science.aat7744

Trávníček, P., Čertner, M., Ponert, J., Chumová, Z., Jersáková, J., & Suda, J. (2019). Diversity in genome size and GC content shows adaptive potential in orchids and is closely linked to partial endoreplication, plant life-history traits and climatic conditions. *New Phytologist*, *224*(4), 1642–1656. https://doi.org/10.1111/nph.15996

Tuskan, G. A., DiFazio, S., Jansson, S., Bohlmann, J., Grigoriev, I., Hellsten, U., Putnam, N., Ralph, S., Rombauts, S., Salamov, A., Schein, J., Sterck, L., Aerts, A., Bhalerao, R. R., Bhalerao, R. P., Blaudez, D., Boerjan, W., Brun, A., Brunner, A., … Rokhsar, D. (2006). The Genome of Black Cottonwood, Populus trichocarpa (Torr. & Gray). *Science*, *313*(5793), 1596–1604. https://doi.org/10.1126/science.1128691

Van Bel, M., Bucchini, F., & Vandepoele, K. (2019). Gene space completeness in complex plant genomes. *Current Opinion in Plant Biology*, *48*, 9–17. https://doi.org/10.1016/j.pbi.2019.01.001

Veeckman, E., Ruttink, T., & Vandepoele, K. (2016). Are We There Yet? Reliably Estimating the Completeness of Plant Genome Sequences. *The Plant Cell*, *28*(8), 1759–1768. https://doi.org/10.1105/tpc.16.00349

Vurture, G. W., Sedlazeck, F. J., Nattestad, M., Underwood, C. J., Fang, H., Gurtowski, J., & Schatz, M. C. (2017). GenomeScope: Fast reference-free genome profiling from short reads. *Bioinformatics*, *33*(14), 2202–2204. https://doi.org/10.1093/bioinformatics/btx153

Wu, T. D., & Watanabe, C. K. (2005). GMAP: A genomic mapping and alignment program for mRNA and EST sequences. *Bioinformatics*, *21*(9), 1859–1875. https://doi.org/10.1093/bioinformatics/bti310

Xia, K., Ou, X., Tang, H., Wang, R., Wu, P., Jia, Y., Wei, X., Xu, X., Kang, S.-H., Kim, S.-K.,

& Zhang, M. (2015). Rice microRNA osa-miR1848 targets the obtusifoliol 14α-demethylase gene OsCYP51G3 and mediates the biosynthesis of phytosterols and brassinosteroids during development and in response to stress. *New Phytologist*, *208*(3), 790–802. https://doi.org/10.1111/nph.13513

Xu, M., Cho, E., Burch-Smith, T. M., & Zambryski, P. C. (2012). Plasmodesmata formation and cell-to-cell transport are reduced in decreased size exclusion limit 1 during embryogenesis in Arabidopsis. *Proceedings of the National Academy of Sciences*, *109*(13), 5098–5103. https://doi.org/10.1073/pnas.1202919109

Yamamoto, Y., Kobayashi, Y., Devi, S. R., Rikiishi, S., & Matsumoto, H. (2003). Oxidative stress triggered by aluminum in plant roots. In J. Abe (Ed.), *Roots: The Dynamic Interface between Plants and the Earth: The 6th Symposium of the International Society of Root Research, 11–15 November 2001, Nagoya, Japan* (pp. 239–243). Springer Netherlands. https://doi.org/10.1007/978-94-017-2923-9_23

Yang, J., Wariss, H. M., Tao, L., Zhang, R., Yun, Q., Hollingsworth, P., Dao, Z., Luo, G., Guo, H., Ma, Y., & Sun, W. (2019). De novo genome assembly of the endangered Acer yangbiense, a plant species with extremely small populations endemic to Yunnan Province, China. *GigaScience*, *8*(7). https://doi.org/10.1093/gigascience/giz085

Yang, W.-Y., Zheng, Y., Bahn, S. C., Pan, X.-Q., Li, M.-Y., Vu, H. S., Roth, M. R., Scheu, B., Welti, R., Hong, Y.-Y., & Wang, X.-M. (2012). The Patatin-Containing Phospholipase A pPLAIIα Modulates Oxylipin Formation and Water Loss in Arabidopsis thaliana. *Molecular Plant*, *5*(2), 452–460. https://doi.org/10.1093/mp/ssr118

Yu, T., Hu, Y., Zhang, Y., Zhao, R., Yan, X., Dayananda, B., Wang, J., Jiao, Y., Li, J., & Yi, X. (2021). Whole-genome sequencing of Acer catalpifolium reveals evolutionary history of

endangered species. *Genome Biology and Evolution*, evab271.

https://doi.org/10.1093/gbe/evab271

Zeng, L., Tu, X.-L., Dai, H., Han, F.-M., Lu, B.-S., Wang, M.-S., Nanaei, H. A., Tajabadipour,

A., Mansouri, M., Li, X.-L., Ji, L.-L., Irwin, D. M., Zhou, H., Liu, M., Zheng, H.-K.,

Esmailizadeh, A., & Wu, D.-D. (2019). Whole genomes and transcriptomes reveal

adaptation and domestication of pistachio. *Genome Biology*, *20*(1), 79.

https://doi.org/10.1186/s13059-019-1686-3

Zhang, S.-W., Li, C.-H., Cao, J., Zhang, Y.-C., Zhang, S.-Q., Xia, Y.-F., Sun, D.-Y., & Sun, Y.

(2009). Altered Architecture and Enhanced Drought Tolerance in Rice via the Down-

Regulation of Indole-3-Acetic Acid by TLD1/OsGH3.13 Activation. *Plant Physiology*,

151(4), 1889–1901. https://doi.org/10.1104/pp.109.146803

Zhao, D., Hamilton, J. P., Bhat, W. W., Johnson, S. R., Godden, G. T., Kinser, T. J., Boachon,

B., Dudareva, N., Soltis, D. E., Soltis, P. S., Hamberger, B., & Buell, C. R. (2019). A

chromosomal-scale genome assembly of Tectona grandis reveals the importance of

tandem gene duplication and enables discovery of genes in natural product biosynthetic

pathways. *GigaScience*, *8*(3). https://doi.org/10.1093/gigascience/giz005

Zhu, W., Li, L., Neuhäuser, B., Thelen, M., Wang, M., Chen, J., Wei, L., Venkataramani, K.,

Exposito-Alonso, M., Liu, C., Keck, J., Barragan, A. C., Schwab, R., Lutz, U., Ludewig,

U., & Weigel, D. (2021). Small peptides modulate the immune function of the ion

channel-like protein ACD6 in Arabidopsis thaliana. *BioRxiv*, 2021.01.25.428077.

https://doi.org/10.1101/2021.01.25.428077

1426

1427   **Figure and Table Captions**

1428

Figure 1. a) Native distributions of *A. saccharum* (blue) and *A. negundo* (orange) in North

America. Leaves indicate location of individuals selected for the reference genomes; *A.*

saccharum from the University of Maryland campus, and *A. negundo* from the Smithsonian

Environmental Research Center. HBEF (Hubbard Brook Experimental Forest) is the location of

the 9 individuals used for RNA-seq. b) All records of occurrence, native, introduced, and

unknown, per BIEN 4.2. Non-native occurrences are predominantly *A. negundo*.

https://gitlab.com/PlantGenomicsLab/AcerGenomes/-

/blob/master/acer/supplemental/figures/acerranges.jpg

1437

Figure 2. Results of assembly testing with *A. saccharum*, comparing fragmentation in terms of

total contigs versus assembly length. The dashed line represents the estimated genome size. Gray

dots are short-read assemblers, shown as highly fragmented. Blue dots are long-read tests of

assembly workflows. Canu refers to the use of reads error-corrected by the Canu pipeline. The

red dot is the selected draft assembly, and the green dot shows scaffolding results following Hi-

C. Detailed assembly statistics are available in File S1.

https://gitlab.com/PlantGenomicsLab/AcerGenomes/-

/blob/master/acer/supplemental/figures/assemblycomparisongraph_busco.jpg

1446

Figure 3. Ks distribution for WGD synteny blocks with a summary of duplication types in (a) *A.*

negundo and (b) *A. saccharum*. Abbreviations for categories of duplication: WGD, whole

genome duplication; TD, tandem duplication; PD, proximal duplication; TRD, transposed

duplication; DSD, dispersed duplication. (c) Circos plot of the thirteen chromosomes ordered

largest to smallest for *A. negundo* (orange bars) and *A. saccharum* (blue bars) with distributions

of gene density (green) and transposable element frequency (purple). Syntenic regions are linked

in gray with darker shades to visually highlight larger recombinations.

https://gitlab.com/PlantGenomicsLab/AcerGenomes/-

/blob/master/acer/supplemental/figures/acsa_acne_manuscript_circos.png

1456

1457 **Figure 4.** a) Differential expression study design showing number of samples collected in fall

1458 and spring from treatment plots at the Hubbard Brook Experimental Forest, Nutrient Perturbation

1459 study. b) Differentially expressed genes (up and downregulated) for each treatment and season

1460 comparison. Charts display both significance and relative expression denoted as log-fold change.

1461 Dotted lines indicate thresholds of significance (0.1 p-adjusted, 1.5 $\log_2$ fold change).

1462 https://gitlab.com/PlantGenomicsLab/AcerGenomes/-

1463 /blob/master/acer/supplemental/figures/hbefplots.jpg

1464

1465 **Figure 5.** a) Gene ontology enrichments for *Acer* (all three species combined), *A. negundo*, and

1466 *A. saccharum*. Abbreviations for gene family dynamics: E, expanded; N, novel; RC, rapidly

1467 contracting. b) Total gene families, shared and unique, among the *Acer*. c) Reconstructed gene

1468 tree showing contracted gene families in red and expanded in green.

1469 https://gitlab.com/PlantGenomicsLab/AcerGenomes/-

1470 /blob/master/acer/supplemental/figures/cafetreevennGO.jpg

1471

1472 **Figure 6.** Orthogroup sizes for aluminum tolerance gene families are presented by species.

1473 Families were selected for inclusion based on documented aluminum tolerance and/or presence

1474 in the HBEF RNA-Seq differential expression results. Color represents the proportion of gene

1475 membership per species, with darker purple equating to more contracted families relative to the

1476 median, and dark green indicating expansion. (H) Family contains HBEF differentially expressed

1477 gene; (E) Expanded in *A. saccharum*; (C) Contracting; (M) Missing; (N) Novel; (*) Rapidly

1478 expanding; Categorization of tolerance is according to literature describing aluminum stress

1479 phenotypes. The undetermined category contains species where tolerance to aluminum or acidic

1480 soils has not been reported. [1]*B. pendula* is undetermined due to high variability in tolerance by

1481 genotype.

1482 https://gitlab.com/PlantGenomicsLab/AcerGenomes/-

1483 /blob/master/acer/supplemental/figures/genefamily_al_comparisons_manuscript_customnorm.ht

1484 ml.jpg

1485

1486 **Figure 7.** *A. negundo* gene families with ontology related to DNA damage and repair, and

1487 secondary enrichments categorized by color. Circles with multiple colors indicate multiple

1488 ontology assignments. Lines indicated known or predicted interactions, or other association via

1489 text-mining, co-expression, or protein homology.

1490 https://gitlab.com/PlantGenomicsLab/AcerGenomes/-

1491 /blob/master/acer/supplemental/figures/acne_dnadamage.jpg

1492

1493 **Supplemental**

1494

Figure S1. Genome size estimation using k-mer distribution analysis

https://gitlab.com/PlantGenomicsLab/AcerGenomes/-

/blob/master/acer/supplemental/figures/Figure_1_SuppInfo.pdf

Figure S2. Hi-C plots https://gitlab.com/PlantGenomicsLab/AcerGenomes/-

/blob/master/acer/supplemental/figures/Figure_2_SuppInfo.pdf

Figure S3. PCA plot

https://gitlab.com/PlantGenomicsLab/AcerGenomes/-

/blob/master/acer/supplemental/figures/Figure_3_SuppInfo.pdf

Figure S4. Syntenic comparisons between the three *Acer* genomes

https://gitlab.com/PlantGenomicsLab/AcerGenomes/-

/blob/master/acer/supplemental/figures/Figure_4_SuppInfo.pdf

1506

Table S1. Illumina, PacBio, and Hi-C sequencing data summaries

https://gitlab.com/PlantGenomicsLab/AcerGenomes/-

/blob/master/acer/supplemental/tables/sequencing_tables.pdf

Table S2. HBEF table of trees

https://gitlab.com/PlantGenomicsLab/AcerGenomes/-

/blob/master/acer/supplemental/tables/hbef_trees.pdf

Table S3. HBEF GO enrichment

https://gitlab.com/PlantGenomicsLab/AcerGenomes/-

/blob/master/acer/supplemental/tables/hbef_functional_enrichment.pdf

Table S4. Orthogroup statistics by species

https://gitlab.com/PlantGenomicsLab/AcerGenomes/-

/blob/master/acer/supplemental/tables/Orthogroup_summary_table.pdf

Table S5. OF dynamics GO enrichment

https://gitlab.com/PlantGenomicsLab/AcerGenomes/-

/blob/master/acer/supplemental/tables/orthogroup_functional_enrichment.pdf

Table S6. CAFE GO enrichment

https://gitlab.com/PlantGenomicsLab/AcerGenomes/-

/blob/master/acer/supplemental/tables/cafe_functional_enrichment.pdf

Table S7. HBEF Nutrient Perturbation Treatment Schedule

https://gitlab.com/PlantGenomicsLab/AcerGenomes/-

/blob/master/acer/supplemental/tables/HBEFNuPertTreatmentTable2015.pdf

1528

File S1. Assembly output stats

https://gitlab.com/PlantGenomicsLab/AcerGenomes/-

/blob/master/acer/supplemental/files/assembly_statistics.xlsx

File S2. Annotation details

https://gitlab.com/PlantGenomicsLab/AcerGenomes/-

/blob/master/acer/supplemental/files/annotation_statistics.xlsx

File S3. Collinearity analysis of recent Ks peak of WGD frequency

https://gitlab.com/PlantGenomicsLab/AcerGenomes/-

/blob/master/acer/supplemental/files/Acer_microsynteny.xlsx

File S4. HBEF differentially expressed genes

https://gitlab.com/PlantGenomicsLab/AcerGenomes/-

/blob/master/acer/supplemental/files/HBEF_DEGs.xlsx

File S5. Orthofinder significant dynamics

https://gitlab.com/PlantGenomicsLab/AcerGenomes/-

/blob/master/acer/supplemental/files/orthofinder_dynamics.xlsx

File S6. CAFE significant rapid evolution

https://gitlab.com/PlantGenomicsLab/AcerGenomes/-

/blob/master/acer/supplemental/files/cafe_rapidly_evolving.xlsx

File S7. *A. negundo* vs *Acer* contracted or missing using longest overall gene as annotation

https://gitlab.com/PlantGenomicsLab/AcerGenomes/-

/blob/master/acer/supplemental/files/acne_contracedmissing_longestoverall.xlsx

File S8. *A. negundo* vs *Acer* contracted or missing using longest *Acer* gene as annotation

https://gitlab.com/PlantGenomicsLab/AcerGenomes/-

/blob/master/acer/supplemental/files/acne_contractedmissing_longestacer.xlsx

File S9. *A. negundo* vs *Acer* expanded or novel using longest overall gene as annotation

https://gitlab.com/PlantGenomicsLab/AcerGenomes/-

/blob/master/acer/supplemental/files/acne_expandednovel_longestacer.xlsx

File S10. *A. negundo* vs *Acer* expanded or novel using longest *Acer* gene as annotation

https://gitlab.com/PlantGenomicsLab/AcerGenomes/-

/blob/master/acer/supplemental/files/acne_expandednovel_longestoverall.xlsx

File S11. *A. saccharum* vs *Acer* contracted or missing using longest *Acer* gene as annotation

https://gitlab.com/PlantGenomicsLab/AcerGenomes/-

/blob/master/acer/supplemental/files/acsa_contractedmissing_longestacer.xlsx

File S12. *A. saccharum* vs *Acer* contracted or missing using longest overall gene as annotation

https://gitlab.com/PlantGenomicsLab/AcerGenomes/-

/blob/master/acer/supplemental/files/acsa_contractedmissing_longestoverall.xlsx

File S13. *A. saccharum* vs *Acer* expanded or novel using longest *Acer* gene as annotation

https://gitlab.com/PlantGenomicsLab/AcerGenomes/-

/blob/master/acer/supplemental/files/acsa_expandednovel_longestacer.xlsx

File S14. *A. negundo* verified missing orthogroups

https://gitlab.com/PlantGenomicsLab/AcerGenomes/-

/blob/master/acer/supplemental/files/acne_missing_verified_string_mapping.tsv

File S15. *A. saccharum* verified missing orthogroups

https://gitlab.com/PlantGenomicsLab/AcerGenomes/-

/blob/master/acer/supplemental/files/acsa_missing_verified_string_mapping.tsv

File S16. Orthogroup comparisons for HBEF DEG and Al tolerance genes

https://gitlab.com/PlantGenomicsLab/AcerGenomes/-

/blob/master/acer/supplemental/files/hbef_orthogroup_Al_comparisons.xlsx