Phylogenomic analyses of echinoid diversification prompt a re-

evaluation of their fossil record

Nicolás Mongiardino Koch[1,2]*, Jeffrey R Thompson[3,4], Avery S Hatch[2], Marina F McCowin[2], A

Frances Armstrong[5], Simon E Coppard[6], Felipe Aguilera[7], Omri Bronstein[8,9], Andreas Kroh[10], Rich

Mooi[5], Greg W Rouse[2]

[1] Department of Earth & Planetary Sciences, Yale University, New Haven CT, USA

[2] Scripps Institution of Oceanography, University of California San Diego, La Jolla CA, USA

[3] Department of Earth Sciences, Natural History Museum, Cromwell Road, SW7 5BD London, UK

[4] University College London Center for Life's Origins and Evolution, London, UK

[5] Department of Invertebrate Zoology and Geology, California Academy of Sciences, San Francisco CA,

USA

[6] Bader International Study Centre, Queen's University, Herstmonceux Castle, East Sussex, UK

[7] Departamento de Bioquímica y Biología Molecular, Facultad de Ciencias Biológicas, Universidad de

Concepción, Concepción, Chile

[8] School of Zoology, Faculty of Life Sciences, Tel Aviv University, Tel Aviv, Israel

[9] Steinhardt Museum of Natural History, Tel-Aviv, Israel

[10] Department of Geology and Palaeontology, Natural History Museum Vienna, Vienna, Austria

* Corresponding author. Email: nicolas.mongiardinokoch@yale.edu

Abstract

Echinoids are key components of modern marine ecosystems. Despite a remarkable fossil record, the emergence of their crown group is documented by few specimens of unclear affinities, rendering their early history uncertain. The origin of sand dollars, one of its most distinctive clades, is also unclear due to an unstable phylogenetic context. We employ seventeen novel genomes and transcriptomes to build a dataset with a near-complete sampling of major lineages and use it to revise the phylogeny and divergence times of echinoids. We introduce the concept of a chronospace—a multidimensional representation of node ages—and use it to explore the effects of using alternative gene samples, models of molecular evolution, and clock priors. We find the choice of clock model to have the strongest impact on divergence times, while gene sampling and the use of site-heterogeneous models show little effects. Crown group echinoids originated in the Permian and diversified rapidly in the Triassic. We clarify the relationships among sand dollars and their close relatives, showing that *Apatopygus* represents a relict lineage with a deep Jurassic origin. Sand dollars are confidently dated to the Cretaceous, implying ghost ranges spanning approximately 50 million years, a remarkable discrepancy with their fossil record.

Keywords: Echinoidea, Sea urchins, Sand dollars, Phylogenomics, Time calibration, Site heterogenous models

Background

The fossil record represents the best source of primary data for constraining the origins of major lineages across the tree of life. However, the fossil record is not perfect, and even for groups with an excellent fossilization potential, constraining their age of origin can be difficult [1, 2]. Furthermore, as many traditional hypotheses of relationships have been revised in light of large-scale molecular datasets, the affinities of fossil lineages and their bearings on inferred times of divergence have also required a reassessment. An exemplary case of this is Echinoidea, a clade comprised of sea urchins, heart urchins, sand dollars, and allies, for which phylogenomic trees have questioned the timing of previously well-constrained nodes [3, 4].

Echinoids are easily recognized by their spine-covered skeletons or tests, composed of numerous tightly interlocking plates. Slightly over 1,000 living species have been described to date [5], a diversity that populates every marine benthic environment from intertidal to abyssal depths [6]. Echinoids are usually subdivided into two morpho-functional groups with similar species-level diversities: "regular" sea urchins, a paraphyletic assemblage of hemispherical, epibenthic consumers protected by large spines; and irregulars (Irregularia), a clade of predominantly infaunal and bilaterally symmetrical forms covered by small and specialized spines. In today's oceans, regular echinoids act as ecosystem engineers in biodiverse coastal communities such as coral reefs [7] and kelp forests [8], where they are often the main consumers. They are first well-known in the fossil record on either side of the Permian-Triassic (P-T) mass extinction event when many species occupied reef environments similar to those inhabited today by their descendants [9, 10]. This extinction event was originally thought to have radically impacted the macroevolutionary history of the clade, decimating the echinoid stem group and leading to the radiation of crown group taxa from a single surviving lineage [11, 12]. However, it is now widely accepted that the origin of crown group Echinoidea (i.e., the divergence between its two

main lineages, Cidaroidea and Euechinoidea) occurred in the Late Permian, as supported by molecular

estimates of divergence [13, 14], as well as the occurrence of Permian fossils with morphologies typical

of modern cidaroids [15, 16]. However, a recent total-evidence study recovered many taxa previously

classified as crown group members along the echinoid stem, while also suggesting that up to three

crown group lineages survived the P-T mass extinction [3]. This result increases the discrepancy between

molecular estimates and the fossil record and renders uncertain the early evolutionary history of crown

group echinoids. Constraining the timing of origin of this clade relative to the P-T mass extinction

(especially in the light of recent topological changes [3, 4]) is further complicated by the poor

preservation potential of stem group echinoids, and the difficulty assigning available disarticulated

remains from the Late Palaeozoic and Early Triassic to specific clades [11, 12, 17-20].

Compared to the morphological conservatism of regular sea urchins, the evolutionary history of

the relatively younger Irregularia was characterized by dramatic levels of morphological and ecological

innovation [21-24]. Within the diversity of irregulars, sand dollars are the most easily recognized (figure

1). The clade includes greatly flattened forms that live in high-energy sandy environments where they

feed using a unique mechanism for selecting and transporting organic particles to the mouth, where

these are crushed using well-developed jaws [25, 26]. Sand dollars (Scutelloida) were long thought to be

most closely related to sea biscuits (Clypeasteroida) given a wealth of shared morphological characters

[19, 25]. The extraordinary fossil record of both sand dollars and sea biscuits suggested their last

common ancestor originated in the early Cenozoic from among an assemblage known as "cassiduloids"

[23, 25], a once diverse group that is today represented by three depauperate lineages: cassidulids (and

close relatives), echinolampadids, and apatopygids [19, 27]. These taxa not only lack the defining

features of both scutelloids and clypeasteroids but have experienced little morphological change since

their origin deep in the Mesozoic [24, 27-29]. However, early molecular phylogenies supported both

cassidulids and echinolampadids as close relatives of sand dollars (e.g., [14, 30]), a topology initially

disregarded for its conflicts with both morphological and paleontological evidence, but later confirmed

using phylogenomic approaches [4]. While many of the traits shared by sand dollars and sea biscuits

have since been suggested to represent a mix of convergences and ancestral synapomorphies

secondarily lost by some "cassiduloids" [3, 4], the strong discrepancy between molecular topologies and

the fossil record remains unexplained. Central to this discussion is the position of apatopygids, a clade so

far unsampled in molecular studies. Apatopygids have a fossil record stretching more than 100 million

years and likely have phylogenetic affinities with even older extinct lineages [3, 19, 28, 29]. Although

current molecular topologies already imply ghost ranges for scutelloids and clypeasteroids that

necessarily extend beyond the Cretaceous-Paleogene (K-Pg) boundary, the phylogenetic position of

apatopygids could impose even earlier ages on these lineages (figure 1). Constraining these divergences

is necessary to understand the timing of origin of the sand dollars, one of the most specialized lineages

of echinoids [24-27]. Resolving some phylogenetic relationships within scutelloids has also been

complicated by their recurrent miniaturization and associated loss of morphological features (figure 1;

[25, 31, 32]).

Figure 1: Neognathostomate diversity and phylogenetic relationships. **A.** *Fellaster zelandiae*, North Island, New Zealand (Clypeasteroida). **B.** Large specimen: *Peronella japonica*, Ryukyu Islands, Japan; Small specimen: *Echinocyamus crispus*, Maricaban Island, Philippines (Laganina: Scutelloida). **C.** Large specimen: *Leodia sexiesperforata*, Long Key, Florida; Small specimen: *Sinaechinocyamus mai*, Taiwan (Scutellina: Scutelloida). **D.** *Rhyncholampas pacificus,* Isla Isabela, Galápagos Islands (Cassidulidae). **E.** *Conolampas sigsbei*, Bimini, Bahamas (Echinolampadidae). **F.** *Apatopygus recens*, Australia (Apatopygidae). **G.** Hypotheses of relationships among neognathostomates. Top: Morphology supports a clade of Clypeasteroida + Scutelloida originating after the K-Pg boundary, subtended by a paraphyletic assemblage of extant (red) and extinct (green) "cassiduloids" [19]. Bottom: A recent total-evidence study split most of cassiduloid diversity into a clade of extant lineages closely related to scutelloids, and an unrelated clade of extinct forms (Nucleolitoida) [3]. Divergence times are much older and conflict with fossil evidence. Cassidulids and apatopygids lacked molecular data in this analysis. Scale bars = 10 mm.

Echinoidea constitutes a model clade in developmental biology and genomics. As these fields embrace a more comparative approach [13, 33, 34], robust and time-calibrated phylogenies are expected to play an increasingly important role. Likewise, the extraordinary fossil record of echinoids and the ease with which echinoid fossils can be incorporated in phylogenetic analyses make them an ideal system to explore macroevolutionary dynamics using phylogenetic comparative methods [3, 31]. In this study, we build upon available molecular resources with 17 novel genome-scale datasets and build the largest molecular dataset for echinoids yet compiled. Our expanded phylogenomic dataset extends sampling to 16 of the 17 currently recognized echinoid orders (plus the unassigned apatopygids) [35], is the first to bracket the extant diversity of both sand dollars and sea biscuits, and include members of all three lineages of living "cassiduloids" (cassidulids, echinolampadids, and apatopygids). We also

125   incorporate a diverse sample of outgroups, providing access to the deepest nodes within the crown

126   groups of all other echinoderm classes (holothuroids, asteroids, ophiuroids, and crinoids). With it, we

127   reconstruct the phylogenetic relationships and divergence times of the major lineages of living echinoids

128   and place their diversification within the broader context of echinoderm evolution.

Methods

(a)   Sampling, bioinformatics, and matrix construction

This study builds upon previous phylogenomic matrices [3, 4], the last of which was augmented

through the addition of eight published datasets (mostly expanding outgroup sampling), as well as 17

novel echinoid datasets (15 transcriptomes and two draft genomes). Final taxon sampling included 12

outgroups and 50 echinoids. For all novel datasets, tissue sampling, DNA/RNA extraction, library

preparation, and sequencing varied by specimen, and are detailed in the supplementary information.

Raw reads for all transcriptomic datasets were trimmed or excluded using quality scores with

Trimmomatic v. 0.36 [36] under default parameters. Further sanitation steps were performed using the

Agalma 2.0 phylogenomic workflow [37], and datasets were assembled *de novo* with Trinity 2.5.1 [38].

For genomic shotgun sequences, adapters were removed with BBDuk

(https://sourceforge.net/projects/bbmap/), and UrQt v. 1.0.18 [39] was used to filter short reads (size <

50) and trim low-quality ends (score < 28). Datasets were then assembled using MEGAHIT v. 1.1.2 [40].

Draft genomes were masked using RepeatMasker v. 4.1.0 [41, 42], before obtaining gene predictions

with AUGUSTUS 3.2.3 [43]. A custom set of universal single-copy orthologs (USCOs) obtained from the

latest *Strongylocentrotus purpuratus* genome assembly (Spur v. 5.0) was employed as the training

dataset. Further details of these analyses, as well as statistics for raw reads and assemblies can be found

in tables S1 and S2.

Multiplexed transcriptomes were sanitized from cross-contaminants using CroCo v. 1.1 [44], and

likely non-metazoan contaminants were removed using alien_index v. 3.0 [45] (removing sequences

with AI values > 45). Datasets were imported back into Agalma, which automated orthology inference

(as described in [37, 46]), gene alignment with MACSE [47], and trimming with GBLOCKS [48]. The amino

acid supermatrix was reduced using a 70% occupancy threshold, producing a final dataset of 1,346 loci

(327,695 sites). As a final sanitation step, gene trees were obtained using ParGenes v. 1.0.1 [49], which

performed model selection (minimizing the Bayesian Information Criterion) and inference using 100

bootstrap (BS) replicates. Trees were then used to remove outlier sequences with TreeShrink v. 1.3.1

[50]. We specified a reduced tolerance for false positives and limited removal to at most three terminals

which had to increase tree diameter by at least 25% (-q 0.01 -k 3 -b 25).

(b)   Phylogenetic inference

Inference was performed under multiple probabilistic and coalescent-aware methods, known to

differ in their susceptibility to model violations. Coalescent-based inference was performed using the

summary method ASTRAL-III [51], estimating support as local posterior probabilities [52]. Among

concatenation approaches, we used Bayesian inference under an unpartitioned GTR+G model in

ExaBayes 1.5 [53]. Two chains were run for 2.5 million generations, samples were drawn every one

hundred, and the initial 25% was discarded as burn-in. We also explored maximum likelihood inference

with partitioned and unpartitioned models. For the former, the fast-relaxed clustering algorithm was

used to find the best-fitting model among the top 10% using IQ-TREE 1.6.12 [54, 55] (-m MFP+MERGE -

rclusterf 10 -rcluster-max 3000), and support was evaluated with 1,000 ultrafast BS replicates [56]. For

the latter, we used the LG4X+R model in RAxML-NG v. 0.5.1 [57] and evaluated support with 200

replicates of BS. Finally, we also implemented the site-heterogenous LG+C60+F+G mixture model using

the posterior mean site frequency (PMSF) approach to provide a fast approximation of the full profile

mixture model [58], allowing the use of 100 BS replicates to estimate support. Given some degree of

topological conflict between the results of the other methods (see below), multiple guide trees were

used to estimate site frequency profiles, but the resulting phylogenies were identical.

Given conflicts between methods in the resolution of one particular node (involving the

relationships among Arbacioida, Salenioida, and the clade of Stomopneustoida + Camarodonta), all

methods were repeated after reducing the matrix to 500 and 100 loci selected for their phylogenetic

usefulness using the approach described in [3, 59] and implemented in the *genesortR* script

(https://github.com/mongiardino/genesortR). This approach relies on seven gene properties routinely

used for phylogenomic subsampling, including multiple proxies for phylogenetic signal—such as the

average BS and Robinson-Foulds (RF) similarity to a target topology—as well as several potential sources

of systematic bias (e.g., rate and compositional heterogeneities). Outgroups were removed before

calculating these metrics. RF similarity was measured to a species tree that had the conflicting

relationship collapsed so as not to bias gene selection in favor of any resolution. A principal component

analysis (PCA) of this dataset resulted in a dimension (PC 2, 17.6% of variance) along which phylogenetic

signal increased while sources of bias decreased (figure S1), and which was used for loci selection. For

the smallest dataset, we also performed inference under the site-heterogeneous CAT+GTR+G model

using PhyloBayes-MPI [60]. Three runs were continued for > 10,000 generations, sampling every two

generations and discarding the initial 25%. Convergence was confirmed given a maximum bipartition

discrepancy of 0.067 and effective sample sizes for all parameters > 150.

Two other approaches were implemented in order to assist in resolving the contentious node.

First, we implemented a likelihood-mapping analysis [61] in IQ-TREE to visualize the phylogenetic signal

for alternative resolutions of the quartet involving these three lineages (Arbacioida, Salenioida, and

Stomopneustoida + Camarodonta) and their sister clade (Irregularia; other taxa were excluded). Second,

we estimated the log-likelihood scores of each site in RAxML (using best-fitting models) for the two most

strongly supported resolutions found through likelihood mapping. These were used to calculate gene-

wise differences in scores, or δ values [62]. In order to search for discernable trends in the signal for

alternative topologies, genes were ordered based on their phylogenetic usefulness (see above) and the

mean per-locus δ values of datasets composed of multiples of 20 loci (i.e., the most useful 20, 40, etc.)

were calculated.

(c)   Time calibration

Node dating was performed using relaxed molecular clocks in PhyloBayes v4.1 using a fixed

topology and a novel set of 22 fossil calibrations corresponding to nodes from our newly inferred

phylogeny (listed in the supplementary information). Depending on the node, we enforced both

minimum and maximum bounds, or either one of these. A birth-death prior was used for divergence

times, which allowed for the implementation of soft bounds [63], leaving 5% prior probability of

divergences falling outside of the specified interval. We explored the sensitivity of divergence time

estimates to gene selection, model of molecular evolution, and clock prior. One hundred loci were

sampled from the full supermatrix according to four different sampling schemes: usefulness (calculated

as explained above, except incorporating all echinoderm terminals), phylogenetic signal (i.e., smallest RF

distance to species tree), clock-likeness (i.e., smallest variance of root-to-tip distances), and level of

occupancy. For clock-likeness, we only considered loci that lay within one standard deviation of the

mean rate (estimated by dividing total tree length by the number of terminals [64]), as this method is

prone to select largely uninformative loci otherwise (figure S2; [59]). A fifth sample of randomly selected

loci was also evaluated. These five datasets were run under two models of molecular evolution, the site-

homogenous GTR+G and the site-heterogenous CAT+GTR+G, and both uncorrelated gamma (UGAM)

and log-normal autocorrelated (LN) clocks were implemented. The combination of these settings (loci

sampled, model of evolution, and clock prior) resulted in 20 analyses. For each, two runs were

continued for 20,000 generations, after which the initial 25% was discarded and the chains thinned to

every two generations (see log-likelihood trace plots in figure S3). To explore the sensitivity of

divergence times to these methodological decisions, 500 random chronograms were sampled from each

analysis (250 from each run), and their node dates were subjected to multivariate analysis of variance

(MANOVA) and between-group PCA (bgPCA) using package *Morpho* [65] in the R statistical environment

[66]. bgPCA involves the use of PCA on the covariance matrix of group means (e.g., the mean ages of all

nodes obtained under UGAM or LN clocks), followed by the projection of original samples onto the

obtained bgPC axes. The result is a multidimensional representation of divergence times—a

chronospace—rotated so as to capture the distinctiveness of observations obtained under different

settings. Separate bgPCAs were performed for loci sampling strategy, clock prior and model of molecular

evolution, and the proportion of total variance explained by bgPC axes was interpreted as an estimate of

the relative impact of these choices on inferred times of divergence. Finally, lineage-through-time plots

were generated using *ape* [67].

Results

Relationships inferred from the full dataset were remarkably stable, with all nodes but one

being identically resolved and fully supported across inference methods (figure 2A). While recovering a

topology similar to those of previous molecular studies [3, 4, 13, 14, 30, 68], this analysis is the first to

sample and confidently place micropygoids and aspidodiadematoids within Aulodonta, as well as resolve

the relationships among all major clades of Neognathostomata (scutelloids, clypeateroids and the three

lineages of extant "cassiduloids"). Our results show that *Apatopygus recens* is not related to the

remaining "cassiduloids" but is instead the sister clade to all other sampled neognathostomates. The

strong support for this placement, as well as for a clade of cassidulids and echinolampadids

(Cassiduloida *sensu stricto*) as the sister group to sand dollars, provides a basis for an otherwise elusive

phylogenetic classification of neognathostomates. Our topology also confirms that *Sinaechinocyamus*

mai, a miniaturized species once considered a plesiomorphic member of Scutelloida based on the

reduction or loss of diagnostic features (figure 1), is in fact a derived paedomorphic lineage closely

related to *Scaphechinus mirabilis* [32].

Figure 2: Phylogenetic relationships among major clades of Echinoidea. **A.** Favored topology. With the exception of a single contentious node within Echinacea (marked with a star; see Discussion), all methods supported the same pattern of relationships. Numbers below major clades correspond to the current numbers of described living species (obtained from [5]). **B.** Likelihood-mapping analysis showing the proportion of quartets supporting different resolutions within Echinacea. While the majority of quartets support the topology depicted in **A** (shown in red), a significant number support an alternative resolution that has been recovered in morphological analyses [19] (shown in blue). **C.** Support for a

clade of Salenioida + (Camarodonta + Stomopneustoida), as depicted in **A**, is predominantly

concentrated among the most phylogenetically useful loci. This signal is attenuated in larger datasets

that contain less reliable genes, eventually favoring an alternative resolution. Only the 584 loci

containing data for the three main lineages of Echinacea were considered. The line corresponds to a

second-degree polynomial regression. **D.** Resolution and bootstrap scores (color coded) of the topology

within Echinacea found using datasets of different sizes and alternative methods of inference.

Salenioida is another major lineage sampled here for the first time, and whose exact position

among regular echinoids proved difficult to resolve. While some methods supported salenioids as the

sister group to a clade of camarodonts, stomopneustoids, and arbacioids (a topology previously

supported by morphology [19]), others recovered a closer relationship of salenioids to Camarodonta +

Stomopneustoida, with arbacioids sister to them all. These results do not stem from a lack of

phylogenetic signal, but rather from the presence of strong and conflicting evidence in the dataset

regarding the position of salenioids (figure 2B). However, a careful dissection of these signals shows that

loci with high phylogenetic usefulness favor the topology shown in figure 2A, with the morphological

hypothesis becoming dominant only after incorporating less reliable loci (figure 2C). In line with these

results, moderate levels of gene subsampling (down to 500 loci) unambiguously support the placement

of arbacioids as sister to the remaining taxa regardless of the chosen method of inference (figure 2D).

More extreme subsampling (down to 100 loci) again results in disagreement among methods. This

possibly stems from the increasing effect of stochastic errors in smaller datasets, as less than half of the

sampled loci in these reduced datasets contain data for all branches of this quartet (see figure 2C). This

result shows the importance of ensuring that datasets (especially subsampled ones) retain appropriate

levels of occupancy for clades bracketing contentious nodes [69]. Despite these disagreements, several

lines of evidence favour the topology shown in figure 2A, including the results of likelihood mapping,

and the increased support for this resolution among the most phylogenetically useful loci and when

using more complex methods of reconstruction, such as partitioned and site-heterogenous models,

which favour this topology regardless of dataset size (figure 2D).

While alternative methods of inference had minor effects on phylogenetic relationships, they

did impact the reconstruction of branch lengths (figure S4). Site-heterogenous models (such as

CAT+GTR+G) returned longer branch lengths overall, but also uncovered a larger degree of molecular

change among echinoderm classes. Branches connecting these clades were stretched to a much larger

extent than those within the ingroup (figure S4), which might affect the inference of node dates. We

tested this hypothesis by exploring the sensitivity of divergence times to the use of alternative models of

molecular evolution, as well as different gene sampling strategies and clock priors, all of which

significantly affected inferred divergence times (MANOVA; all $P < 2.2 \times 10^{-16}$). While nodes connecting

some outgroup taxa were among the nodes most sensitive to these decisions, large effects were also

seen among nodes relating to the origin and diversification of the echinoid clades Cidaroidea and

Aulodonta (figure 3A). All of these nodes varied in age by more than 60 Ma—and up to 115 Ma—among

the consensus topologies of different analyses (figure 3A).

Figure 3: Sensitivity of divergence time estimation to methodological decisions. **A.** The ten most

sensitive ages are found within Cidaroidea, Aulodonta, and among outgroup nodes. For each, the range

shown corresponds to the ages found among the consensus topologies of the 40 time-calibrated runs. **B-**

C. Scores for the between-group PCA (bgPCA) separating chronograms based on the clock model (**B**) or

the model of molecular evolution (**C**) employed. While the first factor has a major impact on inferred

ages, the second one has only minor effects. **D-E.** Change in branch lengths along the bgPCA axis defined

by discriminating based on clock model (**D**) or model of molecular evolution (**E**). Branches are colored

according to their contraction/expansion given a score of − 1 (top) or + 1 (bottom) standard deviations

along the bgPCA axis relative to the mean. Clades/nodes showing age differences larger than 20 Ma are

labelled (none in the case of **E**).

In order to isolate and visualize the impact of these factors on reconstructed ages, their major

effects were extracted with the use of bgPCAs (figures 3B-E, S5). This revealed that the single dimension

of chronospace maximizing the distinctiveness of trees obtained under different clock models explained

53.7% of the total variance in node ages across all analyses. In contrast, the choice of different loci or

models of molecular evolution showed a much lesser effect, explaining only 10.5% and 4.0% of the total

variance, respectively. Even though all of these decisions affected a similar set of sensitive nodes (those

mentioned above, as well as some relationships within Atelostomata; figures 3B-E, S5), the choice of

clock model modified the ages of 17 of these by more than 20 Ma (figure S6), a magnitude of change

that was not induced by either alternative loci or models of evolution on any node (figures S7-S8).

Even when the age of crown Echinodermata was constrained to postdate the appearance of

stereom (the characteristic skeletal microstructure of echinoderms) in the Early Cambrian [70, 71], only

analyses using the most clock-like loci recovered ages concordant with this (i.e., median ages younger

than the calibration enforced; figure S9). Instead, most consensus trees favored markedly older ages for

the clade, in some cases even predating the origin of the Ediacaran biota [72] (figures 3A, 4, S7). Despite

the relative sensitivity of many of the earliest nodes to methodological choices (figure 3), the split

between Crinoidea and all other echinoderms (Eleutherozoa) is always inferred to have predated the

end of the Cambrian (youngest median age = 492.2 Ma), and the divergence among the other major

lineages (classes) of extant echinoderms are constrained to have happened between the Late Cambrian

and Middle Ordovician (see figure S9). Our results also recover an early origin of crown group

Holothuroidea (sea cucumbers; range of median ages = 351.6–383.2 Ma), well before the crown groups

of other extant echinoderm classes. These dates markedly postdate the first records of holothuroid

calcareous rings in the fossil record [73, 74], and imply that this trait does not define the holothuroid

crown group but instead evolved from an echinoid-like jaw-apparatus along its stem [75]. The other

noteworthy disagreement between these results and those of previous studies [76] involves dating

crown group Crinoidea to times that precede the P-T mass extinction (range of median ages = 268.0–

327.7 Ma, although highest posterior density intervals are always wide and include Triassic ages).

Figure 4: Divergence times among major clades of Echinoidea and other echinoderms. **A.** Consensus

chronogram of the two runs using clock-like genes under a CAT+GTR+G model of molecular evolution

and a LN clock prior. Node ages correspond to median values, and bars show the 95% highest posterior

density intervals. **B.** Lineage-through-time plot, showing the rapid divergence of higher-level clades

following the P-T mass extinction (shown with dashed lines, along with the K-Pg boundary). Each line

corresponds to a consensus topology from among the 40 time-calibrated runs performed. **C.** Posterior

distributions of the ages of selected nodes (marked in the chronogram). The effects of models of

molecular evolution are not shown, as they represent the least important factor (see figure 3);

distributions using site-homogenous and heterogenous models are combined for every combination of

targeted loci and clock prior.

Across all of the analyses performed, the echinoid crown group is found to have originated

somewhere between the Pennsylvanian and Cisuralian, with 42% posterior probability falling within the

late Carboniferous and 58% within the early Permian (figure 4C). An origin of the clade postdating the P-

T mass extinction is never recovered, even when such ages were allowed under the joint prior (figure

S10). While the posterior distribution of ages for Euechinoidea spans both sides of the P-T boundary, the

remaining earliest splits within the echinoid tree are constrained to have occurred during the Triassic,

including the origins of Aulodonta, Carinacea, Echinacea, and Irregularia (figures 4, S9). Many echinoid

orders are also inferred to have diverged from their respective sister clades during this period, including

aspidodiadematoids, pedinoids, echinothurioids, arbacioids, and salenioids. LTT plots confirm that

lineage diversification proceeded rapidly throughout the Triassic (figure 4B). Despite the topological

reorganization of Neognathostomata, the clade is dated to a relatively narrow time interval in the Late

to Middle Jurassic (range of median ages = 169.48-179.96 Ma), in agreement with recent estimates [3].

Within this clade, the origins of both scutelloids and clypeasteroids confidently predate the K-Pg mass

extinction (posterior probability of origination before the boundary = 0.9995 and 0.9621, respectively),

despite younger ages being allowed by the joint prior (figure S10).

Discussion

(a)   The echinoid Tree of Life

In agreement with previous phylogenomic studies [3, 4], echinoid diversity can be subdivided

into five major clades (figure 2A). Cidaroids form the sister group to all other crown group echinoids

(Euechinoidea). Some aspects of the relationships among sampled cidaroids are consistent with previous

molecular [77] and morphological studies [19], including an initial split between *Histocidaris* and the

remaining taxa, representing the two main branches of extant cidaroids [5, 35]. Others, such as the

nested position of *Prionocidaris baculosa* within the genus *Eucidaris,* not only implies paraphyly of this

genus but also suggests the need for a taxonomic reorganization of the family Cidaridae. Within

euechinoids, the monophyly of Aulodonta is supported for the first time with sampling of all of its major

groups. The subdivision of these into a clade that includes diadematoids plus micropygoids (which we

propose should retain the name Diadematacea), sister to a clade including echinothurioids and

pedinoids (Echinothuriacea *sensu* [4]) is strongly reminiscent of some early classifications (e.g., [78]).

Our expanded phylogenomic sampling also confirms an aulodont affinity for aspidodiadematoids [3, 35],

and places them within Echinothuriacea as the sister group to Pedinoida.

The remaining diversity of echinoids, which forms the clade Carinacea (figure 2), is subdivided

into Irregularia and their sister clade among regulars, for which we amend the name Echinacea to

include Salenioida. Given the striking morphological gap separating regular and irregular echinoids, the

origin of Irregularia has been shrouded in mystery [19, 23, 78]. Our complete sampling of major regular

lineages determines Echinacea *sensu stricto* to be the sister clade to irregular echinoids. A monophyletic

Echinacea was also supported in a recent total-evidence analysis [3], but the incomplete molecular

sampling of that study resulted in a slightly different topology that placed salenioids as the sister group

to the remaining lineages. However, an overall lack of morphological synapomorphies uniting these

clades had previously been acknowledged [19]. While the relationships within Echinacea proved to be

difficult to resolve even with thousands of loci, multiple lines of evidence lead us to prefer a topology in

which salenioids form a clade with camarodonts + stomopneustoids, with arbacioids sister to all of these

(figure 2).

As has been already established [3, 4, 14, 19, 30], the lineages of irregular echinoids here

sampled are subdivided into Atelostomata (heart urchins and allies) and Neognathostomata (sand

dollars, sea biscuits, and "cassiduloids"). Despite the former being the most diverse of the five main

clades of echinoids (figure 2), its representation in phylogenomic studies remains low, and its internal

phylogeny poorly constrained [35]. On the contrary, recent molecular studies have greatly improved our

understanding of the relationships among neognathosomates [3, 4, 68], revealing an evolutionary

history that dramatically departs from previous conceptions. Even when scutelloids and clypeasteroids

were never recovered as reciprocal sister lineages by molecular phylogenies (e.g., [13, 14, 30]), this

result was not fully accepted until phylogenomic data confidently placed echinolampadids as the sister

lineage to sand dollars [4]. At the same time, this result also rendered the position of the remaining

"cassiduloids", a taxonomic wastebasket with an already complicated history of classification [19, 27, 28,

79], entirely uncertain. An attempt to constrain the position of these using a total-evidence approach [3]

subdivided the "cassiduloids" into three unrelated clades: Nucleolitoida, composed of extinct lineages

and placed outside the node defined by Scutelloida + Clypeasteroida, and two other clades nested

within it (see figure 1). Extant "cassiduloids" were recovered as members of one of the latter clades,

representing the monophyletic sister group to sand dollars. Here, we show that *Apatopygus recens* does

not belong within this clade but is instead the sister group to all other extant neognathostomates. Given

this phylogenetic position, as well as the morphological similarities between *Apatopygus* and the

entirely extinct nucleolitids [19, 28, 79-81], it is likely that the three extant species of apatopygids

represent the last surviving remnants of Nucleolitoida, a clade of otherwise predominantly Mesozoic

neognathostomates [3]. Because of the renewed importance in recognizing this topology, we propose

the name Luminacea for the clade uniting all extant neognathostomates with the exclusion of

Apatopygus (figure 2). This nomenclature refers to the dynamic evolutionary history of the Aristotle's

lantern (i.e., the echinoid jaw-apparatus) within the clade (present in the adults of both clypeasteroids

and scutelloids, but found only in the juveniles of Cassiduloida *sensu stricto*), the inclusion of the so-

called lamp urchins (echinolampadids) within the clade, and the illumination provided by this hitherto

unexpected topology. The previous misplacement of *Apatopygus* within this clade ([3]; see figure 1G) is

likely a consequence of tip-dating preferring more stratigraphically congruent topologies [82], an effect

that can incorrectly resolve taxa on long terminal branches [83]. Given the generally useful phylogenetic

signal of stratigraphic information [84], this inaccuracy further highlights the unusual evolutionary

history of living apatopygids.

(b)   Echinoid diversification

Calibrating phylogenies to absolute time is crucial to understanding evolutionary history, as the

resulting chronograms provide a major avenue for testing hypotheses of diversification. However, the

accuracy and precision of the inferred divergence times hinge upon many methodological choices that

are often difficult or time-consuming to justify (e.g., calibration strategies, prior distributions on node

ages, clock models, etc.) [85-90]. Understanding the impact of each of these sources of uncertainty can

be difficult. Here we explore the sensitivity of node ages to different ways of modelling heterogeneities

among lineages in the rates and patterns of molecular evolution, as well as alternative criteria to sample

molecular loci. To do so, we introduce an approach to visualize and measure the overall effect of these

decisions on inferred dates. Our results reveal that divergence times obtained under site-homogenous

and site-heterogenous models (such as CAT+GTR+G) are broadly comparable. This happens despite

these types of models estimating higher levels of sequence divergence and stretching branches in a non-

isometric manner (figure S3). A similar result was recently found when comparing site-homogenous

models with different numbers of parameters [91], suggesting that relaxed clocks adjust branch rates in

a manner that buffers the effects introduced by using more complex (and computationally-intensive)

models of evolution. Similarly, the choice between different loci also has a small effect on inferred ages,

with little evidence of a systematic difference between the divergence times supported by randomly-

chosen loci and those found using targeted sampling criteria, such as selecting genes for their

phylogenetic signal, usefulness, occupancy, or clock-likeness. A meaningful effect was restricted to a few

ancient nodes (e.g., Echinodermata), for which clock-like genes suggested younger ages that are more

consistent with fossil evidence. While this validates the use of clock-like genes for inferring deep

histories of diversification, the choice of loci had no meaningful effect on ingroup ages. Finally, the

choice between alternative clock models induced differences in ages that were five to ten times

stronger than those of other factors, emphasizing the importance of either validating their choice (e.g.,

[92]) or—as done here—focusing on results that are robust to them.

The origin and early diversification of crown group Echinoidea have always been considered to

have been determined (or at least strongly affected) by the P-T mass extinction [11, 12, 16, 20].

However, estimating the number of crown group members surviving the most severe biodiversity crisis

in the Phanerozoic [93] has been hampered by both paleontological and phylogenetic uncertainties [3,

10, 13-15, 18]. Our results establish with confidence that multiple crown group lineages survived and

crossed this boundary, finding for the first time a null posterior probability of the clade originating after

the extinction event. While the survival of three crown group lineages is slightly favored (figure S11),

discerning between alternative scenarios is still precluded by uncertainties in dating these early

divergences. Echinoid diversification during the Triassic was relatively fast (figure 4B) and involved rapid

divergence among its major clades. Even many lineages presently classified at the ordinal level trace

their origins to this initial pulse of diversification following the P-T mass extinction.

The late Palaeozic and Triassic origins inferred for the crown group and many euechinoid orders

prompts a re-evaluation of fossils from this interval of time. Incompletely known fossil taxa such as the

Pennsylvanian *?Eotiaris meurevillensis*, with an overall morphology akin to that of crown group

echinoids, has a stratigraphic range consistent with our inferred date for the origin of the echinoid

crown group [20]. Additionally, the Triassic fossil record of echinoids has been considered to be

dominated by stem group cidaroids, with the first euechinoids not known until the Late Triassic [17, 94].

However, the Triassic origins of many euechinoid lineages supported by our analyses necessitates that

potential euechinoid affinities should be re-considered for this diversity of Triassic fossils. This is

especially the case for the serpianotiarids and triadocidarids, abundant Triassic families variously

interpreted as cidaroids, euechinoids, or even stem echinoids [3, 17, 94, 95]. A reinterpretation of any of

these as stem euechinoids would suggest that the long-implied gap in the euechinoid record [16, 18] is

caused by our inability to correctly place these key fossils, as opposed to an incompleteness of the fossil

record itself.

While our phylogenomic approach is the first to resolve the position of all major cassiduloid

lineages, the inferred ages for many nodes within Neognathostomata remain in strong disagreement

with the fossil record. No Mesozoic fossil can be unambiguously assigned to either sand dollars or sea

biscuits, a surprising situation given the good fossilization potential and highly distinctive morphology of

these clades [19, 21, 25, 96]. While molecular support for a sister group relationship between scutelloids

and echinolampadids already implied this clade (Echinolampadacea) must have split from clypeasteroids

by the Late Cretaceous [3, 4, 14, 19], this still left open the possibility that the crown groups of sand

dollars and sea biscuits radiated in the Cenozoic. Under this scenario, the Mesozoic history of these

groups could have been comprised of forms lacking their distinctive morphological features,

complicating their correct identification. This hypothesis is here rejected, with the data unambiguously

supporting the radiation of both crown groups preceding the K-Pg mass extinction (figure 4C). While it

remains possible that these results are incorrect even after such a thorough exploration of the time-

calibration toolkit (see for example [90, 97]), these findings also call for a critical reassessment of the

Cretaceous fossil record, and a better understanding of the timing and pattern of morphological

evolution among fossil and extant neognathostomates. For example, isolated teeth with an overall

resemblance to those of modern sand dollars and sea biscuits have been found in Lower Cretaceous

deposits [98], raising the possibility that other overlooked and disarticulated remains might close the

gap between rocks and clocks.

Conclusions

Although echinoid phylogenetics has long been studied using morphological data, the position

of several major lineages (e.g., aspidodiadematoids, micropygoids, salenioids, apatopygids) remained to

be confirmed with the use of phylogenomic approaches. Our work not only greatly expands the

available genomic resources for the clade, but finds novel resolutions for some of these lineages,

improving our understanding of their evolutionary history. The most salient aspect of our topology is the

splitting of the extant "cassiduloids" into two distantly related clades, one of which is composed

exclusively of apatopygids. This result is crucial to constrain the ancestral traits shared by the main

lineages of neognathostomates, helping unravel the evolutionary processes that gave rise to the unique

morphology of the sand dollars and sea biscuits [3, 24, 25].

Although divergence time estimation is known to be sensitive to many methodological

decisions, systematically quantifying the relative impact of these on inferred ages has rarely been done.

Here we propose an approach based on chronospaces (implemented in R and available at

https://github.com/mongiardino/chronospace) that can help visualize key effects and determine the

sensitivity of node dates to different assumptions. Our results shed new light on the early evolutionary

history of crown group echinoids and its relationship with the P-T mass extinction event, a point in time

where the fossil record provides ambiguous answers. They also establish with confidence a Cretaceous

origin for the sand dollars and sea biscuits, preceding their first appearance in the fossil record by at

least 40 to 50 Ma, respectively (and potentially up to 65 Ma). These clades, therefore, join several well-

established cases of discrepancies between the fossil record and molecular clocks, such as those

underlying the origins of placental mammals [99] and flowering plants [100].

Author contributions

514   NMK, RM, and GWR conceived and designed the study. NMK, ASH, MFM, AFA, SEC, FA, OB, and

515   AK performed extractions, prepared libraries, and sequenced transcriptomic/genomic datasets. NMK

516   and AK generated *de novo* assemblies. JRT and RM designed fossil calibrations. NMK run all phylogenetic

517   analyses. NMK and JRT wrote the manuscript with input from all other authors. All authors gave final

518   approval for publication and agree to be held accountable for the work performed therein.

Data, code and materials

521   Raw transcriptomic reads have been deposited in NCBI under Bioproject accession number XXX.

522   Assemblies are available from the authors upon request. R code to plot chronospaces and explore the

523   effects of methodological decisions on divergence time estimation can be found at

524   https://github.com/mongiardino/chronospace.

Competing interests

We declare no competing interests.

Acknowledgments

We are grateful to the crew of R/Vs Antéa, Atlantis, Falkor and Mirai, HOV Alvin, and the crew and PIs of BIOMAGLO deep-sea cruises (Laure Corbari, Karine Olu-Le Roy, Sarah Samadi) for assistance in specimen collection. We would also like to thank Libby Liggins, Wilma Blom, Owen Anderson, Francisco Solís-Marín, Carlos A. Conejeros-Vargas, and Jih-Pai Lin for the collection and shipment of specimens. We gratefully acknowledge assistance from Thomas Saucède (Univ. Burgundy), Marc Eléaume (MNHN), and Charlotte Seid (Scripps) in accessing, cataloging and processing museum specimens. Tim Ravasi provided resources and collection facilities to GWR via King Abdullah University of Science and Technology (KAUST). Institutional support was provided by the Central Research Laboratories and the Department of Geology and Palaeontology at the Natural History Museum Vienna, Austria. This work was supported by a Yale Institute for Biospheric Studies Doctoral Dissertation Improvement Grant and a Society of Systematic Biologists Graduate Student Research Award to NMK. AK received funding from an Austrian Science Fund project (FWF, project number P 29508-B25), FA from Agencia Nacional de Investigación (ANID/PAI Inserción en la Academia, project number PAI79170033). JRT was supported by a Royal Society Newton International Fellowship and a Leverhulme Trust Early Career Fellowship. NMK was supported by a Yale University fellowship.

References

[1] Donoghue PC, Benton MJ. 2007 Rocks and clocks: calibrating the Tree of Life using fossils and

molecules. *Trends Ecol. Evol.* **22**, 424–431.

[2] Smith AB, Peterson KJ. 2002 Dating the time of origin of major clades: molecular clocks and the fossil

record. *Annu. Rev. Earth Planet. Sci.* **30**, 65–88.

[3] Mongiardino Koch N, Thompson JR. 2021 A Total-Evidence Dated Phylogeny of Echinoidea Combining

Phylogenomic and Paleontological Data. *Syst. Biol.* **70**, 421–439.

[4] Mongiardino Koch N, Coppard SE, Lessios HA, Briggs DEG, Mooi R, Rouse GW. 2018 A phylogenomic

resolution of the sea urchin tree of life. *BMC Evol. Biol.* **18**, 189.

[5] Kroh A, Mooi R. 2020 World Echinoidea Database. Accessed at

http://www.marinespecies.org/echinoidea on 2021-07-13. (doi:10.14284/355)

[6] Schultz HA. 2015 *Echinoidea: with pentameral symmetry*. Walter de Gruyter GmbH & Co KG.

[7] Edmunds PJ, Carpenter RC. 2001 Recovery of *Diadema antillarum* reduces macroalgal cover and

increases abundance of juvenile corals on a Caribbean reef. *Proc. Natl. Acad. Sci.* **98**, 5067–5071.

[8] Harrold C, Pearse JS. 1987 The ecological role of echinoderms in kelp forests. *Echinoderm Studies* **2**,

137–233.

[9] Zonneveld JP, Furlong CM, Sanders SC. 2016 Triassic echinoids (Echinodermata) from the Aksala

Formation, north Lake Laberge, Yukon Territory, Canada. *Pap. Palaeontol.* **2**, 87–100.

[10] Thompson JR, Petsios E, Bottjer DJ. 2017 A diverse assemblage of Permian echinoids

(Echinodermata, Echinoidea) and implications for character evolution in early crown group echinoids. *J.*

Paleontol. **91**, 767–780.

[11] Kier PM. 1977 Triassic echinoids. *Smithson. Contrib. Paleobiol.* **30**, 1–88.

[12] Twitchett RJ, Oji T. 2005 Early Triassic recovery of echinoderms. *C. R. Palevol* **4**, 531–542.

[13] Thompson JR, Erkenbrack EM, Hinman VF, McCauley BS, Petsios E, Bottjer DJ. 2017 Paleogenomics

of echinoids reveals an ancient origin for the double-negative specification of micromeres in sea urchins.

Proc. Natl. Acad. Sci. **114**, 5870–5877.

[14] Smith AB, Pisani D, Mackenzie-Dodds JA, Stockley B, Webster BL, Littlewood DTJ. 2006 Testing the

molecular clock: molecular and paleontological estimates of divergence times in the Echinoidea

(Echinodermata). *Mol. Biol. Evol.* **23**, 1832–1851.

[15] Smith AB, Hollingworth N. 1990 Tooth structure and phylogeny of the Upper Permian echinoid

Miocidaris keyserlingi. *Proc. Yorkshire Geol. Soc.* **48**, 47–60.

[16] Thompson JR, Petsios E, Davidson EH, Erkenbrack EM, Gao F, Bottjer DJ. 2015 Reorganization of sea

urchin gene regulatory networks at least 268 million years ago as revealed by oldest fossil cidaroid

echinoid. *Sci. Rep.* **5**, 15541.

[17] Smith AB. 2007 Intrinsic versus extrinsic biases in the fossil record: contrasting the fossil record of

echinoids in the Triassic and early Jurassic using sampling data, phylogenetic analysis, and molecular

clocks. *Paleobiology* **33**, 310–323.

[18] Thompson JR, Hu S-X, Zhang Q-Y, Petsios E, Cotton LJ, Huang J-Y, Zhou C-Y, Wen W, Bottjer DJ. 2018

A new stem group echinoid from the Triassic of China leads to a revised macroevolutionary history of

echinoids during the end-Permian mass extinction. *R. Soc. Open Sci.* **5**, 171548.

[19] Kroh A, Smith AB. 2010 The phylogeny and classification of post-Palaeozoic echinoids. *J. Syst.*

Palaeontol. **8**, 147–212.

[20] Thompson JR, Mirantsev GV, Petsios E, Bottjer DJ. 2020 Phylogenetic analysis of the

Archaeocidaridae and Palaeozoic Miocidaridae (Echinodermata, Echinoidea) and the origin of crown

group echinoids. *Pap. Palaeontol.* **6**, 217–249.

[21] Kier PM. 1982 Rapid evolution in echinoids. *Palaeontology* **25**, 1–9.

[22] Barras CG. 2008 Morphological innovation associated with the expansion of atelostomate irregular

echinoids into fine-grained sediments during the Jurassic. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* **263**,

44–57.

[23] Saucède T, Mooi R, David B. 2007 Phylogeny and origin of Jurassic irregular echinoids

(Echinodermata: Echinoidea). *Geol. Mag.* **144**, 333–359.

[24] Hopkins MJ, Smith AB. 2015 Dynamic evolutionary change in post-Paleozoic echinoids and the

importance of scale when interpreting changes in rates of evolution. *Proc. Natl. Acad. Sci.* **112**, 3758–

3763.

[25] Mooi R. 1990 Paedomorphosis, Aristotle's lantern, and the origin of the sand dollars

(Echinodermata: Clypeasteroida). *Paleobiology* **16**, 25–48.

[26] Nebelsick JH. 2020 Ecology of clypeasteroids. In: *Sea Urchins: Biology and Ecology, 4th edition* (ed.

JM Lawrence), pp. 315–331. Cambridge, Academic Press.

[27] Smith AB. 2001 Probing the cassiduloid origins of clypeasteroid echinoids using stratigraphically

restricted parsimony analysis. *Paleobiology* **27**, 392–404.

[28] Souto C, Mooi R, Martins L, Menegola C, Marshall CR. 2019 Homoplasy and extinction: the

phylogeny of cassidulid echinoids (Echinodermata). *Zool. J. Linn. Soc.* **187**, 622–660.

[29] Kier PM. 1962 Revision of the cassiduloid echinoids. *Smithson. Misc. Collect.* **144**, 1–262.

[30] Littlewood DTJ, Smith AB. 1995 A combined morphological and molecular phylogeny for sea urchins

(Echinoidea: Echinodermata). *Philos. Trans. R. Soc. B* **347**, 213–234.

[31] Mongiardino Koch N. 2021 Exploring adaptive landscapes across deep time: A case study using

echinoid body size. *Evolution* **75**, 1567–1581.

[32] Mooi R. 1990 Progenetic miniaturization in the sand dollar Sinaechinocyamus: implications for

clypeasteroid phylogeny. In: *Echinoderm Research* (eds. C De Ridder, P Dubois, M Lahaye, M Jangoux),

pp. 137–143. Rotterdam, AA Balkema.

[33] Dunn CW, Zapata F, Munro C, Siebert S, Hejnol A. 2018 Pairwise comparisons across species are

problematic when analyzing functional genomic data. *Proc. Natl. Acad. Sci.* **115**, 409–417.

[34] Smith SD, Pennell MW, Dunn CW, Edwards SV. 2020 Phylogenetics is the new genetics (for most of

biodiversity). *Trends Ecol. Evol.* **35**, 415–425.

[35] Kroh A. 2020 Phylogeny and classification of echinoids. In: *Sea Urchins: Biology and Ecology, 4th*

edition (ed. JM Lawrence), pp. 1–17. Cambridge, Academic Press.

[36] Bolger AM, Lohse M, Usadel B. 2014 Trimmomatic: a flexible trimmer for Illumina sequence data.

Bioinformatics **30**, 2114–2120.

[37] Dunn CW, Howison M, Zapata F. 2013 Agalma: an automated phylogenomics workflow. *BMC*

Bioinformatics **14**, 330.

[38] Grabherr MG, Haas BJ, Yassour M, Levin JZ, Thompson DA, Amit I, Adiconis X, Fan L, Raychowdhury

R, Zeng Q, et al. 2011 Full-length transcriptome assembly from RNA-Seq data without a reference

genome. *Nature Biotechnology* **29**, 644–652.

[39] Modolo L, Lerat E. 2015 UrQt: an efficient software for the Unsupervised Quality trimming of NGS

data. *BMC Bioinformatics* **16**, 137.

[40] Li D, Liu C-M, Luo R, Sadakane K, Lam T-W. 2015 MEGAHIT: an ultra-fast single-node solution for

large and complex metagenomics assembly via succinct de Bruijn graph. *Bioinformatics* **31**, 1674–1676.

[41] Smit A, Hubley R, Green P. 2015 RepeatMasker Open-4.0. 2013–2015.

[42] Hoff KJ, Stanke M. 2019 Predicting genes in single genomes with AUGUSTUS. *Curr. Protoc.*

Bioinformatics **65**, e57.

[43] Stanke M, Keller O, Gunduz I, Hayes A, Waack S, Morgenstern B. 2006 AUGUSTUS: ab initio

prediction of alternative transcripts. *Nucleic Acids Res.* **34**, 435–439.

[44] Simion P, Belkhir K, François C, Veyssier J, Rink JC, Manuel M, Philippe H, Telford MJ. 2018 A

software tool 'CroCo' detects pervasive cross-species contamination in next generation sequencing data.

BMC Biol. **16**, 28.

[45] Ryan JF. 2014 Alien Index: identify potential non-animal transcripts or horizontally transferred

genes in animal transcriptomes. (doi: 10.5281/zenodo.21029)

[46] Guang A, Howison M, Zapata F, Lawrence C, Dunn CW. 2021 Revising transcriptome assemblies with

phylogenetic information. *PLoS One* **16**, e0244202.

[47] Ranwez V, Harispe S, Delsuc F, Douzery EJ. 2011 MACSE: Multiple Alignment of Coding SEquences

accounting for frameshifts and stop codons. *PLoS One* **6**, e22594.

[48] Talavera G, Castresana J. 2007 Improvement of phylogenies after removing divergent and

ambiguously aligned blocks from protein sequence alignments. *Syst. Biol.* **56**, 564–577.

[49] Morel B, Kozlov AM, Stamatakis A. 2018 ParGenes: a tool for massively parallel model selection and

phylogenetic tree inference on thousands of genes. *Bioinformatics* **35**, 1771–1773.

[50] Mai U, Mirarab S. 2018 TreeShrink: fast and accurate detection of outlier long branches in

collections of phylogenetic trees. *BMC Genomics* **19**, 272.

[51] Zhang C, Rabiee M, Sayyari E, Mirarab S. 2018 ASTRAL-III: polynomial time species tree

reconstruction from partially resolved gene trees. *BMC Bioinformatics* **19**, 153.

[52] Sayyari E, Mirarab S. 2016 Fast coalescent-based computation of local branch support from quartet

frequencies. *Mol. Biol. Evol.* **33**, 1654–1668.

[53] Aberer AJ, Kobert K, Stamatakis A. 2014 ExaBayes: massively parallel Bayesian tree inference for the

whole-genome era. *Mol Biol Evol* **31**, 2553–2556.

[54] Nguyen L-T, Schmidt HA, von Haeseler A, Minh BQ. 2014 IQ-TREE: a fast and effective stochastic

algorithm for estimating maximum-likelihood phylogenies. *Mol. Biol. Evol.* **32**, 268–274.

[55] Kalyaanamoorthy S, Minh BQ, Wong TK, von Haeseler A, Jermiin LS. 2017 ModelFinder: fast model

selection for accurate phylogenetic estimates. *Nat. Methods* **14**, 587–589.

[56] Hoang DT, Chernomor O, Von Haeseler A, Minh BQ, Vinh LS. 2017 UFBoot2: improving the ultrafast

bootstrap approximation. *Mol. Biol. Evol.* **35**, 518–522.

[57] Kozlov AM, Darriba D, Flouri T, Morel B, Stamatakis A. 2019 RAxML-NG: a fast, scalable and user-

friendly tool for maximum likelihood phylogenetic inference. *Bioinformatics* **35**, 4453-4455.

[58] Wang H-C, Minh BQ, Susko E, Roger AJ. 2018 Modeling site heterogeneity with posterior mean site

frequency profiles accelerates accurate phylogenomic estimation. *Syst. Biol.* **67**, 216–235.

[59] Mongiardino Koch N. 2021 Phylogenomic subsampling and the search for phylogenetically reliable

loci. *Mol. Biol. Evol.* **msab151**. (doi:10.1093/molbev/msab151).

[60] Lartillot N, Rodrigue N, Stubbs D, Richer J. 2013 PhyloBayes MPI. Phylogenetic reconstruction with

infinite mixtures of profiles in a parallel environment. *Syst. Biol.* **62**, 611–615.

[61] Strimmer K, Von Haeseler A. 1997 Likelihood-mapping: a simple method to visualize phylogenetic

content of a sequence alignment. *Proc. Natl. Acad. Sci.* **94**, 6815–6819.

[62] Shen X-X, Hittinger CT, Rokas A. 2017 Contentious relationships in phylogenomic studies can be

driven by a handful of genes. *Nat. Ecol. Evol.* **1**, 0126.

[63] Yang Z, Rannala B. 2006 Bayesian estimation of species divergence times under a molecular clock

using multiple fossil calibrations with soft bounds. *Mol. Biol. Evol.* **23**, 212–226.

[64] Telford MJ, Lowe CJ, Cameron CB, Ortega-Martinez O, Aronowicz J, Oliveri P, Copley RR. 2014

Phylogenomic analysis of echinoderm class relationships supports Asterozoa. *Proc. R. Soc. B* **281**,

20140479.

[65] Schlager S. 2017 Morpho and Rvcg–Shape Analysis in R. In: *Statistical shape and deformation*

analysis (eds. G Zheng, S Li, GJ Székely), pp. 217–256. Academic, London.

[66] R Core Team. 2019 *R: A Language and Environment for Statistical Computing*. Vienna, Austria: R

Foundation for Statistical Computing. See https://www.R-project.org.

[67] Paradis E, Schliep K. 2018 ape 5.0: an environment for modern phylogenetics and evolutionary

analyses in R. *Bioinformatics* **35**, 526–528.

[68] Lin J-P, Tsai M-H, Kroh A, Trautman A, Machado DJ, Chang L-Y, Reid R, Lin K-T, Bronstein O, Lee S-J,

et al. 2020 The first complete mitochondrial genome of the sand dollar *Sinaechinocyamus mai*

(Echinoidea: Clypeasteroida). *Genomics* **112**, 1686–1693.

[69] Dell'Ampio E, Meusemann K, Szucsich NU, Peters RS, Meyer B, Borner J, Petersen M, Aberer AJ,

Stamatakis A, Walzl MG, et al. 2014 Decisive data sets in phylogenomics: lessons from studies on the

phylogenetic relationships of primarily wingless insects. *Mol. Biol. Evol.* **31**, 239–249.

[70] Zamora S, Lefebvre B, Álvaro JJ, Clausen S, Elicki O, Fatka O, Jell P, Kouchinsky A, Lin J-P, Nardin E.

2013 Cambrian echinoderm diversity and palaeobiogeography. *Geol. Soc. Lond. Mem.* **38**, 157–171.

[71] Bottjer DJ, Davidson EH, Peterson KJ, Cameron RA. 2006 Paleogenomics of echinoderms. *Science*

314, 956–960.

[72] Pu JP, Bowring SA, Ramezani J, Myrow P, Raub TD, Landing E, Mills A, Hodgin E, Macdonald FA. 2016

Dodging snowballs: Geochronology of the Gaskiers glaciation and the first appearance of the Ediacaran

biota. *Geology* **44**, 955–958.

[73] Reich M. 2015 Different pathways in early evolution of the holothurian calcareous ring. In: *Progress*

in Echinoderm Palaeobiology (eds. S Zamora, I Rábano), pp. 137–145. Madrid: Instituto Geológico y

Minero de España.

[74] Miller AK, Kerr AM, Paulay G, Reich M, Wilson NG, Carvajal JI, Rouse GW. 2017 Molecular phylogeny

of extant Holothuroidea (Echinodermata). *Mol. Phylogenet. Evol.* **111**, 110–131.

[75] Rahman IA, Thompson JR, Briggs DEG, Siveter DJ, Siveter DJ, Sutton MD. 2019 A new ophiocistioid

with soft-tissue preservation from the Silurian Herefordshire Lagerstätte, and the evolution of the

holothurian body plan. *Proc. R. Soc. B* **286**, 20182792.

[76] Rouse GW, Jermiin LS, Wilson NG, Eeckhaut I, Lanterbecq D, Oji T, Young CM, Browning T, Cisternas

P, Helgen LE, et al. 2013 Fixed, free, and fixed: the fickle phylogeny of extant Crinoidea (Echinodermata)

and their Permian–Triassic origin. *Mol. Phylogenet. Evol.* **66**, 161–181.

[77] Brosseau O, Murienne J, Pichon D, Vidal N, Eléaume M, Ameziane N. 2012 Phylogeny of Cidaroida

(Echinodermata: Echinoidea) based on mitochondrial and nuclear markers. *Org. Divers. Evol.* **12**, 155–

165.

[78] Durham JW, Melville R. 1957 A classification of echinoids. *J. Paleontol.* **31**, 242–272.

[79] Suter SJ. 1994 Cladistic analysis of cassiduloid echinoids: trying to see the phylogeny for the trees.

Biol. J. Linn. Soc. **53**, 31–72.

[80] Kier PM. 1966 Cassiduloids. In: *Treatise on Invertebrate Paleontology, Part U* (ed. R Moore), pp.

492–523. Lawrence, University of Kansas Press.

[81] Mortensen T. 1948 *A Monograph of the Echinoidea. IV, 1 Holectypoida, Cassiduloida*. Copenhagen,

CA Reitzel.

[82] King B. 2020 Bayesian Tip-dated Phylogenetics in Paleontology: Topological Effects and Stratigraphic

Fit. *Syst. Biol.* **70**, 283–294.

[83] Turner AH, Pritchard AC, Matzke NJ. 2017 Empirical and Bayesian approaches to fossil-only

divergence times: a study across three reptile clades. *PLoS One* **12**, e0169885.

[84] Mongiardino Koch N, Garwood RJ, Parry LA. 2021 Fossils improve phylogenetic analyses of

morphological characters. *Proc. R. Soc. B* **288**, 2021004.

[85] Dos Reis M, Gunnell GF, Barba-Montoya J, Wilkins A, Yang Z, Yoder AD. 2018 Using phylogenomic

data to explore the effects of relaxed clocks and calibration strategies on divergence time estimation:

primates as a test case. *Syst. Biol.* **67**, 594–615.

[86] Warnock RC, Yang Z, Donoghue PC. 2011 Exploring uncertainty in the calibration of the molecular

clock. *Biol. Lett.* **8**, 156–159.

[87] Dos Reis M, Thawornwattana Y, Angelis K, Telford MJ, Donoghue PC, Yang Z. 2015 Uncertainty in

the timing of origin of animals and the limits of precision in molecular timescales. *Curr. Biol.* **25**, 2939–

2950.

[88] Sauquet H. 2013 A practical guide to molecular dating. *C. R. Palevol* **12**, 355–367.

[89] Carruthers T, Scotland RW. 2021 Uncertainty in divergence time estimation. *Syst. Biol.* **70**, 855–861.

[90] Carruthers T, Sanderson MJ, Scotland RW. 2020 The implications of lineage-specific rates for

divergence time estimation. *Syst. Biol.* **69**, 660–670.

[91] Tao Q, Barba-Montoya J, Huuki LA, Durnan MK, Kumar S. 2020 Relative efficiencies of simple and

complex substitution models in estimating divergence times in phylogenomics. *Mol. Biol. Evol.* **37**, 1819–

1831.

[92] Tao Q, Tamura K, Battistuzzi FU, Kumar S. 2019 A machine learning method for detecting

autocorrelation of evolutionary rates in large phylogenies. *Mol. Biol. Evol.* **36**, 811–824.

[93] Raup DM, Sepkoski JJ. 1982 Mass extinctions in the marine fossil record. *Science* **215**, 1501–1503.

[94] Kier PM. 1984 Echinoids from the Triassic (St. Cassian) of Italy, their lantern supports, and a revised

phylogeny of Triassic echinoids. *Smithson. Contrib. Paleobiol*. **56**, 1–41.

[95] Smith AB. 1994 Triassic echinoids from Peru. *Palaeontogr. Abt. A* **233**, 177–202.

[96] Kier PM. 1977 The poor fossil record of the regular echinoid. *Paleobiology* **3**, 168–174.

[97] Field D, Berv J, Hsiang A, Lanfear R, Landis M, Dornburg A. 2020 Timing the Extant Avian Radiation:

The Rise of Modern Birds, and the Importance of Modeling Molecular Rate Variation. *Bull. Am. Mus.*

Nat. Hist. **440**, 150–181.

[98] Caramés A, Adamonis S, Concheyro A, Remírez, M. 2019 New Finding of Regular Echinoid Elements

and Microfossils from the Pilmatué Member, Agrio Formation (Early Cretaceous), Neuquén Basin,

Argentina. In: *Advances in South American Micropaleontology* (eds. G Cusminsky, E Bernasconi, G

Concheyro), pp. 1–20. Springer Earth System Sciences. Springer, Cham.

[99] Ronquist F, Lartillot N, Phillips MJ. 2016 Closing the gap between rocks and clocks using total-

evidence dating. *Phil. Trans. R. Soc. B* **371**, 20150136.

[100] Coiro M, Doyle JA, Hilton J. 2019 How deep is the conflict between molecular and fossil evidence

on the age of angiosperms? *New Phytol.* **223**, 83–99.

Supplementary Information for:

N Mongiardino Koch, JR Thompson, AS Hatch, MF McCowin, AF Armstrong, SE Coppard, F Aguilera, O

Bronstein, A Kroh, R Mooi, GW Rouse – Phylogenomic analyses of echinoid diversification prompt a re-

evaluation of their fossil record

Details on tissue sampling, extraction, library preparation, sequencing and bioinformatic

approaches

Apatopygus recens, *Aspidodiadema hawaiiense*, *Fellaster zelandiae*, *Histocidaris variabilis*,

Rhyncholampas pacificus, *Sinaechinocyamus mai*, *Stereocidaris neumayeri*, *Tromikosoma hispidum*.

Tissue subsamples were finely chopped with a scalpel and preserved in RNA*later* (Ambion) buffer

solution for 1 day at 4°C to allow the RNA*later* to effectively penetrate the tissues, followed by long-

term storage at -80°C until RNA extraction. Total RNA was extracted from 1.5 mm Triple-Pure High

Impact Zirconium beads (Benchmark Scientific) in Trizol (Ambion), using Direct-zol RNA Miniprep Kit

(Zymo Research) with in-column DNase I incubation to remove genomic DNA. Prior to mRNA capture,

total RNA concentration was estimated using Qubit RNA HS Assay Kit (Invitrogen; range = 8.33-96

ng/μL), and quality was assessed using either High Sensitivity RNA ScreenTape or RNA ScreenTape with

an Agilent 4200 TapeStation. Mature mRNA was isolated from total RNA and libraries were prepared

using KAPA mRNA HyperPrep Kit (KAPA Biosystems) following the manufacturer's instructions (including

sample customization based on total RNA quantity and quality values), targeting an insert size circa 500

base pairs (bp), and using custom 10-nucleotide Illumina TruSeq style adapters [101]. Post-amplification,

DNA concentration was estimated using Qubit dsDNA BR Assay Kit (Invitrogen; range = 6.06-13.4 ng/μL).

Concentration, quality, and molecular weight distribution of libraries (range = 547-978 bp) was also

assessed using Genomic DNA ScreenTape with an Agilent 4200 TapeStation. Ten libraries (including two

annelids not employed here) were sequenced on one lane of a multiplexed run using NovaSeq S4

platform with 100 bp paired-end reads at the IGM Genomics Center (University of California San Diego).

The assembled transcriptomes were sanitized from cross-contaminant reads product of

multiplexed sequencing using CroCo v. 1.1 [44]. These eight datasets, as well as two annelid

transcriptomes not employed in this study but sequenced in the same lane, were employed. Transcripts

considered over or under-expressed across samples (as defined by default parameters), were kept.

This resulted in an average removal of 1.26% of assembled transcripts (range: 0.4% for

Aspidodiadema to 3.06% for *Histocidaris*).

Clypeaster japonicus, *Encope emarginata*, *Leodia sexiesperforata*, *Peronella japonica*. Eggs were

collected from a single female specimen of each species and immediately placed into RNA later for

preservation. The samples were left in RNAlater at 4°C for at least 1 day and then transferred to a -80°C

freezer until RNA extraction. RNA extraction was performed using Trizol and was then treated with

Ambion's Turbo DNA-free kit. RNA was quantified using both Nanodrop and an Agilent 2100

Bioanalyzer. Only RNA samples with an RNA integrity number (RIN) of > 7 were used. RNA-Seq libraries

were prepared using the NEB Ultra Directional kit using the standard protocol and multiplexed with 6bp

NEB multiplex primers. Paired-end 100 libraries were generated from the resultant libraries at the UC

Berkeley Genome Center using an Illumina HighSeq 2500.

Bathysalenia phoinissa, *Micropyga tuberculata*: The material was collected by the deep-sea

cruise BIOMAGLO [102] conducted in 2017 jointly by the French National Museum of Natural History

(MNHN) as part of the Tropical Deep-Sea Benthos program, the French Research Institute for

Exploitation of the Sea (IFREMER), the "Terres Australes et Antarctiques Françaises" (TAAF), the

Departmental Council of Mayotte and the French Development Agency (AFD), with the financial support

of the European Union (Xe FED). Specimens were sorted into taxonomic bins at class level and

collectively fixed in 70% ethanol at room temperature. Following taxonomic identification, tissue

samples were obtained using sterilized forceps and disposable scalpel blades. Tissue subsamples were

collected in high-purity 96% ethanol and stored at -40°C until DNA extraction. Total genomic DNA was

extracted using the DNeasy Blood and Tissue Kit (QIAGEN) following the manufacturer's instructions,

complemented by the addition of 4 µl RNAse A (QIAGEN, 100 mg/ml) for RNA digestion. Elution buffer

volume was reduced to 60 µl and the elution step repeated twice per sample in order to maximize DNA

concentration and yield. After collection of the first elution, spin-column membranes were rinsed again

twice with fresh elution buffer (40 µl) to further increase yield. Elutions were collected separately, with

the first one used for sequencing and the second for quality control. For the latter, a portion of the

mitochondrial cytochrome c oxidase subunit I (COI) was amplified using PCR. Amplifications were

conducted with TopTaq DNA Polymerase (QIAGEN) using 1 µl of extracted genomic DNA (approx. 10–15

ng). Details on primers and protocols can be found in [103]. PCR products were visualized on a 1%

agarose gel, purified using ExoSAP-IT (Affymetrix), and sequenced at Microsynth GmbH (Vienna, Austria)

with the same primers. Sequences are deposited in GenBank under the accession numbers XXX and XXX.

Prior to library preparation, total genomic DNA concentration was estimated using Qubit DNA

BR Assay Kit (Invitrogen) in order to determine library preparation strategy. This step was carried out by

Macrogen (Korea), using an Illumina TruSeq DNA PCR-free kit (for *Micropyga*) and an Illumina TruSeq

Nano DNA kit (for *Bathysalenia*). Library preparation followed the manufacturer's instructions, targeting

an insert size of 350 base pairs (bp). The libraries were sequenced by Macrogen (Korea) on a shared lane

of a multiplexed run using an Illumina HiSeq X instrument with 150 bp paired-end reads. Sequencing

depth was 134.1 and 161.3 million reads for *Bathysalenia* and *Micropyga*, respectively. Pre-processing of

Illumina shotgun data was carried out by removal of adapters using BBDuk

(https://sourceforge.net/projects/bbmap/) with settings: minlen=50 ktrim=r k=23 mink=11 tpe tbo);

followed by quality trimming of reads using UrQt v. 1.0.18 [39], discarding regions with phred quality

score < 28 and sequences of size < 50. Assembly of the data was carried out with MEGAHIT v. 1.1.2 [40],

in an iterative approach followed by assessment of assembly quality using BUSCO v. 3.0.2. scores [104]

using the metazoan dataset ODB v. 9 [105]. Final BUSCO scores of the draft assemblies were 28.4%

(S:27.8%, D:0.6%, F:48.2%, M:23.4%, n:978) for *Bathysalenia* and 17.2% (S:16.8%, D:0.4%, F:59.0%,

M:23.8%, n:978) for *Micropyga*. As recommended by Hoff and Stanke [42], draft genomes were masked

with RepeatMasker v. 4.1.0 [41] prior to gene prediction, using settings: -norna -xsmall; resulting in 3.82

% and 2.9% masked bases for *Bathysalenia* and *Micropyga*, respectively. Gene prediction was carried

out using AUGUSTUS v. 3.2.3 [43] using settings: --strand=both --singlestrand=false --genemodel=partial

--codingseq=on --sample=0 --alternatives-from-sampling=false --exonnames=on --softmasking=on. This

step relied on a training dataset composed of universal single-copy orthologs (USCOs) derived from the

most recent *Strongylocentrotus purpuratus* genome (Spur v.5).

Echinothrix calamaris, *Tripneustes gratilla*. Tissue samples were taken from live sea urchins

housed in laboratory aquaria and total RNA was immediately extracted from 150 mg of tissue using a

PureLink™ RNA extraction kit (Invitrogen™). The quality of total RNA was assessed on a BioAnalyzer

2100 (Agilent Technologies, Santa Clara, CA, USA) to ensure a RIN > 9 for all samples. Mature mRNA was

extracted from 1ug of total RNA and cDNA libraries were constructed using a TruSeq kit (Illumina).

Quality control of libraries was assessed on a BioAnalyzer and quantification measured using qPCR.

NEBNext Multiplex adaptors were added via ligation, and the cDNA libraries were sequenced at Genome

Quebec, McGill University. Three libraries were multiplex sequenced on one lane of Illumina at a

concentration of 8 pM per cDNA library using HiSeq2500 transcriptome sequencing to generate 125bp

paired-end reads. This resulted in approximately 80 million reads for each transcriptome.

Tetrapygus niger. Adult specimens were collected and transported alive to the University of

Concepcion. Tissue samples (female gonad and tube feet) from one individual were finely chopped with

a scalpel, and total RNA was extracted immediately using TriReagent (Sigma), following the

manufacturer's instructions. Total RNA concentration was estimated using QuantiFluor® RNA System

(111.84-339.04 ng/µL) and quality was assessed using Agilent RNA 6000 Nano kit with an Agilent 2100

TapeStation (RIN: 5.4-8.4). cDNA libraries were prepared at Genoma Mayor SpA (Chile) using TruSeq

Stranded mRNA Kit (Illumina) and sequenced on a multiplexed run using Illumina HiSeq 4000 platform

with 150 paired-end reads, resulting in 54.1 M reads for gonad female and 52.7 M reads for tube feet,

respectively. These two transcriptomic datasets were combined before performing all steps of

bioinformatic processing.

Eucidaris metularia. A single specimen was preserved in RNA*later* after being starved overnight

and kept for 1 day at 4°C before long-term storage at -80°C. Total RNA was extracted using Direct-zol

RNA Miniprep Kit (with in-column DNase treatment; Zymo Research) from Trizol. mRNA was isolated

with Dynabeads mRNA Direct Micro Kit (Invitrogen). RNA concentration was estimated using Qubit RNA

broad range assay kit, and quality was assessed using RNA ScreenTape with an Agilent 4200 TapeStation.

Values were used to customize downstream protocols following manufacturers' instructions. Library

preparation was performed with KAPA-Stranded RNA-Seq kits, targeting an insert size in the range of

200–300 base pairs. The quality and concentration of libraries were assessed using DNA ScreenTape.

The library was sequenced in a multiplexed pair-end runs using Illumina HiSeq 4000 with 7 other

libraries in the same lane (all previously published in [4]). In order to minimize read crossover, 10 bp

sequence tags designed to be robust to indel and substitution errors were employed [101].

The assembled transcriptome was filtered from cross-contaminant reads product of multiplexed

sequencing using CroCo v. 1.1 [44]. Seven other transcriptomes sequenced together were also

employed, and results of this step are reported in [3] (figure S1).

Fossil constraints on node ages

All clade names used below refer to the corresponding crown groups.

Ambulacraria (Echinodermata-Hemichordata divergence) – **Younger bound**: 518 Ma, Middle Atdabanian

(age 3, Series 3 of Cambrian). **Older bound**: 636.1 Ma, Lantian Biota, Ediacaran. **Reference:** [106]. **Notes:**

The root of our tree is constrained with a younger bound concurrent with the earliest occurrence of

stereom in the fossil record (see below). The older bound is based on the maximum age of the Lantian

Biota, a Lagerstätte lacking any trace of eumetazoan fossils.

Echinodermata (Pelmatozoan-Eleutherozoan divergence) – **Younger bound**: Unconstrained. **Older**

bound: 515.5 Ma, Middle Atdabanian (age 3, Series 3 of Cambrian). **Reference**: [106]. **Notes:** This

divergence (which represents the divergence of crown group echinoderms), has an older bound based

upon the earliest occurrence of stereom in the fossil record. Stereom constitutes an echinoderm

synapomorphy readily recognizable in the fossil record due to its unique mesh-like structure [71]. The

first records of stereom point to a sudden and global appearance within the middle Atdabanian [70].

Note that this is also used as younger bound for the divergence between echinoderms and

hemichordates.

Echinozoa (Echinoidea-Holothuroidea divergence) – **Younger bound:** 469.96 Ma, Top of

Pseudoclamacograptus decorates graptolite zone. **Older bound:** 461.95 Ma, Top Floian. **Reference:** [107,

108]. **Notes:** The oldest crown eleutherozoan fossils are disarticulated holothurian calcareous ring

elements which are known from the Red *Orthoceras* limestone of Sweden. This falls within the *P.*

decorates graptolite zone.

Asterozoa (Ophiuroidea-Asteroidea divergence) – **Younger bound**: 521 Ma, Top of *Psigraptus jacksoni*

Zone. **Older bound**: 480.55 Ma, Base of Series 2 Cambrian Period. **Reference**: [107, 108]. **Taxon**:

Maydenia roadsidensis. **Notes**: The divergence of asterozoan classes is calibrated based upon the

stratigraphic occurrence of *Maydenia roadsidensis*, the oldest asterozoan, which is a stem group

ophiuroid [109] and occurs in the *Psigraptus jacksoni* Zone of the Floian.

Asteroidea – **Younger bound:** 239.10 Ma, Top Fassanian, Ladinian, Mo3, *nodosus* biozone, Triassic.

Older bound: 252.16 Ma, P-T Boundary. **Reference**: [110]. **Taxon**: *Trichasteropsis weissmanni*. **Notes**:

Fossil evidence suggests the divergence of the asterozoan crown group took place sometime in the Early

or Middle Triassic [110]. Based upon its phylogenetic position and stratigraphic occurrence as the oldest

crown group asteroid, we use the forcipulatacean *T. weissmanni* from the Middle Triassic Muschelkalk

as the soft bound on this divergence.

Crinoidea – **Younger bound**: 247.06 Ma, Top Spathian (Paris Biota, Lower Shale unit, Thaynes Group,

Spathian, Triassic). **Older bound**: Unconstrained. **Reference**: [111]. **Taxon**: *Holocrinus*. **Notes**: Crown

group Crinoidea is difficult to define, though a rapid post-Palaeozoic morphological diversification is

supported by fossil evidence. We use *Holocrinus* from the Early Triassic of the Thaynes group to calibrate

the younger bound on the divergence of the crown group (i.e., the split between Isocrinida and all other

extant crinoids).

Ophiuroidea (Euryophiurida-Ophintegrida divergence) – **Younger bound**: 247.06 Ma, Top Spathian (Paris

Biota, Lower Shale unit, Thaynes Group, Spathian, Triassic). **Older bound**: Unconstrained. **Reference**:

[112]. **Taxon**: *Shoshonura brayardi*. **Notes**: *S. brayardi* from the Spathian Thaynes Group of the Western

USA is a member of the crown group ophiuroid clade Ophiodermatina. It is thus the currently oldest

described member of the ophiuroid crown group, setting its minimum age of divergence.

Pneumonophora (Holothuriida-Neoholothuriida divergence) – **Younger bound**: 259.8 Ma, S*pinosus* zone

of Early Ladinian (used base Ladinian). **Older bound**: 241.5 Ma, Base Wuchiapingian. **Reference**: [74,

107]. **Notes**: The oldest (undescribed) holothuriid calcareous ring ossicles are from the *Spinosus* zone of

the Ladinian of Germany and calibrate its divergence from neoholothuriids.

Echinoidea (Cidaroidea-Euechinoidea divergence) – **Younger bound**: 298.9 Ma, Base Carnian. **Older**

bound: 237 Ma, Base Permian. **Reference**: [113]. **Taxon**: *Triassicidaris? ampezzana*. **Notes**: The recent

phylogenetic analysis of Mongiardino Koch and Thompson [3] found many traditional early crown group

echinoids as members of the stem group. Given this result, *Triassicidaris? ampezzana*, a cidaroid which

is known from the St. Cassian beds of Italy, becomes the current oldest crown group echinoid.

Pedinoida-Aspidodiadematoida – **Younger bound**: 209.46 Ma, Top of Norian. **Older bound**: 252.16 Ma,

P-T boundary. **Reference**: See discussion in supplement of Thompson et al. [13]. **Taxon**: *Diademopsis* ex.

gr. *heberti*. **Notes**: The oldest euechinoid, *Diademopsis* ex. gr. *heberti*, which is a pedinoid, is known

from the Norian of Peru. This fossil calibrates the younger bound on the pedinoid-aspidodiadematoid

divergence.

Carinacea (Echinacea-Irregularia divergence) – **Younger bound**: 234.5 Ma, Top Sinemurian. **Older**

bound: 190.8 Ma, Top Julian 1 ammonoid zone of Norian. **Reference**: Li et al. [114]. **Taxon**:

Jesionekechinus. **Notes**: The divergence of Carinacea is calibrated based upon the oldest irregular

echinoid, *Jesionekechinus*, which occurs above the Sinemurian of New York Canyon, Nevada. For a more

detailed discussion of this divergence see the supplement of Thompson et al. [13].

Salenioida-(Camarodonta+Stomopneustoida) divergence– **Younger bound**: 228.35 Ma, Base Hettangian.

Older bound: 201.3 Ma, Top Carnian. **Reference**: [14]. **Taxon**: *Acrosalenia chartroni*. **Notes**: The stem

group salenioid *Acrosalenia chartroni* from the Hettangian of France is the earliest representative of

Echinacea. Given the topology recovered by our analyses, this fossil was used to calibrate the divergence

between salenioids and the clade composed of stomopneustoids and camarodonts.

Stomopneustoida-Camarodonta divergence – **Younger bound**: 201.3 Ma, Top Pliensbachian. **Older**

bound: 182.7 Ma, Base Jurassic. **Reference**: See discussion in supplement of Thompson et al. [13].

Taxon: Stomechinids from Morocco like *Diplechinus hebbriensis* [115]. **Notes**: The oldest stomechinids,

which are stem group stomopneustoids, are from the Early Jurassic of Morocco.

Neognathostomata-Atelostomata divergence – **Younger bound**: 234.5 Ma, Base of Toarcian. **Older**

bound: 182.7 Ma, St. Cassian Beds, bottom of Julian 2 ammonoid Zone. **Reference**: [14, 114]. **Taxon**:

Younger bound is set by *Galeropygus sublaevis* (older bound is relatively uninformative). **Notes**: The

base of the Toarcian (which contains the oldest neognathostomate *G. sublaevis*) is used as the lower

bound on the divergence between the neognathostomata and all other crown irregular echinoids. Note

that depending on the position of the unsampled echinoneoids, this split might also correspond to the

origin of crown group Irregularia (see recent topologies recovered by Lin et al. [68] and Mongiardino

Koch & Thompson [3]).

Atelostomata (Holasteroida-Spatangoida divergence) – **Younger bound**: 137.68 Ma, Bottom of

Campylotoxus zone in Valanginian. **Older bound**: 201.3 Ma, Base of Jurassic. **Reference**: [116]. **Taxon**:

Younger bound is based on *Toxaster* (older bound is loosely uninformative). **Notes**: The toxasterids are

stem group spatangoids that calibrate the divergence between spatangoids and holasteroids. The oldest

Toxaster are from the *Campylotoxus* zone in the Valanginian, and set the younger bound on the

divergence.

Echinolampadacea (Cassiduloida-Scutelloida divergence) – **Younger bound**: 113 Ma, Base Albian. **Older**

bound: 145 Ma, Base Cretaceous. **Reference**: [117]. **Taxon**: Younger bound is based on *Eurypetalum*

rancheriana (older bound is loosely uninformative). **Notes**: The oldest member of Echinolampadacea is

the faujasiid *Eurypetalum rancheriana* (originally placed in the genus *Faujasia* and later transferred, see

[28, 29]) from the Late Albian of Colombia.

Scutelloida (Scutelliformes-Laganiformes divergence) – **Younger bound**: 56.0 Ma, Bottom of Eocene.

Older bound: Unconstrained. **Reference**: [14, 118]. **Taxon**: Younger bound is based on *Eoscutum*

doncieuxi. **Notes**: We use the base of the Eocene as the younger bound on the divergence between

scutelliforms and laganiforms. This is based upon the occurrence of *E. doncieuxi*, the oldest scutelliform,

which is known from the Lower Eocene. The older bound is left unconstrained in order to account for

the current uncertainty in the origin of the clade (i.e., allow for Mesozoic ages).

Leodia-(*Encope+Mellita*) divergence – **Younger bound**: 23.03 Ma, Base Miocene. **Older bound**: 33.9 Ma,

Base of Oligocene. **Reference**: [119, 120]. **Taxon**: Younger bound is based on *Encope michoacanensis*

(older bound is loosely uninformative). **Notes**: This node is constrained using the oldest representative

of these three genera of mellitids.

Clypeasteroida – **Younger bound**: 47.8 Ma, Base Lutetian. **Older bound**: Unconstrained. **Reference**:

[121, 122]. **Taxon**: Younger bound is based on *Clypeaster calzadai* and *Clypeaster moianensis*. **Notes**:

The oldest known fossil species of Clypeasteroida (the clypeasterids *C. calzadai* and *C. moianensis*) are

from the middle Eocene, upper Lutetian (Biarritzian) of Cataluña, Spain. The older bound is left

unconstrained in order to account for the current uncertainty in the origin of the clade (i.e., allow for

Mesozoic ages).

<u>Strongylocentrotidae-Toxopneustidae divergence</u> – **Younger bound**: 44.4 Ma, Eocene Planktonic Zones

E9-E12. **Older bound**: 56.0 Ma, Base Eocene. **Reference**: [123, 124]. **Taxon**: Younger bound is based on

Lytechinus axiologus (older bound is loosely uninformative)*.* **Notes**: The oldest member of the clade

comprising toxopneustids and strongylocentrotids is the toxopneustid *Lytechinus axiologus.* This taxon is

known from the Eocene Yellow Limestone Group of Jamaica. The Yellow Limestone group represents the

Eocene Planktonic Zones E9-E12.

Supplementary Figures and Tables

Figure S1: Ordering of loci enforced using *genesortR* [59] and its relationship to the seven gene

properties employed. High ranking loci (i.e., the most phylogenetically useful) show low root-to-tip

variances (or high clock-likeness), low saturation and compositional heterogeneity, as well as high

average bootstrap and Robinson-Foulds similarity to a target topology (in this case, with the contentious

relationship among major lineages of Echinacea collapsed).

1000

Figure S2: Relationship between the root-to-tip variance (a proxy for the clock-likeness of loci) and the

rate of evolution (estimated as the total tree length divided by the number of terminals). The most

clock-like loci (shown in red), which are often favored for the inference of divergence times (e.g., [90,

125]), are among the most highly conserved and can provide little information for constraining node

ages [59]. Highly clock-like genes with a higher information content were used instead by choosing the

loci with the lowest root-to-tip variance from among those that were within 1 standard deviation from

the mean evolutionary rate (shown in blue).

1008

LN-CATGTR-1 LN-CATGTR-2 UGAM-CATGTR-1 UGAM-CATGTR-2

1009 **Figure S3:** Trace plots of the log-likelihood of different time-calibration runs. All runs show evidence of

1010 reaching convergence and stationarity before our imposed burn-in fraction of 10,000 generations. For

1011 simplicity, only runs under the CAT+GTR+G model are plotted. Those run under GTR+G converged much

1012 faster.

1013

Figure S4: Estimated branch lengths across different models of molecular evolution. Different site-

homogenous models (left) infer similar levels of divergence, and the choice between them induces little

distortion in the general tree structure. Site-heterogenous models on the other hand, not only infer a

larger degree of divergence between terminals relative to site-homogenous ones (center and left), but

they also distort the tree (i.e., impose a non-isometric stretching), with branch lengths connecting

outgroup taxa expanding much more than those within the ingroup clade.

1020

Figure S5: Effect of gene sampling strategy on inferred times of divergence. **A.** A chronospace, depicting

the differences in inferred ages between chronograms obtained using different loci. Axes constitute the

first two dimensions of the bgPCA (capturing 8.8% out of a total of 10.5% of variances summarized by all

four dimensions). The inset shows the centroid for each dataset, and the width of the lines connecting

them are scaled to the inverse of the Euclidean distances that separates them (as a visual summary of

overall similarity). The ages most different to those obtained under random loci sampling were found

when selecting clock-like genes. **B.** Change in branch lengths along the first two bgPC axes. Branches are

colored according to their contraction/expansion given a score of − 1 (top) or + 1 (bottom) standard

deviations, relative to the mean. Clades/nodes showing age differences larger than 20 Ma are labelled.

1030

Figure S6: Distribution of posterior probabilities for node ages that show an average difference larger

than 20 Ma depending on the choice of clock prior.

1033

Figure S7: Distribution of posterior probabilities for node ages that show a maximum difference larger

than 20 Ma depending on the choice of clock prior (no node showed average differences larger than 20

Ma). The largest differences can be seen in the relatively younger ages of Ambulacraria and

Echinodermata when using clock-like genes, and in the relatively older ages for some nodes within

cidaroids and starfish when using loci with high occupancy. Other sampling criteria largely agree on

inferred node ages, as can also be seen in figure S5 as short distances between their centroids in

chronospace.

1041

1042 **Figure S8:** Distribution of posterior probabilities for node ages that are the most affected by the choice

1043 of model of molecular evolution. No node showed average differences larger than 20 Ma, so those with

1044 the largest effect are shown.

1045

Figure S9: Median ages for selected clades across the consensus trees of the 40 time-calibrated runs

performed.

1048

Figure S10: Prior distributions of all constrained nodes. Five hundred replicates were sampled from the

joint prior, showing appropriately broad distributions of node ages. Blue lines show minima and yellow

ones maxima (when enforced); dotted lines show the age of the Permian-Triassic (251.9 Ma) and

Cretaceous-Paleogene (66 Ma) mass extinction events. Nodes whose ages are of special interest

(Echinoidea, Scutelloida, and Clypeasteroida) are shown in pink, revealing large prior probabilities of the

divergence occurring at either side of mass extinction events.

1055

Figure S11: Number of lineages inferred to have crossed the Permian-Triassic (P-T) boundary. The

probabilities of each scenario are estimated from the inferred divergence times of the 10,000

chronograms sampled across all of the analyses performed (250 for the two runs of each combination of

sampled loci, model of molecular evolution, and clock prior). The probability of three or more crown

group lineages surviving the Permian-Triassic extinction is 58.47%.

Table S1: Transcriptomic/genomic datasets added in this study relative to the taxon sampling of

Mongiardino Koch et al. [4] and Mongiardino Koch & Thompson [3]. Taxa with citations were taken from

the literature, and details can be found in the corresponding studies and associated NCBI BioProjects.

Taxa are shown in alphabetical order; those identified with "OG" are outgroup taxa (i.e., non-echinoids).

Voucher specimens are deposited at the Benthic Invertebrate Collection, Scripps Institution of

Oceanography (SIO-BIC), and the Echinoderm Collection, Muséum National d'Histoire Naturelle (MNHN-

IE). If multiple accession numbers are shown for a given taxon, these datasets were merged before

assembly. Similar details for all other specimens can be found in [3, 4].

Taxon	Citation	Tissue type	Collector	Locality (depth)	Voucher	GenBank accession number
Amphiura filiformis (OG)	[126]	-	-	-	-	SRX2255774
Apatopygus recens	This study	Mixed	Owen Anderson	Foveaux Strait, South Island, New Zealand	SIO-BIC E7142	
Aspidodiadema hawaiiense	This study	Mixed	Greg Rouse, Avery Hatch	Seamount 4, Costa Rica, Pacific Ocean (1,003 m)	SIO-BIC E7363	
Asterias rubens (OG)	[127]	-	-	-	-	SRX445860
Astrophyton muricatum (OG)	[128, 129]	-	-	-	-	SRX1391908
Bathysalenia phoinissa	This study	Tube feet, spine muscles	BIOMAGLO Cruise Team	Mozambique Channel, Mayotte (295-336 m)	MNHN-IE-2016-23	

Clypeaster japonicus	This study	Eggs	Frances Armstrong	Misaki Marine Biological Station, Kanagawa, Japan	-	
Echinothrix calamaris	This study	Male gonad	-	Phillipines	-	
Encope emarginata	This study	Eggs	Gulf Specimen Marine Lab	Apalachee Bay, Florida, USA	-	
Eucidaris metularia	This study	Mixed	Greg Rouse	Al-Fahal Reef, Red Sea, Makkah, Saudi Arabia	SIO-BIC 2017-008	
Fellaster zelandiae	This study	Spines, tube feet, gut, joining tissue of Aristotle's lantern	Wilma Blom	Western end of Cornwallis Beach, Auckland, New Zealand	SIO-BIC E7920	
Histocidaris variabilis	This study	Gonad	Greg Rouse, Avery Hatch	Las Gemelas Seamount, near Isla del Coco, Costa Rica, Pacific Ocean (571 m)	SIO-BIC E7350	
Lamberticrinus messingi (OG)	[129, 130]	-	-	-	-	SRX1397823
Leodia sexiesperforata	This study	Eggs	Frances Armstrong	Bocas del Toro, Panama	-	
Metacrinus rotundus (OG)	[131]	-	-	-	-	DRX021520
Micropyga tuberculata	This study	Tube feet, spine muscles	BIOMAGLO Cruise Team	Mozambique Channel, Îles Glorieuses (385-410 m)	MNHN-IE-2016-39	

Species	Reference	Tissue	Collector	Location	Voucher	SRA
Ophiocoma echinata (OG)	[127]	-	-	-	-	SRX445856
Peronella japonica	This study	Eggs	Frances Armstrong	Kanazawa, Japan	-	
Rhyncholampas pacificus	This study	Mixed	Carlos A. Conejeros-Vargas	Panteón Beach, Puerto Ángel Bay, Oaxaca, Mexico	SIO-BIC E7918	
Scaphechinus mirabilis	[132]	-	-	-	-	DRX187887 DRX187888
Sinaechinocyamus mai	This study	Gregory's diverticulum	Jih-Pai Lin	Miaoli County, Taiwan	SIO-BIC E7919	
Sterechinus neumayeri	[133]	-	-	-	-	ERX3498697 ERX3498698 ERX3498699 ERX3498700 ERX3498701
Stereocidaris nascaensis	This study	Muscle surrounding Aristotle's lantern	Charlotte Seid	Off San Ambrosio, Desventuradas Islands, Chile (215 m)	SIO-BIC E7154	
Tetrapygus niger	This study	Female gonad, tube feet	Felipe Aguilera	Dichato Bay, Chile	-	
Tripneustes gratilla	This study	Pedicellaria	-	Phillipines	-	

| *Tromikosoma hispidum* | This study | Tube feet | Lisa Levin, Todd Litke | Quepos Plateau, Costa Rica, Pacific Ocean (2067 m) | SIO-BIC E7251 |

1069

Table S2: Details of molecular datasets and supermatrix. Statistics for raw reads and assemblies are

shown for datasets incorporated in this study relative to the sampling of [3, 4] (where similar statistics

can be found for the other datasets). Taxa are shown in alphabetical order; those identified with "OG"

are outgroup taxa (i.e., non-echinoids). Novel datasets correspond to Illumina pair-end transcriptomes,

except for two draft genomes (*Bathysalenia phoinissa* and *Micropyga tuberculata*). Mean insert size is

expressed in number of base pairs. For transcriptomes, read pairs (initial) shows numbers input into

Agalma [37], (i.e., after processing with Trimmomatic [36] or UrQt [39]), while read pairs (retained)

shows those that passed the Agalma curation checks (including ribosomal, adapter, quality, and

compositional filters), and represent the final number of read pairs used for assembly. For genomes, see

information in the bioinformatic details above. Assemblies were further sanitized with either

alien_index [45] alone or in combination with CroCo [44] (see bioinformatic details), and the number of

assembled transcripts/contigs shows the size of datasets after these curation steps. The number of loci

shows the occupancy of terminals in the supermatrix output by Agalma (1,346 loci at 70% occupancy),

after which loci were further removed by TreeShrink [50], resulting in the final occupancy scores.

Species	Mean insert size	Read pairs (initial)	Read pairs (retained)	Assembled transcripts/contigs	Number of loci	Removed by TreeShrink	Final occupancy
Abatus agassizii	-	-	-	-	522	5	38.4
Abatus cordatus	-	-	-	-	497	2	36.8
Acanthaster planci (OG)	-	-	-	-	864	8	63.6
Amphiura filiformis (OG)	180.1	61,558,173	52,206,727	416,946	761	8	55.9
Apatopygus recens	415.3	74,315,687	56,898,850	274,380	1152	5	85.2
Apostichopus japonicus (OG)	-	-	-	-	849	9	62.4
Araeosoma leptaleum	-	-	-	-	1184	2	87.8
Arbacia lixula	-	-	-	-	1234	1	91.6
Aspidodiadema hawaiiense	228.7	109,716,219	104,518,714	311,032	1060	11	77.9
Asterias rubens (OG)	230.4	31,890,613	25,495,009	103,090	805	9	59.1
Asthenosoma varium	-	-	-	-	1254	7	92.6

Species							
Astrophyton muricatum (OG)	195.9	25,191,954	22,281,829	149,146	478	5	35.1
Bathysalenia phoinissa	-	-	-	154,120	605	6	44.5
Brissus obesus	-	-	-	-	773	0	57.4
Caenopedina hawaiiensis	-	-	-	-	1094	2	81.1
Clypeaster japonicus	198.6	10,505,520	9,298,316	74,743	829	1	61.5
Clypeaster rosaceus	-	-	-	-	1242	2	92.1
Clypeaster subdepressus	-	-	-	-	1233	3	91.4
Colobocentrotus atratus	-	-	-	-	1158	0	86.0
Conolampas sigsbei	-	-	-	-	1001	5	74.0
Cystechinus c.f. *giganteus*	-	-	-	-	661	0	49.1
Dendraster excentricus	-	-	-	-	337	1	25.0
Diadema setosum	-	-	-	-	305	1	22.6
Echinarachnius parma	-	-	-	-	1187	3	88.0
Echinocardium cordatum	-	-	-	-	1012	2	75.0
Echinocardium mediterraneum	-	-	-	-	1185	1	88.0
Echinocyamus crispus	-	-	-	-	709	6	52.2
Echinometra mathaei	-	-	-	-	1055	0	78.4
Echinothrix calamaris	203.9	68,871,087	60,443,904	251,788	1252	4	92.7
Encope emarginata	206.2	12,015,888	10,461,870	66,241	1076	1	79.9
Eucidaris metularia	321.6	38,802,159	29,047,401	83,318	412	2	30.5
Eucidaris tribuloides	-	-	-	-	414	0	30.8
Evechinus chloroticus	-	-	-	-	1196	1	88.8
Fellaster zelandiae	313.9	62,619,791	51,742,748	168,598	1269	1	94.2
Heliocidaris erythrogramma	-	-	-	-	931	0	69.2
Histocidaris variabilis	329.2	84,103,672	73,884,180	144,044	1189	5	88.0
Holothuria forskali (OG)	-	-	-	-	743	8	54.6
Lamberticrinus messingi (OG)	195.0	22,989,820	20,234,597	59,049	351	3	25.9
Leodia sexiesperforata	203.8	16,211,182	14,174,475	68,964	1064	1	79.0
Leptosynapta clarki (OG)	-	-	-	-	403	4	29.6
Lissodiadema lorioli	-	-	-	-	770	0	57.2
Loxechinus albus	-	-	-	-	1265	1	93.9

Species							
Lytechinus variegatus	-	-	-	-	1257	2	93.2
Mellita tenuis	-	-	-	-	1002	1	74.4
Meoma ventricosa	-	-	-	-	1053	0	78.2
Mesocentrotus nudus	-	-	-	-	1247	0	92.6
Metacrinus rotundus (OG)	198.5	23,791,832	20,900,345	83,718	642	7	47.2
Micropyga tuberculata	-	-	-	170,514	415	4	30.5
Ophiocoma echinata (OG)	278.5	28,427,026	24,836,025	130,153	712	7	52.4
Paracentrotus lividus	-	-	-	-	1266	0	94.1
Patiria miniata (OG)	-	-	-	-	740	8	54.4
Peronella japonica	208.3	17,696,287	15,707,931	106,110	1043	6	77.0
Prionocidaris baculosa	-	-	-	-	794	3	58.8
Psammechinus miliaris	-	-	-	-	1173	1	87.1
Rhyncholampas pacificus	346.8	63,116,413	52,160,741	234,910	1070	2	79.3
Saccoglossus kowalevskii (OG)	-	-	-	-	668	6	49.2
Scaphechinus mirabilis	401.7	10,169,195	8,796,067	127,664	974	4	72.1
Sinaechinocyamus mai	297.2	60,208,172	52,265,201	164,646	1233	1	91.5
Sphaerechinus granularis	-	-	-	-	1214	1	90.1
Sterechinus neumayeri	207.3	26,376,279	21,308,840	122,001	1097	1	81.4
Stereocidaris nascaensis	327.0	69,962,495	60,316,050	121,508	1200	6	88.7
Stomopneustes variolaris	-	-	-	-	908	0	67.5
Strongylocentrotus purpuratus	-	-	-	-	1266	5	93.7
Tetrapygus niger	202.5	63,040,541	52,761,671	163,084	1287	2	95.5
Tripneustes gratilla	154.6	62,962,239	57,015,692	130,376	1309	1	97.2
Tromikosoma hispidum	300.4	57,208,357	47,909,038	234,502	1238	7	91.5

1084

Supplementary References (references 1-100 in main text)

[101] Faircloth BC, Glenn TC. 2012 Not all sequence tags are created equal: designing and validating sequence identification tags robust to indels. *PLoS One* **7**, e42543.

[102] Corbari L, Samadi S, Olu K. 2017 BIOMAGLO cruise, RV Antea. (doi: doi.org/10.17600/17004000)

[103] Bronstein O, Kroh A, Miskelly AD, Smith SD, Dworjanyn SA, Mos B, Byrne M. 2019 Implications of range overlap in the commercially important pan-tropical sea urchin genus *Tripneustes* (Echinoidea: Toxopneustidae). *Mar. Biol.* **166**, 34.

[104] Seppey M, Manni M, Zdobnov EM. 2019 BUSCO: assessing genome assembly and annotation completeness. Methods Mol. Biol. **1962**, 227–245.

[105] Kriventseva EV, Tegenfeldt F, Petty TJ, Waterhouse RM, Simao FA, Pozdnyakov IA, Ioannidis P, Zdobnov EM. 2015 OrthoDB v8: update of the hierarchical catalog of orthologs and the underlying free software. *Nucleic Acids Res.* **43**, 250–256.

[106] Benton MJ, Donoghue PC, Asher RJ, Friedman M, Near TJ, Vinther J. 2015 Constraints on the timescale of animal evolutionary history. *Palaeontol. Electron.* **18**, 1–106.

[107] Erkenbrack EM, Thompson JR. 2019 Cell type phylogenetics informs the evolutionary origin of echinoderm larval skeletogenic cell identity. *Comm. Biol.* **2**, 160.

[108] Cooper R, Sadler P, Hammer O, Gradstein F. 2012 The Ordovician Period. In: *The geologic time scale* (eds. FM Gradstein, JG Ogg, M. Schmitz, G. Ogg), pp. 489–523. Elsevier.

[109] Hunter AW, Ortega-Hernández J. 2021 A new somasteroid from the Fezouata Lagerstätte in Morocco and the Early Ordovician origin of Asterozoa. *Biol. Lett.* **17**, 20200809.

[110] Villier L, Brayard A, Bylund KG, Jenks JF, Escarguel G, Olivier N, Stephen DA, Vennin E, Fara E. 2018

Superstesaster promissor gen. et sp. nov., a new starfish (Echinodermata, Asteroidea) from the Early

Triassic of Utah, USA, filling a major gap in the phylogeny of asteroids. *J. Syst. Palaeontol.* **16**, 395–415.

[111] Saucède T, Vennin E, Fara E, Olivier N, Team TPB. 2019 A new holocrinid (Articulata) from the Paris

Biota (Bear Lake County, Idaho, USA) highlights the high diversity of Early Triassic crinoids. *Geobios* **54**,

45–53.

[112] Thuy B, Escarguel G. 2019 A new brittle star (Ophiuroidea: Ophiodermatina) from the Early Triassic

Paris Biota (Bear Lake County, Idaho, USA). *Geobios* **54**, 55–61.

[113] Kroh A. 2011 Echinoids from the Triassic of St. Cassian-A review. *Geo. Alp.* **8**, 136–140.

[114] Li Z, Chen Z-Q, Zhang F, Ogg JG, Zhao L. 2020 Global carbon cycle perturbations triggered by

volatile volcanism and ecosystem responses during the Carnian Pluvial Episode (late Triassic). *Earth-Sci.*

Rev. **211**, 103404.

[115] Smith AB. 2011 *Gymnodiadema* and the Jurassic roots of the Arbacioida (stirodont echinoids).

Swiss J. Palaeontol. **130**, 155–171.

[116] François É, Marchand D, David B. 2003 Fluctuations morphologiques et hétérochronies chez

Toxaster (échinides, Crétacé inférieur). *C. R. Palevol* **2**, 597–605.

[117] Cooke CW. 1955 Some Cretaceous echinoids from the Americas. *US Geol. Surv. Prof. Pap.* **264**, 87–

112.

[118] Roman J. 1989 L'ancêtre éocène des Scutellidae (Echinoidea; Clypeasteroida). In: *Echinoderm*

Research. Proceedings of the Second European Echinoderm Conference, Brussels (ed. C De Ridder), pp.

18–21. Rotterdam, AA Balkema.

[119] Durham JW. 1994 Fossil *Encope* (Echinoidea) from the Pacific coast of southern Mexico. *Rev. Mex.*

Cienc. Geol. **11**, 113–116.

[120] Coppard SE, Lessios HA. 2017 Phylogeography of the sand dollar genus *Encope*: implications

regarding the Central American Isthmus and rates of molecular evolution. *Sci. Rep.* **7**, 11520.

[121] Ali MSM. 1983 The paleogeographic distribution of *Clypeaster* (Echinoidea) during the Cenozoic

Era. *Neues Jahrb. Geol. Paläontol.* **8**, 449–464.

[122] Via L, Padreny J. 1970 Dos nuevas especies de *Clypeaster* del Eoceno de Cataluña. *Publ. Inst. Invest.*

Geol. Diput. Prov. **24**, 89–118.

[123] Wade BS, Pearson PN, Berggren WA, Pälike H. 2011 Review and revision of Cenozoic tropical

planktonic foraminiferal biostratigraphy and calibration to the geomagnetic polarity and astronomical

time scale. *Earth-Sci. Rev.* **104**, 111–142.

[124] Gold DP, Fenton JP, Casas-Gallego M, Novak V, Pérez-Rodríguez I, Cetean C, Price R, Nembhard N,

Thompson H. 2018 The biostratigraphic record of Cretaceous to Paleogene tectono-eustatic relative sea-

level change in Jamaica. *J. S. Am. Earth Sci.* **86**, 140–161.

[125] Smith SA, Brown JW, Walker JF. 2018 So many genes, so little time: A practical approach to

divergence-time estimation in the genomic era. *PLoS One* **13**, e0197433.

[126] Dylus DV, Czarkwiani A, Stångberg J, Ortega-Martinez O, Dupont S, Oliveri P. 2016 Large-scale gene

expression study in the ophiuroid Amphiura filiformis provides insights into evolution of gene regulatory

networks. *Evodevo* **7**, 1–17.

[127] Reich A, Dunn CW, Akasaka K, Wessel, G. 2015 Phylogenomic analyses of Echinodermata support

the sister groups of Asterozoa and Echinozoa. *PLoS One* **10**, e0119627.

[128] Linchangco Jr GV, Foltz DW, Reid R, Williams J, Nodzak C, Kerr AM, Miller AK, Hunter R, Wilson NG,

Nielsen WJ, et al. 2017 The phylogeny of extant starfish (Asteroidea: Echinodermata) including *Xyloplax*,

based on comparative transcriptomics. *Mol. Phylogenet. Evol* **115**, 161–170.

[129] Janies DA, Witter Z, Linchangco GV, Foltz DW, Miller AK, Kerr AM, Jay J, Reid RW, Wray GA. 2016

EchinoDB, an application for comparative transcriptomics of deeply-sampled clades of echinoderms.

BMC Bioinformatics **17**, 17–48.

[130] Clouse RM, Linchangco Jr GV, Kerr AM, Reid RW, Janies DA. 2015 Phylotranscriptomic analysis

uncovers a wealth of tissue inhibitor of metalloproteinases variants in echinoderms. *R. Soc. Open Sci.* **2**,

150377.

[131] Koga H, Fujitani H, Morino Y, Miyamoto N, Tsuchimoto J, Shibata TF, Nozawa M, Shigenobu S,

Ogura A, Tachibana K, et al. 2016 Experimental approach reveals the role of *alx1* in the evolution of the

echinoderm larval skeleton. *PLoS One* **11**, e0149067.

[132] Simakov O, Kawashima T, Marlétaz F, Jenkins J, Koyanagi R, Mitros T, Hisata K, Bredeson J,

Shoguchi E, Gyoja F, et al. 2015 Hemichordate genomes and deuterostome origins. *Nature* **527**, 459–

465.

[133] Collins M, Peck LS, Clark MS. 2021 Large within, and between, species differences in marine

cellular responses: Unpredictability in a changing environment. *Sci. Total Environ.* **794**, 148594.