Phylogenomic analyses of echinoid diversification prompt a re-

evaluation of their fossil record

Short title: Phylogeny and diversification of sea urchins

Nicolás Mongiardino Koch[1,2]*, Jeffrey R Thompson[3,4], Avery S Hatch[2], Marina F McCowin[2], A

Frances Armstrong[5], Simon E Coppard[6], Felipe Aguilera[7], Omri Bronstein[8,9], Andreas Kroh[10], Rich

Mooi[5], Greg W Rouse[2]

[1] Department of Earth & Planetary Sciences, Yale University, New Haven CT, USA. [2] Scripps Institution of

Oceanography, University of California San Diego, La Jolla CA, USA. [3] Department of Earth Sciences,

Natural History Museum, Cromwell Road, SW7 5BD London, UK. [4] University College London Center for

Life's Origins and Evolution, London, UK. [5] Department of Invertebrate Zoology and Geology, California

Academy of Sciences, San Francisco CA, USA. [6] Bader International Study Centre, Queen's University,

Herstmonceux Castle, East Sussex, UK. [7] Departamento de Bioquímica y Biología Molecular, Facultad de

Ciencias Biológicas, Universidad de Concepción, Concepción, Chile. [8] School of Zoology, Faculty of Life

Sciences, Tel Aviv University, Tel Aviv, Israel. [9] Steinhardt Museum of Natural History, Tel-Aviv, Israel. [10]

Department of Geology and Palaeontology, Natural History Museum Vienna, Vienna, Austria

* Corresponding author. Email: nicolas.mongiardinokoch@yale.edu

Abstract

Echinoids are key components of modern marine ecosystems. Despite a remarkable fossil record, the emergence of their crown group is documented by few specimens of unclear affinities, rendering much of their early history uncertain. The origin of sand dollars, one of its most distinctive clades, is also unclear due to an unstable phylogenetic context and discrepancies between molecular divergence times and fossil evidence. We employ seventeen novel genomes and transcriptomes to build a phylogenomic dataset with a near-complete sampling of major lineages. With it, we revise the phylogeny and divergence times of echinoids, and place their history within the broader context of echinoderm evolution. We also introduce the concept of a chronospace—a multidimensional representation of node ages—and use it to explore the effects of using alternative gene samples, models of molecular evolution, and clock priors. We find the choice of clock model to have the strongest impact on divergence times, while the use of site-heterogeneous models shows little effects. The choice of loci shows an intermediate impact, affecting mostly deep Paleozoic nodes, for which clock-like genes recover dates more congruent with fossil evidence. Our results reveal that crown group echinoids originated in the Permian and diversified rapidly in the Triassic, despite the relative lack of fossil evidence for this early diversification. We also clarify the relationships among sand dollars and their close relatives, showing that the genus *Apatopygus* represents a relict lineage with a deep Jurassic origin. Surprisingly, the origin of sand dollars is confidently dated to the Cretaceous, implying ghost ranges spanning approximately 50 million years, a remarkable discrepancy with their rich fossil record.

Keywords: Echinoidea, Sea urchins, Sand dollars, Phylogenomics, Time calibration, Site heterogeneous models

Introduction

The fossil record represents the best source of primary data for constraining the origins of major lineages across the tree of life. However, the fossil record is not perfect, and even for groups with an excellent fossilization potential, constraining their age of origin can be difficult [1, 2]. Furthermore, as many traditional hypotheses of relationships have been revised in light of large-scale molecular datasets, the affinities of fossil lineages and their bearings on inferred times of divergence have also required a reassessment. An exemplary case of this is Echinoidea, a clade comprised of sea urchins, heart urchins, sand dollars, and allies, for which phylogenomic trees have questioned the timing of previously well-constrained nodes [3, 4].

Echinoids are easily recognized by their spine-covered skeletons or tests, composed of numerous tightly interlocking plates. Slightly over 1,000 living species have been described to date [5], a diversity that populates every marine benthic environment from intertidal to abyssal depths [6]. Echinoids are usually subdivided into two morpho-functional groups with similar species-level diversities: "regular" sea urchins, a paraphyletic assemblage of hemispherical, epibenthic consumers protected by large spines; and irregulars (Irregularia), a clade of predominantly infaunal and bilaterally symmetrical forms covered by small and specialized spines. In today's oceans, regular echinoids act as ecosystem engineers in biodiverse coastal communities such as coral reefs [7] and kelp forests [8], where they are often the main consumers. They are first well-known in the fossil record on either side of the Permian-Triassic (P-T) mass extinction event when many species occupied reef environments similar to those inhabited today by their descendants [9, 10]. This extinction event was originally thought to have radically impacted the macroevolutionary history of the clade, decimating the echinoid stem group and leading to the radiation of crown group taxa from a single surviving lineage [11, 12]. However, it is now widely accepted that the origin of crown group Echinoidea (i.e., the divergence between its two

main lineages, Cidaroidea and Euechinoidea) occurred in the Late Permian, as supported by molecular

estimates of divergence [13, 14], as well as the occurrence of Permian fossils with morphologies typical

of modern cidaroids [15, 16]. However, a recent total-evidence study recovered many taxa previously

classified as crown group members along the echinoid stem, while also suggesting that up to three

crown group lineages survived the P-T mass extinction [3]. This result increases the discrepancy between

molecular estimates and the fossil record and renders uncertain the early evolutionary history of crown

group echinoids. Constraining the timing of origin of this clade relative to the P-T mass extinction

(especially in the light of recent topological changes [3, 4]) is further complicated by the poor

preservation potential of stem group echinoids, and the difficulty assigning available disarticulated

remains from the Late Paleozoic and Early Triassic to specific clades [11, 12, 17-20].

Compared to the morphological conservatism of regular sea urchins, the evolutionary history of

the relatively younger Irregularia was characterized by dramatic levels of morphological and ecological

innovation [21-24]. Within the diversity of irregulars, sand dollars are the most easily recognized (Fig. 1).

The clade includes greatly flattened forms that live in high-energy sandy environments where they feed

using a unique mechanism for selecting and transporting organic particles to the mouth, where these

are crushed using well-developed jaws [25, 26]. Sand dollars (Scutelloida) were long thought to be most

closely related to sea biscuits (Clypeasteroida) given a wealth of shared morphological characters [19,

25]. The extraordinary fossil record of both sand dollars and sea biscuits suggested their last common

ancestor originated in the early Cenozoic from among an assemblage known as "cassiduloids" [23, 25], a

once diverse group that is today represented by three depauperate lineages: cassidulids (and close

relatives), echinolampadids, and apatopygids [19, 27]. These taxa not only lack the defining features of

both scutelloids and clypeasteroids but have experienced little morphological change since their origin

deep in the Mesozoic [24, 27-29]. However, early molecular phylogenies supported both cassidulids and

echinolampadids as close relatives of sand dollars (e.g., [14, 30]), a topology initially disregarded for its

conflicts with both morphological and paleontological evidence, but later confirmed using phylogenomic

approaches [4]. While many of the traits shared by sand dollars and sea biscuits have since been

suggested to represent a mix of convergences and ancestral synapomorphies secondarily lost by some

"cassiduloids" [3, 4], the strong discrepancy between molecular topologies and the fossil record remains

unexplained. Central to this discussion is the position of apatopygids, a clade so far unsampled in

molecular studies. Apatopygids have a fossil record stretching more than 100 million years and likely

have phylogenetic affinities with even older extinct lineages [3, 19, 28, 29]. Although current molecular

topologies already imply ghost ranges for scutelloids and clypeasteroids that necessarily extend beyond

the Cretaceous-Paleogene (K-Pg) boundary, the phylogenetic position of apatopygids could impose even

earlier ages on these lineages (Fig. 1). Constraining these divergences is necessary to understand the

timing of origin of the sand dollars, one of the most specialized lineages of echinoids [24-27]. Resolving

some phylogenetic relationships within scutelloids has also been complicated by their recurrent

miniaturization and associated loss of morphological features (Fig. 1; [25, 31, 32]).

Figure 1: Neognathostomate diversity and phylogenetic relationships. **A.** *Fellaster zelandiae*, North

Island, New Zealand (Clypeasteroida). **B.** Large specimen: *Peronella japonica*, Ryukyu Islands, Japan;

Small specimen: *Echinocyamus crispus*, Maricaban Island, Philippines (Laganina: Scutelloida). **C.** Large

specimen: *Leodia sexiesperforata*, Long Key, Florida; Small specimen: *Sinaechinocyamus mai*, Taiwan

(Scutellina: Scutelloida). **D.** *Rhyncholampas pacificus,* Isla Isabela, Galápagos Islands (Cassidulidae). **E.**

Conolampas sigsbei, Bimini, Bahamas (Echinolampadidae). **F.** *Apatopygus recens* , Australia

(Apatopygidae). **G.** Hypotheses of relationships among neognathostomates. Top: Morphology supports

a clade of Clypeasteroida + Scutelloida originating after the K-Pg boundary, subtended by a paraphyletic

assemblage of extant (red) and extinct (green) "cassiduloids" [19]. Bottom: A recent total-evidence

study split most of cassiduloid diversity into a clade of extant lineages closely related to scutelloids, and

an unrelated clade of extinct forms (Nucleolitoida) [3]. Divergence times are much older and conflict

with fossil evidence. Cassidulids and apatopygids lacked molecular data in this analysis. Scale bars = 10

mm.

Echinoidea constitutes a model clade in developmental biology and genomics. As these fields

embrace a more comparative approach [13, 33, 34], robust and time-calibrated phylogenies are

expected to play an increasingly important role. Likewise, the extraordinary fossil record of echinoids

and the ease with which echinoid fossils can be incorporated in phylogenetic analyses make them an

ideal system to explore macroevolutionary dynamics using phylogenetic comparative methods [3, 31]. In

this study, we build upon available molecular resources with 17 novel genome-scale datasets and build

the largest molecular matrix for echinoids yet compiled. Our expanded phylogenomic dataset extends

sampling to 16 of the 17 currently recognized echinoid orders (plus the unassigned apatopygids) [35],

and is the first to bracket the extant diversity of both sand dollars and sea biscuits and include members

of all three lineages of living "cassiduloids" (cassidulids, echinolampadids, and apatopygids). We also

incorporate a diverse sample of outgroups, providing access to the deepest nodes within the crown

groups of all other echinoderm classes (holothuroids, asteroids, ophiuroids, and crinoids). With it, we

reconstruct the phylogenetic relationships and divergence times of the major lineages of living echinoids

and place their diversification within the broader context of echinoderm evolution.

Results

133   **(a)   Phylogeny of Echinoidea**

134   Phylogenetic relationships supported by the full dataset were remarkably stable, with all nodes

but one being identically resolved and fully supported across a wide range of inference methods (Fig.

2A). While recovering a topology similar to those of previous molecular studies [3, 4, 13, 14, 30, 36], this

analysis is the first to sample and confidently place micropygoids and aspidodiadematoids within

Aulodonta, as well as resolve the relationships among all major clades of Neognathostomata

(scutelloids, clypeateroids and the three lineages of extant "cassiduloids"). Our results show that

Apatopygus recens is not related to the remaining "cassiduloids" but is instead the sister clade to all

other sampled neognathostomates. The strong support for this placement, as well as for a clade of

cassidulids and echinolampadids (Cassiduloida *sensu stricto*) as the sister group to sand dollars, provides

a basis for an otherwise elusive phylogenetic classification of neognathostomates. Our topology also

confirms that *Sinaechinocyamus mai*, a miniaturized species once considered a plesiomorphic member

of Scutelloida based on the reduction or loss of diagnostic features (Fig. 1), is in fact a derived

paedomorphic lineage closely related to *Scaphechinus mirabilis* [32].

Figure 2: Phylogenetic relationships among major clades of Echinoidea. **A.** Favored topology. With the exception of a single contentious node within Echinacea (marked with a star), all methods supported the same pattern of relationships. Numbers below major clades correspond to the current numbers of described living species (obtained from [5]). **B.** Likelihood-mapping analysis showing the proportion of quartets supporting different resolutions within Echinacea. While the majority of quartets support the topology depicted in **A** (shown in red), a significant number support an alternative resolution that has been recovered in morphological analyses [19] (shown in blue). **C.** Support for a clade of Salenioida + (Camarodonta + Stomopneustoida), as depicted in **A**, is predominantly concentrated among the most

phylogenetically useful loci. This signal is attenuated in larger datasets that contain less reliable genes,

eventually favoring an alternative resolution. Only the 584 loci containing data for the three main

lineages of Echinacea were considered. The line corresponds to a second-degree polynomial regression.

D. Resolution and bootstrap scores (color coded) of the topology within Echinacea found using datasets

of different sizes and alternative methods of inference.

Salenioida is another major lineage sampled here for the first time, and whose exact position

among regular echinoids proved difficult to resolve. While some methods supported salenioids as the

sister group to a clade of camarodonts, stomopneustoids, and arbacioids (a topology previously

supported by morphology [19]), others recovered a closer relationship of salenioids to Camarodonta +

Stomopneustoida, with arbacioids sister to them all (as shown in Fig. 2A). As revealed using likelihood

mapping, these results do not stem from a lack of phylogenetic signal, but rather from the presence of

strong and conflicting evidence in the dataset regarding the position of salenioids (Fig. 2B). However, a

careful dissection of these signals shows that loci with high phylogenetic usefulness (as defined by [3,

37]; see Methods) favor the topology shown in Fig. 2A, with the morphological hypothesis becoming

dominant only after incorporating less reliable loci (Fig .2C). In line with these results, moderate levels of

gene subsampling (down to 500 loci) targeting the most phylogenetically useful loci unambiguously

support the placement of arbacioids as sister to the remaining taxa, regardless of the chosen method of

inference (Fig. 2D). More extreme subsampling (down to 100 loci) again results in disagreement among

methods. This possibly stems from the increasing effect of stochastic errors in smaller datasets, as less

than half of the sampled loci in these reduced datasets contain data for all branches of this quartet (see

Fig. 2C). This result shows the importance of ensuring that datasets (especially subsampled ones) retain

appropriate levels of occupancy for clades bracketing contentious nodes [38]. Despite these

disagreements, several lines of evidence favor the topology shown in Fig. 2A, including the results of

likelihood mapping, and the increased support for this resolution among the most phylogenetically

useful loci and when using more complex methods of reconstruction, such as partitioned and site-

heterogeneous models, which always favor this topology regardless of dataset size (Fig. 2D).

(b)   Sensitivity of node ages

While alternative methods of inference had minor effects on phylogenetic relationships, they

did impact the reconstruction of branch lengths (Fig. S1). Site-heterogeneous models (such as

CAT+GTR+G) returned longer branch lengths overall, but also uncovered a larger degree of molecular

change among echinoderm classes. Branches connecting these clades were stretched to a much larger

extent than those within the ingroup (Fig. S1), which might affect the inference of node dates. We

tested this hypothesis by exploring the sensitivity of divergence times to the use of alternative models of

molecular evolution (site-homogeneous vs. site-heterogeneous), as well as different clock priors

(autocorrelated vs. uncorrelated), and gene sampling strategies (using five different approaches; see

Methods). All combinations of these factors were explored, resulting in 20 different time-calibration

settings that were run using Bayesian approaches under a constrained tree topology (that shown in Fig.

2A). Posterior distributions of node ages were then obtained from each of these time-calibrated

analyses, confirming that inferred divergence times were significantly affected by all of these factors

(multivariate analysis of variance; all $P < 2.2 \times 10^{-16}$). While nodes connecting some outgroup taxa were

among those most sensitive to these decisions, large effects were also seen among nodes relating to the

origin and diversification of the echinoid clades Cidaroidea and Aulodonta. All of these nodes varied in

age by more than 60 Ma—and up to 115 Ma—among the consensus topologies of different analyses

(Fig. S2).

In order to isolate and visualize the impact of each of these factors on divergence time

estimation, chronograms were represented in a multidimensional space of node dates, with each axis

representing the age of a given node. We term this type of graph a chronospace given its similarities to

the treespaces commonly used to explore topological differences between phylogenetic trees [39]. Each

observation (chronogram) was classified as obtained under a specific clock prior, model of molecular

evolution, and gene sampling strategy, and the major effects of each of these choices were extracted

with the use of between-group principal component analyses (bgPCAs). This method of ordination

allows for the isolation of axes that capture differences between group means, and has been used

before with other high-dimensional datasets such as obtained from microarrays and geometric

morphometric approaches [40, 41]. The single dimension of chronospace maximizing the distinctiveness

of chronograms obtained under different clock models explained 53.7% of the total variance in node

ages across all analyses (Fig. 3). In contrast, the choice of different loci or models of molecular evolution

showed a much lesser effect, explaining only 10.5% and 4.0% of the total variance, respectively. Even

though all of these decisions affected a similar set of sensitive nodes (those mentioned above, as well as

some relationships within Atelostomata), the choice of clock model modified the ages of 17 of these by

more than 20 Ma (Fig. S3). This degree of change was induced on only 5 nodes by selecting alternative

loci (Fig. S4), and was not induced on any node by enforcing different models of evolution (Fig. S5).

Regarding gene choice, the ages most different to those obtained under random loci selection were

found when using the most clock-like genes (Fig. 3C).

Figure 3: Sensitivity of divergence time estimation to methodological decisions. Between-group principal

component analysis (bgPCA) was used to retrieve axes that separate chronograms based on the clock

model (**A**), model of molecular evolution (**B**), and gene sampling strategy (**C**) employed. In the latter

case, only the first two out of four bgPCA dimensions are shown. The inset shows the centroid for each

loci sampling strategy, and the width of the lines connecting them are scaled to the inverse of the

Euclidean distances that separates them (as a visual summary of overall similarity). The proportions of

total variance explained are shown on the axis labels. The impact of the clock model is such that a

bimodal distribution of chronograms can be seen even when bgPCA are built to discriminate based on

other factors (as in **C**).

(c)   Echinoid (and echinoderm) divergence times

       Even when the age of crown Echinodermata was constrained to postdate the appearance of

stereom (the characteristic skeletal microstructure of echinoderms) in the Early Cambrian [42, 43], only

analyses using the most clock-like loci recovered ages concordant with this (i.e., median ages younger

than the calibration enforced; Fig. S4). Instead, most consensus trees favored markedly older ages for

the clade, in some cases even predating the origin of the Ediacaran biota [44] (Fig. S6). Despite the

relative sensitivity of many of the earliest nodes to methodological choices (Fig. S2), the split between

Crinoidea and all other echinoderms (Eleutherozoa) is always inferred to have predated the end of the

Cambrian (youngest median age = 492.2 Ma), and the divergence among the other major lineages

(classes) of extant echinoderms are constrained to have happened between the Late Cambrian and

Middle Ordovician (Fig. S6). Our results also recover an early origin of crown group Holothuroidea (sea

cucumbers; range of median ages = 351.6–383.2 Ma), well before the crown groups of other extant

echinoderm classes. These dates markedly postdate the first records of holothuroid calcareous rings in

the fossil record [45, 46], and imply that this trait does not define the holothuroid crown group but

instead evolved from an echinoid-like jaw-apparatus along its stem [47]. The other noteworthy

disagreement between our results and those of previous studies [48] involves dating crown group

Crinoidea to times that precede the P-T mass extinction (range of median ages = 268.0–327.7 Ma,

although highest posterior density intervals are always wide and include Triassic ages).

Across all of the analyses performed, the echinoid crown group is found to have originated

somewhere between the Pennsylvanian and Cisuralian, with 42% posterior probability falling within the

late Carboniferous and 58% within the early Permian (Fig. 4). An origin of the clade postdating the P-T

mass extinction is never recovered, even when such ages were allowed under the joint prior (Fig. S7).

While the posterior distribution of ages for Euechinoidea spans both sides of the P-T boundary, the

remaining earliest splits within the echinoid tree are constrained to have occurred during the Triassic,

including the origins of Aulodonta, Carinacea, Echinacea, and Irregularia (Figs. 4 and S6). Many echinoid

orders are also inferred to have diverged from their respective sister clades during this period, including

aspidodiadematoids, pedinoids, echinothurioids, arbacioids, and salenioids. LTT plots confirm that

lineage diversification proceeded rapidly throughout the Triassic (Fig. 4B). Despite the topological

reorganization of Neognathostomata, the clade is dated to a relatively narrow time interval in the Late

to Middle Jurassic (range of median ages = 169.48-179.96 Ma), in agreement with recent estimates [3].

Within this clade, the origins of both scutelloids and clypeasteroids confidently predate the K-Pg mass

extinction (posterior probability of origination before the boundary = 0.9995 and 0.9621, respectively),

despite younger ages being allowed by the joint prior (Fig. S7).

Figure 4: Divergence times among major clades of Echinoidea and other echinoderms. **A.** Consensus

chronogram of the two runs using clock-like genes under a CAT+GTR+G model of molecular evolution

and a autocorrelated log-normal (LN) clock prior. Node ages correspond to median values, and bars

show the 95% highest posterior density intervals. **B.** Lineage-through-time plot, showing the rapid

divergence of higher-level clades following the P-T mass extinction (shown with dashed lines, along with

the K-Pg boundary). Each line corresponds to an individual consensus topology from among the time-

calibrated runs performed. **C.** Posterior distributions of the ages of selected nodes (identified in **A**). The

effects of models of molecular evolution are not shown, as they represent the least important factor

(see Fig. 3); distributions using site-homogeneous and heterogeneous models are combined for every

combination of targeted loci and clock prior.

Discussion

(a)   The echinoid Tree of Life

283   In agreement with previous phylogenomic studies [3, 4], echinoid diversity can be subdivided

into five major clades (Fig. 2A). Cidaroids form the sister group to all other crown group echinoids

(Euechinoidea). Some aspects of the relationships among sampled cidaroids are consistent with previous

molecular [49] and morphological studies [19], including an initial split between *Histocidaris* and the

remaining taxa, representing the two main branches of extant cidaroids [5, 35]. Others, such as the

nested position of *Prionocidaris baculosa* within the genus *Eucidaris,* not only implies paraphyly of this

genus but also suggests the need for a taxonomic reorganization of the family Cidaridae. Within

euechinoids, the monophyly of Aulodonta is supported for the first time with sampling of all of its major

groups. The subdivision of these into a clade that includes diadematoids plus micropygoids (which we

propose should retain the name Diadematacea), sister to a clade including echinothurioids and

pedinoids (Echinothuriacea *sensu* [4]) is strongly reminiscent of some early classifications (e.g., [50]).

Our expanded phylogenomic sampling also confirms an aulodont affinity for aspidodiadematoids [3, 35]

and places them within Echinothuriacea as the sister group to Pedinoida.

In agreement with previous phylogenomic studies [3, 4], echinoid diversity can be subdivided

into five major clades (Fig. 2A). Cidaroids form the sister group to all other crown group echinoids

(Euechinoidea). Some aspects of the relationships among sampled cidaroids are consistent with previous

molecular [47] and morphological studies [19], including an initial split between *Histocidaris* and the

remaining taxa, representing the two main branches of extant cidaroids [5, 35]. Others, such as the

nested position of *Prionocidaris baculosa* within the genus *Eucidaris,* not only implies paraphyly of this

genus but also suggests the need for a taxonomic reorganization of the family Cidaridae. Within

euechinoids, the monophyly of Aulodonta is supported for the first time with sampling of all of its major

groups. The subdivision of these into a clade that includes diadematoids plus micropygoids (which we

propose should retain the name Diadematacea), sister to a clade including echinothurioids and

pedinoids (Echinothuriacea *sensu* [4]) is strongly reminiscent of some early classifications (e.g., [48]).

Our expanded phylogenomic sampling also confirms an aulodont affinity for aspidodiadematoids [3, 35]

and places them within Echinothuriacea as the sister group to Pedinoida.

The remaining diversity of echinoids, which forms the clade Carinacea (Fig. 2), is subdivided into

Irregularia and their sister clade among regulars, for which we amend the name Echinacea to include

Salenioida. Given the striking morphological gap separating regular and irregular echinoids, the origin of

Irregularia has been shrouded in mystery [19, 23, 50]. Our complete sampling of major regular lineages

determines Echinacea *sensu stricto* to be the sister clade to irregular echinoids. A monophyletic

Echinacea was also supported in a recent total-evidence analysis [3] , but the incomplete molecular

sampling of that study resulted in a slightly different topology that placed salenioids as the sister group

to the remaining lineages. However, an overall lack of morphological synapomorphies uniting these

clades had previously been acknowledged [19]. While the relationships within Echinacea proved to be

difficult to resolve even with thousands of loci, multiple lines of evidence lead us to prefer a topology in

which salenioids form a clade with camarodonts + stomopneustoids, with arbacioids sister to all of these

(Fig. 2).

As has been already established [3, 4, 14, 19, 30], the lineages of irregular echinoids here

sampled are subdivided into Atelostomata (heart urchins and allies) and Neognathostomata (sand

dollars, sea biscuits, and "cassiduloids"). Despite the former being the most diverse of the five main

clades of echinoids (Fig. 2), its representation in phylogenomic studies remains low, and its internal

phylogeny poorly constrained [35]. On the contrary, recent molecular studies have greatly improved our

understanding of the relationships among neognathosomates [3, 4, 36], revealing an evolutionary

history that dramatically departs from previous conceptions. Even when scutelloids and clypeasteroids

were never recovered as reciprocal sister lineages by molecular phylogenies (e.g., [13, 14, 30]), this

result was not fully accepted until phylogenomic data confidently placed echinolampadids as the sister

lineage to sand dollars [4]. At the same time, this result rendered the position of the remaining

"cassiduloids", a taxonomic wastebasket with an already complicated history of classification [19, 27, 28,

51], entirely uncertain. An attempt to constrain the position of these using a total-evidence approach [3]

subdivided the "cassiduloids" into three unrelated clades: Nucleolitoida, composed of extinct lineages

and placed outside the node defined by Scutelloida + Clypeasteroida, and two other clades nested

within it (see Fig. 1). Extant "cassiduloids" were recovered as members of one of the latter clades,

representing the monophyletic sister group to sand dollars. Here, we show that *Apatopygus recens* does

not belong within this clade but is instead the sister group to all other extant neognathostomates. Given

this phylogenetic position, as well as the morphological similarities between *Apatopygus* and the

entirely extinct nucleolitids [19, 28, 51-53], it is likely that the three extant species of apatopygids

represent the last surviving remnants of Nucleolitoida, a clade of otherwise predominantly Mesozoic

neognathostomates [3]. Because of the renewed importance in recognizing this topology, we propose

the name Luminacea for the clade uniting all extant neognathostomates with the exclusion of

Apatopygus (Fig. 2A). This nomenclature refers to the dynamic evolutionary history of the Aristotle's

lantern (i.e., the echinoid jaw-apparatus) within the clade (present in the adults of both clypeasteroids

and scutelloids, but found only in the juveniles of Cassiduloida *sensu stricto*), the inclusion of the so-

called lamp urchins (echinolampadids) within the clade, and the illumination provided by this hitherto

unexpected topology. The previous misplacement of *Apatopygus* within this clade ([3]; see Fig. 1G) is

likely a consequence of tip-dating preferring more stratigraphically congruent topologies [54], an effect

that can incorrectly resolve taxa on long terminal branches [55]. Given the generally useful phylogenetic

signal of stratigraphic information [56], this inaccuracy further highlights the unusual evolutionary

history of living apatopygids.

(b)   Chronospaces: a statistical exploration of time-calibration strategies

Calibrating phylogenies to absolute time is crucial to understanding evolutionary history, as the

resulting chronograms provide a major avenue for testing hypotheses of diversification, character

evolution and other macroevolutionary processes. However, the accuracy and precision of the inferred

divergence times hinge upon many methodological choices that are often difficult or time-consuming to

justify (e.g., calibration strategies, prior distributions on node ages, clock models, etc.) [57-62].

Understanding the impact of each of these sources of uncertainty can be difficult. The sensitivity of

inferred ages is commonly explored by running analyses under different settings and summarizing the

results in tables or by stacking chronograms in order to visualize the relative position of nodes (see for

example [58, 59]). This approach discards a lot of information contained in Bayesian posterior

distributions of time-calibrated topologies. For example, patterns of correlation among branch lengths

(i.e., ways in which changes in a given branch length are accommodated by other regions of the tree)

are lost, and alternative ways in which node ages are attained (e.g., which of all descendant branches

expand as a given node becomes older) are not usually explored, even though they can potentially

represent very different evolutionary scenarios.

Here we analyze the sensitivity of node ages to alternative criteria to sample loci from

phylogenomic datasets, as well as different ways of modelling the patterns of molecular evolution

across sites and the rates of evolution among lineages. To do so, we introduce an approach to visualize

the distribution of chronograms in a multidimensional space of node ages, a chronospace, and measure

the overall effect of these decisions on inferred dates using multivariate statistical methods. Our results

reveal that divergence times obtained under site-homogeneous and site-heterogeneous models (such as

CAT+GTR+G) are broadly comparable. This happens despite the latter estimating higher levels of

sequence divergence and stretching branches in a non-isometric manner (Fig. S1). Site-heterogeneous

models have become common for the analysis of phylogenomic datasets spanning deep divergences, as

they exhibit reduced sensitivity to saturation and are thus more robust to long-branch attraction

artifacts [63-65]. These models are also central to the debate regarding contentious nodes in the tree of

life, including the position of ctenophores [66, 67] and the monophyly of arachnids [68, 69]. While the

use of CAT models for time calibration has become equally popular, the degree to which they impact

estimates of node ages has received less scrutiny. The lack of a meaningful effect uncovered here,

coupled with their high computational burden [70], questions their usefulness for time-scaling

phylogenies. A similar result was recently found when comparing site-homogeneous models with

different numbers of parameters [71], suggesting that relaxed clocks adjust branch rates in a manner

that buffers the effects introduced by using more complex (and computationally-intensive) models of

sequence evolution.

Similarly, the choice between different loci has a small effect on inferred ages, with little

evidence of a systematic difference between the divergence times supported by randomly-chosen loci

and those found using targeted sampling criteria, such as selecting genes for their phylogenetic signal,

usefulness, occupancy, or clock-likeness. A meaningful effect was restricted to a few ancient nodes (e.g.,

Echinodermata), for which clock-like genes suggested younger ages that are more consistent with fossil

evidence. This result provides direct empirical evidence that validates the selection of clock-like genes

for inferring deep histories of diversification, an approach so far supported mostly by simulations [62,

72]. At the same time, the choice of loci seems to be less relevant for younger nodes, at least when

employing large numbers of characters as done here. Finally, the choice between alternative clock

models induced differences in ages that were five to ten times stronger than those of other factors,

emphasizing the importance of either validating their choice (e.g., [73]) or—as done here—focusing on

results that are robust to them.

(c)   Echinoid origins and diversification

The origin and early diversification of crown group Echinoidea have always been considered to

have been determined (or at least strongly affected) by the P-T mass extinction [11, 12, 16, 20].

However, estimating the number of crown group members surviving the most severe biodiversity crisis

in the Phanerozoic [74] has been hampered by both paleontological and phylogenetic uncertainties [3,

10, 13-15, 18]. Our results establish with confidence that multiple crown group lineages survived and

crossed this boundary, finding for the first time a null posterior probability of the clade originating after

the extinction event. While the survival of three crown group lineages is slightly favored (Fig. S8),

discerning between alternative scenarios is still precluded by uncertainties in dating these early

divergences. Echinoid diversification during the Triassic was relatively fast (Fig. 4B) and involved rapid

divergences among its major clades. Even many lineages presently classified at the ordinal level trace

their origins to this initial pulse of diversification following the P-T mass extinction.

412  The late Paleozoic and Triassic origins inferred for the crown group and many euechinoid orders

413 prompts a re-evaluation of fossils from this interval of time. Incompletely known fossil taxa such as the

414 Pennsylvanian *?Eotiaris meurevillensis*, with an overall morphology akin to that of crown group

415 echinoids, has a stratigraphic range consistent with our inferred date for the origin of the echinoid

416 crown group [20]. Additionally, the Triassic fossil record of echinoids has been considered to be

417 dominated by stem group cidaroids, with the first euechinoids not known until the Late Triassic [17, 75].

418 However, the Triassic origins of many euechinoid lineages supported by our analyses necessitates that

419 potential euechinoid affinities should be re-considered for this diversity of Triassic fossils. This is

420 especially the case for the serpianotiarids and triadocidarids, abundant Triassic families variously

421 interpreted as cidaroids, euechinoids, or even stem echinoids [3, 17, 75, 76]. A reinterpretation of any of

422 these as euechinoids would suggest that the long-implied gap in the euechinoid record [16, 18] is caused

423 by our inability to correctly place these key fossils, as opposed to an incompleteness of the fossil record

424 itself.

425  While our phylogenomic approach is the first to resolve the position of all major cassiduloid

426 lineages, the inferred ages for many nodes within Neognathostomata remain in strong disagreement

427 with the fossil record. No Mesozoic fossil can be unambiguously assigned to either sand dollars or sea

428 biscuits, a surprising situation given the good fossilization potential and highly distinctive morphology of

429 these clades [19, 21, 25, 77]. While molecular support for a sister group relationship between scutelloids

430 and echinolampadids already implied this clade (Echinolampadacea) must have split from clypeasteroids

431 by the Late Cretaceous [3, 4, 14, 19], this still left open the possibility that the crown groups of sand

432 dollars and sea biscuits radiated in the Cenozoic. Under this scenario, the Mesozoic history of these

433 groups could have been comprised of forms lacking their distinctive morphological features,

434 complicating their correct identification. This hypothesis is here rejected, with the data unambiguously

435 supporting the radiation of both crown groups preceding the K-Pg mass extinction (Fig. 4C). While it

remains possible that these results are incorrect even after such a thorough exploration of the time-

calibration toolkit (see for example [62, 78]), these findings call for a critical reassessment of the

Cretaceous fossil record, and a better understanding of the timing and pattern of morphological

evolution among fossil and extant neognathostomates. For example, isolated teeth with an overall

resemblance to those of modern sand dollars and sea biscuits have been found in Lower Cretaceous

deposits [79], raising the possibility that other overlooked and disarticulated remains might close the

gap between rocks and clocks.

(d)   Conclusions

Although echinoid phylogenetics has long been studied using morphological data, the position

of several major lineages (e.g., aspidodiadematoids, micropygoids, salenioids, apatopygids) remained to

be confirmed with the use of phylogenomic approaches. Our work not only greatly expands the

available genomic resources for the clade, but finds novel resolutions for some of these lineages,

improving our understanding of their evolutionary history. The most salient aspect of our topology is the

splitting of the extant "cassiduloids" into two distantly related clades, one of which is composed

exclusively of apatopygids. This result is crucial to constrain the ancestral traits shared by the main

lineages of neognathostomates, helping unravel the evolutionary processes that gave rise to the unique

morphology of the sand dollars and sea biscuits [3, 24, 25].

Although divergence time estimation is known to be sensitive to many methodological

decisions, systematically quantifying the relative impact of these on inferred ages has rarely been done.

Here we propose an approach based on chronospaces (implemented in R [80] and available at

https://github.com/mongiardino/chronospace) that can help visualize key effects and determine the

sensitivity of node dates to different assumptions. Our results shed new light on the early evolutionary

history of crown group echinoids and its relationship with the P-T mass extinction event, a point in time

where the fossil record provides ambiguous answers. They also establish with confidence a Cretaceous

origin for the sand dollars and sea biscuits, preceding their first appearance in the fossil record by at

least 40 to 50 Ma, respectively (and potentially up to 65 Ma). These clades, therefore, join several well-

established cases of discrepancies between the fossil record and molecular clocks, such as those

underlying the origins of placental mammals [81] and flowering plants [82].

Methods

(a)   Sampling, bioinformatics, and matrix construction

This study builds upon previous phylogenomic matrices [3, 4], the last of which was augmented

through the addition of eight published datasets (mostly expanding outgroup sampling), as well as 17

novel echinoid datasets (15 transcriptomes and two draft genomes). For all novel datasets, tissue

sampling, DNA/RNA extraction, library preparation, and sequencing varied by specimen, and are

detailed in the Supplementary Information. Raw reads have been deposited in NCBI under Bioproject

accession numbers XXX and XXX. Final taxon sampling included 12 outgroups and 50 echinoids

(accession numbers and sampling details can be found in Table S1).

Raw reads for all transcriptomic datasets were trimmed or excluded using quality scores with

Trimmomatic v. 0.36 [83] under default parameters. Further sanitation steps were performed using the

Agalma 2.0 phylogenomic workflow [84], and datasets were assembled *de novo* with Trinity 2.5.1 [85].

For genomic shotgun sequences, adapters were removed with BBDuk

(https://sourceforge.net/projects/bbmap/), and UrQt v. 1.0.18 [86] was used to filter short reads (size <

50) and trim low-quality ends (score < 28). Datasets were then assembled using MEGAHIT v. 1.1.2 [87].

Draft genomes were masked using RepeatMasker v. 4.1.0 [88, 89], before obtaining gene predictions

with AUGUSTUS 3.2.3 [90]. A custom set of universal single-copy orthologs (USCOs) obtained from the

latest *Strongylocentrotus purpuratus* genome assembly (Spur v. 5.0) was employed as the training

dataset. Settings and further details of these analyses can be found in the Supplementary Information.

Multiplexed transcriptomes were sanitized from cross-contaminants using CroCo v. 1.1 [91], and

likely non-metazoan contaminants were removed using alien_index v. 3.0 [92] (removing sequences

with AI values > 45). Datasets were imported back into Agalma, which automated orthology inference

(as described in [84, 93]), gene alignment with MACSE [94], and trimming with GBLOCKS [95]. The amino

acid supermatrix was reduced using a 70% occupancy threshold, producing a final dataset of 1,346 loci

(327,695 sites). As a final sanitation step, gene trees were obtained using ParGenes v. 1.0.1 [96], which

performed model selection (minimizing the Bayesian Information Criterion) and inference using 100

bootstrap (BS) replicates. Trees were then used to remove outlier sequences with TreeShrink v. 1.3.1

[97]. We specified a reduced tolerance for false positives and limited removal to at most three terminals

which had to increase tree diameter by at least 25% (-q 0.01 -k 3 -b 25). Statistics for the supermatrix

and all assemblies can be found in Table S2.

(b)   Phylogenetic inference

Inference was performed under multiple probabilistic and coalescent-aware methods, known to

differ in their susceptibility to model violations. Coalescent-based inference was performed using the

summary method ASTRAL-III [98], estimating support as local posterior probabilities [99]. Among

concatenation approaches, we used Bayesian inference under an unpartitioned GTR+G model in

ExaBayes 1.5 [100]. Two chains were run for 2.5 million generations, samples were drawn every one

hundred, and the initial 25% was discarded as burn-in. We also explored maximum likelihood inference

with partitioned and unpartitioned models. For the former, the fast-relaxed clustering algorithm was

used to find the best-fitting model among the top 10% using IQ-TREE 1.6.12 [101, 102] (-m MFP+MERGE

-rclusterf 10 -rcluster-max 3000), and support was evaluated with 1,000 ultrafast BS replicates [103]. For

the latter, we used the LG4X+R model in RAxML-NG v. 0.5.1 [104] and evaluated support with 200

replicates of BS. Finally, we also implemented the site-heterogeneous LG+C60+F+G mixture model using

the posterior mean site frequency (PMSF) approach to provide a fast approximation of the full profile

mixture model [105], allowing the use of 100 BS replicates to estimate support. Given some degree of

topological conflict between the results of the other methods (see below), multiple guide trees were

used to estimate site frequency profiles, but the resulting phylogenies were identical.

Given conflicts between methods in the resolution of one particular node (involving the

relationships among Arbacioida, Salenioida, and the clade of Stomopneustoida + Camarodonta), all

methods were repeated after reducing the matrix to 500 and 100 loci selected for their phylogenetic

usefulness using the approach described in [3, 37] and implemented in the *genesortR* script

(https://github.com/mongiardino/genesortR). This approach relies on seven gene properties routinely

used for phylogenomic subsampling, including multiple proxies for phylogenetic signal—such as the

average BS and Robinson-Foulds (RF) similarity to a target topology—as well as several potential sources

of systematic bias (e.g., rate and compositional heterogeneities). Outgroups were removed before

calculating these metrics. RF similarity was measured to a species tree that had the conflicting

relationship collapsed so as not to bias gene selection in favor of any resolution. A principal component

analysis (PCA) of this dataset resulted in a dimension (PC 2, 17.6% of variance) along which phylogenetic

signal increased while sources of bias decreased (Fig. S9), and which was used for loci selection. For the

smallest dataset, we also performed inference under the site-heterogeneous CAT+GTR+G model using

PhyloBayes-MPI [106]. Three runs were continued for > 10,000 generations, sampling every two

generations and discarding the initial 25%. Convergence was confirmed given a maximum bipartition

discrepancy of 0.067 and effective sample sizes for all parameters > 150.

Two other approaches were implemented in order to assist in resolving the contentious node.

First, we implemented a likelihood-mapping analysis [107] in IQ-TREE to visualize the phylogenetic signal

for alternative resolutions of the quartet involving these three lineages (Arbacioida, Salenioida, and

Stomopneustoida + Camarodonta) and their sister clade (Irregularia; other taxa were excluded). Second,

we estimated the log-likelihood scores of each site in RAxML (using best-fitting models) for the two most

strongly supported resolutions found through likelihood mapping. These were used to calculate gene-

wise differences in scores, or δ values [108]. In order to search for discernable trends in the signal for

alternative topologies, genes were ordered based on their phylogenetic usefulness (see above) and the

mean per-locus δ values of datasets composed of multiples of 20 loci (i.e., the most useful 20, 40, etc.)

were calculated.

(c)   Time calibration

Node dating was performed using relaxed molecular clocks in PhyloBayes v4.1 using a fixed

topology and a novel set of 22 fossil calibrations corresponding to nodes from our newly inferred

phylogeny (listed in the Supplementary Information). Depending on the node, we enforced both

minimum and maximum bounds, or either one of these. A birth-death prior was used for divergence

times, which allowed for the implementation of soft bounds [109], leaving 5% prior probability of

divergences falling outside of the specified interval. We explored the sensitivity of divergence time

estimates to gene selection, model of molecular evolution, and clock prior. One hundred loci were

sampled from the full supermatrix according to four targeted sampling schemes: usefulness (calculated

as explained above, except incorporating all echinoderm terminals), phylogenetic signal (i.e., smallest RF

distance to species tree), clock-likeness (i.e., smallest variance of root-to-tip distances), and level of

occupancy. For clock-likeness, we only considered loci that lay within one standard deviation of the

mean rate (estimated by dividing total tree length by the number of terminals [110]), as this method is

otherwise prone to selecting largely uninformative loci (Fig. S10; [37]). A fifth sample of randomly

chosen loci was also evaluated. These five datasets were run under two models of molecular evolution,

the site-homogeneous GTR+G and the site-heterogeneous CAT+GTR+G, and both uncorrelated gamma

(UGAM) and autocorrelated log-normal (LN) clocks were implemented. The combination of these

settings (loci sampled, model of evolution, and clock prior) resulted in 20 analyses. For each, two runs

were continued for 20,000 generations, after which the initial 25% was discarded and the chains thinned

to every two generations (see log-likelihood trace plots in the electronic supplementary material, Fig.

S11). To explore the sensitivity of divergence times to these methodological decisions, 500 random

chronograms were sampled from each analysis (250 from each run), and their node dates were

subjected to multivariate analyses of variance (MANOVA) and between-group PCA (bgPCA) using

package *Morpho* [111] in the R statistical environment [80]. bgPCA involves the use of PCA on the

covariance matrix of group means (e.g., the mean ages of all nodes obtained under UGAM or LN clocks),

followed by the projection of original samples onto the obtained bgPC axes. The result is a

multidimensional representation of divergence times—a chronospace—rotated so as to capture the

distinctiveness of observations obtained under different settings. Separate bgPCAs were performed for

loci sampling strategy, clock prior and model of molecular evolution, and the proportion of total

variance explained by bgPC axes was interpreted as an estimate of the relative impact of these choices

on inferred times of divergence. Finally, lineage-through-time plots were generated using *ape* [112].

Author contributions

NMK, RM, and GWR conceived and designed the study. NMK, ASH, MFM, AFA, SEC, FA, OB, and

AK performed extractions, prepared libraries, and sequenced transcriptomic/genomic datasets. NMK

and AK generated *de novo* assemblies. JRT and RM designed fossil calibrations. NMK run all phylogenetic

analyses. NMK and JRT wrote the manuscript with input from all other authors. All authors gave final

approval for publication and agree to be held accountable for the work performed therein.

Data, code and materials

Raw transcriptomic reads have been deposited in NCBI under Bioproject accession number XXX.

Assemblies are available from the authors upon request. R code to plot chronospaces and explore the

effects of methodological decisions on divergence time estimation can be found at

https://github.com/mongiardino/chronospace.

Competing interests

We declare no competing interests.

Acknowledgments

We are grateful to the crew of R/Vs Antéa, Atlantis, Falkor and Mirai, HOV Alvin, and the crew

and PIs of BIOMAGLO deep-sea cruises (Laure Corbari, Karine Olu-Le Roy, Sarah Samadi) for assistance in

specimen collection. We would also like to thank Libby Liggins, Wilma Blom, Owen Anderson, Francisco

Solís-Marín, Carlos A. Conejeros-Vargas, and Jih-Pai Lin for the collection and shipment of specimens.

We gratefully acknowledge assistance from Thomas Saucède (Univ. Burgundy), Marc Eléaume (MNHN),

and Charlotte Seid (Scripps) in accessing, cataloging and processing museum specimens. Tim Ravasi

provided resources and collection facilities to GWR via King Abdullah University of Science and

Technology (KAUST). Institutional support was provided by the Central Research Laboratories and the

Department of Geology and Palaeontology at the Natural History Museum Vienna, Austria. This work

was supported by a Yale Institute for Biospheric Studies Doctoral Dissertation Improvement Grant and a

Society of Systematic Biologists Graduate Student Research Award to NMK. AK received funding from an

Austrian Science Fund project (FWF, project number P 29508-B25), FA from Agencia Nacional de

Investigación (ANID/PAI Inserción en la Academia, project number PAI79170033), and GWR and RM

from NSF grant DEB-2036186. JRT was supported by a Royal Society Newton International Fellowship

and a Leverhulme Trust Early Career Fellowship. NMK was supported by a Yale University fellowship.

References

1.   Donoghue PC, Benton MJ (2007) **Rocks and clocks: calibrating the Tree of Life using fossils and**

molecules *Trends in Ecology & Evolution* 22:424-431.

2.   Smith AB, Peterson KJ (2002) **Dating the time of origin of major clades: molecular clocks and**

the fossil record *Annual Review of Earth and Planetary Sciences* 30:65-88.

3.   Mongiardino Koch N, Thompson JR (2021) **A total-evidence dated phylogeny of Echinoidea**

combining phylogenomic and paleontological data *Systematic Biology* 70:421-439.

4.   Mongiardino Koch N, Coppard SE, Lessios HA, Briggs DE, Mooi R, Rouse GW (2018) **A**

phylogenomic resolution of the sea urchin tree of life *BMC Evolutionary Biology* 18:189.

5.   Kroh A, Mooi R (2021) **World Echinoidea Database**. Visited 13/July/2021.

http://www.marinespecies.org/echinoidea.

6.   Schultz HA (2015) **Echinoidea: with pentameral symmetry** Walter de Gruyter GmbH & Co KG.

7.   Edmunds PJ, Carpenter RC (2001) **Recovery of *Diadema antillarum* reduces macroalgal cover**

and increases abundance of juvenile corals on a Caribbean reef *Proceedings of the National Academy*

of Sciences 98:5067-5071.

8.   Harrold C, Pearse JS (1987) **The ecological role of echinoderms in kelp forests** In: Jangoux M,

Lawrence JM, editors. *Echinoderm Studies 2*. Rotterdam: AA Balkema. pp. 137–233.

9.   Zonneveld JP, Furlong CM, Sanders SC (2016) **Triassic echinoids (Echinodermata) from the**

Aksala Formation, north Lake Laberge, Yukon Territory, Canada *Papers in Palaeontology* 2:87-100.

10.   Thompson JR, Petsios E, Bottjer DJ (2017) **A diverse assemblage of Permian echinoids**

(Echinodermata, Echinoidea) and implications for character evolution in early crown group echinoids

Journal of Paleontology 91:767-780.

11.   Kier PM (1977) **Triassic echinoids** *Smithsonian Contributions to Paleobioliology* 30:1-88.

12.   Twitchett RJ, Oji T (2005) **Early Triassic recovery of echinoderms** *Comptes Rendus Palevol* 4:531-

542.

13.   Thompson JR, Erkenbrack EM, Hinman VF, McCauley BS, Petsios E, Bottjer DJ (2017)

Paleogenomics of echinoids reveals an ancient origin for the double-negative specification of

micromeres in sea urchins *Proceedings of the National Academy of Sciences* 114:5870-5877.

14.   Smith AB, Pisani D, Mackenzie-Dodds JA, Stockley B, Webster BL, Littlewood DTJ (2006) **Testing**

the molecular clock: molecular and paleontological estimates of divergence times in the Echinoidea

(Echinodermata) *Molecular Biology and Evolution* 23:1832-1851.

15.   Smith A, Hollingworth N (1990) **Tooth structure and phylogeny of the Upper Permian echinoid**

Miocidaris keyserlingi *Proceedings of the Yorkshire Geological Society* 48:47-60.

16.   Thompson JR, Petsios E, Davidson EH, Erkenbrack EM, Gao F, Bottjer DJ (2015) **Reorganization**

of sea urchin gene regulatory networks at least 268 million years ago as revealed by oldest fossil

cidaroid echinoid *Scientific Reports* 5:15541.

17.   Smith AB (2007) **Intrinsic versus extrinsic biases in the fossil record: contrasting the fossil**

record of echinoids in the Triassic and early Jurassic using sampling data, phylogenetic analysis, and

molecular clocks *Paleobiology* 33:310-323.

18.   Thompson JR, Hu S-x, Zhang Q-Y, Petsios E, Cotton LJ, Huang J-Y, et al (2018) **A new stem group**

echinoid from the Triassic of China leads to a revised macroevolutionary history of echinoids during

the end-Permian mass extinction *Royal Society Open Science* 5:171548.

19.   Kroh A, Smith AB (2010) **The phylogeny and classification of post-Palaeozoic echinoids** *Journal*

of Systematic Palaeontology 8:147-212.

20.   Thompson JR, Mirantsev GV, Petsios E, Bottjer DJ (2020) **Phylogenetic analysis of the**

Archaeocidaridae and Palaeozoic Miocidaridae (Echinodermata, Echinoidea) and the origin of crown

group echinoids *Papers in Palaeontology* 6:217-249.

21.   Kier PM (1982) **Rapid evolution in echinoids** *Palaeontology* 25:1-9.

22.   Barras CG (2008) **Morphological innovation associated with the expansion of atelostomate**

irregular echinoids into fine-grained sediments during the Jurassic *Palaeogeography,*

Palaeoclimatology, Palaeoecology 263:44-57.

23.   Saucède T, Mooi R, David B (2007) **Phylogeny and origin of Jurassic irregular echinoids**

(Echinodermata: Echinoidea) *Geological Magazine* 144:333-359.

24.   Hopkins MJ, Smith AB (2015) **Dynamic evolutionary change in post-Paleozoic echinoids and the**

importance of scale when interpreting changes in rates of evolution *Procedings of the National*

Academy of Sciences 112:3758-3763.

25.   Mooi R (1990) **Paedomorphosis, Aristotle's lantern, and the origin of the sand dollars**

(Echinodermata: Clypeasteroida) *Paleobiology* 16:25-48.

26.   Nebelsick JH (2020) **Ecology of clypeasteroids**. In: Lawrence JM, editor. *Sea Urchins: Biology and*

Ecology, 4th edition. Cambridge: Academic Press. pp. 315-331.

27.   Smith AB (2001) **Probing the cassiduloid origins of clypeasteroid echinoids using**

stratigraphically restricted parsimony analysis *Paleobiology* 27:392-404.

28.   Souto C, Mooi R, Martins L, Menegola C, Marshall CR (2019) **Homoplasy and extinction: the**

phylogeny of cassidulid echinoids (Echinodermata) *Zoological Journal of the Linnean Society* 187:622-

660.

29.   Kier PM (1962) **Revision of the cassiduloid echinoids** *Smithsonian Miscellaneous Collections*

144:1-262.

30.   Littlewood D, Smith A (1995) **A combined morphological and molecular phylogeny for sea**

urchins (Echinoidea: Echinodermata) *Philosophical Transactions of the Royal Society B: Biological*

Sciences 347:213-234.

31.   Mongiardino Koch N (2021) **Exploring adaptive landscapes across deep time: A case study**

using echinoid body size *Evolution* 75:1567-1581.

32.   Mooi R (1990) **Progenetic miniaturization in the sand dollar *Sinaechinocyamus*: implications**

for clypeasteroid phylogeny. In: De Ridder C, Dubois P, Lahaye M, Jangoux M, editors. *Echinoderm*

Research. Rotterdam: AA Balkema. pp. 137-143.

33.   Dunn CW, Zapata F, Munro C, Siebert S, Hejnol A (2018) **Pairwise comparisons across species**

are problematic when analyzing functional genomic data *Proceedings of the National Academy of*

Sciences 115:409-417.

34.   Smith SD, Pennell MW, Dunn CW, Edwards SV (2020) **Phylogenetics is the new genetics (for**

most of biodiversity) *Trends in Ecology & Evolution* 35:415-425.

35.   Kroh A (2020) **Phylogeny and classification of echinoids**. In: Lawrence J, editor. *Sea Urchins:*

Biology and Ecology, 4th edition. Cambridge: Academic Press. pp. 1-17.

36.   Lin J-P, Tsai M-H, Kroh A, Trautman A, Machado DJ, Chang L-Y, et al (2020) **The first complete**

mitochondrial genome of the sand dollar Sinaechinocyamus mai (Echinoidea: Clypeasteroida)

Genomics 112:1686-1693.

37.   Mongiardino Koch N (2021) **Phylogenomic subsampling and the search for phylogenetically**

reliable loci *Molecular Biology and Evolution* msab151. doi: 10.1093/molbev/msab151.

38.   Dell'Ampio E, Meusemann K, Szucsich NU, Peters RS, Meyer B, Borner J, et al (2014) **Decisive**

data sets in phylogenomics: lessons from studies on the phylogenetic relationships of primarily

wingless insects *Molecular Biology and Evolution* 31:239-249.

39.   Hillis DM, Heath TA, John KS (2005) **Analysis and visualization of tree space** *Systematic Biology*

54:471-482.

40.   Culhane AC, Perriere G, Considine EC, Cotter TG, Higgins DG (2002) **Between-group analysis of**

microarray data *Bioinformatics* 18:1600-1608.

41.   Cardini A, O'Higgins P, Rohlf FJ (2019) **Seeing distinct groups where there are none: spurious**

patterns from between-group PCA *Evolutionary Biology* 46:303-316.

42.   Zamora S, Lefebvre B, Álvaro JJ, Clausen S, Elicki O, Fatka O, et al (2013) **Cambrian echinoderm**

diversity and palaeobiogeography *Geological Society, London, Memoirs*. 38:157-171.

43.   Bottjer DJ, Davidson EH, Peterson KJ, Cameron RA (2006) **Paleogenomics of echinoderms**

Science 314:956-960.

44.   Pu JP, Bowring SA, Ramezani J, Myrow P, Raub TD, Landing E, et al (2016) **Dodging snowballs:**

Geochronology of the Gaskiers glaciation and the first appearance of the Ediacaran biota *Geology*

44:955-958.

45.   Reich M (2015) **Different pathways in early evolution of the holothurian calcareous ring**. In:

Zamora S, Rábano I, editors. *Progress in Echinoderm Palaeobiology*. Madrid: Instituto Geológico y

Minero de España. pp. 137-145.

46.   Miller AK, Kerr AM, Paulay G, Reich M, Wilson NG, Carvajal JI, et al (2017) **Molecular phylogeny**

of extant Holothuroidea (Echinodermata) *Molecular Phylogenetics and* Evolution 111:110-131.

47.   Rahman IA, Thompson JR, Briggs DE, Siveter DJ, Siveter DJ, Sutton MD (2019) **A new**

ophiocistioid with soft-tissue preservation from the Silurian Herefordshire Lagerstätte, and the

evolution of the holothurian body plan *Proceedings of the Royal Society* B 286:20182792.

48.   Rouse GW, Jermiin LS, Wilson NG, Eeckhaut I, Lanterbecq D, Oji T, et al (2013) **Fixed, free, and**

fixed: the fickle phylogeny of extant Crinoidea (Echinodermata) and their Permian–Triassic origin.

Molecular Phylogenetics and Evolution 66:161-181.

49.   Brosseau O, Murienne J, Pichon D, Vidal N, Eléaume M, Ameziane N (2012) **Phylogeny of**

Cidaroida (Echinodermata: Echinoidea) based on mitochondrial and nuclear markers *Organisms*

Diversity & Evolution 12:155-165.

50.   Durham JW, Melville R (1957) **A classification of echinoids** *Journal of Paleontology* 31:242-272.

51.   Suter SJ (1994) **Cladistic analysis of cassiduloid echinoids: trying to see the phylogeny for the**

trees *Biological Journal of the Linnean Society* 53:31-72.

52.   Kier PM (1966) **Cassiduloids**. In: Moore R, editor. *Treatise on Invertebrate Paleontology, Part U*.

Lawrence: University of Kansas Press. pp. 492-523.

53.   Mortensen T (1948) **A Monograph of the Echinoidea. IV, 1 Holectypoida, Cassiduloida**.

Copenhagen: CA Reitzel.

54.   King B (2020) **Bayesian tip-dated phylogenetics in paleontology: topological effects and**

stratigraphic fit *Systematic Biology* 70:283-294.

55.   Turner AH, Pritchard AC, Matzke NJ (2017) **Empirical and Bayesian approaches to fossil-only**

divergence times: a study across three reptile clades *PLoS One* 12:e0169885.

56.   Mongiardino Koch N, Garwood RJ, Parry LA (2021) **Fossils improve phylogenetic analyses of**

morphological characters *Proceedings of the Royal Society B: Biological Sciences* 288:2021004.

57.   Dos Reis M, Gunnell GF, Barba-Montoya J, Wilkins A, Yang Z, Yoder AD (2018) **Using**

phylogenomic data to explore the effects of relaxed clocks and calibration strategies on divergence

time estimation: primates as a test case *Systematic Biology* 67:594-615.

58.   Warnock RC, Yang Z, Donoghue PC (2011) **Exploring uncertainty in the calibration of the**

molecular clock *Biology Letters* 8:156-159.

59.   Dos Reis M, Thawornwattana Y, Angelis K, Telford MJ, Donoghue PC, Yang Z (2015) **Uncertainty**

in the timing of origin of animals and the limits of precision in molecular timescales *Current Biology*

25:2939-2950.

60.   Sauquet H (2013) **A practical guide to molecular dating**. *Comptes Rendus Palevol* 12:355-367.

61.   Carruthers T, Scotland RW (2021) **Uncertainty in divergence time estimation** *Systematic Biology*

70:855-861.

62.   Carruthers T, Sanderson MJ, Scotland RW (2020) **The implications of lineage-specific rates for**

divergence time estimation *Systematic Biology* 69:660-670.

63.   Lartillot N, Brinkmann H, Philippe H (2007) **Suppression of long-branch attraction artefacts in**

the animal phylogeny using a site-heterogeneous model *BMC Evolutionary Biology* 7:S4.

64.   Wang H-C, Minh BQ, Susko E, Roger AJ (2017) **Modeling site heterogeneity with posterior mean**

site frequency profiles accelerates accurate phylogenomic estimation Systematic Biology 67:216-235.

65.   Wang H-C, Li K, Susko E, Roger AJ (2008) **A class frequency mixture model that adjusts for site-**

specific amino acid frequencies and improves inference of protein phylogeny *BMC Evolutionary*

Biology 8:1-13.

66.   Li Y, Shen X-X, Evans B, Dunn CW, Rokas A (2021) **Rooting the animal tree of life** *Molecular*

Biology and Evolution msab170. doi: 10.1093/molbev/msab170.

67.   Pisani D, Pett W, Dohrmann M, Feuda R, Rota-Stabelli O, Philippe H, et al (2015) **Genomic data**

do not support comb jellies as the sister group to all other animals *Proceedings of the National*

Academy of Sciences 112:15402-15407.

68.   Howard RJ, Puttick MN, Edgecombe GD, Lozano-Fernandez J (2020) **Arachnid monophyly:**

Morphological, palaeontological and molecular support for a single terrestrialization within

Chelicerata *Arthropod Structure & Development* 59:100997.

69.   Ballesteros JA, Sharma PP (2019) **A critical appraisal of the placement of Xiphosura**

(Chelicerata) with account of known sources of phylogenetic error *Systematic Biology* 68:896-917.

70.   Whelan NV, Halanych KM (2016) **Who let the CAT out of the bag? Accurately dealing with**

substitutional heterogeneity in phylogenomic analyses *Systematic Biology* 66:232-255.

71.   Tao Q, Barba-Montoya J, Huuki LA, Durnan MK, Kumar S (2020) **Relative efficiencies of simple**

and complex substitution models in estimating divergence times in phylogenomics *Molecular Biology*

and Evolution 37:1819-1831.

72.   Smith SA, Brown JW, Walker JF (2018) **So many genes, so little time: A practical approach to**

divergence-time estimation in the genomic era *PLoS One* 13:e0197433.

73.   Tao Q, Tamura K, U. Battistuzzi F, Kumar S (2019) **A machine learning method for detecting**

autocorrelation of evolutionary rates in large phylogenies *Molecular Biology and Evolution* 36:811-824.

74.   Raup DM, Sepkoski JJ (1982) **Mass extinctions in the marine fossil record** *Science* 215:1501-

1503.

75.   Kier PM (1984) **Echinoids from the Triassic (St. Cassian) of Italy, their lantern supports, and a**

revised phylogeny of Triassic echinoids *Smithsonian Contributions to Paleobiology* 56:1-41.

76.   Smith AB (1994) **Triassic echinoids from Peru** *Palaeontographica Abteilung A* 233:177-202.

77.   Kier PM (1977) **The poor fossil record of the regular echinoid** *Paleobiology* 3:168-174.

78.   Field D, Berv J, Hsiang A, Lanfear R, Landis M, Dornburg A (2020) **Timing the extant avian**

radiation: the rise of modern birds, and the importance of modeling molecular rate variation *Bulletin*

of the American Museum of Natural History 440:150-181.

79.   Caramés A, Adamonis S, Concheyro A, Remírez M (2019) **New finding of regular echinoid**

elements and microfossils from the Pilmatué Member, Agrio Formation (Early Cretaceous), Neuquén

Basin, Argentina. In: Cusminsky G, Bernasconi E, Concheyro G, editors. *Advances in South American*

micropaleontology. Springer. pp. 1-20.

80.   R Core Team (2019) **R: A Language and Environment for Statistical Computing**. Vienna, Austria:

R Foundation for Statistical Computing.

81.   Ronquist F, Lartillot N, Phillips MJ (2016) **Closing the gap between rocks and clocks using total-**

evidence dating *Philosophical Transactions of the Royal Society B* 371:20150136.

82.   Coiro M, Doyle JA, Hilton J (2019) **How deep is the conflict between molecular and fossil**

evidence on the age of angiosperms? *New Phytologist* 223:83-99.

83.   Bolger AM, Lohse M, Usadel B (2014) **Trimmomatic: a flexible trimmer for Illumina sequence**

data *Bioinformatics* 30:2114-2120.

84.   Dunn CW, Howison M, Zapata F (2013) **Agalma: an automated phylogenomics workflow** *BMC*

Bioinformatics 14:330.

85.   Grabherr MG, Haas BJ, Yassour M, Levin JZ, Thompson DA, Amit I, et al (2011) **Full-length**

transcriptome assembly from RNA-Seq data without a reference genome *Nature Biotechnology*

29:644-652.

86.   Modolo L, Lerat E (2015) **UrQt: an efficient software for the Unsupervised Quality trimming of**

NGS data *BMC Bioinformatics* 16:1-8.

87.   Li D, Liu C-M, Luo R, Sadakane K, Lam T-W (2015) **MEGAHIT: an ultra-fast single-node solution**

for large and complex metagenomics assembly via succinct de Bruijn graph *Bioinformatics* 31:1674-

1676.

88.   Smit A, Hubley R, Green P (2013-2015) **RepeatMasker Open-4.0**.

89.   Hoff KJ, Stanke M (2019) **Predicting genes in single genomes with AUGUSTUS**. *Current Protocols*

in Bioinformatics. 65:e57.

90.   Stanke M, Keller O, Gunduz I, Hayes A, Waack S, Morgenstern B (2006) **AUGUSTUS: ab initio**

prediction of alternative transcripts *Nucleic Acids Research* 34:435-439.

91.   Simion P, Belkhir K, François C, Veyssier J, Rink JC, Manuel M, et al (2018) **A software tool**

'CroCo'detects pervasive cross-species contamination in next generation sequencing data. *BMC*

Biology 16:28.

92.   Ryan JF (2014) **Alien Index: identify potential non-animal transcripts or horizontally**

transferred genes in animal transcriptomes. doi: 10.5281/zenodo.21029.

93.   Guang A, Howison M, Zapata F, Lawrence C, Dunn CW (2021) **Revising transcriptome**

assemblies with phylogenetic information *PLos One* 16:e0244202.

94.   Ranwez V, Harispe S, Delsuc F, Douzery EJ (2011) **MACSE: Multiple Alignment of Coding**

SEquences accounting for frameshifts and stop codons *PLoS One* 6:e22594.

95.   Talavera G, Castresana J (2007) **Improvement of phylogenies after removing divergent and**

ambiguously aligned blocks from protein sequence alignments *Systematic Biology* 56:564-577.

96.   Morel B, Kozlov AM, Stamatakis A (2018) **ParGenes: a tool for massively parallel model**

selection and phylogenetic tree inference on thousands of genes *Bioinformatics* 35:1771-1773.

97.   Mai U, Mirarab S (2018) **TreeShrink: fast and accurate detection of outlier long branches in**

collections of phylogenetic trees *BMC Genomics* 19:272.

98.   Zhang C, Rabiee M, Sayyari E, Mirarab S (2018) **ASTRAL-III: polynomial time species tree**

reconstruction from partially resolved gene trees *BMC Bioinformatics* 19:153.

99.   Sayyari E, Mirarab S (2016) **Fast coalescent-based computation of local branch support from**

quartet frequencies *Molecular Biology and Evolution* 33:1654-1668.

100.   Aberer AJ, Kobert K, Stamatakis A (2014) **ExaBayes: massively parallel Bayesian tree inference**

for the whole-genome era *Molecular Biology and Evolution* 31:2553-2556.

101.   Nguyen L-T, Schmidt HA, von Haeseler A, Minh BQ (2014) **IQ-TREE: a fast and effective**

stochastic algorithm for estimating maximum-likelihood phylogenies *Molecular Biology and Evolution*

32:268-274.

102.   Kalyaanamoorthy S, Minh BQ, Wong TK, von Haeseler A, Jermiin LS (2017) **ModelFinder: fast**

model selection for accurate phylogenetic estimates *Nature Methods* 14:587.

103.   Hoang DT, Chernomor O, Von Haeseler A, Minh BQ, Vinh LS (2017) **UFBoot2: improving the**

ultrafast bootstrap approximation *Molecular Biology and Evolution* 35:518-522.

104.   Kozlov AM, Darriba D, Flouri T, Morel B, Stamatakis A (2019) **RAxML-NG: a fast, scalable and**

user-friendly tool for maximum likelihood phylogenetic inference *Bioinformatics* 35:4453-4455.

105.   Wang H-C, Minh BQ, Susko E, Roger AJ (2018) **Modeling site heterogeneity with posterior mean**

site frequency profiles accelerates accurate phylogenomic estimation *Systematic Biology* 67:216-235.

106.   Lartillot N, Rodrigue N, Stubbs D, Richer J (2013) **PhyloBayes MPI. Phylogenetic reconstruction**

with infinite mixtures of profiles in a parallel environment *Systematic Biology* 62:611-615.

107.   Strimmer K, Von Haeseler A (1997) **Likelihood-mapping: a simple method to visualize**

phylogenetic content of a sequence alignment *Proceedings of the National Academy of Sciences*

94:6815-6819.

108.   Shen X-X, Hittinger CT, Rokas A (2017) **Contentious relationships in phylogenomic studies can**

be driven by a handful of genes *Nature Ecology &* Evolution 1:0126.

109.   Yang Z, Rannala B (2006) **Bayesian estimation of species divergence times under a molecular**

clock using multiple fossil calibrations with soft bounds *Molecular Biology and Evolution* 23:212-226.

110.   Telford MJ, Lowe CJ, Cameron CB, Ortega-Martinez O, Aronowicz J, Oliveri P, et al (2014)

Phylogenomic analysis of echinoderm class relationships supports Asterozoa *Proceedings of the Royal*

Society of London B: Biological Sciences 281:20140479.

111.   Schlager S (2017) **Morpho and Rvcg–Shape Analysis in R**. In: Zheng G, Li S, Székely GJ, editors.

Statistical shape and deformation analysis. Elsevier. pp. 217-256.

112.   Paradis E, Schliep K (2018) **ape 5.0: an environment for modern phylogenetics and**

evolutionary analyses in R *Bioinformatics* 35:526-528.

Supplementary Information for:

860   N Mongiardino Koch, JR Thompson, AS Hatch, MF McCowin, AF Armstrong, SE Coppard, F Aguilera, O

861   Bronstein, A Kroh, R Mooi, GW Rouse – Phylogenomic analyses of echinoid diversification prompt a re-

862   evaluation of their fossil record

Sequencing and further bioinformatic details

Apatopygus recens, *Aspidodiadema hawaiiense*, *Fellaster zelandiae*, *Histocidaris variabilis*,

Rhyncholampas pacificus, *Sinaechinocyamus mai*, *Stereocidaris neumayeri*, *Tromikosoma hispidum*.

Tissue subsamples were finely chopped with a scalpel and preserved in RNA*later* (Ambion) buffer

solution for 1 day at 4°C to allow the RNA*later* to effectively penetrate the tissues, followed by long-

term storage at -80°C until RNA extraction. Total RNA was extracted from 1.5 mm Triple-Pure High

Impact Zirconium beads (Benchmark Scientific) in Trizol (Ambion), using Direct-zol RNA Miniprep Kit

(Zymo Research) with in-column DNase I incubation to remove genomic DNA. Prior to mRNA capture,

total RNA concentration was estimated using Qubit RNA HS Assay Kit (Invitrogen; range = 8.33-96

ng/μL), and quality was assessed using either High Sensitivity RNA ScreenTape or RNA ScreenTape with

an Agilent 4200 TapeStation. Mature mRNA was isolated from total RNA and libraries were prepared

using KAPA mRNA HyperPrep Kit (KAPA Biosystems) following the manufacturer's instructions (including

sample customization based on total RNA quantity and quality values), targeting an insert size circa 500

base pairs (bp), and using custom 10-nucleotide Illumina TruSeq style adapters [113]. Post-amplification,

DNA concentration was estimated using Qubit dsDNA BR Assay Kit (Invitrogen; range = 6.06-13.4 ng/μL).

Concentration, quality, and molecular weight distribution of libraries (range = 547-978 bp) was also

assessed using Genomic DNA ScreenTape with an Agilent 4200 TapeStation. Ten libraries (including two

annelids not employed here) were sequenced on one lane of a multiplexed run using NovaSeq S4

platform with 100 bp paired-end reads at the IGM Genomics Center (University of California San Diego).

The assembled transcriptomes were sanitized from cross-contaminant reads product of

multiplexed sequencing using CroCo v. 1.1 [91]. These eight datasets, as well as two annelid

transcriptomes not employed in this study but sequenced in the same lane, were incorporated.

Transcripts considered over or under-expressed across samples (as defined by default parameters),

were kept. This resulted in an average removal of 1.26% of assembled transcripts (range: 0.4% for

Aspidodiadema to 3.06% for *Histocidaris*).

Clypeaster japonicus, *Encope emarginata*, *Leodia sexiesperforata*, *Peronella japonica*. Eggs were

collected from a single female specimen of each species and immediately placed into RNA later for

preservation. The samples were left in RNAlater at 4°C for at least 1 day and then transferred to a -80°C

freezer until RNA extraction. RNA extraction was performed using Trizol and was then treated with

Ambion's Turbo DNA-free kit. RNA was quantified using both Nanodrop and an Agilent 2100

Bioanalyzer. Only RNA samples with an RNA integrity number (RIN) of > 7 were used. RNA-Seq libraries

were prepared using the NEB Ultra Directional kit using the standard protocol and multiplexed with 6bp

NEB multiplex primers. Paired-end 100 libraries were generated from the resultant libraries at the UC

Berkeley Genome Center using an Illumina HighSeq 2500.

Bathysalenia phoinissa, *Micropyga tuberculata*: The material was collected by the deep-sea

cruise BIOMAGLO [114] conducted in 2017 jointly by the French National Museum of Natural History

(MNHN) as part of the Tropical Deep-Sea Benthos program, the French Research Institute for

Exploitation of the Sea (IFREMER), the "Terres Australes et Antarctiques Françaises" (TAAF), the

Departmental Council of Mayotte and the French Development Agency (AFD), with the financial support

of the European Union (Xe FED). Specimens were sorted into taxonomic bins at class level and

collectively fixed in 70% ethanol at room temperature. Following taxonomic identification, tissue

samples were obtained using sterilized forceps and disposable scalpel blades. Tissue subsamples were

collected in high-purity 96% ethanol and stored at -40°C until DNA extraction. Total genomic DNA was

extracted using the DNeasy Blood and Tissue Kit (QIAGEN) following the manufacturer's instructions,

complemented by the addition of 4 µl RNAse A (QIAGEN, 100 mg/ml) for RNA digestion. Elution buffer

volume was reduced to 60 µl and the elution step repeated twice per sample in order to maximize DNA

concentration and yield. After collection of the first elution, spin-column membranes were rinsed again

twice with fresh elution buffer (40 µl) to further increase yield. Elutions were collected separately, with

the first one used for sequencing and the second for quality control. For the latter, a portion of the

mitochondrial cytochrome c oxidase subunit I (COI) was amplified using PCR. Amplifications were

conducted with TopTaq DNA Polymerase (QIAGEN) using 1 µl of extracted genomic DNA (approx. 10–15

ng). Details on primers and protocols can be found in [115]. PCR products were visualized on a 1%

agarose gel, purified using ExoSAP-IT (Affymetrix), and sequenced at Microsynth GmbH (Vienna, Austria)

with the same primers. Sequences are deposited in GenBank under the accession numbers XXX and XXX.

Prior to library preparation, total genomic DNA concentration was estimated using Qubit DNA

BR Assay Kit (Invitrogen) in order to determine library preparation strategy. This step was carried out by

Macrogen (Korea), using an Illumina TruSeq DNA PCR-free kit (for *Micropyga*) and an Illumina TruSeq

Nano DNA kit (for *Bathysalenia*). Library preparation followed the manufacturer's instructions, targeting

an insert size of 350 base pairs (bp). The libraries were sequenced by Macrogen (Korea) on a shared lane

of a multiplexed run using an Illumina HiSeq X instrument with 150 bp paired-end reads. Sequencing

depth was 134.1 and 161.3 million reads for *Bathysalenia* and *Micropyga*, respectively. Pre-processing of

Illumina shotgun data was carried out by removal of adapters using BBDuk

(https://sourceforge.net/projects/bbmap/) with settings: `minlen=50 ktrim=r k=23 mink=11`

`tpe tbo`); followed by quality trimming of reads using UrQt v. 1.0.18 [86], discarding regions with

phred quality score < 28 and sequences of size < 50. Assembly of the data was carried out with MEGAHIT

v. 1.1.2 [87], in an iterative approach followed by assessment of assembly quality using BUSCO v. 3.0.2.

scores [116] using the metazoan dataset ODB v. 9 [117]. Final BUSCO scores of the draft assemblies

were 28.4% (S:27.8%, D:0.6%, F:48.2%, M:23.4%, n:978) for *Bathysalenia* and 17.2% (S:16.8%, D:0.4%,

F:59.0%, M:23.8%, n:978) for *Micropyga*. As recommended by Hoff and Stanke [89], draft genomes were

masked with RepeatMasker v. 4.1.0 [88] prior to gene prediction, using settings: `-norna -xsmall`;

resulting in 3.82 % and 2.9% masked bases for *Bathysalenia* and *Micropyga*, respectively. Gene

prediction was carried out using AUGUSTUS v. 3.2.3 [90] using settings: `--strand=both --`

`singlestrand=false --genemodel=partial --codingseq=on --sample=0 --`

`alternatives-from-sampling=false --exonnames=on --softmasking=on`. This step

relied on a training dataset composed of universal single-copy orthologs (USCOs) derived from the most

recent *Strongylocentrotus purpuratus* genome (Spur v.5).

Echinothrix calamaris, *Tripneustes gratilla*. Tissue samples were taken from live sea urchins

housed in laboratory aquaria and total RNA was immediately extracted from 150 mg of tissue using a

PureLink™ RNA extraction kit (Invitrogen™). The quality of total RNA was assessed on a BioAnalyzer

2100 (Agilent Technologies, Santa Clara, CA, USA) to ensure a RIN > 9 for all samples. Mature mRNA was

extracted from 1ug of total RNA and cDNA libraries were constructed using a TruSeq kit (Illumina).

Quality control of libraries was assessed on a BioAnalyzer and quantification measured using qPCR.

NEBNext Multiplex adaptors were added via ligation, and the cDNA libraries were sequenced at Genome

Quebec, McGill University. Three libraries were multiplex sequenced on one lane of Illumina at a

concentration of 8 pM per cDNA library using HiSeq2500 transcriptome sequencing to generate 125bp

paired-end reads. This resulted in approximately 80 million reads for each transcriptome.

Tetrapygus niger. Adult specimens were collected and transported alive to the University of

Concepcion. Tissue samples (female gonad and tube feet) from one individual were finely chopped with

a scalpel, and total RNA was extracted immediately using TriReagent (Sigma), following the

manufacturer's instructions. Total RNA concentration was estimated using QuantiFluor® RNA System

(111.84-339.04 ng/µL) and quality was assessed using Agilent RNA 6000 Nano kit with an Agilent 2100

TapeStation (RIN: 5.4-8.4). cDNA libraries were prepared at Genoma Mayor SpA (Chile) using TruSeq

Stranded mRNA Kit (Illumina) and sequenced on a multiplexed run using Illumina HiSeq 4000 platform

with 150 paired-end reads, resulting in 54.1 M reads for gonad female and 52.7 M reads for tube feet,

respectively. These two transcriptomic datasets were combined before performing all steps of

bioinformatic processing.

Eucidaris metularia. A single specimen was preserved in RNA*later* after being starved overnight

and kept for 1 day at 4°C before long-term storage at -80°C. Total RNA was extracted using Direct-zol

RNA Miniprep Kit (with in-column DNase treatment; Zymo Research) from Trizol. mRNA was isolated

with Dynabeads mRNA Direct Micro Kit (Invitrogen). RNA concentration was estimated using Qubit RNA

broad range assay kit, and quality was assessed using RNA ScreenTape with an Agilent 4200 TapeStation.

Values were used to customize downstream protocols following manufacturers' instructions. Library

preparation was performed with KAPA-Stranded RNA-Seq kits, targeting an insert size in the range of

200–300 base pairs. The quality and concentration of libraries were assessed using DNA ScreenTape.

The library was sequenced in a multiplexed pair-end runs using Illumina HiSeq 4000 with 7 other

libraries in the same lane (all previously published in [4]). In order to minimize read crossover, 10 bp

sequence tags designed to be robust to indel and substitution errors were employed [113].

The assembled transcriptome was filtered from cross-contaminant reads product of multiplexed

sequencing using CroCo v. 1.1 [91]. Seven other transcriptomes sequenced together were also

employed, and results of this step are reported in Fig. S1 of Mongiardino Koch & Thompson [3].

 Fossil constraints

All clade names used below refer to the corresponding crown groups.

<u>Ambulacraria (Echinodermata-Hemichordata divergence)</u> – **Younger bound**: 518 Ma, Middle Atdabanian

(age 3, Series 3 of Cambrian). **Older bound**: 636.1 Ma, Lantian Biota, Ediacaran. **Reference:** [118]. **Notes:**

The root of our tree is constrained with a younger bound concurrent with the earliest occurrence of

stereom in the fossil record (see below). The older bound is based on the maximum age of the Lantian

Biota, a Lagerstätte lacking any trace of eumetazoan fossils.

<u>Echinodermata (Pelmatozoan-Eleutherozoan divergence)</u> – **Younger bound**: Unconstrained. **Older**

bound: 515.5 Ma, Middle Atdabanian (age 3, Series 3 of Cambrian). **Reference**: [118]. **Notes:** This

divergence (which represents the divergence of crown group echinoderms), has an older bound based

upon the earliest occurrence of stereom in the fossil record. Stereom constitutes an echinoderm

synapomorphy readily recognizable in the fossil record due to its unique mesh-like structure [43]. The

first records of stereom point to a sudden and global appearance within the middle Atdabanian [42].

Note that this is also used as younger bound for the divergence between echinoderms and

hemichordates.

<u>Echinozoa (Echinoidea-Holothuroidea divergence)</u> – **Younger bound:** 469.96 Ma, Top of

Pseudoclamacograptus decorates graptolite zone. **Older bound:** 461.95 Ma, Top Floian. **Reference:** [119,

120]. **Notes:** The oldest crown eleutherozoan fossils are disarticulated holothurian calcareous ring

elements which are known from the Red *Orthoceras* limestone of Sweden. This falls within the *P.*

decorates graptolite zone.

<u>Asterozoa (Ophiuroidea-Asteroidea divergence)</u> – **Younger bound**: 521 Ma, Top of *Psigraptus jacksoni*

Zone. **Older bound**: 480.55 Ma, Base of Series 2 Cambrian Period. **Reference**: [119, 120]. **Taxon**:

Maydenia roadsidensis. **Notes**: The divergence of asterozoan classes is calibrated based upon the

stratigraphic occurrence of *Maydenia roadsidensis*, the oldest asterozoan, which is a stem group

ophiuroid [121] and occurs in the *Psigraptus jacksoni* Zone of the Floian.

Asteroidea – **Younger bound:** 239.10 Ma, Top Fassanian, Ladinian, Mo3, *nodosus* biozone, Triassic.

Older bound: 252.16 Ma, P-T Boundary. **Reference**: [122]. **Taxon**: *Trichasteropsis weissmanni*. **Notes**:

Fossil evidence suggests the divergence of the asterozoan crown group took place sometime in the Early

or Middle Triassic [122]. Based upon its phylogenetic position and stratigraphic occurrence as the oldest

crown group asteroid, we use the forcipulatacean *T. weissmanni* from the Middle Triassic Muschelkalk

as the soft bound on this divergence.

Crinoidea – **Younger bound**: 247.06 Ma, Top Spathian (Paris Biota, Lower Shale unit, Thaynes Group,

Spathian, Triassic). **Older bound**: Unconstrained. **Reference**: [123]. **Taxon**: *Holocrinus*. **Notes**: Crown

group Crinoidea is difficult to define, though a rapid post-Palaeozoic morphological diversification is

supported by fossil evidence. We use *Holocrinus* from the Early Triassic of the Thaynes group to calibrate

the younger bound on the divergence of the crown group (i.e., the split between Isocrinida and all other

extant crinoids).

Ophiuroidea (Euryophiurida-Ophintegrida divergence) – **Younger bound**: 247.06 Ma, Top Spathian (Paris

Biota, Lower Shale unit, Thaynes Group, Spathian, Triassic). **Older bound**: Unconstrained. **Reference**:

[124]. **Taxon**: *Shoshonura brayardi*. **Notes**: *S. brayardi* from the Spathian Thaynes Group of the Western

USA is a member of the crown group ophiuroid clade Ophiodermatina. It is thus the currently oldest

described member of the ophiuroid crown group, setting its minimum age of divergence.

Pneumonophora (Holothuriida-Neoholothuriida divergence) – **Younger bound**: 259.8 Ma, S*pinosus* zone

of Early Ladinian (used base Ladinian). **Older bound**: 241.5 Ma, Base Wuchiapingian. **Reference**: [46,

119]. **Notes**: The oldest (undescribed) holothuriid calcareous ring ossicles are from the *Spinosus* zone of

the Ladinian of Germany and calibrate its divergence from neoholothuriids.

Echinoidea (Cidaroidea-Euechinoidea divergence) – **Younger bound**: 298.9 Ma, Base Carnian. **Older**

bound: 237 Ma, Base Permian. **Reference**: [125]. **Taxon**: *Triassicidaris? ampezzana*. **Notes**: The recent

phylogenetic analysis of Mongiardino Koch and Thompson [3] found many traditional early crown group

echinoids as members of the stem group. Given this result, *Triassicidaris? ampezzana*, a cidaroid which

is known from the St. Cassian beds of Italy, becomes the current oldest crown group echinoid.

Pedinoida-Aspidodiadematoida – **Younger bound**: 209.46 Ma, Top of Norian. **Older bound**: 252.16 Ma,

P-T boundary. **Reference**: See discussion in supplement of Thompson et al. [13]. **Taxon**: *Diademopsis* ex.

gr. *heberti*. **Notes**: The oldest euechinoid, *Diademopsis* ex. gr. *heberti*, which is a pedinoid, is known

from the Norian of Peru. This fossil calibrates the younger bound on the pedinoid-aspidodiadematoid

divergence.

Carinacea (Echinacea-Irregularia divergence) – **Younger bound**: 234.5 Ma, Top Sinemurian. **Older**

bound: 190.8 Ma, Top Julian 1 ammonoid zone of Norian. **Reference**: Li et al. [126]. **Taxon**:

Jesionekechinus. **Notes**: The divergence of Carinacea is calibrated based upon the oldest irregular

echinoid, *Jesionekechinus*, which occurs above the Sinemurian of New York Canyon, Nevada. For a more

detailed discussion of this divergence see the supplement of Thompson et al. [13].

Salenioida-(Camarodonta+Stomopneustoida) divergence– **Younger bound**: 228.35 Ma, Base Hettangian.

Older bound: 201.3 Ma, Top Carnian. **Reference**: [14]. **Taxon**: *Acrosalenia chartroni*. **Notes**: The stem

group salenioid *Acrosalenia chartroni* from the Hettangian of France is the earliest representative of

Echinacea. Given the topology recovered by our analyses, this fossil was used to calibrate the divergence

between salenioids and the clade composed of stomopneustoids and camarodonts.

Stomopneustoida-Camarodonta divergence – **Younger bound**: 201.3 Ma, Top Pliensbachian. **Older**

bound: 182.7 Ma, Base Jurassic. **Reference**: See discussion in supplement of Thompson et al. [13].

Taxon: Stomechinids from Morocco like *Diplechinus hebbriensis* [127]. **Notes**: The oldest stomechinids,

which are stem group stomopneustoids, are from the Early Jurassic of Morocco.

Neognathostomata-Atelostomata divergence – **Younger bound**: 234.5 Ma, Base of Toarcian. **Older**

bound: 182.7 Ma, St. Cassian Beds, bottom of Julian 2 ammonoid Zone. **Reference**: [14, 126]. **Taxon**:

Younger bound is set by *Galeropygus sublaevis* (older bound is relatively uninformative). **Notes**: The

base of the Toarcian (which contains the oldest neognathostomate *G. sublaevis*) is used as the lower

bound on the divergence between the neognathostomata and all other crown irregular echinoids. Note

that depending on the position of the unsampled echinoneoids, this split might also correspond to the

origin of crown group Irregularia (see recent topologies recovered by Lin et al. [36] and Mongiardino

Koch & Thompson [3]).

Atelostomata (Holasteroida-Spatangoida divergence) – **Younger bound**: 137.68 Ma, Bottom of

Campylotoxus zone in Valanginian. **Older bound**: 201.3 Ma, Base of Jurassic. **Reference**: [128]. **Taxon**:

Younger bound is based on *Toxaster* (older bound is loosely uninformative). **Notes**: The toxasterids are

stem group spatangoids that calibrate the divergence between spatangoids and holasteroids. The oldest

Toxaster are from the *Campylotoxus* zone in the Valanginian, and set the younger bound on the

divergence.

Echinolampadacea (Cassiduloida-Scutelloida divergence) – **Younger bound**: 113 Ma, Base Albian. **Older**

bound: 145 Ma, Base Cretaceous. **Reference**: [129]. **Taxon**: Younger bound is based on *Eurypetalum*

rancheriana (older bound is loosely uninformative). **Notes**: The oldest member of Echinolampadacea is

the faujasiid *Eurypetalum rancheriana* (originally placed in the genus *Faujasia* and later transferred, see

[28, 29]) from the Late Albian of Colombia.

Scutelloida (Scutelliformes-Laganiformes divergence) – **Younger bound**: 56.0 Ma, Bottom of Eocene.

Older bound: Unconstrained. **Reference**: [14, 130]. **Taxon**: Younger bound is based on *Eoscutum*

doncieuxi. **Notes**: We use the base of the Eocene as the younger bound on the divergence between

scutelliforms and laganiforms. This is based upon the occurrence of *E. doncieuxi*, the oldest scutelliform,

which is known from the Lower Eocene. The older bound is left unconstrained in order to account for

the current uncertainty in the origin of the clade (i.e., allow for Mesozoic ages).

Leodia-(*Encope+Mellita*) divergence – **Younger bound**: 23.03 Ma, Base Miocene. **Older bound**: 33.9 Ma,

Base of Oligocene. **Reference**: [131, 132]. **Taxon**: Younger bound is based on *Encope michoacanensis*

(older bound is loosely uninformative). **Notes**: This node is constrained using the oldest representative

of these three genera of mellitids.

Clypeasteroida – **Younger bound**: 47.8 Ma, Base Lutetian. **Older bound**: Unconstrained. **Reference**:

[133, 134]. **Taxon**: Younger bound is based on *Clypeaster calzadai* and *Clypeaster moianensis*. **Notes**:

The oldest known fossil species of Clypeasteroida (the clypeasterids *C. calzadai* and *C. moianensis*) are

from the middle Eocene, upper Lutetian (Biarritzian) of Cataluña, Spain. The older bound is left

unconstrained in order to account for the current uncertainty in the origin of the clade (i.e., allow for

Mesozoic ages).

Strongylocentrotidae-Toxopneustidae divergence – **Younger bound**: 44.4 Ma, Eocene Planktonic Zones

E9-E12. **Older bound**: 56.0 Ma, Base Eocene. **Reference**: [135, 136]. **Taxon**: Younger bound is based on

Lytechinus axiologus (older bound is loosely uninformative). **Notes**: The oldest member of the clade

comprising toxopneustids and strongylocentrotids is the toxopneustid *Lytechinus axiologus.* This taxon is

known from the Eocene Yellow Limestone Group of Jamaica. The Yellow Limestone group represents the

Eocene Planktonic Zones E9-E12.

Supplementary Figures and Tables

1092

Figure S1: Estimated branch lengths across different models of molecular evolution. Different site-

homogeneous models (left) infer similar levels of divergence, and the choice between them induces

little distortion in the general tree structure. Site-heterogeneous models on the other hand, not only

infer a larger degree of divergence between terminals relative to site-homogeneous ones (center and

left), but they also distort the tree (i.e., impose a non-isometric stretching), with branch lengths

connecting outgroup taxa expanding much more than those within the ingroup clade.

1099

Figure S2: The ten most sensitive node dates are found within Cidaroidea, Aulodonta, and among

outgroup nodes. For each, the range shown spans the interval between the minimum and maximum

ages found among the consensus topologies of all time-calibrated runs.

1103

1104 **Figure S3:** Distribution of posterior probabilities for node ages that show an average difference larger

1105 than 20 Ma depending on the choice of clock prior.

1106

Figure S4: Distribution of posterior probabilities for node ages that show a maximum difference larger

than 20 Ma depending on the choice of clock prior. The largest differences can be seen in the relatively

younger ages of Ambulacraria and Echinodermata when using clock-like genes, and in the relatively

older ages for some nodes within cidaroids and starfish when using loci with high occupancy. Other

sampling criteria largely agree on inferred node ages, as can also be seen in Fig. 3C as short distances

between their centroids in chronospace.

1113

1114 **Figure S5:** Distribution of posterior probabilities for node ages that are the most affected by the choice

1115 of model of molecular evolution. No node showed average differences larger than 20 Ma, so those with

1116 the largest effect are shown.

1117

1118   **Figure S6:** Median ages for selected clades across the consensus trees of the 40 time-calibrated

1119   experiments performed (20 analyses x 2 runs each).

1120

Figure S7: Prior distributions of all constrained nodes. Five hundred replicates were sampled from the

joint prior, showing appropriately broad distributions of node ages. Blue lines show minima and yellow

ones maxima (when enforced); dotted lines show the age of the Permian-Triassic (251.9 Ma) and

Cretaceous-Paleogene (66 Ma) mass extinction events. Nodes whose ages are of special interest

(Echinoidea, Scutelloida, and Clypeasteroida) are shown in pink, revealing large prior probabilities of the

divergence occurring at either side of mass extinction events.

1127

1128 **Figure S8:** Number of lineages inferred to have crossed the Permian-Triassic (P-T) boundary. The

1129 probabilities of each scenario are estimated from the inferred divergence times of the 10,000

1130 chronograms sampled across all of the analyses performed (250 for the two runs of each combination of

1131 sampled loci, model of molecular evolution, and clock prior). The probability of three or more crown

1132 group lineages surviving the Permian-Triassic extinction is 58.47%.

1133

Figure S9: Ordering of loci enforced using *genesortR* [37] and its relationship to the seven gene

properties employed. High ranking loci (i.e., the most phylogenetically useful) show low root-to-tip

variances (or high clock-likeness), low saturation and compositional heterogeneity, as well as high

average bootstrap and Robinson-Foulds similarity to a target topology (in this case, with the contentious

relationship among major lineages of Echinacea collapsed).

1139

Figure S10: Relationship between the root-to-tip variance (a proxy for the clock-likeness of loci) and the

rate of evolution (estimated as the total tree length divided by the number of terminals). The most

clock-like loci (shown in red), which are often favored for the inference of divergence times (e.g., [62,

72]), are among the most highly conserved and can provide little information for constraining node ages

[37]. Highly clock-like genes with a higher information content were used instead by choosing the loci

with the lowest root-to-tip variance from among those that were within 1 standard deviation from the

mean evolutionary rate (shown in blue).

1147

Figure S11: Trace plots of the log-likelihood of different time-calibration runs. All runs show evidence of

reaching convergence and stationarity before our imposed burn-in fraction of 10,000 generations. For

simplicity, only runs under the CAT+GTR+G model are plotted. Those run under GTR+G converged much

faster.

Table S1: Transcriptomic/genomic datasets added in this study relative to the taxon sampling of

Mongiardino Koch et al. [4] and Mongiardino Koch & Thompson [3]. Taxa with citations were taken from

the literature, and details can be found in the corresponding studies and associated NCBI BioProjects.

Taxa are shown in alphabetical order; those identified with "OG" are outgroup taxa (i.e., non-echinoids).

Voucher specimens are deposited at the Benthic Invertebrate Collection, Scripps Institution of

Oceanography (SIO-BIC), and the Echinoderm Collection, Muséum National d'Histoire Naturelle (MNHN-

IE). If multiple accession numbers are shown for a given taxon, these datasets were merged before

assembly. Similar details for all other specimens can be found in [3, 4].

Taxon	Citation	Tissue type	Collector	Locality (depth)	Voucher	GenBank accession number
Amphiura filiformis (OG)	[137]	-	-	-	-	SRX2255774
Apatopygus recens	This study	Mixed	Owen Anderson	Foveaux Strait, South Island, New Zealand	SIO-BIC E7142	
Aspidodiadema hawaiiense	This study	Mixed	Greg Rouse, Avery Hatch	Seamount 4, Costa Rica, Pacific Ocean (1,003 m)	SIO-BIC E7363	
Asterias rubens (OG)	[138]	-	-	-	-	SRX445860
Astrophyton muricatum (OG)	[139, 140]	-	-	-	-	SRX1391908
Bathysalenia phoinissa	This study	Tube feet, spine muscles	BIOMAGLO Cruise Team	Mozambique Channel, Mayotte (295-336 m)	MNHN-IE-2016-23	

Species	Source	Tissue	Collector	Location	Voucher	Accession
Clypeaster japonicus	This study	Eggs	Frances Armstrong	Misaki Marine Biological Station, Kanagawa, Japan	-	
Echinothrix calamaris	This study	Male gonad	-	Phillipines	-	
Encope emarginata	This study	Eggs	Gulf Specimen Marine Lab	Apalachee Bay, Florida, USA	-	
Eucidaris metularia	This study	Mixed	Greg Rouse	Al-Fahal Reef, Red Sea, Makkah, Saudi Arabia	SIO-BIC 2017-008	
Fellaster zelandiae	This study	Spines, tube feet, gut, joining tissue of Aristotle's lantern	Wilma Blom	Western end of Cornwallis Beach, Auckland, New Zealand	SIO-BIC E7920	
Histocidaris variabilis	This study	Gonad	Greg Rouse, Avery Hatch	Las Gemelas Seamount, near Isla del Coco, Costa Rica, Pacific Ocean (571 m)	SIO-BIC E7350	
Lamberticrinus messingi (OG)	[140, 141]	-	-	-	-	SRX1397823
Leodia sexiesperforata	This study	Eggs	Frances Armstrong	Bocas del Toro, Panama	-	
Metacrinus rotundus (OG)	[142]	-	-	-	-	DRX021520

Micropyga tuberculata	This study	Tube feet, spine muscles	BIOMAGLO Cruise Team	Mozambique Channel, Îles Glorieuses (385-410 m)	MNHN-IE-2016-39
Ophiocoma echinata (OG)	[138]	-	-	-	-	SRX445856
Peronella japonica	This study	Eggs	Frances Armstrong	Kanazawa, Japan	-
Rhyncholampas pacificus	This study	Mixed	Carlos A. Conejeros-Vargas	Panteón Beach, Puerto Ángel Bay, Oaxaca, Mexico	SIO-BIC E7918
Scaphechinus mirabilis	[143]	-	-	-	-	DRX187887 DRX187888
Sinaechinocyamus mai	This study	Gregory's diverticulum	Jih-Pai Lin	Miaoli County, Taiwan	SIO-BIC E7919
Sterechinus neumayeri	[144]	-	-	-	-	ERX3498697 ERX3498698 ERX3498699 ERX3498700 ERX3498701
Stereocidaris nascaensis	This study	Muscle surrounding Aristotle's lantern	Charlotte Seid	Off San Ambrosio, Desventuradas Islands, Chile (215 m)	SIO-BIC E7154

Tetrapygus niger	This study	Female gonad, tube feet	Felipe Aguilera	Dichato Bay, Chile	-
Tripneustes gratilla	This study	Pedicellaria	-	Phillipines	-
Tromikosoma hispidum	This study	Tube feet	Lisa Levin, Todd Litke	Quepos Plateau, Costa Rica, Pacific Ocean (2067 m)	SIO-BIC E7251

1160

Table S2: Details of molecular datasets and supermatrix. Statistics for raw reads and assemblies are

shown for datasets incorporated in this study relative to the sampling of [3, 4] (where similar statistics

can be found for the other datasets). Taxa are shown in alphabetical order; those identified with "OG"

are outgroup taxa (i.e., non-echinoids). Novel datasets correspond to Illumina pair-end transcriptomes,

except for two draft genomes (*Bathysalenia phoinissa* and *Micropyga tuberculata*). Mean insert size is

expressed in number of base pairs. For transcriptomes, read pairs (initial) shows numbers input into

Agalma [84], (i.e., after processing with Trimmomatic [83] or UrQt [86]), while read pairs (retained)

shows those that passed the Agalma curation checks (including ribosomal, adapter, quality, and

compositional filters), and represent the final number of read pairs used for assembly. For genomes, see

information in the bioinformatic details above. Assemblies were further sanitized with either

alien_index [92] alone or in combination with CroCo [91] (see bioinformatic details), and the number of

assembled transcripts/contigs shows the size of datasets after these curation steps. The number of loci

shows the occupancy of terminals in the supermatrix output by Agalma (1,346 loci at 70% occupancy),

after which loci were further removed by TreeShrink [97], resulting in the final occupancy scores.

Species	Mean insert size	Read pairs (initial)	Read pairs (retained)	Assembled transcripts/contigs	Number of loci	Removed by TreeShrink	Final occupancy
Abatus agassizii	-	-	-	-	522	5	38.4
Abatus cordatus	-	-	-	-	497	2	36.8
Acanthaster planci (OG)	-	-	-	-	864	8	63.6
Amphiura filiformis (OG)	180.1	61,558,173	52,206,727	416,946	761	8	55.9
Apatopygus recens	415.3	74,315,687	56,898,850	274,380	1152	5	85.2
Apostichopus japonicus (OG)	-	-	-	-	849	9	62.4
Araeosoma leptaleum	-	-	-	-	1184	2	87.8
Arbacia lixula	-	-	-	-	1234	1	91.6
Aspidodiadema hawaiiense	228.7	109,716,219	104,518,714	311,032	1060	11	77.9
Asterias rubens (OG)	230.4	31,890,613	25,495,009	103,090	805	9	59.1
Asthenosoma varium	-	-	-	-	1254	7	92.6

Astrophyton muricatum (OG)	195.9	25,191,954	22,281,829	149,146	478	5	35.1
Bathysalenia phoinissa	-	-	-	154,120	605	6	44.5
Brissus obesus	-	-	-	-	773	0	57.4
Caenopedina hawaiiensis	-	-	-	-	1094	2	81.1
Clypeaster japonicus	198.6	10,505,520	9,298,316	74,743	829	1	61.5
Clypeaster rosaceus	-	-	-	-	1242	2	92.1
Clypeaster subdepressus	-	-	-	-	1233	3	91.4
Colobocentrotus atratus	-	-	-	-	1158	0	86.0
Conolampas sigsbei	-	-	-	-	1001	5	74.0
Cystechinus c.f. *giganteus*	-	-	-	-	661	0	49.1
Dendraster excentricus	-	-	-	-	337	1	25.0
Diadema setosum	-	-	-	-	305	1	22.6
Echinarachnius parma	-	-	-	-	1187	3	88.0
Echinocardium cordatum	-	-	-	-	1012	2	75.0
Echinocardium mediterraneum	-	-	-	-	1185	1	88.0
Echinocyamus crispus	-	-	-	-	709	6	52.2
Echinometra mathaei	-	-	-	-	1055	0	78.4
Echinothrix calamaris	203.9	68,871,087	60,443,904	251,788	1252	4	92.7
Encope emarginata	206.2	12,015,888	10,461,870	66,241	1076	1	79.9
Eucidaris metularia	321.6	38,802,159	29,047,401	83,318	412	2	30.5
Eucidaris tribuloides	-	-	-	-	414	0	30.8
Evechinus chloroticus	-	-	-	-	1196	1	88.8
Fellaster zelandiae	313.9	62,619,791	51,742,748	168,598	1269	1	94.2
Heliocidaris erythrogramma	-	-	-	-	931	0	69.2
Histocidaris variabilis	329.2	84,103,672	73,884,180	144,044	1189	5	88.0
Holothuria forskali (OG)	-	-	-	-	743	8	54.6
Lamberticrinus messingi (OG)	195.0	22,989,820	20,234,597	59,049	351	3	25.9
Leodia sexiesperforata	203.8	16,211,182	14,174,475	68,964	1064	1	79.0
Leptosynapta clarki (OG)	-	-	-	-	403	4	29.6
Lissodiadema lorioli	-	-	-	-	770	0	57.2
Loxechinus albus	-	-	-	-	1265	1	93.9

Species							
Lytechinus variegatus	-	-	-	-	1257	2	93.2
Mellita tenuis	-	-	-	-	1002	1	74.4
Meoma ventricosa	-	-	-	-	1053	0	78.2
Mesocentrotus nudus	-	-	-	-	1247	0	92.6
Metacrinus rotundus (OG)	198.5	23,791,832	20,900,345	83,718	642	7	47.2
Micropyga tuberculata	-	-	-	170,514	415	4	30.5
Ophiocoma echinata (OG)	278.5	28,427,026	24,836,025	130,153	712	7	52.4
Paracentrotus lividus	-	-	-	-	1266	0	94.1
Patiria miniata (OG)	-	-	-	-	740	8	54.4
Peronella japonica	208.3	17,696,287	15,707,931	106,110	1043	6	77.0
Prionocidaris baculosa	-	-	-	-	794	3	58.8
Psammechinus miliaris	-	-	-	-	1173	1	87.1
Rhyncholampas pacificus	346.8	63,116,413	52,160,741	234,910	1070	2	79.3
Saccoglossus kowalevskii (OG)	-	-	-	-	668	6	49.2
Scaphechinus mirabilis	401.7	10,169,195	8,796,067	127,664	974	4	72.1
Sinaechinocyamus mai	297.2	60,208,172	52,265,201	164,646	1233	1	91.5
Sphaerechinus granularis	-	-	-	-	1214	1	90.1
Sterechinus neumayeri	207.3	26,376,279	21,308,840	122,001	1097	1	81.4
Stereocidaris nascaensis	327.0	69,962,495	60,316,050	121,508	1200	6	88.7
Stomopneustes variolaris	-	-	-	-	908	0	67.5
Strongylocentrotus purpuratus	-	-	-	-	1266	5	93.7
Tetrapygus niger	202.5	63,040,541	52,761,671	163,084	1287	2	95.5
Tripneustes gratilla	154.6	62,962,239	57,015,692	130,376	1309	1	97.2
Tromikosoma hispidum	300.4	57,208,357	47,909,038	234,502	1238	7	91.5

1175

Supplementary References (references 1-112 in main text)

1176

113.   Faircloth BC, Glenn TC (2012) **Not all sequence tags are created equal: designing and validating**

sequence identification tags robust to indels *PLoS One* 7:e42543.

114.   BIOMAGLO cruise, RV Antea (2017). doi: 10.17600/17004000.

115.   Bronstein O, Kroh A, Miskelly AD, Smith SD, Dworjanyn SA, Mos B, et al (2019) **Implications of**

range overlap in the commercially important pan-tropical sea urchin genus *Tripneustes* **(Echinoidea:**

Toxopneustidae) *Marine Biology* 166:34.

116.   Seppey M, Manni M, Zdobnov EM (2019) **BUSCO: assessing genome assembly and annotation**

completeness *Methods in Molecular Biology* 1962:227-245.

117.   Kriventseva EV, Tegenfeldt F, Petty TJ, Waterhouse RM, Simao FA, Pozdnyakov IA, et al (2015)

OrthoDB v8: update of the hierarchical catalog of orthologs and the underlying free software *Nucleic*

acids research 43:250-256.

118.   Benton MJ, Donoghue PC, Asher RJ, Friedman M, Near TJ, Vinther J (2015) **Constraints on the**

timescale of animal evolutionary history *Palaeontologica Electronica* 18:1-106.

119.   Erkenbrack EM, Thompson JR (2019) **Cell type phylogenetics informs the evolutionary origin of**

echinoderm larval skeletogenic cell identity *Communications Biology* 2:160.

120.   Cooper R, Sadler P, Hammer O, Gradstein F (2012) **The Ordovician Period**. In: Gradstein FM, Ogg

JG, Schmitz M, Ogg G, editors. *The geologic time scale 2012*. Elsevier. pp. 489-523.

121.   Hunter AW, Ortega-Hernández J (2021) **A new somasteroid from the Fezouata Lagerstätte in**

Morocco and the Early Ordovician origin of Asterozoa *Biology Letters* 17:20200809.

122.   Villier L, Brayard A, Bylund KG, Jenks JF, Escarguel G, Olivier N, et al (2018) *Superstesaster*

promissor **gen. et sp. nov., a new starfish (Echinodermata, Asteroidea) from the Early Triassic of Utah,**

USA, filling a major gap in the phylogeny of asteroids *Journal of Systematic Palaeontology* 16:395-415.

123.   Saucède T, Vennin E, Fara E, Olivier N, Team TPB (2019) **A new holocrinid (Articulata) from the**

Paris Biota (Bear Lake County, Idaho, USA) highlights the high diversity of Early Triassic crinoids

Geobios 54:45-53.

124.   Thuy B, Escarguel G (2019) **A new brittle star (Ophiuroidea: Ophiodermatina) from the Early**

Triassic Paris Biota (Bear Lake County, Idaho, USA) *Geobios* 54:55-61.

125.   Kroh A (2011) **Echinoids from the Triassic of St. Cassian-A review** *Geo.Alp* 8:136-140.

126.   Li Z, Chen Z-Q, Zhang F, Ogg JG, Zhao L (2020) **Global carbon cycle perturbations triggered by**

volatile volcanism and ecosystem responses during the Carnian Pluvial Episode (late Triassic) *Earth-*

Science Reviews 211:103404.

127.   Smith AB (2011) ***Gymnodiadema* and the Jurassic roots of the Arbacioida (stirodont echinoids)**

Swiss Journal of Palaeontology 130(1):155-171.

128.   François É, Marchand D, David B (2003) **Fluctuations morphologiques et hétérochronies chez**

Toxaster (échinides, Crétacé inférieur) *Comptes Rendus Palevol* 2:597-605.

129.   Cooke CW (1955) **Some Cretaceous echinoids from the Americas** *US Geological Survey*

Professional Papers 264:87-112.

130.   Roman J (1989) **L'ancêtre éocène des Scutellidae (Echinoidea; Clypeasteroida)**. In: De Ridder C,

editor. *Echinoderm Research. Proceedings of the Second European Echinoderm Conference, Brussels*.

Rotterdam: AA Balkema. pp. 18-21.

131.   Durham JW (1994) **Fossil *Encope* (Echinoidea) from the Pacific coast of southern Mexico**

Revista Mexicana de Ciencias Geológicas 11(1):113-116.

132.   Coppard SE, Lessios HA (2017) **Phylogeography of the sand dollar genus *Encope*: implications**

regarding the Central American Isthmus and rates of molecular evolution *Scientific Reports* 7(1):11520.

133.   Ali MSM (1983) **The paleogeographic distribution of *Clypeaster* (Echinoidea) during the**

Cenozoic Era *Neues Jahrb für Geologie und Paläontologie Monatshefte* 8:449-464.

134.   Via L, Padreny J (1970) **Dos nuevas especies de *Clypeaster* del Eoceno de Cataluña**.

Publicaciones del Instuto de Investigaciones Geológicas de la Diputación Provincial 24:89-118.

135.   Wade BS, Pearson PN, Berggren WA, Pälike H (2011) **Review and revision of Cenozoic tropical**

planktonic foraminiferal biostratigraphy and calibration to the geomagnetic polarity and astronomical

time scale *Earth-Science Reviews* 104:111-142.

136.   Gold DP, Fenton JP, Casas-Gallego M, Novak V, Pérez-Rodríguez I, Cetean C, et al (2018) **The**

biostratigraphic record of Cretaceous to Paleogene tectono-eustatic relative sea-level change in

Jamaica *Journal of South American Earth Sciences* 86:140-161.

137.   Dylus DV, Czarkwiani A, Stångberg J, Ortega-Martinez O, Dupont S, Oliveri P (2016) **Large-scale**

gene expression study in the ophiuroid Amphiura filiformis provides insights into evolution of gene

regulatory networks *Evodevo* 7:1-17.

138.   Reich A, Dunn C, Akasaka K, Wessel G (2015) **Phylogenomic analyses of Echinodermata support**

the sister groups of Asterozoa and Echinozoa *PLoS One* 10:e0119627.

139.   Linchangco Jr GV, Foltz DW, Reid R, Williams J, Nodzak C, Kerr AM, et al (2017) **The phylogeny of**

extant starfish (Asteroidea: Echinodermata) including Xyloplax, based on comparative transcriptomics

Molecular Phylogenetics and Evolution 115:161-170.

140.   Janies DA, Witter Z, Linchangco GV, Foltz DW, Miller AK, Kerr AM, et al (2016) **EchinoDB, an**

application for comparative transcriptomics of deeply-sampled clades of echinoderms *BMC*

Bioinformatics 17:17-48.

141.   Clouse RM, Linchangco Jr GV, Kerr AM, Reid RW, Janies DA (2015) **Phylotranscriptomic analysis**

uncovers a wealth of tissue inhibitor of metalloproteinases variants in echinoderms *Royal Society*

Open Science 2:150377.

142. Koga H, Fujitani H, Morino Y, Miyamoto N, Tsuchimoto J, Shibata TF, et al (2016) **Experimental**

approach reveals the role of alx1 in the evolution of the echinoderm larval skeleton *PLoS One*

11:e0149067.

143. Simakov O, Kawashima T, Marlétaz F, Jenkins J, Koyanagi R, Mitros T, et al (2015) **Hemichordate**

genomes and deuterostome origins *Nature* 527:459-465.

144. Collins M, Peck LS, Clark MS (2021) **Large within, and between, species differences in marine**

cellular responses: Unpredictability in a changing environment *Science of The Total Environment*

148594.