Ecosystem services in connected catchment to coast ecosystems: monitoring

to detect emerging trends

Christopher J. Brown* [1], Chantal Saint Ange [2], Rod M. Connolly [1], Syezlin Hasan [2], Sue Jackson

[2], Joseph M. McMahon [2], James C. R. Smart [2]

1. Coastal and Marine Research Centre, Australian Rivers Institute, School of Environment and

Science, Griffith University, Gold Coast, QLD 4222, Australia

2. Australian Rivers Institute, School of Environment and Science, Griffith University, Nathan,

QLD, Australia

*Corresponding author: chris.brown@griffith.edu.au 170 Kessels Road Nathan, Queensland,

Australia, 4111.

Abstract

There is an increasing need for long-term monitoring of ecosystems and their services to inform on-

ground management and policy direction. The supply of many ecosystem services relies on

connections that span multiple ecosystems. Monitoring the underlying condition of interconnected

ecosystems, using established indicators, is therefore required to track effectiveness of past

interventions and, ideally, identify impending change. Here we conduct performance testing of

ecological indicators for a catchment-to-coast system with the aim of identifying the time-scales

over which they respond to change. We chose a case-study of a coastal fishery in Northern Australia

that exhibits strong catchment-to-coast connectivity, has long-term available data and is under

threat from water resource development. We developed a novel approach to performance testing.

Our model drew on state-space modelling to capture ecological dynamics, and structural equation

modelling to capture covariation in indictors timeseries. We first quantified covariation among three

established ecological indicators: pasture biomass, vegetation greenness and barramundi catch per

unit effort. Covariation in the indicators was driven by river flow, with higher values of all indicators

occurring in years with greater river flow. We then defined reference bounds for each indicator that

accounted for natural variation in river flow. We predicted the emergence times for each indicator,

as the time taken for each indicator to emerge from the background of natural variation. Emergence

times quantified at 80% and higher confidence levels were >10 years in all cases. Past trends and

current status of ecosystem service flows are often used by decision makers to directly inform near-

term actions, particularly provisioning services (such as barramundi catch) due to their important

contribution to regional economies. We found that the ecological indicators should be used to

assess historical performance over decadal timespans, but not as short-term indicators of recent

change. More generally, we offer an approach to performance testing of indicators. This approach

could be useful for quantifying time-scales of ecosystem response in other systems where cross-

ecosystem connections are important.

Keywords: *Lates calcarifer*, power analysis, state-space model, catchment to coast, estuary,

emergence time, integrated coastal management.

Introduction

The supply of ecosystem services can depend on ecological processes that operate across multiple

ecosystems (Barbier et al. 2011; Lowe et al. 2022). Interconnections among ecosystems can occur

through the movement of plant propagules, migration of animals and the transport of physical

matter in air and water. In particular, the flow of water from catchments to coast creates many

challenges for the management of human activities that may impact upon the supply of ecosystem

services. For example, the capture of water in dams for agriculture, power or municipal uses

upstream interferes with the supply of ecosystem services all the way to the ocean, potentially

affecting ecosystem inputs to agriculture on floodplains, the productivity of estuarine and coastal

marine fisheries (Gillanders et al. 2011) and the capture of carbon in estuarine habitats (Macreadie

et al. 2017). Therefore, monitoring of multiple ecological indicators that together span

interconnected systems is important to characterize the processes that support ecosystem services

in catchments and inform the management of human activities. This monitoring of ecological

processes plays multiple roles in the decision-making. First, it informs decision makers about trends

in downstream ecosystem processes so they can respond proactively to address causal changes in

hydrology upstream. Second, it helps to detect change in ecosystems and attribution to

management actions. Finally, monitoring can hold decision makers accountable for actions that

affect the integrity of ecosystems downstream (Stevenson et al. 2021).

A challenge for monitoring that spans a whole catchment is to attribute trends in ecological

processes to changes in water flow, because the causes of changes in water flow are separated in

time and space from their impacts (Alvarez-Romero et al. 2011; Brown et al. 2019). Interannual

variation in flow linked to long-term climate cycles (e.g. El Nino southern oscillation) is a common

property of many systems, and this variation also contributes to masking of long-term trends caused

by climate change or human interventions. Monitoring design, such as 'before/after, control/impact'

approaches can be used to resolve the attribution problem (Underwood 1992). However, in whole of

catchment monitoring programs, commonly no controls exist; monitoring programs are not

designed with a specific intervention such as water extraction for irrigation as the focus. In such

cases we need to know how long underlying trends in hydrology/flow will take to manifest as trends

in the ecological indicators.

The testing of indicator performance, where the ability to detect trends in indicators is explored

through simulation models, is a popular method in fisheries science for evaluating indicators (Fulton

et al. 2005; Blanchard et al. 2014; Nickols et al. 2019), but it is only just beginning to be applied to

ecosystem indictors (Collen & Nicholson 2014; Watermeyer et al. 2021). Performance testing

involves constructing a system model that represents the fundamental system dynamics, and then

running simulations that represent the stages of underlying trends, data collection and

interpretation of data by decision-makers (Fulton et al. 2005). Simulations are run multiple times so

that we can define error bounds on the probability of a trend of a given magnitude being detected

with a given level of confidence. This testing can be evaluated through univariate indicators, such as

binary indicators of whether a threshold value is reached, or via power analyses that consider the

sensitivity and specificity of indicators for different decision thresholds (i.e. risk of missed detections

versus false alarms - Mapstone 1995). The development of performance testing methods in fisheries

science and biodiversity monitoring has run parallel to the development of the emergence time

concept in climate studies (Hawkins & Sutton 2012). Climate models can be used to quantify when

human induced trends in climate caused by greenhouse gas emissions emerge from background

variation in climate. Thus, models of trends underpin the attribution of climate trends to human

causes and provide an easily communicable statistic about lags between human actions and

detection of changes in the system under study. The emergence time concept could also be applied

to the testing of indicators for interconnected catchment ecosystems, where high interannual

variation in river flows may mask long-term trends in multiple downstream ecological processes.

However, development of such performance testing is often held back by the requirement to

develop complex system models. Developing system models for catchment to coast systems is a

particularly complex task because the modelling needs to span multiple disciplines (Brown et al.

2019). As such, catchment to coast systems models are rare. Simpler approaches to performance

testing would facilitate more widespread testing of indicators, and therefore inform how managers

can use indicators.

Here we develop an approach to quantifying emergence time of multivariate catchment processes.

We use a case-study of a riverine-estuarine-coastal marine system in the Gulf of Carpentaria in

tropical northern Australia. The Mitchell catchment is earmarked for potential water resource

development, which may provide agricultural benefits, but will also disrupt freshwater flows (CSIRO

2018) and impact the downstream ecosystems and fisheries that depend on them (Turschwell et al.

2019; Broadley et al. 2020). Similarly, climate change is causing long-term shifts in weather patterns

and flows in the region (Karim et al. 2015). We first selected commonly used monitoring indicators

for the interconnected ecosystems of the Mitchell catchment. We then sought to fit a process model

that quantified the relationship among indicators and that allowed the effect of uncertainty within

those indicators to be assessed (Aim 1). From this model we inferred natural bounds of variation for

each indicator (Aim 2). We finally quantified the sensitivity of indicators to long-term trends in water

flow via their emergence time from background variation (Aim 3). The emergence time for each

indicator represents when the impacts of change will become apparent to human observers

(Hawkins & Sutton 2012). Overall, our approach provides a method for performance testing of

multiple indicators. Our approach could be used to support development of indicator sets for

attribution of trends in human impacts, ecological condition accounts (Czúcz et al. 2021) and

monitoring of human impacts.

Methods

Indicators for the Mitchell catchment

The Mitchell catchment covers an area of over 72,000 km^2 and discharges into the Gulf of

Carpentaria. This catchment is subject to highly seasonal flow regimes. During the dry season many

of the ephemeral rivers become disconnected water holes and in the wet season a large area

becomes inundated. There is high inter-annual variability in rainfall, which drives inter-annual

variation in flow and floodplain productivity (Ndehedehe et al. 2021). We therefore selected

variables and indicators for the catchment with the aim of capturing variation in ecological states

related to river flow.

We focused on indicators of capacity to supply ecosystem services that included (i) fish biomass that

supports barramundi (*Lates calcarifer*) catch in a commercial fishery and resource harvesting of

barramundi by Indigenous Australians, and (ii) pasture biomass that supports commercial livestock

production. In particular, the catadromous barramundi life-cycle exemplifies cross-system

dependence (Barramundi migrate from freshwater to estuaries to spawn, Milton and Chenery 2005).

Variability in barramundi biomass may be correlated to variability in pasture growth on flood plains

which supports livestock production, because both are dependent on regional cycles in river flow.

Time-series data (Table S1) were collected for the ecosystem service capacity indicators, and

additional indicators where long-term data were available. Stream flow, a hydrological indicator, was

included and sourced from the Northern Australia Water Resource Assessment (NAWRA).

Yearly data on barramundi (*Lates calcarifer*) catch and fishing effort were used to derive an indicator

of the barramundi population state. Data were sourced from the Queensland Fisheries 'QFish' data

cube (State of Queensland Department of Agriculture Fisheries and Forestry 2018) for the Mitchell

region. Barramundi are an important fishery species in northern Australia, and they are caught in

commercial, Indigenous, recreational and charter fisheries. They also play an important cultural role

within the ecosystem (Jackson et al. 2012). Barramundi spawn in estuaries, with juveniles then

moving upstream to freshwater habitats, returning to saltwater as adults (Robins et al. 2005).

138 Seasonal floodplains provide connection to off-stream freshwater habitats as well as highly

139 productive, short-term habitats. Individuals residing in upstream habitats return to the estuary and

140 saltwater to spawn as adults. Barramundi are impacted by flow regimes through changes in

141 connectivity throughout the catchment which benefit growth and survival (Robins et al. 2005;

142 Balston 2009; Roberts et al. 2019; Leahy & Robins 2021). Barramundi opportunistically exploit

143 multiple aquatic habitats at different stages of their life cycle. Therefore, if connectivity between

144 these different ecosystem assets is diminished, this can impact their harvestable biomass in

145 following seasons. Barramundi catch per unit effort (CPUE) is a suitable indicator of the interlinked

146 aquatic ecosystems, because barramundi are a near-apex predator in the system so their production

147 aggregates multiple trophic levels, and they depend on persistent freshwater in lagoons and

148 waterholes to survive dry periods (Robins et al. 2005).

149 Normalized difference vegetation index (NDVI) is an indicator of vegetation greenness (Pettorelli et

150 al. 2005) and was obtained from the Bureau of Meteorology (Table S1). Pasture biomass was

151 included as an indicator of supply of forage for grazing. Pasture biomass was obtained as the

152 predicted output from Queensland Government's AussieGRASS model. AussieGRASS also uses

153 stream flow as input data; however, since other variables (i.e. rainfall, soil typ, temperature and local

154 grazing pressure) are also inputs to AussieGRASS (State of Queensland Department of Environment

155 and Science, 2021), pasture biomass and stream flow are not highly correlated.

Aim 1: Quantify interrelations among condition indicators

We combined two common modelling approaches to develop a model of indicator interrelationships

(fig. 1). The first was a state-space model of barramundi catch per unit effort (CPUE) dynamics

(Millar & Meyer 2000). State space models are widely applied for modelling fisheries dynamics,

informing on reference points for fisheries management and for estimating environmental effects on

fishery dynamics (Aeberhard et al. 2018), but have rarely been applied to model whole ecosystem

change (Auger-Méthé et al. 2021). The state-space model was used to fit catches (a measure of

ecosystem service flow), then estimate latent states for stock biomass (a measure of ecosystem

capacity to deliver services) and environmental process variation (fig. 1).

The second modelling approach was a structural equation model that related the observed indicator

variables to a latent (unobserved) state, which we termed 'ecological condition index' (fig. 1).

Structural equation models are widely used to develop indicators, because they can model the

correlations between observed indicators and a latent state that is hypothesized to be equivalent to

the joint condition of different components of an ecosystem (Brown & Hamilton 2018). We linked

the two modelling approaches by assuming that the observed indicators of NDVI, pasture biomass,

and CPUE were all dependent on an unobserved ecosystem condition index. Ecosystem condition, in

turn, was assumed to be driven by freshwater flow (fig 1).

The state space model had the form:

175
$$B_t = \left(B_{t-1} + rB_{t-1}\left(1 - \frac{B_{t-1}}{K}\right) - C_t \right) u_t \qquad \text{(eqn 1)}$$

176
$$ln\,(CPUE_t) \sim normal(ln\,(qB_t), \sigma^{CPUE}) \qquad \text{(eqn 2)}$$

where B_t was the latent estimate of biomass, r was the intrinsic growth parameter, K was the

carrying capacity, C_t was catches (assumed fixed), q is the catchability coefficient, u_t were process

errors and σ^{CPUE} was the standard deviation for CPUE errors (we use superscripts to demarcate

parameters relating to different variables, subscripts to demarcate indices). B_1 was assumed a fixed

fraction of 20% of the estimate of K (Millar & Meyer 2000; Streipert et al. 2019). We tested the

effects of the assumptions that B_1 was 20% of K and that q was constant over time in additional

sensitivity analyses (Supplementary material).

The state space model was linked to the structural equation model through a dependence of the

process errors on a latent variable we interpreted as the ecosystem condition index:

$$ln\,(u_t) \sim normal(\beta^u v_t, \sigma^u) \qquad \text{(eqn 3)}$$

where β^u was a regression coefficient and v_t was the latent condition index. The process errors

were assumed to be log-normally distributed so they had a multiplicative effect on biomass. This

formulation is consistent with population theory because changes in survival have multiplicative

effects on population size.

The state-space model assumed a one-year lag between flow and changes in barramundi biomass.

Flow likely impacts barramundi catches and biomass at a number of different time-scales, as

described above, but we assumed the single year impact for simplicity. To further verify this

assumption, we confirmed that residuals from the modelled CPUE relationship had no

autocorrelation. Future modelling could potentially separate the different types of flow effects on

barramundi populations. But note this would require additional data types, because biomass,

productivity and catchability parameters are close to being interchangeable in unstructured surplus

production models, so they are difficult to estimate simultaneously.

NDVI and pasture biomass were also related directly to the condition index:

$$y_t \sim normal(a^y + \beta^y v_t, \sigma^y) \qquad \text{(eqn 4)}$$

where y_t refered to either NDVI or pasture biomass and a^y is the intercept.

NDVI and pasture biomass were mean centred and standardized prior to analysis.

Finally, ecosystem condition was estimated as a regression on freshwater flow:

$$v_t \sim normal(\beta^v \ln(F_t), 1) \qquad \text{(eqn 5)}$$

Where F_t was freshwater flow and the standard deviation was arbitrarily set at 1, because v_t is scale

invariant. In our regression we used the log of flow, to reflect hypothesized multiplicative effects of

flow on ecological condition.

The NDVI data started in 1992, so the condition index was estimated only from flow, CPUE and

pasture biomass in 1990 and 1991. The model also interpolated the missing NDVI values. We

presented results as marginal predictions for each data type and the latent states for barramundi

biomass and ecological condition.

We obtained parameter and prediction probabilities within a Bayesian estimation framework. We

set priors based on prior information for each parameter and prior expectations for the variance in

NDVI and CPUE. It was also important to set weakly-informative priors to aid algorithm convergence

and identification of parameters. In particular, the latent state v_t and the β coefficients can 'sign-

switch' (Hui et al. 2015). Prior choices and algorithm settings are explained in the Supplementary

material. The model assumptions were checked by confirming independent normally distributed

residuals, where residuals were defined as the observed values of CPUE, NDVI and pasture biomass

minus their posterior predicted median value. Residual independence was confirmed by plotting the

auto-correlation function with the R function 'acf' (R Development Core Team 2019). Further, we

calculated the cross-correlation of residuals across indicators to confirm the model had captured all

dependencies between indicator values. We further verified the model through a simulation study,

focussing on the model's ability to interpolate missing data (Supplementary material).

We determined the strength of the relationship between each indicator variable, ecological

condition, and flow as the direct effect of condition on each indicator (simply the marginal posterior

distributions for β^y) and the total effect of flow (logged) on the condition index $= \beta^v . \beta^y$. For

barramundi we calculated the direct and total effect sizes on surplus production, where surplus

production was defined as the sum of change in biomass and catch (i.e. $\mathrm{SP}_t = B_t - B_{t-1} + C_t$). We

then plotted the total effect of a doubling of flow on each indicator and the direct effect of a one

S.D. change in ecological condition on each indicator (see Supplemental methods). We also

calculated one-tailed probabilities that the total effect was greater than zero. We note that given the

prior choices (Table S2) the prior probability that $\beta^v . \beta^y >0$ was 0.5.

Aim 2: Bounds of natural variation

We used the fitted model to define the bounds of natural variation for each indicator. The bounds

were defined to represent the typical ranges of indicator values given historical variation in flow. The

historical condition of flow was characterized by fitting an autoregressive moving average model to

the historical flow series (1990-2015), then simulating 1000 random flow series from the fitted

model (Supplementary material). To define bounds for the indicators we simulated their values for

each of the 1000 flow series. For NDVI and pasture biomass, the bounds were then defined as the

marginal probability quantiles integrating over measurement noise (also called predictive intervals)

and the flow simulations. The bounds for barramundi CPUE also depended on biomass, so for

simulations of barramundi we assumed a biomass fixed at the biomass that gives maximum

sustainable yield ($B_1 = 50\%$ of B_0). This is a common reference point used in fishery management

(Mace 1994). For the simulations of bounds, we also assumed that biomass was initially at the

biomass that gives maximum sustainable yield, doing so avoided transient dynamics affecting the

simulations.

Aim 3: Emergence time for indicators

We then estimated each indicator's emergence time, given an assumed long-term trend in flow.

Climate change is causing long-term changes in flows in the region (Karim et al. 2015), which may

affect interconnected catchment ecosystem services, including barramundi catches and pasture

growth. Thus, our scenarios represent the ability to detect this long-term change under an assumed

trend magnitude. We simulated flow series and added a linear decline in flow, as an example of a

situation where environmental change or water resource development could put increasing

constraints on ecosystem services. We simulated 1000 by 30-year time-series for each indicator and

each trend in flow. We then defined emergence times as the time taken for the mean indicator value

to cross the lower 5%, 20% and 40% probability quantiles. The different quantiles represent different

degrees of confidence for emergence, the lower the quantile the greater the confidence the trend

has emerged from background variation.

Results

Aim 1: Interrelationships among indicators

Verification of the model fits indicated that there was no autocorrelation in the residuals for the

fitted values for any of the indicators (Fig S1). Overall, the R^2 was highest for log CPUE (0.91),

moderate for NDVI (0.43) and lowest for pasture biomass (0.09) (Fig S2). Sensitivity analyses in which

we varied the assumptions regarding constant catchability, the initial biomass ratio and higher

fishing effort indicated the main results were robust to alternative assumptions (Figs S2, S3 and S4).

Consequently, below we present only the main results assuming stable catchability and an initial

biomass ratio of 20%.

Model fits indicated an overall increase in CPUE from 1990 to 2012, and then a slight decrease from

2012-2015. NDVI and pasture biomass had no long-term trends, but varied year-to-year (Fig. 2).

NDVI and barramundi surplus production were strongly related to river flow (Fig. 3A, pr($\beta^v . \beta^y$.)>0 =

0.99 and 0.96 respectively), whereas pasture biomass had a somewhat weaker relationship with

river flow (figure 3A, pr($\beta^v . \beta^y$.)>0 = 0.93). Therefore, the ecological condition variable primarily

captured the correlation between flow, NDVI and barramundi biomass change (Fig 2E, Fig 3B).

Ecological condition showed peaks in 1991, 1998-99 and around 2010 that were driven by increased

flow in those years. But overall ecological condition did not vary strongly from the long-term average

(95% CIs generally overlap zero). Barramundi biomass was predicted to have increased from 0.2 to

0.5 as a fraction of B_0 over 1990-2012, and to have declined to ~0.3 by 2015. The decline in 2012-

2015 corresponded to a decline in ecological condition and a decline in flow.

A simulation study where we varied the number of years of missing NDVI data suggested the

model's predictions for the variables and ecological condition were not biased by missing data, but

the predictions for NDVI were less accurate with a greater amount of missing data (Fig. S5).

Coverage by the 95% C.I.s was always >0.95 (Table S3). The parameter estimate for the effect of

ecological condition on NDVI was most affected by missing data, and the parameter estimates for

pasture biomass and barramundi biomass were least affected.

Aim 2: Bounds of natural variation

The bounds for each indicator had broad intervals (Fig. 4), for instance, for CPUE the 80% probability

interval ranged from 39-65 tonnes per fishing year, compared to a historical range of 19-50 tonnes

per fishing year. This broad variation in the probability bounds was caused by the high variation in

year-to-year flow.

Aim 3: Emergence times

Emergence times were greatest for pasture biomass and lowest for CPUE (Figs 4 & 5). For instance, if

flow decreased by multiples of 0.9 (10% per year), the time required for mean CPUE to fall below the

5% quantile (90% probability intervals) of historical variation was 16 years, but >30 years for pasture

biomass and NDVI. Emergence times were lower for higher quantiles, reflecting the reduced level of

confidence that the trend had emerged from background variation. The differences in emergence

times among indicators became smaller as the quantiles were increased; for example at a quantile of

40% the indicators had similar emergence times for all trends in flow.

Code and data to repeat the analyses is available at: https://github.com/cbrown5/ecological-

condition-latent-model.

Discussion

We identified relationships among all indicators and flow, and particularly for barramundi catch per

unit effort and NDVI. There is high interannual variation in the Mitchell catchment, driven by

interannual variation in rainfall. Wet years drive increased vegetation growth and barramundi

production, which was reflected in correlations among river flow, NDVI and barramundi catch. Thus,

ecological condition, defined here as covariation in the indicators, largely reflected year-to-year

variation in flow. This variation underscores the importance of quantifying interconnections

between assets via river flow and among ecosystem services. The implication is that change in one

variable (flow) can cause changes in multiple processes, and consequently there would be correlated

changes in the annual capacity of the system to deliver different ecosystem services.

By defining the ecological condition index as a latent variable we were able to partition out

environmental effects shared by all indicators (captured in the ecological condition index) from

those that were unique to specific indicators (variation in indicators that is not explained by the

condition index). The ability to identify shared trends may be desirable in an indicator, because it

facilitates the attribution of changes that affect only specific indicators versus those that impact the

overall condition of the ecosystem (Sutherland et al. 2016; Kupschus et al. 2016). Over time there

was an increasing trend in barramundi CPUE from the 1990s to 2015. This was driven by high flows

2005-2015 and management changes in the fishery in the late 1990s and early 2000s, which reduced

fishing effort and put increased restrictions on fishing gear (Streipert et al. 2019). NDVI had a strong

correlation with flow and CPUE, but in contrast to CPUE, NDVI did not show as strong a long-term

trend, because it was not affected by fishery management changes. Pasture biomass had the

weakest correlation with the other indicators, perhaps because it was more strongly driven by

rainfall that is local to a region and less by inundation caused by increased river flow.

The typical approach to defining joint indices of multiple ecosystem components has been spatial

and thematic aggregation by summing spatially-specific counts or by averaging and then normalising

spatially-specific indicators (Maes et al. 2020). The aggregation procedure requires the use of some

form(s) of aggregation functions and weights. Our model-based approach to defining a condition

index can also be interpreted as a weighted sum of the indicators, where the weights were

estimated empirically. Each approach has its own advantages and disadvantages. Our approach

requires time-series data, which are not available for many indicators or in some regions (Dvarskas

2018). Expert weights do not require historical data and are simple to compute, but can also be

arbitrary. Arbitrary weights are not repeatable across different groups of experts and may mis-

represent management priorities (Game et al. 2013). Further, our approach partitioned variation in

CPUE caused by flow from changes in management of the fishery. Simply summing weighted CPUE

with other indicators would tend to overinflate the influence of barramundi fishery management on

ecological condition as a whole. Thus, a key advantage of our approach is that it can partition

changes unique to individual indicators, versus changes that are shared across the ecosystem.

Identifying redundancy and complementarities among different indicators is an important step in

ensuring that monitoring programs can focus on a parsimonious set of indicators that is

comprehensive of the ecosystem's dynamics (Czúcz et al. 2021).

A number of modelling toolboxes have been previously developed to quantify connections among

the condition of ecosystems and ecosystem service flows, but none of these directly address the

issue of how effectively one can detect trends in indicators. Examples include the soil and water

assessment tool (Bieger et al. 2017), the Integrated valuation of ecosystem services tradeoff

(INVEST) tool (Guerry et al. 2012) and globally applicable artificial intelligence tools (Martínez-López

et al. 2019). These tools have been applied for measuring connected ecosystem services across

catchments, in particular regulating ecosystem services relating to water quantity, water quality and

sediment erosion (Bieger et al. 2017; Hamel et al. 2017). All share a common approach, based on

parameterising models of soil erosion and run-off and then analysing predicted service flows

spatially in a geographic information system. However, these toolboxes are designed for making

service flow projections, given sufficient information exists to parameterize their models. The utility

of our approach is as a performance evaluation tool, which is an understudied aspect of biodiversity

indicator development (Rowland et al. 2020; Watermeyer et al. 2021). Our approach, which blends

state-space and structural equation modelling, could be extended to model multivariate change in

other ecosystems. This type of model warrants further development for assessing ecosystem level

change, particularly where there is a need to differentiate the different sources of stochasticity in

observed time-series (Auger-Méthé et al. 2021)

The high interannual variation in flow and ecological condition will create challenges for detecting

change in the Mitchell catchment. Our estimates of the interannual variation for each indicator were

broad and thus our model predicted that emergence times across all indicators at typical confidence

bounds (10% or 5%) were >10 years and sometimes >30 years for all but the most extreme trends in

flow. These long emergence times mean the indicators will not be useful for managers aiming to use

the model to pre-emptively respond to change, because the changes will not be detected on a

management relevant timespan (Stevenson et al. 2021). Rather, the indicators will capture long-

term low-bandwidth changes in the ecosystem. For comparison, proposed water developments in

the Mitchell catchment are expected to cause much smaller reductions in flow, in order of 1-5% per

year (Petheram et al. 2018), than most of the trends we explored here. Such water developments

however are predicted to impact estuarine ecosystem services (Broadley et al. 2020; Leahy & Robins

2021). Thus, our findings suggest management planning should not rely on indicators of ecological

condition. Instead, management should use pressure indicators, that may better pre-empt change in

ecological condition (Czúcz et al. 2021). Predictive modelling of the response to ecosystem services

to future changes in land-use and water flow is also needed to inform pre-emptive management in

the Mitchell catchment (McMahon et al. 2022).

There are some important caveats to consider in our analysis. First, there was missing data in some

of the indicator time-series. Missing data in time-series is a common problem in modelling and will

reduce the power to detect long-term change (Dvarskas 2018). The latent variable approach enabled

interpolating missing data, on the strength of correlated variables (Hui et al. 2015). A simulation

study suggested accuracy of predictions for a variable was not sensitive to missing data in other

variables. Thus, we suggest our approach could have value more generally for creating indices of

ecological condition when there are gaps in time-series.

The analysis of barramundi used a simplified stock assessment model. For many fisheries additional

data may be available, such as data on age structure, and this could improve the accuracy of the

modelling. Our simple stock assessment did compare well with a more quantitatively complex stock

assessment, with the stock status estimates being similar (Streipert et al. 2019) – this is the

consequence of both assessment models being driven primarily by trends in CPUE and the

assumption of an depletion of 0.2 B_0 in 1990. More importantly, the availability of barramundi to

estuarine fisheries, as well as productivity, is also likely to be affected by flow (Robins et al. 2005).

Future analyses could incorporate this effect if more data were available to separate catchability

from productivity effects.

An important caveat is that our indicators are an incomplete representation of the processes

supporting capacity to supply ecosystem services in the Mitchell catchment. We focussed on assets

that support commercial fish harvest and commercial livestock grazing, because there were

sufficient time-series data for these indicators to assess emergence times. However, Indigenous uses

of ecosystems are also an important component of the ecosystem services in Northern Australian

catchments (Scheepers & Jackson 2012; Jackson et al. 2012, 2014), which may be particularly

sensitive to development (Stoeckl et al. 2013). Our modelling did not include condition directly

relating to Indigenous services, due to lack of time-series data. It is likely that our conclusions

regarding emergence time would also apply to those assets supporting traditional uses, such as

water holes, given the interconnections in this catchment. Future modelling studies could integrate

Indigenous knowledge to enhance the predictive power of ecosystem service models. For example,

Indigenous knowledge on the spatial distribution of barramundi (Scheepers & Jackson 2012) could

be incorporated into ecosystem service models (e.g. Guerry et al. 2012). This would enhance our

ability to predict change in ecosystem services, as well as quantify their value in a broader range

cultural contexts.

Conclusion

We analyzed change in indicators for interconnected ecosystem assets and found a high level of

variation. This high level of variation means long time-series are required to detect change in the

capacity of the ecosystem to supply ecosystem services. Our quantification of timescales to detect

change provides important context for policy-makers when they are considering indicators

(Stevenson et al. 2021). For instance, we urge caution when using indicators like the ones analysed

here in cost-benefit analysis when evaluating water resource development proposals. Economic

variables may respond rapidly and positively to development (Stevenson et al. 2021), but the

consequent degradation of ecological condition and ecosystem's capacity to supply ecosystem

services may take much longer to be noticed. These temporal differences in response times may not

be adequately captured in cost benefit analysis, which may lead to decisions that are biased towards

development opportunities that bring short-term benefits but have long-term environmental costs.

Literature cited

Aeberhard WH, Mills Flemming J, Nielsen A. 2018. Review of State-Space Models for Fisheries
Science. Annual Review of Statistics and Its Application **5**:215–235. Annual Reviews Inc.
Alvarez-Romero JG, Pressey RL, Ban NC, Vance-Borland K, Willer C, Klein CJ, Gaines SD. 2011.
Integrated land-sea conservation planning: the missing links. Annual Review of Ecology,
Evolution, and Systematics **42**:381–409.

Auger-Méthé M, Newman K, Cole D, Empacher F, Gryba R, King AA, Leos-Barajas V, Mills Flemming J,
Nielsen A, Petris G. 2021. A guide to state–space modeling of ecological time series.
Ecological Monographs. Wiley Online Library.
Balston J. 2009. An analysis of the impacts of long-term climate variability on the commercial
barramundi (Lates calcarifer) fishery of north-east Queensland, Australia. Fisheries Research
99:83–89.
Barbier EB, Hacker SD, Kennedy C, Koch EW, Stier AC, Silliman BR. 2011. The value of estuarine and
coastal ecosystem services. Ecological Monographs **81**:169–193. John Wiley & Sons, Ltd.
Bieger K, Arnold JG, Rathjens H, White MJ, Bosch DD, Allen PM, Volk M, Srinivasan R. 2017.
Introduction to SWAT+, a completely restructured version of the soil and water assessment
tool. JAWRA Journal of the American Water Resources Association **53**:115–130. Wiley Online
Library.
Blanchard JL, Andersen KH, Scott F, Hintzen NT, Piet G, Jennings S. 2014. Evaluating targets and
trade-offs among fisheries and conservation objectives using a multispecies size spectrum
model. Journal of Applied Ecology **51**:612–622. John Wiley & Sons, Ltd.
Broadley A, Stewart-Koster B, Kenyon RA, Burford MA, Brown CJ. 2020. Impact of water
development on river flows and the catch of a commercial marine fishery. Ecosphere
11:e03194. John Wiley & Sons, Ltd.
Brown CJ, Hamilton R. 2018. Estimating the footprint of pollution on coral reefs with models of
species turnover. Conservation Biology.
Brown CJ, Jupiter SD, Albert S, Anthony KR, Hamilton RJ, Fredston-Hermann A, Halpern BS, Lin H-Y,
Maina J, Mangubhai S. 2019. A guide to modelling priorities for managing land-based
impacts on coastal ecosystems. Journal of Applied Ecology **56**:1106–1116.
Collen B, Nicholson E. 2014. Taking the measure of change. Science **346**:166–167. American
Association for the Advancement of Science.
CSIRO. 2018. Water resource assessment for the Mitchell catchment. An overview report to the
Australian Government from the CSIRO Northern Australia Water Resource Assessment.
Pages 24–24. Australia.
Czúcz B, Keith H, Maes J, Driver A, Jackson B, Nicholson E, Kiss M, Obst C. 2021. Selection criteria for
ecosystem condition indicators. Ecological Indicators **133**:108376.
Dvarskas A. 2018. Mapping ecosystem services supply chains for coastal Long Island communities:
Implications for resilience planning. Ecosystem Services **30**:14–26. Elsevier.
Fulton EA, Smith AD, Punt AE. 2005. Which ecological indicators can robustly detect effects of
fishing? ICES Journal of Marine Science: Journal du Conseil **62**:540–551.
Game ET, Kareiva P, Possingham HP. 2013. Six common mistakes in conservation priority setting.
Conservation Biology **27**:480–485. Wiley Online Library.
Gillanders BM, Elsdon TS, Halliday IA, Jenkins GP, Robins JB, Valesini FJ. 2011. Potential effects of
climate change on Australian estuaries and fish utilising estuaries: a review. Marine and
Freshwater Research **62**:1115–1131.
Guerry AD et al. 2012. Modeling benefits from nature: using ecosystem services to inform coastal
and marine spatial planning. International Journal of Biodiversity Science, Ecosystem
Services & Management **8**:107–121. Taylor & Francis.
Hamel P, Falinski K, Sharp R, Auerbach DA, Sánchez-Canales M, Dennedy-Frank PJ. 2017. Sediment
delivery modeling in practice: Comparing the effects of watershed characteristics and data
resolution across hydroclimatic regions. Science of the Total Environment **580**:1381–1388.
Elsevier.
Hawkins E, Sutton R. 2012. Time of emergence of climate signals. Geophysical Research Letters **39**.
John Wiley & Sons, Ltd. Available from https://doi.org/10.1029/2011GL050087 (accessed
May 19, 2021).

Hui FKC, Taskinen S, Pledger S, Foster SD, Warton DI. 2015. Model-based approaches to
unconstrained ordination. Methods in Ecology and Evolution **6**:399–411. British Ecological
Society.
Jackson S, Finn M, Featherston P. 2012. Aquatic Resource Use by Indigenous Australians in Two
Tropical River Catchments: The Fitzroy River and Daly River. Human Ecology **40**:893–908.
Springer.
Jackson S, Finn M, Scheepers K. 2014. The use of replacement cost method to assess and manage
the impacts of water resource development on Australian indigenous customary economies.
Journal of Environmental Management **135**:100–109. Elsevier.
Karim F, Dutta D, Marvanek S, Petheram C, Ticehurst C, Lerat J, Kim S, Yang A. 2015. Assessing the
impacts of climate change and dams on floodplain inundation and wetland connectivity in
the wet–dry tropics of northern Australia. Journal of Hydrology **522**:80–94.
Knape J. 2016. Decomposing trends in Swedish bird populations using generalized additive mixed
models. Journal of Applied Ecology **53**:1852–1861. John Wiley & Sons, Ltd.
Kupschus S, Schratzberger M, Righton D. 2016. Practical implementation of ecosystem monitoring
for the ecosystem approach to management. Journal of Applied Ecology **53**:1236–1247. John
Wiley & Sons, Ltd.
Leahy SM, Robins JB. 2021. River flows affect the growth of a tropical finfish in the wet-dry rivers of
northern Australia, with implications for water resource development. HydrobiologiaDOI:
10.1007/s10750-021-04641-7. Available from https://doi.org/10.1007/s10750-021-04641-7.
Lowe EC, Steven R, Morris RL, Parris KM, Aguiar AC, Webb CE, Bugnot AB, Dafforn KA, Connolly RM,
Mayer Pinto M. 2022. Supporting urban ecosystem services across terrestrial, marine and
freshwater realms. Science of The Total Environment **817**:152689.
Mace PM. 1994. Relationships between common biological reference points used as thresholds and
targets of fisheries management strategies. Canadian Journal of Fisheries and Aquatic
Sciences **51**:110–122.
Macreadie PI, Nielsen DA, Kelleway JJ, Atwood TB, Seymour JR, Petrou K, Connolly RM, Thomson AC,
Trevathan-Tackett SM, Ralph PJ. 2017. Can we manage coastal ecosystems to sequester
more blue carbon? Frontiers in Ecology and the Environment **15**:206–213. John Wiley &
Sons, Ltd.
Maes J, Czúcz B, Keith H, Jackson B, Nicholson E, Dasoo M. 2020. A review of ecosystem condition
accounts: lessons learned and options for further development. One Ecosystem **5**:e53485.
Mapstone BD. 1995. Scalable decision rules for environmental impact studies: Effect size, type I, and
type II errors. Ecological Applications **5**:401–410. Ecological Society of America.
Martínez-López J, Bagstad KJ, Balbi S, Magrach A, Voigt B, Athanasiadis I, Pascual M, Willcock S, Villa
F. 2019. Towards globally customizable ecosystem service models. Science of the Total
Environment **650**:2325–2336. Elsevier B.V.
McMahon JM, Hasan S, Brooks A, Curwen G, Dyke J, Ange CS, Smart JCR. 2022. Challenges in
modelling the sediment retention ecosystem service to inform an ecosystem account –
Examples from the Mitchell catchment in northern Australia. Journal of Environmental
Management **314**:115102.
Millar RB, Meyer R. 2000. Non-linear state space modelling of fisheries biomass dynamics by using
Metropolis-Hastings within-Gibbs sampling. Journal of the Royal Statistical Society: Series C
(Applied Statistics) **49**:327–342. Wiley Online Library.
Milton DA, Chenery SR. 2005. Movement patterns of barramundi Lates calcarifer, inferred from
$^{87}Sr/^{86}Sr$ and Sr/Ca ratios in otoliths, indicate non-participation in spawning. Marine
Ecology Progress Series **301**:279–291. Inter-Research.
Ndehedehe CE, Onojeghuo AO, Stewart-Koster B, Bunn SE, Ferreira VG. 2021. Upstream flows drive
the productivity of floodplain ecosystems in tropical Queensland. Ecological Indicators
125:107546.

Nickols KJ, White JW, Malone D, Carr MH, Starr RM, Baskett ML, Hastings A, Botsford LW. 2019.
Setting ecological expectations for adaptive management of marine protected areas. Journal
of Applied Ecology **56**:2376–2385.
Petheram C, Watson I, Bruce C. 2018. Water resource assessment for the Mitchell catchment. A
report to the Australian Government from the CSIRO Northern Australia. Water Resource
Assessment, part of the National Water Infrastructure Development Fund: Water Resource
Assessments. CSIRO, Canberra, Australia.
Pettorelli N, Vik JO, Mysterud A, Gaillard J-M, Tucker CJ, Stenseth NC. 2005. Using the satellite-
derived NDVI to assess ecological responses to environmental change. Trends in Ecology &
Evolution **20**:503–510.
R Development Core Team. 2019. R: A language and environment for statistical computing. R
Foundation for Statistical Computing, Vienna, Austria. URL: www.R-project.org.
Roberts BH, Morrongiello JR, King AJ, Morgan DL, Saunders TM, Woodhead J, Crook DA. 2019.
Migration to freshwater increases growth rates in a facultatively catadromous tropical fish.
Oecologia **191**:253–260.
Robins JB, Halliday IA, Staunton-Smith J, Mayer DG, Sellin MJ. 2005. Freshwater-flow requirements
of estuarine fisheries in tropical Australia: a review of the state of knowledge and application
of a suggested approach. Marine and Freshwater Research **56**:343–360.
Rowland JA, Lee CKF, Bland LM, Nicholson E. 2020. Testing the performance of ecosystem indices for
biodiversity monitoring. Ecological Indicators **116**:106453.
Scheepers K, Jackson S. 2012. Indigenous people's socio-economic values and river flows in the
Mitchell River delta, Cape York. CSIRO: Water for a Healthy Country Flagship Report.
State of Queensland Department of Agriculture Fisheries and Forestry. 2018. QFish data cube.
State of Queensland Department of Environment and Science,. 2021. The Long Paddock. Available
from https://www.longpaddock.qld.gov.au/about/resources/.
Stevenson SL, Watermeyer K, Caggiano G, Fulton EA, Ferrier S, Nicholson E. 2021. Matching
biodiversity indicators to policy needs. Conservation Biology **35**:522–532. Wiley Online
Library.
Streipert S, Filar J, Robins J, Whybird O. 2019. Stock assessment of the barramundi (Lates calcarifer)
fishery in Queensland, Australia. May 2019. Queensland Government. State of Queensland.
Available from http://era.daf.qld.gov.au/id/eprint/7003/ (accessed July 8, 2021).
Sutherland GD, Waterhouse FL, Smith J, Saunders SC, Paige K, Malt J. 2016. Developing a systematic
simulation-based approach for selecting indicators in strategic cumulative effects
assessments with multiple environmental valued components. Ecological Indicators **61**:512–
525. Elsevier.
Turschwell MP, Stewart-Koster B, King AJ, Pusey B, Crook D, Boone E, Douglas M, Allsop Q, Jackson S,
Kennard MJ. 2019. Flow-mediated predator–prey dynamics influence fish populations in a
tropical river. Freshwater Biology **64**:1453–1466.
Underwood AJ. 1992. Beyond BACI: the detection of environmental impacts on populations in the
real, but variable, world. Journal of Experimental Marine Biology and Ecology **161**:145–178.
Watermeyer KE, Guillera-Arroita G, Bal P, Burgass MJ, Bland LM, Collen B, Hallam C, Kelly LT,
McCarthy MA, Regan TJ. 2021. Using decision science to evaluate global biodiversity indices.
Conservation Biology **35**:492–501. Wiley Online Library.

Figures

Figure 1 Overview of the approach showing the measured variables, the latent variables and the

scope of the model.

Figure 2 Model fits to data for CPUE (A), NDVI (B); pasture biomass (C); flow (D) and the estimated

latent variables for ecological condition (variable v_t in the model) (E) and barramundi biomass

relative to its carrying capacity as estimated by the model (F). For model predictions, lines show

median and shaded areas show 95% CIs. Darker areas in E-F indicate 50% CIs. The solid horizontal

line in (E) indicates the mean value, in (F) it indicates BMSY of 50% carrying capacity.

Figure 3 Strength of the relationships between each indicator and (A) a doubling of flow; and (B) a 1

S.D. increase in the ecological condition index. Note that NDVI and pasture are differences on a scale

standardized to 1 S.D., whereas surplus production is on a multiplicative scale. Horizontal bars show

median, thick solid lines show 75% probability quantiles and thin grey lines show 95% probability

quantiles.

Figure 4 Bounds for catch per unit effort (A) NDVI (B) and pasture biomass (C). Grey zones show 40%

(dark grey), 60% (medium grey) and 90% (light grey) probability intervals. Examples of trend lines

shown in blue shades, as proportion of the mean.

595 **Figure 5** Emergence times for the three indicators, each panel represents a different lower bound:

596 2.5% = 95% probability intervals, 5% = 90% probability intervals and 10% = 80% probability intervals.