Title: Spatio-temporal patterns of multi-trophic biodiversity and food-web characteristics

across a river catchment

Author list:

Rosetta C. Blackman[1, 2, *]

Hsi-Cheng Ho[1]

Jean-Claude Walser[3]

Florian Altermatt[1, 2, *]

Author affiliation:

[1] Eawag, Swiss Federal Institute of Aquatic Science and Technology, Department of Aquatic

Ecology, Überlandstrasse 133, CH-8600 Dübendorf, Switzerland

[2] Department of Evolutionary Biology and Environmental Studies, University of Zurich,

Winterthurerstr. 190, CH-8057 Zürich, Switzerland

[3] Genetic Diversity Centre (GDC), Department of Environmental Systems Sciences (D-USYS),

Federal Institute of Technology (ETH), Rämistrasse 101, 8092 Zurich, Switzerland

* Corresponding Authors:

Email: rosiecblackman@gmail.com or florian.altermatt@eawag.ch

ORCID IDs:

Rosetta C. Blackman - https://orcid.org/0000-0002-6182-8691

Hsi-Cheng Ho - https://orcid.org/0000-0002-0734-0249

Jean-Claude Walser - https://orcid.org/0000-0003-1513-0783

Florian Altermatt - https://orcid.org/0000-0002-4831-6958

Abstract

Accurate characterisation of ecological communities with respect to their biodiversity and food-

web structure is essential for conservation. However, combined empirical study of biodiversity

and multi-trophic food webs at a large spatial and temporal resolution has been prohibited by the

lack of appropriate access to such data from natural systems. Here, we assessed biodiversity and

food-web characteristics across a 700 km^2 riverine network through time using environmental

DNA. We find contrasting biodiversity patterns, with richness (α-diversity) of fish increasing

towards downstream positions within the catchment, while freshwater bacteria and invertebrates

having an invariant and minimal decrease in richness, respectively, with downstream position.

Food-web characteristics, such as link density and nestedness, however, were relatively

conserved across space, but varied over season. Patterns of biodiversity across major taxonomic

groups are thus not directly scalable to food-web structures at the same spatial and temporal

scales, indicating that effective conservation measures must consider them jointly.

Introduction

The study of biodiversity patterns[1–3] and the characterisation of food-web structures[4,5] are

essential, yet often disconnected goals in ecology. Understanding these patterns is not only of

fundamental interest, but also needed to predict stability, functioning and resilience of natural

ecosystems and to bend the curve of biodiversity loss in the context of anthropogenic pressures

including contemporary global change[6].

Studies on biodiversity predominantly focus on analyses of α-, β- and γ -diversity and possible

underlying fundamental drivers of their spatial or temporal patterns[7]. Freshwater rivers are

highly spatially structured systems[8–10] in which theoretical and empirical studies have identified

characteristic patterns of biodiversity for specific groups. For example, fish α-diversity has been

found to increase with distance downstream[11], whereas headwaters often show high endemic

bacterial species richness[12]. Aquatic invertebrate biodiversity exhibits more complicated overall

patterns with disproportionately high biodiversity being found in headwaters[13] and a significant

increase in biodiversity linked to catchment size[14]. However, these group-specific biodiversity

patterns have been mostly studied in isolation from one another, although species are present

within the same system and trophically interact with each other. Indeed, recent theoretical work

shows that contrasting patterns driven by species' resource competition are possible[15]. Therefore,

to ensure optimal strategies for conservation and understanding of biodiversity patterns across

different organismal groups, an ensemble approach integrating major taxonomic and trophic

groups is crucial to reveal how species are linked through trophic interactions[16].

Trophic interactions and food webs by definition encompass multiple groups of organisms.

Individual freshwater food webs are well-resolved[17], and often exhibiting distinct features, such

as highly nested structures[18] and prevalent omnivory[17,19]. Nevertheless, food-web studies often

have a localised perspective due to methodological limitations of sampling food-web interactions

and organismal occurrence in a standardized and comparable manner across different places and

organismal groups[20–22]. Due to the same reason, these studies also tend to focus on simple spatial

and environmental gradients[23] or temporal change[24] when spatio-temporal influences should be

considered in conjunction[25,26]. This is particularly problematic in freshwater riverine ecosystems,

that are characterised by a high spatio-temporal structure: they exhibit characteristic spatial

structures, and have strong seasonal variations driven by changing abiotic conditions[27] and

pronounced life-cycle changes of key taxa inhabiting these systems. The variation in dynamics

over the course of a year remains a significant gap in our understanding of freshwater food

webs[17,28].

To effectively conserve riverine biodiversity, we must encompass spatio-temporal variation of

multiple trophic levels to understand the underlying dynamics of both biodiversity patterns and

food-web characteristics[4,25,26]. In particular, molecular monitoring techniques may now provide a

suitable solution to break through the above-mentioned methodological constraints caused by

sampling based primarily on species sight or capture. Environmental DNA, or eDNA, is the

collection of DNA extracted from an environmental sample such as water, air or sediment[29]. By

collecting eDNA we can screen samples for multiple taxonomic groups via metabarcoding[30],

thereby creating a biodiversity assessment suitable for food-web reconstruction.

Here, we use eDNA metabarcoding to assess patterns of biodiversity and reconstruct local food

webs via a metaweb-based approach[31]. The metaweb approach is suitable to capture food-web

patterns based on taxa co-occurrence data[24,32] in both terrestrial and aquatic systems. We used

high-throughput sequencing and a multi-marker approach to examine three taxonomically

relevant groups, namely fish (via the 12S barcode region), invertebrates (cytochrome *c* oxidase I

- COI) and bacteria (16S) in a large-scale river network over the course of three seasons (spring,

summer and autumn). This approach allowed us to test for association of biodiversity patterns

and food-web structures with network location and seasonal change. We found contrasting

effects on biodiversity patterns and food-web structure from spatial and temporal influences,

respectively, providing insight into the underlying changing ecosystem dynamics and indicating

that effective and targeted conservation measures must consider them jointly.

Results

Data collection and community construction

We collected water samples from the upper Thur catchment, Switzerland, which covers

approximately 700 km^2 and is made up of three sub-catchments: River Thur, Necker and Glatt.

Seventy-three field sites were selected to allow for maximum coverage across the catchment

area, comprising of a broad size range of upstream drainage sizes (i.e., the drainage area size

indicates the location of the site within the catchment: upstream sites having a small drainage

area and lowland sites having a larger drainage, see Fig. 1). Water samples were filtered on site

and DNA extracted in a specialist clean lab environment at Eawag, Switzerland. A total of 12.29

million, 14.32 million and 14.02 million raw reads were produced from the 12S, COI and 16S

libraries, respectively. After bioinformatic processing the average sequencing read depth per

sample was: 43,176, 50,704 and 49,857 for 12S, COI and 16S, respectively. This included 159

Zero-radius Operational Taxonomic Units (ZOTUs), 3179 ZOTUs and 11,320 ZOTUs for 12S,

COI and 16S respectively (see Methods and Supplementary Information Table S1 for details on

laboratory and bioinformatic analysis). For further analysis, ZOTUs were merged to genus level

and only fish genera, invertebrate genera with an aquatic life stage and bacteria associated with

freshwater were kept for further analysis (See Methods for further details). Analysis was

performed on presence/absence data to merge the three libraries for the complete freshwater

community and exclude a possible influence of uneven sample read depth generated from

multiple markers (see also [33]).

To quantify biodiversity, we calculated α-diversity (local richness) at genus level at each site and

compared β-diversity (variation in community composition between sites) by using Jaccard

dissimilarity. Jaccard dissimilarity was also partitioned into taxon replacement (turnover) and

taxon loss (nestedness) components to assess the mechanisms contributing to the variation in

community assemblage across the catchment.

To measure food webs and functional characteristics of the community, we constructed a

metaweb based on known interactions of genera classified into different functional feeding

groups (Fig. 2; see also Methods and Supplementary Information Table S2). We defined local

food webs at each field site based on genera co-occurrence and the corresponding subset of

interactions from the metaweb, to determine broad changes in the community over time and

space. With this approach, variation in food-web structures and function feeding groups emerge

from the spatio-temporal differences in genus composition (See Methods for further details).

To assess the relationship between α-diversity of each group (fish, invertebrates and bacteria)

and food-web structures with site location within the catchment, we ran linear mixed model

analysis. Drainage area (km^2) was log transformed to fit model assumptions. For each dependent

variable, drainage area and season were the fixed effects, while site was the random effect. We

determined the overall effect of both factors using analysis of variance and contrast testing of

estimated marginal means to determine the influence of seasonal changes on all α-diversity and

food-web elements (see Methods for details). To examine the effect of river distance on β-

diversity we constructed a matrix of pairwise distances for sites that were connected along the

fluvial network and to examine the effect of river distance on β-diversity we performed a Mantel

test (see Methods and Supplementary information).

Spatial and temporal biodiversity patterns

In total we detected 374 genera across all organismal groups associated with freshwater,

including 12 fish genera, 80 invertebrate genera and 282 bacteria genera. When combining all

seasons, α-diversity (genus richness) ranged between 8–96 genera with all taxonomic groups

combined (Fig. 1). Over the different seasons, mean local α-diversity was 70 (range 10–92) in

spring, 48 (range 8–85) in summer, and 63 (range 19–96) in autumn (See Supporting Information

Fig. S1 and Table S3). We used mixed models to assess the influence of drainage area and

season on the local α-diversity of each group (Fig. 3, for model output see Supporting

Information Table S4–S6). Of the three taxonomic groups, only fish α-diversity significantly

increased with the size of the drainage area (p < 0.001, Fig. 3a and Supporting Information for

model output Table S4– S6), while for bacteria and invertebrates there was no significant

relationship (p = 0.670 and p = 0.239, respectively, Fig. 3c and 3e). There was, however, a

significant seasonal effect in α-diversity on invertebrate and bacteria genera (p < 0.001 for both

groups, Fig. 3d and 3f, see Supporting Information Table S4–S6). The influence of seasons can

be seen further in the contrast testing (Fig. 3b, d and f and Supporting Information Table S7 for

full contrast testing results), which shows invertebrate α-diversity in spring was significantly

higher compared to summer and autumn (p < 0.001, Fig. 3d), and bacteria α-diversity in summer

was significantly lower compared to spring and autumn (p < 0.001, Fig. 3f). Fish α-diversity

however did not vary significantly between seasons, with the mean number of 2 (range 0–6) fish

genera remaining constant (Fig. 3b).

Regarding the β-diversity across the catchment, the Jaccard's dissimilarity significantly

increased with the river distance among sites in all organismal groups across all seasons, apart

from bacteria in spring and autumn (fish Mantel statistics: Spring 0.143, p < 0.01, Summer

0.171, p = 0.001, Autumn 0.321, p = 0.001; invertebrate Mantel statistics: Spring 0.095, p < 0.05,

Summer 0.169, p = 0.001 and Autumn 0.114, p < 0.05; bacteria Mantel statistics: Spring -0.058,

p = 0.834, Summer 0.179, p < 0.005 and Autumn 0.069, p = 0.12, See Supplementary

Information Table S8). Further partitioned analyses on taxon replacement and loss revealed

contrasting patterns. Taxon replacement between sites increased over river distance for all

groups in most seasons (significant or marginally significant, see Table S8), apart from fish in

spring (Mantel statistics 0.07, p = 0.891) invertebrates in spring (Mantel statistics 0.066, p =

0.051) and bacteria in spring and summer (Mantel statistics 0.016, p = 0.34 and Mantel statistics

0.07, p = 0.068, respectively). In contrast, taxon loss was only found to significantly increase

over river distance for fish in spring and autumn (Mantel statistics 0.219, p = 0.001 and 0.059, p

= 0.05, respectively) and bacteria in summer (Mantel statistics 0.089, p < 0.05).

Spatial and temporal changes in food-web structure and functional characteristics

We examined commonly used food-web structural characteristics (link density, connectance,

nestedness, omnivory, coherence, number of links, modularity, and robustness), as well as

functional characteristics (functional diversity and redundancy). Functional characteristics were

examined by using the designated functional feeding groups (e.g., shredders and omnivorous

fish), based on specialised feeding behaviours (Fig. 2 & Supporting Information Fig. S2). Here

we describe the results of link density, connectance, nestedness and omnivory as the most

ecologically important food-web descriptors in our study, while the results from the remaining

descriptors (coherence, number of links, modularity, and robustness) are presented in the

Supplementary Information (See Fig. S3). Drainage area did not significantly influence any of

the food-web structures looked at (Fig. S3 and S4). However, season had a significant influence

on the change in food-web structures (Fig. 4 and S4), namely, link density (p < 0.001, Fig. 4a),

connectance (p < 0.001, Fig. 4b), nestedness (p < 0.001, Fig. 4c), and omnivory (p < 0.001, Fig.

4c and see Supporting Information Table S4 - S6 for full results), indicating that seasonal

variation is of more importance in food-web dynamics than site location within the riverine

network. To examine seasonal variation further, we carried out contrast testing, which showed a

range of seasonal change in food-web dynamics. In particular, spring had significantly higher

link density than summer and autumn (p < 0.001 and p < 0.001, respectively, Fig. 4a); autumn

had significantly lower connectance than spring and summer (p < 0.001 and p < 0.001,

respectively, Fig. 4b), and the same is true for nestedness ($p < 0.01$ and $p < 0.01$, respectively,

Fig. 4c); finally, summer had significantly higher omnivory than autumn ($p = 0.001$, Fig 4d, see

Supporting Information Table S7 for full contrast testing results).

Functional diversity (defined as the change in function feeding groups used in this study)

significantly decreased with drainage area and was also significantly different across seasons (p

< 0.05 Fig. 5a and $p < 0.001$ Fig. 5b, respectively, see Supporting Information Table S4 - S6 for

full results), whereas only season has a significant influence on the functional redundancy ($p <$

0.001, Fig. 5d). To further examine the changes in seasonal influence we compared seasonal

values for all functional characteristics. Functional diversity in spring was significantly higher

than in summer and autumn ($p = 0.001$ and $p = 0.001$, respectively, Fig. 5b), while functional

redundancy in summer was significantly lower than in spring and autumn ($p < 0.001$ and $p <$

0.001, respectively, Fig. 5d and Supporting Information Table S7).

Discussion

Studies of biodiversity and food-web assemblages in riverine networks are often constrained to

local scales or aggregated to a single time point, which in essence fails to capture the spatial

processes and the temporal fluctuations that together play a key role in community dynamics

present within a river network[8]. Our study is a first assessment utilising data derived from eDNA

metabarcoding to detect patterns of biodiversity and food-web characteristics across three major

taxonomic groups, namely fish, invertebrates and bacteria, in a whole river network at a spatial

and temporally large scale. In our study we find contrasting patterns of biodiversity across these

groups, which indicates different mechanisms may shape these organismal communities. By

using a metaweb approach, we showed a strong signature of seasonality in food-web structures

across the river network. We also showed that functional characteristics are influenced by both

spatial and temporal changes. Overall, our study supports the need to include both spatial and

temporal scales in order to understand changes in ecosystems, particularly as we see increased

effects of contemporary global change.

Aquatic biodiversity is subject to fluvial influences within a dendritic network, whereby spatial

patterns of α-diversity and β-diversity are known for key groups, such as fish, invertebrates and

microbes[11–14,34]. Our data is congruent with previous studies on fish diversity[11,33], in that both α-

diversity significantly increased downstream, and β-diversity (community dissimilarity)

significantly increased with river distance in all three seasons. However, patterns for

invertebrates and bacteria differed from the predictions, and exhibited no significant influence of

drainage size on α-diversity but were influenced by seasonal change. For invertebrates, seasonal

variation can be linked directly to the emergence of the non-aquatic adult life stage of several

macroinvertebrate genera we detected (Diptera, Ephemeroptera, Plecoptera and Trichoptera),

which takes place in the late spring and summer months[27], and literally removes these organisms

from the aquatic food-web as they become air-bound and often end up in terrestrial food-webs.

For the spatial patterns, past studies often looked at spatial scales much larger or much smaller,

and thus disparity may be due to a mismatch in scale looked at here. For example, Finn and

colleagues[13] suggested headwaters harbour disproportionately high invertebrates compared to

lower points in the catchment, but their studied focused on small (1–2) and mid (3–4) stream

orders only, whereas the sites in our study ranged from small to much larger rivers (stream

orders 1–7, [35]). Contrastingly, when looking at scales about 50-fold larger, Altermatt and

colleagues[14] showed the number of key aquatic invertebrate taxa (Ephemeroptera, Plecoptera and

Trichoptera) increased with catchment size, however they also found a combination of local

factors (catchment areas, drainage area, elevation and network centrality) had the greatest

influence on local invertebrate α-diversity. Possibly, the scale at which we study invertebrate α-

diversity and contributing local factors falls between such small-scale vs large-scale perspective,

but instead we detected regional patterns where α-diversity remains relatively constant

throughout the catchment. Similarly, drainage area did not have a significant effect on bacterial

α-diversity, and β-diversity increased with river distance in summer only. We can therefore

postulate that aquatic bacteria are able to persist as they disperse through the catchment;

however, the number of bacteria genera (the α-diversity) that do persist are subject to seasonal

influences. Our findings expand on past studies, highlighting the influences of scale and

temporal variation on different groups, which is of utmost importance when trying to conserve

biodiversity and understand the differing drivers behind these patterns in a river network.

By resolving the fundamental trophic relationships among broad feeding groups, we established

a trophic interaction metaweb for the three taxonomic groups examined in this study. The meta-

web with genus co-occurrence data thus allowed our investigation of both spatial and temporal

influences on the characteristics of local food webs. We found that the most significant factor

driving freshwater riverine food-web characteristics was season. All food-web structural

characteristics (apart from the number of links and robustness) were lowest in autumn, despite

overall α-diversity being lowest in summer. This indicates that the food-web structural variation

we detected is not merely reflecting the genera richness dynamics over seasons, but the genera

composition dynamics instead. In other words, the structures of the food webs are more

influenced by those genera absent in autumn. Indeed, previous studies on freshwater food webs

that have examined temporal changes have often found reduced productivity as the main driver

behind a declined food-web structure in winter vs summer[20,36]. Thus, the decrease in structure we

captured in this study could be the start of the productivity restriction seen in food webs over the

winter months. The less-connected and less-nested food-web structure also implies weaker

resource competition among consumers, which may be necessary for their coexistence in the

food web[37], especially when competition becomes more costly with lower resource productivity

in autumn.

Moving from broad food-web patterns to examining functional feeding groups enables us to

study fundamental changes within the community and possible effects at the ecosystem level[17].

Interestingly, and contrary to the biodiversity patterns and food-web dynamics, we see

significant effects of both drainage area and season on the functional diversity, which we found

to be higher in smaller drainage areas and in spring. The spatial part of such a pattern along the

hierarchical river network is demonstrated in the River Continuum Concept, which shows

increased diversity of invertebrate functional feeding groups at low stream orders[38,39]. Whereas

both drainage area and season influences functional diversity, season is the only significant

influence on functional redundancy, with fewer genera per functional feeding group observed in

summer rather than in autumn. These results indicate that in summer the genera absent are across

functional groups, whereas in autumn the genera absent include whole functional feeding groups

leading to the constriction of local food webs. The inconsistent temporal patterns of food-web

structure versus functional characteristics further imply that multiple feeding groups share

similar trophic roles in the food web (Fig. S2), though they each adopt specialised feeding

behaviour within the catchment and thus perform distinctive ecological functions (e.g.,

shredders, collectors, filterers). In other words, the structure and function of a food web does not

necessarily match and synchronise (*sensu stricto*[22,40]). These results are particularly encouraging,

because the patterns of both food-web structure and functional diversity are known to be

important for ecosystem health assessment and identifying potential vulnerability to

perturbation[17,26,41]. Addressing the consistency of their patterns across spatial and temporal scales

will likely lead to novel and comprehensive understanding of biodiversity and ecosystem

function loss due to environmental change.

In our study, the functional feeding groups were defined at the level of genus, while we expect

investigations at finer resolutions will be promising for future work that can reveal not only more

accurate patterns, but also the influences of sampling taxonomic resolution. Similarly, our

selection of focal taxa may have influenced the food-web patterns we detected. For example,

algae become an increasingly important resource when moving from allochthonous inputs in

headwaters to larger streams with increased light levels further down the catchment[38,39]. With the

selected three key taxonomic groups in riverine ecosystems, we present a broad and relevant

view capturing trophic roles from basal resources (cyanobacteria) to top consumers (piscivorous

fish), captured by three relatively broad metabarcoding markers. However, we also by default

excluded some further groups, such as algae or terrestrial taxa, which could have been relevant

as primary producers and as terrestrial-aquatic linkages, respectively. The choice of taxonomic

groups looked at was both methodologically defined, as well as driven by the goal to have an

overseeable and clearly aquatic-focused view on food webs. Thereby, it captures by default a

subset from the real-world food web in which more species are involved.

By using the eDNA technique, we gain three notable advantages: its scalability for monitoring

complex and large systems, its reusable nature, and its being a non-invasive method of collecting

biodiversity information[42,43]. Our understanding of how the information we ascertain from eDNA

sample collection has greatly improved in recent years due to studies on the hydrological

influences[44] and a general understanding of the rate of eDNA persistence in lotic ecosystems[45].

However, the successful detection of taxa with eDNA is also linked to the ecology of individual

species[45], and some seasonal variation in the detection of several taxonomic groups is known[46–

49]. Therefore, it is possible that some taxa that were not detected in the colder season (autumn) in

this study were false-negative records. However, these non-detections are likely linked to low

abundance or low metabolic rates, and thus these species are, while not physically absent, at least

"relatively absent" in ecological terms.

In summary, our work showed we can construct comprehensible food webs at a large scale and

over time by using eDNA sampling and combining multiple markers. Based on the biodiversity

patterns we observed that spatial and temporal influences are different across groups.

Furthermore, temporal influences were the significant driver of change for commonly used food

web descriptors. Our approach is a first demonstration of the application of eDNA to a complex

river network for the reconstruction of food web patterns and can be easily replicated in other

systems worldwide. As biodiversity in freshwater systems face huge threats from anthropogenic

pressures, including global climate change, establishing vital information on the changes in

biodiversity and food web composition over spatio-temporal scales is essential for the detection

of these stressors in order to protect and conserve river systems.

Methods

Site selection and eDNA sample collection

Environmental DNA samples were collected from the edge of the waterbody using single use

disposable 50 ml syringes. At each site on each sampling event, a total of 1 L of water was

filtered through two 0.22 μm sterivex filters (Merck Millipore, Merck KgaA, Darmstadt,

Germany), sealed with luer caps and placed in individually labelled bags. Sampling was carried

out over five consecutive days in May (spring), August (summer) and October (autumn) in 2018.

During the summer sampling campaign, four sites were dry and could not be sampled (total

samples across all seasons is n = 215). Negative field control samples were collected on each

sampling day by filtering 500 ml of ddH$_2$O through a sterivex filter in the same way as field

samples were collected (n = 15). Negative field samples were processed alongside field samples.

All samples were placed in labelled bags and cooled in a cool box until frozen at -20 °C on

return to the laboratory.

Extraction and library preparation

DNA extraction and first round PCR set up was carried out in a specialist eDNA clean lab, with

separate lab facilities for post PCR workflows. Samples were extracted using DNeasy

PowerWater Sterivex Kit (Qiagen, Hilden, Germany) following the manufacturer's protocol.

Prior to extraction, each sterivex was defrosted at room temperature and then wiped with 10%

bleach, then 70% ethanol solution prior to extraction to remove any DNA from the outside of the

filters. Extraction controls were carried out alongside sample extractions and consisted of blank

sterivex filters (n = 5). Samples selected at random were analysed using a QuBit 3.0 fluorometer

for double stranded DNA concentration, values measured between 0.317 - 27.5 ng/μl. All

negative controls (field and extraction) were tested and recorded below detection limits.

Samples were sequenced with the following markers: a 106 bp fragment of the mitochondrial

12S marker ([50], hereafter referred to as 12S) used to amplify vertebrate DNA, a 313 bp fragment

of the mitochondrial cytochrome oxidase I marker ([51] and [52], hereafter referred to as COI) used to

amplify metazoan DNA and a 450 bp fragment of the V3-V4 region of the 16S marker

(hereafter, 16S) used to amplify bacteria and archaea DNA (See Supporting Information Table

S8 for primer sequences). Positive controls (n = 6 per library prep, see Supporting Information

Table S9) and PCR negative controls (2 μl of ddH$_2$O, n = 11 per library prep) were included in

each library. Each library consisted of 252 samples in total (including positive and negative

controls).

Library preparation followed a two-step PCR process for both the 12S and COI markers, the 16S

library was carried out using a three-step PCR[53] (See Supporting Information Methods for full

details), all samples were amplified in triplicate. After the initial amplification, where Nextera®

transposase sequences (Microsynth, AG, Balgach, Switzerland) are added to the PCR product, all

samples were tested for amplification success with the AM320 method on the QiAxcel Screening

Cartridge (Qiagen, Germany). PCR products were cleaned using ZR-96 Plate clean-up Kit

(Zymo Research) following the manufacturers protocol with the minor modification by which

the elution step was prolonged to 2 minutes at 4000 g. The clean amplicons were indexed using

the Illumina Nextera XT Index Kit A, C and D following the manufacturer's protocol (Illumina,

Inc., San Diego, CA, USA). A reaction contained 25 µL 2x KAPA HIFI HotStart ReadyMix

(Kapa Biosystems, Inc., USA), 5 µL of each of the Nextera XT Index adaptors and 15 µL of the

DNA templates. The final PCR products were then cleaned using the Thermo MG Magjet bead

clean up kit (Thermo Fisher Scientific Inc., MA, USA) and a customized programme for the

KingFisher Flex Purification System (Thermo Fisher Scientific Inc., MA, USA) in order to

remove excessive Nextera XT adaptors. The cleaned product of 50 µl was then eluted into a new

plate and stored at 4 °C prior to quantification. For each library, the clean indexed amplicons

were quantified in duplicate on a Spark Multimode Microplate Reader (Tecan, US Inc. USA)

prior to equimolar pooling using the BRAND Liquid Handling Station (BRAND GMBH + CO

KG, Wertheim, GE). Negative extraction and PCR controls were added according to their

concentration. Library concentration was quantified on the Agilent 4200 TapeStation (Agilent

Technologies, Inc., USA) and verified by the Qubit the HS Assay Kit. Paired-end sequencing

was performed on an Illumina MiSeq (Illumina, Inc. San Diego, CA, USA) at the Genetic

Diversity Centre at ETH (See supporting information Table S1 for library loading information).

Bioinformatics

After each of the libraries were sequenced, the data was demultiplexed and reads were quality

checked using usearch v11.0.667 [54] and FastQC[55]. Raw reads were first end-trimmed, merged

and full-length primer sites were removed using usearch v11.0.667 [54] (16S reads were merged

using Flash v1.2.11, [56]). The merged and primer trimmed reads were quality filtered using

prinseq-lite (0.20.4). The UNOISE3 (usearch v11.0.667) workflow with an additional clustering

of 99% (usearch v11.0.667) identity was applied to obtain error corrected and chimera-filtered

sequence variants ZOTUs. The ZOTUs were then clustered using a 97 %, similarity approach

and taxonomic assignment with a 0.85 confidence threshold was performed using SINTAX in

the usearch software v11.0.667 [54] with following databases: 12S: NCBI BLAST (v200416), COI:

Custom reference database (Including MIDORI un-trimmed (V20180221) and 16S: GreenGenes

(V13.5), RDP (160411) and SILVA (V128). See Supporting Information Methods for all

bioinformatic parameters and reference database information used for each library. Prior to

downstream analysis, 0.1% of reads were removed from each sample to reduce errors caused by

tag-jumping or minor contamination in library preparation[57].

Biodiversity patterns

We calculated α-diversity (genus richness) at each site and compared β-diversity between sites

by using Jaccard dissimilarity using the betapart R-package[58]. We constructed a matrix of

pairwise distances between sites along the fluvial network with the igraph R-package[59] and

compared the similarity between β-diversity and river distance using the Mantel test with the

vegan R-package[60]. To further partition β-diversity into species loss and replacement, sites which

did not record genus from a target group were removed from the analysis of that group only.

Feeding group assignment

Feeding groups were determined based on literature, species inventories and targeted expert

knowledge[61,62] for fish and invertebrates. Genera of bacteria were included if the phyla they

belong to has a strong association with freshwater habitat[63], bacteria were then broadly divided

into heterotrophic and cyanobacteria, the latter constituting a basal resource. A constant basal

resource of detritus was also included in all food webs (See Supporting Information Table S2 for

assignment of genera to feeding groups).

Food-web structure

We constructed a metaweb based on known trophic relationships among feeding groups

(Supplementary Table S2), then defined local food webs using this metaweb at each field site,

based on co-occurrence of genera (nodes) and their interactions (links). The metaweb approach

has been applied to identify food-web characteristic change across spatial gradients in terrestrial

ecosystems[32], and temporal changes in aquatic ecosystems[24], but has yet to be used at a large

spatial scale in an aquatic ecosystem over time.

We then quantified common food-web structural characteristics for each local food web

generated. These included fundamental node-link composition features, i.e., number of links (L),

link density (L/S, the number of feedings links per taxa divided by the genus α-diversity at each

site) and connectance (L/S^2, the proportion of realised interactions). To further explore the

holistic topology of these food webs and their change over space and time, we adopted the

following indices to determine some more key features. Nestedness is an indication that the diets

of specialist taxa are subsets of generalist taxa's diets, and was calculated using the nestedness

metric based on Overlap and Decreasing Fill (NODF) function in vegan[60]. Modularity is

indication of a less nested network[64] in that the nodes are characterised into modules which,

unlike nested structures, do not overlap, and was calculated using the multilevel.community()

function from the igraph package[59]. Omnivory is defined as the proportion of taxa with a mixed

trophic level diet. Thus, the more omnivores a community has, the less coherent the food web[65].

Omnivory is often debated as a measure of food-web stability, and weak omnivorous interactions

are likely to lead to a more stable food web[66]. Trophic level of genera and omnivory were

calculated using the methods stated in William and Martinez[21]. Coherence is defined as the

overall degree of homogeneity of the difference between trophic levels of every consumer-

resource pair within the local food web. As described by [65], the coherence of a network can

reliably be used to establish the stability of a network by looking at the distribution of trophic

distances over all links in each network. For example, a perfectly coherent network ($q = 0$)

implies that each taxon within the food-web only occupies a single trophic level. Coherence was

calculated using the methods stated in Johnson et al[65]. Robustness was defined by Dunne and

Williams[5] as the "*proportion of species subjected to primary removals that resulted in a total*

loss … of some specified proportion of the species". In our study we used 50% as the proportion

of species lost and excluded basal resources for primary removal. Robustness was calculated

using the methods stated in Dunne and Williams[5].

To explore the change in functional characteristics within the food webs, we used the broad

feeding groups as described when constructing the interaction network (See Fig. 1 and Fig. S2).

Functional diversity was calculated as the number of feeding groups that had at least one genus

present in the sample. Functional redundancy was calculated as α-diversity divided by functional

diversity, reflecting the average number of genera within a feeding group.

Data analysis

We ran linear mixed model analysis to assess the relationship between group genus α-diversity

(bacteria, invertebrates and fish), food-web structural characteristics (coherence, connectance,

link density, number of links, omnivory, modularity, nestedness and robustness) and functional

characteristics (functional diversity and redundancy) with site location within the catchment.

Drainage area (km^2) was log transformed to fit model assumptions. For each dependent variable,

drainage area and season were the fixed effects. We included site as a random effect to account

for repeated sampling of sites (e.g., genus α-diversity ~ drainage area + season + (1 | Site)).

Significance was calculated using the lmerTest R-package[67], which applies Satterthwaite's

method to estimate degrees of freedom and generate p-values for mixed models (Table S4 and

S5). We then used anova() to see the overall effect of both factors: Drainage area and Season

(Table S6) and follow up contrast testing were carried out using the emmeans R-package [68] to

test for significant differences between seasons (Table S7).

Acknowledgments: We thank Samuel Hürlemann and Jeanine Brantschen for their help in the

laboratory and field. We also thank Xing Xing, Silvana Kaeser, Elvira Mächler and Roman

Alther for their assistance with field sample collection and Silvia Kobel and Aria Minder (GDC),

for their help with library preparation. The data generation and analysis in this paper was

generated in collaboration with the Genetic Diversity Centre (https://gdc.ethz.ch), ETH Zurich.

Funding is from the University of Zurich Research Priority Programme Global Change and

Biodiversity and the Swiss National Science Foundation (grants No. PP00P3_179089 and

31003A_173074) to FA.

Author contributions:

R.C.B and F.A conceived the study. R.C.B performed the genomic lab analyses and J.C.W.

performed the bioinformatics. R.C.B assigned trophic levels to all taxa and constructed the

metaweb, and H.H performed food-web analysis. R.C.B performed all statistical analyses. R.C.B

produced all figures and illustrations. All authors contributed to the interpretation of the

networks and discussed and commented on the paper.

Competing interests:

The authors confirm there are no competing interests.

Materials & Correspondence:

Correspondence to Rosetta C. Blackman or Florian Altermatt

Data availability

Sequencing data generated during this study will be made available in a public repository upon

publication.

References

References

1.   Whittaker, R. H. Vegetation of the Siskiyou mountains, Oregon and California. *Ecol.*

Monogr. **30**, 279–338 (1960).

2.   Wilson, R. J., Thomas, C. D., Fox, R., Roy, D. B. & Kunin, W. E. Spatial patterns in

species distributions reveal biodiversity change. *Nature* **432**, 393–396 (2004).

3.   Newbold, T. *et al.* Global effects of land use on local terrestrial biodiversity. *Nature* **520**,

45–50 (2015).

4.   Ings, T. C. *et al.* Ecological networks--beyond food webs. *J. Anim. Ecol.* **78**, 253–269

(2009).

5.   Dunne, J. A. & Williams, R. J. Cascading extinctions and community collapse in model

food webs. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **364**, 1711–1723 (2009).

6.   Leclère, D. *et al.* Bending the curve of terrestrial biodiversity needs an integrated strategy.

Nature **585**, 551–556 (2020).

7.   Vellend, M. *The Theory of Ecological Communities*. vol. 57 229 (Princeton University

Press., 2016).

8.   Altermatt, F. Diversity in riverine metacommunities: a network perspective. *Aquat. Ecol.*

47, 365–377 (2013).

9.   Peterson, E. E. *et al.* Modelling dendritic ecological networks in space: an integrated

network perspective. *Ecol. Lett.* **16**, 707–719 (2013).

10.   Tonkin, J. D. *et al.* The role of dispersal in river network metacommunities: Patterns,

processes, and pathways. *Freshw. Biol.* **63**, 141–163 (2018).

11.   Muneepeerakul, R. *et al.* Neutral metacommunity models predict fish diversity patterns in

Mississippi-Missouri basin. *Nature* **453**, 220–222 (2008).

12.   Besemer, K. *et al.* Headwaters are critical reservoirs of microbial diversity for fluvial

networks. *Proc. Biol. Sci.* **280**, 20131760 (2013).

13. Finn, D. S., Bonada, N., Múrria, C. & Hughes, J. M. Small but mighty: headwaters are vital

to stream network biodiversity at two levels of organization. *jnbs* **30**, 963–980 (2011).

14. Altermatt, F., Seymour, M. & Martinez, N. River network properties shape α-diversity and

community similarity patterns of aquatic insect communities across major drainage basins.

J. Biogeogr. **40**, 2249–2260 (2013).

15. Harvey, E., Gounand, I., Fronhofer, E. A. & Altermatt, F. Disturbance reverses classic

biodiversity predictions in river-like landscapes. *Proceedings of the Royal Society B:*

Biological Sciences **285**, 20182441 (2018).

16. Jason M. Tylianakis, Etienne Laliberté, Anders Nielsen, Jordi Bascompte. Conservation of

species interaction networks. *Biol. Conserv.* **143**, 2270–2279 (2010).

17. Thompson, R. M. *et al.* Food webs: reconciling the structure and function of biodiversity.

Trends Ecol. Evol. **27**, 689–697 (2012).

18. Woodward, G. & Hildrew, A. G. Food web structure in riverine landscapes. *Freshw. Biol.*

47, 777–798 (2002).

19. Williams, R. J. & Martinez, N. D. Limits to trophic levels and omnivory in complex food

webs: theory and data. *Am. Nat.* **163**, 458–468 (2004).

20. Thompson, R. M. & Townsend, C. R. The Effect of Seasonal Variation on the Community

Structure and Food-Web Attributes of Two Streams: Implications for Food-Web Science.

Oikos **87**, 75–88 (1999).

21. Wood, S. A., Russell, R., Hanson, D., Williams, R. J. & Dunne, J. A. Effects of spatial scale

of sampling on food web structure. *Ecol. Evol.* **5**, 3769–3782 (2015).

22. J. M. Tylianakis & R.J. Morris. Ecological networks across environmental gradients.

Annual Review of Ecology, Evolution, and Systematics **48**, 25–48 (2017).

23. Romanuk, T. N. *et al.* The Structure of Food Webs along River Networks. *Ecography* **29**,

3–10 (2006).

24. Olivier, P. *et al.* Exploring the temporal variability of a food web using long-term

biomonitoring data. *Ecography* **42**, 2107–2121 (2019).

25. Poisot, T., Canard, E., Mouillot, D., Mouquet, N. & Gravel, D. The dissimilarity of species

interaction networks. *Ecol. Lett.* **15**, 1353–1361 (2012).

26. Delmas, E. *et al.* Analysing ecological networks of species interactions. *Biol. Rev. Camb.*

Philos. Soc. (2018) doi:10.1111/brv.12433.

27. Tavares-Cromar, A. F. & Williams, D. D. The importance of temporal resolution in food

web analysis: Evidence from a detritus-based stream. *Ecol. Monogr.* **66**, 91–113 (1996).

28. Poisot, T., Stouffer, D. B. & Gravel, D. Beyond species: why ecological interaction

networks vary through space and time. *Oikos* **124**, 243–251 (2015).

29. Thomsen, P. F. & Willerslev, E. Environmental DNA – An emerging tool in conservation

for monitoring past and present biodiversity. *Biol. Conserv.* **183**, 4–18 (2015).

30. Deiner, K., Fronhofer, E. A., Mächler, E., Walser, J.-C. & Altermatt, F. Environmental

DNA reveals that rivers are conveyer belts of biodiversity information. *Nat. Commun.* **7**,

12544 (2016).

31. Dunne, J. A. The network structure of food webs. in *Ecological Networks: Linking*

Structure and Dynamics (ed. Pascual, J. A. & Dunne, J. A.) 27–86 (University Press,

Oxford., 2006).

32. Neff, F. *et al.* Changes in plant-herbivore network structure and robustness along land-use

intensity gradients in grasslands and forests. *Sci Adv* **7**, (2021).

33.   Blackman, R. C. *et al.* Mapping biodiversity hotspots of fish communities in subtropical

streams through environmental DNA. *Scientific Reports* **4**, e65352 (2021).

34.   Widder, S. *et al.* Fluvial network organization imprints on microbial co-occurrence

networks. *Proc. Natl. Acad. Sci. U. S. A.* **111**, 12799–12804 (2014).

35.   Mächler, E. *et al.* Assessing different components of diversity across a river network using

eDNA. *Environmental DNA* **1**, 290–301 (2019).

36.   Peralta-Maraver, I., López-Rodríguez, M. J. & de Figueroa, J. M. T. Structure, dynamics

and stability of a Mediterranean river food web. *Mar. Freshwater Res.* **68**, 484–495 (2017).

37.   Kondoh, M., Kato, S. & Sakato, Y. Food webs are built up with nested subwebs. *Ecology*

91, 3123–3130 (2010).

38.   Vannote, R. L., Minshall, G. W., Cummins, K. W., Sedell, J. R. & Cushing, C. E. The River

Continuum Concept. *Can. J. Fish. Aquat. Sci.* **37**, 130–137 (1980).

39.   Power, M. E. & Dietrich, W. E. Food webs in river networks. *Ecol. Res.* (2002)

doi:10.1046/j.0912-3814.2002.00503.x.

40.   Montoya, D., Yallop, M. L. & Memmott, J. Functional group diversity increases with

modularity in complex food webs. *Nat. Commun.* **6**, 7379 (2015).

41.   Gravel, D., Albouy, C. & Thuiller, W. The meaning of functional trait composition of food

webs for ecosystem functioning. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **371**, (2016).

42.   Ruppert, K. M., Kline, R. J. & Rahman, M. S. Past, present, and future perspectives of

environmental DNA (eDNA) metabarcoding: A systematic review in methods, monitoring,

and applications of global eDNA. *Global Ecology and Conservation* **17**, e00547 (2019).

43.   Altermatt, F. *et al.* Uncovering the complete biodiversity structure in spatial networks: the

example of riverine systems. *Oikos* **129**, 607–618 (2020).

44.   Carraro, L., Mächler, E., Wüthrich, R. & Altermatt, F. Environmental DNA allows

605   upscaling spatial patterns of biodiversity in freshwater ecosystems. *Nat. Commun.* **11**, 3585

606   (2020).

607   45.   Barnes, M. A. & Turner, C. R. The ecology of environmental DNA and implications for

608   conservation genetics. *Conserv. Genet.* **17**, 1–17 (2016).

609   46.   Bista, I. *et al.* Annual time-series analysis of aqueous eDNA reveals ecologically relevant

610   dynamics of lake ecosystem biodiversity. *Nat. Commun.* **8**, 14087 (2017).

611   47.   Erickson, R. A., Merkes, C. M., Jackson, C. A., Goforth, R. R. & Amberg, J. J. Seasonal

612   trends in eDNA detection and occupancy of bigheaded carps. *J. Great Lakes Res.* **43**, 762–

613   770 (2017).

614   48.   Troth, C. R., Sweet, M. J., Nightingale, J. & Burian, A. Seasonality, DNA degradation and

615   spatial heterogeneity as drivers of eDNA detection dynamics. *Sci. Total Environ.* **768**,

616   144466 (2021).

617   49.   Thalinger, B. *et al.* The Effect of Activity, Energy Use, and Species Identity on

618   Environmental DNA Shedding of Freshwater Fish. *Frontiers in Ecology and Evolution* **9**,

619   73 (2021).

620   50.   Kelly, R. P., Port, J. A., Yamahara, K. M. & Crowder, L. B. Using environmental DNA to

621   census marine fishes in a large mesocosm. *PLoS One* **9**, e86175 (2014).

622   51.   Leray, M. *et al.* A new versatile primer set targeting a short fragment of the mitochondrial

623   COI region for metabarcoding metazoan diversity: application for characterizing coral reef

624   fish gut contents. *Front. Zool.* **10**, 34 (2013).

625   52.   Geller, J., Meyer, C., Parker, M. & Hawk, H. Redesign of PCR primers for mitochondrial

626   cytochrome c oxidase subunit I for marine invertebrates and application in all-taxa biotic

627  surveys. *Mol. Ecol. Resour.* **13**, 851–861 (2013).

628 53. Mansfeldt, C. *et al.* Microbial community shifts in streams receiving treated wastewater

629  effluent. *Sci. Total Environ.* **709**, 135727 (2020).

630 54. Edgar, R. C. Search and clustering orders of magnitude faster than BLAST. *Bioinformatics*

631  **26**, 2460–2461 (2010).

632 55. Andrews, S. FASTQC A Quality Control tool for High Throughput Sequence Data.

633  *Babraham Institute* (2015).

634 56. Magoč, T. & Salzberg, S. L. FLASH: fast length adjustment of short reads to improve

635  genome assemblies. *Bioinformatics* **27**, 2957–2963 (2011).

636 57. Hänfling, B. *et al.* Environmental DNA metabarcoding of lake fish communities reflects

637  long-term data from established survey methods. *Mol. Ecol.* **25**, 3101–3119 (2016).

638 58. Baselga, A. & Orme, C. D. L. betapart : an R package for the study of beta diversity :

639  Betapart package. *Methods Ecol. Evol.* **3**, 808–812 (2012).

640 59. Csárdi, G. & Nepusz, T. The igraph software package for complex network research. *Inter*

641  *Journal, Complex Systems* **1695**, 1–9 (2006).

642 60. Oksanen, J. *et al.* vegan: Community Ecology Package 2.5-6. *https://CRAN.R-*

643  *project.org/package=vegan* (2019).

644 61. Tachet H., Bournaud M., Richoux P. & Usseglio-Polatera P. *Invertébrés d'eau douce -*

645  *systématique, biologie, écologie.* (CNRS Editions, Paris., 2010).

646 62. Schmidt-Kloiber, A. & Hering, D. www.freshwaterecology.info – An online tool that

647  unifies, standardises and codifies more than 20,000 European freshwater organisms and

648  their ecological preferences. *Ecol. Indic.* **53**, 271–282 (2015).

649 63. Newton, R. J., Jones, S. E., Eiler, A., McMahon, K. D. & Bertilsson, S. A guide to the

natural history of freshwater lake bacteria. *Microbiol. Mol. Biol. Rev.* **75**, 14–49 (2011).

64. Fortuna, M. A. *et al.* Nestedness versus modularity in ecological networks: two sides of the

same coin? *J. Anim. Ecol.* **79**, 811–817 (2010).

65. Johnson, S., Domínguez-García, V., Donetti, L. & Muñoz, M. A. Trophic coherence

determines food-web stability. *Proc. Natl. Acad. Sci. U. S. A.* **111**, 17923–17928 (2014).

66. Wootton, K. L. Omnivory and stability in freshwater habitats: Does theory match reality?

Freshw. Biol. **62**, 821–832 (2017).

67. Kuznetsova, A., Brockhoff, P. B. & Christensen, R. H. B. lmerTest Package: Tests in Linear

Mixed Effects Models. *Journal of Statistical Software, Articles* **82**, 1–26 (2017).

68. Lenth, R. V. Estimated Marginal Means, aka Least-Squares Means [R package emmeans

version 1.6.1]. (2021).

Figures:

Figure 1: Genus α-diversity of bacteria, invertebrates and fish collected from eDNA

samples in the Thur catchment in (a) spring, (b) summer and (c) autumn. The circles are

scaled proportionate to the α-diversity of each group. Grey lines give the sub-catchments Glatt,

Necker, and Thur, respectively and the blue arrow shows the direction of flow.

Figure 2: Food-web network generated from eDNA data. All genera were categorised into

feeding groups dependant on their dominant feeding behaviour, we then used this information to

construct a trophic interaction metaweb using these links. Number of genera in each group is

shown in circles, arrows indicate energy flow.

672
673

Figure 3: Genus α-diversity in space and time. Plots a, c and e show α-diversity in each group

as a function of drainage area. Line indicates linear mixed model with shaded area showing 95%

confidence intervals as calculated using the model predictions and standard error. Plots b, d and f

show change in α-diversity in each group over season. Grey lines indicate link sample points

over the three sampling seasons. In all plots colour represents season: yellow – Spring, green –

Summer and orange – Autumn.

Figure 4: Food-web structural characteristics: Box plots showing the change in each food-

web structural characteristic over the three seasons: a: Link Density, b: Connectance, c:

Nestedness and d: Omnivory. Grey lines indicate link sample points over the three sampling

seasons and colour represents season: yellow – Spring, green – Summer and orange – Autumn.

Figure 5: Functional diversity and functional redundancy. a and b: Functional Diversity, c

and d: Functional Redundancy. Plots a and c show the change over drainage area. Line indicates

linear mixed model with shaded area showing 95% confidence intervals as calculated using the

model predictions and standard error. Plots b and d show change over season. Grey lines indicate

link sample points over the three sampling seasons and colour represents season: yellow –

Spring, green – Summer and orange – Autumn.