Machine learning models based on molecular descriptors to predict human and environmental toxicological factors in continental freshwater

Rémi Servien[a,b,*], Eric Latrille[a,b], Dominique Patureau[a], Arnaud Hélias[c,d]

[a]INRAE, Univ. Montpellier, LBE, 102 Avenue des étangs, F-11000 Narbonne, France

[b]ChemHouse Research Group, Montpellier, France

[c]ITAP, Univ Montpellier, INRAE, Institut Agro, Montpellier, France

[d]ELSA, Research group for environmental life cycle sustainability assessment and ELSA-Pact industrial chair, Montpellier, France

*corresponding author : remi.servien@inrae.fr

Highlights:

- Characterization factors (for human health and ecotoxicological impacts) were predicted using molecular descriptors.

- Several linear or non-linear machine learning methods were compared.

- The non-linear methods tend to outperform the linear ones using a train and test procedure. Cluster-then-predict approaches often show the best performances, highlighting their usefulness.

- This methodology was then used to derive characterization factors that were missing for more than a hundred chemicals in USEtox®.

Abstract: It is a real challenge for life cycle assessment practitioners to identify all relevant substances contributing to the ecotoxicity. Once this identification has been made, the lack

of corresponding ecotoxicity factors can make the results partial and difficult to interpret. So,

it is a real and important challenge to provide ecotoxicity factors for a wide range of

compounds. Nevertheless, obtaining such factors using experiments is tedious, time-

consuming, and made at a high cost. A modeling method that could predict these factors

from easy-to-obtain information on each chemical would be of great value. Here, we present

such a method, based on machine learning algorithms, that used molecular descriptors to

predict two specific endpoints in continental freshwater for ecotoxicological and human

impacts. The different tested machine learning algorithms show good performances on a

learning database and the non-linear methods tend to outperform the linear ones. The

cluster-then-predict approaches usually show the best performances which suggests that

these predicted models must be derived for somewhat similar compounds. Finally,

predictions were derived from the validated model for compounds with missing

toxicity/ecotoxicity factors.

Graphical abstract:

Keywords: machine learning, Life Cycle Assessment, characterisation factors, toxicity,

ecotoxicity, continental freshwater.

1.   Introduction

Recent legislations such as the Registration, Evaluation, Authorization and restriction of

Chemicals (REACH) regulation in the EU requires that manufacturers of substances and

formulators register to provide toxicological and ecotoxicological data for substances with

volume higher than one metric ton per year. Furthermore, the U.S. Environmental Protection

Agency (EPA) has more than 85,000 chemicals listed under the Toxic Substances Control

Act (Hinds and Weller, 2016). Therefore, robust toxicological and ecotoxicological data are

quickly needed to make informed decisions on how to regulate new chemicals. These data

must also be coupled with environmental exposures and sources data, to better understand

the impact on the environment.

To address the cause-effect relationships between the flow of molecules emitted by human

activities and the consequences for ecosystems and humans, Life Cycle Assessment (LCA)

offers a structured, operational, and standardized (Finkbeiner et al., 2006) methodological

framework. Two main steps are at the core of this approach:

• Quantification of the masses of substances emitted into the environment through the

Life Cycle Inventory (LCI). While it is possible to rely on databases that facilitate this

inventory work for the background of the system under study, this task must

nevertheless be carried out on a case-by-case basis to represent all the specificities

of the foreground elements. This is the task of the LCA practitioner.

• Calculation of the impacts on ecosystems and human health of these emitted

masses. Due to the complexity of environmental mechanisms, it is not possible to

(re)model impact pathways on a case-by-case basis. Therefore, LCA uses

characterization factors (CF) to assess the potential impacts of a compound.

Concretely, if two compounds are emitted with the same mass, the one with the

higher CFs will have the higher impact. Then, CFs are multiplied by the emitted

masses of each compound to determine the impacts. CFs are not recalculated for

each study but provided within a Life Cycle Impact Assessment (LCIA) method.

For a given impact, the LCIA method designer refers to the knowledge of the scientific

community to model the mechanisms involved. For human toxicity and freshwater

ecotoxicity, USEtox® (Rosenbaum et al., 2008), was developed by life cycle initiative under

the United Nations Environmental Programme (UNEP) and the Society for Environmental

Toxicology and Chemistry (SETAC) (Henderson et al. 2011) to produce a transparent and

consensus characterization model. USEtox® is also used for the European Product

Environmental Footprint (PEF) (Saouter et al., 2020). This model gathers in one single

characterization factor the chemical fate, the exposure, and the effect for each of the several

thousands of organic and inorganic compounds. Then, the USEtox® model results can be

extended to determine endpoint effects expressed as total (i.e. cancer and non-cancer)

disability-adjusted life years (DALY) for human health impacts and potentially disappeared

fraction of species (PDF) for ecotoxicological impacts. The PDF represents an increase in

the fraction of species potentially disappearing as a consequence of emission in a

compartment while the DALY represents an increase in adversely affected life years. These

endpoints are now consensual at an international level (Verones et al., 2017).

If the structure of the USEtox® multimedia model is always the same, to determine the CF of

a molecule, numerous physicochemical parameters (such as solubility, hydrophobicity,

degradability) and detailed toxicological and ecotoxicological data must be provided. For

example, EC50 values (i.e. the effective concentration required to have a 50% effect) for at

least three species from three different trophic levels are required for the ecotoxocological

effect factor.

Over the past few decades, thousands of tests (in laboratory and field) have been carried out to evaluate the potential hazard effects of chemicals (He et al., 2017). Usually, toxicity testing has relied on *in vivo* animal models, which is extremely costly and time-consuming (Xia et al., 2008). In recent years, under societal pressures, there has been a significant paradigm shift in toxicity testing of chemicals from traditional *in vivo* tests to less expensive and higher throughput *in vitro* methods (National Research Council, 2007). However, it is still extremely difficult to test existing chemicals due to their large and ever increasing number, which leaves their impacts largely unknown. For example, in a recent study, Aemig et al. (2021) studied the potential impacts on Human health and aquatic environment of the release of 286 organic and inorganic micropollutants at the scale of France. One of their conclusion was that, due to a lack of characterization factors, these impacts could be assessed only for 1/3 of these molecules. That is why more computational models are needed to complement experimental approaches to decrease the experimental cost and determine the prioritization for those chemicals which may need further *in vivo* studies. Such models already exist, like QSAR models that are mostly linear models based on the chemical structure of compounds (Danish QSAR database (DTU, 2015), ECOSAR (Mayo-Bean et al., 2011), VEGA (Benfenati et al., 2013)) and are used to predict ecotoxicological data (LC50) needed for REACH for example. Recently, machine learning algorithms have been used to predict some midpoints based on molecular descriptors and environmental parameters (Marvuglia et al., 2014 and 2015; Song et al., 2017; Lysenko et al 2018) and a first review on this subject could be found in Wu and Wang (2018). After these first works, predictions of hazardous concentration 50% (HC50) based on 14 physicochemical characteristics (Hou et al., 2020a) or on 691 more various variables (Hou et al., 2020b) were carried out. Nevertheless, their input variables need some experiments and could be difficult to collect. This problem was tackled by Song et al. (2021) who predicted Lethal Concentration 50 (LC50) based on 2000 easy-to-obtain molecular descriptors. In the case of USEtox®, despite its wide use in LCA, it only offers characterization factors for

approximately 3000 chemicals and even for this limited number of compounds, 19% of ecotoxicity CFs and 67% of human toxicity CFs are missing.

The objective of this article is thus to propose a new way of calculating CFs using machine learning approaches to solve the problem of nonlinearity that could affect a linear QSAR method. This makes it possible, when the CFs are not determined due to lack of time or lack of data, to propose values based solely on easily identifiable molecular descriptors. Here, the main differences with the above-cited methods are twofold: first, our input variables are only molecular descriptors that could be easily collected for any newly available compounds; second, our output variables are directly the CFs that are closer to the endpoints (DALY and PDF) than the HC50 or the LC50 (i.e. the acute aquatic toxicity experimental threshold). These two specific endpoints will be studied in the present paper through the emission of compounds in continental freshwater and will be named CF_{ET} for ecotoxicological impacts and CF_{HT} for human ones. To address this aim, we will test different methods (linear and non-linear) and assess their performances, to build a robust model that could predict CFs that are currently lacking.

2. Materials & Methods

2.1. USEtox® database

The last version of the USEtox® database was downloaded, namely the corrective release 2.12 (USEtox®, 2020). The whole USEtox® 2.12 database contains 3076 compounds. The CFs were computed using the default landscape.

2.2. TyPol database

We recently developed TyPol (Typology of Pollutants), a classification method based on statistical analyses combining several environmental parameters (i.e., sorption coefficient,

degradation half-life, Henry constant) and an ecotoxicological parameter (bioconcentration

factor BCF), and structural molecular descriptors (i.e., number of atoms in the molecule,

molecular surface, dipole moment, energy of orbitals). Molecular descriptors are calculated

using an *in silico* approach (combining Austin Model1 and Dragon software). In the present

paper, we only extracted and used the molecular descriptors from the TyPol database, as

this information could be easily collected for any new compound. The 40 descriptors

included in the TyPol database have been selected based on a literature review on QSAR

equations used to predict the main environmental processes as degradation, sorption,

volatilization. These 40 descriptors were the ones most frequently used in the equations,

meaning describing the best the behaviour of organic compounds in the environment. By

consequence, even if no environmental parameters were directly incorporated as input in our

model, some information that is directly linked to them were included in the 40 molecular

descriptors. These descriptors are constitutional, geometric, topological, and quantum-

chemical descriptors (see Table 1); 35 described the 2D-structure of the compound while the

other five are linked to its 3D-structure. An important advantage of the unique use of

molecular descriptors is that they are easily and quickly computable for not yet synthesized

compounds. For more details, we refer the interested reader to Servien et al. (2014) and to

Mamy et al. (2015) where the choice of the 40 molecular descriptors is described in details.

Now, TyPol gathers 526 compounds, including pesticides, persistent chemicals,

pharmaceuticals and their transformation products (Benoit et al. 2017, Traoré et al. 2018).

Table 1 – List of the 40 molecular descriptors in TyPol

Category	Molecular descriptors		
Constitutional	Number of atoms	Number of non-H atoms	Number of hydrogen atoms
	Number of chlorine atoms	Number of carbon atoms	Number of nitrogen atoms

	Number of oxygen atoms	Number of phosphorus atoms	Number of sulfur atoms
	Number of fluorine atoms	Number of circuits	Number of halogen atoms
	Number of bonds	Number of non-H bonds	Number of double bonds
	Number of triple bonds	Number of multiple bonds	Number of rotatable bonds
	Number of aromatic bonds	Sum of conventional bond order	Number of rings
	Molecular weight		
Geometric	Connolly molecular surface area		
Topological	Connectivity index of order 0	Connectivity index of order 1	Connectivity index of order 2
	Connectivity index of order 3	Connectivity index of order 4	Connectivity index of order 5
	Valence connectivity index of order 0	Valence connectivity index of order 1	Valence connectivity index of order 2
	Valence connectivity index of order 3	Valence connectivity index of order 4	Valence connectivity index of order 5
Quantum-chemical	Polarizability	Electric dipole moment	HOMO energy
	LUMO energy	Total energy	

4   **2.3.   Machine learning methods**

To predict the CFs using the molecular descriptors we used three modeling methods

combined. The first method is a linear well-known prediction method namely the Partial

Least Squares (PLS) (Wold, 1985). It finds the multidimensional directions in the observable

variable (molecular descriptor) space that explains the maximum multidimensional variance

direction in the predicted variable (CF) space. That provides a linear regression model based

on the observable variables to predict the predicted variable. We also chose to compare two

machine learning methods adapted to non-linear problems: the random forest (Breiman

2001) and the support vector machines (SVM) (Drucker et al. 1996). Random forests are a

machine learning method, for classification or, in our case, regression, that operate by

constructing a multitude of decision trees that uses a random subset of the training data and

limits the number of variables used at each split and outputting the mean prediction

(regression) of the individual trees. SVM constructs a hyperplane or set of hyperplanes in a

high- or infinite-dimensional space in which the problem is linearly separable.

These choices allowed us to compare several ideas. The PLS is a simple linear method that

will not exhibit good performances if the underlying relationship is not linear. The SVM and

RF methods are well-known non-linear machine learning algorithms that used to show good

results in this kind of problem (Hou et al., 2020a).

All the models were computed in the freeware R (R core team, 2019). The PLS has been

computed using the package mixOmics (Rohart et al., 2017), the random forests using the

package randomForest (Liaw et al., 2002), and the SVM using the package e1071 (Meyer et

al., 2019). These 3 modeling methods have some parameters that needed to be fixed: the

number of latent components for the PLS (fixed using the tune.pls function), the number of

variables randomly sampled as candidates at each split for the random forests (selected

using the tune.randomForest function) and, for the SVM, the gamma parameter of the radial

kernel and the cost of constraints violation (using the tune.svm function). All these different

tune functions are based on cross-validation (i.e. a training/test procedure to find the best

value for the parameters) using default function values.

2.4.   Clustering-based model

A recent popular way to make predictions is to use a cluster-then-predict approach. That is,

clustering is used for pre-classification which is to arrange a given collection of input patterns

into natural meaningful clusters. Then, the clustering results are used to construct a predictor

in each cluster. The main idea of the cluster-then-predict approach is that if the clustering

performs well, the prediction will be easier by modeling only similar compounds. If a new

compound with no CF_{ET} and/or CF_{HT} is investigated, the clustering can easily be applied to it

before the prediction model itself. The cluster-then-predict approach has already been

applied with success in various domains such as sentiment prediction (Sony et al., 2015),

finance (Tsai et al., 2014), chemometrics (Minh Maï Le et al., 2018). So we decided to use

the clustering given by the TyPol application (more details in Servien et al., 2014). Note that

the TyPol clustering has already been shown relevant on various occasion: in combination

with mass spectrometry to categorize tebuconazole products in soil (Storck et al., 2016), to

explore the potential environmental behaviour of putative chlordecone transformation

products (Benoit et al., 2017) or to classify pesticides with similar environmental behaviors

(Traore et al., 2018; Mamy et al., 2021). So, the clustering procedure of TyPol was applied

on the whole database of 526 compounds using the 40 molecular descriptors. This approach

provided us a global clustering based on all the available information contained in the TyPol

database. It is based on PLS, hierarchical clustering and an optimal choice of the number of

clusters and is detailed in Servien et al. (2014). The obtained clustering is given in

Supplementary Figure S1 and relies on 5 different clusters. The Supplementary Figure S2

represents this clustering restricted to the common molecules between TyPol and USEtox®.

We could see that, as the cluster 5 is only constituted of one compound, the cluster-then-

predict-models cannot be applied.

Based on this clustering, we then defined three other competing methods. For these

methods, a different model (with different parameters) was derived for the compounds in

each cluster. Consequently, six different models were calibrated and tested for each CF

prediction: global PLS, global SVM, global random forest, cluster-then-PLS, cluster-then-

SVM and cluster-then-Random Forest.

2.5.   Comparison procedure of the models

To assess the performances of the different models we used the following procedure:

1. Split each cluster (the whole dataset if the model is global, only the data lying in the

11   dedicated cluster if that is a cluster-then-predict model) between a training set (85%

12   of the dataset) and a test set (15%) (Pareto principle). The test set is not used for any

13   step of the procedure (such as the imputation of the missing data, the calibration of

14   the parameters …).

15   2. Imputation of the NA (Not Available, *i.e.* missing) values (less than 1%) in the

16   descriptor matrix using the NIPALS algorithm (Wold, 1985).

17   3. Tune the parameters and train the specific models by performing cross-validation on

18   the training set. We have 3 global models to train (PLS, random forest, and SVM)

19   and the cluster-then-test models (PLS, random forest and SVM for each cluster).

4. Test the different models on the test set. Compute the absolute error.

5. Back to step 1.

The whole algorithm was repeated 200 times. All the performances were compared in terms

of absolute error. The absolute error is the absolute difference between the prediction and

the true value. It has been shown to be the most natural and unambiguous measure of error

(Willmott et Matsuura, 2005) and is chosen to be easily comparable to the assumed error on

the experimental CFs (2-3 logs, see Rosenbaum, 2008). For each cluster, we chose the

model with the lowest median absolute error.

2.6 Predictions

Then, the best model was calibrated and computed on the whole cluster. Finally, it was

applied to the compounds, according to their clusters, with a CF_{ET} (or a CF_{HT}) equals to NA

to provide a prediction. For the compounds in cluster 5, this best model cannot be a cluster-

then-predict one and, by consequence, is a global one. To assess the robustness of our

prediction we derived a 95% prediction interval for each prediction. The type of model and its

corresponding parameters were fixed during this process, according to the best model of the

cluster. For example, if the best model of cluster 1 was the random forest approach, random

forest models are used with the parameters optimized during the previous step. Then, we

performed a leave-one-out bootstrap on the dataset that was used to compute the model

(the whole dataset if the model is global, only the data lying in the dedicated cluster if that is

a cluster-then-predict model) and a new model was computed on this leave-one-out sample.

A prediction was carried for each leave-one-out model (*i.e.* n-1 models if n is the number of

compounds of the, eventually global, cluster) and the 2.5% and 97.5% quantile of these

predictions were computed and considered as the prediction interval (Hou et al., 2020a). The

whole modeling process is summarized in the following Figure 1.

Figure 1 Schematic representation of the modeling procedure adopted in the paper.

The five molecular descriptors contributing the most to the prediction were then derived for

each chosen model to assess the differences between models and to interpret their

relevance. For a random forest model, these descriptors are calculated using variable

permutations (Breiman, 2001), for the SVM they are the descriptors with the higher

coefficients in absolute value.

3.   Results

3.1.   Descriptive analysis of the intersection of the TyPol and the USEtox®

databases

As the objective of this proof-of-concept study was to predict USEtox® CF_{ET} and CF_{HT} using

the molecular descriptors contained in TyPol, we could only use the compounds that are

present in both databases. This resulted in 274 compounds that are detailed in Table S1 in

supplementary material and the range of their CF_{ET} and CF_{HT} values are summarized in the

boxplots in Figures 2 and S3. Note that for the 274 common compounds there are 15 NA

values for the CF_{ET} and 102 for the CF_{HT}.

Figure 2 Boxplots of the CF_{HT} for the USEtox® database and the common molecules

between the USEtox® and the TyPol databases. This CF_{HT} is equal to $\log_{10}((DALY+\varepsilon).kg^{-1})$.

The ε is needed as some values of the DALY are exactly equal to zero. ε has been chosen

equal to 1e-10 to be below the minimum of the USEtox® database (5e-9).

We could see on these two figures that the common compounds present higher CF_{ET} and

CF_{HT} values than the one of the complete USEtox® database: it focuses on the more

dangerous compounds as their boxplots are above the USEtox® counterparts and they

cover the whole order of magnitude of the CFs of the USEtox® database.

3.2.   Clustering of the compounds

The global Typol clustering of Supplementary Figure S1 with only the common compounds

is plotted in Supplementary Figure S2 and the boxplots of each molecular descriptor per

cluster are given in Supplementary Figure S4 with different indicators in Table S2. We could

see that they are clustered in 5 groups with different sizes (respectively 33 compounds in the

first black cluster, 122 compounds in the second red cluster, 91 compounds in the third

green cluster, 27 compounds in the fourth blue cluster, and one compound in the fifth brown

cluster). Cluster 1 grouped compounds with a high number of aromatic bonds, double bonds,

rotatable bonds, and multiple bonds. Cluster 2 is an intermediate one between clusters 1

and 3, with less extreme values. Cluster 3 is made of compounds with the lowest molecular

mass. Cluster 4 gathered compounds presenting a high number of halogens, rings, and

circuits. The unique compound in the fifth cluster is erythromycin (highest molecular mass

and number of H and C, lowest number of rings) and, obviously, no cluster-then-predict

model could be built for this cluster

As a first analysis of the clustering given by TyPol, we could see in Figure 3 below the

boxplots of the CF_{ET} and CF_{HT} within the 5 clusters.

Figure 3- Boxplot by cluster for the CF_{ET} and CF_{HT} values. Note that the unique compound

of Cluster 5 has no CF_{HT} value. The size of the clusters and the numbers of NA are gathered

in the legend.

The predictions will be made difficult for the CF_{ET} of cluster 1 as it covers a wide range

whereas it includes a relatively small number of compounds. On the contrary, cluster 3

covers a small range with no extreme values and includes a high number of compounds, for

this cluster the cluster-then-predict approach could produce interesting results.

3.3.   Performances of the machine learning methods

1   **3.3.1.**   **For the prediction of the CF_{ET}**

The methodology described in the previous section was applied to our dataset and gave the

results gathered in Figure 4 for each cluster.

Figure 4 - Performances of the different methods (RF: random forest, PLS: partial least

squares, SVM: support vector machines) in terms of the log of the absolute error of the CF_{ET}

with respect to the different clusters for 200 repetitions. In each cluster, the models are

coloured from dark blue (best) to clear blue (worst) according to their median of the absolute

error. The red dotted line represents an absolute error of 1 log that is considered as

acceptable (Rosebaum et al. 2008 ; Douziech et al., 2019).

The performances were not similar from one cluster to another. For example, performances

of all methods for cluster 1 were very poor (median absolute error above 1) whereas

performances for cluster 4 seemed good despite its smallest size (median absolute error

around 0.6). Therefore, a future prediction of an unknown compound which lies in cluster 1 will be less reliable than in other clusters. Note that we could not test this in the next section as no NA value is present in this cluster 1.

The cluster-then-predict methods seemed more appropriate in each cluster. The cluster-then-RF approach had the best performances (with a global median absolute error equals to 0.64 and the best performances on clusters 2 and 3), even if there was not a big difference between the different methods. The cluster-then-SVM was also the best method for the two clusters 1 and 4. The linear methods (PLS and cluster-then-PLS) had higher absolute errors but were competitive. The individual predictions of the best method in each cluster are reported in Figure S5.

3.3.2.   Prediction for the CF_{HT}

Let us recall that we have more NA values for the CF_{HT} (102) than for the CF_{ET} (15). The performances of the methods are illustrated in the following figure.

Figure 5- Performances of the different methods (RF: random forest, PLS: partial least

squares, SVM: support vector machines) in terms of the log of the absolute error of the CF_{HT}

with respect to the different clusters for 200 repetitions. In each cluster, the models are

coloured from dark blue (best) to clear blue (worst) according to their median of the absolute

error. The red dotted line represents an absolute error of 1 log that is considered as

acceptable (Rosebaum et al. 2008 ; Douziech et al., 2019).

We observed that, despite its small size (11 compounds), the CF_{HT} of the first cluster were

well predicted (with the best performance for the cluster-then-RF approach). It could be

explained by the small range of the CF_{HT} values of this cluster, as illustrated on the boxplot

in Figure 3. The performances of all the methods were comparable on clusters 2 and 3

where the best method was the SVM. Cluster 4 seemed to be the most difficult to predict: all

the methods had their worst results on this cluster and, if the SVM had an acceptable

median absolute error of 0.82, all the medians of the other methods were above 1.3. Global

performances of the different methods were given in Supplementary Figure S6. Note that, as

for CF_{ET}, the linear methods based on PLS were outperformed by the other ones.

3.4.   Best model predictions

3.4.1.   Best models for the CF_{ET}

Using our methodology, we could exhibit a median absolute error of 0.62 log for the

prediction of the CF_{ET} on the whole dataset using the best models. If we looked closer on

Table S3, we could see that the median of our estimations is below 0.6 log except for cluster

1 (above 1 log).

Then we calibrated the best models on the whole dataset of each cluster: a cluster-then-

predict approach using SVM for clusters 1 and 4 and using random forest for clusters 2 and

3. To compare the different models in each cluster and give an idea of what were the

important molecular descriptors we provided the five most important molecular descriptors

for each cluster in the Table S4. We could see in this table that the important molecular

descriptors strongly differ from one cluster to another.

Then the models were used to predict the missing CF_{ET} of the common compounds between

USEtox® and TyPol databases. These values were by consequence new estimations of the

CF_{ET} for compounds on which we had no information. The prediction intervals were relatively

small: less than 0.5 log_{10} in a log scale which highlighted the robustness of the estimation.

They are given in Table S5. No NA value was present in cluster 1 with no prediction for this

cluster. For cluster 2 gathering molecules with intermediate molecular mass, 9 CF_{ET} values

were predicted for various kinds of compounds. One value concerns the antibiotic

sulfamethazine and its value is quite near to the one of sulfamethoxazole and sulfadiazine of

the same sulphonamide antibiotic family constituted of the sulphonamide group ($-S(=O)_2$-

NR_2R_3). Cluster 3 grouped compounds with the lowest molecular mass and the lowest

median CF_{ET} like ibuprofen, phthalates, cresol constituted of monoaromatic ring substituted

with methyl, carboxylic groups. The CF_{ET} prediction for acetylsalicylic acid seemed coherent

with the value of the nearest compounds (herbicides mecoprop) of this group. Cluster 4

gathered compounds with the highest median CF_{ET} and that presented a high number of

rings halogenated or not, like PAH and hormones. The 5 CF_{ET} predicted concerned 4 PAHs

and 1 hormone. By comparison to the 2 other PAHs present in this cluster, the 4 predicted

CF_{ET} were quite similar and higher. Concerning the prediction for the hormone, the CF_{ET} was

intermediate between the CF_{ET} of the 3 other hormones in the cluster. It seems that all these

5 predicted values were very closed, falling near the median value of this cluster.

11   **3.4.2.   Best models for the CF_{HT}**

The best models had a global median of 0.75 log for the prediction of the CF_{HT}. We could

see on the results gathered on Table S6 that the best performances are for cluster 1 and

that they are comparable for the other clusters.

Then, the global SVM model was calibrated and computed on the whole dataset. It was used

to predict the compound of clusters 2, 3, 4, and 5. Let us recall that there was a single

molecule in cluster 5 and, as it has a NA value for its CF_{HT}, the best global model (SVM) was

used. For cluster 1, a cluster-then-RF model was computed. The most important descriptors

of these two models are gathered in the Table S7 and, as for CF_{ET}, were strongly different

between the different best models.

Then, this model was used to predict the CF_{HT} value for the 102 common compounds

without a CF_{HT} value. These predictions are reported in Supplementary Table S8. As for the

CF_{ET}, the small width of the prediction interval (less than a log_{10} in a log scale) highlighted

the robustness of the approach even with a relatively small number like estimations made for

compounds that lie in cluster 1. In this cluster 1, CF_{HT} for a phthalate (DEHP) was already

known, but the one for diisodecyl and diisononyl phthalate was predicted with value in the

same range. The 3 cyclines (tetracycline, aureomycin, and oxytetracycline) grouped in cluster 1, presented also similar predicted CF_{HT}. This was also the case for triclosan and triclocarban in cluster 2. Similar predicted and known CF_{HT} were found for four herbicides from the substituted urea family (linuron, diuron, monolinuron, isoproturon) in cluster 3. Cluster 4 gathered a small number of molecules but with the highest median CF_{HT}, the predicted CF_{HT} of the organochlorine insecticide isodrin was similar to another congener of the same family, aldrin.

4.   Discussion

It is a real and important challenge to provide characterization factors for a wide range of compounds. Obviously, it is expected that these new calculated factors have an acceptable margin of error. As reported in UNEP/SETAC (2019), it is commonly assumed that the uncertainty of the characterization factors can vary by approximately 2-3 orders of log-magnitude (Rosenbaum et al. 2008) or significantly higher (up to 7 orders) if all sources of uncertainty are considered (Douziech et al. 2019). The results obtained in the previous section are very promising as they are below the level of uncertainty commonly assumed and as they are based on molecular descriptors that could be easily obtained for each compound without ecotoxicity factor. Based on this fact we could already provide 15 new CF_{ET} and 102 new CF_{HT} for the common molecules between USEtox® and TyPol without a previous value.

The idea of predicting ecotoxicity characterization factors for chemicals using machine learning algorithms has already been used (Hou et al., 2020a and 2020b). But, here, our findings go further. Indeed, we show that we could directly obtain accurate estimations of endpoint values from easy-to-obtain molecular descriptors. This will open the door to the fast characterization of each new unknown compound that appears, including transformation products. We also show that the cluster-then-predict approach can give better performances

than the approach without the clustering step. This local (*i.e.* cluster-then-predict) approach

confirms that local models could be an efficient prediction method when heterogeneity of

data generates nonlinear relations between the response and the explanatory variables

(Lesnoff et al., 2020).

Across the clusters and models, there is a general trend that the non-linear models tend to

outperform the linear ones. This suggests that a linear model is not fully adequate to capture

the complexity of the relationship between the molecular descriptors and the CFs. However,

the use of linear model for e.g. a QSAR is likely due to the ease of interpreting its

coefficients, while interpretation is much more challenging for machine learning approaches

such as random forest or SVM. Thus, the advantages or drawbacks of linear/non-linear

approaches must be balanced according to the final goal of each study. Here, as the main

goal is to calculate the most accurate CFs, non-linear models seem more suited. We must

also mention that a new emerging field is developing tools needed to help making black-box

models (e.g. random forest) more interpretable (Bénard et al., 2021).

The difficult interpretability of the machine learning models used in this study can thus be

viewed as a limitation. On another side, even if we already had an acceptable number of

compounds in our training datasets, the model accuracies would benefit of the inclusion of

new compounds. These compounds could be carefully chosen to improve the models where

there is a clear need (i.e. where the performances of the models are not good enough), for

example in the cluster 1 for CF_{ET} or in the cluster 4 for CF_{HT}.

One of the interests of USEtox® and its three-step structure (fate - exposure - effect) is that

it can be adapted to some specific contexts (a more accurate and spatialized fate model, a

different exposure...) while keeping the steps that are not modified. However, these

adaptations of USEtox® are not widely used and are reserved for advanced users. Our

approach does not allow this, with a direct one-step estimation of CFs. It was designed to

provide default CF values for molecules where information is missing. We have chosen to

directly predict the CF by simplicity, as the first tests revealed that doing three models (for

the three steps) and then calculating the CFs produced less accurate results. It would

however be an interesting perspective to estimate only some of the stages by these learning

approaches and to combine them with stages modelled in a classical way in USEtox®

5.   Conclusion

This paper presents a modeling method to derive characterization factors from easily obtainable molecular descriptors. The results presented here show that models that can handle non-linearity and that could be adapted to a small number of compounds (using the cluster-then-predict approaches) are the best suited. The cluster-then-predict approaches could also be more accepted by the users, as they allow to consider mainly the compounds similar to the one under investigation. By consequence, the missing characterization factors, as well as those of new molecules, could now be quickly estimated with an overall good precision as performed in Servien et al. (2021). More generally, one of the key factors in the evaluation of toxicity and ecotoxicity in LCA lies in the construction of the characterization factors: a task requiring a large amount of data and a consequent investment of time. The use of machine learning allows us to go beyond these constraints and to propose a new methodology in the LCA framework. This makes it possible to obtain characterization factor values in a fast and simple way, which can be used as long as conventionally established CFs are not available.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgments

The authors are grateful to Pierre Benoit, Laure Mamy, and Virginie Rossard for their work

on TyPol.

Funding

This research did not receive any specific grant from funding agencies in the public,

commercial, or not-for-profit sectors.

Supplementary materials

Supplementary material associated with this article can be found, in the online version.

Scripts and data used are provided.

References

Aemig, Q., Hélias, A., Patureau, D., 2021. Impact assessment of a large panel of organic

and inorganic micropollutants released by wastewater treatment plants at the scale of

France, Water Research, 188, 116524, https://doi.org/10.1016/j.watres.2020.116524.

Bénard, C., Biau, G., da Veiga, S., Scornet, E, 2021. Interpretable random forests via rule

extraction. In International Conference on Artificial Intelligence and Statistics, vol. 130 of

Proceedings of Machine Learning Research, 937–945 (PMLR, 2021).

Benfenati, E., Manganaro, A., Gini, G.C., 2013. VEGA-QSAR: AI Inside a Platform for

Predictive Toxicology. CEUR Workshop Proceedings, 21-28.

Benoit, P., Mamy, L., Servien, R., Li, Z., Latrille, E., Rossard, V., Bessac, F., Patureau, D.,

Martin-Laurent, F., 2017. Categorizing chlordecone potential degradation products to explore

their environmental fate, Science of the Total Environnement, 574, 781–795. https://doi.org/10.1016/j.scitotenv.2016.09.094.

Breiman, L., 2001. Random Forests, Machine Learning, 45 (1), 5–32. https://doi.org/10.1023/A:1010933404324.

Cortes, C., Vapnik, V., 1995. Support-vector networks, Machine Learning, 20 (3), 273–297. https://doi.org/10.1007/BF00994018.

Douziech, M., Oldenkamp, R., van Zelm, R., King, H., Hendriks, A.J., Ficheux, A.-S., Huijbregts, M.A.J., 2019. Confronting variability with uncertainty in the ecotoxicological impact assessment of down-the-drain products, Environment International, 126, 37-45, https://doi.org/10.1016/j.envint.2019.01.080.

Drucker, H., Burges, C.C., Kaufman, L., Smola, A.J., Vapnik, V., 1997, Support Vector Regression Machines, Advances in Neural Information Processing Systems 9, NIPS, 155–161, MIT Press. https://dl.acm.org/doi/10.5555/2998981.2999003.

DTU, 2015. Danish QSAR database. Danish QSAR group, National Food Institute, Technical University of Denmark.

Finkbeiner, M., Inaba, A., Tan, R., Christiansen, K., Klüppel, H.-J., 2006. The New International Standards for Life Cycle Assessment: ISO 14040 and ISO 14044. The International Journal of Life Cycle Assessment, 11 (2), 80–85. https://doi.org/10.1065/lca2006.02.002.

He, J., Tang, Z., Zhao, Y., Fan, M., Dyer, S. D., Belanger, S. E., Wu, F., 2017. The combined QSAR-ICE models: practical application in ecological risk assessment and water

quality criteria, Envronnemental Science & Technology, 51, 8877. https://doi.org/10.1021/acs.est.7b02736.

Henderson, A.D., Hauschild, M.Z., Van De Meent, D., Huijbregts, M.A.J., Larsen, H.F., Margni, M., McKone, T.E., Payet, J., Rosenbaum, R.K., Jolliet O., 2011. USEtox® fate and ecotoxicity factors for comparative assessment of toxic emissions in life cycle analysis: sensitivity to key chemical properties, The International Journal of Life Cycle Assessment, 16, pp. 701-709 https://doi.org/10.1007/s11367-011-0294-6.

Hinds, R.d.C., Weller, J.L., 2016. Toxic Substances Control Act. Environmental Law Practice Guide, vol. 4.

Hou, P., Jolliet, O., Zhu, J., Xu, M., 2020a. Estimate ecotoxicity characterization factors for chemicals in life cycle assessment using machine learning models. Environment International, 135, 105393. https://doi.org/10.1016/j.envint.2019.105393.

Hou, P., Zhao, B., Jolliet, O., Zhu, J., Wang, P., Xu, M., 2020b. Rapid Prediction of Chemical Ecotoxicity Through Genetic Algorithm Optimized Neural Network Models, ACS Sustainable Chemistry & Engineering, 8 (32), 12168-12176. https://dx.doi.org/10.1021/acssuschemeng.0c03660.

Lesnoff, M., Metz, M., Roger, JM., 2020. Comparison of locally weighted PLS strategies for regression and discrimination on agronomic NIR data, Journal of Chemometrics, 34(5), e3209, https://doi.org/10.1002/cem.3209.

Liaw, A., Wiener, M., 2002. Classification and Regression by randomForest, R News, 2(3), 18-22. http://CRAN.R-project.org/doc/Rnews/.

Lysenko, A., Sharma, A., Boroevich, K., Tsunoda, T., 2018. An integrative machine learning

approach for prediction of toxicity-related drug safety, Life Science Alliance, 1(6),

e201800098. https://doi.org/10.26508/lsa.201800098.

Mamy, L., Patureau, D., Barriuso, E., Bedos, C., Bessac, F., Louchart, X., Martin-Laurent, F.,

Miège, C., Benoit, P., 2015. Prediction of the fate of organic compounds in the environment

from their molecular properties: A review, Critical Reviews in Environmental Science and

Technology, 45:1277-1377. http://doi.org/10.1080/10643389.2014.955627.

Mamy, L., Bonnot, K., Benoit, P., Bockstaller, C., Latrille, E., Rossard, V., Servien, R.,

Patureau, D., Prevost, F., Pierlot, F., and Bedos, C., 2021. Assessment of pesticides

volatilization potential based on their molecular properties using the TyPol tool. Journal of

Hazardous Materials, 415, 125613. https://doi.org/10.1016/j.jhazmat.2021.125613.

Marvuglia, A., Kanevski, M., Leuenberger, M., Benetto, E., 2014. Variables selection for

ecotoxicity and human toxicity characterization using Gamma Test. In: B. Murgante et al.

(Eds.): ICCSA 2014, Part III, LNCS 8581, pp. 640–652, 2014. Proceedings of the 14th

International Conference on Computational Science and Applications (ICCSA 2014),

University of Minho, Guimaraes, Portugal. https://doi.org/10.1007/978-3-319-09150-1_47

Marvuglia, A., et al. 2015. Random Forest for toxicity of chemical emissions: features

selection and uncertainty quantification. Journal of Environmental Accounting and

Management 3(3): 229-241. https://doi.org/10.5890/JEAM.2015.09.003.

Mayo-Bean, K., Nabholz, J., Clements, R., Zeeman, M., Henry, T., Rodier, D., Moran, K.,

Meylan, B., Ranslow, P., 2011. Methodology document for the ECOlogical Structure-Activity

Relationship Model (ECOSAR) class program: estimating toxicity of industrial chemicals to

aquatic organisms using ECOSAR class program (Ver. 1.1). In: US Environmental Protection Agency, Office of Chemical Safety and Pollution Prevention, Office of Pollution Prevention and Toxics, Washington, DC.

Meyer, D., Dimitriadou, E., Hornik, K., Weingessel, A., Leisch, F., 2019. e1071: Misc Functions of the Department of Statistics, Probability Theory Group (Formerly: E1071), TU Wien. R package version 1.7-2. https://CRAN.R-project.org/package=e1071.

Minh Maï Le, L., Kégl, B., Gramfort, A., Marini, C., Nguyen, D., Cherti, M., Tfaili, S., Tfayli, A., Baillet-Guffroy, A., Prognon, P., Chaminade, P., Caudron, E., 2018. Optimization of classification and regression analysis of four monoclonal antibodies from Raman spectra using collaborative machine learning approach, Talanta, 184, 260-265, https://doi.org/10.1016/j.talanta.2018.02.109.

National Research Council, 2007. Toxicity Testing in the 21st Century: A Vision and a Strategy; National Academies Press, https://doi.org/10.17226/11970.

R Core Team, 2019. R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria. http://www.r-project.org/index.html.

Rohart, F., Gautier, B., Singh, A., Le Cao, K.-A., 2017. mixOmics: An R package for omics feature selection and multiple data integration, PLoS computational biology, 13(11), e1005752. https://doi.org/10.1371/journal.pcbi.1005752.

Rosenbaum, R.K., Margni, M., Jolliet, O., 2007. A flexible matrix algebra framework for the multimedia multipathway modelling of emission to impacts, Environment International, 33(5),624-634. https://doi.org/10.1016/j.envint.2007.01.004.

Rosenbaum, R.K., Bachmann, T. M., Gold, L. S., Huijbregts, M.A.J., Jolliet, O., Juraske, R., Koehler, A., Larsen, H.F., MacLeod, M., Margni, M., McKone, T.E., Payet, J., Schuhmacher, M., van de Meent, D., Hauschild, M.Z., 2008. USEtox®—the UNEP-SETAC Toxicity Model: Recommended Characterisation Factors for Human Toxicity and Freshwater Ecotoxicity in Life Cycle Impact Assessment, The International Journal of Life Cycle Assessment, 13 (7), 532–546. https://doi.org/10.1007/s11367-008-0038-4.

Saouter, E., Biganzoli, F., Ceriani, L., Versteeg, D., Crenna, E., Zampori, L., Sala, S., Pant, R., 2020. Environmental Footprint: Update of Life Cycle Impact Assessment Methods – Ecotoxicity freshwater, human toxicity cancer, and non-cancer, Publications Office of the European Union, Luxembourg, https://doi.org/10.2760/300987.

Servien, R., Mamy, L., Li, Z., Rossard, V., Latrille, E., Bessac, F., Patureau, D., Benoit, P., 2014. TyPol - a new methodology for organic compounds clustering based on their molecular characteristics and environmental behaviour, Chemosphere, 111, 613–622. https://doi.org/10.1016/j.chemosphere.2014.05.020.

Servien, R., Leenknecht, C., Bonnot, K., Rossard, V., Latrille, E., Mamy, L., Benoit, P., Hélias, A., Patureau, D., 2021. Improved impact assessment of micropollutants release from WWTPs, Case Studies in Chemical and Environmental Engineering, 100172. https://doi.org/10.1016/j.cscee.2021.100172.

Soni, R., Mathai, K.J., 2016. An Innovative 'Cluster-then-Predict' Approach for Improved Sentiment Prediction. In: Choudhary R., Mandal J., Auluck N., Nagarajaram H. (eds) Advanced Computing and Communication Technologies. Advances in Intelligent Systems and Computing, vol 452. Springer, Singapore. https://doi.org/10.1007/978-981-10-1023-1_13.

Song, R., Keller, A.A. and Suh, S., 2017. Rapid life-cycle impact screening using artificial

neural networks, Environmental science & technology, 51(18), pp.10777-10785.

https://doi.org/10.1021/acs.est.7b02862.

Song, R., Li, D., Chang, A., Tao, M., Qin, Y., Keller, A., Suh, S., 2021. Accelerating the pace

of   ecotoxicological   assessment   using   artificial   intelligence.   Ambio.

https://doi.org/10.1007/s13280-021-01598-8.

Storck, V., Lucini, L., Mamy, L., Ferrari, F., Papadopoulou, E.S., Nikolaki, S., Karas, P.A.,

Servien, R., Karpouzas, D.G., Trevisan, M., Benoit, P., and Martin-Laurent, F, 2016.

Identification and characterization of tebuconazole transformation products in soil by

combining suspect screening and molecular typology, Environmental Pollution, 208 B, 537-

545. https://doi.org/10.1016/j.envpol.2015.10.027.

Traore, H., Crouzet, O., Mamy, L., Sireyjol, C., Rossard, V., Servien, R., Latrille, E., Martin-

Laurent, F., Patureau, D., Benoit, P., 2018. Clustering pesticides according to their molecular

properties, fate and effects by considering additional ecotoxicological parameters in the

TyPol   method, Environmental Science and Pollution Research, 25(5), 4728-4738.

https://doi.org/10.1007/s11356-017-0758-8.

Tsai, C.-F., 2014. Combining cluster analysis with classifier ensembles to predict financial

distress, Information Fusion, 16, 46-58. https://doi.org/10.1016/j.inffus.2011.12.001.

UNEP-SETAC, 2019. Global Guidance for Life Cycle ImpactAssessment Indicators: Volume

2. https://www.lifecycleinitiative.org/training-resources/global-guidance-for-life-cycle-impact-

assessment-indicators-volume-2/ (accessed Nov 22, 2020).

USEtox® 2020: USEtox® database system, https://USEtox®.org/model/download.

Verones, F., Bare, J., Bulle, C., Frischknecht, R., Hauschild, M., Hellweg, S., Henderson, A.,

Jolliet, O., Laurent, A., Liao, X., et al., 2017. LCIA Framework and Cross-Cutting Issues

Guidance within the UNEP-SETAC Life Cycle Initiative, Journal of Cleaner Production, 161,

957–967. https://doi.org/10.1016/j.jclepro.2017.05.206.

Willmott, C., Matsuura, K., 2005. Advantages of the Mean Absolute Error (MAE) over the

Root Mean Square Error (RMSE) in Assessing Average Model Performance, Climate

Research, 30, 79. https://doi.org/10.3354/cr030079.

Wold, H., 1985. Partial least squares, In Kotz, Samuel; Johnson, Norman L. (eds.),

Encyclopedia of statistical sciences, vol 6, New York, Wiley.

Wu, Y., Wang, G., 2018. Machine Learning Based Toxicity Prediction: From Chemical

Structural Description to Transcriptome Analysis. International Journal of Molecular Science,

19, 2358. https://doi.org/10.3390/ijms19082358.

Xia, M., Huang, R., Witt, K.L., Southall, N., Fostel, J., Cho, M.-H., Jadhav, A., Smith, C.S.,

Inglese, J., Portier, C.J., Tice, R.R., Austin, C.P., 2008. Compound cytotoxicity profiling

using quantitative high-throughput screening, Environmental Health Perspectives, *116* (3),

284– 291, https://doi.org/10.1289/ehp.10727.