A chimeric KaiA-like regulator extends the nonstandard KaiB3-KaiC3 clock system in bacteria

Christin Köbler[a,1], Nicolas M. Schmelling[b,1], Alice Pawlowski[b,1], Philipp Spät[c], Nina M. Scheurer[a], Lutz Berwanger[b], Boris Maček[c], Ilka M. Axmann[b], Annegret Wilde[a,*]

[a]Institute of Biology III, Faculty of Biology, University of Freiburg, 79104 Freiburg, Germany;
[b]Institute for Synthetic Microbiology, Biology Department, Heinrich Heine University Düsseldorf, 40225 Düsseldorf, Germany; [c]Department of Quantitative Proteomics, Interfaculty Institute for Cell Biology, Eberhard Karls University Tübingen, 72076 Tübingen, Germany

[1]C.K., N.M.S and A.P. contributed equally to this work.

*Prof. Dr. Annegret Wilde, Albert-Ludwigs-Universität Freiburg, Institut für Biologie III, Schänzlestr. 1, 79104 Freiburg, Germany, Phone: +49 (0) 761-20397828

Email: annegret.wilde@biologie.uni-freiburg.de

Author Contributions: C.K., I.M.A. and A.W. designed the study. C.K., N.M.S., A.P., P.S., N.M.Sche. and L.B. performed and analyzed the experiments. All authors interpreted and discussed the data. C.K., N.M.S., A.P., P.S., B.M., I.M.A and A.W. wrote the paper.

Competing Interest Statement: The authors declare no competing interest.

Keywords: cyanobacteria, circadian clock, *Synechocystis* 6803, KaiA

This PDF file includes:

      Main Text
      Figures 1 to 4

Abstract

The rotation of the Earth results in predictable environmental changes that pose challenges for organisms and force them to adapt. To address this daily rhythm, organisms from all kingdoms of life have evolved diverse timing mechanisms. In the cyanobacterium *Synechococcus elongatus* PCC 7942, the three proteins KaiA, KaiB, and KaiC constitute the central timing mechanism that drives circadian oscillations. In addition to the standard oscillator, named KaiAB1C1, *Synechocystis* sp. PCC 6803 harbors several, diverged clock homologs. The nonstandard KaiB3-KaiC3 system was suggested to impact the metabolic switch in response to darkness. Here, we demonstrate the direct interaction of KaiC3 with Sll0485, which is a potential new chimeric KaiA homolog that we named KaiA3. The existence of a functional link between these proteins is further supported by the co-occurrence of genes encoding KaiA3 with the KaiB3–KaiC3-like gene products in 10 cyanobacterial and five other bacterial species. KaiA3 is annotated as a NarL-type response regulator due to its similarity to the response regulator receiver domains. However, its similarity to canonical NarL drastically decreases in the C-terminal domain, which resembles the circadian clock protein KaiA. In line with this, we detected the stimulation of KaiC3 phosphorylation by KaiA3 *in vitro*. Furthermore, we showed that deletion of the *kaiA3* gene led to growth defects during mixotrophic growth conditions and, like a *kaiC3*-deficient mutant, viability was impaired during chemoheterotrophic growth in complete darkness. In summary, we suggest KaiA3 as a novel, nonstandard KaiA homolog within the cyanobacterial phylum, extending the KaiB3-KaiC3 system in Cyanobacteria and other prokaryotes.

Significance Statement

Many organisms that are subjected to day and night cycles are able to predict daily changes using a circadian clock system. Cyanobacteria use a relatively simple timekeeper, a protein-based oscillator consisting of KaiA, KaiB and KaiC. Several cyanobacteria and other prokaryotes harbor additional sets of oscillator components that, however, lack the KaiA protein. Our study identified a divergent KaiA homolog that is functionally and genetically linked to potential KaiB3-KaiC3 timekeepers. The data show that the nonstandard KaiA homolog affects the phosphorylation of KaiC3 from the cyanobacterium *Synechocystis*, suggesting that a similar function exists in other prokaryotes.

Main Text

Introduction

Most organisms on Earth are subjected to predictable environmental changes, such as light and darkness or temperature variations during the day and night cycle. To adapt to these ever-changing conditions, eukaryotic and prokaryotic organisms have evolved endogenous biological clock systems.
To date, cyanobacteria are the only prokaryotes with a well-understood and robust circadian rhythm. The model organism for cyanobacterial circadian clocks, *Synechococcus elongatus* PCC 7942 (hereafter *Synechococcus*), encodes three genes, *kaiA*, *kaiB,* and *kaiC,* that constitute the core oscillator (1). The KaiC protein forms a homohexamer and has autokinase, autophosphatase, ATP synthase, and ATPase activities (1–4). Through binding to KaiC, KaiA stimulates the autokinase and ATPase activities of KaiC and thus the protein is phosphorylated (5–7). Upon phosphorylation of two neighboring residues (Ser431 and Thr432), KaiC undergoes structural rearrangements, exposing a binding site for KaiB (8–10). After binding, KaiB sequesters KaiA from KaiC, promoting KaiC's autophosphatase activity and the protein reverts back to its unphosphorylated state (8, 9, 11). The interplay of KaiA and KaiB is crucial for the KaiC phosphorylation pattern, which in turn confers the clock phase and rhythmicity to the cell (12, 13).

In *Synechococcus*, the KaiA protein functions as a homodimer and harbors two distinct domains
connected by a linker sequence (14–16). The N-terminal domain is similar to bacterial response
regulators but lacks the aspartate residue crucial for phosphorylation; hence, it is designated as a
pseudoreceiver domain (PsR domain) (14). This domain was shown to bind the oxidized form of
quinones and is therefore able to directly sense the onset of darkness and forward signals to the
C-terminal domain (14, 17). The C-terminus has a four-helix bundle secondary structure and is
highly conserved within Cyanobacteria. The domain harbors the KaiA dimer interface and the KaiC
binding site and is necessary to stimulate the autophosphorylation activity of KaiC (14, 16).
Mutations of *kaiA*, resulting in altered periodicity, were mapped throughout both domains, showing
their importance for rhythmicity (16, 18). For a more detailed review on the KaiABC oscillator and
its regulatory network, see Cohen and Golden (19), Swan *et al.* (20) and Snijder and Axmann (21).
The circadian timing mechanism of *Synechococcus* is well understood; however, the composition
of Kai systems can vary largely in other cyanobacteria and different clock sets, and their functions
are less elucidated (22–25). *Synechocystis* sp. PCC 6803 (hereafter *Synechocystis*) harbors, in
addition to the canonical *kaiAB1C1* gene cluster, two further *kaiB* and *kaiC* homologs (26). The
proteins KaiA, KaiB1, and KaiC1 seem to constitute the main timing mechanism, occupying the
canonical SasA-RpaA output-signaling pathway (27, 28). Deletion of either *kaiAB1C1* or *rpaA* led
to decreased viability in light-dark cycles compared to that of the wild-type (WT) strain (27, 29).
This phenotype was aggravated under photomixotrophic conditions and in complete darkness. In
contrast, the function of the additional Kai homologs has rarely been investigated, and multiple
copies of *kaiA* have not been identified thus far in any cyanobacterial strain (22).
For the *Synechocystis* KaiB3-KaiC3 timing system, Aoki and Onai suggested a function in the fine-
tuning of the core oscillator KaiAB1C1 by modulating its amplitude and period (30). This idea was
supported by Wiegard *et al.,* who recently investigated the characteristics of the KaiC3 protein and
proposed interplay between the KaiB3-KaiC3 system and the proteins of the standard clock system
(31). Phenotypical analysis hinted at similar cellular functions of both timing systems. While deletion
of *kaiC3* had no effect on growth in light-dark cycles, the cells displayed growth defects under
chemoheterotrophic conditions compared to that of the WT (29, 31). The impairment was less
severe in comparison with a Δ*kaiAB1C1*-deficient strain, which completely lost the ability to grow in
the dark. Furthermore, Wiegard *et al.* verified KaiA-independent autophosphorylation,
autodephosphorylation, and ATPase activity of *Synechocystis* KaiC3; enzymatic activities
suggested to be conserved in all cyanobacterial KaiC proteins (22, 31). However, compared to
Synechococcus KaiC, KaiC3 ATPase activity was found to be diminished and lacked temperature
compensation, an essential feature of true circadian oscillations (2, 31).
To date, the regulatory network of the KaiB3-KaiC3 system has remained enigmatic, as it does not
interact with KaiA and does not utilize the SasA-RpaA output pathway, suggesting alternative, not
yet identified, components for KaiB3-KaiC3-based signal transduction (27). In a large-scale protein-
protein interaction screen, a potential interaction partner of KaiC3 was identified (32). This protein,
Sll0485, was categorized as a NarL-type response regulator and could be a potential element in
the KaiB3-KaiC3 signaling pathway (33).
Within this study, we characterized Sll0485 bioinformatically, highlighting a resemblance of the
protein's N-terminal domain with the receiver domain of NarL-type response regulators, yet the
C-terminal domain shared similarities to KaiA homologs. Furthermore, we showed computationally
strong co-occurrences of the KaiB3-KaiC3 system with Sll0485 in the genomic context of
Cyanobacteria and other bacteria. We verified that the physical interaction of KaiC3 with Sll0485
increased the phosphorylation activity of KaiC3. Thus, we propose that Sll0485 is a novel KaiA-like
homolog linked to the KaiB3-KaiC3 system. Deletion of *sll0485* led to impaired viability during
nonphotoautotrophic growth, in line with previous studies on the KaiB3-KaiC3 system (31). Hence,
we introduced the name KaiA3 for Sll0485 and will use it from here onwards. Furthermore, to

facilitate the distinction of the KaiA homologs, we will use the name KaiA1 for the *Synechocystis*
KaiA core clock homolog Slr0756.
126
127
128
129   **Results**
130
KaiA3 is a chimeric protein harboring a NarL-type response regulator domain at the N-terminus and
a conserved KaiA-like motif at the C-terminus
The canonical clock genes *kaiABC* and *kaiA1B1C1* form a cluster in *Synechococcus* and
Synechocystis, respectively. In contrast, the *kaiB3* and *kaiC3* genes of *Synechocystis* are localized
in different regions of the chromosome (Fig. 1B, Fig. S1). Here, the *kaiB3* gene forms a
transcriptional unit with the open reading frame *sll0485,* encoding a hypothetical protein that we
named KaiA3. This hypothetical protein is annotated as a NarL-type response regulator (33) yet
displays a chimeric structure.
Using reciprocal BLAST analyses, we detected KaiA3 orthologs in 15 cyanobacterial species,
mainly belonging to the order *Chroococcales* ((34); Data S1), and in five bacterial genera outside
of Cyanobacteria, namely, *Roseiflexus*, *Chloroflexus*, *Chloroherpeton*, *Rhodospirillum*, and
Bradyrhizobium.
Analysis of the KaiA3 amino acid sequences from all cyanobacterial species encoding this protein
revealed that the N-terminus is conserved and indeed shows high homology to NarL-type response
regulators (Fig. S2). The canonical NarL protein consists of an N-terminal receiver domain, a linker
and a C-terminal DNA-binding domain with a helix-turn-helix motif (33, 35). For KaiA3, the
similarities to the NarL protein decreased in the C-terminus (Fig. S2), and concurrently,
conservation between KaiA3 and the KaiA protein family increased (Fig. 1A). The conserved
residues in the C-terminus correspond to structurally important features in the *Synechococcus* KaiA
protein, such as α-helical secondary structures, the KaiA dimer interface or residues critical for the
KaiA-KaiC interaction (Fig. 1A, 14, 15). Additionally, the lack of conservation in the N-terminus
compared to that seen in known KaiA orthologs is consistent with the results of Dvornyk and Mei,
which propose that different N-terminal domains exist for KaiA homologs for functional
diversification (36).
Multiple sequence alignments of the C-termini of the canonical KaiA and KaiA3 were performed to
construct a gene tree to further elucidate the relationship of KaiA3 and KaiA orthologs. The resulting
gene tree (Fig. 1A) distinctly separated KaiA3 from canonical KaiA orthologs. To further investigate
the evolutionary relationship of KaiA3, multiple sequence alignments of the C-terminus of orthologs
of KaiA3, KaiA, and Slr1783 (Rre1) as a reference for NarL orthologs in Cyanobacteria (37) were
used to construct a phylogenetic tree (Fig. S3). Here, KaiA3 orthologs form a distinct clade at the
basis of the KaiA orthologs when compared to both orthologous groups of Slr1783 (Rre1)/NarL (*E.*
coli, UniProtKB - P0AF28) and KaiA simultaneously (Fig. S3). In summary, these findings
strengthen the idea that the C-terminus of KaiA3 has a similar function to KaiA.
We further constructed three-dimensional models of KaiA3 to gain a better understanding of its
potential functions. To date, no structure is available for KaiA3, and it was impossible to generate
a reliable three-dimensional model covering the full-length KaiA3 sequence due to the enigmatic
structure of the linker region for which no significant similarities could be detected. Thus, we
modeled the N-terminus (residues 1-140) and the remaining part of the sequence separately
(residues 141-299). For the N-terminus, numerous hits for response regulator domains were
obtained, confirming previous findings. The 3D-model structures of KaiA3 are highly similar and
display the canonical fold of response regulator domains: a central five-stranded parallel β-sheet
flanked on both faces by five amphipathic α-helices and a phosphorylatable aspartate residue in
the β3-strand (Fig. S2, Fig. S4). This aspartate residue (D65), important for phosphorylation in
NarL-type response regulators (Fig. S2, blue stars), is conserved in all but *Pleurocapsa* and
Microcystis species. Thus, most KaiA3 homologs, including the *Synechocystis* protein, harbor a
potential phosphorylation site. Furthermore, the structure superimposes well on the PsR domain of
KaiA, even though the PsR domain lacks the aspartate residue that can be phosphorylated and the

α4-helix between the β4- and β5-strands (Fig. S4). The amino acid sequence between the β4- and
β5-strands shows the least conservation between KaiA and KaiA3, yet the level of sequence
conservation in this region is generally low for KaiA and its homologs (16). In contrast to the
N-terminal response regulator domain, the C-terminal domain of KaiA3 revealed a unique fold,
which has only been detected in KaiA thus far (38) and the N-terminal domain of the phosphoserine
phosphatase RsbU from *Bacillus subtilis* (39), namely, a unique four α-helix bundle constituting the
KaiA-like motif (Fig. S5). In conclusion, we propose that KaiA3 consists of two protein modules with
i) the N-terminal domain, resembling an NarL-type response regulator receiver domain, including
its phosphorylation site, and ii) the C-terminal domain displaying features of a KaiA-like motif. This
is particularly intriguing since putative *kaiA* orthologs outside of Cyanobacteria have not been
identified until recently (36).
189
Conserved synteny and co-occurrence of KaiA3 and the KaiB3-KaiC3 system among prokaryotes
As in *Synechocystis*, we found the *kaiA3* gene upstream of *kaiB3* in all analyzed cyanobacterial
genomes. Furthermore, the *kaiA3B3* cluster is usually extended by *kaiC3* with only two exceptions
(*Synechocystis* and *Microcystis aeruginosa* NIES-843), resembling the structure of the canonical
kaiABC gene cluster (Fig. 1B). Interestingly, *kaiA3B3C3* synteny is also found in other prokaryotic
genomes that harbor orthologs of *kaiA3*, except for *Chloroflexus aggregans* DMS 9485 (Fig. 1B).
Furthermore, we detected strong significant co-occurrences between KaiA3 and KaiB3 (p<0.0001),
as well as KaiA3 and KaiC3 (p<0.0001; Fig. 1C). The co-occurrence of KaiB3 and KaiC3 has
already been shown previously (22). Thus, KaiA3 forms a distinct set of proteins with KaiB3 and
KaiC3 that shows no further significant co-occurrence with other clock components (Fig. 1C, (22).
Altogether, both data sets suggest a functional relationship between KaiA3 and the KaiB3-KaiC3
system.
202
KaiC3 interacts with the hypothetical protein KaiA3
Using yeast two-hybrid experiments, we verified an interaction between the clock proteins KaiC3
and KaiA3 (Fig. 2A), consistent with the previous protein-protein interaction analysis by Sato *et al.*
(32). Although KaiA3 clearly interacts with KaiC3, an interaction with KaiB3, the second element of
the KaiB3-KaiC3 clock system, was not detected (Fig. S6). To further characterize the interaction
of the proteins *in vitro*, we expressed them recombinantly in *E. coli* and analyzed the complex
formation using native PAGE (Fig. 2B). KaiA3-His6 (monomer: 35 kDa) migrated as one single
band of approximately 100 kDa size, confirming the capability of KaiA3 to form homodimers.
Recombinant KaiB3 (monomer: 12 kDa) was recently shown to form dimers and tetramers of 13
and 41 kDa size, respectively, after size exclusion chromatography (31). KaiB3 displayed three
distinct bands in native gels. The two lower bands most likely represent the dimeric and tetrameric
forms, while the uppermost band (~ 67 kDa) is probably an impurity in the protein preparation.
Recombinant KaiC3 was produced with an N-terminal Strep-tag (31). Strep-KaiC3 (monomer:
58 kDa) migrated as one band at ~ 400 kDa, in line with the size of the hexamer (348 kDa).
Interestingly, this band vanishes completely when KaiC3 is incubated with KaiA3 (Fig. 2B). Instead,
protein accumulates in the well of the gel, indicating precipitation of the KaiA3/KaiC3 complex. On
the other hand, the formation of a large protein complex was observed when all three clock
components, KaiA3, KaiB3 and KaiC3, were incubated together for 16 h at 30 °C. The size matches
a complex consisting of one KaiC3 hexamer, six KaiA3 dimers and six KaiB3 monomers (840 kDa).
The presence of KaiA3 in the complex was validated by Western blotting using an anti-His antibody
(Fig. 2C). As expected, no such complex formed when KaiA3 was replaced by *Synechococcus*
KaiA (Fig. 2B). The interaction of KaiA3 and KaiC3 could, in addition, be validated using
immunoprecipitation-coupled liquid chromatography-mass spectrometry (LC-MS) analysis of
FLAG-tagged KaiC3 (Fig. S8). Supplementary experiments did not reveal any interaction between
KaiA3 and KaiC1 or KaiC2 (Fig. S6, Fig. S9), implying specificity for KaiC3 exists.
228
KaiA3 stimulates the autokinase activity of KaiC3
Previous studies showed that KaiC3 has autokinase activity and that this activity is independent of
KaiA (28, 31). Since our studies revealed an interaction of KaiC3 with KaiA3, we were interested in

probing the influence of KaiA3 on the phosphorylation of KaiC3. For this purpose, the above-
described recombinant Kai proteins were used. KaiC3 was incubated at 30 °C in the presence or
absence of other Kai proteins, and its phosphorylation state was analyzed by SDS-PAGE or
LC-MS/MS (Fig. 3). KaiC3 purified from *E. coli* was predominantly phosphorylated (Fig. 3A). During
the course of incubation, KaiC3 autodephosphorylates as is typical for KaiC proteins (39).
KaiC3-dephosphorylation is enhanced by KaiB3. In contrast, the addition of KaiA3 keeps KaiC3 in
the phosphorylated state, similar to the *Synechococcus* KaiC/KaiA core oscillator (6). The KaiA3-
mediated stimulation of KaiC3 phosphorylation was even more obvious when dephosphorylated
KaiC3 (NP-KaiC3) was used in the assay (Fig. 3B). Analysis of KaiC3 phosphorylation by
LC-MS/MS identified the neighboring residues Ser423 and Thr424 as phosphorylation sites, which
are conserved across KaiC homologs. However, only singly phosphorylated peptides could be
identified, with either Ser423 or Thr424 being phosphorylated. Overall, phosphorylation site
localization probabilities were higher for Ser423 (84 %) compared to Thr424 (57 %), defining
Ser423 as the primary phosphorylation site at present assay conditions (Fig. S10). In line with SDS-
PAGE analyses, consecutive phosphorylation occupancy analysis of the corresponding peptides
revealed low phosphorylation levels throughout most incubation conditions, except for enhanced
phosphorylation when KaiC3 was coincubated with KaiA3 for 16 h (Fig. 3C, Data S3). Interestingly,
an additional, low abundant phosphorylation event could be identified on Thr321, which however
remained at low occupancy (<1 %) at all incubation conditions.
As expected, *Synechococcus* KaiA did not influence the phosphorylation of KaiC3 (Fig. S11).
Based on these analyses, we conclude that KaiA3 likely has a KaiA-like function in promoting the
phosphorylation of KaiC3.
254
Deletion of kaiA3 impacts growth and viability during mixotrophic and chemoheterotrophic growth
As KaiA3 shows significant co-occurrence with KaiB3 and KaiC3, as well as *in vitro* interaction with
KaiC3, an interconnected function of the three proteins is more than likely. Deletion of *kaiC3* led to
growth impairments in complete darkness on glucose, but not in light-dark cycles (31); thus, the
kaiA3 knockout mutant (Δ*kaiA3*) was analyzed under these growth conditions. Cells were grown in
liquid culture under constant light, plated on agar in different dilutions and grown
photoautotrophically and photomixotrophically in continuous light and in 12-h light/12-h dark cycles
(Fig. 4A) or chemoheterotrophically (Fig. 4B). As the strains grow very slowly in chemoheterotrophic
conditions, cells were spotted at higher concentrations under these conditions. There were no
differences in viability of the mutant strains in comparison to the WT under photoautotrophic
conditions in continuous light and in light/dark cycles. Under photomixotrophic conditions, the
Δ*kaiA3* strain showed less viability, and viability was partly restored by re-insertion of *kaiA*. A mutant
strain, lacking all three alternative *kai* genes (Δ*kaiA3B3C3)* grew similar to the WT. However, in
light/dark cycles this strain seemed to have some advantage in comparison to the WT (Fig. 4A).
Spot assays under chemoheterotrophic conditions gave a clearer picture: the mutant strain lacking
kaiA3 and the triple knockout showed a similar phenotype. They were unable to grow in complete
darkness and in the complementation strain this ability was fully restored (Fig. 4B). These results
coincide with the previously detected impairments displayed by the Δ*kaiC3* strain during
chemoheterotrophic growth (31), strengthening the idea that the alternative *kai3* genes form a
regulatory complex with a common function.
275
276
277
278 **Discussion**
279
280 Despite increasing knowledge on circadian rhythms in Cyanobacteria and other prokaryotes,
281 information on the function, composition, and network of (cyano)bacterial nonstandard clock
282 systems remains scarce. By identifying KaiA3 and verifying its interaction with the KaiB3-KaiC3
283 complex, we could add another component to nonstandard clock systems.
284
Alternative KaiA orthologs outside of Cyanobacteria

Recently, new putative KaiA orthologs have been bioinformatically identified in prokaryotes outside
of Cyanobacteria (36). Even though multiple copies of the core clock proteins KaiB and KaiC have
been found to be encoded in cyanobacterial genomes, canonical KaiA was only found as a single
copy. In phylogenetic reconstructions, KaiA3 forms a distinct clade at the basis of the KaiA clade.
Together with the results of the structural analyses, interesting questions are raised about the
evolution of the KaiABC-based circadian clock system. It remains elusive whether KaiA3 together
with KaiB3 and KaiC3 represents an ancestral system for the well-studied *Synechococcus*
circadian clock, and further in-depth analyses are needed to elucidate the evolution of this new
putative timing system. Furthermore, together with KaiB3 and KaiC3, which have already been
found in other bacteria, including *Rhodospirillum rubrum,* where rhythmic processes have been
described (41), KaiA3 homologs could represent an overlooked component of Kai-based systems
outside of Cyanobacteria.
298
The two-domain architecture of KaiA3 and complex formation
KaiA3 displays a modular structure with an N-terminal response regulator domain connected via a
linker sequence to a C-terminal KaiA-like domain. The high structural similarity of the N-terminal
domain to numerous response regulator domains from various organisms indicates that the core
structure and activity are maintained, while adaptivity and variation from this provide specificity for
acting in distinct pathways (30). Within KaiA3, the aspartate residue crucial for phosphorylation is
conserved; thus, theoretically, the protein could receive an input signal from a cognate histidine
kinase, which has not been identified thus far.
The nearly unique structure of the KaiA C-terminus is, in addition to KaiA, solely found in the
N-terminal domain of the phosphatase RsbU *of Bacillus subtilis* (39). RsbU acts as a positive
regulator of the alternative sigma factor B, which is involved in the general stress response (42).
Interestingly, the N-terminal domain of RsbU forms dimers similar to KaiA, and the proposed
binding site for its corresponding activator RsbT is in an equivalent location to the KaiC-binding site
on KaiA (39). These findings may reflect how protein domains change during evolution while their
original functions are conserved. Compared to KaiA, in KaiA3, only five out of the 13 residues for
KaiC interaction are conserved (Fig. 1). However, KaiC3 also shows, in comparison to the KaiC1
homolog, less conservation of the residues important for the interaction with KaiA (28), perhaps
due to coevolution.
The KaiA3 structures obtained from models based on NarL and KaiA superimpose neatly on the
crystal structure of the KaiA dimer (Fig. S5, D), yet the composition of the linker subsequent to the
α5-helix (E141-D178) remains cryptic. In KaiA, the canonical linker, represented by residues
G165-K173, carries a β-sheet (β_6) crucial for the KaiA-KaiB interaction (43, 44). Native PAGE
analysis indicates that a complex forms when all three KaiA3, KaiB3 and KaiC3 clock components
are present but there is no complex between KaiA3 and KaiB3 (Fig. 2B). However, *Synechococcus*
CikA also lacks such a β-sheet but is able to form a complex with KaiB via its PsR domain (44).
Our analyses clearly imply that KaiA3 binds KaiC3 (Fig. 2, Fig. S8), which induces KaiC3
phosphorylation (Fig. 3). Therefore, the proteins form a stable complex only together with KaiB3,
otherwise aggregation occurs (Fig. 2B). Taken together, the data are consistent with a model in
which KaiA3 is able to fulfill the functions of a KaiA homolog, such as dimerization, by binding to
the KaiC homolog and enhancing autophosphorylation. Other mechanistic processes, such as
sequestration to the CI ring by binding to a KaiB homolog, remain to be investigated but are clearly
possible.
331
The function of KaiA3 within the cell
The physiological function of KaiA3 seems to be related to the different metabolic modes of
Synechocystis. This cyanobacterial strain is able to use glucose as an energy and carbon source
in the light as well as in the dark. It should be noted that the *Synechocystis* WT strain used in this
study was able to grow in complete darkness when supplemented with glucose. This is different
from an earlier study that showed that *Synechocystis* needs a 5 min blue-light pulse at least once
a day to be able to grow heterotrophically in the dark (45). The authors described this kind of
behavior as light-activated heterotrophic growth. There are no studies that explain why cells need

this 5 min light pulse, but it is also clear that our laboratory strain grows fully chemoheterotrophically
(29). Mutants deficient in *kaiA3* lose the ability to grow chemoheterotrophically on glucose, which
is an aggravated effect, compared to *kaiC3*-deficient mutants, which merely showed reduced
growth rates during heterotrophy (31). Similarly, in *Synechococcus,* the disruption of *kaiA* led to
one of the most severe mutations regarding activity loss and was traced back to an unbalanced
output signaling of the circadian clock (46). Related phenotypes, such as impaired viability during
light-dark cycles, including the complete loss of chemoheterotrophic growth on glucose, were
observed for the *Synechocystis* mutants Δ*kaiA1B1C1*, Δ*sasA*, and Δ*rpaA* (29, 47). For Δ*sasA,* it
was shown that the mutant strain was able to accumulate glycogen but was incapable of utilizing
the storage compound to grow heterotrophically, probably due to the inability to catabolize glucose
(47). A recent metabolomics study suggested that the growth inhibition of the Δ*kaiA1B1C1* and
Δ*rpaA* mutants in a light-dark cycle might be at least partly related to a defect in the inhibition of the
RuBisCo enzyme in the dark and increased photorespiration leading to the accumulation of the
potentially toxic product 2-phosphoglycolate (48). This previous study also revealed an enhanced
growth defect of the Δ*kaiA1B1C1* and Δ*rpaA* mutants under photomixotrophic conditions in
light/dark cycles, similar to the Δ*kaiA3* strain in the current study. This further supports the idea that
one of the functions of the KaiA3B3C3 system is to fine-tune the core clock system KaiA1B1C1.
Based on the conclusions from our studies, we propose that KaiA3 is a novel, nonstandard KaiA
homolog and element of the KaiC3-based signaling pathway. KaiA3 most likely adopts the function
of a KaiA homolog. The N-terminal half of KaiA3 might still have a response regulator function, yet
the exact mechanism remains unclear. Among other actions, KaiA3 needs to be placed within the
regulatory and metabolic network.
362
363
364
365 **Materials and Methods**
366
367 *Reciprocal BLAST of Sll0485 (KaiA3) and Slr1783 (Rre1)*
368 Reciprocal BLAST was performed as described in Schmelling *et al.*, 2017 (22). For comparison
369 with existing data about other circadian clock proteins, the 2017 database was used. The protein
370 sequences of Sll0485 (KaiA3) and Slr1783 (Rre1), as a reference for NarL response regulators
371 (37) from *Synechocystis* were used as query sequences for this reciprocal BLAST.
372
373 *Synteny analyses using SyntTax*
374 The conservation of gene order was analyzed with the web tool 'SyntTax' (53;
375 *https://archaea.i2bc.paris-saclay.fr/synttax/*). If not mentioned otherwise default settings (Best
376 match, 10 % norm. BLAST) were applied. Chromosomes were selected manually according to the
377 results of Schmelling *et al.* (2017).
378
Multiple sequence alignments with Mafft and Jalview
Sequence alignments, visualization, and analysis were performed with 'Jalview' (50). The
sequences were aligned with Mafft, and if not mentioned otherwise, default settings (L-INS-i,
pairwise alignment computation method -- localpair using Smith-Waterman algorithm, gap opening
penalty: 1.53, gap opening penalty at local pairwise alignment: -2.00, group-to-group gap extension
penalty: 0.123, matrix: BLOSUM62) were applied (51). For the analyses of the C-terminus, the
alignments were trimmed to position 168 in the KaiA reference sequence of *Synechococcus*. After
trimming, the alignment was recalculated with Mafft using the abovementioned default parameters.
387
2D and 3D structure predictions
The alignments generated in Jalview were then used with 'Ali2D' for secondary structure prediction
(56; https://toolkit.tuebingen.mpg.de). The identity cutoff to invoke a new PSIPRED run was set to
30%. Three-dimensional protein structures were modeled with either Phyre2 or SWISS-MODEL
(58, 59; http://www.sbg.bio.ic.ac.uk/phyre2/html/page.cgi?id=index;

https://swissmodel.expasy.org/). The resulting structures were then analyzed and illustrated using
UCSF Chimera (60; https://www.cgl.ucsf.edu/chimera/)
395
Phylogenetic reconstruction of protein trees
Phylogenetic reconstruction of the protein trees of Sll0485 (KaiA3), Slr1783 (Rre1)/NarL (*E. coli*,
UniProtKB - P0AF28), and KaiA was achieved with MEGA X (56, 57) using the above constructed
alignments. For all alignments, a neighbor-joining tree and a maximum likelihood tree were
constructed and compared. For the construction of the neighbor-joining trees, 1000 bootstrap
iterations with a p-distance substitution model and a gamma distribution with three gamma
parameters were used. For the construction of the maximum likelihood trees, an initial tree was
constructed using the maximum parsimony algorithm. Further trees were constructed using 1000
bootstrap iterations with an LG-G substitution model, a gamma distribution with three gamma
parameters and nearest-neighbor-interchange (NNI) as the heuristic method.
406
Yeast two-hybrid assay
AH109 yeast cells (Clontech) were used for the yeast two-hybrid experiments. Transformation of
yeast cells was performed according to the manufacturer's guidelines using the Frozen-EZ Yeast
Transformation kit (Zymo Research). The construction of plasmids used for transformation is
described in the Supplementary Information. Successfully transformed cells were selected on
complete supplement mixture (CSM) lacking leucine and a tryptophan (-Leu -Trp) dropout medium
(MP Biochemicals) at 30 °C for 3 - 4 days. Cells containing bait and prey plasmids were streaked
on CSM lacking leucine, tryptophan and histidine (-Leu -Trp -His) dropout medium (MP
Biochemicals) with the addition of 12.5 mM 3-amino-1,2,4-triazole (3-AT, Roth) and incubated for
6 days at 30 °C to screen for interaction.
417
Expression and purification of recombinant Kai proteins
Synechocystis KaiB3 and *Synechococcus* KaiA (plasmids kindly provided by T. Kondo, Nagoya
University, Japan) were produced as GST-fusion proteins in *E. coli* BL21(DE3) as described (30;
https://www.protocols.io/view/expression-and-purification-of-gst-tagged-kai-prot-48ggztw). Briefly,
proteins were purified with affinity chromatography using glutathione-agarose 4B (Macherey and
Nagel), and the N-terminal GST-tag was removed by PreScission Protease (Cytiva) prior to elution
of the untagged proteins from the glutathione resin. *Synechocystis* KaiC3 was produced with an
N-terminal Strep-tag (Strep-KaiC3) in *E. coli* Rosetta gami B (DE3) cells and purified via affinity
chromatography   using   Strep-tactin   XT   superflow   (IBA-Lifesciences)   (30;
https://www.protocols.io/view/heterologous-expression-and-affinity-purification-meac3ae).
Synechocystis ORF *sll0485,* encoding KaiA3, was inserted into vector pET22b to create a
C-terminal His6-fusion. KaiA3-His6 was expressed in *E. coli* Tuner(DE3) cells and purified with
immobilized metal affinity chromatography (IMAC) using PureProteome™ Nickel Magnetic Beads
(Millipore). For a detailed protocol see at protocols.io (dx.doi.org/10.17504/protocols.io.bu5bny2n).
Recombinant proteins were stored at -80 °C in buffer containing 20 mM Tris, pH 8.0, 150 mM NaCl,
0.5 mM EDTA, 5 mM MgCl$_2$, and 1 mM ATP.
434
KaiC3 phosphorylation assay
Recombinant Strep-KaiC3 purified from *E. coli* mainly exists in its phosphorylated form (KaiC3-P).
Fully dephosphorylated Strep-KaiC3 (KaiC3-NP) was generated by incubating the protein for 18 h
at 30 °C in assay buffer (20 mM Tris, pH 8.0, 150 mM NaCl, 0.5 mM EDTA, 5 mM MgCl2, and 1
mM ATP). The autokinase activity of KaiC3-NP was investigated by incubating 4 µg KaiC3 for 16
h at 30 °C in 20 µl assay buffer in the presence or absence of 2 µg KaiA3-His6, KaiB3 and
Synechococcus KaiA. Ten-microliter aliquots were taken before and after incubation at 30 °C, and
the reaction was stopped with SDS sample buffer. Samples were applied to a LowC SDS gel (10%
T, 0.67% C) (63; https://www.protocols.io/view/lowc-sds-page-gysbxwe) and stained with
Coomassie Blue R.
For analysis of KaiC3 phosphorylation occupancy levels in *in vitro* coincubation by LC-MS/MS,
proteins were digested in solution by chymotrypsin (Fig. 3C) or trypsin (Fig. S10), and measured

on a Q Exactive HF mass spectrometer (ThermoFisher Scientific) as described before (59). Raw
data was processed as described in supplementary methods with the following exceptions:
MaxQuant version 1.6.8.0 was used and phosphorylation of Ser, Thr and Tyr was defined as a
variable modification. Phosphorylation occupancy levels for each condition were calculated based
on intensity sums of modified/(modified + unmodified) peptides containing phosphorylation sites
Ser423 or Thr424 (Data S3)
453
Native protein PAGE and immunodetection of KaiA3
Kai proteins (10 µl samples containing 2 µg dephosphorylated Strep-KaiC3, 1 µg KaiA3-His6, 1 µg
KaiB3, 1 µg *Synechococcus* KaiA) were incubated for 16 h at 30 °C in phosphorylation assay buffer,
followed by separation of the native proteins in 4-16% native PAGE at 4 °C using a clear native
buffer system (Serva). Protein bands were visualized with Coomassie staining (ROTIBlue Quick,
Carl Roth) or immunodetected with a monoclonal anti-His antibody conjugated to HRP (MA1-
21315-HRP, Thermo Fisher, 1:2000 diluted). A detailed protocol is found at protocols.io
(dx.doi.org/10.17504/protocols.io.bu67nzhn).
462
Strains and growth conditions
Wild type *Synechocystis* (PCC-M, resequenced, (60)), the deletion strains Δ*rpaA* (27), Δ*kaiC3* (31),
Δ*kaiA3* and Δ*kaiA3B3C3* (SI, Fig. S1) and the complementation strain Δ*kaiA3*/*kaiA3* (SI, Fig. S1)
were cultured photoautotrophically in BG11 medium (61) supplemented with 10 mM TES buffer
(pH 8) under constant illumination with 75 µmol photons $m^{-2} s^{-1}$ of white light (Philips TLD Super
80/840) at 30 °C. Cells were grown either in Erlenmeyer flasks with constant shaking (140 rpm) or
on plates (0.75 % Bacto-Agar, Difco) supplemented with 0.3 % thiosulfate. For photomixotrophic
experiments, 0.2 % glucose was added to the plates. For chemoheterotrophic growth experiments
in complete darkness, *Synechocystis* cells were spotted in different dilutions on BG11 agar plates
containing 0.2 % glucose and incubated either mixotrophically for 3 days with continuous
illumination or chemoheterotrophically in the dark for 26 days.
474
475
476 **Data availability**
477

484
485
486 **Acknowledgments**
487

We thank Research Unit FOR2816 'SCyCode' (The Autotrophy-Heterotrophy Switch in
Cyanobacteria: Coherent Decision-Making at Multiple Regulatory Layers) funded by the German
Research Foundation for fruitful discussions and financial support. This work was financially
supported by grants (WI2014/5-3; 10-1; AX 84/1-3 and MA 4918/4-1) from the German Research
Foundation to A.W., I.M.A. and B.M. respectively. We thank Pauline Morys, Annika Klopp, and
Werner Bigott for technical assistance.

494
495
496 **References**
497

498 1.   M. Nakajima, *et al.*, Reconstitution of circadian oscillation of cyanobacterial KaiC
499   phosphorylation in vitro. *Science.* **308**, 414–415 (2005).
500 2.   K. Terauchi, *et al.*, ATPase activity of KaiC determines the basic timing for circadian clock

501  of cyanobacteria. *Proc. Natl. Acad. Sci. U. S. A.* **104**, 16377–16381 (2007).

502 3. M. Egli, *et al.*, Dephosphorylation of the Core Clock Protein KaiC in the Cyanobacterial
503  KaiABC Circadian Oscillator Proceeds via an ATP Synthase Mechanism. *Biochemistry* **51**,
504  1547–1558 (2012).

505 4. R. Pattanayek, *et al.*, Visualizing a Circadian Clock Protein: Crystal Structure of KaiC and
506  Functional Insights. *Mol. Cell* **15**, 375–388 (2004).

507 5. Y. I. Kim, G. Dong, C. W. Carruthers, S. S. Golden, A. LiWang, The day/night switch in KaiC,
508  a central oscillator component of the circadian clock of cyanobacteria. *Proc. Natl. Acad. Sci.*
509  *U. S. A.* **105**, 12825–12830 (2008).

510 6. H. Iwasaki, T. Nishiwaki, Y. Kitayama, M. Nakajima, T. Kondo, KaiA-stimulated KaiC
511  phosphorylation in circadian timing loops in cyanobacteria. *Proc. Natl. Acad. Sci. U. S. A.*
512  **99**, 15788–93 (2002).

513 7. R. Pattanayek, M. Egli, Protein–Protein Interactions in the Cyanobacterial Circadian Clock:
514  Structure of KaiA Dimer in Complex with C-Terminal KaiC Peptides at 2.8 Å Resolution.
515  *Biochemistry* **54**, 4575–4578 (2015).

516 8. Y.-G. Chang, R. Tseng, N.-W. Kuo, A. LiWang, Rhythmic ring-ring stacking drives the
517  circadian oscillator clockwise. *Proc. Natl. Acad. Sci. U. S. A.* **109**, 16847–51 (2012).

518 9. Y.-G. Chang, N.-W. Kuo, R. Tseng, A. LiWang, Flexibility of the C-terminal, or CII, ring of
519  KaiC governs the rhythm of the circadian clock of cyanobacteria. *Proc. Natl. Acad. Sci. U.*
520  *S. A.* **108**, 14431–6 (2011).

521 10. T. Nishiwaki, *et al.*, Role of KaiC phosphorylation in the circadian clock system of
522  *Synechococcus elongatus* PCC 7942. *Proc. Natl. Acad. Sci. U. S. A.* **101**, 13927–13932
523  (2004).

524 11. Y. Kitayama, H. Iwasaki, T. Nishiwaki, T. Kondo, KaiB functions as an attenuator of KaiC
525  phosphorylation in the cyanobacterial circadian clock system. *EMBO J.* **22**, 2127–2134
526  (2003).

527 12. R. Tseng, *et al.*, Cooperative KaiA–KaiB–KaiC Interactions Affect KaiB/SasA Competition
528  in the Circadian Clock of Cyanobacteria. *J. Mol. Biol.* **426**, 389–402 (2014).

529 13. M. J. Rust, J. S. Markson, W. S. Lane, D. S. Fisher, E. K. O'Shea, Ordered phosphorylation
530  governs oscillation of a three-protein circadian clock. *Science* **318**, 809–12 (2007).

531 14. S. B. Williams, I. Vakonakis, S. S. Golden, A. C. LiWang, Structure and function from the
532  circadian clock protein KaiA of *Synechococcus elongatus*: a potential clock input
533  mechanism. *Proc. Natl. Acad. Sci. U. S. A.* **99**, 15357–62 (2002).

534 15. I. Vakonakis, *et al.*, NMR structure of the KaiC-interacting C-terminal domain of KaiA, a
535  circadian clock protein: implications for KaiA-KaiC interaction. *Proc. Natl. Acad. Sci. U. S.*
536  *A.* **101**, 1479–84 (2004).

537 16. S. Ye, I. Vakonakis, T. R. Ioerger, A. C. LiWang, J. C. Sacchettini, Crystal structure of
538  circadian clock protein KaiA from *Synechococcus elongatus*. *J. Biol. Chem.* **279**, 20511–8
539  (2004).

540 17. Y. I. Kim, D. J. Vinyard, G. M. Ananyev, G. C. Dismukes, S. S. Golden, Oxidized quinones
541  signal onset of darkness directly to the cyanobacterial circadian oscillator. *Proc. Natl. Acad.*
542  *Sci. U. S. A.* **109**, 17765–17769 (2012).

543 18. H. Nishimura, *et al.*, Mutations in KaiA, a clock protein, extend the period of circadian rhythm
544  in the cyanobacterium *Synechococcus elongatus* PCC 7942. *Microbiology* **148**, 2903–2909
545  (2002).

546 19. S. E. Cohen, S. S. Golden, Circadian Rhythms in Cyanobacteria. *Microbiol. Mol. Biol. Rev.*
547  **79**, 373–385 (2015).

548 20. J. A. Swan, S. S. Golden, A. LiWang, C. L. Partch, Structure, function, and mechanism of
549  the core circadian clock in cyanobacteria. *J. Biol. Chem.* **293**, 5026–5034 (2018).

550 21. J. Snijder, I. M. Axmann, "The Kai-Protein Clock—Keeping Track of Cyanobacteria's Daily
551  Life" in *Subcellular Biochemistry*, (Springer, 2019), pp. 359–391.

552 22. N. M. Schmelling, *et al.*, Minimal tool set for a prokaryotic circadian clock. *BMC Evol. Biol.*
553  **17**, 169 (2017).

554 23. V. Dvornyk, O. Vinogradova, E. Nevo, Origin and evolution of circadian clock genes in

prokaryotes. *Proc Natl Acad Sci U S A* **100**, 2495–2500 (2003).

24.   I. M. Axmann, S. Hertel, A. Wiegard, A. K. Dörrich, A. Wilde, Diversity of KaiC-based timing
systems in marine Cyanobacteria. *Mar. Genomics* **14**, 3–16 (2014).

25.   V. Dvornyk, "The circadian clock gear in cyanobacteria: Assembled by evolution" in *Bacterial
Circadian Programs*, (Springer-Verlag Berlin Heidelberg, 2009), pp. 241–258.

26.   Y. Kanesaki, *et al.*, Identification of Substrain-Specific Mutations by Massively Parallel
Whole-Genome Resequencing of *Synechocystis* sp. PCC 6803. *DNA Res.* **19**, 67–79
(2012).

27.   C. Köbler, S. J. Schultz, D. Kopp, K. Voigt, A. Wilde, The role of the *Synechocystis* sp. PCC
6803 homolog of the circadian clock output regulator RpaA in day–night transitions. *Mol.*
Microbiol. **110**, 847–861 (2018).

28.   A. Wiegard, *et al.*, Biochemical analysis of three putative KaiC clock proteins from
Synechocystis sp. PCC 6803 suggests their functional divergence. *Microbiology* **159**, 948–
958 (2013).

29.   A. K. Dörrich, J. Mitschke, O. Siadat, A. Wilde, Deletion of the *Synechocystis* sp. PCC 6803
kaiAB1C1 gene cluster causes impaired cell growth under light-dark conditions.
Microbiology **160**, 2538–2550 (2014).

30.   S. Aoki, K. Onai, "Circadian clocks of *Synechocystis* sp. strain PCC 6803,
Thermosynechococcus elongatus, *Prochlorococcus* spp., *Trichodesmium* spp. and other
species" in *Bacterial Circadian Programs*, (Springer-Verlag Berlin Heidelberg, 2009), pp.
259–282.

31.   A. Wiegard, *et al.*, *Synechocystis* KaiC3 displays temperature- And KaiB-dependent
ATPase activity and is important for growth in darkness. *J. Bacteriol.* **202**, 1–36 (2020).

32.   S. S. Sato, *et al.*, A large-scale protein protein interaction analysis in *Synechocystis* sp. PCC
6803. *DNA Res.* **14**, 207–16 (2007).

33.   I. Baikalov, *et al.*, Structure of the *Escherichia coli* response regulator NarL. *Biochemistry*
35, 11053–11061 (1996).

34.   J. Komarek, J. Kaštovský, J. Mares, J. Johansen, Taxonomic classification of
cyanoprokaryotes (cyanobacterial genera) 2014, using a polyphasic approach. *Preslia -*
Praha- **86**, 295–335 (2014).

35.   M. Y. Galperin, Structural classification of bacterial response regulators: Diversity of output
domains and domain combinations. *J. Bacteriol.* **188**, 4169–4182 (2006).

36.   V. Dvornyk, Q. Mei, Evolution of *kaiA*, a key circadian gene of cyanobacteria. *Sci. Rep.* **11**,
9995 (2021).

37.   M. K. Ashby, J. Houmard, Cyanobacterial Two-Component Proteins: Structure,Diversity,
Distribution, and Evolution. *Microbiol. Mol. Biol. Rev.* **70**, 472–509 (2006).

38.   I. Vakonakis, A. C. LiWang, Structure of the C-terminal domain of the clock protein KaiA in
complex with a KaiC-derived peptide: implications for KaiC regulation. *Proc. Natl. Acad. Sci.*
U. S. A. **101**, 10925–30 (2004).

39.   O. Delumeau, *et al.*, Functional and structural characterization of RsbU, a stress signaling
protein phosphatase 2C. *J. Biol. Chem.* **279**, 40927–40937 (2004).

40.   T. Nishiwaki, T. Kondo, Circadian autodephosphorylation of cyanobacterial clock protein
KaiC occurs via formation of ATP as intermediate. *J. Biol. Chem.* **287**, 18030–18035 (2012).

41.   E. Van Praag, R. Degli Agosti, R. Bachofen, Rhythmic Activity of Uptake Hydrogenase in
the Prokaryote *Rhodospirillum rubrum*. *J. Biol. Rhythms* **15**, 218–224 (2000).

42.   X. Yang, C. M. Kang, M. S. Brody, C. W. Price, Opposing pairs of serine protein kinases
and phosphatases transmit signals of environmental stress to activate a bacterial
transcription factor. *Genes Dev.* **10**, 2265–2275 (1996).

43.   J. Snijder, *et al.*, Structures of the cyanobacterial circadian oscillator frozen in a fully
assembled state. *Science.* **355**, 1181–1184 (2017).

44.   R. Tseng, *et al.*, Structural basis of the day-night transition in a bacterial circadian clock.
Science. **355**, 1174–1180 (2017).

45.   S. L. Anderson, L. McIntosh, Light-activated heterotrophic growth of the cyanobacterium
Synechocystis sp. strain PCC 6803: A blue-light-requiring process. *J. Bacteriol.* **173**, 2761–

2767 (1991).

46.   D. G. Welkie, *et al.*, Genome-wide fitness assessment during diurnal growth reveals an
expanded role of the cyanobacterial circadian clock protein KaiA. *Proc. Natl. Acad. Sci. U.*
S. A. **115**, E7174–E7183 (2018).

47.   A. K. Singh, L. A. Sherman, Pleiotropic effect of a histidine kinase on carbohydrate
metabolism in *Synechocystis* sp. strain PCC 6803 and its requirement for heterotrophic
growth. *J. Bacteriol.* **187**, 2368–76 (2005).

48.   N. M. Scheurer, *et al.*, Homologs of Circadian Clock Proteins Impact the Metabolic Switch
Between Light and Dark Growth in the Cyanobacterium Synechocystis sp. PCC 6803. *Front.*
Plant Sci. **12**, 675227 (2021).

49.   J. Oberto, SyntTax: A web server linking synteny to prokaryotic taxonomy. *BMC*
Bioinformatics **14** (2013).

50.   A. M. Waterhouse, J. B. Procter, D. M. A. Martin, M. Clamp, G. J. Barton, Jalview Version
2-A multiple sequence alignment editor and analysis workbench. *Bioinformatics* **25**, 1189–
1191 (2009).

51.   K. Katoh, D. M. Standley, MAFFT multiple sequence alignment software version 7:
Improvements in performance and usability. *Mol. Biol. Evol.* **30**, 772–780 (2013).

52.   L. Zimmermann, *et al.*, A Completely Reimplemented MPI Bioinformatics Toolkit with a New
HHpred Server at its Core. *J. Mol. Biol.* **430**, 227–2243 (2018).

53.   A. Waterhouse, *et al.*, SWISS-MODEL: homology modelling of protein structures and
complexes. *Nucleic Acids Res.* **46** (2018).

54.   L. A. Kelley, S. Mezulis, C. M. Yates, M. N. Wass, M. J. E. Sternberg, The Phyre2 web portal
for protein modeling, prediction and analysis. *Nat. Protoc.* **10**, 845–858 (2015).

55.   E. F. Pettersen, *et al.*, UCSF Chimera - A visualization system for exploratory research and
analysis. *J. Comput. Chem.* **25**, 1605–1612 (2004).

56.   S. Kumar, G. Stecher, M. Li, C. Knyaz, K. Tamura, MEGA X: Molecular evolutionary
genetics analysis across computing platforms. *Mol. Biol. Evol.* **35**, 1547–1549 (2018).

57.   G. Stecher, K. Tamura, S. Kumar, Molecular Evolutionary Genetics Analysis (MEGA) for
macOS. *Mol. Biol. Evol.* **37**, 1237–1239 (2020).

58.   T. Nishiwaki, *et al.*, A sequential program of dual phosphorylation of KaiC as a basis for
circadian rhythm in cyanobacteria. *EMBO J.* **26**, 4029–4037 (2007).

59.   P. Spät, T. Barske, B. Maček, M. Hagemann, Alterations in the CO_2 availability induce
alterations in the phosphoproteome of the cyanobacterium *Synechocystis* sp. PCC 6803.
New Phytol. **231**, 1123–1137 (2021).

60.   D. Trautmann, B. Voß, A. Wilde, S. Al-Babili, W. R. Hess, Microevolution in cyanobacteria:
Re-sequencing a motile substrain of *Synechocystis* sp. PCC 6803. *DNA Res.* **19**, 435–448
(2012).

61.   R. Rippka, J. Deruelles, M. Herdman, J. B. Waterbury, R. Y. Stanier, Generic Assignments,
Strain Histories and Properties of Pure Cultures of Cyanobacteria. *Microbiology* **111**, 1–61
(1979).

62.   Y. Perez-Riverol, *et al.*, The PRIDE database and related tools and resources in 2019:
improving support for quantification data. *Nucleic Acids Res.* **47**, D442–D450 (2019).

Figures and Tables

Fig. 1. Bioinformatic analyses of KaiA3. (A) Multiple sequence alignment and maximum likelihood-inferred phylogenetic reconstruction of KaiA3 and selected KaiA orthologs. The sequences were aligned with Mafft (L-INS-i default parameters, Jalview), trimmed to position 168 of the C-terminus of *Synechococcus* KaiA and are represented in the Clustalx color code with conservation visibility set to 25%. Marks above the alignment refer to *Synechococcus* KaiA as a reference. Light green bars and dots indicate residues critical for KaiC interaction, light pink bars and dots represent residues important for dimerization, and light gray blocks outline residues forming α-helices as secondary structures. Aligned sequences were used to infer a maximum likelihood protein tree. The scale bar indicates 1 substitution per position. Bootstrap values (n=1000) are displayed at branches. Bootstrap values less than 50 are not shown. (B) Synteny analysis of KaiABC compared to KaiA3, KaiB3, and KaiC3 for selected bacterial species. Analysis was performed with the online tool SyntTax, a prokaryotic synteny and taxonomy explorer (https://archaea.i2bc.paris-saclay.fr/synttax/; 2020-06-08). Default settings were used for analysis (best match, 10% norm. Blast). (C) Co-occurrence of KaiA3 using pairwise Fisher's exact tests with circadian clock proteins in Cyanobacteria. Network of significant co-occurring circadian clock factors from Schmelling *et al.* (22) including KaiA3 in Cyanobacteria. The line color corresponds to the level of significance resulting from the pairwise Fisher's exact test. Missing links are those that had a p-value higher than 0.01. Node size is proportional to the degree of that node.

676
677
Fig. 2. Analysis of KaiA3 protein interactions. (A) Yeast two-hybrid interaction analysis of KaiA3
with the KaiC3 protein. As a positive control, KaiA1 dimer interaction was used. Yeast two-hybrid
reporter strains carrying the respective bait, and prey plasmids were selected by plating on
complete supplement medium (CSM) lacking leucine and tryptophan (-Leu -Trp). AD, GAL4
activation domain; BD, GAL4 DNA-binding domain; empty, bait and prey plasmids without protein
sequence (only AD/BD domain). The physical interaction between bait and prey fusion proteins
was determined by growth on complete medium lacking leucine, tryptophan and histidine
(-Leu -Trp -His) and the addition of 12.5 mM 3-amino-1,2,4-triazole (3-AT). BD was fused to the
N-terminus of KaiA3. For clear presentation, spots were assembled from several assays (original
scans are shown in Fig. S7). (B), (C) Interaction analysis of recombinant Kai proteins in native
polyacrylamide gels. Proteins were incubated for 16 h at 30 °C and subsequently subjected to
4-16% clear native PAGE. Gels were either stained with Coomassie Blue (B) or blotted and
immunodecorated with a monoclonal anti-His antibody for the detection of recombinant KaiA3-His6
(C). Recombinant *Synechococcus* KaiA was used for comparison.

692
693

694
695
Fig. 3. KaiA3-stimulated phosphorylation of KaiC3. KaiC3 was incubated at 30 °C with or without
KaiA3 and KaiB3. Aliquots were taken at 0 h and 16 h followed by separation of phosphorylated
(P) and nonphosphorylated (NP) KaiC3 on SDS-PAGE (10% T, 0.67% C). (A) Phosphorylated
KaiC3 was used as the starting material for the assay. (B) Nonphosphorylated KaiC3 generated by
an 18 h incubation at 30 °C prior to the start point was used as the material for the assay. (C)
Analysis of KaiC3 phosphorylation levels by LC-MS/MS. Samples were prepared as in (B) and
digested in solution with chymotrypsin. Phosphorylation occupancy was calculated for each sample
(in %) based on intensity sums of modified/(modified + unmodified) peptides containing
phosphorylation sites Ser423 or Thr424 (marked in bold; phosphopeptides exclusively singly
phosphorylated). Sequences and start/end positions of detected peptides are indicated. *detection
of nonmodified peptides only.
707
708

709
710
Fig. 4. Deletion of *kaiA3* results in growth defects during mixotrophic and chemoheterotrophic
growth. Proliferation of the WT, the *kaiA3* and *kaiA3B3C3* deletion mutant, and the Δ*kaiA3*/*kaiA3*
complementation strain was tested under phototrophic (continuous light, - glucose)
photomixotrophic (continuous light, + glucose) and heterotrophic (darkness, + glucose) conditions.
Strains were grown in liquid culture under constant light, different dilutions were spotted on agar
plates and incubated in the indicated light conditions with light phase corresponding to 75 µmol
photons m^{-2} s^{-1} white light. A representative result of three independent experiments is shown.
(A) Cultures were diluted to OD_{750nm} value 0.4 and dilution series were spotted on agar plates with
or without the addition of glucose. Plates were analyzed after 6 or 8 days of continuous light and
12h/12h light-dark cycles, respectively. (B) Cultures were diluted to OD_{750nm} values of 1.2, 0.8 and
0.4 and spotted on agar plates supplemented with 0.2% glucose. Plates were analyzed after 3 or
26 days of continuous light and darkness, respectively.
723
724
725