1

Discovery of genes that modulate flavivirus replication in an interferon-dependent manner

Sarah Lesage[1,2]*, Maxime Chazal[1]*, Guillaume Beauclair[1,3], Damien Batalie [4], Elodie Couderc[1,5], Aurianne Lescure[6], Elaine Del Nery[6], Frédéric Tangy[7], Annette Martin[4], Nicolas Manel[8#] and Nolwenn Jouvenet[1#]

* These authors contributed equally
# corresponding authors

[1] Institut Pasteur, Virus Sensing and Signaling Unit, UMR3569 CNRS, Paris, France
[2] Université de Paris, Paris, France
[3] Université Paris-Saclay, CEA, CNRS, Institute for Integrative Biology of the Cell (I2BC), Gif-sur-Yvette, France
[4] Institut Pasteur, Molecular Genetics of RNA Viruses Unit, UMR3569 CNRS, Université de Paris, Paris, France
[5] Institut Pasteur, Insect-Virus Interactions Unit, Department of Virology, UMR2000 CNRS, Paris, France
[6] Institut Curie, PSL Research University, Department of Translational Research-Biophenics High-Content Screening Laboratory, Cell and Tissue Imaging Facility (PICT-IBiSA), Paris, France
[7] Institut Pasteur, Viral Genomics and Vaccination Unit, UMR3569 CNRS, Paris, France
[8] Institut Curie, PSL Research University, INSERM U932, Paris, France.

Short title: Modulation of flavivirus replication by interferon effectors

Summary

Establishment of the interferon (IFN)-mediated antiviral state provides a crucial initial line of defense against viral infection. Numerous genes that contribute to this antiviral state remain to be identified. Using a loss-of-function strategy, we screened an original library of 1156 siRNAs targeting 386 individual curated human genes in stimulated microglial cells infected with Zika virus (ZIKV), an emerging RNA virus that belongs to the flavivirus genus. The screen recovered twenty-one potential host proteins that modulate ZIKV replication in an IFN-dependent manner, including the previously known IFITM3 and LY6E. Further characterization contributed to delineate the spectrum of action of these genes towards other pathogenic RNA viruses, including Hepatitis C virus and SARS-CoV-2. Our data revealed that APOL3 acts as a proviral factor for ZIKV and several other related and unrelated RNA viruses. In addition, we showed that MTA2, a chromatin remodeling factor, possesses potent flavivirus-specific antiviral functions. Our work identified previously unrecognized genes that modulate the replication of RNA viruses in an IFN-dependent way, opening new perspectives to target weakness points in the life cycle of these viruses.

Author summary

The interferon (IFN) response, which is a powerful arm of the antiviral immune program, protects cells from invading viral pathogens by transcriptionally inducing the expression of interferon-stimulated genes (ISGs). ISGs effectively establish the antiviral state by directly blocking the viral life cycle by targeting specific stages of replication. Some ISGs are specific to a virus or a viral family, while others are broad-spectrum. Identifying which ISGs exert antiviral function against which virus is crucial because it enables the subsequent understanding of weakness points in the life cycle of viruses. Using a loss-of-fonction screening strategy, we investigated the ability of 386 unique ISGs to inhibit the replication of Zika virus (ZIKV), a virus that recently emerged in the Pacific and the Americas. Using this approach, we identified novel modulators of ZIKV replication in IFN-treated human cells. Our work contributes to a better understanding of the molecular basis behind the complexity the antiviral program. Furthermore, taking advantage of naturally occurring viral inhibitors may be an effective strategy for the development of novel drugs that activate, amplify or mimic their action.

Introduction

Viruses are high on the list of global public health concerns, as illustrated by recent epidemics caused by Ebola, Zika (ZIKV) and Nipah viruses, as well as by the ongoing SARS-CoV-2 pandemic. The vast majority of these emerging RNA viruses have zoonotic origins and have recently crossed host species barrier [1]. In order to establish itself in a host species, one of the first and most restrictive barriers that a virus needs to overcome is the antiviral innate immune system. This response has evolved to rapidly control viral replication and limit virus spread *via* detection of viral nucleic acids by pathogen recognition receptors (PRRs) [2]. These PRRs can be membrane-associated, such as Toll-like receptor (TLRs), or cytosolic, such as retinoic acid inducible gene I (RIG-I)-like receptor (RLRs). Upon binding to viral nucleic acids, these PRRs interact with adaptor proteins and recruit signaling complexes. These events lead to the expression of type I interferons (IFNs). Secreted IFNs will then bind to their heterodimeric receptor (IFNAR1/IFNAR2) and activate the canonical JAK/STAT pathway in infected and surrounding cells. This activation triggers the assembly of the interferon-stimulated gene 3 (ISGF3) complex (composed of STAT1, STAT2 and IRF-9 proteins), which subsequently induces the expression of up to approximately 2000 IFN-stimulated genes (ISGs) [3,4], effectively establishing the antiviral state. ISGs comprise a core of genes that are induced at high levels essentially in all cell types, as well as cell-type specific genes that are the result of transcriptome remodeling [5,6], highlighting the importance of studying ISGs in relevant cell types. Some of these ISGs have been well characterized. They directly block the viral life cycle by targeting specific stages of virus replication, including entry into host cells, protein translation, replication or assembly of new viral particles [3,7]. Some ISGs are specific to a virus or a viral family, while others are broad-spectrum. They can also be negative or positive regulators of IFN signaling and thus facilitate, or not, the return to cellular homeostasis. However, the contribution of most ISGs to the antiviral state remains poorly understood.

Over the last decades, flaviviruses have provided some of the most important examples of emerging or resurging diseases, including ZIKV, dengue virus (DENV), Yellow fever virus (YFV) and West Nile virus (WNV) [8]. These flaviviruses are arthropod-borne viruses transmitted to vertebrate hosts by mosquitoes. They cause a spectrum of potentially severe diseases including hepatitis, vascular shock syndrome, encephalitis, acute flaccid paralysis, congenital abnormalities and fetal death [8]. They are now globally distributed and infect up to 400 million people annually. Lesser-known flaviviruses are beginning to emerge in different parts of the world, as illustrated by the recent incursion of Usutu virus (USUV) in the Mediterranean basin [9].

ZIKV was isolated in 1947 in a macaque from the Zika Forest in Uganda [10]. For decades, it remained in Africa and Asia where it sparked local epidemics characterized by a mild self-limiting disease in humans. In recent years, Asian lineage viruses have emerged as a global public health threat with widespread epidemics in the Pacific Islands and Americas, where over 35 countries have reported local transmission in 2016. An estimated 1 million individuals were affected by ZIKV in Brazil in 2015-16. Infection by ZIKV has been linked to several neurological disorders, including Guillain-Barré syndrome (GBS),

meningoencephalitis, myelitis and congenital microcephaly, fetal demise and abortion [10]. Children
exposed to ZIKV in utero may present neurocognitive deficits, regardless of head size at birth. ZIKV
infection is now identified as a sexually-transmitted illness as well [11]. As all flaviviruses, ZIKV is an
enveloped virus containing a positive-stranded RNA genome of ~11 kb. Upon viral entry, the viral
genome is released and translated by the host cell machinery into a large polyprotein precursor. The
latter is processed by host and viral proteases into three structural proteins, including C (core), prM
(precursor of the M protein) and E (envelope) glycoproteins, and seven non-structural proteins (NS)
called NS1, NS2A, NS2B, NS3, NS4A, NS4B and NS5 [8]. The structural proteins constitute the viral
particle, while NS proteins coordinate RNA replication, viral assembly and modulate innate immune
responses.

The importance of IFN signaling in mediating host restriction of ZIKV is illustrated by the severe
pathogenicity in IFNAR1-/- and STAT2-/- but not in immunocompetent mice [12–14]. Moreover, the
Zika strain that is responsible for the recent epidemics has accumulated mutations that increase
neurovirulence via the ability to evade the immune response [15]. Microglial cells, which are the resident
macrophages of the brain, represent ZIKV targets and potential reservoirs for viral persistence [16].
Moreover, they may play a role in ZIKV transmission from mother to fetal brain [17] and affect the
proliferation and differentiation of neuronal progenitor cells [18]. In order to comprehend the molecular
bases behind the efficacy of the IFN response to ZIKV replication, we set up a high throughput assay to
identify genes that are modulating viral replication in human microglial cells (HMC3) stimulated with
IFN.

Results

A loss of function screen identified genes modulating ZIKV replication in IFN-stimulated
human microglial cells

We first performed pilot experiments to determine the feasibility of conducting large-scale loss-of-
function studies to identify novel genes regulating ZIKV replication in stimulated HMC3 cells. Five
hundreds cells were seeded in 384-well microplates on day 1, transfected with individual siRNA 6 hours
later, treated with IFNα2 at day 2, infected with ZIKV at day 3 and fixed 24 hours later (Fig. 1A).
Percentages of infected cells were determined by confocal analysis by measuring the number of cells
expressing the viral E protein, using the pan-flavivirus anti-E antibody 4G2 (Fig. 1A). Nuclei were
identified with DAPI staining for imaging and segmentation purposes. We optimized IFNα2
concentration and viral multiplicity of infection (MOI) to obtain a significant decrease of E-positive
cells upon IFNα2-treatment (Fig. S1A). We used siRNAs targeting IFNAR1, which are expected to
neutralize IFN signaling, as positive controls (Fig. S1A). siRNAs against IFITM3, an ISG known to
potently inhibit ZIKV replication in several human cell lines and primary fibroblasts [19,20], were used
as additional positive controls (Fig. S1A). Negative controls were non-targeting siRNAs. As expected,

in the presence of siRNAs targeting IFNAR1, IFN signaling was neutralized and the level of infection
was almost rescued to the level of non-treated cells (Fig. S1A). In cells silenced for IFITM3 expression,
the number of infected cells was partly restored to the level of non-treated cells (Fig. S1A). Such partial
rescue was expected since the antiviral state requires the concerted action of numerous ISGs [21]. These
data also revealed that IFITM3 is a potent anti-ZIKV ISG in microglial cells.

We scaled up the experiment by screening an arrayed library containing 1158 siRNAs targeting 386
human genes (Table S1). These genes were selected based on a gene signature defined by clustering and
correlation of expression with MX1, a well-described ISG, in a dataset of gene expression in primary
human $CD4^+$ T cells (Cerboni et al., in preparation). 36% of the identified genes overlapped with
previous ISG libraries [21,22], ensuring that the screen would be simultaneously capable of identifying
expected positive hits and find new genes of interest. To overcome potential off-target effect and a
limited efficacy of transcript knockdown, each gene was targeted by 3 different siRNAs. Numerous
siRNAs targeting IFNAR1 and IFNAR2 were used as positive controls. Negative controls were non-
targeting siRNAs. Transfection efficiency was evaluated using siRNAs against KIF11, a protein
essential for cell survival [23]. The same experimental protocol than in small-scale experiments was
applied (Fig. 1A). Three images were acquired per condition with an INCell2200 automated wide-field
system. The mean cell count and the percentages of infected cells were extracted from quantification.
The screen was performed in duplicate.

For quality control purposes, we first compared the number of cells in each well in the 2 replicates.
We observed an expected distribution of the number of cells in 3 fields with a median close to 1000
cells per well for the two replicates (Fig. S1B). The number of cells per condition was slightly higher in
the first replicate than in the second one. However, the R^2 coefficient of determination of the linear
regression was close to 0.7 (Fig. S1C), indicating that the reproducibility of the experiment was correct.
As expected [23], siRNAs against KIF11 were lethal, validating the transfection protocol (Fig. S1B, C).
The 2 screens were first analyzed by taking into consideration the intensity of the E signal per cell. The
number of cells expressing the viral protein E distributed as predicted, with a median close to 15% for
the 2 screens (Fig. S1D). As expected from pilot experiments (Fig. S1A), siRNAs targeting IFNAR-1
and -2 rescued ZIKV replication in IFN-treated cells (Fig. S1D, E). The reproducibility of the infection
status of the cells between the 2 screens, with a R^2 greater than 0.8, was satisfactory (Fig. S1E). The
data were then analysis using a second approach that identified infected cells based on the E expression
independently of the intensity of the signal. The 2 methods identified similar number of infected cells
(Fig. S1F). Results were expressed as robust Z-scores for each siRNA (Fig. 1B, C). Genes were defined
as hits when at least two over three of their robust Z scores had an absolute value superior to 2 in the
two replicates, in at least one of the analysis. The screen identified 9 antiviral genes and 12 proviral ones
(Fig. 1D). Some hits were previously described as modulators of ZIKV replication, such as IFITM3
[19,20] and LY6E [24], thus validating our loss-of-function screening approach. These twenty-one hits
were selected for further validation.

IFNα2-treated HMC3 cells were transfected with pool of 3 siRNAs against each candidate, and not
by individual ones as in the primary screening. Twenty-four hours post-ZIKV infection, intracellular
viral RNA production was quantified by RT-qPCR and the number of cells positive for the viral protein
E was assessed by flow cytometry analysis. The same samples were used to assess the efficacy of the
siRNAs. RT-qPCR analyses revealed that 15 out of the 21 siRNA pools were reducing the expression
of their respective targets in IFN-treated cells (Fig. S1G). mRNAs levels of C1R, XCL1, GBP3, NADK,
C22orf39 and RUBCN were below the detection limit in IFNα2-treated HMC3 (Fig. S1G). These genes
were thus excluded from further anaysis. Reduced expression of IRF9, IFITM3, MTA2 and GPD2
significantly enhanced both viral RNA yield and the number of infected cells as compared to IFNα2-
treated cells transfected with control siRNAs (Fig. 1E and F). Both IRF9, which belongs to the ISGF3
complex [25], and IFITM3 [26] are well-known broadly-acting IFN effectors. The activities of MTA2
have, so far, not been linked to antiviral immunity. MTA2 is a component of the nucleosome remodeling
deacetylase NuRD complex, which exhibits ATP-dependent chromatin remodeling activity in addition
to histone deacetylase activity [27]. Ten times more viral RNA copies were recovered in cells silenced
for MTA2 expression than in control cells (Fig. 1E) and four times more cells were positive for the viral
E protein (Fig. 1F). These effects were comparable to the ones induced by IFNAR1 silencing (Fig. 1E
and F). Transfection with siRNA against the other 3 antiviral candidates (PXK, NMI and IFI16) had no
significant effect on ZIKV replication in these assays (Fig. 1E and 1F), suggesting that they may be
false positive candidates. Reducing the expression of LY6E, ISG15 and APOL3 significantly decreased
both viral RNA production and the number of cells positive for the E protein (Fig. 1G and 1H), validating
the pro-viral activities of these 3 candidates. The pro-viral function of USP18 and NAPA were also
validated since reducing their expression led to a significant reduction of the number of infected cells
as compared to control cells (Fig. 1H). Reduced expression of ISG20 or CCND3 had no significant
effect on ZIKV replication (Fig. 1G and 1H). IRF2, which was identified as a pro-viral hit by the screen,
behaved like an antiviral gene in the validation experiments (Fig. 1G). Together, these experiments
validated the antiviral function of IRF9, IFITM3, MTA2 and GPD2 and the pro-viral function of LY6E,
USP18, ISG15, APOL3 and NAPA in IFNα2-treated HMC3 cells infected with ZIKV.
196
Effect of a selection of candidate genes on HCV and SARS-CoV-2 replication
We next explored the ability of 10 candidate genes (IRF9, IFITM3, MTA2, GPD2, LY6E, USP18,
ISG15, APOL3, GBP3 and NAPA) to modulate the replication of two other pathogenic RNA viruses:
Hepatitis C virus (HCV) and Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2), which
are, respectively, related and unrelated to ZIKV. HCV, which is a member of the Hepacivirus genus
within the *Flaviviridae* family, has a tropism for hepatocytes. SARS-CoV-2 belongs to the
Coronaviridae family and has a tropism for pneumocytes and enterocytes.

HCV infections were conducted in hepatoma Huh-7.5 cells, which support well viral replication [28].
Huh-7.5 cells are unable to induce IFN expression since they express an inactive form of RIG-I [29] but
they possess an intact JAK/STAT pathway and do thus respond to IFN treatment [30]. RT-qPCR
analyses revealed that 8 out 10 siRNA pools efficiently reduced the expression of their respective targets
in stimulated Huh-7.5 cells (Fig. S2A). Since LY6E and APOL3 mRNA levels were under the limit of
detection of the assays in IFNα2-treated Huh-7.5 cells (Fig. S2A), they were excluded from further
analysis. As expected, reduced expression of IFNAR and IRF9 significantly enhanced HCV RNA yield
and the production of infectious particles in IFNα2-treated Huh-7.5 cells, as compared to control cells
(Fig. 2A, B). Reduced expression of IFITM3, MTA2 or GPD2, which significantly enhanced ZIKV
replication in HCM3 cells (Fig. 1E, F), did not affect HCV RNA production (Fig. 2A). However,
surprisingly, their reduced expression triggered a significant decrease in the release of infectious HCV
particles as compared to control cells (Fig. 2B). This suggests that they might favor a late stage of HCV
replication cycle. RT-qPCR analysis and titration assays identified USP18 and ISG15 as pro-HCV
factors in IFNα2-treated Huh-7.5 cells (Fig. 2C, D), validating previous results [31–34]. Of note, HCV
RNA production and infectious particle release were significantly increased in cells with reduced NAPA
levels (Fig. 2C, D), suggesting that NAPA may exert an antiviral effect on HCV, while it was not the
case for ZIKV (Fig. 1G, H).
SARS-CoV-2 replication was assessed in A549 alveolar epithelial cells expressing the viral receptor
ACE2 (A549-ACE2) by RT-qPCR and flow cytometry analysis using an antibody against the viral
protein Spike (S). Of note, silencing GBP3 in A549-ACE2 cells triggered cell death. RT-qPCR analyses
showed that all siRNA pools were reducing the expression of their respective targets in stimulated A549-
ACE2 cells (Fig. S2B). These analyses revealed the ability of IRF9 to act as an anti-SARS-CoV-2 gene
(Fig. 2E, F). Unexpectedly, GPD2 and IFITM3, which we identified as genes possessing anti-ZIKV
activities (Fig. 1), tended to behave like pro-viral genes in the context of SARS-CoV-2 infection (Fig.
2E and 2F). Viral RNA yields decreased significantly in cells silenced for USP18, ISG15 and NAPA
expression, as compared to cells transfected with control siRNAs (Fig. 2G), suggesting that these 3
genes promote viral replication in stimulated A549-ACE2 cells. By contrast to what we observed in
ZIKV infected cells (Fig. 1G, H), LY6E seemed to restrict SARS-CoV-2 (Fig. 2H). These results are in
accordance with a recent report [35]. Reducing MTA2 or APOL3 expression did not affect SARS-CoV-
2 replication.
These results suggest that some genes are broadly-acting IFN effectors, such as IRF9 and ISG15.
Other genes appeared to have evolve modulatory function toward a specific viral family or genus, such
as APOL3 and MTA2. Finally, some genes, including LY6E, IFITM3 and GPD2, exhibited opposite
modulatory functions towards different viral species.
238
ZIKV, DENV-2, WNV, VSV and MeV, but not MVA, require the expression of APOL3 for
optimal replication in IFN-treated cells

LY6E, ISG15 and APOL3 exhibited significant pro-ZIKV activities in stimulated cells, as measured

by cell-associated viral RNA levels (Fig. 1G) and percentage of E-positive cells (Fig. 1H). Among these

3 genes, APOL3 is the least described and was thus selected for further characterization. APOL3 is one

of the 6 members of the apolipoprotein L gene family. Apolipoproteins are typically associated with the

transport of lipids in the organism and were originally described as members of the high-density

lipoprotein family, which are involved in cholesterol transport [36]. In human cells, the expression of

the 6 members of the APOL gene family are up-regulated by multiple pro-inflammatory signaling

molecules, including IFNs and TNFα [36,37]. These regulations suggest a link between APOL proteins

and the innate immune system. siRNA targeting APOL2, APOL3, APOL4, APOL5 and APOL6 were

present in our library (Table S1). Among these 5 APOLs, only APOL3 was identified as a facilitator of

ZIKV infection by our screen (Fig. 1). We decided to test the ability of APOL1 to modulate ZIKV

replication since it was previously identified in a high-throughput overexpression screen as an ISG able

to increase YFV infection in $STAT1^{-/-}$ fibroblasts and Huh-7cells [21].

Analysis of mRNA levels of APOL1 and APOL3 revealed that the genes were upregulated by

IFNα2 treatment in HMC3 cells (Fig. 3A). Both genes thus qualify as genuine ISGs in these cells. The

implication of APOL1 and APOL3 in ZIKV replication was investigated using loss-of-function

approaches. siRNA-silencing reduced the levels of APOL1 and APOL3 mRNAs by ~80% and ~85%,

respectively, when compared to cells expressing scrambled control siRNAs (Fig. 3B). USP18, which is

known to negatively regulates the JAK-STAT pathway, and, as such, is a broad-spectrum pro-viral

factor [38], was identified during our screen as a pro-ZIKV candidate in HMC3 cells (Fig. 1D). Since

its pro-ZIKV function was validated in our system (Fig. 1H), siRNAs specific for USP18 were used as

positive controls. siRNA-silencing reduced the abundance of USP18 mRNAs by ~80% when compared

to cells expressing control siRNAs (Fig. 3B). Viral replication was assessed by flow cytometry by

measuring the number of cells positive for the viral protein E in cells silenced for APOL1, APOL3 or

USP18, treated or not with IFNα2. Since ZIKV is sensitive to IFNα2-treatment (Fig. S1A), higher MOIs

were used in IFNα2-treated cells than in untreated ones to compensate for its antiviral effects. As

expected (Fig. 1H), reduced expression of USP18 significantly decreased the number of IFN-treated

cells positive for the viral protein E, as compared to cells transfected with control siRNAs (Fig. 3C).

Extinction of APOL1 and APOL3 resulted in a modest, but reproducible, decrease in the number of E-

positive HMC3 cells pre-treated with IFNα2 (Fig. 3C). A pro-viral effect of APOL1 was also observed

in unstimulated cells (Fig. 3C). The efficacy of the siRNAs against APOL1 and USP18 were further

validated by Western Blot analysis using specific antibodies (Fig. 3D). APOL3 levels in cell lysates

could not be assessed due to the lack of available antibodies. Levels of expression of the viral proteins

NS5 and E were slightly decreased in IFNα2-cells expressing reduced levels of APOL1 or APOL3,

compared to control cells (Fig 3D). Together, these results suggest that ZIKV requires the expression of

APOL1 and APOL3 for optimal replication in HMC3 cells. By contrast to APOL1, the pro-viral action

of APOL3 was dependent on IFNα2-treatment.

To ensure that the APOL1- and APOL3-mediated modulation of viral replication was not restricted

to HMC3 cells, silencing experiments were performed in ZIKV-infected human podocytes treated or

not with IFNα2. Podocytes are physiologically relevant for ZIKV infection since viral RNA was

detected in kidneys of infected patients [39]. Assessing the mRNA abundance of APOL1 and APOL3

by RT-qPCR analysis of cells treated or not with IFNα2 revealed that both genes qualify as ISGs in

podocytes (Fig. 3E). siRNA-mediated silencing of APOL1, APOL3 and USP18 was efficient in

podocytes (Fig. 3F). Reduced expression of APOL1 or APOL3 resulted in a significant decrease of the

percentage of infected cells in IFNα2-treated podocytes, but not in unstimulated cells (Fig. 3G). Western

blot analysis performed in IFNα2-treated podocytes revealed that cells expressing little APOL1/3 were

producing less viral proteins than controls cells (Fig. 3H), confirming the pro-ZIKV activity of the two

APOLs. These data revealed that APOL3 and APOL1 facilitate the replication of ZIKV in podocytes

treated with IFNα2.

We tested whether APOL1 and APOL3 were active against DENV-2 or WNV, which are mosquito-

borne flaviviruses closely related to ZIKV. HMC3 cells were treated or not with IFNα2 and the MOIs

were adapted to the IFNα2 treatment. Flow cytometry analysis using anti-E antibodies revealed that

both DENV-2 and WNV replication were significantly decreased in IFNα2-treated cells silenced for

APOL1 or APOL3 expression (Fig. 3I). Reducing APOL1 and APOL3 expression in non-treated cells

also significantly reduced WNV replication (Fig. 3I). Thus, APOL1/3 may well have flavivirus genus-

specific proviral activities since they seems to contribute to ZIKV, WNV and DENV replication (Fig.

3C, D, G, H and I) but not to SARS-CoV-2 replication (Fig. 2G, H). To further delineate the spectrum

of action of these two genes towards other viruses, we tested the effect of APOL1/3 silencing on the

replication of Vesicular Stomatitis virus (VSV) and Measles virus (MeV), which are negative-strand

RNA viruses belonging to the *Rhabdoviridae* and *Paramyxoviridae* families, respectively. Experiments

were performed with a MeV strain modified to express GFP [40]. We also included in the analysis

Modified Vaccinia Ankara virus (MVA), a DNA virus belonging to the *poxviridae* family, that was

engineered to express GFP (MVA-GFP). Flow cytometry analysis using an antibody against the viral

protein G revealed that VSV was highly dependent on APOL1 and APOL3 expression for efficient

replication in IFNα2-treated HMC3 cells (Fig. 3I). Optimal replication of MeV-GFP in stimulated

HMC3 cells also required APOL1 and APOL3 expression (Fig. 3I). APOL1 proviral activity was also

observed in unstimulated cells (Fig. 3I). By contrast, MVA-GFP replication was not affected by reduced

expression of either APOL1 or APOL3 (Fig. 3I).

Together, these experiments suggest that APOL1 and APOL3 could favor a replication process

shared by ZIKV, DENV-2, WNV, VSV and MeV. Unlike APOL1 in HMC3 cells, APOL3 pro-viral

activities were dependent on IFN-treatment.

APOL1 and APOL3 likely promote viral replication independently of their interaction with phosphoinositides

Recent data revealed that APOL1 and APOL3 play a role in lipid metabolism in podocytes and, more specifically, in the regulation of the production of phosphatidylinositol-4-phosphate (PI(4)P), via an indirect interaction with the PI4KB kinase [41]. PI(4)P is involved in Golgi secretory functions by facilitating the recruitment of proteins that promote vesicular transport [42]. PI(4)P is also essential for the establishment of efficient viral replication via the formation of membranes which serve as platforms for the production of viral RNA [43,44]. APOL1 and/or APOL3 could thus impact ZIKV replication via their ability to regulate the production of PI(4)P. To test this hypothesis, we first assessed APOL1 and APOL3 localization in HMC3 cells. In the absence of specific antibodies for APOL1 and APOL3 validated for immunofluorescence assays, we investigated the localization of GFP-tagged versions of APOL3 and APOL1 in HMC3 cells, together with markers for the Golgi apparatus (Fig. 4A), early or late endosomes (Fig. S3). APOL1-GFP and GFP-APOL3 localized in closed proximity to the cis-Golgi (Fig. 4A), where PI(4)P and PI4KB localize [45], and not in late nor early endosomes (Fig. S3). In line with this, APOL1-GFP and GFP-APOL3 associated with PI4KB in HMC3 cells (Fig. 4B). Of note, APOL1-GFP was also detected in vesicle-like structures whose identity could not be established (Fig. 4A, white arrow). They may represent lipid droplets or fragmented Golgi. The localization of APOL1-GFP and GFP-APOL3 could not be investigated in ZIKV-infected cells since we observed that transfection rendered cells non-permissive to viral infection.

We then performed experiments with a well-characterized PI4KB kinase inhibitor that decreases PI(4)P expression [46]. We first analyzed by immunofluorescence the intensity of the PI(4)P signal in cells treated for 24 h with different concentrations of the drug in HCM3 cells. The presence of the PI4KB inhibitor triggered a dose-dependent decrease of the PI(4)P signal (Fig. 4C), suggesting that the drug is efficient in HMC3 cells. We then infected cells with ZIKV in the presence of different concentration of the inhibitor. Since the effect of APOL3 on ZIKV replication is dependent on IFNα2 (Fig. 3), the analysis was also performed in stimulated cells. Coxsackie B3 virus (CVB3), an enterovirus that replicates in a PI(4)P-dependent manner, was used as a positive control since its replication is sensitive to the drug [47]. As negative controls, we used cells infected with WNV, whose replication is not affected by the PI4KB inhibitor [48]. As previously shown in HeLa cells [47], a dose-dependent reduction of the number of cells positive for the CVB3 viral protein 1 (VP1) was triggered by the inhibitor treatment (Fig. 4D). As shown previously in monkey cells [48], WNV replication was unaffected by the PI4KB inhibitor in HCM3 cells (Fig. 4D). ZIKV protein production was not sensitive to the treatment with the PI4KB kinase inhibitor, independently of the presence of IFNα2 (Fig. 4D). These experiments suggest that the pro-viral activities of APOL1 and APOL3 are not related to their interaction with PI4KB or PI(4)P in microglial cells.

MTA2 restricts ZIKV replication in IFNα2-stimulated human cells.

MTA2 was identified in our screen as a gene with potent anti-ZIKV activities (Fig. 1). MTA2 shows

a very broad expression pattern and is strongly expressed in many tissues. It belongs to the NuRD

complex, which establishes transcriptional modulation of a number of target genes in vertebrates,

invertebrates and fungi[27]. Since its function has, so far, not been linked to viral infection, we decided

to investigate its potential antiviral activities further. To ensure that the MTA2-mediated inhibition of

viral replication was not restricted to HMC3 cells, experiments were also performed in Huh-7 hepatoma

cells, which are physiologically relevant for flavivirus infection and are thus extensively used in

Flaviviridae research. siRNAs targeting IFNAR1 were used as positive controls in these experiments.

siRNA-silencing reduced the levels of MTA2 and IFNAR1 mRNAs by at least 80%, when compared to

cells expressing scrambled control siRNAs, independently of the stimulation or infection status of

HMC3 and Huh-7 cells (Fig. 5A-D). MTA2 was included in our gene list because its expression

clustered with MX1 in T cells (Cerboni et al. in preparation). However, MTA2 mRNA abundance, as

measured by RT-qPCR analysis, remained unchanged upon IFNα2 treatment in both cells types (Fig.

5A and 5C), indicating that MTA2 is not an ISG in these cells.

Assessment of viral replication by RT-qPCR revealed that cell-associated viral RNA yields were

significantly higher in IFN-treated HMC3 cells silenced for MTA2 expression, as compared to controls

cells (Fig. 5E). This is in line with previous results (Fig. 1E). Cytometry analysis using anti-E antibodies

confirmed that MTA2 anti-ZIKV activities were dependent on the presence of IFN in HCM3 cells (Fig.

5F). Since MTA2 is not an ISG in HMC3 cells (Fig. 5A), these results suggest that MTA2 may require

an active IFN signaling to exert its anti-ZIKV activities in these cells. As in HMC3 cells, reduced

expression of MTA2 triggered a significant increase of intracellular viral RNA production in stimulated

Huh-7 cells (Fig. 5G). Reducing MTA2 expression had a more pronounced effect on the percentage of

infected cells that reducing IFNAR1 expression in stimulated Huh-7 cells (Fig. 5H). Albeit to a lesser

extent than in stimulated cells, MTA2 anti-ZIKV activity was also observed in non-stimulated Huh-7

cells by flow cytometry and RT-qPCR analysis (Fig. 5G and H).

The effect of MTA2 on viral protein production was further assessed by Western blot analysis using

anti-E and anti-NS5 antibodies in stimulated and unstimulated HMC3 and Huh-7 cells. These

experiments validated further the efficacy of the siRNAs against MTA2 in both cell lines (Fig. 5I and

5J). Expression of the viral proteins NS5 and E were increased in stimulated HMC3 and Huh-7 cells

expressing reduced levels of MTA2 or IFNAR1, as compared to control cells (Fig. 5I and 5J). In

agreement with the flow cytometry analysis (Fig. 5H), MTA2 anti-ZIKV activity was less dependent of

IFN-treatment in Huh-7 cells than in HMC3 cells (Fig. 5I and 5J). As observed in flow cytometry

analysis (Fig. 5H), MTA2 effect on viral protein production was more potent than the one of IFNAR1

in stimulated Huh-7 cells (Fig. 5J).

These results represent the first evidence of the ability of MTA2 to restrict the replication of any

virus.

MTA2 restricts YFV and WNV replication in IFNα2-stimulated Huh-7 cells.

We tested whether MTA2 was active against WNV and YFV in Huh-7 cells, which are permissive to these 2 flaviviruses. As in previous experiments, higher MOIs were used in the presence of IFNα2. Cytometry analysis revealed that MTA2 silencing significantly enhanced the replication of these 2 flaviviruses in an-IFN dependent manner (Fig. 6A and 6B), indicating that MTA2 antiviral activity is broader that ZIKV. We then tested the effect of MTA2 silencing on the replication of VSV and MeV in Huh-7 cells. Reduced expression of MTA2 decreased the number of cells infected with VSV and MeV (Fig. 6C and 6D), independently of the IFN stimulation. This is consistent with the pro-HCV activity of MTA2, as measured by titration in stimulated Huh-7.5 cells (Fig. 2B). MTA2 may thus possesses a flavivirus genus-specific antiviral function.

Discussion

Several gain-of-fonction screens have been performed to identify ISGs that modulate flavivirus infection. Pionner screens tested the activities of relatively small amounts of ISGs by overexpression [49,50]. The first comprehensive overexpression screen in which more than 380 ISGs were evaluated for antiviral activity against six viruses, including the *Flaviviridae* HCV, WNV and YFV, was published in 2011 by Schoggins and collaborators [21]. To avoid potential physiological irrelevance induced by gene overexpression, we opted for a silencing approach to identify genes that modulate ZIKV replication in an IFN-induced state. We used a siRNA library which was designed in the context of an HIV project. A limitation of our library is that targeted genes were selected based on a transcriptomic analysis of primary T cells stimulated by contacts with activated monocytes (Cerboni et al. in preparation), and not on ZIKV-target cells. Nevertheless, it contains a high fraction of core ISGs that overlaps with previous screens [21,22]. Furthermore, most arrayed screens designed to identify cellular factors that modulate ZIKV replication, including ours, monitored viral replication after a single round of infection, often by assessing viral protein expression. Therefore, only genes that inhibit early stages of viral replication, up to protein production, can be identified. Quantifying viral titers in supernatants collected from individual wells of the first round of screening should identify genes that affect late stages of viral replication, such as viral assembly, maturation and release, as well as viral infectivity. Alternatively, viral replication could be monitored after several rounds of infection. Nevertheless, despite these two main limitations, our screening strategy identified 21 genes affecting the number of cells positive for the viral protein E in IFN-treated microglial cells.

Some hits were previously described as ISGs able to modulate ZIKV replication, such as IFITM3 [19,20] and LY6E [24], thus validating our screening approach. Despite being in our gene list, Viperin, IFI6, PARP-12 and C19orf66, which are known to affect ZIKV replication in human cells [51–55], were not identified as viral modulators by our strategy. They may have a weaker influence on viral replication in HMC3 cells than in the cells in which their role was previously established [51–55]. In line with this

hypothesis, Viperin restricts the replication of several neurotropic flaviviruses in a cell type-dependent

manner [56]. One can also envisage that the expression levels of Viperin, IFI6, PARP-12 and C19orf66

are low in HMC3 cells, even when stimulated by IFNα2, and therefore are poorly, if at all downregulated

by specific siRNAs.

We validated the role of 5 hits as genes contributing to an optimal ZIKV replication in stimulated

HMC3 cells using RT-qPCR and flow cytometry analysis: LY6E, USP18, ISG15, APOL3 and NAPA.

The identification of LY6E as a gene enhancing ZIKV replication was expected, since it was previously

shown to promote the internalization of flaviviruses in U2OS human osteosarcoma cells [24]. Since

ISG15 and USP18 negatively regulate IFN signaling pathway [57,58], they are expected to act as broad

pro-viral ISGs. NAPA interacts with SNARE protein complexes to trigger their disassembly [59].

SNARE proteins belong to a superfamily of membrane fusion proteins that localize at the plasma

membrane, the Golgi apparatus and on different endocytic vesicles. They regulate the traffic of these

vesicles between the plasma membrane and the Golgi [60]. Several viruses, including influenza A virus

and VSV, hijack SNARE proteins to enter host cells [61]. SNARE complexes may thus contribute to

NAPA pro-viral activities. We validated the anti-viral functions of 4 screen hits in stimulated HMC3

cells infected with ZIKV: IRF9, IFITM3, MTA2 and GPD2. Identification of IRF9, which plays a key

role in ISG expression [25], and IFITM3, which restricts early stages of ZIKV infection [19,20],

validates our screening strategy. GPD2 is a mitochondrial protein involves in the metabolism of

glycerol. No link between GPD2 and viral infections has been established yet. However, it regulates

inflammatory response in macrophages [62]. Further experiments will be required to confirm that NAPA

and GPD2 have the ability to modulate ZIKV replication in stimulated human cells.

Experiments performed on cells infected with HCV or SARS-CoV-2 contributed to delineate the

spectrum of action of a selection of the screen hits. Our data illustrate once again that some ISGs have

virus-specific antiviral activities [63]. For instance, we found that ZIKV, but not the related HCV, was

sensitive to IFITM3 expression, confirming (HCV) and extending (ZIKV) recently reported data

[20,64]. Our data also confirm that some ISGs exert opposite effect on different viruses. For instance,

as described previously, LYE6 promotes the replication of ZIKV [24] but restricts that of SARS-CoV-

2 [35]. Viruses have developed numerous innovative strategies to evade ISG-mediated restriction

[65,66]. Hijacking individual ISG for promoting their replication is one of them. This hypothesis may

explain why, to our surprise, our screen recovered more pro-viral genes that antiviral ones.

Our results identified APOL3 and APOL1 as ISGs required for optimal ZIKV replication in HMC3

cells. The proviral activity of APOL1 was less dependent on IFN that the one of APOL3, which suggest

that both proteins act via different mechanisms. Reduced expression of APOL1 and APOL3 also

restricted the replication of WNV and DENV in stimulated HMC3 cells. In line with these findings,

over-expression of APOL1 was previously reported to increase YFV replication in Huh-7 cells [21].

VSV replication was highly reduced in the absence of one of these 2 genes. APOL1 has been reported

to act as an antiviral ISG in the context of infection with alphaviruses (Sindbis virus and Venezuelan

equine encephalitis virus) and human parainfluenza virus [21,67]. Its over-expression also inhibits HIV-

1 infection in monocytes [68]. Thus, APOL1 and APOL3 seem to behave like pro- or anti-viral ISG

depending on the virus, or have no obvious role (HCV). In line with our data on flaviviruses, a recent

report found that overexpression of APOL1 promoted infection with ZIKV and DENV-2, confirming a

proviral role for this factor [69]. However in this study, an increase of ZIKV, DENV-2 and HCV

replication was also reported in Huh-7.5 cells expressing siRNAs targeting APOL1 and APOL3 in the

absence of IFN treatment [69], yielding conflicting data that will deserve further investigation.

From our data, we formulated the hypothesis that APOL1 and APOL3 pro-viral activities could be

linked to their ability to bind to anionic phospholipids, including several phosphoinositides, in particular

PI(4)P [41]. Both APOL1 and APOL3 were detected in PI(4)P-containing liposomes [41]. Moreover,

reduced expression of APOL3 resulted in reduction of PI(4)P levels in podocytes [41]. PI(4)P plays a

pivotal role in the Golgi secretory functions by facilitating recruitment of proteins that promote vesicular

transport [42]. Our immunofluorescence data revealed that GFP-tagged version of APOL1 and APOL3

localize mainly in the Golgi, where PI(4)P localizes [42]. Numerous RNA viruses, including various

Picornaviridae, HCV, coronaviruses and parainfluenza type 3, rely on PI(4)P to build membranous

replication platform, where viral replication and assembly take place [70]. Flaviviruses are no exception

and also trigger intracellular membrane remodelling for the building of membranous replication

platforms [71]. Experiments conducted with a well-characterized PI4KB kinase inhibitor excluded the

possibility that APOL1 and APOL3 pro-viral effects depend on PI(4)P and its synthetizing protein,

PI4KB. In line with these results, PI(4)P are not important for the replication of WNV and the related

Usutu virus [48]. Other avenues should be thus explored to understand how APOL1 and APOL3

modulate the replication of several RNA viruses. Since APOL1 and APOL3 impact the replication of

unrelated viruses, they may be negative regulators of IFN signalling pathway, by, for instance,

contributing to the proper routing of members of the JAK/STAT pathway.

Our data revealed that MTA2 possesses potent antiviral function in the context of ZIKV, WNV and

YFV in stimulated cells, whereas its exhibited a proviral role for HCV, VSV and MeV. Reducing MTA2

expression in the presence of IFNα2 enhanced flaviviral replication to a level comparable to the

inhibition of IFNAR1. Despite being part of our gene list, MTA2 is not induced by IFN in HMC3 Huh-7

cells but its antiviral activity was dependent on IFN in these cells. MTA2 may thus interact with an ISG

to act on viral replication. MTA2 is a component of the NuRD complex, an unusual complex which

exhibits ATP-dependent chromatin remodeling activity in addition to histone deacetylase activity [27].

The complex establishes transcriptional modulation of a number of target genes in vertebrates,

invertebrates and fungi [27]. MTA2 related activities have not, so far, been linked to innate immunity

in virus-infected cells. However, a link between the NuRD complex and STAT1-mediated IFN response

was established in the context of infection with the protozoan parasite *Toxoplasma gondii* [72]. A

Toxoplasma protein, named TgIST, translocates to the host cell nucleus where it recruits the complex

NuRD to STAT1-dependent promoters, resulting in altered chromatin and blocked STAT1-mediated

transcription [72]. Moreover, HDAC1, which is also a member of the NuRD complex, associates with both STAT1 and STAT2 in human cells [73]. Furthermore, specific reduction of HDAC1 expression inhibits IFNβ-induced transcription whereas HDAC1 overexpression enhances IFNβ-induced transcription [73]. Finally, HDAC inhibitors block the formation of ISGF3 and this was associated with impairment of STAT2 nuclear accumulation in mouse L929 cells [74]. These findings indicate a fundamental role for deacetylase activity and HDAC1 in transcriptional control in response to IFN. One could thus envisage that MTA2, within the NuRD complex, also interacts with STAT1 in cells stimulated with IFN and favors its action locally. This interaction could restrict flavivirus infection, possibly *via* enhancing the expression of a subset of flavivirus-specific ISGs.

Our work identified previously unrecognized genes that modulate the replication of RNA viruses in an IFN-dependent way. Future studies combining transcriptomic analysis of IFN-treated cells and high throughput loss-of-function screens will help define the interferome of cell types relevant for viral infection. Such studies are primordial to continue investigating the complexity the IFN-mediated antiviral program.

Materials and Methods

Cells

Human microglial cells (HMC3) were purchased from the American Type Culture Collection (ATCC, CRL-3304). They were maintained in Dulbecco's modified Eagle's medium (DMEM) containing GlutaMAX I and sodium pyruvate (Gibco), supplemented with 10% fetal bovine serum (FBS) and 1% penicillin-streptomycin (P/S) (final concentration of 100 units/mL and 100 μg/mL, respectively) (Sigma) and non-essential amino acids (GibcoTM NEAA 100X MEM, Life Technologies). Podocytes were described previously [75]. They were grown at 33°C in Roswell Park Memorial Institute medium (RPMI) containing GlutaMAX I (Gibco) and supplemented with 10% FBS and P/S. Before any experiments, cells were differentiated during 7 days at 37°C. Human hepatocellular carcinoma Huh-7 cells [76], which were kindly given by Cinzia Traboni (IRBM, Pomezia, Italy), were maintained in DMEM supplemented with 10% FBS and 1% P/S. Huh-7.5 cells (Apath, LLC), a subclone of Huh-7 cells [76] were cultured in DMEM supplemented with non-essential amino acids, 1mM sodium pyruvate, 10% FBS and P/S. Vero NK cells, which are African green monkey kidney epithelial cells, were purchased from ATCC and used for viral titration assays. They were maintained in DMEM containing GlutaMAX I and sodium pyruvate (Gibco), supplemented with 10% FBS and P/S. *Aedes albopictus* C6-36 cells were maintained in Leibovitz's L-15 medium containing 10% FBS, 1% P/S, 1% Non-Essential Amino Acids Solution (Gibco) and 2% Tryptose Phosphate Browth (Gibco). Human lung epithelial A549 cells were modified to stably express hACE2 using the pLenti6-hACE2 lentiviral

transduction, as described previously [77]. Cell cultures were verified to be mycoplasma free with the

MycoAlertTM Mycoplasma Detection Kit (Lonza).

Virus stocks, titration and infection

The Zika strain PF13 (kindly provided by V. M. Cao-Lormeau and D. Musso, Institut Louis Malardé,

Tahiti Island, French Polynesia) was isolated from a viremic patient in French Polynesia in 2013. Stocks

were produced on C6-36 cells. The Dengue 2 virus (DENV-2) strain Malaysia SB8553 was obtained

from the Centro de Ingeniería Genética y Biotecnología (CIGB), Cuba. The YFV Asibi strain and the

WNV Israeli strain IS-98-STI were provided by the Biological resource Center of the Institut Pasteur.

Stocks of DENV-2, YFV and WNV were produced on Vero NK cells. Viruses were concentrated by

polyethylene glycol 6000 precipitation and purified by centrifugation in a discontinued gradient of

sucrose. Flaviviruses were titrated on Vero NK cells by plaque assay as previously described [78] and

titers were expressed in plaque-forming units (PFU)/ml. The Measles Schwarz strain expressing GFP

(MeV-GFP) was described previously [40]. VSV Indiana and the CVB3 Nancy strain were kindly

provided by N. Escriou (Institut Pasteur) and M. Bessaud, respectively (Institut Pasteur). Modified

Vaccinia Ankara virus (MVA) expressing eGFP (MVA-GFP) was kindly provided by the ANRS via O.

Schwartz (Institut Pasteur). It was manufactured by Transgene (Illkirch-Graffenstaden, France).

The fluorescent marker, eGFP, is expressed under the control of the early promotor p11K7.5 and viral

preparations were purified by tangential flow filtration. HMC3 cells were infected at the following

MOIs: a MOI of 10 with DENV-2, 0,5 with WNV, 0,005 with VSV, 1 with MeV-GFP and 0,05 with

MVA-GFP. IFN-treated HMC3 cells were infected at a MOI of 20 with DENV-2, 5 with WNV, 0,01

with VSV, 2 with MeV-GFP and 0,25 with MVA-GFP. Huh-7 were infected at a MOI of 1 with YFV

and 0,25 with WNV. IFN-treated Huh-7 cells were infected at a MOI of 10 with YFV and WNV. Highly

cell culture-adapted HCV Jad strain was obtained following transfection of Huh-7.5 cells with *in vitro*

transcribed genome-length RNA as described previously [79–81]. Large volumes of HCV stocks were

prepared following infection at a MOI of 0.01 50% tissue culture infectious doses 50 (TCID50) per cell

with supernatants collected post-RNA transfection [82]. HCV infectious titers were determined by

TCID50 assays in Huh-7.5 cells as described previously [79]. IFN-treated Huh-7.5 cells were infected

at MOI of 3 TCID50/cell with HCV. The SARS-CoV-2 strain BetaCoV/France/IDF0372/2020

(historical) was supplied by the French National Reference Centre for Respiratory Viruses hosted by

Institut Pasteur (Paris, France) and headed by Pr. S. van der Werf. The human samples from which the

strain was isolated were provided by Dr. X. Lescure and Pr. Y. Yazdanpanah from the Bichat Hospital,

Paris, France and Dr. Vincent Foissaud, HIA Percy, Clamart, France, respectively. A549-ACE2 cells

were infected with SARS-CoV-2 at a MOI of 2.

High throughput Screen

Five hundreds HMC3 cells were seeded in 384-well microplates in the morning of day 1 using a

MultiDrop combi liquid dispenser (Thermo Fisher Scientific), in 40 μL of cell culture media. Cells were

allowed to adhere for 4 hours (+/-1h) before transfection with individual siRNAs (10nM) diluted in a

mix of OptiMEM (Life Technologies) and 0.05μL of Interferin reagent (Polyplus Transfection). siRNAs

were transfected using an Evo 150 with MCA384 (Tecan). The library contained 1158 siRNA targeting

386 genes. siRNA targeting KIF11 was used to assess the transfection efficiency. On day 2, cells were

treated with 1000U/ml of IFNα2a. Interferon was diluted into cell culture media and 10μL of the mix

was robotically transferred to each well (except non-treated controls). 24 hours after IFN treatment, cell

media was removed from siRNA-transfected plates and 40μL of the ZIKV PF13 strain, diluted to a final

concentration of 7,500 particles/well, was added to the plates with the MCA384 head (Tecan). ZIKV

titer was $6.5.10^8$ PFU/ml. Cells were then incubated 24 hours prior to fixation. Cells were fixed with 4%

of formaldehyde (Sigma-Aldrich) for 20 min, plates were then washed once with PBS and quenched

with NH4Cl (50mM) solution. Cells were then blocked with 1% BSA solution and permeabilized with

0.5% Triton X-100. Cells were next incubated for 60 min with mouse primary antibody anti-4G2 (1:500)

which reacts with flavivirus E proteins. Cells were then washed twice in PBS solution and incubated

with Alexa Fluor 488-coupled secondary antibodies (ThermoFisher Scientific). Nuclei were stained with

0.2 μg/ml Hoechst (Sigma). Images were acquired with an INCell2200 automated wide-field system

(GE Healthcare,) using a Nikon 10X/0.45, Plan Apo, CFI/60. Three fields per well were analyzed using

the INCell Analyzer 3.7 Workstation software. Two independent screens were performed. The mean

cell count and the percentages of infected cells were extracted from quantification.

Data analysis and hit calling

In the first analysis, data were processed using a software developed internally at the Biophenics

platform. For hit identification, the robust Z-score method was used under the assumption that most

siRNAs are inactive against ZIKV and can serve as controls [83,84]. Raw values were log-transformed

for cell count only to make the data more symmetric and close to normal distribution. In order to correct

for plate positional effects, median polishing [84] was applied to each analyzed feature. It iteratively

subtracts row, column and well median, computed from all plates within one screen. Hits for each

compound were identified as follows: sample median and median absolute deviation (MAD) were

calculated from the population of screening data points (named as sample) and used to compute Robust

Z-scores (RZ-scores) according to a formula, in which the reference population corresponds to the

siRNA-treated wells, and MAD is defined as the median of the absolute deviation from the median of

the corresponding wells:

$$rZ\text{-}Score = \frac{x - med(pop\ réf)}{1.4826\ x\ med(|pop\ réf - med(pop\ réf)|)}$$

A gene was identified as a 'hit', if the RZ-score was < -2 or > 2 pointing in the same direction for 2
siRNAs targeting the same gene in both screens. Final values in the hit table correspond to the RZ-score
of the second strongest siRNA. In the second analysis, data were process using an homemade script and
CellProfiler [85]. Nucleus and viral assembly sites detected by the E signal were counted. As in the first
analysis, rZ-Score and percentages of infected cells were quantified. Considering that each gene was
targeted by three individual siRNA, genes were clusterized as hits, if at least two over three of their
robust Z score absolute value were superior to 2. Genes were defined as hits when they were identified
in at least one of the analysis.
612
Antibodies, plasmids and reagents
The following primary antibodies were used in the study: anti-E MAb 4G2 hybridoma cells, anti-NS5-
ZIKV [86], anti-VSV-G (IE9F9, Kerafast), anti-CVB3 VP1 (M7064, Agilent), anti-SARS-CoV-2 S
protein mAb10 (1 μg/ml, a kind gift from H. Mouquet, Institut Pasteur, Paris, France), APOL1
(HPA018885, Sigma), MTA2 (8106, abcam), GM130 (12480, cell signaling), PI(4)P (Z-P004), PI(4)KB
(06-578, Milipore) and anti-actin (A5316, Sigma). Secondary antibodies were as followed: anti-mouse
Alexa 488 (A11001, Life Technologies), anti-mouse Alexa 680 (A21058, Life Technologies) and anti-
rabbit DyLight 800 (SA5-35571, TermoScientific). The PI4KB inhibitor (1881233-39-1,
MedChemExpress) and IFNα2a (Sigma-Aldrich, SRE0013) were used at the indicated concentration.
GFP-APOL3 et APOL3-GFP were subcloned into pcDNA.3.1 from templates previously described
[41].
624
siRNA transfection
626   HMC3 cells were transfected using INTERFERin transfection reagent (Polyplus Transfection). Huh-7
627   cells, Huh-7.5 cells and podocytes were transfected with siRNAs at 10 nM final concentration using
628   Lipofectamine RNAiMax (Life Technologies). All siRNAs were obtained from Dharmacon
629   (siGENOME-SMARTpool).
630
RNA extraction and RT-qPCR assays
Total RNAs were extracted from cell lysates using the NucleoSpin RNA II Kit (Macherey-Nagel)
following the manufacturer's protocol and were eluted in nuclease-free water. First-strand
complementary DNA synthesis was performed with the RevertAid H Minus M-MuLV Reverse
Transcriptase (Thermo Fisher Scientific). Quantitative real-time PCR was performed on a real-time PCR
system (QuantStudio 6 Flex, Applied Biosystems) with SYBR Green PCR Master Mix (Life
Technologies). Data were analyzed with the ΔΔCT method, with all samples normalized to GAPDH.
All experiments were performed in technical triplicate. Viral genome equivalents concentrations
(GE/ml) were determined by extrapolation from a standard curve generated from serial dilutions of the
plasmid encoding the full-length genome of the Zika strain MR766 [87] or plasmids encoding a

fragment of the RNA-dependent RNA polymerase (RdRp)-IP4 of SARS-CoV-2. HCV RNA was quantified by one-step reverse transcription-quantitative PCR using 50 ng of total intracellular RNA and TaqMan® Fast Virus 1-Step Master Mix (Applied Biosystems) with primers and probe targeting the HCV 5′ nontranslated region as described previously [82]. Viral RNA levels were normalized with respect to 18S RNA levels quantified in parallel using TaqMan ribosomal RNA control reagents (Applied Biosystems). The products were analysed on a 7500 Fast Real-Time PCR system (Applied Biosystems). Serial dilutions of a genome-length *in vitro* transcribed HCV RNA served to establish standard curves and calculate HCV GE/μg total RNA concentrations. Primers and probe used for RT-qPCR analysis are given in Table S2.

Western blot analysis

Cells were collected in RIPA buffer (Sigma) containing protease inhibitors (Roche Applied Science). Cell lysates were normalized for protein content with Pierce 660nm Protein Assay (Thermo Scientific), boiled in NuPAGE LDS sample buffer (Thermo Fisher Scientific) in non-reducing conditions. Samples were separated by SDS-PAGE (NuPAGE 4–12% Bis-Tris Gel, Life Technologies) with MOPS running buffer. Separated proteins were transferred to a nitrocellulose membrane (Bio-Rad). After blocking with PBS-Tween-20 0.1% (PBST) containing 5% milk for 1 h at RT, the membrane was incubated overnight at 4°C with primary antibodies diluted in blocking buffer. Finally, the membranes were incubated for 1 h at RT with secondary antibodies diluted in blocking buffer, washed, and scanned using an Odyssey CLx infrared imaging system (LI-COR Bioscience).

Flow cytometry

Infected cells were fixed with cytofix/cytoperm kit (BD Pharmingen) and stained using the indicated primary and secondary antibodies. Non-infected, antibody-stained samples served as controls for signal background. Data were acquired using Attune NxT Acoustic Focusing Cytometer (Life Technologies) and analyzed using FlowJo software.

Immunofluorescence assay

Cells were fixed with PFA 4% (Sigma) during 20min. Cells were permeabilized with PBS Triton X-100 (0.5%) for 15min at RT. After washing with PBS, they were incubated for 30 min with PBS + 0.05% Tween 20 + 5% BSA. The slides were then incubated overnight at 4°C with primary antibodies diluted in PBS. After washing with PBS, they were incubated for 1 h with secondary antibodies and washed with PBS. Nuclei were stained using PBS/NucBlue (Life Technologies, R37606). The mounting medium used is the Prolong gold (Life Technologies, P36930). All preparations were observed with a confocal microscope (ZEISS LSM 700 inverted) and images were acquired with the ZEN software.

Statistical analysis.

Data are presented as means ± SD and were analyzed using GraphPad Prism 7. Statistical analysis of
percentage values or fold enrichment values were performed on logit or log-transformed values,
respectively. Statistical analysis was performed with two tailed paired t-test or by one- or two-way
analysis of variance (ANOVA) with Dunnet's multiple comparisons test. Each experiment was
performed at least twice, unless otherwise stated. Statistically significant differences are indicated as
follows: *: $p < 0.05$, **: $p < 0.01$ and ***: $p < 0.001$; ns, not significant.

Acknowledgments
We thank Dr. M.A. Saleem (University of Bristol, UK) for generously providing the podocytes (via E.
Pays, Université Libre de Bruxelles, Belgium); C.M. Rice (Rockefeller University, New York, USA)
for Huh-7.5 cells; Cinzia Traboni (IRBM, Pomezia, Italy) for Huh-7 cells; M. Bessaud (Institut Pasteur)
for the CVB3 Nancy strain; N. Escriou (Institut Pasteur) for the VSV Indiana strain and anti-VSV-G
antibodies; V. M. Cao-Lormeau and D. Musso (Institut Louis Malardé, Tahiti Island, French Polynesia)
for the ZIKV-PF13 strain; L. Hermida and G. Enrique Guillen Nieto from the Centro de Ingeniería
Genética y Biotecnología (CIGB), Cuba, for the DENV-2 strain Malaysia SB8553; T. Wakita (NIID,
Tokyo, Japan) for pJFH1 HCV cDNA; R. Bartenschlager (University of Heidelberg, Germany) for
pJFH1-2EI3-adapt cDNA; the French National Reference Centre for Respiratory Viruses hosted by
Institut Pasteur (France) and headed by S. van der Werf for providing the historical SARS-CoV-2 viral
strains; A. Merits (University of Tartu, Estonia) for anti-ZIKV NS5 antibodies; P. Desprès (Université
de la Réunion, PIMIT) for 4G2 hybridoma cells; H. Mouquet (Institut Pasteur) for anti-SARS-CoV-2 S
antibodies; M. Evans (Icahn School of Medicine at Mount Sinai, New York, USA) for the plasmid
encoding the full-length Zika MR766 genome; F. Porrot and O. Schwartz (Institut Pasteur) for
producing and sharing the MVA-GFP, C. Combredet (Institut Pasteur) for producing MeV-GFP; as well
as E. Pays and S. Uzureau (Université Libre de Bruxelles, Belgium) for APOL1 and APOL3 plasmids
and for stimulating APOL-focused discussions. We are grateful to the members of our laboratories for
helpful discussions and technical advice. Finally, we thank Emeline Perthame (Bioinformatics and
Biostatistics HUB, Institut Pasteur) for her help in statistical analysis.

Declaration of interests. The authors declare no competing interests.

References

1.   Woolhouse M, Scott F, Hudson Z, Howey R, Chase-Topping M. Human viruses:
discovery and emergence. Philos Trans R Soc Lond B Biol Sci. 2012;367: 2864–71.
doi:10.1098/rstb.2011.0354
2.   Streicher F, Jouvenet N. Stimulation of Innate Immunity by Host and Viral RNAs.
Trends in Immunology. 2019;40: 1134–1148. doi:10.1016/j.it.2019.10.009

3.   Schoggins JW. Recent advances in antiviral interferon-stimulated gene biology.
F1000Res. 2018;7. doi:10.12688/f1000research.12450.1
4.   Rusinova I, Forster S, Yu S, Kannan A, Masse M, Cumming H, et al. Interferome
v2.0: an updated database of annotated interferon-regulated genes. Nucleic Acids Res.
2013;41: D1040-6. doi:10.1093/nar/gks1215
5.   Touzot M, Grandclaudon M, Cappuccio A, Satoh T, Martinez-Cingolani C, Servant N,
et al. Combinatorial flexibility of cytokine function during human T helper cell
differentiation. Nat Commun. 2014;5: 3987. doi:10.1038/ncomms4987
6.   Mostafavi S, Yoshida H, Moodley D, LeBoité H, Rothamel K, Raj T, et al. Parsing the
interferon transcriptional network and its disease associations. Cell. 2016;164: 564–578.
doi:10.1016/j.cell.2015.12.032
7.   Schneider WM, Chevillotte MD, Rice CM. Interferon-Stimulated Genes: A Complex
Web of Host Defenses. Annu Rev Immunol. 2014;32: 513–545. doi:10.1146/annurev-
immunol-032713-120231
8.   Pierson TC, Diamond MS. The continued threat of emerging flaviviruses. Nat
Microbiol. 2020. doi:10.1038/s41564-020-0714-0
9.   Ashraf U, Ye J, Ruan X, Wan S, Zhu B, Cao S. Usutu virus: an emerging flavivirus in
Europe. Viruses. 2015;7: 219–38. doi:10.3390/v7010219
10.   Pierson TC, Diamond MS. The emergence of Zika virus and its new clinical
syndromes. Nature. 2018;560: 573–581. doi:10.1038/s41586-018-0446-y
11.   Matusali G, Houzet L, Satie A-P, Mahé D, Aubry F, Couderc T, et al. Zika virus
infects human testicular tissue and germ cells. J Clin Invest. 2018;128: 4697–4710.
doi:10.1172/JCI121735
12.   Rossi SL, Tesh RB, Azar SR, Muruato AE, Hanley KA, Auguste AJ, et al.
Characterization of a Novel Murine Model to Study Zika Virus. Am J Trop Med Hyg.
2016;94: 1362–9. doi:10.4269/ajtmh.16-0111
13.   Lazear HM, Govero J, Smith AM, Platt DJ, Fernandez E, Miner JJ, et al. A Mouse
Model of Zika Virus Pathogenesis. Cell Host Microbe. 2016;19: 720–30.
doi:10.1016/j.chom.2016.03.010
14.   Tripathi S, Balasubramaniam VR, Brown JA, Mena I, Grant A, Bardina SV, et al. A
novel Zika virus mouse model reveals strain specific differences in virus pathogenesis and
host inflammatory immune responses. PLoS Pathog. 2017;13: e1006258.
doi:10.1371/journal.ppat.1006258
15.   Xia H, Luo H, Shan C, Muruato AE, Nunes BTD, Medeiros DBA, et al. An
evolutionary NS1 mutation enhances Zika virus evasion of host interferon induction. Nat
Commun. 2018;9: 414. doi:10.1038/s41467-017-02816-2
16.   Meertens L, Labeau A, Dejarnac O, Cipriani S, Sinigaglia L, Bonnet-Madin L, et al.
Axl Mediates ZIKA Virus Entry in Human Glial Cells and Modulates Innate Immune
Responses. Cell Rep. 2017;18: 324–333. doi:10.1016/j.celrep.2016.12.045
17.   Xu P, Shan C, Dunn TJ, Xie X, Xia H, Gao J, et al. Role of microglia in the
dissemination of Zika virus from mother to fetal brain. PLoS Negl Trop Dis. 2020;14.
doi:10.1371/journal.pntd.0008413
18.   Wang J, Liu J, Zhou R, Ding X, Zhang Q, Zhang C, et al. Zika virus infected primary
microglia impairs NPCs proliferation and differentiation. Biochem Biophys Res Commun.
2018;497: 619–625. doi:10.1016/j.bbrc.2018.02.118
19.   Savidis G, Perreira JM, Portmann JM, Meraner P, Guo Z, Green S, et al. The IFITMs
Inhibit Zika Virus Replication. Cell Rep. 2016;15: 2323–30. doi:10.1016/j.celrep.2016.05.074
20.   Monel B, Compton AA, Bruel T, Amraoui S, Burlaud-Gaillard J, Roy N, et al. Zika
virus induces massive cytoplasmic vacuolization and paraptosis-like death in infected cells.
EMBO J. 2017;36: 1653–1668. doi:10.15252/embj.201695597

21.   Schoggins JW, Wilson SJ, Panis M, Murphy MY, Jones CT, Bieniasz P, et al. A
diverse range of gene products are effectors of the type I interferon antiviral response. Nature.
2011;472: 481–5. doi:10.1038/nature09907
22.   Liu S-Y, Sanchez DJ, Aliyari R, Lu S, Cheng G. Systematic identification of type I
and type II interferon-induced antiviral factors. PNAS. 2012;109: 4239–4244.
23.   Groth-Pedersen L, Aits S, Corcelle-Termeau E, Petersen NHT, Nylandsted J, Jäättelä
M. Identification of cytoskeleton-associated proteins essential for lysosomal stability and
survival of human cancer cells. PLoS One. 2012;7: e45381.
doi:10.1371/journal.pone.0045381
24.   Hackett BA, Cherry S. Flavivirus internalization is regulated by a size-dependent
endocytic pathway. Proc Natl Acad Sci U S A. 2018;115: 4246–4251.
doi:10.1073/pnas.1720032115
25.   Paul A, Tang TH, Ng SK. Interferon Regulatory Factor 9 Structure and Regulation.
Front Immunol. 2018;9. doi:10.3389/fimmu.2018.01831
26.   Smith S, Weston S, Kellam P, Marsh M. IFITM proteins-cellular inhibitors of viral
entry. Curr Opin Virol. 2014;4: 71–77. doi:10.1016/j.coviro.2013.11.004
27.   Lai AY, Wade PA. NuRD: A multi-faceted chromatin remodeling complex in
regulating cancer biology. Nat Rev Cancer. 2011;11: 588–596. doi:10.1038/nrc3091
28.   Blight KJ, McKeating JA, Rice CM. Highly permissive cell lines for subgenomic and
genomic hepatitis C virus RNA replication. J Virol. 2002;76: 13001–14.
29.   Sumpter R, Loo YM, Foy E, Li K, Yoneyama M, Fujita T, et al. Regulating
intracellular antiviral defense and permissiveness to hepatitis C virus RNA replication
through a cellular RNA helicase, RIG-I. J Virol. 2005;79: 2689–99.
doi:10.1128/JVI.79.5.2689-2699.2005
30.   Mukherjee A, Di Bisceglie AM, Ray RB. Hepatitis C virus-mediated enhancement of
microRNA miR-373 impairs the JAK/STAT signaling pathway. J Virol. 2015;89: 3356–65.
doi:10.1128/JVI.03085-14
31.   Li Y, Ma MX, Qin B, Lin L-T, Richardson CD, Feld J, et al. The Ubiquitin-Specific
Protease 18 Promotes Hepatitis C Virus Production by Increasing Viral Infectivity. Mediators
of Inflammation. 2019;2019: e3124745. doi:10.1155/2019/3124745
32.   Randall G, Chen L, Panis M, Fischer AK, Lindenbach BD, Sun J, et al. Silencing of
USP18 Potentiates the Antiviral Activity of Interferon Against Hepatitis C Virus Infection.
Gastroenterology. 2006;131: 1584–1591. doi:10.1053/j.gastro.2006.08.043
33.   Arnaud N, Dabo S, Akazawa D, Fukasawa M, Shinkai-Ouchi F, Hugon J, et al.
Hepatitis C virus reveals a novel early control in acute immune response. PLoS Pathog.
2011;7: e1002289. doi:10.1371/journal.ppat.1002289 PPATHOGENS-D-11-00682 [pii]
34.   Chen L, Sun J, Meng L, Heathcote J, Edwards AM, McGilvray ID. ISG15, a
ubiquitin-like interferon-stimulated gene, promotes hepatitis C virus production in vitro:
implications for chronic infection and response to treatment. J Gen Virol. 2010;91: 382–388.
doi:10.1099/vir.0.015388-0
35.   Pfaender S, Mar KB, Michailidis E, Kratzel A, Boys IN, V'kovski P, et al. LY6E
impairs coronavirus fusion and confers immune control of viral disease. Nature Microbiology.
2020;5: 1330–1339. doi:10.1038/s41564-020-0769-y
36.   Vanhollebeke B, Pays E. The function of apolipoproteins L. Cell Mol Life Sci.
2006;63: 1937–1944. doi:10.1007/s00018-006-6091-x
37.   Monajemi H, Fontijn RD, Pannekoek H, Horrevoets AJG. The apolipoprotein L gene
cluster has emerged recently in evolution and is expressed in human vascular tissue.
Genomics. 2002;79: 539–546. doi:10.1006/geno.2002.6729
38.   Malakhova OA, Kim KI, Luo J-K, Zou W, Kumar KGS, Fuchs SY, et al. UBP43 is a
novel regulator of interferon signaling independent of its ISG15 isopeptidase activity. EMBO

J. 2006;25: 2358–2367. doi:10.1038/sj.emboj.7601149

39.   Alcendor DJ. Zika Virus Infection of the Human Glomerular Cells: Implications for
Viral Reservoirs and Renal Pathogenesis. J Infect Dis. 2017;216: 162–171.
doi:10.1093/infdis/jix171

40.   Combredet C, Labrousse V, Mollet L, Lorin C, Delebecque F, Hurtrel B, et al. A
Molecularly Cloned Schwarz Strain of Measles Virus Vaccine Induces Strong Immune
Responses in Macaques and Transgenic Mice. Journal of Virology. 2003;77: 11546–11554.
doi:10.1128/JVI.77.21.11546-11554.2003

41.   Uzureau S, Lecordier L, Uzureau P, Hennig D, Graversen JH, Homblé F, et al.
APOL1 C-Terminal Variants May Trigger Kidney Disease through Interference with APOL3
Control of Actomyosin. Cell Rep. 2020;30: 3821-3836.e13. doi:10.1016/j.celrep.2020.02.064

42.   Waugh MG. The Great Escape: how phosphatidylinositol 4-kinases and PI4P promote
vesicle exit from the Golgi (and drive cancer). Biochem J. 2019;476: 2321–2346.
doi:10.1042/BCJ20180622

43.   Delang L, Paeshuyse J, Neyts J. The role of phosphatidylinositol 4-kinases and
phosphatidylinositol 4-phosphate during viral replication. Biochem Pharmacol. 2012;84:
1400–1408. doi:10.1016/j.bcp.2012.07.034

44.   Strating JR, van Kuppeveld FJ. Viral rewiring of cellular lipid metabolism to create
membranous replication compartments. Curr Opin Cell Biol. 2017;47: 24–33.
doi:10.1016/j.ceb.2017.02.005

45.   Daboussi L, Costaguta G, Ghukasyan R, Payne GS. Conserved role for Gga proteins
in phosphatidylinositol 4-kinase localization to the trans-Golgi network. PNAS. 2017;114:
3433–3438.

46.   Rutaganira FU, Fowler ML, McPhail JA, Gelman MA, Nguyen K, Xiong A, et al.
Design and Structural Characterization of Potent and Selective Inhibitors of
Phosphatidylinositol 4 Kinase IIIβ. J Med Chem. 2016;59: 1830–1839.
doi:10.1021/acs.jmedchem.5b01311

47.   Hsu N-Y, Ilnytska O, Belov G, Santiana M, Chen Y-H, Takvorian PM, et al. Viral
reorganization of the secretory pathway generates distinct organelles for RNA replication.
Cell. 2010;141: 799–811. doi:10.1016/j.cell.2010.03.050

48.   Martín-Acebes MA, Blázquez A-B, Jiménez de Oya N, Escribano-Romero E, Saiz J-
C. West Nile virus replication requires fatty acid synthesis but is independent on
phosphatidylinositol-4-phosphate lipids. PLoS One. 2011;6: e24970.
doi:10.1371/journal.pone.0024970

49.   Jiang D, Weidner JM, Qing M, Pan XB, Guo H, Xu C, et al. Identification of five
interferon-induced cellular proteins that inhibit west nile virus and dengue virus infections. J
Virol. 2010;84: 8332–41. doi:10.1128/JVI.02199-09

50.   Itsui Y, Sakamoto N, Kurosaki M, Kanazawa N, Tanabe Y, Koyama T, et al.
Expressional screening of interferon-stimulated genes for antiviral activity against hepatitis C
virus replication. J Viral Hepat. 2006;13: 690–700. doi:10.1111/j.1365-2893.2006.00732.x

51.   Van der Hoek KH, Eyre NS, Shue B, Khantisitthiporn O, Glab-Ampi K, Carr JM, et
al. Viperin is an important host restriction factor in control of Zika virus infection. Sci Rep.
2017;7: 4475. doi:10.1038/s41598-017-04138-1

52.   Richardson RB, Ohlson MB, Eitson JL, Kumar A, McDougal MB, Boys IN, et al. A
CRISPR screen identifies IFI6 as an ER-resident interferon effector that blocks flavivirus
replication. Nat Microbiol. 2018. doi:10.1038/s41564-018-0244-1

53.   Li L, Zhao H, Liu P, Li C, Quanquin N, Ji X, et al. PARP12 suppresses Zika virus
infection through PARP-dependent degradation of NS1 and NS3 viral proteins. Sci Signal.
2018;11: eaas9332. doi:10.1126/scisignal.aas9332

54.   Wu Y, Yang X, Yao Z, Dong X, Zhang D, Hu Y, et al. C19orf66 interrupts Zika virus

replication by inducing lysosomal degradation of viral NS3. PLOS Neglected Tropical
Diseases. 2020;14: e0008083. doi:10.1371/journal.pntd.0008083

55. Vanwalscappel B, Gadea G, Desprès P. A Viperin Mutant Bearing the K358R
Substitution Lost its Anti-ZIKA Virus Activity. Int J Mol Sci. 2019;20: E1574.
doi:10.3390/ijms20071574

56. Lindqvist R, Kurhade C, Gilthorpe JD, Överby AK. Cell-type- and region-specific
restriction of neurotropic flavivirus infection by viperin. Journal of Neuroinflammation.
2018;15: 80. doi:10.1186/s12974-018-1119-3

57. Perng Y-C, Lenschow DJ. ISG15 in antiviral immunity and beyond. Nat Rev
Microbiol. 2018;16: 423–439. doi:10.1038/s41579-018-0020-5

58. Honke N, Shaabani N, Zhang D-E, Hardt C, Lang KS. Multiple functions of USP18.
Cell Death Dis. 2016;7: e2444–2444. doi:10.1038/cddis.2016.326

59. Huang X, Sun S, Wang X, Fan F, Zhou Q, Lu S, et al. Mechanistic insights into the
SNARE complex disassembly. Science Advances. 2019;5: eaau8164.
doi:10.1126/sciadv.aau8164

60. Jahn R, Scheller RH. SNAREs — engines for membrane fusion. Nat Rev Mol Cell
Biol. 2006;7: 631–643. doi:10.1038/nrm2002

61. Pirooz SD, He S, Zhang T, Zhang X, Zhao Z, Oh S, et al. UVRAG is required for
virus entry through combinatorial interaction with the class C-Vps complex and SNAREs.
Proc Natl Acad Sci U S A. 2014;111: 2716–2721. doi:10.1073/pnas.1320629111

62. Langston PK, Nambu A, Jung J, Shibata M, Aksoylar HI, Lei J, et al. Glycerol
phosphate shuttle enzyme GPD2 regulates macrophage inflammatory responses. Nat
Immunol. 2019;20: 1186–1195. doi:10.1038/s41590-019-0453-7

63. Schoggins JW. Interferon-Stimulated Genes: What Do They All Do? Annual Review
of Virology. 2019;6: 567–584. doi:10.1146/annurev-virology-092818-015756

64. Tartour K, Nguyen XN, Appourchaux R, Assil S, Barateau V, Bloyet LM, et al.
Interference with the production of infectious viral particles and bimodal inhibition of
replication are broadly conserved antiviral properties of IFITMs. PLoS Pathog. 2017;13:
e1006610. doi:10.1371/journal.ppat.1006610

65. García-Sastre A. Ten Strategies of Interferon Evasion by Viruses. Cell Host Microbe.
2017;22: 176–184. doi:10.1016/j.chom.2017.07.012

66. Schulz KS, Mossman KL. Viral Evasion Strategies in Type I IFN Signaling – A
Summary of Recent Developments. Front Immunol. 2016;7. doi:10.3389/fimmu.2016.00498

67. Schoggins JW, Macduff DA, Imanaka N, Gainey MD, Shrestha B, Eitson JL, et al.
Pan-viral specificity of IFN-induced genes reveals new roles for cGAS in innate immunity.
Nature. 2013. doi:10.1038/nature12862

68. Taylor HE, Khatua AK, Popik W. The innate immune factor apolipoprotein L1
restricts HIV-1 infection. J Virol. 2014;88: 592–603. doi:10.1128/JVI.02828-13

69. Yamane D, Feng H, Rivera-Serrano EE, Selitsky SR, Hirai-Yuki A, Das A, et al.
Basal expression of interferon regulatory factor 1 drives intrinsic hepatocyte resistance to
multiple RNA viruses. Nat Microbiol. 2019;4: 1096–1104. doi:10.1038/s41564-019-0425-6

70. Burke JE, McPhail JA, Rathinaswamy M, Jenkins ML. Defining how viruses
manipulate lipid phosphoinositides through activation of PI4P kinases to mediate viral
replication. The FASEB Journal. 2020;34: 1–1. doi:10.1096/fasebj.2020.34.s1.02060

71. Gillespie LK, Hoenen A, Morgan G, Mackenzie JM. The Endoplasmic Reticulum
Provides the Membrane Platform for Biogenesis of the Flavivirus Replication Complex.
Journal of Virology. 2010;84: 10438–10447. doi:10.1128/JVI.00986-10

72. Olias P, Etheridge RD, Zhang Y, Holtzman MJ, Sibley LD. Toxoplasma Effector
Recruits the Mi-2/NuRD Complex to Repress STAT1 Transcription and Block IFN-γ-
Dependent Gene Expression. Cell Host Microbe. 2016;20: 72–82.

doi:10.1016/j.chom.2016.06.006

73.   Nusinzon I, Horvath CM. Interferon-stimulated transcription and innate antiviral
immunity require deacetylase activity and histone deacetylase 1. PNAS. 2003;100: 14742–
14747. doi:10.1073/pnas.2433987100

74.   Vlasáková J, Nováková Z, Rossmeislová L, Kahle M, Hozák P, Hodný Z. Histone
deacetylase inhibitors suppress IFNα-induced up-regulation of promyelocytic leukemia
protein. Blood. 2007;109: 1373–1380. doi:10.1182/blood-2006-02-003418

75.   Saleem MA, O'Hare MJ, Reiser J, Coward RJ, Inward CD, Farren T, et al. A
conditionally immortalized human podocyte cell line demonstrating nephrin and podocin
expression. J Am Soc Nephrol. 2002;13: 630–638. doi:10.1681/ASN.V133630

76.   Nakabayashi H, Taketa K, Miyano K, Yamane T, Sato J. Growth of human hepatoma
cells lines with differentiated functions in chemically defined medium. Cancer Res. 1982;42:
3858–3863.

77.   Buchrieser J, Dufloo J, Hubert M, Monel B, Planas D, Rajah MM, et al. Syncytia
formation by SARS-CoV-2-infected cells. EMBO J. 2021;40: e107405.
doi:10.15252/embj.2020107405

78.   Chazal M, Beauclair G, Gracias S, Najburg V, Simon-Loriere E, Tangy F, et al. RIG-I
Recognizes the 5' Region of Dengue and Zika Virus Genomes. Cell Rep. 2018;24: 320–328.
doi:10.1016/j.celrep.2018.06.047

79.   Boukadida C, Marnata C, Montserret R, Cohen L, Blumen B, Gouttenoire J, et al. NS2
proteins of GB virus B and hepatitis C virus share common protease activities and membrane
topologies. J Virol. 2014;88: 7426–44. doi:10.1128/JVI.00656-14

80.   Wakita T, Pietschmann T, Kato T, Date T, Miyamoto M, Zhao Z, et al. Production of
infectious hepatitis C virus in tissue culture from a cloned viral genome. Nat Med. 2005;11:
791–6. doi:10.1038/nm1268

81.   Kaul A, Woerz I, Meuleman P, Leroux-Roels G, Bartenschlager R. Cell culture
adaptation of hepatitis C virus and in vivo viability of an adapted variant. J Virol. 2007;81:
13168–79. doi:10.1128/JVI.01362-07

82.   Aicher S, Kakkanas A, Cohen L, Blumen B, Oprisan G, Njouom R, et al. Differential
regulation of the Wnt/beta-catenin pathway by hepatitis C virus recombinants expressing core
from various genotypes. Sci Rep. 2018;8: 11185. doi:10.1038/s41598-018-29078-2

83.   Malo N, Hanley JA, Cerquozzi S, Pelletier J, Nadon R. Statistical practice in high-
throughput screening data analysis. Nat Biotechnol. 2006;24: 167–175. doi:10.1038/nbt1186

84.   Birmingham A, Selfors LM, Forster T, Wrobel D, Kennedy CJ, Shanks E, et al.
Statistical Methods for Analysis of High-Throughput RNA Interference Screens. Nat
Methods. 2009;6: 569–575. doi:10.1038/nmeth.1351

85.   McQuin C, Goodman A, Chernyshev V, Kamentsky L, Cimini BA, Karhohs KW, et
al. CellProfiler 3.0: Next-generation image processing for biology. PLOS Biology. 2018;16:
e2005970. doi:10.1371/journal.pbio.2005970

86.   Mutso M, Saul S, Rausalu K, Susova O, Žusinaite E, Mahalingam S, et al. Reverse
genetic system, genetically stable reporter viruses and packaged subgenomic replicon based
on a Brazilian Zika virus isolate. J Gen Virol. 2017;98: 2712–2724. doi:10.1099/jgv.0.000938

87.   Schwarz MC, Sourisseau M, Espino MM, Gray ES, Chambers MT, Tortorella D, et al.
Rescue of the 1947 Zika Virus Prototype Strain with a Cytomegalovirus Promoter-Driven
cDNA Clone. mSphere. 2016;1. doi:10.1128/mSphere.00246-16

   Figure legends

Figure 1. **A loss of function screen identified genes modulating ZIKV replication in IFN-**
stimulated human microglial cells. (A) Scheme summarizing the screen conditions. (B, C) Scatter
plots showing the rZ-score obtained in the 1st (B) and second analysis (C) of the 2 screens. The green
line represents the linear regression, as compared to the expected perfect correlation (dotted black line).
Antiviral and proviral hits are depicted in red and blue, respectively. (D) List of the antiviral and proviral
hits as identified by the 2 analysis of the 2 screens. Assessment of the antiviral (E, F) and proviral (G,
H) activities of some hits. HMC3 cells were transfected with either pool of 3 siRNAs against the
indicated candidate gene or non-targeting (NT) siRNAs, treated with IFNα2 (200U/mL) for 24 hours
and infected with ZIKV (at an MOI of 5 PFU/cell) for 24 hours. Control cells transfected with NT
siRNA in the absence of IFNα2 treatment (NT-IFN) are included. (E, G) Cell-associated viral RNA was
quantified by RT-qPCR and expressed as genome equivalents (GE) per μg of total cellular RNA. (F, H)
The number of cells positive for viral protein E was assessed by flow cytometry and are expressed
relatively to the NT+IFN control of each experiment. Data are means ± SD of three or four independent
experiments, *p<0.05, **p<0.01,***p<0.001, paired t-tests.

Figure 2. Effect of a selection of candidate genes on HCV and SARS-CoV-2 replication. (A-D).
Huh-7.5 cells were transfected with a pool of 3 siRNAs against selected candidates (antiviral and
proviral genes, as identified in the ZIKV screen (Fig. 1), are in red and blue, respectively) or non-
targeting (NT) siRNAs, treated with IFNα2 (200U/mL) for 24 hours, then infected with HCV at a MOI
of 3 TCID50/cell for 48 hours. (A, C) Cell-associated viral RNA was quantified by RT-qPCR and
expressed as genome equivalents (GE) per μg of total cellular RNA. (B, D) Release of infectious HCV
particles was determined by TCID50 assays. Data are expressed relatively to the NT+IFN control of
each experiment. Plotted values are expressed relative to mean NT+IF across experiements and
represent means ± SD of two independent experiments each in duplicates, *p<0.05,
p<0.01,*p<0.001, ****p<0.0001 paired t-tests. (E-H). A549-ACE2 cells were transfected with a
pool of 3 siRNAs against selected candidates (antiviral genes in red, proviral genes in blue) or non-
targeting (NT) siRNAs, treated with IFNα2 (200U/mL) for 24 hours and infected with SARS-CoV-2 at
a MOI of 2 for 24 hours. (E, G) Cell-associated viral RNA was quantified by RT-qPCR and expressed
as genome equivalents (GE) per μg of total cellular RNA. (F, H) The number of cells positive for the
viral protein spike (S) was assessed by flow cytometry Data are expressed relatively to the NT+IFN
control of each experiment. Data are means ± SD of triplicates of one experiment, *p<0.05,
p<0.01,*p<0.001, paired t-tests.

Figure 3. Effect of reduced expression of APOL1 and APOL3 on the replication of a panel of viruses in HMC3 cells and podocytes. (A, E) *APOL1* mRNA and *APOL3* mRNA abundance were quantified by RT-qPCR analysis in HMC3 cells or podocytes treated or not with IFNα2 (200U/mL) for 24 hours and expressed as copy numbers per μg of total cellular RNA. HMC3 cells (B) or podocytes (F) were transfected with pool of 3 siRNAs targeting *APOL1, APOL3* and *USP18* mRNAs or with non-targeting (NT) control siRNAs. The relative amounts of *APOL1, APOL3* and *USP18* mRNAs were determined by RT-qPCR analysis and were normalized to that of GAPDH mRNA. They are expressed as compared to abundance relative to cells transfected with control NT siRNAs. HMC3 cells (C) or podocytes (G) were transfected with the indicated siRNAs, treated, or not, with IFNa2 (200U/mL) for 24 hours, and infected with ZIKV for 24 hours. HMC3 cells were infected at a MOI of 2 and podocytes at a MOI of 1. The percentages of cells that were positive for the viral E proteins were determined by flow cytometric analysis. Data are expressed relatively to the siRNA NT control of each experiment. HMC3 cells (D) or podocytes (H) were treated with IFNα2 (200U/mL), transfected with the indicated siRNAs pools and subjected to Western blotting analysis with antibodies against the indicated proteins. (I) HMC3 cells were transfected with the indicated siRNAs pools, treated with IFNa2 (200U/mL) for 24 hours and infected with the indicated viruses for 18 to 24 hours, at the MOI indicated in the MM section. The percentages of the cells positive for viral proteins or GFP were determined by flow cytometric analysis. Data are means ± SD of three or four independent experiments, *p<0.05, **p<0.01,***p<0.001, paired t-tests.

Figure 4. APOL1 and APOL3 promote viral replication independently of their interaction with phosphoinositides. HMC3 cells were transfected with GFP-tagged versions of APOL1 and APOL3. Thirty hours later, they were stained with antibodies recognizing GM130 (A) or PI4KB (B) and with NucBlue to detect nuclei. Images are representative of numerous observations over 2 independent experiments. The white arrow shows an APOL1-GFP-positive vesicle. (C) HCM3 cells were treated with different concentrations of the PI4KB inhibitor and were stained for PI(4)P. (D) HMC3 cells treated with different doses of PI4KB inhibitor were infected with CVB3, WNV or ZIKV, in the presence or absence of IFNα2 (200U/mL). The percentages of the cells positive for viral proteins were determined by flow cytometric analysis. Data are means ± SD of three independent experiments, *p<0.05, **p<0.01,***p<0.001, one-way ANOVA.

Figure 5. MTA2 restricts ZIKV replication in IFNα2-stimulated cells. HMC3 (A, B, E, F, I) and Huh-7 cells (C, D, G, H, J) were transfected with pool of 3 siRNAs targeting *MTA2* or *IFNAR1* mRNAs or non-targeting (NT) control siRNAs, treated or not with IFNα2 (100U/mL) for 24 hours, and infected with ZIKV (MOI of 1 for HMC3 cells, MOI of 5 for Huh-7 cells) for 24 hours. (A-D) The relative amounts of *MTA2* and *IFNAR1* mRNAs were determined by RT-qPCR analysis and normalized to that

of *GAPDH* mRNA and siRNA-NT without IFN. (E, G) Cell-associated viral RNA was quantified by

RT-qPCR and expressed as genome equivalents (GE) per µg of total cellular RNA. (F, H) Number of

infected cells was assessed by staining of viral protein E and flow cytometry analysis. (I, J) Cells were

treated with IFNα2 (200U/mL) or left untreated, transfected with the indicated siRNAs pools and

subjected to Western blotting analysis with antibodies against the indicated proteins. Data are means ±

SD of three independent experiments, *p<0.05, **p<0.01,***p<0.001, paired t-tests.

1041

Figure 6. Effect of reduced expression of MTA2 on the replication of YFV, WNV, VSV and MeV-

GFP. Huh-7 cells were transfected with the indicated siRNAs pool, treated or not with IFNα2

(200U/mL) for 24 hours and infected with WNV (A) or YFV (B) for 24 hours, at the MOIs indicted in

the MM section. The percentages of the cells positive for viral protein Env was determined by flow

cytometric analysis. HMC3 cells were transfected with the indicated siRNAs pool, treated or not with

IFNα2 (200U/mL) for 24 hours and infected with VSV for 18 hours (C) or MeV-GFP for 24 hours (D),

at the MOIs indicted in the MM section. The percentages of the cells positive for viral protein G or GFP

were determined by flow cytometric analysis. Data are expressed relatively to the siRNA NT control of

each experiment. Data are means ± SD of three or four independent experiments, *p<0.05,

p<0.01,*p<0.001, paired t-tests.

1052

1053   **Supplementary figure legends**

1054

Figure S1. Quality control and reproducibility of the screens. (A) HMC3 cells were transfected with

either pool of 3 siRNAs against IFNAR or IFITM3 or non-targeting (NT) siRNAs, treated with IFNα2

(1000U/mL) for 24 hours and infected with ZIKV PF13 at a MOI of 7 PFU/cell for 24 hours. The

number of cells positive for viral protein E was assessed by confocal analysis using the pan-flavivirus

anti-E antibody 4G2. (B) Distribution of the "number of cells per 3 fields" parameter for each screen.

The values of the control wells (cells transfected with siRNA targeting KIF11) are shown in dark gray.

(C) Representation of the number of cells per 3 fields of screen 1 as a function of the screen 2. The green

line represents the linear regression as compared to the expected perfect correlation (dotted black line).

(D) Distribution of the "percentage of infected cells" parameter for each screen in the first analysis. The

values of the control conditions (cells transfected with siRNAs targeting IFNAR1 or IFNAR2) are

shown in dark grey. (E) Representation of the percentage of infected cells per well of screen 1 as a

function of screen 2, as identified by the first analysis. The green line represents the linear regression as

compared to the expected perfect correlation (dotted black line). (F) Representation of the percentage

of infected cells per well in the analysis 1 as a function of analysis 2. The green line represents the linear

regression. (G) HMC3 cells were transfected with pools of 3 siRNAs targeting the indicated genes or

with non-targeting (NT) control siRNAs. The relative abundances of the mRNAs of the candidate genes

were determined by RT-qPCR analysis and were normalized with respect to GAPDH mRNA level.

They are expressed relatively to abundance in cells transfected with NT siRNAs set at 1. Data are means

± SD of three or four independent experiments. ND: not determined due to mRNA levels below assay

threshold. The samples are the same than in Fig. 1E-H.

1075

Figure S2. Efficacy of the specific siRNAs. Huh-7.5 cells (A) or A549-ACE2 cells (B) were transfected

with pool of 3 siRNAs targeting the indicated genes or with non-targeting (NT) control siRNAs. The

relative abundances of the mRNAs of the candidate genes were determined by RT-qPCR analysis and

were normalized with respect to GAPDH mRNA level. Values are expressed relatively to abundance in

cells transfected with NT siRNA in each experiment, set at 1. Data are means ± SD of three or four

independent experiments. ND: not determined due to mRNA levels below assay threshold. The samples

are the same than in Fig. 2.

1083

1084 **Figure S3. Localization of GFP-tagged version of APOL1 and APOL3 in HMC3 cells**. Cells were

1085 transfected with GFP-tagged versions of APOL1 and APOL3. Thirty hours later, they were stained with

1086 antibodies recognizing EEA1 or CD63 and with NucBlue to detect nuclei. Images are representative of

1087 numerous observations over 2 independent experiments.

A

APOL1-GFP   GM130   merge   zoom

GFP-APOL3   GM130   merge   zoom

B

APOL1-GFP   PI4KB   merge   zoom

GFP-APOL3   PI4KB   merge   zoom

C

PI(4)P   DMSO   10nM   40nM   160nM

D

Ratio of infected cells

1.5

1.0

0.5

0.0

CVB3   WNV   ZIKV -IFN   ZIKV +IFN

DMSO
10 nM
40 nM
160 nM