Measuring cystic fibrosis drug responses in organoids derived from 2D differentiated

nasal epithelia

Gimano D. Amatngalim [1,2, *], Lisa W. Rodenburg [1,2], Bente L. Aalbers [3], Henriette H. M.

Raeven [1,2], Ellen M. Aarts [1,2], Iris A.L. Silva [4], Wilco Nijenhuis [5], Sacha Vrendenbarg [1,2],

Evelien Kruisselbrink [1,2], Jesse E. Brunsveld [1,2], Cornelis M. van Drunen [6], Sabine Michel [1],

Karin M. de Winter-de Groot [1], Harry G. Heijerman [3], Lukas C. Kapitein [5], Magarida D.

Amaral [4], Cornelis K. van der Ent [1], Jeffrey M. Beekman [1,2]

[1] Department of Pediatric Pulmonology, Wilhelmina Children's Hospital, University Medical

Center Utrecht, Utrecht University, member of ERN-LUNG, Utrecht, the Netherlands.

[2] Regenerative Medicine Center Utrecht, University Medical Center Utrecht, Utrecht

University, Utrecht, the Netherlands.

[3] Department of Pulmonology, University Medical Center Utrecht, Utrecht, the Netherlands.

[4] BioISI-Biosystems & Integrative Sciences Institute, Faculty of Sciences, University of

Lisboa, Lisboa, Portugal.

[5] Cell Biology, Neurobiology and Biophysics, Department of Biology, Faculty of Science,

Utrecht University, Utrecht, Netherlands.

[6] Department of Otorhinolaryngology, Amsterdam University Medical Centers, University of

Amsterdam, Amsterdam.

*Correspondence: G.D.Amatngalim@umcutrecht.nl

Running title: CFTR drug assay with airway organoids

Keywords: Cystic fibrosis, Cystic Fibrosis Transmembrane Conductance Regulator, Nasal

epithelium, organoids, Personalized medicine

Abstract

Cystic Fibrosis (CF) is caused by genetic defects that impair the cystic fibrosis

transmembrane conductance regulator (CFTR) channel in airway epithelial cells. These

defects may be overcome by specific CFTR modulating drugs, for which the efficacy can be

predicted in a personalized manner using 3D nasal-brushing-derived airway organoids in a

forskolin-induced swelling assay. Despite of this, previously described application of 3D

airway organoids in CFTR function assays have not been fully optimal. In this report we

therefore describe an alternative method of culturing nasal brushing-derived airway

organoids, which are created from an equally differentiated airway epithelial monolayer of a

2D air-liquid interface culture. In addition, we have defined organoid culture conditions, with

the growth factor/cytokine combination neuregulin-1β and interleukine-1β, which enabled

consistent detection of CFTR modulator responses in nasal airway organoids cultures from

subjects with CF.

Introduction

Cystic fibrosis (CF) is a monogenic epithelial disease caused by mutations in the cystic fibrosis conductance transmembrane regulator (*CFTR*) gene [1]. This defect impairs CFTR-dependent anion conductance in airway epithelia [2], which leads to a severe respiratory disease [3]. CFTR modulators are target-specific drugs that may restore CFTR function in individuals with CF [4]. However, the efficiency of modulators largely depends on the CFTR genotype of an individual with CF. More than 2,000 distinct CFTR mutations have been reported (http://www.genet.sickkids.on.ca//) with variable effects on CFTR expression or function. In addition to common mutations, such as the F508del allele, approximately 1,000 rare mutations have been identified that each affect less than 5 individuals worldwide. This low prevalence makes it unfeasible to determine CFTR modulator drug efficacy in large cohort clinical studies.

As an alternative of determining drug efficacy directly in individuals with CF, the effects of CFTR modulators can be predicted using patient-derived epithelial cultures in functional CFTR assays [5]. This is traditionally done with 2D air-liquid interface (ALI)-differentiated airway epithelia by assessment of CFTR-dependent chloride (Cl⁻) conductance via electrophysiology [6–8]. However, a major disadvantage of the ALI-culture model system is the limited scalability. In contrast, CFTR-expressing epithelial organoids from various tissues, i.e. airway, intestine, kidney, are emerging as a novel model system in which drug efficacy can be tested more efficiently in a mid-to high-throughput fashion [9–11]. Previously, we and others have shown that intestinal organoids from subjects with CF can be used to predict drug responses in a forskolin-induced swelling (FIS) assay, reflecting CFTR-dependent fluid secretion [12–15]. Nevertheless, based on the origin of CF respiratory disease, it remains

postulated that airway epithelial models are more predictive for determining CFTR modulator

responses.

Indeed, we previously reported CFTR modulator response measurements in long-term

expanded distal airway organoids using FIS [11]. However, we observed large variations in FIS

measurements, as swelling was limited to well-differentiated spherical structures. Others have

successfully used 3D nasal airway organoids (NAOs) derived from minimal-invasive nasal

brushings in functional CFTR assays [16–18], which are a more suitable option for personalized

drug testing compared to cultures derived from invasive (tracheo)bronchial and intestinal

tissues. However, previously described functional CFTR assays using NAOs were either low-

in throughput or required CFTR function measurements over large time periods [16–19].

Altogether, there is a remaining need for a further optimized and scalable FIS assay using

airway organoids, especially derived from nasal brushings, which enables efficient CFTR

modulator response measurements in subjects with CF. In this report, we describe an

alternative organoid culture method, in which NAOs are derived from 2D differentiated

human nasal epithelial cell (HNEC) monolayers. We furthermore describe optimized airway

organoid culture condition to improve CFTR modulator response measurements, by including

the growth factor/cytokine combination neuregulin-1β and IL-1β. Validation studies using

this culture condition showed consistent detection of genotype-specific responses to CFTR

modulators in FIS assays, including repairing effects of the FDA-approved CFTR triple

modulator therapy VX-661/VX-445/VX-770 [20].

Results and Discussion

We previously reported CFTR function measurements in distal airway organoids using a FIS

assay [11]. However, in this study we observed large variation in swelling between individual

organoids, which was restricted to large spherical structures. This is likely caused by the

unsynchronized differentiation of individual organoids, which may influence differentiation-

dependent CFTR function (Figure S1). To overcome this unsynchronized differentiation of

organoids within a culture, we set up a novel approach in which evenly differentiated airway

organoids are established from a 2D differentiated human nasal epithelial cells (HNEC)

monolayer (Figure 1A). In the organoid culture procedure, HNEC derived from nasal

brushings are first isolated and expanded in regular 2D cell cultures (Figure 1B). HNEC

stained positive for p63 and cytokeratin 5 (KRT5), confirming a basal stem cell phenotype

(Figure 1C). After expansion, HNEC were differentiation in conventional 2D air-liquid

interface (ALI) transwell-cultures, to recapitulate the mucociliary airway epithelium. Indeed,

well-differentiated ALI-HNEC cultures resembled the native nasal epithelial tissue, by

displaying a similar pseudostratified morphology and consisting of MUC5AC/CC10$^+$

secretory cell, β-tubulin IV$^+$ ciliated cells, and p63/KRT5$^+$ basal cells (Figure 1D and E).

CFTR function and effects of modulators were confirmed in ALI-HNEC from a healthy

control (HC) and individual with CF and a F508del/F508del genotype (Figure S2). During

isolation of airway organoids from resected airway tissues, based on the method described by

Sachs et al.[11], we observed that large epithelial fragments, obtained after collagenase

treatment, self-organized into well differentiated organoids within a few days after gel

embedding. Based on this observation and the assumption that ALI-cultures reflect the

differentiated airway epithelia tissue, we proposed that epithelial fragments from 2D-cultures

could be converted into 3D organoids as well. Indeed, embedding of differentiated ALI-

derived epithelial fragments in a 3D extracellular matrix led to formation of organoids with

clear lumens within 48 h (Figure 1F). From a single 12 mm^2 transwell insert we are able to

generate a yield of organoids that is sufficient for 48 independent wells (approx. 25-50

organoids/well) of a 96 well plate. In terms of scalability, this demonstrates a major advantage

of our method compared to the conventional use of 2D ALI-cultures. ALI-culture-derived

airway organoids directly displayed a differentiated phenotype. This was confirmed by visual

observation of beating cilia and accumulation of mucus (Video S1), as well as by

immunofluorescence imaging, demonstrating β-tubulin IV$^+$ cilia and MUC5AC$^+$ secretory

cells inside of the organoids, and p63$^+$/KRT5$^+$ basal cells at the basal side (Figure 1G, H).

Next, we determined whether ALI-derived NAOs could be used to measure CFTR function in

FIS assays in a 96 well plate format (Figure 2A). Forskolin stimulation of organoids from HC

subjects increased swelling in time, which was dose-dependent and reached a plateau at 5 μM

forskolin (Figure 2B-D). Organoid swelling could be attenuated with the NKKC1 inhibitor

bumetanide, demonstrating chloride-dependence of forskolin-induced organoid fluid secretion

(Figure S3A and B). Moreover, chemical CFTR inhibitors significantly reduced FIS in HC

NAOs (Figure 2E and F), indicating CFTR dependence. Upon comparison of cultures from

HC subjects and subject with CF, we observed that both HC and CF NAOs displayed cystic

lumens, which were not significant different in size (Figure 2G, H). This corresponds with

observations made in distal airway organoids [11], and suggests intrinsic CFTR-independent

fluid secretion mediated by alternative chloride channels. In support of this, stimulation with

the Ca^{2+}-activated Cl$^-$ channel (CaCC) activator E$_{act}$ induced organoid swelling in CF NAOs,

which was significant higher compared to HC cultures (Figure S3C and D). This suggests

dominance of CFTR-independent fluid secretion in CF NAOs, also observed by others [21]. In

contrast, FIS measured in NAOs from HC subjects was significantly higher compared to

subjects with CF (Figure 2I and J), reflecting differences in CFTR-dependent fluid secretion.

Although CF and HC NAOs could be distinguished phenotypically based on FIS, we initially

did not observe repairing effects of the CFTR corrector and potentiator combination VX-

809/VX-770 in CF F508del/F508del NAOs from multiple subjects (Figure S3E and F).

Therefore, we aimed to optimize the CFTR modulator response measurements, by modifying

the organoid culture conditions. Recent studies suggested that secretory cells are the primary

airway epithelial cell type mediating CFTR function [22,23]. Therefore, we examining a panel of

growth factors and cytokines that could modulate secretory cell functions, which were added

after plating of the epithelial fragments and during organoid culturing (Figure 3A). The

examined factors included, neuregulin-1β (NR), which has been reported to enhance epithelial

polarization and differentiation of secretory cells in ALI-cultures [24,25]. Moreover, the effect of

the pro-inflammatory cytokine IL-1β was examined, which has been shown to enhance goblet

cell differentiation, CFTR mRNA expression, chloride conductance, and CFTR modulator

responses in ALI-cultures [26–30]. In contrast to independent factors, a combination of NR/IL-1β

led in detection of VX-809/VX-770 modulator responses in CF F508del/F508del NAOs

(Figure 3B and Figure S4A). NR/IL-1β-cultured HC NAOs displayed a similar swelling

response compared to control culture conditions (Figure 3C and S4B). In support of reducing

CFTR-independent fluid secretion, NR/IL-1β attenuated lumen formation in CF NAOs, as

quantitated by organoids size measurements (Figure S4C and D). Moreover, in line with

enhanced specificity of CFTR function, NR/IL-1β increased and reducing the expression of

CFTR and the CaCC ANO1 respectively (Figure S4E). The Cl⁻ channel SLC26A9 was also

increased upon stimulation with NR/IL-1β, which may support CFTR function, as proposed

by others [31]. NR/IL-1β did not affect the expression of mucociliary differentiation markers

(Figure S4F and G).

We further validated CFTR function and modulator response measurements in the NR/IL-1β

culture condition. HC NAOs displayed a significant higher swelling response when compared

to cultures from CF F508del/F508del subjects (Figure 3D, F, and S5A). Moreover, the CFTR

modulator combination VX-809/VX-770 consistently enhanced FIS measured in CF

F508del/F508del NAOs from 7 independent subjects (Figure 3E, F, and S5B). In addition to

VX-809, FIS responses in CF F508del/F508del NAOs were modulated upon treatment with

other CFTR correctors (Figure S5C). We were also able to detect VX-770 potentiator

responses in NAOs from subjects with CF and a S1251N gating mutation (Figure 3G, H, and

S5D). Moreover, the highly effective CFTR triple modulator therapy VX-445/VX-661/VX-

770 [20], induced a high increase in FIS measurements in CF F508del/F508del NAOs cultured

with NR/IL-1β (Figure 3I, J, and S5E). VX-445 by itself did not increase FIS, whereas

chemical CFTR inhibition completely diminished increases in swelling by VX-445/VX-

661/VX-770, demonstrating CFTR-dependence of the modulator treatment effect (Figure S5F

and G).

In summary, we described a new method of culturing nasal brushing-derived airway

organoids.

Previously we have shown that long-term expanded airway organoids can be cultured as 2D

differentiated ALI-cultures [11]. In this report, we demonstrate further flexibility between 2D

and 3D airway culture models, by showing the possibility to convert 2D differentiated ALI-

cultures into 3D airway organoids. We furthermore described organoid culture conditions that

improved quantification of CFTR modulating drug responses in FIS assays. The NR/IL-1β

culture condition reflects the chronically inflamed airway epithelium of individuals with CF,

and therefore may act as physiological conditions for testing CFTR modulator responses in

NAOs. Further research is required to determine whether NR/IL-1β also improve CFTR

modulator responses in other airway model systems, such as distal airway organoids, or other

CFTR-expressing epithelial cells. In both the initially tested organoid culture conditions and

the optimized conditions with NR/IL-1β, we could not discriminate HC from CF NAOs based

on lumen size, which is in contrast to the CFTR-dependent intestinal organoid model [9,13].

Furthermore, due to limited numbers of donors with diverse CFTR genotypes, we were

unable to determine whether residual FIS correlated to disease severity, as shown by others [18].

In recent studies, other have shown that CFTR modulator responses in 3D nasal cultures

correlate with drug responsiveness in individuals with CF [19,32]. Whether CFTR drug responses

measurements in our model correlate with clinical outcome measurements will be determined

in follow up studies. CFTR function measurements in ALI-culture derived NAOs may also be

used to examine novel target-specific therapies for subjects with CF with unmet, such as

assessment of CRISPR-gene editing, read-through agents, or compounds targeting nonsense

mediated decay. Moreover, complementary to the widely used 2D ALI-cultures, airway

organoids derived from this model may be further used to study other respiratory tract

disorders.

Materials and Methods

Reagents and Tools Table

Reagent or Resource	Source	Identifier
Antibodies		
Mouse IgG1 anti-β-tubulin IV	Emergo Biogenex	#MU178-UC
Rabbit anti-β-tubulin IV	Abcam	#ab179509
Rabbit anti-MUC5AC	Abcam	#ab198294
Rabbit anti-P63	Abcam	#ab124762
Mouse IgG1 anti cytokeratin 5	Abcam	#ab17130
Mouse IgG1 anti-CC10	Acris, Origene	#AM26360PU-N
Goat anti-Mouse IgG1, Alexa Fluor 488	Invitrogen	#A-21121
Goat anti-Mouse IgG1, Alexa Fluor	Invitrogen	#A-21240

647		
Goat anti-Rabbit IgG, Alexa Fluor 488	Invitrogen	#A-11034
Goat anti-Rabbit IgG, Alexa Fluor 546	Invitrogen	#A-11035
Biological Samples		
Nasal brushings	UMC Utrecht	Protocol ID: 16/586, NL54885.041.16
Nasal turbinate tissue	Academic Medical Center Amsterdam	N/A
Chemicals, Peptides, and Recombinant Proteins		
TrypLE express enzyme	Fisher scientific	# 12605010
Sputolysin	Calbiochem	#560000-10
Collagen IV	Sigma-Aldrich	#C7521
CryoStor® CS10	StemCell Technologies	#07930
PureCol	Advanced BioMatrix	#5005
Bronchial epithelial cell medium-basal (BEpiCM-b)	Sciencell	#3211
Advanced DMEM F12	Thermo Fisher	#12634-028
B-27 Supplement, serum free	Thermo Fisher	#12587010

GlutaMAX Supplement	Thermo Fisher	#35050-061
HEPES	Thermo Fisher	#15630080
(±)-Epinephrine hydrochloride	Sigma-Aldrich	#E4642
Hydrocortisone	Sigma-Aldrich	#H0888
3,3',5-Triiodo-L-thyronine sodium salt	Sigma-Aldrich	#T6397
N-Acetyl-L-cysteine	Sigma-Aldrich	#A9165
Nicotinamide	Sigma-Aldrich	#N0636
SB02190	Sigma-Aldrich	#S7067
DMH-1	Selleck Chemicals	#S7146
A83-01	Tocris	#2939/10
Y-27632	Selleck Chemicals	#S1049
DAPT	Fisher Scientific	#15467109
TTNPB	Cayman	#16144-1
Recombinant human FGF-7	Peprotech	#100-19
Recombinant human FGF-10	Peprotech	#100-26
Recombinant human EGF	Peprotech	#AF-100-15
Recombinant human HGF	Peprotech	#100-39H
Recombinant Neuregulin-1β	Peprotech	#100-03

Recombinant Interleukin 1β	Peprotech	#200-01B
Penicillin-Streptomycin	Thermo Fisher	#15070-063
Primocin	Invivogen	#ant-pm-2
Amphotericin B	Thermo Fisher	#15290018
Gentamicin	Sigma-Aldrich	#G1397
Vancomycin	Sigma-Aldrich	#SBR00001
Collagenase type II	Thermo Fisher	#17101-015
CFTRinh-172	Sigma-Aldrich	#C2992
GlyH101	Sigma-Aldrich	#219671
VX-809	Selleck Chemicals	# S1565
VX-661	Selleck Chemicals	#S7059
VX-770	Selleck Chemicals	#S1144
Correctors C1-18	Cystic Fibrosis Foundation Therapeutics	https://www.cff.org/Research/Researcher-Resources/Tools-and-Resources/CFTR-Chemical-Compound-Program/
Forskolin	Sigma-Aldrich	#F3917
3-Isobutyl-1-methylxanthine	Sigma-Aldrich	# I5879
Amiloride	Sigma-Aldrich	#1019701
Calcein green acetoxymethyl (AM)	Invitrogen	#C34852
Alexa Fluor 555 Phalloidin	Invitrogen	#A34055
Phalloidin-iFluor	Abcam	#ab176752

405		
Prolong Gold antifade reagent	Thermo Fischer	#P36934
Prolong Gold antifade reagent with DAPI	Thermo Fischer	#P36935
iQ SYBR Green Supermix	Bio-Rad	#1708880
Critical Commercial Assays		
RNeasy Mini Kit	Qiagen	#74104
iScript cDNA synthesis kit	Bio-Rad	#1708891
Experimental Models: Cell Lines		
Human nasal airway epithelial cells (HNEC)	This paper	N/A
Hek293T – R-spondin-1 mFc cell line	Trevigen Cat# 3710-001-	N/A
Oligonucleotides		
Primers used for this manuscript		Table S4
Equipment		
Zeiss LSM800 confocal microscopy	Zeiss	N/A
Leica SP8X confocal microscope	Leica	N/A

Leica THUNDER imager	Leica	N/A
Leica TCS SP8 STED 3X microscope	Leica	N/A
Ussing chamber system	Physiologic Instruments	N/A
Voltage clamp	World Precision Instruments	#DVC-1000
PowerLab	AD Instruments	#8/30
Epithelial voltohmmeter EVOM2	World Precision Instruments	N/A
Nanodrop spectrophotometer	Thermo Fisher	N/A
CFX96 real-time detection machine	Bio-Rad	N/A
Software and Algorithms		
Zen Blue Software	Zeiss	https://www.zeiss.com/microscopy/int/products/microscope-software/zen.html
Prism 8	GraphPad Software Inc.	https://www.graphpad.com/scientific-software/prism/
Microsoft Excel	Microsoft Corporation	https://office.microsoft.com/excel
Adobe Illustrator	Adobe	https://www.adobe.com/nl/products/illustrator.html
LabChart 6	AD Instruments	https://labchart.software.informer.com/6.0/
CFX Manager 3.1	BioRad	https://www.bio-rad.com/en-us/sku/1845000-cfx-manager-software?ID=1845000
LAS X software	Leica	https://www.leica-microsystems.com/products/microscope-

		software/p/leica-las-x-ls/
ImageJ/FIJI		https://imagej.net/Fiji/Downloads
Others		
Cytological brush	CooperSurgical	#C0004
12 mm Transwell® with 0.4 µm Pore Polyester Membrane Insert	Corning	#3460
6.5 mm Transwell® with 0.4 µm Pore Polyester Membrane Insert	Corning	#3470
Matrigel	Corning	#354230
BC isolation and expansion medium		Table S1
Air-liquid interface differentiation medium		Table S2
Airway organoid culture medium		Table S3

Methods and Protocols

Patient materials and sample collection

Nasal brushings were collected from healthy volunteers without respiratory tract symptoms (n=22 independent donors), and subjects with CF (n=22 independent donors) by a trained research nurse or physician, essentially as previously described [33]. Sampling of adults was

conducted using a cytological brush and without anesthetics. Nasal brushings of infants (<6

years old) were collected with a modified interdental brush [34]. Samples were taken from the

inferior turbinates of both left and right nostrils and stored in collection medium, consisting of

advanced (ad)DMEM/F12, with GlutaMAX (1% v/v), HEPES (10 mM), Penicillin-

Streptomycin (1% v/v), and Primocin (100 µg/mL). Nasal brushings were collected and stored

with informed consent of all participants and was approved by a specific ethical board for the

use of biobanked materials TcBIO (Toetsingscommissie Biobanks), an institutional Medical

Research Ethics Committee of the University Medical Center Utrecht (protocol ID: 16/586).

Nasal samples from infants with CF were collected as part of the Precision study (protocol ID:

NL54885.041.16), which was approved by the Medical Research Ethics Committee of the

University Medical Center Utrecht (Utrecht, The Netherlands). Resected inferior nasal

turbinate tissue was obtained from a subject who underwent corrective surgery for turbinate

hypertrophy at the Academic Medical Center in Amsterdam, the Netherlands. The tissue was

accessible for research within the framework of patient care, in accordance with the "Human

Tissue and Medical Research: Code of conduct for responsible use" (2011) (www.federa.org),

describing the no-objection system for coded anonymous further use of such tissue without

necessary written or verbal consent.

Isolation and expansion of human nasal airway epithelial cells as 2D-cultures

Nasal cells were dissociated from the brush in the collection medium by scraping through a

sterile P1000 pipette tip with the top cut off. After centrifugation (400 g for 5 min), the pellet

was treated with TrypLE express enzyme, supplemented with 1x Sputolysin. The cells were

incubated for 10 min at 37 ºC and strained using a 100 µm strainer. After centrifugation (400

g for 5 min), the remaining pellet was used for isolation of human nasal epithelial cells

(HNEC). HNEC were isolated and expanded on 6-well culture plates coated with collagen IV

(50 µg/mL) and using BC isolation and expansion medium (Table S1) respectively. BC

isolation medium contained additional antibiotics to suppress microbial outgrowth and was

used during the first week of epithelial cell isolation, before switching to BC expansion

medium. BC expansion medium included the gamma-secretase inhibitor DAPT, which

reduces outgrowth of squamous cells at late passages. Growth factors (FGF7, FGF10, EGF,

and HGF) were added freshly to the culture medium. Cultures were refreshed three times a

week (Monday, Wednesday, Friday). After reaching 80-90% confluency, cells were passaged

with TrypLE express enzyme and further expanded for one passage before freezing with

CryoStor® CS10, supplemented with Y-27632 (5 µM) to create a master cell bank. For

further use, a work cell bank was created of passage 2 cells, which was cryo-stored.

Differentiation of 2D ALI-HNEC cultures

Expanded HNEC (passage =4-6) were cultured on transwell inserts (0.4 µm pore size

polyester membrane), which were coated with PureCol (30 µg/mL). Cells were seeded in a

density of 0.2 or $0.5*10^6$ cells on 24 or 12 wells inserts respectively and cultured in

submerged conditions in BC expansion medium until reaching confluency. Afterwards,

culture medium was changed with air-liquid interface (ALI)-differentiation medium (Table

S2) supplemented with A83-01 (500 nM), and cells were additionally cultured in submerged

condition for 1-2 days. Subsequently, culture medium at the apical side was removed and

cells were further differentiated as ALI-cultures. After 3-4 days, cells were refreshed with

ALI-diff medium without additional A83-01 and differentiated for at least 14 additional days

at ALI-conditions. Medium was refreshed twice a week (Monday and Thursday or Tuesday

and Friday), and the apical side of the cultures was washed with PBS once a week.

Conversion of 2D differentiated ALI cultures into airway organoids

Differentiated ALI-cultures were washed at the apical surface with PBS and subsequently treated at the basolateral side with collagenase type II (1 mg/mL) diluted in adDMEM/F12. Cultures were incubated at 37 ºC and 5% CO_2 for 45-60 min until the epithelium detaches from the transwell insert. Next, the dissociated epithelial layer was transferred to a 15 ml tube in 1 ml adDMEM/F12 + 10 % (v/v) Fetal bovine serum (FBS), mechanically disrupted into smaller fragment by pipetting, and strained with a 100 µm filter. After centrifugation (at 400 g, 5 min), the epithelial pellet was resuspended in ice-cold 75% growth factor reduced Matrigel (v/v in airway organoid (AO) medium Table S3). Next, epithelial fragments were embedded in 30 µl Matrigel droplets on pre-warmed 24-well suspension plates. Droplets were solidified at 37 ºC and 5% CO_2 for 20-30 min, before adding 0.5 ml AO medium (Table S3). In optimized conditions for measuring CFTR modulator responses, AO culture medium was further supplemented with neuregulin-1β (NR, 0.5 nM) and interleukin-1β (IL-1β; 10 ng/ml). Besides, NR/IL-1β we furthermore examined the effects of culturing with other interleukins i.e. IL-13, IL-4, IL-10, and the growth factors: fibroblast growth factor 2, 7, 10, hepatocyte growth factor, and insulin-like growth factor 1, which did not improve the detection of CFTR modulator responses. AO medium was refreshed twice a week (Monday and Thursday or Tuesday and Friday).

Forskolin-induced swelling (FIS) assay

ALI-derived NAOs were used in forskolin-induced swelling (FIS) assays, essentially as previously described with minor adaptation [11,35]. In short, organoids were transferred 3-5 days after conversion in 96-well plates in 4 µl droplets of 75% matrigel (v/v in AO medium),

containing approx. 25-50 structures. After solidification of droplets, 100 µl AO medium was

added to each well. In optimized conditions for measuring CFTR modulator responses, AO

culture medium was supplemented with neuregulin-1β (0.5 nM) and interleukin-1β (10

ng/ml). organoid swelling was conducted in quadruplicates. In indicated experiments, NAOs

were pre-treated with CFTRinh-172 and GlyH101 (CFTRi, both 50 µM) or vehicle as

negative control, for 4 h. CFTR correctors: VX-809, VX-661 (both 10 µM), VX-445 (5 µM),

C1-18 (all 10 µM) or vehicle were pre-treated for 48 h. Before assessment of FIS, NAOs were

stained with Calcein green AM (3 µM) for 30 min. Afterwards, organoids were stimulated

with forskolin with indicated concentration. In cultures from subjects with CF, the CFTR

potentiator VX-770 (10 µM) or vehicle was added together with forskolin. Swelling of NAOs

was quantitated by measuring the increase of the total area of calcein green AM-stained

organoids in a well during 15-min time intervals for a period of 2 h. Images were acquired

with a Zeiss LSM800 confocal microscopy, using a 2.5 or 5x objective and experiments were

conducted at 37°C and 95% O_2/5% CO_2 to maintain a pH of 7.4. Data was analysed using Zen

Blue Software and Prism 8.

Ussing chamber experiments

For open circuit ussing chamber measurements, transwell inserts (⌀ 12 mm) were mounted in

the chamber device and continuously perfused at the apical and basal side with a Ringer

solution of the following composition (mmol/L) 145 NaCl, 1.6 K_2HPO_4, 1 $MgCl_2$, 0.4

KH_2PO_4, 1.3 Ca^{2+} Gluconate and 5 Glucose, and pH adjusted to 7.4. Following a 20 min

stabilization period amiloride (20µM) was added to the apical side to block epithelial Na^+

channel-mediated currents, followed by forskolin/IBMX (2µM/100µM), VX-770 (3µM) and

CFTRInh-172 (30µM) were added sequentially. Transepithelial Voltage (V_{te}) values were

recorded at all times with Power Lab software (AD Instruments Inc.). Values for V_{te} were

referred to the basal side of the epithelium and transepithelial resistance (R_{te}) was determined

by applying short intermittent pulses (0.5 μA/s), measuring pulsated deviations in V_{te} and

accounting for the area of the inserts. An empty insert was previously recorded to correct the

measured values. Short-circuit currents (I_{eq-sc}) were calculated according to Ohm's law from

V_{te} and R_{te} ($I_{eq-sc} = V_{te} / R_{te}$).

RNA extraction, cDNA synthesis, and quantitative real time PCR

Total RNA was extracted from ALI-cultures using the RNeasy Mini Kit according to the

manufacturer's protocol. RNA yield was determined by a Nanodrop spectrophotometer and

subsequently cDNA was synthesized by use of the iScript cDNA synthesis kit according to

the manufacturer's protocol. Quantitative real-time PCR (qPCR) was performed with specific

primers (Table S4) using the iQ SYBR Green Supermix and a CFX96 real-time detection

machine. CFX Manager 3.1 software was used to calculate relative gene expression

normalized to the housekeeping genes *ATP5B* and *RPL13A* according to the standard curve

method.

Immunofluorescence staining and microscopy

2D expanded HNEC and ALI-HNEC cultures were fixed in 4% paraformaldehyde for 15 min,

permeabilized in 0.25% (v/v) Triton-X in PBS for 30 min and treated with blocking buffer,

consisting of 1% (w/v) BSA and 0.25% (v/v) Triton-X in PBS, for 60 min. Next, primary

antibodies (1:500) in blocking buffer were added at the apical side and incubated for 1-2 h or

overnight. Afterwards, cells were washed three times with PBS and incubated with secondary

antibodies and Phalloidin (1:500) in blocking buffer for 30 min in dark, followed by three

washings in PBS. Transwell membranes were subsequently cut from the inserts and placed on

slides. All samples were mounted with Prolong Gold antifade reagent with or without DAPI.

Resected nasal tissue and ALI-HNEC cultures were fixed with 4 % paraformaldehyde and

embedded in paraffin after dehydration. After deparaffinization followed by antigen retrieval

using citrate buffer (pH = 6) for 20 min, the 5 µm sections were permeabilized in 0.25% (v/v)

Triton-X in PBS for 15 min, then treated with blocking buffer, consisting of 5% (w/v) BSA

and 0.025% (v/v) Triton-X in PBS, for 30 min. Primary antibodies in blocking buffer, were

incubated for 2 h, followed by incubation of secondary antibodies for 1 h. Samples were

mounted with Prolong Gold reagent with DAPI. Organoids plated in 4 µl droplets of 75%

matrigel (v/v) in a 96 wells plate, were fixed with 4 % paraformaldehyde for 10 min, and

stained as previously described [11,36], using indicated primary antibodies. Images were

acquired with a Leica SP8X confocal microscope, Leica THUNDER imager, and Leica TCS

SP8 STED 3X microscope. Images were processed using LAS X software and ImageJ/FIJI.

Quantification and statistical analysis

Swelling assays were conducted in quadruplicates for each experimental condition, and

results are shown as mean ± SEM of independent subjects. Increases in the total surface area

of all organoids in a single well is calculated as normalized swelling, relative to t=0, which is

set as baseline of 100%. For swelling assays, statistical analysis was assessed with area under

the curve (AUC) values (t=120 min). Analysis of differences was determined with a one/two-

way repeated measurements ANOVA and Bonferroni *post-hoc* test or (un)paired Student's t-

test as indicated in the figure legends. Normal distribution was tested using the Shapiro Wilk

test. Differences were considered significant at $p < 0.05$. Statistical analysis was conducted

using Prism 8 (GraphPad Software Inc.).

Acknowledgments

This work was supported by grants of the Dutch Cystic Fibrosis Foundation (NCFS, HIT-CF

grant); Netherlands Organization for Health Research and Development (ZonMw); Health

Holland (grant no 40-41200-98-9296); SRC 013 from CF Trust-UK; UIDB/04046/2020 and

UIDP/04046/2020 centre grants (to BioISI), both from FCT/MCTES Portugal; and "HIT-CF"

(H2020-SC1-2017-755021) from EU. This work is supported by the European Research

Council (ERC Consolidator Grant 819219 to L.C.K.).

Author contribution

Conception and/or design: GA, LR, CE, JMB; Acquisition, analysis, or interpretation of data:

GA, LR, HR, EA, IS, WN, SV, BA, EK, JEB, SM, KW, HH, LK, MA, CE, JMB; Drafting the

work or revising it critically for important intellectual content: GA, CD, MA, CE, JMB; Final

approval of the version submitted for publication; GA, LR, HR, EA, IS, WN, SV, BA, EK,

JEB, CD, SM, KW, HH, LK, MA, CE, JMB.

Conflict of interest

JMB has a patent granted (10006904) related to CFTR function measurements in organoids

and received personal fees from HUB/ Royal Dutch academy of sciences, during the conduct

of the study; non-financial support from Vertex Pharmaceuticals, and personal fees and non-

financial support from Proteostasis Therapeutics, outside the submitted work. CE reports

grants from GSK, Nutricia, TEVA, Gilead, Vertex, ProQR, Proteostasis, Galapagos NV,

Eloxx pharmaceuticals, outside the submitted work; In addition, CE has a patent related to

CFTR function measurements in organoids (10006904) with royalties paid. MA reports grants

and personal fees from Vertex Pharmaceuticals, grants from Gilead Sciences, Inc., grants and

personal fees from Proteostasis Therapeutics, personal fees from Translate Bio MA, Inc,

during the conduct of the study.

Data availability

All data is provided with the manuscript.

Reference list

1. Riordan, J.R., Rommens, J.M., Kerem, B., Alon, N., Rozmahel, R., Grzelczak, Z., Zielenski, J., Lok, S., Plavsic, N., and Chou, J.L. (1989). Identification of the cystic fibrosis gene: cloning and characterization of complementary DNA. Science *245*, 1066–1073.

2. Knowles, M.R., Stutts, M.J., Spock, A., Fischer, N., Gatzy, J.T., and Boucher, R.C. (1983). Abnormal ion permeation through cystic fibrosis respiratory epithelium. Science *221*, 1067–1070.

3. Stoltz, D.A., Meyerholz, D.K., and Welsh, M.J. (2015). Origins of Cystic Fibrosis Lung Disease. N. Engl. J. Med. *372*, 351–362.

4. Boyle, M.P., and De Boeck, K. (2013). A new era in the treatment of cystic fibrosis: correction of the underlying CFTR defect. Lancet Respir. Med. *1*, 158–163.

5. Clancy, J.P., Cotton, C.U., Donaldson, S.H., Solomon, G.M., VanDevanter, D.R., Boyle, M.P., Gentzsch, M., Nick, J.A., Illek, B., Wallenburg, J.C., et al. (2019). CFTR modulator theratyping: Current status, gaps and future directions. J. Cyst. Fibros. Off.

J. Eur. Cyst. Fibros. Soc. *18*, 22–34.

6.   Gentzsch, M., Boyles, S.E., Cheluvaraju, C., Chaudhry, I.G., Quinney, N.L., Cho, C.,

Dang, H., Liu, X., Schlegel, R., and Randell, S.H. (2017). Pharmacological Rescue of

Conditionally Reprogrammed Cystic Fibrosis Bronchial Epithelial Cells. Am. J.

Respir. Cell Mol. Biol. *56*, 568–574.

7.   Awatade, N.T., Uliyakina, I., Farinha, C.M., Clarke, L.A., Mendes, K., Solé, A.,

Pastor, J., Ramos, M.M., and Amaral, M.D. (2015). Measurements of Functional

Responses in Human Primary Lung Cells as a Basis for Personalized Therapy for

Cystic Fibrosis. EBioMedicine *2*, 147–153.

8.   Pranke, I.M., Hatton, A., Simonin, J., Jais, J.P., Le Pimpec-Barthes, F., Carsin, A.,

Bonnette, P., Fayon, M., Stremler-Le Bel, N., Grenet, D., et al. (2017). Correction of

CFTR function in nasal epithelial cells from cystic fibrosis patients predicts

improvement of respiratory function by CFTR modulators. Sci. Rep. *7*, 7375.

9.   Dekkers, J.F., Wiegerinck, C.L., de Jonge, H.R., Bronsveld, I., Janssens, H.M., de

Winter-de Groot, K.M., Brandsma, A.M., de Jong, N.W.M., Bijvelds, M.J.C., Scholte,

B.J., et al. (2013). A functional CFTR assay using primary cystic fibrosis intestinal

organoids. Nat. Med. *19*, 939–945.

10.   Schutgens, F., Rookmaaker, M.B., Margaritis, T., Rios, A., Ammerlaan, C., Jansen, J.,

Gijzen, L., Vormann, M., Vonk, A., Viveen, M., et al. (2019). Tubuloids derived from

human adult kidney and urine for personalized disease modeling. Nat. Biotechnol. *37*,

303–313.

11.   Sachs, N., Papaspyropoulos, A., Zomer-van Ommen, D.D., Heo, I., Böttinger, L., Klay,

D., Weeber, F., Huelsz-Prince, G., Iakobachvili, N., Amatngalim, G.D., et al. (2019).

431   Long-term expanding human airway organoids for disease modeling. EMBO J. *38*,

432   e100300.

433 12. Berkers, G., van Mourik, P., Vonk, A.M., Kruisselbrink, E., Dekkers, J.F., de Winter-

434   de Groot, K.M., Arets, H.G.M., Marck-van der Wilt, R.E.P., Dijkema, J.S.,

435   Vanderschuren, M.M., et al. (2019). Rectal Organoids Enable Personalized Treatment

436   of Cystic Fibrosis. Cell Rep. *26*, 1701-1708.e3.

437 13. Dekkers, J.F., Berkers, G., Kruisselbrink, E., Vonk, A., de Jonge, H.R., Janssens, H.M.,

438   Bronsveld, I., van de Graaf, E.A., Nieuwenhuis, E.E.S., Houwen, R.H.J., et al. (2016).

439   Characterizing responses to CFTR-modulating drugs using rectal organoids derived

440   from subjects with cystic fibrosis. Sci. Transl. Med. *8*, 344ra84.

441 14. van Mourik, P., Beekman, J.M., and van der Ent, C.K. (2019). Intestinal organoids to

442   model cystic fibrosis. Eur. Respir. J. *54*.

443 15. Ramalho, A.S., Fürstová, E., Vonk, A.M., Ferrante, M., Verfaillie, C., Dupont, L.,

444   Boon, M., Proesmans, M., Beekman, J.M., Sarouk, I., et al. (2020). Correction of

445   CFTR function in intestinal organoids to guide treatment of Cystic Fibrosis. Eur.

446   Respir. J.

447 16. Guimbellot, J.S., Leach, J.M., Chaudhry, I.G., Quinney, N.L., Boyles, S.E., Chua, M.,

448   Aban, I., Jaspers, I., and Gentzsch, M. (2017). Nasospheroids permit measurements of

449   CFTR-dependent fluid transport. JCI insight *2*.

450 17. Brewington, J.J., Filbrandt, E.T., LaRosa, F.J. 3rd, Ostmann, A.J., Strecker, L.M.,

451   Szczesniak, R.D., and Clancy, J.P. (2018). Detection of CFTR function and modulation

452   in primary human nasal cell spheroids. J. Cyst. Fibros. Off. J. Eur. Cyst. Fibros. Soc.

453   *17*, 26–33.

454   18.   Liu, Z., Anderson, J.D., Deng, L., Mackay, S., Bailey, J., Kersh, L., Rowe, S.M., and

455   Guimbellot, J.S. (2020). Human Nasal Epithelial Organoids for Therapeutic

456   Development in Cystic Fibrosis. Genes (Basel). *11*.

457   19.   Sette, G., Lo Cicero, S., Blaconà, G., Pierandrei, S., Bruno, S.M., Salvati, V., Castelli,

458   G., Falchi, M., Fabrizzi, B., Cimino, G., et al. (2021). Theratyping cystic fibrosis in

459   vitro in ALI-culture and organoid models generated from patient-derived nasal

460   epithelial Conditionally Reprogrammed Stem Cells. Eur. Respir. J.

461   20.   Keating, D., Marigowda, G., Burr, L., Daines, C., Mall, M.A., McKone, E.F., Ramsey,

462   B.W., Rowe, S.M., Sass, L.A., Tullis, E., et al. (2018). VX-445-Tezacaftor-Ivacaftor in

463   Patients with Cystic Fibrosis and One or Two Phe508del Alleles. N. Engl. J. Med.

464   *379*, 1612–1620.

465   21.   Clarke, L.L., and Boucher, R.C. (1992). Chloride secretory response to extracellular

466   ATP in human normal and cystic fibrosis nasal epithelia. Am. J. Physiol. *263*, C348-

467   56.

468   22.   Okuda, K., Dang, H., Kobayashi, Y., Carraro, G., Nakano, S., Chen, G., Kato, T.,

469   Asakura, T., Gilmore, R.C., Morton, L.C., et al. (2021). Secretory Cells Dominate

470   Airway CFTR Expression and Function in Human Airway Superficial Epithelia. Am.

471   J. Respir. Crit. Care Med. *203*, 1275–1289.

472   23.   Carraro, G., Langerman, J., Sabri, S., Lorenzana, Z., Purkayastha, A., Zhang, G.,

473   Konda, B., Aros, C.J., Calvert, B.A., Szymaniak, A., et al. (2021). Transcriptional

474   analysis of cystic fibrosis airways at single-cell resolution reveals altered epithelial cell

475   states and composition. Nat. Med. *27*, 806–814.

24.   Kettle, R., Simmons, J., Schindler, F., Jones, P., Dicker, T., Dubois, G., Giddings, J.,

Van Heeke, G., and Jones, C.E. (2010). Regulation of neuregulin 1beta1-induced

MUC5AC and MUC5B expression in human airway epithelium. Am. J. Respir. Cell

Mol. Biol. *42*, 472–481.

25. Vermeer, P.D., Panko, L., Karp, P., Lee, J.H., and Zabner, J. (2006). Differentiation of

human airway epithelia is dependent on erbB2. Am. J. Physiol. Lung Cell. Mol.

Physiol. *291*, L175-80.

26. Gray, T., Coakley, R., Hirsh, A., Thornton, D., Kirkham, S., Koo, J.-S., Burch, L.,

Boucher, R., and Nettesheim, P. (2004). Regulation of MUC5AC mucin secretion and

airway surface liquid metabolism by IL-1beta in human bronchial epithelia. Am. J.

Physiol. Lung Cell. Mol. Physiol. *286*, L320-30.

27. Abdullah, L.H., Coakley, R., Webster, M.J., Zhu, Y., Tarran, R., Radicioni, G.,

Kesimer, M., Boucher, R.C., Davis, C.W., and Ribeiro, C.M.P. (2018). Mucin

Production and Hydration Responses to Mucopurulent Materials in Normal versus

Cystic Fibrosis Airway Epithelia. Am. J. Respir. Crit. Care Med. *197*, 481–491.

28. Gentzsch, M., Cholon, D.M., Quinney, N.L., Boyles, S.E., Martino, M.E.B., and

Ribeiro, C.M.P. (2018). The cystic fibrosis airway milieu enhances rescue of F508del

in a pre-clinical model. Eur. Respir. J. *52*.

29. Chen, G., Sun, L., Kato, T., Okuda, K., Martino, M.B., Abzhanova, A., Lin, J.M.,

Gilmore, R.C., Batson, B.D., O'Neal, Y.K., et al. (2019). IL-1β dominates the

promucin secretory cytokine profile in cystic fibrosis. J. Clin. Invest. *129*, 4433–4450.

30. Brouillard, F., Bouthier, M., Leclerc, T., Clement, A., Baudouin-Legros, M., and

Edelman, A. (2001). NF-kappa B mediates up-regulation of CFTR gene expression in

Calu-3 cells by interleukin-1beta. J. Biol. Chem. *276*, 9486–9491.

31.   Balázs, A., and Mall, M.A. (2018). Role of the SLC26A9 Chloride Channel as Disease

501   Modifier and Potential Therapeutic Target in Cystic Fibrosis. Front. Pharmacol. *9*,

502   1112.

503   32.   Anderson, J.D., Liu, Z., Odom, L.V., Kersh, L., and Guimbellot, J.S. (2021). CFTR

504   function and clinical response to modulators parallel nasal epithelial organoid

505   swelling. Am. J. Physiol. Lung Cell. Mol. Physiol. *321*, L119–L129.

506   33.   Harris, C.M., Mendes, F., Dragomir, A., Doull, I.J.M., Carvalho-Oliveira, I., Bebok, Z.,

507   Clancy, J.P., Eubanks, V., Sorscher, E.J., Roomans, G.M., et al. (2004). Assessment of

508   CFTR localisation in native airway epithelial cells obtained by nasal brushing. J. Cyst.

509   Fibros. Off. J. Eur. Cyst. Fibros. Soc. *3 Suppl 2*, 43–48.

510   34.   Stokes, A.B., Kieninger, E., Schögler, A., Kopf, B.S., Casaulta, C., Geiser, T.,

511   Regamey, N., and Alves, M.P. (2014). Comparison of three different brushing

512   techniques to isolate and culture primary nasal epithelial cells from human subjects.

513   Exp. Lung Res. *40*, 327–332.

514   35.   Boj, S.F., Vonk, A.M., Statia, M., Su, J., Vries, R.R.G., Beekman, J.M., and Clevers,

515   H. (2017). Forskolin-induced Swelling in Intestinal Organoids: An In Vitro Assay for

516   Assessing Drug Response in Cystic Fibrosis Patients. J. Vis. Exp.

517   36.   Dekkers, J.F., Alieva, M., Wellens, L.M., Ariese, H.C.R., Jamieson, P.R., Vonk, A.M.,

518   Amatngalim, G.D., Hu, H., Oost, K.C., Snippert, H.J.G., et al. (2019). High-resolution

519   3D imaging of fixed and cleared organoids. Nat. Protoc. *14*, 1756–1771.

Figure Legends

Figure 1) Conversion of differentiated ALI-HNEC cultures into nasal airway organoids.

(A) Graphic illustration showing workflow of culturing nasal airway organoids (NAOs) from

2D expanded and differentiated human nasal epithelial cells (HNEC). (B) Brightfield images

of a 2D expanded HNEC. (C) IF staining of HNEC with basal cell markers p63 (red) and

cytokeratin 5 (KRT5, green). (D) Sections of nasal tissue (top) and 18 days differentiated air-

liquid interface (ALI-) cultured HNEC (bottom), demonstrating IF staining of MUC5AC

(goblet cells, red), β-tubulin IV (ciliated cells, green), p63 and KRT5 (basal cell markers,

green and red respectively). (E) Whole-mount IF confocal images of ALI-HNEC, showing

maximal projections of the apical side of the secretory cell markers MUC5AC and Club Cell

protein 10 (CC10), and ciliated cell marker β-tubulin IV (Z-stack at the bottom). The

projection of the basal side shows the basal cell marker p63 (Z-stack at the bottom). All

markers are shown in green. (F) Time course showing self-organization of differentiated ALI-

HNEC-derived epithelial fragments into organoids. (G) IF staining of NAOs for β-tubulin IV,

MUC5AC, and p63 (all in green). (H) Confocal images showing in the top panels staining

with DAPI (blue), β-tubulin IV (green), and MUC5AC (red). The bottom panels show

staining with DAPI (blue), p63 (green), and KRT5 (red). Data information: DAPI (blue) was used as nuclear staining (C-E, G, and H). Phalloidin (red) was used as actin cytoskeleton staining (E and G).

Figure 2) Forskolin-induced swelling assay with ALI-culture derived nasal airway organoids. (A) Graphic illustration showing FIS assay with ALI-derived nasal airway organoids (NAOs). (B) Representative brightfield images (top) and images of calcein green AM esters-strained (bottom) organoids from a HC subject, which were unstimulated (Control) or treated with forskolin (Fsk, 5 µM). Images were taken at t=0, 60, and 120 min after stimulation. (C and D) NAOs from HC subjects (n=5 independent donors) were stimulated with different concentrations of Fsk (0-10 µM), followed by quantification of FIS. (E and F) HC NAOs (n=3 independent donors) were pre-treated with CFTR inhibitors or vehicle, followed by assessment of FIS. (G) Representative brightfield images of NAOs from HC subjects and subjects with CF. (H) Quantification of the mean organoid area (μm^2, means ± SD) of HC and CF NAOs (both n=6 independent donors). (I and J) Comparison of FIS between HC and CF NAOs (both n=8 independent donors) after Fsk stimulation. Data information: Results are depicted as the percentage change in surface area relative to t=0 (normalized area) measured at 15-minute time intervals for 2 h (means ± SD) (C, E, I), and area under the curve (AUC) plots (t=120 min, means ± SD) (D, F, J). Analysis of differences was determined with a one-way ANOVA and Bonferroni *post-hoc* test (D), and unpaired t-test (F, H, J). ** $p<0.01$, *** $p<0.001$, **** $p< 0.0001$.

Figure 3) Validation of CFTR modulator responses in NR/IL-1β-cultured NAOs.
(A) Illustration showing culturing of NAOs with neuregulin 1β (NR) and interleukin-1β (IL-1β) (B) CF F508del/F508del NAOs (n=3 independent donors) were cultured at control

conditions, or with NR, IL-1β, or combination (NR+IL-1β). Afterwards, FIS was determined

in response to VX-809/VX-770. (C) Assessment of FIS in HC NAOs (n=3 independent

donors) cultured under control conditions or NR+IL-1β. (D) FIS measured in HC NAOs (n=5

independent donors) and (E) CF F508del/F508del NAOs (n=7 independent donors) cultured

with NR/IL-1β. FIS responses in CF NAOs were determined in response to VX-809/VX-770.

(F) Comparison of FIS measured in HC (Figure 3F) and CF F508del/F508del NAOs (Figure

3E). (G and H) FIS in NR+IL-1β cultured NAOs from CF F508del/S1251N subjects (n=4

independent subjects), stimulated with Fsk and VX-770 or vehicle. (I and J) CF F508del

homozygous NAOs cultured with NR/IL-1β (n=4 independent donors) were pre-treated with

vehicle or VX-661/VX-445. Swelling was determined afterwards following acute stimulation

with Fsk together with VX-770 or vehicle. Data information: Swelling results are depicted as

(D, E, G, I) the percentage change in surface area relative to t=0 (normalized area) measured

at 15-minute time intervals for 2 h (means ± SD), and (B, C, F, H, J) area under the curve

(AUC) plots (t=120 min, means ± SD). Analysis of differences was determined with a paired

t-test (B, C, F; within groups, H, J), unpaired t-test (F; HC compared to CF). * $p<0.05$, **

$p<0.01$, *** $p<0.001$, **** $p< 0.0001$.

Figure 1

Figure 2

Figure 3