Mutations in coral soma and sperm imply lifelong stem cell differentiation

Elora H. López-Nandam[1,2]*, Rebecca Albright[2], Erik A. Hanson[1], Elizabeth A. Sheets[1,†], Stephen

R. Palumbi[1]

[1]Biology Department, Hopkins Marine Station of Stanford University, Pacific Grove, California

93950 USA

[2] Institute for Biodiversity and Sustainability Science, California Academy of Sciences, San

Francisco, California, 94118, USA

[†] Current address: Genomics Institute, University of California, Santa Cruz, 95064, USA

*corresponding author

In many animals, the germline differentiates early in embryogenesis, so only

mutations that accumulate in germ cells are inherited by offspring[1]. Exceptions to this

developmental process may indicate that other mechanisms have evolved to limit the effects

of deleterious mutation accumulation[2]. Stony corals are animals that can live for hundreds

of years[3] and have long been thought to produce gametes from somatic tissue[4]. To clarify

conflicting evidence about germline-soma distinction in corals, we sequenced high

coverage, full genomes with technical replicates for parent coral branches and their sperm

pools. We identified single nucleotide variants (SNVs) unique to each parent branch, then

checked if each SNV was shared by the respective sperm pool: 26% of post-embryonic

SNVs were shared by the sperm and 74% were not. We also identified germline SNVs,

those that present in the sperm but not in the parent. These data suggest that self-renewing

stem cells in corals differentiate into germ and soma throughout the adult life of the colony,

with SNV rates and patterns differing markedly in stem, soma, and germ lineages. In addition

25 to informing germline evolution, these insights inform how corals may generate adaptive

26 diversity necessary in the face of global climate change.

27   In many animals, germ and somatic cells differentiate early in the embryonic stage,

28 leading to separate lineages. Weismann first hypothesized this process in 1889 to explain why

29 mutations that accumulate in somatic tissues during an animal's lifetime– including those that

30 cause cancer- are not inherited by that animal's offspring. Instead, only mutations in germ cells,

31 which undergo fewer cell divisions and have lower mutation rates, are inherited[5,6]. Since

32 Weismann, embryonic germ-soma separation has been shown in vertebrates and many other

33 animal taxa, but not in plants or in some animal groups, including cnidarians, sponges, tunicates,

34 and platyheminths[7,8].

35   Animal exceptions to Weismann's Germ Plasm Theory are intriguing because they may

36 have novel mechanisms to reduce the number of deleterious mutations inherited by sexually

37 produced offspring. Moreover, such exceptions may signal the potential existence of stem cell

38 lineage types not seen in vertebrates. The model cnidarians *Hydra* and *Hydractinia* possess

39 interstitial stem cells, denoted i-cells, that can differentiate into both germ and soma during adult

40 life[9,10]. A few models have hypothesized how heritable post-embryonic mutations may affect the

41 gamete pool[11–13], but there is very little data on the pattern of somatic mutations and their

42 inheritance in long-lived animals[14].

43   Clonal, colonial corals can live for hundreds to thousands of years, and were long thought

44 to generate gametes from the somatic cells of clonal polyps[4]. Coral colonies accumulate somatic

45 mutations at a rate similar to noncancerous human tissues[14]. If these mutations are inherited by

46 the coral's gametes, they must increase the heritable mutational load of these animals. Some

47 previous studies identified putative somatic mutations in the gametes or juvenile offspring of

48 mutant parents[15,16], but others have reported absence of somatic mutations in the gametes[17].

These studies tracked few mutations, ranging from 9 to 170, and none detected germline

mutations in gametes or offspring. Only one verified that their putative mutations were not PCR

or sequencing error[17]. Here, we interrogated full genomes of multiple branches from multiple

coral colonies and their sperm. We identified germline variants in the sperm as well as post-

embryonic variants in the parent. The data reject the hypothesis that somatic cells give rise to

germ cells in corals, and reject the hypothesis that corals possess embryonic germline

differentiation. We hypothesize that both parent tissue and sperm arise from a common stem cell

lineage that proliferates and differentiates throughout the long lives of these animals.

To clarify the inheritance of mutations and the presence of germline-soma distinction in

Acropora hyacinthus, we removed branches from soon-to-spawn adult coral colonies and placed

them into individual cups of seawater (Extended Data Fig. 1). Each branch released gamete

bundles into its respective cup 20 minutes later (Fig 1a, Extended Data Fig. 1b). We extracted

DNA from each branch and each sperm pool, then constructed two replicate full genome

libraries from each DNA extraction (Fig 1b). To be verified, a SNV had to be present in both

replicate libraries of a given sample. The technical replicates eliminated over 90% of putative

SNVs that would have been called if we had used one library per sample, although the exact

number varied by SNV category (Extended Data Figs. 2, 3).

We identified four different types of SNVs: those that were unique to the polyps from a

single parent branch in a colony but were not detected in the sperm from that branch (Parent

Only) (1c), those that were found in just a single parent branch in a colony and were also shared

by the sperm from that branch (Parent and Sperm) (1d), those that were unique to a single sperm

pool in a colony and not present in any branch of the colony (Single Sperm Pool) (1e), and those

that were shared by all sperm pools in a colony but had never been seen in the polyps from any

branch (All Sperm Pools) (1f).

We assayed nine parent polyp samples, and the respective sperm pools for seven of those

samples, across three different colonies. The average depth of coverage across the genome was

40.6 ±3.1 (1 s.e.m) for the parent polyp libraries and 65.2 ±6.9 (1 s.e.m) for the sperm pool

libraries (Supp. Table 1). Across the full dataset we identified 2,356 SNVs, and all but one were

at unique sites, indicating that the SNVs called were not a result of consistent mapping error or

bias (Supp. Table 2). Each SNV was classified as a Gain of Heterozygosity (GoH), in which the

aberrant sample was a new heterozygote and all others were homozygous, or a Loss of

Heterozygosity (LoH), in which the aberrant sample was homozygous and the other samples

were heterozygous.

We identified 146-351 post-embryonic SNVs per parent branch (Supp. Table 1), and we

found that on average 25.7% ±3.7% (1 s.e.m.) SNVs were shared between a parent branch and

its respective sperm pool (Parent and Sperm, labelled P+S in Fig. 2a), whereas 74.3% ±3.7% (1

s.e.m.) post-embryonic SNVs found in a branch were not in the sperm (Parent Only, labelled PO

in Fig. 2a). These findings contradict the hypothesis from the Germ Plasm Theory that post-

embryonic mutations would not be found in the sperm at all, as well as the common assumption

that all coral somatic cells can produce gametes.

We found 50-145 post-embryonic SNVs in every sperm pool (Supp. Table 1). Of these

39.2% ±3.5% were shared with its parent branch (Parent and Sperm), and 52.2% ±5.4% were

found only in the sperm pool (Single Sperm Pool Only) (Fig. 2b). A small number of sperm

SNVs, 8.5% ±2.1%, were found in all sperm pools from that colony but none of the parent

samples that spawned them (All Sperm Pools, labelled ASP, Fig. 2b). That 2 out of every 5

SNVs present in a given sperm pool are post-embryonic, non-germline variants indicates that the

lack of an embryonic germline increases the number of SNVs in a colony's gametes by 66%,

compared to what the diversity would have been if the germline were segregated at the

embryonic stage. This may help explain the high degree of heterozygosity in many stony coral

species, though it is not yet known what fraction of these SNVs are too deleterious to survive

into adulthood.

The rate of SNVs per bp was significantly higher across the full callable genome than the

rate of SNVs in the callable coding regions of the genome for all SNV types (Fig. 2c) (see Table

1 for all means and Wilcoxon signed-rank test results). This may indicate that there is stronger

purifying selection against SNVs in coding regions than in non-coding regions of the genome, or

it may be a result of higher mismatch repair in exons[18].

We examined the spectrum of mutations, the relative proportions of mutations in

different classes, and found no significant differences in spectra among parent only, shared, and

germ line specific mutations (Extended Data Fig. 5). These data confirm lack of a signature of

UV-associated mutations in corals[13], which is intriguing considering that these colonies grow in

high UV conditions, and in highly oxygenated warm water[19].

Losses of heterozygosity tend to arise as a result of gene conversion due to homologous

recombination, a form of double stranded DNA break repair[20]. Consistent with previous

findings[14], 38.3 % ±3.0 % (1 s.e.m.) of all parent SNVs being GoH and 61.7 ±3.0% (1 s.e.m.)

being LoH. SNVs that were shared by the parent branch tissues and the sperm had a much higher

fraction of GoH and lower fraction of LoH (73.8 ±3.6% and 26.2 ±3.6%, respectively) than did

parent SNVs that were not found in the sperm (25.2 ±2.4% and 74.8 ±2.4%, respectively)

(Wilcoxon signed-rank test, V=28, p = 0.015) (Fig. 2d). SNVs found in just a single sperm pool

had approximately equal proportions of each type, 51.1% ±2.5% GOH and 48.9 ±2.5% LOH

SNVs. High LOH in soma that is not inherited by the sperm could be due to high incidence of

double-strand breaks in somatic cells exposed to high light and photosynthetically derived

oxidation, or high LOH levels in the parent-only SNVs may reflect stronger selection against

GOH than LOH in differentiated somatic cells.

To explore the role of selection on patterns of genome change, we compared the rates of

missense and synonymous SNVs across four classes: all somatic SNVs, parent-only SNVs,

shared parent and sperm SNVs, and SNVs found in a single sperm pool only. There were no

coding SNVs in the all sperm pool category. The average rate of coding mutations was highest in

parent only SNVs (6.0×10^{-7} ±1.6×10^{-7}, Extended Data Fig. 6a). The percent of coding

mutations that were missense was higher in single sperm pool SNVs (73.7 ±11.8%) than in the

other categories (55.2 ±6.8% all somatic, 51.5 ±10.7% parent only, 47.6 ±14.6% parent and

sperm, Extended Data Fig. 6b). The higher mean percent missense in SSPO was not statistically

significant, likely attributable to the small number of coding mutations in each category. Like

most studies on somatic mutations to date, the small number of coding mutations in this study

(94) leaves us underpowered to detect selection[21]. However, the fairly consistent pattern of more

missense mutations in sperm pool samples than somatic samples provides a first hint that the

SNVs in the soma may experience stronger negative selection than germline SNVs.

If we had found only separate parent and sperm SNVs this would have shown that

Acropora corals have classical Weismannian germ and somatic cell lineage differentiation at the

embryonic stage, which has been suggested previously[17]. Likewise, if we had found that all

parent tissue SNVs were also in the sperm, we would have concluded that *Acropora* corals

developed gametes directly from those tissues[15,16]. However, 74% of the SNVs that we identified

in parent tissue were also present in the sperm spawned from that branch, and 26% were not.

Based on these data, we hypothesize that in colonial corals, shared parent and sperm SNVs

derive from mutations in a common ancestor stem cell lineage that self-renews and proliferates

through the colony, and that eventually differentiates into both germ and soma throughout the

colony's adult life. This type of lineage (i-cells) has been described in Hydrozoan cnidarians[9,10].

Although i-cells have not yet been identified in corals, cells that look like the i-cells are present

in larvae of the coral *Acropora millepora*[22]. Our data suggest that branch-specific SNVs shared

in germ and somatic cells first arose in an i-cell lineage proliferating in that branch, and then

differentiated into germ and soma (Fig. 3a). SNVs found only in the parent but not in the sperm

would have arisen in terminally-differentiated somatic cells that cannot produce gametes, and

SNVs found only in the sperm would have arisen from differentiated germ cells (Figure 3a).

We hypothesize that the program of sequential germ line differentiation during adult life

shown in Hydrozoans is likely a conserved trait across Cnidaria and was present in the cnidarian

common ancestor. Germ and soma differentiation appears to happen locally, resulting in evident

mosaicism in every branch (Fig. 2a, Fig. 3b). Solana[8] made a similar prediction for planarians–

when stem cells can differentiate into both soma and germ cells, mutations that appear in both

the soma and the germline are derived from mutations in those stem cells.

Weismann's original germ cell theory proposed a mechanism whereby mutations

accumulated during an individual's lifetime would not be inherited by its offspring. In the case of

corals, our data show that some post-embryonic SNVs that occur in different branches within a

colony are inherited by its sperm. If i-cell SNVs are subject to selection, then the selection

regime that growing i-cell lines face could select for novel beneficial changes as well as select

against deleterious ones [2,23]. Reef building corals are extremely sensitive to small increases in

temperature, but these environmental changes frequently result in the death of just part of a

colony. If partial survival of a colony is the result of selection for post-embryonic SNVs in

different parts of the colony, then adaptation to environmental change may occur over the

lifetime of a single colony. If some of those post-embryonic SNVs are inherited by the surviving

polyps' gametes, then this may be an alternative, rapid route to adaptation for corals.

Our data suggest that anthozoans have i-cells that self-renew and remain multipotent

throughout the adult lifespan, which has previously been described in medusozoans. We also

show, for the first time, the genome-level consequences of SNVs in i-cells on the mutation load

of a long-lived animal species that lacks an embryonic germline. SNVs in the stem cell lines of a

coral colony increase the number of SNVs in the sperm by 66%. This may help to explain the

high degree of heterozygosity and adaptive polymorphism in many stony coral species.

Mechanisms that corals use to avoid mutational meltdown in long-lived cell lineages might

include consistent screening by natural selection in proliferating cell lines, or yet-to-be

discovered controls on coding gene mutation rates.

Acknowledgements

EHLN was funded by NSF GRFP and a Stanford Graduate Fellowship in Science and

Engineering (Morgridge Family Foundation). Research funded by grants from the Chan-

Zuckerberg BioHub and NSF OCE-1736736. We thank N. Neff at CZ-BioHub for sequencing

power and A. Voskoboynik P. Bump, and B. Cornwell for stem cell insights. Thanks to R.

Ritson-Williams, R. Ross, and BIOTA for field collection support, and thanks to the staff of the

Steinhart Aquarium for taking care of the corals.

References

1.   Saffman, E. E. & Lasko, P. Germline development in vertebrates and invertebrates. *Cell.*

186   *Mol. Life Sci. C. 1999 558* **55**, 1141–1163 (1999).

187 2. Buss, L. W. Evolution, development, and the units of selection. *Proc. Natl. Acad. Sci. U.*

188   *S. A.* **80**, 1387–91 (1983).

189 3. Bythell, J. C., Brown, B. E. & Kirkwood, T. B. L. Do reef corals age? *Biol. Rev* 0–0

190   (2017). doi:10.1111/brv.12391

191 4. van Oppen, M. J. H., Souter, P., Howells, E. J., Heyward, A. & Berkelmans, R. Novel

192   Genetic Diversity Through Somatic Mutations: Fuel for Adaptation of Reef Corals?

193   *Diversity* **3**, 405–423 (2011).

194 5. Lynch, M. Evolution of the mutation rate. *Trends Genet.* (2010).

195   doi:10.1016/j.tig.2010.05.003

196 6. Moore, L. *et al.* The mutational landscape of human somatic and germline cells 1 Author

197   List 2. *bioRxiv* 2020.11.25.398172 (2020). doi:10.1101/2020.11.25.398172

198 7. Whitham, T. G. & Slobodchikoff, C. N. Evolution by individuals, plant-herbivore

199   interactions, and mosaics of genetic variability: The adaptive significance of somatic

200   mutations in plants. *Oecologia* **49**, 287–292 (1981).

201 8. Solana, J. Closing the circle of germline and stem cells: The Primordial Stem Cell

202   hypothesis. *Evodevo* **4**, (2013).

203 9. DuBuc, T. Q. *et al.* Transcription factor AP2 controls cnidarian germ cell induction.

204   *Science (80-.).* **367**, 757–762 (2020).

205 10. Juliano, C. E., Swartz, S. Z. & Wessel, G. M. A conserved germline multipotency

206   program. *Development* **137**, 4113–4126 (2010).

207 11. Orive, M. E. Somatic Mutations in Organisms with Complex Life Histories. *Theor. Popul.*

208   *Biol.* **59**, 235–249 (2001).

209   12.   Klekowski, E. J. & Godfrey, P. J. Ageing and mutation in plants. *Nature* **340**, 389–391
210   (1989).

211   13.   Hanlon, V. C. T., Otto, S. P. & Aitken, S. N. Somatic mutations substantially increase the
212   per-generation mutation rate in the conifer *Picea sitchensis*. *Evol. Lett.* (2019).
213   doi:10.1002/evl3.121

214   14.   López, E. H. & Palumbi, S. R. Somatic Mutations and Genome Stability Maintenance in
215   Clonal Coral Colonies. *Mol. Biol. Evol.* **37**, 828–838 (2020).

216   15.   Schweinsberg, M., González Pech, R. A., Tollrian, R. & Lampert, K. P. Transfer of
217   intracolonial genetic variability through gametes in Acropora hyacinthus corals. *Coral*
218   *Reefs* **33**, 77–87 (2014).

219   16.   Vasquez Kuntz, K. L. *et al.* Title: Juvenile corals inherit mutations acquired during the
220   parent's lifespan Running Title: Juvenile corals inherit mutations acquired during the
221   parent's 2 lifespan 3 4. *bioRxiv* 2020.10.19.345538 (2020).
222   doi:10.1101/2020.10.19.345538

223   17.   Barfield, S., Aglyamova, G. V. & Matz, M. V. Evolutionary origins of germline
224   segregation in Metazoa: evidence for a germ stem cell lineage in the coral *Orbicella*
225   *faveolata* (Cnidaria, Anthozoa). *Proc. R. Soc. B Biol. Sci.* **283**, 20152128 (2016).

226   18.   Frigola, J. *et al.* Reduced mutation rate in exons due to differential mismatch repair. *Nat.*
227   *Genet.* ng.3991 (2017). doi:10.1038/ng.3991

228   19.   Shick, J. M., Lesser, M. P. & Jokiel, P. L. Effects of ultraviolet radiation on corals and
229   other coral reef organisms. *Glob. Chang. Biol.* **2**, 527–545 (1996).

230   20.   Takata, M. *et al.* Homologous recombination and non-homologous end-joining pathways
231   of DNA double-strand break repair have overlapping roles in the maintenance of

chromosomal integrity in vertebrate cells. *EMBO J.* **17**, 5497–508 (1998).

21.   Martincorena, I. *et al.* Universal Patterns of Selection in Cancer and Somatic Tissues. *Cell*

171, 1029-1041.e21 (2017).

22.   Reyes-Bermudez, A. & Miller, A. D. J. In vitro culture of cells derived from larvae of the

staghorn coral Acropora millepora. doi:10.1007/s00338-009-0527-3

23.   Otto, S. P. & Orive, M. E. Evolutionary consequences of mutation and selection within an

individual. *Genetics* **141**, (1995).

Figures and Legends

Figure 1. Data collection (a-b) and mutation classification (c-f). a.) Twenty minutes prior to
spawning, 3-4 branches were broken off of 3 parent colonies and placed into individual cups of
seawater, for a total of 10 branches in cups. Branches then released eggs and sperm into each
cup, and sperm was collected from the cup. Both the sperm pool and the parent samples were
stored in RNAlater and frozen. b.) Genomic DNA was extracted from each parent branch and
sperm pool (see Methods). For each genomic DNA extraction we constructed two full genome
libraries (see Methods) for technical replication. Classification of mutation types: c.) A mutation
unique to a single branch of the colony, but the sperm from the branch does not share the mutant
genotype. d.) A mutation unique to a single branch of the colony, and the sperm from the branch
shares the mutant genotype. e.) A mutation unique to just one sperm pool in the colony, not

 shared by other sperm pools or the parent branches. f.) A mutant genotype shared by all sperm
252 pools from a particular colony, but none of the parent branches in that colony.
253

254
Figure 2. SNV rates and proportions across different classifications. a.) The rate of SNVs per bp
for two SNV types: shared by parent and sperm (gray) and found in the parent only (yellow) for
the seven parent-sperm pairs from the three colonies. b.) The average percentage of SNV type
(parent and sperm, P+S), (parent only, PO), and all sperm pools (ASP) found in each sperm pool
sample (N=7). c.) The average rate of SNVs per bp per sample (N=7) across the full genome and
for the coding regions only, for three SNV types: parent and sperm (P+S), parent only (PO), and
single sperm pool only (SSPO). For each SNV type, each subtype (GoH and LoH). d.) The
average percentage of SNVs that were GoH and LoH for each of the four SNV types found in
each sample (N=7). For b, c, and d, the mean for each category is shown as a large point with
error bars extending out; error bars represent ±1 s.e.m. Each individual data point (N=7 for P+S,
SSPO, PO, and All Parent, N=3 for ASP) is shown as a smaller point for each category.
266

267
268
269

Figure 3. Schematics for how SNVs that arise at different stages of cell lineage development and differentiation proliferate. a.) SNVs that arise in i-cell may be found in both germ and somatic cells later, if the mutant i-cell lineage differentiates into both germ and soma. SNVs that arise in soma or germ cell lineage post-differentiation will only be found in those differentiated lineages. b.) The framework laid out in a) overlaid onto a branching, colonial coral structure suggests how SNVs in one branch of the colony may not be present in other branches.

Methods

Sample collection

Gravid coral colonies of *Acropora hyacinthus* were collected in Palau (Bureau of Marine

Resources permit number RE-19-07 and CITES permit PW19-018) in February 2019 and

transported to the Coral Spawning Lab at the California Academy of Sciences where they were

kept on a Palauan lunar and day/night cycle until spawning, with methods adapted from [24].

Colonies were monitored for spawning activity on nights 6 – 9 after the simulated full moon in

March 2019 (from 27 March to 30 March 2019). Prior to spawning, pliers were used to break off

2-3 cm branches that were "set," or showed visual signs of impending gamete release: three

branches from each of two colonies, and four branches from a third. Each branch was placed in a

labeled 5 mL vial of seawater where they spawned approximately 20 minutes later (Fig. 1a).

After the gamete bundles were released, they were transferred to labeled 1.5 mL tubes and left to

dissociate into eggs and sperm. Upon dissociation, eggs were removed via pipet, leaving a

concentrated sperm pool. Each concentrated sperm pool was pipetted into a 1.5 mL tube of

RNAlater. Each coral branch was placed in a 5 mL tube of RNAlater. Sperm pools in RNAlater

were stored at -20° C and coral branches in RNAlater were stored at -80° C until time for DNA

extraction.

DNA Extraction and Library Preparation

For each coral branch, the top layer of tissue was scraped from the coral skeleton with a

razor blade. DNA was then extracted from tissue using the NucleoSpin Tissue Mini kit columns

and corresponding protocol for extraction from animal tissue (Macherey-Nagel, Duren,

Germany). For each sperm pool, the tube containing RNAlater and sperm was vortexed

vigorously, then 200-400 ul of the sperm solution was pipetted out and mixed with 2x volume of

D.I. water. The sperm pools were then centrifuged for 3 minutes at 13,000 rpm. The supernatant

was pipetted off, leaving just the pelleted sperm at the bottom of the tube. DNA was extracted

from sperm pellets using the same Macherey-Nagel NucleoSpin Tissue Mini kit columns and

protocol as the parent tissue. Nextera full genome libraries were generated using a modified,

low-volume protocol optimized for coral DNA (Supplementary Methods). We constructed two

replicate libraries for each DNA extraction (Fig. 1b). Libraries were sequenced first on an iSeq

100 for quality control and then on a NovaSeq 6000 S4 at the Chan-Zuckerberg Biohub

Sequencing facility in San Francisco, CA, USA.

Reference genome assembly

In May 2020 we collected sperm from an additional *Acropora hyacinthus* colony for the

construction of a high quality *Acropora hyacinthus* reference genome assembly. This colony

originated in Palau and spawned at the California Academy of Sciences, where the sperm was

collected. Sperm was collected by pipetting, then it was rinsed and spun in seawater 3 times at

13,000 rpm for 3 minutes each spin (following methods from [25]). The cleaned sperm pellet was

then flash frozen in liquid nitrogen. The frozen sperm pellet was shipped to Dovetail Genomics

(Scott's Valley, CA, USA) for DNA extraction, sequencing, and genome assembly. The initial *de*

novo assembly was produced through a combination of Illumina short-read sequencing and

PacBio long-read sequencing. Proximity ligation was achieved with Dovetail™ Omni-C™

Technology, which uses a sequence-independent endonuclease approach to chromatin

fragmentation. The final genome assembly is made up of 908 scaffolds, of which 14 represent

full chromosome-length scaffolds, the same number of chromosomes as is in the *Acropora*

millepora genome [26]. The complete assembly is 446,422,234 nucleotides, with $N_{50} = 26,527,962$

nucleotides.

Reference genome annotation

Genome annotation was performed using MAKER2[27] in a *de novo*, iterative approach

based on https://gist.github.com/darencard/bb1001ac1532dd4225b030cf0cd61ce2.

Transcriptome evidence from *Acropora hyacinthus*[28] (and https://matzlab.weebly.com/data--

code.html), *Acropora millepora*[26,29], and *Acropora tenuis* (https://matzlab.weebly.com/data--

code.html) was provided for the initial round of annotation. Additionally, proteome evidence

from *Acropora digitifera*[30] and *Acropora millepora*[26] was utilized for the first round. Genome

wide repeat families were annotated by RepeatModeler2.0.1[31] and used as evidence for the initial

round. The *ab initio* gene predictors AUGUSTUS v3.2.3[32] and SNAP[33] were trained with the

gene models annotated by the previous round of annotation. The second round was then

conducted with these trained prediction models along with repeat, transcript, and protein

evidence annotated during the previous round. A third round of annotation was then performed

following the same procedures as round two. Following the final round, the completeness and

quality of the annotated transcriptome was assessed with BUSCOv5[34] and the OrthoDB v10[35]

eukaryota and metazoan datasets. The BUSCO score against the metazoan dataset was 71.3%

complete, 13.6% fragmented, and 15.1% missing (Supp. Table 3).

Read mapping and SNP calling

Adapters were trimmed from reads using Trimmomatic version 0.39. Trimmed reads

were mapped to the *Acropora hyacinthus* v1 genome using hisat2 with the parameters --very-

sensitive --no-spliced-alignment. Duplicate reads were removed with Picardtools

MarkDuplicates. Haplotype calling was performed with the Genome Analysis Toolkit version

4.1.0.0 Haplotypecaller tool[36]. We combined GVCFS from the same coral colony into a multi-

sample GVCF using CombineGVCFs. Joint genotype calling was then performed on each mutli-

sample GVCF using GenotypeGVCFs with the option –all-sites to produce genotypes for both

variant and nonvariants sites[37]. The genotype-called multi-sample VCFs were filtered with

SelectVariants to filter files by depth, with minimum depth and maximum depth determined by a

Poisson distribution of the average depth for a given sample, with p <0.0001[38]. The filtered files

resulting from these steps were considered the "callable" regions of the genome, and were used

as the denominator for mutation frequency calculations. We filtered for just biallelic single

nucleotide polymorphisms (SNPs) using VCFtools. For the complete read mapping and SNP

calling pipeline see **https://github.com/eloralopez/CoralGermline**

Identifying post-embryonic single nucleotide variants (SNVs)

Single nucleotide variants from the genotyped colony VCFs using custom Python3 and R

scripts (**https://github.com/eloralopez/CoralGermline).** Putative post-embryonic SNVs were

identified by comparing the parent branch genotype calls from a given colony. A SNP was called

a putative post-embryonic SNV if the SNP a.) appeared in just one branch of the colony, and b.)

the SNP had the same genotype call in both replicate libraries from that mutant branch (Fig 1c,

d).

Germline mutations were identified by comparing the sperm genotype calls from a given

colony. A SNP was called a putative unique germline mutation if the SNP a.) appeared in just

one sperm pool spawned by the colony, b.) the SNP had the same genotype call in both replicate

libraries from the sperm pool, and c.) the genotype in the mutant sperm pool did not match the

genotype of the parent branch that spawned it (Fig. 1e). Alternatively, we called a SNP a

putative global germline mutation if the SNP a.) appeared in every replicate library from every

sperm pool spawned by the colony and b.) the genotype in the sperm pools did not match the

genotypes of any of the parent branches in that colony (Fig. 1f).

Classifying putative SNVs

Once we had generated a set of putative somatic and germline mutations, we classified

each SNV as either a Gain of Heterozygosity (GoH) or Loss of Heterozygosity (LoH) mutation,

and classified the directionality of the change (A to T, etc.) as described in [14].

Final filtering of putative mutations to arrive at final set of mutations

The final filtering step was to eliminate putative mutations that had been classified GoH

if the putatively mutant allele had been seen before in any of the other libraries. This step was

necessary because the filtering threshold to call a heterozygotes was 10%. This means that if the

putatively mutant allele was present at <10% allele frequency in other samples, it was not truly a

mutant, but present in the colony at low levels. Similar, we eliminated putative mutations

classified as LoH if the mutant sample was heterozygous at a level <10%. The mutations that

passed this filter is the final set of mutations upon which all subsequent analyses were

performed.

Are parent SNVs shared by their sperm pool or not?

Once we arrived at a set of somatic mutation with all filters applied, we checked to see if

the genotype of the mutant parent matched the genotype of the sperm pool that came from that

branch. GoH SNVs were considered SHARED if one or both genotypes of the two sperm pool

replicates spawned by the mutant branch matched the mutant genotype and NOT SHARED if

neither of the genotypes of the two sperm pool replicates spawned by the mutant branch matched

the mutant genotype (Fig. 1c, d). LoH SNVs were considered SHARED if both genotypes of the

two sperm pool replicates spawned by the mutant branch matched the mutant genotype and NOT

SHARED if one or neither of the genotypes of the two sperm pool replicates spawned by the

mutant branch matched the mutant genotype (Fig. 1c, d).

Designating SNV effects on codons

We classified each mutation by the genomic region (intron, exon, etc.) it fell in and, if it

fell in a coding region, the type (synonymous, missense) using the program snpEff[39] configured

with the *Acropora hyacinthus* version 1 genome. We calculated the rate of missense and

synonymous SNVs per bp in the coding region by dividing the number of SNVs by the total

callable coding region. We calculated the average percent of coding SNVs that were missense

per sample for three categories: Parent Only, Parent and Sperm, and Single Sperm Pool Only.

There were no coding SNVs in the All Sperm Pools category (Extended Data Fig. 6).

Selection on mutations

We calculated dN/dS, a measure of selection on the genome, using the R package

dndscv[21]. We first created a reference CDS file for the *Acropora hyacinthus* genome using the

Acroporal hyacinthus genome annotations (gff3) file[26] and a custom Python3 script. We then

created *dndscv* input files from the filtered set of mutations found in each coral colony. We ran

dndscv with theta = 0.

Mutation rates

To catalogue post-embryonic SNVs, we compared sequences from 3 parent branches in

each colony and recorded cases in which one branch showed a genotype different from all the

others. This catalogue includes true somatic mutations in that branch, but also includes any

sequencing or PCR errors injected during sample preparation and sequencing. To limit these

errors, we compared sequences from each technical replicate and catalogued a mutation only

when it was visible in both replicates of a branch and in none of the replicates of any other

branch. Comparing technical replicates was highly successful at reducing noise from sequencing

error, eliminating over 90% of putative SNVs that would have been identified using the same

pipeline and filters without technical replicates (Extended Data Figs. 2, 3). To find the average

frequency per nucleotide of somatic mutations unique to a given branch in a coral colony, we

divided the number of SNVs in a sample by the number of total callable nucleotides sequenced

for that sample (Supplementary Table 1).

$$\text{\# SNV per Mb} = (\text{\#SNVs} / \text{Callable region (bp)}) * 1{,}000{,}000$$

The callable genome sizes were 1.2×10^8 bp, 1.2×10^8 bp, and 0.80×10^8 bp and the

callable coding region sizes were 1.3×10^7 bp, 1.3×10^7 bp, and 0.91×10^7 for each of the three

colonies, CA56, CA60, and CA65, respectively (Supp. Table 1).

Tissue mixtures

Because DNA was extracted from tissue scrapings encompassing multiple polyps per

parent branch, we were initially concerned that the parent sample might be a heterogenous mix

of somatic and germ tissues. Germ and stem cells in other cnidarians tend to reside at the base of

the polyp, so the scraping method was intended to take off just the somatic tissue and not the

germ or stem cells. To check this, we plotted the average variant allele frequency (VAF) of the

mutant parent (that is, the average VAF for the two technical replicate libraries) against the

average VAF of the mutant sperm pool.

$$\text{VAF} = \text{\# of reads supporting SNV} / \text{total \# of reads at locus}$$

$$\text{Average VAF} = (\text{VAF}_1 + \text{VAF}_2)/2$$

If the parent samples were a mix of somatic and germ and/or stem cells, then a germline

mutation may have erroneously been called a somatic mutation. If that were the case, then the

VAF of the parent would be considerably lower than the VAF of the sperm pool, because the

sperm pool's mutant would not be diluted by nonmutant somatic tissue. In that case, we would

expect the trendline of the parent VAF: sperm VAF linear model to have a slope significantly

greater than 1. In reality, the slope of the trendline was less than 1, suggesting that we

undercounted the number of mutant reads in the sperm. This gives us confidence that the parent

samples were in fact all or almost all somatic tissue, and that the mutations found in the parents

were in the soma. We used a linear regression model to check the relationship between the

variant allele frequency of the mutant parent and the variant allele frequency of the mutant sperm

for all inherited GoH mutations, where variant allele frequency equals the number of reads with

mutation divided by the total number of reads. The slope of the trendline was 0.36, with $R = 0.3$

and $p = 5.9e\text{-}08$ (Extended Data Fig. 7). This suggests that there may have been some

undercounting of the number of mutant reads in the sperm. The distribution of average VAF in

the parents for SNVs classified as "Parent Only" has a larger leftward skew, which also suggests

that some SNVs found at low frequency in the parent samples may have been missed in the

sperm (Extended Data Fig. 8). This gives us confidence that the parent samples were in fact all

or almost all somatic tissue, and that the mutations found in the parents were in the soma.

Types of mutations

For Gain of Heterozygote mutations (where the mutant genotype is a novel heterozygous

SNP) we classified the somatic mutation as inherited by the sperm if at least one of the technical

replicate libraries of the sperm contained the mutant allele. If neither sperm library contained the

mutant allele, then it was considered not inherited by the sperm. For Loss of Heterozygosity

mutations (where the mutant is a homozygous genotype at a site for which the parent colony is

heterozygous) to be classified as inherited by the sperm both sperm libraries had to be entirely

homozygous for the allele of the mutant parent. For these cases, if any read for a parent sample

showed the minor allele also seen in the putative GoH mutation, we did not call these as

mutations. Likewise, if any of the sperm reads showed the minor allele also seen in the parent

branch, we did not call this a LoH mutation. These stringent filters reduced the number of

mutations we could characterize but led to high confidence in our characterization of variants.

Data availability

All raw fastq files, as well as the *Acropora hyacinthus* genome version 1 assembly, are

accessioned under BioProject PRJNA707502 at NCBI. The accession numbers for the fastqs are

SAMN18207983-SAMN18208014 and the accession number for the assembly FASTA is

SAMN20335437.

Code availability

The code used for this study can be found at **https://github.com/eloralopez/CoralGermline**.

Methods References

24.   Craggs, J. *et al.* Inducing broadcast coral spawning ex situ: Closed system mesocosm

472   design and husbandry protocol. *Ecol. Evol.* **7**, 11066–11078 (2017).

473   25.   Shumaker, A. *et al.* Genome analysis of the rice coral Montipora capitata. *Sci. Rep.* **9**, 1–

474   16 (2019).

475   26.   Fuller, Z. L. *et al.* Population genetics of the coroal Acropora millepora: Toward genomic

476   prediction of bleaching. *Science (80-.).* **369**, (2020).

477   27.   Holt, C. & Yandell, M. MAKER2: An annotation pipeline and genome-database

478   management tool for second-generation genome projects. *BMC Bioinformatics* **12**, 1–14

479   (2011).

480   28.   Barshis, D. J. *et al.* Genomic basis for coral resilience to climate change. *Proc. Natl. Acad.*

481   *Sci. U. S. A.* **110**, 1387–92 (2013).

482   29.   Moya, A. *et al.* Whole Transcriptome Analysis of the Coral Acropora millepora Reveals

483   Complex Responses to CO 2-driven Acidification during the Initiation of Calcification.

Mol. Ecol. **21**, 2440–2454 (2012).

30.   Shinzato, C. *et al.* Using the Acropora digitifera genome to understand coral responses to

environmental change. (2011). doi:10.1038/nature10249

31.   Flynn, J. M. *et al.* RepeatModeler2 for automated genomic discovery of transposable

element families. *Proc. Natl. Acad. Sci. U. S. A.* **117**, 9451–9457 (2020).

32.   Stanke, M. & Waack, S. Gene prediction with a hidden Markov model and a new intron

submodel. in *Bioinformatics* **19**, 215–225 (Oxford Academic, 2003).

33.   Korf, I. Gene finding in novel genomes. *BMC Bioinformatics* **5**, 1–9 (2004).

34.   Simão, F. A., Waterhouse, R. M., Ioannidis, P., Kriventseva, E. V. & Zdobnov, E. M.

BUSCO: Assessing genome assembly and annotation completeness with single-copy

orthologs. *Bioinformatics* **31**, 3210–3212 (2015).

35.   Kriventseva, E. V. *et al.* OrthoDB v10: Sampling the diversity of animal, plant, fungal,

protist, bacterial and viral genomes for evolutionary and functional annotations of

orthologs. *Nucleic Acids Res.* **47**, D807–D811 (2019).

36.   Poplin, R. *et al.* Scaling accurate genetic variant discovery to tens of thousands of

samples. *bioRxiv* 201178 (2017). doi:10.1101/201178

37.   Van der Auwera, G. A. *et al.* From fastQ data to high-confidence variant calls: The

genome analysis toolkit best practices pipeline. *Curr. Protoc. Bioinforma.* **43**, 11.10.1-

11.10.33 (2013).

38.   Wu, F. L. *et al.* A comparison of humans and baboons suggests germline mutation rates

do not track cell divisions. *PLoS Biol.* **18**, e3000838 (2020).

39.   Cingolani, P. *et al.* A program for annotating and predicting the effects of single

nucleotide polymorphisms, SnpEff: SNPs in the genome of Drosophila melanogaster strain

w1118; iso-2; iso-3. *Fly (Austin).* **6**, 80–92 (2012).

Acknowledgements

EHLN was funded by NSF GRFP and a Stanford Graduate Fellowship in Science and

Engineering (Morgridge Family Foundation). Research funded by Chan-Zuckerberg BioHub,

NSF OCE-1736736, and the Hope for Reefs Initiative. We thank N. Neff at CZ-BioHub for

sequencing power and A. Voskoboynik, P. Bump, and B. Cornwell for stem cell insights. Thanks

to R. Ritson-Williams, R. Ross, and BIOTA for field collection support, and thanks to the staff

of the Steinhart Aquarium for taking care of the corals.

Author contributions

EHLN, RA, and SRP designed the study. RA kept corals alive and healthy and facilitated lab

spawning. EHLN and RA collected parent and sperm samples for this study. EAS optimized the

library preparation protocol used in this study. EHLN extracted DNA and constructed libraries

for all samples. EHLN designed and performed all bioinformatic analyses for the study, except

for genome annotation, which was performed by EAH. EHLN wrote the manuscript and all

authors provided input and feedback on drafts.

Competing interest declaration

The authors have no competing interests to declare.

Additional Information

Supplementary information is available for this paper:

Supplementary Table 1. Summary of sequencing depth and SNV counts (.txt file)

Supplementary Table 2. List of verified SNVs and their metadata (.txt file)

Supplementary Table 3. Summary statistics for *Acropora hyacinthus* version 1 genome

annotations (.xlsx file)

Supplementary Methods. The optimized full genome library preparation protocol for

corals (.docx file).

Correspondence and requests for materials should be addressed to Elora H. López-Nandam,

elopez-nandam@calacademy.org

Extended data figure/table legends

535
536

537
538   **Extended Data Figure 1.** Photos of the experimental setup. a.) Colony 60, with sampled
539   branches denoted by numbered washers. For scale, the outside diameter of each washer is 1.59
540   cm. b.) Two of the branches (CAP6 and CAP8) sampled from Colony 56, releasing gamete
541   bundles (indicated by arrows) into their respective 5 ml tubes of seawater.
542

543
544 **Extended Data Figure 2.** The mean number of mutations identified in each parent branch in the
545 Parent Only (PO) or Parent and Sperm (PO), or single sperm pool only (SSPO) when looking at
546 one library per sample (aka without technical replicates, pink) or two libraries per sample (with
547 technical replicates, gray). Error bars represent ± 1 s.e.m. Each data point (N=7) is shown as a
548 smaller point for each category. Note that the y axis is on a log-10 scale.

549
Extended Data Figure 3. The percentage of mutations identified when replicate libraries are
included for each of the four categories of SNVs: Parent Only (PO), Parent and Sperm (P+S),
Single Sperm Pool Only (SSPO), and All Sperm Pools (ASP). Error bars represent ± 1 s.e.m.
Each data point (N=7) is shown as a smaller point for each category. The percentage is expressed
as a fraction of the number of mutations identified when no technical replicates are included,
such that:
% = (# of mutations identified using technical replicates*100) / (# of mutations identified
without technical replicates)

559
560   **Extended Data Figure 4**. Distribution of average GQ score and average read depth for each
561   mutation in the colony (top) as well as distributions of the lowest GQ score and lowest read
562   depth for every mutation identified in each colony (bottom). Mutations found when using no
563   technical replicates are shown in pink, and the mutations found when including technical
564   replicates shown in gray.
565

566
567   **Extended Data Figure 5.** SNV spectra from various subsets of SNVs. None of the SNV spectra
568   were significantly different each other for the different data subsets (see X^2 tests below). Error
569   bars represent ± 1 s.e.m.
570   X^2 tests of independence results for Extended Data Figure 5:

Comparison	X^2
Parent only vs. Shared Parent and Sperm	df = 6, N=1715, X^2 = 7.3495, p= 0.2897
Single Sperm Pool Only vs. All Sperm Pools	df = 6, N=604, X^2 = 4.2981, p= 0.6364
Parent GOH SNVs vs Parent LOH SNVs	df = 6, N=1715, X^2 = 3.9411 p= 0.6846

Extended Data Figure 6. Rates and percentages of missense and synonymous mutations across
sample and category. a.) The number of missense (circle) and synonymous (triangle) SNVs per
bp of the coding region for each category (all somatic, P+S, PO, and SSPO) for each sample.
Dashed lines indicate the mean number of coding SNVs (the sum of missense and synonymous)
for each category across all samples. b.) The mean percent missense for each category, with error
bars indicating ± 1 s.e.m. Each data point (N=7) is shown as a smaller point for each category.

Extended Data Figure 7. Variant allele frequencies for parents and their respective sperm pools.
a.) GOH mutations for which sperm VAF =0 were classified as Parent Only (black), likewise
LOH variants in the parent for which the VAF < 1 in the sperm were classified as Parent Only.
All other SNVs were classified as being shared by both a parent branch and its corresponding
sperm pool (red). B.) For GOH SNVs that are shared by a parent branch and its sperm, average
variant allele frequency of the two replicate parent libraries is positively correlated with the
average variant allele frequency of the two replicate sperm pool libraries. The slope of the
relationship is 0.36 and R=0.3 (red line). The black line shows a 1:1 line for reference.
589

590
591   **Extended Data Figure 8.** The distribution of variant allele frequencies in the parent branches for
592   GOH SNVs in the parents only (orange) and in the parent and sperm samples (gray). The mean
593   parent variant allele frequency for each category is marked by dashed lines.
594
595
596   **Extended Data Table 1.** Comparison of SNV frequencies in the coding region vs. across the full
597   genome for six SNV types (shown in Figure 2c). Error shown is ±1 s.e.m.
598

Type	Average frequency per bp (Full genome, coding)	Wilcox test (two-tailed, paired)
Parent Only, GOH	Full: $4.4 \times 10^{-7} \pm 0.6 \times 10^{-7}$ Coding: $9.2 \times 10^{-8} \pm 3.1 \times 10^{-8}$	V=28, P= 0.016
Parent Only, LOH	Full: $1.32 \times 10^{-6} \pm 0.19 \times 10^{-6}$ Coding: $5.2 \times 10^{-7} \pm 1.4 \times 10^{-7}$	V=28, P= 0.016
Parent and Sperm, GOH	Full: $4.8 \times 10^{-7} \pm 0.9 \times 10^{-7}$ Coding: $1.8 \times 10^{-7} \pm 0.6 \times 10^{-7}$	V=27, P= 0.031
Parent and Sperm, LOH	Full: $2.6 \times 10^{-7} \pm 1.1 \times 10^{-7}$	V=28, P= 0.016

	Coding: $1.6 \times 10^{-8} \pm 1.6 \times 10^{-8}$	
Single Sperm Pool Only, GOH	Full: $3.8 \times 10^{-7} \pm 0.5 \times 10^{-7}$ Coding: $1.5 \times 10^{-7} \pm 0.5 \times 10^{-7}$	V=28, P= 0.016
Single Sperm Pool Only, LOH	Full: $3.6 \times 10^{-7} \pm 0.4 \times 10^{-7}$ Coding: $1.6 \times 10^{-7} \pm 0.6 \times 10^{-7}$	V=27, P= 0.031

602
603

604