Mixed culture metagenomics of the microbes making sour beer

Renan Eugênio Araujo Piraine [1,2], Fábio Pereira Leivas Leite [2] and Matthew L. Bochman [1,3]*

[1] Bochman Lab, Department of Molecular and Cellular Biochemistry, Indiana University, Bloomington, 47405,
IN, United States; renanbiotec@gmail.com (R.E.A.P); bochman@indiana.edu (M.L.B.)
[2] Laboratório de Microbiologia, Centro de Desenvolvimento Tecnológico, Universidade Federal de Pelotas,
Pelotas, 96010-610, RS, Brazil; fleivasleite@gmail.com (F.P.L.L)
[3] Wild Pitch Yeast, Bloomington, IN 47405, USA.
* Correspondence: bochman@iu.edu; Tel.: +1-812-856-2095

Abstract: Mixed microbial cultures create sour beers, but many brewers do not know which mi-
crobes comprise their cultures. The objective of this work was to use deep sequencing to identify
microorganisms in sour beers brewed by spontaneous and non-spontaneous methods. Twenty
samples were received from brewers, which were processed for microbiome analysis by next gen-
eration sequencing. For bacteria, primers were used to amplify the V3-V4 region of the 16S rRNA
gene; fungal DNA detection was performed using primers to amplify the entire internal tran-
scribed spacer region. The sequencing results were then used for taxonomy assignment, sample
composition, and diversity analyses, as well as nucleotide BLAST searching. We identified 60
genera and 140 species of bacteria, of which the most prevalent were *Lactobacillus acetotolerans*,
Pediococcus damnosus, and *Ralstonia pickettii/mannitolilytica*. In fungal identification, 19 genera and 26
species were found, among which the most common yeasts were *Brettanomyces bruxellensis* and
Saccharomyces cerevisiae. In some cases, genetic material from more than 60 microorganisms was
found in a single sample. In conclusion, we were able to determine the microbiomes of various
mixed cultures used to produce beer, providing useful information to better understand the sour
beer fermentation process and brewing techniques.

Keywords: microbiome, mixed-fermentation, sour, beer, yeast, bacteria

1. Introduction

Traditionally, beer production methods are divided into two categories: (a) fer-
mentation at the bottom of the fermenter, performed by *Saccharomyces pastorianus* in lager
beer production; and (b) top-fermented beers, in which *S. cerevisiae* yeast ferments at the
top of the wort, producing ales. Extending the concept to mixed fermentations, two new
categories can be included: (c) non-spontaneous fermentation, carried out by an in-house
starter culture, which consists of yeast and lactic acid bacteria (LAB); and (d) spontane-
ous fermentation, in which microorganisms such as enterobacteria, yeasts, mold, LAB,
and acetic acid-producing bacteria (AAB) among others are inoculated through exposure
to ambient air or external sources (*e.g.*, wood, flowers, or fruits) to ferment these beers
[1,2].
Mixed fermentations are generally performed by yeasts and LAB in the process of
creating sour beers, forming a complex microbiota that acts through their interaction and
cooperation [3]. Microorganisms and their enzymes are used through biotechnological
processes for acidification, alcoholization, proteolysis, lipolysis, and amino acid conver-
sion in beer worts [1]. Various phases of fermentation can be characterized in mixed
fermentation beers, in which different bacteria and yeasts are isolated at specific periods
[4]. Changes in the presence and concentrations of various microorganisms suggest the
existence of a microenvironment regulated according to substrate conversion and growth
limiting factors, such as pH, carbohydrate concentration, oxygen, temperature, and al-

cohol concentration.

The global beer market is experiencing a resurgence in the interest in sour beers because new products and more complex flavors are being obtained by large and small production breweries around the world [5]. However, most brewers do not know exactly which microorganisms are present in their mixed culture fermentations, as well as the relative proportions of these microbes in their starter cultures. Even though wild yeasts, LAB, and some Gram-positive bacteria are considered contaminants in the vast majority of beer fermentations, these same microorganisms are often highly desired for the production of specific sour and wild beer styles. As an example, Lambics and American Coolship Ales are beverages with unique flavor profiles generated by "spoilage" organisms [6]. Thus, it is of interest to know the microorganisms present in these fermentative processes, seeking to characterize the microbial diversity, parameters that influence sensory parameters, and to deepen the knowledge of mixed fermentation beers. The objective of this study was to identify bacteria and yeasts present in different beers and mixed culture samples from several locations, produced by homebrewers and craft breweries, using spontaneous and non-spontaneous fermentation methods.

2. Materials and Methods

2.1. Samples

The samples used in this study were obtained through the crowdfunding project "Mixed culture metagenomics of the microbes making sour beer" (DOI 10.18258/13495) hosted on the Experiment platform (www.experiment.com). Each brewer collected and sent their own sample in glass beer bottles or plastic vials containing beer or slurry, which were stored at 4°C until their analysis.

During July-December 2019, 20 samples were received, including spontaneous and non-spontaneous fermentation beers produced by homebrewers and craft breweries, as well as some samples from house cultures propagated to ferment these beers. Samples were obtained from different regions, including countries such as Canada, the United States, and Israel. Furthermore, these samples had different maintenance times, ranging from a few months of storage to more than 5 years of use and propagation by the brewer. Non-spontaneous fermentation samples originated from commercial blends, bottle dregs, or from cultures that were propagated and maintained by brewers, of which the exact microbial composition was not known. Spontaneous fermentation samples originated in their great majority through the process of exposing the beer wort to the ambient air using open fermenters. In some of these samples, it was observed that the brewers inoculated the beer using an external source with the possible presence of microorganisms, such as wood, flowers, or fruits. Details concerning the individual samples can be found in the Supplementary Information file S1.

In the case of beer, 50 mL was processed by centrifugation at 2,000 × g for 10 min at 4°C, and the supernatant was decanted. For mixed culture samples, a small volume (≤5 mL) was resuspended in 50 mL sterile water and treated in the same manner. Then, the pelleted cells and debris were resuspended in 500 µL of 2x DNA/RNA Shield (Zymo Research (Irvine, CA) and stored at –20°C. Samples were then submitted for microbiome analysis through the ZymoBIOMICS® Targeted Sequencing Service for Microbiome Analysis by the Zymo Research company.

2.2. DNA extraction and sequencing

DNA extraction was performed using a ZymoBIOMICS® -96 MagBead DNA kit (Zymo Research, Irvine, CA) or ZymoBIOMICS® DNA Miniprep kit (Zymo Research, Irvine, CA) with an automated extraction platform. Bacterial and fungal identifications were performed with 10% PhiX spike-in using next generation sequencing (NGS) on an Illumina® MiSeq™ system, with a v3 reagent kit (600 cycles). First, targeted libraries were prepared for both groups. For bacteria, 16S ribosomal RNA gene-targeted se-

quencing was performed using a Quick-16S™ NGS Library Prep Kit (Zymo Research, Irvine, CA), in which 16S primers were used to amplify the V3/V4 region of the 16S rRNA gene, maintaining good coverage and high sensitivity. Fungal internal transcribed spacer (ITS)-targeted gene sequencing was performed using the same kit described above, though ITS2 primers were used instead 16S primers, which amplify the entire ITS region and allow molecular phylogenetic sequence identification for many fungi.

PCR reactions were performed to prepare the sequencing library, controlling cycles and limiting PCR chimera formation. The quantification of final PCR products was performed with qPCR fluorescence readings and pooled together based on equal molarity. Cleaning and quantification of final pooled libraries was performed using Select-a-Size DNA Clean & Concentrator™ (Zymo Research, Irvine, CA), TapeStation® (Agilent Technologies, Santa Clara, CA), and Qubit® (Thermo Fisher Scientific, Waltham, WA). Negative controls (blanks) were used during all processes, as well as the ZymoBIOMICS® Microbial Community Standard (Zymo Research, Irvine, CA) as a positive control, which mimics a mixed microbial community of well-defined composition, containing Gram-negative and Gram-positive bacteria and yeasts. Additional information can be found at the ZymoBiomics™ Service website (https://www.zymoresearch.com/pages/microbiome-analysis-services).

2.3. Bioinformatics analyses

Sequencing results were used for taxonomy assignment, sample composition visualization, and alpha- and beta-diversity analyses. Unique amplicon sequence variants (ASVs) were inferred from raw reads using the DADA2 pipeline [7]. The ASVs of bacteria and fungi were used to create a 3-dimensional principle component analysis (PCoA) plot using the matrix of paired-wise distances between samples, calculated by the Bray-Curtis dissimilarity. Uclust from Qiime v. 1.9.1 was used for taxonomy assignment using the Zymo Research Database, a 16S and ITS database that was internally designed and curated, as a reference. Results were re-analyzed by amplified sequence alignment using the nucleotide collection database from NCBI (National Center for Biotechnology Information, Rockville, MD) and nucleotide BLAST tool. Taxonomy nomenclature and classification was analyzed using the Taxonomy Browser tool on the NCBI website (http://www.ncbi.nlm.nih.gov/Taxonomy/Browser) [8].

Sequencing results were used to separately analyze the phylogenetic relationships between groups of bacteria and fungi with MEGA v.10.1.7 software for alignment, construction, and visualization of phylogenetic trees. Sequence alignment was performed using ClustalW, which was also used to construct neighbor-joining (N-J) phylogenetic trees with 1000 bootstrap trials. Circular trees were used as templates for final figures, which had schemes and colors added using Gimp v.2.10.18 software. The microbial composition for each sample was evaluated using GraphPad Prism v.7 software, which was also used to plot the data. Samples were also analyzed for taxonomy, visualization, and interactive reporting using the Knomics-Biota system [9].

3. Results

Based on the different pipelines used by ZymoResearch and Knomics-Biota, with ASVs examined through the analysis of each amplified fragment using the NCBI nucleotide database, we were able to determine the metagenomes of 20 different beer samples and starter cultures that originated from mixed fermentations made by both spontaneous and non-spontaneous methods. Details about the collected samples are shown in Table 1 and Supplementary Information file S1.

150        **Table 1**. Identification and characteristics of samples of mixed cultures received for the study.

Sample	Origin	Material	Spontaneous fermentation	Fruits, woods, flowers or another microbe source added	Culture maintenance time	Commercial strains inoculated
1	Jerusalem (IL)	Culture pre-pitch	YES	NO	6-12 months	-
2	Alberta (CA)	Beer/slurry	NO	YES	1-6 months	-
3	Alberta (CA)	Beer	YES	YES	6-12 months	New W. Saison (Escarpment Labs) Brett 'M' (Escarpment Labs)
4	Alberta (CA)	Beer	YES	NO	6-12 months	-
5	Alberta (CA)	Beer	YES	YES	6-12 months	-
6	Alberta (CA)	Beer	YES	YES	6-12 months	-
7	Alberta (CA)	Beer	YES	YES	6-12 months	-
8	Alberta (CA)	Beer	YES	YES	6-12 months	-
9	Washington (US)	Culture pre-pitch	YES	NO	2-3 year	-
10	Washington (US)	Culture pre-pitch	YES	NO	6-12 months	-
11	Ohio (US)	Beer/slurry	NO	NO	6-12 months	-
12	Nevada (US)	Beer	NO	YES	7-8 years	Sour Solera (Bootleg Biology); Mélange (The Yeast Bay); BugCounty (East Coast Yeast); dregs from beer bottles
13	Nevada (US)	Beer	YES	YES	3-4 years	Dregs from beer bottles
14	California (US)	Beer	NO	NO	6-12 months	Dregs from beer bottles
15	California (US)	Culture pre-pitch	YES	NO	5-6 years	-
16	California (US)	Beer	NO	NO	4-5 years	-
17	California (US)	Beer	YES	NO	1-6 months	-
18	California (US)	Beer	NO	NO	2-3 years	-
19	California (US)	Beer	NO	NO	1-2 years	WY3763 (Wyeast); WY3711 (Wyeast); WLP650 (White Labs)
20	Michigan (US)	Beer	YES	NO	1-6 months	WLP565 (White Labs); dregs from beer bottles

153 * IL, Israel; CA, Canada; and US, United States.

156       *3.1. Bacteria - 16S rRNA V3-V4 region analysis*

157       3.1.1 Bacterial composition

The analysis of bacteria present in the various samples revealed 60 genera and 120 different species, with some samples containing only one species of bacteria (*e.g.,* samples 15, 16, 17, and 18) and others containing >50 species (*e.g.,* sample 1) (Figure 1). Based on ASV analysis, the most prevalent bacteria were *Lactobacillus acetotolerans*, which was identified in 60% of the analyzed samples (n=12), followed by *Pediococcus damnosus* and *Ralstonia picketii/mannitolilytica*, both identified in 35% of the samples (n=7). Although a large number of bacteria were identified at the species level, we observed that several ASVs did not allow the differentiation of species belonging to the same genus, leading to designation of two different species for the same ASV, such as for *L. collinoides/paracollinoids*. Only six ASVs could not be identified at the genus level, allowing only their classification at the level of phylum, order, or family, as in the case of Myxococales bacteria classification. The microbial composition and raw reads of each sample are available in detail in Supplementary Information file S1, while the total number of bacterial and fungal species identified is shown in Figure 2.

The significant presence of bacteria from the Enterobacteriaceae family was observed in different samples, which could be >15% of the ASV total composition, as in samples 1 and 14, and even surpass 90% of the bacterial composition, as in samples 6, 7, and 20 (Figure A1, Appendix A). However, even though many of these beers were made by spontaneous fermentation and were inoculated with external sources of microorganisms, the presence of this bacterial group was <2% of the total composition of ASVs identified in most samples (n=14).

Figure 1: Bacterial composition of the 20 samples analyzed

. Bacterial identification was performed from ASVs originating from NGS using the V3/V4 region of the 16S rRNA gene. Due to the large number of species found, colors for identification were designated according to genera, with the exception of the Enterobacteriaceae family, which is identified in the graph with the red color only.

Figure 2: Number of species identified in each sample, for bacteria (left) and fungi (right). Metagenome identification of mixed fermentation samples made it possible to identify samples containing from just one to >55 species of bacteria, while for fungi, it was possible to detect samples with only one yeast participating in fermentation, up to more than eight different species acting together during fermentation.

Based on the differences in microbial composition among samples, beta diversity demonstrated through PCoA plots revealed some similarities among samples according to the identified genera (Figure 3A). We highlight four main groups: group A (samples 4, 5, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, and 18), in which the genus with the highest prevalence was *Lactobacillus*; group B (samples 6, 7, and 20), in which samples were composed mainly by Enterobacteriaceae bacteria; group C (samples 1 and 2), in which *Pediococcus* (*Pediococcus damnosus* exclusively) was the genus with the highest proportion; and group D (samples 3 and 19), in which the microbial composition revealed a large presence of other ASVs detected, such as *Gluconobacter* spp. and *Ralstonia* spp. Similarities observed in bacterial composition could also be seen in the phylogenetic tree generated through the Knomics-Biota pipeline , which grouped samples in a similar way (Figure 3B). Note that the similarity between samples 1 and 11 was due to specific ASVs of *P. damnosus* present only in these two samples.

(a) (b)

Figure 3: **Sample similarity based on bacterial composition**. (**a**) PCoA tridimensional plot created using the matrix of pair-wise distances between samples calculated by the Bray-Curtis dissimilarity using genera found in the microbial composition. 3D images, schematical representation, and visualization obtained using the EMPeror tool. Below the PCoA plot is a schematic representation based on sample composition, where arrows show different directions for locating the samples on the graph, according to the presence and concentration of Enterobacteriaceae bacteria, *Lactobacillus* spp., *P. damnosus* and other bacteria such as *Ralstonia* spp. and *Gluconobacter* spp. (**b**) Phylogenetic tree showing clustering of the samples by similarity of their taxonomic composition, calculated by the Bray-Curtis dissimilarity at the ASV level, using Ward's method.

3.1.2. Phylogenetic analysis

The construction of phylogenetic trees (Figure 4a) showed that *Lactobacillus* spp. and *Pediococcus* spp. were the main microbes from the LAB, a group of great importance in the production of sour beers. At least nine different *Lactobacillus* spp. were detected, including *L. brevis*, *L. plantarum*, *L. casei/paracasei*, *L. backii*, *L. buchneri*, *L. lindneri*, and *L. delbrueckii* in addition to those mentioned above. For *Pediococcus* spp., only *P. damnosus* was identified, but the presence of possible subspecies could be responsible for the multiple different ASVs found. Also belonging to the LAB group, bacteria in the genera *Leuconostoc*, *Weissela*, and *Aerococcus* were likewise detected (with lower prevalence), as well as bacteria with potential probiotic activity, such as *Bifidobacterium* spp. and *Bacillus* spp. Other phylogenetically distinct groups of bacteria also demonstrate important participation in spontaneous and non-spontaneous fermentations, such as the AAB *Gluconobacter* spp., and *Acetobacter* spp., especially *G. oxydans*, and *A. pasteurianus*.

Several species of the genus *Pseudomonas* were also detected, mainly in sample 20, in which it was possible to identify eight different ASVs related to this genus, corresponding mainly to *P. fluorescens*. Although *Pseudomonas* spp. are mainly known because of the pathogenic bacterium *P. aeruginosa*, this organism was not detected in the studied samples. A special emphasis was given to the Enterobacteriaceae family in the phylogenetic tree, as they correspond to a large number of identified ASVs and thus represent a vast

number of species found at different concentrations. The samples' metagenomes showed the presence of a large number of genera included in this family, among them *Enterobacter* spp., *Salmonella* spp., *Klebsiella* spp., and *Pantoea* spp. During the identification of the ASVs corresponding to these bacteria, it was possible to observe a high similarity among the genera, which made specific identification difficult in many cases, especially at the species level.

Analyzing the metagenomes discovered (Figure 4b), three main phyla were identified with high prevalence: Proteobacteria (66.67% of the identified bacteria belonged to this phylum), Firmicutes (19.38%), and Actinobacteria (11.63%). ASVs from other phyla were also detected, though at smaller proportions: Deinococcus-Thermus (0.78%), Cyanobacteria (0.78%), and Planctomycetes (0.78%). Almost half of the identified bacteria belonged to the Gammaproteobacteria class, which includes families such as Enterobacteriaceae, Pseudomonadaceae, and Acetobacteriaceae among others, totaling 17 different bacterial families. Though most studies and applications involving mixed fermentation beers focus on the Lactobacillaceae family, it was related to only 7.75% of ASVs, representing a small portion of the variety of microorganisms identified.

(b)

Figure 4: Phylogenetic analysis and abundance of bacterial taxa. (**a**) Phylogenetic tree constructed based on the ASVs of 120
bacterial species found in the mixed culture samples. Phylogenetic relationships were made using MEGA v.10.1.7 software for
alignment, construction, and visualization of the phylogenetic tree. Sequence alignment was performed using ClustalW. (**b**)
The image shows the prevalence of taxonomic classifications according to genera and species found in the bacterial
microbiomes. Higher taxonomic classifications are shown in circles near the image center, while lower classifications are
shown toward the outer edge. Images were constructed using MEGA and GIMP 2.1 software.

3.2. Fungi – ITS2 region analysis

3.2.1. Fungal composition

NGS targeting the ITS2 region was able to detect 19 genera and 26 different species
of fungi in the 20 analyzed samples (Figure 2). Among them, yeasts were the most iden-
tified microorganisms, while ASVs from filamentous fungi, molds, and more complex
fungi (*e.g.,* mushrooms) were also be identified in smaller proportions. Samples were
basically dominated by two main yeasts: *Brettanomyces bruxellensis* and *S. cerevisiae*, which
were present in 75% (n=15) and 65% (n=13) of the samples, respectively (Figure 5). Mi-
crobial composition analyses revealed that in the vast majority samples (n=16), fermen-
tation was carried out by at least two different species of fungi. Of these, an association
between *Brettanomyces* spp. and *Saccharomyces* spp. was shown to be recurrent (n=7), in
which both exceeded at least 5% concentration in the ASVs found in each sample.
Other yeasts, such as *Issatchenkia orientalis* (also known as *Pichia kludriavzevii, Can-*
dida krusei, and *Candida glycerinogenes*) and *Wickerhamomyces anomalus* (or *Pichia anomala*)
also participated in these mixed fermentation beers, which could be detected in 30% (n=6)
and 25% (n=5) of the samples, respectively, with a large presence in samples 3, 6, 7, and
10. It is interesting to note that not only were yeasts identified but also more complex
fungi such as *Penicillium* spp., which were identified in most of the ASVs found in sample
14. Overall, we observed that several fungal species were often present in mixed culture
samples, with more than seven species found in samples 11 and 18, and over 11 different
fungi detected in sample 3.
405

Figure 5: **Fungal composition of the 20 samples analyzed**. Fungal identification was performed from ASVs originating from NGS of the ITS2 region, amplifying the entire ITS region. Colors for identification were designed according to fungal species as shown below the plot.

The beta diversity of the samples was evaluated according to the genera identified in the microbial composition of each sample. In these analyses, we could verify the distribution of samples into four groups (Figure 6a) by similarity between identified microorganisms: group A (samples 4, 9, 15, 16, and 20) where > 95% of the sample was composed of *S. cerevisiae*; group B (samples 1, 12, 13, 17, 18, and 19) in which > 90% of the identified ASVs corresponded to *Brettanomyces* spp.; group C (samples 2, 5, 8, and 11) in which approximately 70% of the sample was represented by *S. cerevisiae*, 25% *Brettanomyces* spp., and 5% other fungi; and group D (samples 3, 6, 7, 10, and 14) in which other genera such as *Wickerhamomyces, Issatchenkia, Penicillium,* and *Lachancea* dominated.

The tree constructed based on the hierarchical clustering between ASVs detected for ITS2 (Figure 6b) reveals that samples tend to follow the same relationship observed in Figure 6a, though some samples such as 13 and 14 had greater hierachical approximation because similar ASVs referring to *Brettanomyces custersianus* were found in significant proportions (86% and 10% of the ASVs identified in these samples, respectively). Similarly, samples 9 and 11, though organized in different groups in Figure 6a, presented a phylogenetic relationship of proximity in this analysis because there were specific ASVs corresponding to *B. bruxellensis* and *Brettanomyces anomalus* found exclusively in their microbial composition. The other samples displayed phylogenetic relationships in accordance with those observed in the PCoA analysis, confirming similarities in fungi composition among the samples evaluated in this study.

(a)   (b)

Figure 6: **Sample similarity based on fungal composition**. (**a**) PCoA tridimensional plot created using the matrix of pairwise distances between samples calculated by the Bray-Curtis dissimilarity using the genera found in the microbial composition. Below the PCoA plot is a schematic representation based on sample composition, where arrows demonstrate different directions for sample localization according to the presence and concentration of *Brettanomyces* spp., *Saccharomyces* spp., *Penicillium* spp., *Issatchenkia* spp., and *Wickerhamomyces* spp. (**b**) The tree shows clustering of the samples by similarity of their taxonomic composition, calculated by the Bray-Curtis dissimilarity at ASV level, using Ward's method.

3.2.2. Phylogenetic analysis

The construction of a phylogenetic tree using the fungal species detected in this study reveals different subspecies of both *B. bruxellensis* and *S. cerevisiae* (Figure 7a). Their excellent growth and adaptability in beer wort favors the presence of these yeasts, as well as the presence of other *Brettanomyces* spp. such as *B. anomalus, B. naardenensis,* and *B. custersianus,* all of which were identified in the spontaneous and non-spontaneous fermentation beers. Also, multiple species from the *Pichia* genus were detected, such as *P. fermentans* and *P. membranifaciens,* with emphasis on four subspecies of *I. orientalis* that were identified in large proportion in some samples. We observed that different ASVs referring to the fungus *Penicillium* spp. were present in these samples, as well as more complex fungi such as *Perenniporia japonica* and *Fomitopsis palustres* (ASVs identified only in samples 3 and 18, respectively), whose presence is generally related to wood colonization.

Fungi from the Ascomycota phylum, which contains yeasts, accounted for 88.46% of the ASVs found, and to a lesser extent (11.54%) from the Basidiomycota phylum (Figure 7b). The Pichiaceae family, which contains yeasts of the genera *Brettanomyces, Pichia, Kregervanrija,* and *Issatchenkia,* displayed the highest proportion among the families classified (34.62%) in this study. The *Saccharomyces, Torulaspora, Lachancea,* and *Kluyveromyces* genera belong to the Saccharomycetaceae family, a taxonomic classification that com-

prises genera of great importance in mixed fermentation beer production, and such microbes were responsible for 19.23% of the fungi ASVs classified. Other families were responsible for the classification of only one genus, representing 3.85% each of the total yeasts found.

(a)

(**b**)

Figure 7: **Phylogenetic analysis and abundance of fungal taxa**. (**a**) Phylogenetic tree constructed based on the ASVs of 26 fungal species found in the mixed fermentation beer samples. (**b**) The image shows the prevalence of taxonomic classifications according to genera and species found in the fungal microbiomes. Higher taxonomic classifications are shown in circles near the image center, while lower classifications are shown toward the outer edge.

4. Discussion

4.1. Exploring the microbiomes of mixed fermentation beers

Elucidating the microbiomes of mixed fermentation beers is important to understand the participation of microorganisms in this fermentation niche. The production of sour beers from spontaneous fermentations is associated with inconsistencies in product quality, unpredictability in fermentation results, and extra time to consumption [10], as it is often not known which microorganisms will act in the beer wort exposed to the thousands of bacteria and fungi present in the environment. Thus, identifying and characterizing the metagenomes present in mixed fermentation beers and mixed cultures samples allows one to predict which microorganisms typically participate in these fermentations and formulate new bacterial and yeast blends to mimic this fermentation process in a controlled and reproducible way.

Preliminary studies performed by our research group concerning the metagenomes of commercial beers made by mixed fermentation and barrels used in their maturation corroborate the data presented here, where *L. acetotolerans*, *L. brevis*, *L. buchneri*, *B. bruxellensis*, and *S. cerevisiae* were commonly observed in samples (data not shown). Though different microorganisms have been detected at lower concentrations, the metagenome data in this study are in accordance with observations by other researchers, such as Bokulich et al. (2012), Bokulich et al. (2015), De Roos et al. (2019) and Tyakht et al. (2021). We observed in our work that dozens of bacterial species can be found through NGS using the V3/V4 region of the 16S rRNA, revealing that beers produced spontane-

ously and non-spontaneously represent an interesting source for the identification of new microorganisms that cooperate or compete among themselves in the consumption of substrates present in beer wort [13]. Their presence is important in mixed fermentation beers, as they are able to enhance the production of specific flavors and acidify the wort through lactic acid production, typical characteristics in these beer styles [2].

Fungal ASV detection by sequencing the ITS2 region revealed microbiomes based on two main genera: *Brettanomyces* spp. and *Saccharomyces* spp., which were found in the vast majority of samples corresponding to > 90% of the ASVs identified per sample. However, there is still a little-known world of unconventional yeasts to be discovered and characterized, which comprises less common genera such as *Issatchenkia* and *Wickerhamomyces*. The fungal microbiomes presented a lower variability in the number of species when compared to bacteria, which may be related to avidity in glucose consumption by commercial strains (reducing the concentration of carbohydrates with easy assimilation in the medium) and the competition among yeasts, which can present killer characteristics, producing secondary metabolites that aim to stop the multiplication of cross-feed competitors [14,15]. Even so, 12 different yeast genera were detected participating in these fermentations.

4.2. Lactobacillus spp. and Pediococcus spp. are often identified in mixed fermentation beers

In recent years, the application of LAB has been explored in sour beer production, with emphasis on bacteria in the *Lactobacillus* genus as not just contaminants in beer worts but as interesting tools for acidification and production of new flavors [16]. Although the vast majority of these bacteria are not tolerant to hop alpha-acid concentrations up to 20 International Bitterness Units (IBUs), some species and subspecies adapt to the adverse conditions that exist in beers, becoming tolerant to this and others selective pressures, such as hydrostatic pressure, alcohol, and low pH [17]. In our study, we observed nine *Lactobacillus* spp., including *L. acetotolerans* (with eight identified subspecies), *L. backii*, and *L. plantarum*, a genus present in 18 of the 20 samples analyzed. Several authors highlight the presence of *Lactobacillus* spp. in the microbiomes of mixed fermentation beers, such as Tyakht et al. (2021) and Spitaels et al. (2014), who detected these bacteria in American Coolship Ales, Wild Ales, Sour Ales, and Belgian Lambic Beers.

P. damnosus was the only identified species from the *Pediococcus* genus, but 18 subspecies were detected in this study. Similar to *Lactobacilllus* spp., *Pediococcus* spp. are commonly considered as beer and manufacturing environment contaminants, a characteristic that encourages research into the identification of resistance factors, mainly acquired through specific plasmids and genes [13,18]. In Lambic-style beers (produced by spontaneous fermentation), *P. damnosus* is one of the main isolated species and can be easily identified as part of the microbiota of the interior surface of wooden barrels [1] or in the air and on other brewery equipment surfaces [19]. In our study, ASVs corresponding to these bacteria were found in seven samples, representing up to 70% of the bacterial microbiome, as in sample 2.

Biological acidification during the brewing process by the action of *Lactobacillus* spp. or *Pediococcus* spp. has several benefits: flavor stability, greater zinc bioavailability, fast final attenuation, lower wort viscosity, and smoother hop bitterness, among others [2]. In pre-fermentation tests using these bacteria, impacts are observed on volatile compounds and organic acid production, which result in significant differences in the sensorial characteristics of sour beers [16]. These bacteria can also use carbohydrates that are not metabolized by conventional yeasts, such as maltotriose, maltotetraose, and cellobiose, resulting in beer overattenuation [10,20]. Although *Pediococcus* spp. are historically reported as responsible for high levels of diacetyl production and causing viscous "sick beers" (through exopolysaccharide secretion), these bacteria have been intentionally combined with *Brettanomyces* spp. to generate a deeper acidity and mouthfeel in various mixed fermentation beers [20].

4.3. Bacteria of the Enterobacteriaceae family and their presence in spontaneous and
non-spontaneous fermentations

Enterobacteriaceae bacteria are well-known contaminants in spontaneous fermen-
tation beers, participating in the initial stages of fermentation (also known as the
Enterobacteriaceae phase) that starts on day one and can last approximately 1 month [1].
In our work, ASVs corresponding to different bacteria in this family were identified, such
as *Rahnella* spp., *Klebsiella* spp., *Enterobacter* spp., *Providencia* spp., *Escherichia* spp.,
Erwiniae spp., and *Pantoea* spp. In samples 6, 7, and 20, we found that almost the entire
bacterial microbiome corresponded to Enterobacteriaceae, with emphasis on the species
R. aquatilis, K. oxytoca, E. asburiae, and *E. billingiae,* which had concentrations >20% of the
ASVs found in these samples. Using methods for the isolation of microorganisms in
specific culture media, Spitaels et al. (2014) could verify that after 2 months of matura-
tion, only *P. damnosus* was isolated in beers from which Enterobacteriaceae were found
during the initial fermentation period. Thus, it is noteworthy that although we detected
ASVs corresponding to these bacteria in the samples in our study, this does not imply
that these microbes were viable at the time of sample collection.

Different studies have successfully isolated and identified Enterobacteriaceae in
spontaneous fermentation beers, whether American Coolship Ales, Belgian Lambics, or
Wild Ales [4,5,9]. In such cases, these bacteria were highly abundant (>8% of ASVs) in six
beer samples. Although they are responsible for off-flavor production and can be harm-
ful to health through biogenic amine production [1,5], these bacteria are also linked to the
production of specific flavors related to young Lambics, such as 2,3-butanediol, ethyl ac-
etate, higher alcohols, acetic acid, lactic acid, succinic acid, and fatty acids [6,20].

4.4. Other bacteria

The metagenomes of our mixed culture samples were not restricted to bacteria from
the Enterobacteriaceae and Lactobacillaceae families. Several different species were often
present, mainly AAB from the genera *Gluconobacter* and *Acetobacter.* Indeed, *G. oxydans*
corresponded to 64.23% of the ASVs found in sample 3, and different *Acetobacter* spp.
were present in concentrations >20% in samples 2, 4, and 5. Their presence is likely due to
the fact that they are able to tolerate hop alpha-acids and ethanol concentrations up to
10%, also being responsible for ethanol oxidation with the formation of organic acids,
mainly acetic acid (one of the flavors commonly associated with spontaneous fermented
beers) [21].

Similar to our findings, Tyakht et al. (2021) detected the presence of the
Leuconostocacaceae and Acetobactereaceae families at abundant levels (>5%) in the
microbiomes of two wild beers. Not only are LAB important in the flavor bouquet con-
struction of sour beers, but other less conventional bacteria can also be highlighted, such
as *Acetobacter* spp., which are related to the production of beers with high contents of
5-methyl-furfural, flavonoids, and 2- and 3-methyl butanol [5]. When associated with the
presence of lactic acid, acetic acid addition by these bacteria (at adequate amounts) can
increase the sensory complexity of the beer, resulting in the construction of "layered
flavors" so important in traditional Belgian sour beers [20].

Bacteria of the genus *Pseudomonas* were also detected. Even though a large variabil-
ity of ASVs for *Pseudomonas* spp. was found only in sample 20, *P. fluorescens* was identi-
fied in 20% of the studied samples (n=4), with concentrations ranging from 0.3-8% in
samples in which it was identified (samples 1, 3, 12, and 19). This genus was also identi-
fied by Rodhouse (2017), mainly in malt samples, with a decrease in its detection during
the brewing process (mashing, boiling, and bottling). Because spontaneously fermented
beers are usually exposed to the environment, it is possible that these bacteria are trans-
ported from the raw brewing ingredients (mainly malt) to the wort during the open-air
fermentation process [23].

Aside from *Pseudomonas* spp., other Gram-negative bacteria found in significant concentrations were *Ralstonia* spp., representing 17% of bacterial ASVs found in sample 19 and 8% in sample 3. Both genera were also detected in small concentrations (<10^3 cell/mL) by other researchers, such as Takahashi et al. (2015) who observed the presence of these bacteria in the early and intermediate stages of beer-like beverage fermentation, and Bokulich et al. (2012), who detected them in American Coolship Ales. Other bacteria such as *Acinetobacter* spp., *Stenotrophomonas* spp., *Sphingomonas* spp., and *Staphylococcus* spp. were also detected in small amounts in the metagenomes of mixed fermentation beers by Bossaert et al. (2021). However, De Roos et al. (2019) detected a high relative abundance of the same bacteria on the surfaces of wooden barrels used in beer fermentation.

4.5. Fungi besides Brettanomyces spp. and Saccharomyces spp.

In addition to the traditional and well-known *Brettanomyces* spp. and *Saccharomyces* spp., other yeasts were present in the microbiomes of our mixed culture samples. Genera such as *Debaryomyces*, *Lachancea*, *Pichia*, and mainly *Issatchenkia* and *Wickerhamomyces* were identified as part of the samples' fungal microbiomes. *I. orientalis*, a yeast known for its industrial application in bioethanol production and participation in the construction of the aroma bouquet in wines [25], has been used in co-fermentations with *S. cerevisiae* for the production of beers with higher levels of fruity esters [26]. In our study, we identified this yeast at high concentrations (>45% of the ASVs found) in the fungal microbiomes of samples 3 and 14, demonstrating its adaptability to beer wort and its participation in the fermentation process.

Wickerhamomyces anomalus use has already been studied in a controlled way in the production of beers, classifying this yeast as a potential organism for the primary souring technique with contribution to fruity profiles [27,28]. *W. anomalus* was detected in our study, comprising almost the entire fungal microbiome (>98%) in samples 6 and 7, as well as being relatively abundant in sample 3 (9% of the ASVs). Although Tyakht et al. (2021) did not detect the presence of *W. anomalus* in the beer samples analyzed, Spitaels et al. (2014) confirmed its participation in Lambic beer fermentation, identifying relevant levels (>20%) in the microbiomes of beers after 24 months and detecting its presence in the air of the brewery environment and on the external surface of casks.

Varied *Pichia* spp. such as *P. manshurica* and *P. fermentans*, were also found in the analyzed samples, generally in concentrations <2%. However, *P. membranifaciens* was the only species of this genus that could be identified above this rate in the microbiome of a sample, representing 7% of the ASVs found in sample 2. Although in our study the concentration of *Pichia* spp. was considered low, other studies such as Bossaert et al. (2021) identified *P. membranifaciens* comprising up to 98% of the OTUs (Operational Taxonomic Units) found in beers at specific maturation periods, also revealing that the beer evolution leads to changes in the microbiome. *Lachancea* spp. are also reported to participate in mixed fermentation beers, responsible mainly for pH decrease in sour beers through lactic acid production [27,29]. *L. fermentati/dasiensis* was detected in our work in sample 3, with a relatively abundant concentration of 12% of the fungal microbiome.

The beer microbiome in our study revealed that among the most abundant families of fungi, Aspergillaceae represented around 7% of the identified ASVs. Cason et al. (2020) observed a similar relative abundance (6.7% of OTUs) in Sesotho beer samples, a traditional South African style produced by spontaneous fermentation. Fungi of the *Penicilllium* genus, which is part of this family, are related to gushing, and their presence has been detected inside beer and wine barrels, mainly participating in biofilm formation on porous surfaces [11,31]. These fungi are capable of producing enzymes responsible for the degradation of lignocellulose and beta-glycans, contributing to changes in beer color and aroma [32]. Other researchers such as Bossaert et al. (2021) and De Roos et al. (2019) have also identified *Penicillium* spp. in their samples. However, we found only one sam-

ple (sample 14) with >80% of fungal ASVs related to *Penicillium carneum/roqueforti*. ASVs referring to fungi with advanced and more complex structures (*e.g.*, *P. japonica*) were also detected in samples, at levels between 0.05% and 0.5%, indicating little relevance in the microbiomes of the analyzed beers.

4.6. The importance of traditional yeasts in mixed fermentation beers

Brettanomyces (also known as *Dekkera*), perhaps the second most important genus of yeast in mixed fermentation beers, can be found in large proportions in spontaneous and non-spontaneous fermentation beers, being associated with the production of phenolic and esterified volatile compounds [10]. The ability of *Brettanomyces* ssp. to metabolize complex sugars, produce acetic acid, generate a characteristic aromatic profile, and cause "super-attenuation" in beer wort has aroused interest in their use [28,33], which has led to several production laboratories around the world distributing these yeasts commercially.

Here, *B. bruxellensis* was the species with highest proportions found in the ASVs of this genus, totaling >90% of the fungal microbiome in some samples, such as 12, 17, and 19. Sobel et al. (2017) also highlighted the large presence of *B. bruxellensis* in traditional beers from countries such as Belgium, Italy, and Switzerland, detecting it in the metagenomes of 36 samples (n=39). Not only did we find *B. bruxellensis* in large proportions in our samples, but also *B. anomalus* (15% of ASVs in sample 1) and *B. custersianus*, which comprised >85% of ASVs in sample 13. These species present differences in sugar metabolism, aggregation, and flavor production, leading to interest in their presence during the fermentation process [35]. It is noteworthy that the ASVs corresponding to the *Brettanomyces* spp. and *Saccharomyces* spp. found in samples 2, 11, and 18 are possibly not related to wild strains but rather to the use of commercial blends inoculated at the beginning of fermentation.

The identification of the yeasts in the metagenomes of our samples revealed only one species in the *Saccharomyces* genus, *S. cerevisiae*, which together with *S. pasteurianus* are the brewer's yeasts most used for beer fermentation and are widely commercially available through propagation laboratories [36]. Wild strains of *S. cerevisiae* can offer interesting characteristics, such as extracellular production of different secondary metabolites (related to the production of new aromas) and tolerance to different stress conditions (*e.g.*, salinity, temperature, and high levels of ethanol) [15]. These strains have mainly been isolated and characterized from spontaneous fermentations of traditional fermented beverages, such as Kveik strains, isolated from traditional beers named Norwegian Kveik Beer [37]. In seven samples, in which there was no commercial *S. cerevisiae* inoculum, ASVs corresponding to *S. cerevisiae* were detected, which may be related to the presence of wild strains acting during fermentation. However, because the detection of this yeast may be related to cross-contamination of utensils and fermenters by the presence of commercial *S. cerevisiae* on such equipment [12], further studies must be conducted to confirm these ASVs as originating from wild strains.

Among the eight strains of *S. cerevisiae* detected in the metagenomes here, only one was identified as a hybrid of this yeast with another species (*S. cerevisiae* × *S. eubayanus* × *S. uvarum*), found at low concentration (<2% of the ASVs) in just one sample (#20), a beer produced through spontaneous fermentation. Yeasts with hybrid genotypes are generally reported in beer and wine, where their presence is caused by interspecific hybridization during diversification and adaptation of yeasts to the industrial niche [38].

5. Conclusions

The microbiome analysis of 20 samples of mixed fermentation beers with different origins allowed us to deepen our knowledge of the metagenomes of mixed cultures used for beer production, identifying the distribution and concentration of bacteria and fungi in these samples. Exploring new microorganisms and their strains that adapt to and

ferment beer wort is an important key factor in the rational development of new blends for brewers, so that beers and their flavors become reliably reproducible. Based on these data, we conclude this is a vast field that needs to be further explored, with potential for industrial applications and in the development of basic and applied science. Future work connecting the bacterial compositions of mixed cultures used for fermentation to metadata on the base wort, kinetics of fermentation, and organoleptic compounds produced in various sour beers will enable brewers to construct bespoke mixed cultures the generate desired sensorial results.

Supplementary Materials: The following are available online at www.mdpi.com/xxx/s1, Supplementary Information S1: Raw reads, ASVs sequences, microorganisms' identification and prevalence in each sample – Microsoft Excel File.

Funding: This research was funded by donors to the Experiment.com project "Mixed culture metagenomics of the microbes making sour beer" (DOI 10.18258/13495).

Data Availability Statement: The data presented in this study are available in Supplementary Information.

Conflicts of Interest: The authors declare no conflict of interest.

Appendix A

Figure A1: **Enterobacteriaceae family presence in the bacterial composition of the analyzed samples**. Bacteria from the Enterobacteriaceae family are commonly detected in the first fermentation phase of mixed fermentation beers produced by spontaneous fermentation (De Roos and De Vuyst, 2019), and in our study could be identified in relevant proportions in samples 1, 6, 7, 14, 19, and 20 (>8% of the ASVs detected in each sample).

References

1.   De Roos, J.; De Vuyst, L. Microbial acidification, alcoholization, and aroma production during spontaneous lambic beer production. *J. Sci. Food Agric.* **2019**, *99*, 25–38, doi:10.1002/jsfa.9291.

2.   Vriesekoop, F.; Krahl, M.; Hucker, B.; Menz, G. 125th Anniversary review: Bacteria in brewing: The good, the bad and the ugly. *J. Inst. Brew.* **2012**, *118*, 335–345, doi:10.1002/jib.49.

3.   Faria-Oliveira, F.; Diniz, R.H.S.; Godoy-Santos, F.; Piló, F.B.; Mezadri, H.; Castro, I.M.; Brandão, R.L. The Role of Yeast and

Lactic Acid Bacteria in the Production of Fermented Beverages in South America. In *Food Production and Industry*; InTech,
2015; Vol. i, p. 13.

4.   Spitaels, F.; Wieme, A.D.; Janssens, M.; Aerts, M.; Daniel, H.M.; Van Landschoot, A.; De Vuyst, L.; Vandamme, P. The
microbial diversity of traditional spontaneously fermented lambic beer. *PLoS One* **2014**, *9*, doi:10.1371/journal.pone.0095384.

5.   Bossaert, S.; Winne, V.; Van Opstaele, F.; Buyse, J.; Verreth, C.; Herrera-Malaver, B.; Van Geel, M.; Verstrepen, K.J.;
Crauwels, S.; De Rouck, G.; et al. Description of the temporal dynamics in microbial community composition and beer
chemistry in sour beer production via barrel ageing of finished beers. *Int. J. Food Microbiol.* **2021**, *339*,
doi:10.1016/j.ijfoodmicro.2020.109030.

6.   Bokulich, N.A.; Bamforth, C.W.; Mills, D.A. Brewhouse-resident microbiota are responsible for multi-stage fermentation of
American coolship ale. *PLoS One* **2012**, *7*, doi:10.1371/journal.pone.0035507.

7.   Callahan, B.J.; McMurdie, P.J.; Rosen, M.J.; Han, A.W.; Johnson, A.J.A.; Holmes, S.P. DADA2: High-resolution sample
inference from Illumina amplicon data. *Nat. Methods* **2016**, *13*, 581–583, doi:10.1038/nmeth.3869.

8.   Schoch, C.L.; Ciufo, S.; Domrachev, M.; Hotton, C.L.; Kannan, S.; Khovanskaya, R.; Leipe, D.; McVeigh, R.; O'Neill, K.;
Robbertse, B.; et al. NCBI Taxonomy: A comprehensive update on curation, resources and tools. *Database* **2020**, *2020*, 1–21,
doi:10.1093/database/baaa062.

9.   Tyakht, A.; Kopeliovich, A.; Klimenko, N.; Efimova, D.; Dovidchenko, N.; Odintsova, V.; Kleimenov, M.; Toshchakov, S.;
Popova, A.; Khomyakova, M.; et al. Characteristics of bacterial and yeast microbiomes in spontaneous and
mixed-fermentation beer and cider. *Food Microbiol.* **2021**, *94*, 103658, doi:10.1016/j.fm.2020.103658.

10.   Dysvik, A.; La Rosa, S.L.; De Rouck, G.; Rukke, E.-O.; Westereng, B.; Wicklund, T. Microbial Dynamics in Traditional and
Modern Sour Beer Production. *Appl. Environ. Microbiol.* **2020**, *86*, 1–14, doi:10.1128/AEM.00566-20.

11.   De Roos, J.; Van der Veken, D.; De Vuyst, L. The interior surfaces of wooden barrels are an additional microbial inoculation
source for lambic beer production. *Appl. Environ. Microbiol.* **2019**, *85*, 1–18, doi:10.1128/AEM.02226-18.

12.   Bokulich, N.A.; Bergsveinson, J.; Ziola, B.; Mills, D.A. Mapping microbial ecosystems and spoilage-gene flow in breweries
highlights patterns of contamination and resistance. *Elife* **2015**, *4*, e04634, doi:10.7554/eLife.04634.

13.   Kajala, I.; Bergsveinson, J.; Friesen, V.; Redekop, A.; Juvonen, R.; Storgårds, E.; Ziola, B. *Lactobacillus backii* and *Pediococcus*
damnosus isolated from 170-year-old beer recovered from a shipwreck lack the metabolic activities required to grow in
modern lager beer. *FEMS Microbiol. Ecol.* **2018**, *94*, 1–10, doi:10.1093/femsec/fix152.

14.   Rodhouse, L.; Carbonero, F. Overview of craft brewing specificities and potentially associated microbiota. *Crit. Rev. Food*
Sci. Nutr. **2019**, *59*, 462–473, doi:10.1080/10408398.2017.1378616.

15.   Molinet, J.; Cubillos, F.A. Wild Yeast for the Future: Exploring the Use of Wild Strains for Wine and Beer Fermentation.
Front. Genet. **2020**, *11*, 1–8, doi:10.3389/fgene.2020.589350.

16.   Dysvik, A.; Liland, K.H.; Myhrer, K.S.; Westereng, B.; Rukke, E.O.; de Rouck, G.; Wicklund, T. Pre-fermentation with lactic
acid bacteria in sour beer production. *J. Inst. Brew.* **2019**, *125*, 342–356, doi:10.1002/jib.569.

17.   Sakamoto, K.; Konings, W.N. Beer spoilage bacteria and hop resistance. *Int. J. Food Microbiol.* **2003**, *89*, 105–124,
doi:10.1016/S0168-1605(03)00153-3.

18.   Garcia-Garcia, J.H.; Damas-Buenrostro, L.C.; Cabada-Amaya, J.C.; Elias-Santos, M.; Pereyra-Alférez, B. *Pediococcus*
damnosus strains isolated from a brewery environment carry the horA gene. *J. Inst. Brew.* **2017**, *123*, 77–80,
doi:10.1002/jib.397.

19.   Garcia-Garcia, J.H.; Galán-Wong, L.J.; Pereyra-Alférez, B.; Damas-Buenrostro, L.C.; Pérez, E.; Carlos Cabada, J. Distribution
of *Lactobacillus* and *Pediococcus* in a brewery environment. *J. Am. Soc. Brew. Chem.* **2017**, *75*, 312–317,
doi:10.1094/ASBCJ-2017-4294-01.

20.   Bossaert, S.; Crauwels, S.; Lievens, B.; De Rouck, G. The power of sour - A review: Old traditions, new opportunities.

BrewingScience **2019**, *72*, 78–88, doi:10.23763/BrSc19-10bossaert.

21. Ashtavinayak, P.; Elizabeth, H.A. Review: Gram Negative Bacteria in Brewing. *Adv. Microbiol.* **2016**, *06*, 195–209, doi:10.4236/aim.2016.63020.

22. Rodhouse, L. Phylogenetic Diversity of the Bacterial Communities in Craft Beer. **2017**.

23. Justé, A.; Malfliet, S.; Lenaerts, M.; De Cooman, L.; Aerts, G.; Willems, K.A.; Lievens, B. Microflora during malting of barley: Overview and impact on malt quality. *BrewingScience* **2011**, *64*, 22–31.

24. Takahashi, M.; Kita, Y.; Kusaka, K.; Mizuno, A.; Goto-Yamamoto, N. Evaluation of microbial diversity in the pilot-scale beer brewing process by culture-dependent and culture-independent method. *J. Appl. Microbiol.* **2015**, *118*, 454–469, doi:10.1111/jam.12712.

25. Zhao, Y.; Sun, Q.; Zhu, S.; Du, F.; Mao, R.; Liu, L.; Tian, B.; Zhu, Y. Biodiversity of non-*Saccharomyces* yeasts associated with spontaneous fermentation of Cabernet Sauvignon wines from Shangri-La wine region, China. *Sci. Rep.* **2021**, *11*, 1–10, doi:10.1038/s41598-021-83216-x.

26. van Rijswijck, I.M.H.; Wolkers - Rooijackers, J.C.M.; Abee, T.; Smid, E.J. Performance of non-conventional yeasts in co-culture with brewers' yeast for steering ethanol and aroma production. *Microb. Biotechnol.* **2017**, *10*, 1591–1602, doi:10.1111/1751-7915.12717.

27. Osburn, K.; Amaral, J.; Metcalf, S.R.; Nickens, D.M.; Rogers, C.M.; Sausen, C.; Caputo, R.; Miller, J.; Li, H.; Tennessen, J.M.; et al. Primary souring: A novel bacteria-free method for sour beer production. *Food Microbiol.* **2018**, *70*, 76–84, doi:10.1016/j.fm.2017.09.007.

28. Capece, A.; Romaniello, R.; Siesto, G.; Romano, P. Conventional and Non-Conventional Yeasts in Beer Production. *Fermentation* **2018**, *4*, 38, doi:10.3390/fermentation4020038.

29. Zdaniewicz, M.; Satora, P.; Pater, A.; Bogacz, S. Low lactic acid-producing strain of *Lachancea thermotolerans* as a new starter for beer production. *Biomolecules* **2020**, *10*, doi:10.3390/biom10020256.

30. Cason, E.D.; Mahlomaholo, B.J.; Taole, M.M.; Abong, G.O.; Vermeulen, J.G.; de Smidt, O.; Vermeulen, M.; Steyn, L.; Valverde, A.; Viljoen, B. Bacterial and Fungal Dynamics During the Fermentation Process of Sesotho, a Traditional Beer of Southern Africa. *Front. Microbiol.* **2020**, *11*, 1–14, doi:10.3389/fmicb.2020.01451.

31. Vogt, E.I.; Kupfer, V.M.; Vogel, R.F.; Niessen, L. Evidence of gushing induction by *Penicillium oxalicum* proteins. *J. Appl. Microbiol.* **2017**, *122*, 708–718, doi:10.1111/jam.13366.

32. Nigam, D.; Asthana, M.; Kumar, A. *Penicillium□: A fungus in the wine and beer industries*; Elsevier B.V., 2017; ISBN 9780444635013.

33. Serra Colomer, M.; Funch, B.; Forster, J. The raise of *Brettanomyces* yeast species for beer production. *Curr. Opin. Biotechnol.* **2019**, *56*, 30–35, doi:10.1016/j.copbio.2018.07.009.

34. Sobel, J.; Henry, L.; Rotman, N.; Rando, G. BeerDeCoded: the open beer metagenome project. *F1000Research* **2017**, *6*, 1676, doi:10.12688/f1000research.12564.1.

35. Colomer, M.S.; Chailyan, A.; Fennessy, R.T.; Olsson, K.F.; Johnsen, L.; Solodovnikova, N.; Forster, J. Assessing Population Diversity of *Brettanomyces* Yeast Species and Identification of Strains for Brewing Applications. *Front. Microbiol.* **2020**, *11*, 1–21, doi:10.3389/fmicb.2020.00637.

36. Stewart, G. *Saccharomyces* species in the Production of Beer. *Beverages* **2016**, *2*, 34, doi:10.3390/beverages2040034.

37. Lengeler, K.B.; Stovicek, V.; Fennessy, R.T.; Katz, M.; Förster, J. Never Change a Brewing Yeast? Why Not, There Are Plenty to Choose From. *Front. Genet.* **2020**, *11*, doi:10.3389/fgene.2020.582789.

38. Gallone, B.; Steensels, J.; Mertens, S.; Dzialo, M.C.; Gordon, J.L.; Wauters, R.; Theßeling, F.A.; Bellinazzo, F.; Saels, V.; Herrera-Malaver, B.; et al. Interspecific hybridization facilitates niche adaptation in beer yeast. *Nat. Ecol. Evol.* **2019**, *3*, 1562–1575, doi:10.1038/s41559-019-0997-9.

909