VAL1 as an assembly platform co-ordinating co-transcriptional repression and chromatin regulation at Arabidopsis *FLC*

Pawel Mikulski [1,4], Philip Wolff [1], Tiancong Lu [1,2], Danling Zhu [1,3] & Caroline Dean [1*]

[1] Cell and Developmental Biology, John Innes Centre, Norwich, UK

[2,3,4] Present addresses: [2]State Key Laboratory of Plant Genomics and National Center for Plant Gene Research, Institute of Genetics and Developmental Biology, Chinese Academy of Sciences, Beijing, China; [3]SUSTech-PKU Institute of Plant and Food Science, Department of Biology, Southern University of Science and Technology, Shenzhen, China; [4]Department of Biochemistry, University of Oxford, Oxford, UK

Abstract

Polycomb (PcG) silencing is crucial for development across eukaryotes, but how PcG targets are regulated is still incompletely understood. The slow timescale of cold-induced PcG silencing at *Arabidopsis thaliana FLOWERING LOCUS C* (*FLC*) makes it an excellent system to dissect this mechanism. Binding of the DNA binding protein VAL1 to an *FLC* intronic RY motif within the PcG nucleation region is an early step in the silencing process. VAL1 interacts with APOPTOSIS AND SPLICING ASSOCIATED PROTEIN (ASAP) complex and POLYCOMB REPRESSIVE COMPLEX 1 (PRC1). Here, we show that ASAP and PRC1 function as co-repressors that quantitatively regulate *FLC* transcription. Upon the shift to cold PRC1-mediated H2Aub accumulates only at the nucleation region, is transiently maintained after transfer back to warm, but unlike the PRC2-delivered H3K27me3 does not spread across the locus. H2K27me3 thus provides long-term epigenetic silencing, whereas H2Aub is a transient repression signal. Overall, our work highlights how a DNA sequence-specific binding protein can act as an assembly platform co-ordinating the co-transcriptional repression and chromatin regulation necessary for Polycomb silencing.

Introduction

From elegant genetic and molecular analysis in Drosophila the Polycomb group (PcG) proteins have been shown to play a central role in developmental gene regulation[1]. They function in distinct complexes to maintain epigenetic silencing of genomic targets[2]. The most well-studied complexes are POLYCOMB REPRESSIVE COMPLEX 1 (PRC1), which monoubiquitinates the core of histone H2A (H2Aub), and POLYCOMB REPRESSIVE COMPLEX 2 (PRC2), which methylates histone H3 tails at residue 27 (H3K27me). Initially thought to be sequentially recruited (PRC2, then PRC1) many studies have now demonstrated a much more complex scenario with variant and canonical PRC1 [3,4], interdependent recruitment of PRC1 and PRC2 and co-operative interactions mediating association to Polycomb Response Elements (PREs)[5–8]. There is also a lack of clarity about target site specificity as Polycomb complexes have been proposed to directly 'sample' transcriptionally permissive chromatin sites, rather than be recruited by developmental regulators[9]. Furthermore, the relationship between transcription and chromatin silencing appears to be able to switch between instructive and responsive[10]. Further dissection of the molecular events underpinning Polycomb silencing is therefore necessary to define the core mechanistic principles.

Arabidopsis *FLOWERING LOCUS C (FLC)* is a PcG target central to the process of vernalization ensuring flowering in favourable conditions[11,12]. *FLC* encodes a floral repressor whose expression is high in winter annual Arabidopsis accessions through the activity of FRIGIDA[13,14]. Short cold exposure in autumn/winter results in a relatively rapid transcriptional repression of *FLC*, in a process

involving cold-induced *FLC* antisense transcripts called *COOLAIR*[15]. Over the longer timescale of
winter, *FLC* is epigenetically silenced through a PRC2-dependent epigenetic switch[16]. This switch
occurs at individual *FLC* alleles with a low probability[17], leading to progressive silencing over the
whole plant. A cold-induced step in this switch is the VIN3/VRN5-dependent nucleation of
H3K27me3 at an intragenic site that covers three nucleosomes between the transcription start site and
the beginning of intron 1[18–20]. When plants return to warm temperatures, H3K27me3 spreads to cover
the entire gene, a feature required for the long-term epigenetic silencing throughout the rest of
development[17]. The *FLC* system thus provides the opportunity to dissect the components required for
a PRC2 epigenetic switch.
58
Previously, we had shown that the transcriptional repressor VIVIPAROUS1/ABI3-LIKE (VAL1) was
a central factor in the switching mechanism, necessary for H3K27me3 nucleation at *FLC*[21]. VAL1
recognizes conserved RY motifs in *FLC* intron 1 with a one bp mutation in the first RY motif
sufficient to completely block the vernalization process. The *val1* mutation attenuated *FLC* cold-
induced transcriptional repression, but not the longer-term Polycomb memory. VAL1 was shown to
interact with components of the APOPTOSIS AND SPLICING ASSOCIATED PROTEIN (ASAP),
HDAC complex and PRC1, with components of ASAP interacting with the PRC2 accessory proteins
VIN3 and VRN5. This then provided a connection between the sequence at the nucleation region and
the machinery for long-term PcG silencing. However, the molecular role of the VAL1 interactors was
not known.
69
Here, we investigate how VAL1 and its interactors ASAP and PRC1 function in *FLC* regulation.
Overall, our results suggest that ASAP and PRC1 co-ordinate different steps in *FLC* regulation
determining whether and how plants respond to cold, with the sequence-specific transcription factor
VAL1 acting as the common recruitment platform. ASAP and PRC1 link co-transcriptional
processing with chromatin regulation with PRC1-deposited H2AUb accumulating at the nucleation
region during cold exposure. However, this is only maintained at the nucleation region, unlike the
PRC2-delivered H3K27me3 which spreads across the locus for long-term epigenetic silencing. Our
work reveals how VAL1 functions as an assembly platform to co-ordinate co-transcriptional
repression and chromatin regulation at Arabidopsis *FLC*.
79
Results
Mutations of the ASAP complex affect *FLC* expression in warm but not cold-treated plants
To analyse whether ASAP had a role in VAL1-dependent cold-induced FLC silencing we crossed
asap mutants into a genotype carrying an active *FRI* allele in order to confer a vernalization
requirement. The generated FRI *asap* genotypes were analysed for *FLC* expression over a
vernalization time course. Plants were harvested before any cold exposure (non-vernalized; NV),
immediately after 2, 4 or 6 weeks of cold treatment (2WT0, 4WT0, 6WT0) and after 6 weeks of cold
followed by a warm period for 7 days (6WT7). Interestingly, the major effect of the *asap* mutants
both *FRI* and *fri* genotypes was in warm (NV) conditions (Fig. 1a, b). The *asap* mutants did not
influence the rate of *FLC* silencing after different cold and post-cold treatments (Fig. 1a). To
understand the relationship of VAL1 and ASAP function we generated double mutant combinations in
a Col-0 background and analysed in NV conditions. We observed a strong increase in *FLC* expression
in warm-grown double mutants of *val1 asap* (Fig. 1b), consistent with VAL1 interacting with many
complexes to effect *FLC* silencing. We also found a stronger release of *FLC* silencing in
combinations of *asap* mutants (Fig. 1b). As expected with the general correlation of sense and
antisense transcription at *FLC*[22], *COOLAIR* was also upregulated in single *asap* mutants in warm
conditions (Fig. S1a). These data suggest that a major role of ASAP at *FLC* is in quantitative

transcriptional regulation that sets the level of *FLC* expression in the warm. Consistently, we found an interaction in yeast two hybrid between VAL1 and SIN3A-ASSOCIATED PROTEIN 18 (SAP18), (Fig. 1c,d, S1b), as similarly found in IP-MS with VAL1::VAL1-HA used as a bait[21]. This result is also matching the functional connection between SAP18 and VAL1 reported recently[23]. Given the physical interaction of ASAP components with the PRC2 accessory proteins VIN3 and VRN5, which mediate the cold-induced H3K27me3 nucleation[21], the lack of an *asap* mutant phenotype during cold and post-cold steps likely indicates functional redundancy with other processes during the cold phase.

SR45-dependent co-transcriptional repression at *FLC*

A role for ASAP outside of Polycomb nucleation at *FLC* led us to analyse a link between co-transcriptional and chromatin regulation, as previously suggested for ASAP at other targets[24]. We selected the *sr45* mutant as an ASAP representative mutant with strong *FLC* upregulation (Fig. 1b) and profiled the co-transcriptional and chromatin changes at *FLC*. *sr45* increased relative *FLC* transcript levels both in total (Fig, 1b) and chromatin-associated, nascent RNA (Fig. 1e) and *COOLAIR* increased co-ordinately (Fig. S1a). We did not observe changes in *FLC* splicing efficiency at introns 1-3 caused by loss of SR45 or other ASAP components (Fig. S2). However, *sr45* caused higher levels of RNA PolII phosphorylated on serine-2 and serine-5 on its C-terminal domain (CTD) (Fig. 2a), which are associated with active transcription and increased the RNA PolII peak around the *FLC* nucleation region. Furthermore, the balance of active and repressive chromatin modifications was changed in warm grown *sr45* plants, with increased H3K36me3 and decreased H3K27me3 at *FLC* (Fig. 2b), in agreement with the transcriptional upregulation in the mutant. ASAP-dependent co-transcriptional regulation thus correlated with a variety of chromatin features at the *FLC* locus.

VAL1 affects specific nucleosome dynamics

A role for ASAP on co-transcriptional processes implied that its interactor, VAL1, may also have a similar function, especially given its impact on the active/repressive chromatin modifications[21]. To test this, we analysed chromatin-association of *FLC* RNA in warm-grown *val1* seedlings as a readout for bona fide *FLC* transcription. Like the relative increase in total RNA[21], and similar to *sr45* (Fig. 1e), *val1* increased chromatin associated *FLC* RNA (Fig. 3a). Given the VAL1 binding site is downstream of the *FLC* promoter we asked if such transcriptional changes might be accompanied by changes in nucleosome dynamics[25]. We adapted the histone salt fractionation protocol that preferentially extracts nucleosomes recently disrupted by either transcription or remodelling[25] and analysed *FLC*. The fraction of low-salt extractable nucleosomes decreased at the +1 and +2 nucleosomes in cold-treated samples (Fig. S3), agreeing with the previous observation of cold-induced nucleosome stabilization[19]. Furthermore, we observed that *val1* increased the low-salt extractable fraction for nucleosomes +1 and +3, but not for nucleosome +2 (Fig. 3b, c). Nucleosomes +1 and +3 correspond to the edges of the H3K27me3-enriched nucleation region found in *FLC* after cold exposure[26]. These VAL1-dependent changes were confined to chromatin extracted from warm-grown *val1* seedings and were not found in cold-treated samples (Fig. S3). Interestingly, they were not connected to gross changes in chromatin accessibility as judged by FAIRE (formaldehyde-assisted isolation of regulatory elements, a technique using nuclease hypersensitivity to identify open and therefore potentially regulatory chromatin) (Fig. 3d). Overall, these data connect VAL1-induced co-transcriptional quantitative regulation to nucleosome dynamics at the edges of the *FLC* nucleation region, a region where RNA PolII occupancy is high.

H2Aub accumulates during cold at the *FLC* nucleation region but does not spread like H3K27me3 upon return to warm

To begin to investigate the functional significance of the VAL1-PRC1 interaction we undertook
analysis of H2AUb accumulation at *FLC* on the vernalization-sensitive genotype ColFRI after
different cold exposure. Reduced *FLC* expression correlated with increasing H2Aub levels as plants
were exposed to progressive cold (Fig. 4a). H2Aub accumulated at the *FLC* nucleation region with
dynamics similar to PRC2-induced H3K27me3 nucleation. To establish whether the H2Aub
correlated with *FLC* transcriptional activity we then analysed lines with or without FRIGIDA (FRI),
which is the main determinant distinguishing vernalization requirement or rapid-cycling habit. In the
rapid-cycling Col-0, H2Aub levels at *FLC* were much higher than in Col-FRI reflecting the ~ 20x
reduction in *FLC* transcription (Fig. 4b). These results suggest that H2Aub is inversely correlated with
transcription and PRC1 has a typical transcription repression function at *FLC*, rather than a context-
dependent role in transcriptional activation[3,4].

We then tested how H2Aub changes in warm conditions post-cold, when *FLC* chromatin transitions
into a stably repressed state. We measured H2Aub and H3K27me3 at *FLC* during the vernalization
timecourse[26]. As expected, H3K27me3 accumulated in the nucleation region during cold exposure
(6W+day 0) and spread across the whole gene body after transfer back to warm (6W+day 10/14/20)
(Fig. 4c). H2Aub accumulated in the nucleation region during cold, similar to the H3K27me3
dynamics, and continued to increase for 10 and 14 days after transfer back to the warm but did not
spread across the locus (Fig. 4c). Interestingly, 20 days after transfer back to the warm, the H2Aub
peak at the nucleation decreased back to levels seen at the end of 6W cold. These dynamics closely
match the occupancy of the VRN5 accessory protein at the nucleation region[17].

VAL1, PRC1 and NDX physically interact and are required for H2Aub accumulation at the
***FLC* nucleation region**
Genetic dissection of the role of H2AUb at *FLC* is complex as the multiple Arabidopsis PRC1
components are functionally redundant and single PRC1 mutants do not show a changed *FLC*
expression relative to wildtype[27,28] (Fig. 5a). Loss of all three BMI subunits (A, B, C) de-represses
FLC expression in warm-grown Col-0 plants up to 25 fold[29]. Higher order PRC1 mutant combinations
are seedling lethal or show severe developmental aberrations[27,30,31]. In addition, Arabidopsis PRC1 has
a range of partners central to its involvement in different developmental pathways[32]. However, to
pursue the consequence of the VAL1-PRC1 interaction we analysed *FLC* expression differences in
val1 bmi1B line. *val1 bmi1B* did not increase *FLC* expression above *val1* alone in any condition tested
(Fig. 5b, c) or further influence *FLC* H2Aub levels in warm-grown plants (Fig. 5d). In addition, *val1*
bmi1B did not show *FLC/COOLAIR* reactivation post-cold (Fig. S4). This was also true for the
combination *val1 ring1A* (Fig. S5). We did find an additive effect of the *val1 bmi1B* combination on
COOLAIR expression (Fig. S6), so for this aspect of *FLC* regulation other PRC1 components do not
completely cover loss of BMI1B function.

We undertook further IP-MS experiments with VAL1-HA as a bait and found a robust interaction
between VAL1 with other proteins (Table S1), including PRC1 members, as well as NODULIN
HOMEOBOX (NDX) (Fig. 6a). Interestingly, NDX has been previously implicated in single-stranded
DNA recognition and stabilization of the R-loop at *FLC*[33]. Using yeast-two-hybrid analysis we found
that NDX does not interact directly with VAL1 but instead interacts with RING1A and RING1B
(PRC1 components) (Fig. 6b, c, S7), as also found recently[34]. Genetic analysis between *val1*, *prc1*
mutants and *ndx1* revealed that NDX is required for the *FLC* upregulation caused by *val1*, and loss of
PRC1 does not affect the consequences of *NDX* deficiency (Fig. 5a); *COOLAIR* was upregulated
similarly to *FLC* in the mutants (Fig. S8). Furthermore, we also showed that NDX, VAL1 or VAL1-
recognized RY motifs are necessary for full H2Aub enrichment at *FLC*, as well as H3K27me3 (Figs.

6d, e, S9 and as published in[21]). NDX has been shown to associate with the 3' region of *FLC* and
stabilize R-loops[33,35], so it will be important to establish if these genetic interactions point to an
additional NDX/R-loop interaction in the *FLC* nucleation region or are the result of the gene loop[36].

H2Aub enrichment at *FLC* was unaffected in mutants of the core PRC2 component
VERNALIZATION 2, or the PRC2-accessory protein VERNALIZATION 5 in non-vernalized
conditions but was slightly reduced after 6 weeks cold exposure (Fig. S10). The lack of difference in
NV conditions is in agreement with H2Aub being generally independent of LHP1 and PRC2 function
in Arabidopsis, as previously reported[37]. However, the small effect after cold potentially connects
close functioning of VAL1, NDX and PRC1 in *FLC* co-transcriptional repression to PRC2 nucleation
and chromatin silencing.

Discussion
How transcriptional repression links to epigenetic silencing is far from understood. Cell type-specific
or developmentally induced transcriptional repressors bind different Polycomb complexes but direct
recruitment models are too simplistic[9,10]; linear genetic hierarchies have not been established and
functional output of PREs has been shown to be heavily dependent on the chromatin context[38]. Our
findings that VAL1 acts as an assembly platform co-ordinating ASAP and PRC1 activities that occur
at different stages in the *FLC* regulation helps explain some of this complexity. It also raises
interesting parallels to the non-linear recruitment activities of the PcG proteins at Drosophila
PREs[7,39,40], with ASAP co-transcriptional repressors and PRC1 components likely redundantly
establishing the chromatin environment required for robust PRC2 silencing (Fig. 7).

ASAP and PRC1 have been documented to have very different roles in gene regulation in different
organisms. ASAP has previously been shown to play a role as part of the spliceosome and exon
junction complexes in many organisms[24,41], whereas, PRC1 functionality is associated with H2AUb
and transcriptional silencing[42]. In plants, ASAP has been associated with regulation of innate
immunity[43] and abiotic responses[44] and considered to function through altering RNA metabolism or
alternative splicing[45,46]. However, we did not detect any effect of a *sr45* mutation on alterative
splicing or splicing efficiency of the *FLC* sense strand introns. Instead, we found changes linked to
transcriptional repression – RNA PolII post-translational modifications and classic histone
modifications associated with Polycomb target ON or OFF states. We thus propose that SR45, as an
ASAP component, influences a co-transcriptional step that may function to influence RNA PolII
complex composition and thus the chromatin modifications deposited during transcription. Since there
is physical linkage of VIN3 and VRN5[21] to ASAP components, the ASAP complex is likely to
function at *FLC* both during growth in the warm, as judged by mutant phenotypes, and redundantly
with other functions in *FLC* silencing during cold exposure. Changed co-transcriptional regulation
with RNA PolII both involved in histone modification deposition, but also sensitive to histone
modifications on the chromatin template, would seem a probable mechanism underlying the
instructive/responsive transcription-chromatin relationship.

The PRC1-mediated H2AUb modification has been associated with transcriptional regulation and
epigenetic stability in Drosophila and mammalian cells[42]. At *FLC*, the PRC1-H2AUb is associated
with transcriptional repression, either in low *FLC* expressing genotypes (Col-0) in warm conditions,
or in response to cold in the high *FLC* expressing Col-FRI genotype. In this respect it appears to
function at *FLC* more as a variant PRC1 rather than a canonical PRC1. The timescale of the VAL1-
PRC1-mediated H2Aub accumulation at the *FLC* nucleation region during cold exposure suggests an
intermediary role between the transcriptional silencing determined by ASAP-VAL1 and the long-term

PRC2 epigenetically stable spread state. However, the genetic redundancy makes PRC1 functionality
hard to tease apart so we were unable to dissect the interdependency of PRC1-delivered H2AUb and
PRC2-delivered H3K27me3 in the epigenetic switch at *FLC*. Therefore, how VAL1 interaction with
ASAP and PRC1 is co-ordinated spatially and temporally at the *FLC* nucleation region remains to be
fully elaborated.
245
VAL1 interacts with ASAP via SAP18 and PRC1 but has also been shown to interact with LHP1[32],
histone deacetylases (HDA9, HDA6)[47,48] and Mediator complex component (MED13)[47]. It will be
important in the future to understand how one protein can mediate all these interactions and whether
they are reinforced through multivalent interactions with other proteins at the nucleation region, like
the Drosophila PRE situation. The original intronic mutation in the RY motif caused relatively little
warm mis-regulation of *FLC*[21], just attenuation of the cold-induced transcriptional silencing without a
subsequent effect on stability of the silenced state. The stronger phenotype of the VAL1 interactors in
the warm argues that they also function independently of VAL1. The lack of phenotype in the cold
suggests that functionally redundant, cold-specific regulators are still to be identified. These are likely
to involve the antisense transcripts at *FLC* (*COOLAIR*) that contribute to transcriptional silencing
during cold exposure, which associate with the chromatin at the 5' end of *FLC,* and in which natural
variation has a functional consequence[49,50].
258
Overall, we interpret our data as showing that VAL1 functions as an assembly platform to co-ordinate
co-transcriptional repression and chromatin regulation at Arabidopsis *FLC*. We envisage that
transcription is downregulated by multiple repressor complexes (ASAP and PRC1) either linked to
RNA PolII and interacting with VAL1 at the nucleation region or assembled at the nucleation region.
These appear to function independently in the warm, with H2AUb playing a major role in the cold
associated with *FLC* silencing, alongside PRC2 deposition of H3K27me3. Given the parallels with the
Drosophila PcG mechanism the activities of these many protein complexes are likely to function
cooperatively to influence the different steps in the mechanism. Further elucidation will require
detailed analysis of the co-transcriptional silencing, chromatin modification and nucleosome
remodelling.
269
Material and methods
Plant material
All mutants and transgenic lines were either in Col-0 or FRI[sf2] background[21,50]. Unless otherwise
stated, experiments in non-vernalized conditions used Col-0 as a wildtype. The analyses in
vernalization timecourse used Col FRI[sf2] (Col FRI) as a wildtype to capture dynamics of cold-induced
FLC expression and chromatin changes. Mutant alleles were used as described previously: *bmi1B*
(*drip1-1*), *bmi1A* (*drip2-1*), *ring1A* (GK-293A04), *ring1B* (SALK_117958), *ndx1-4*, *flc* (*flc-2*), RY
motif (*vrn8*), *val1-2*, *vin3-4*, *vrn2-1*, *vrn5-8*, *sap18* (SALK_02363C), *sr45* (SALK_018273), *acinus*
(SALK_078554)[17,21,33,34,51]. Double mutants were generated by crosses between homozygous single
mutant lines and selected by PCR-based genotyping using primers listed in Table S2.
Growth conditions
Seeds were surface sterilized and sown on MS-GLU (MS without glucose) media plates and stratified
at 4°C in the dark for 2 days. For non-vernalized (NV) conditions, seedlings were grown for 14 days
in long-day conditions (16 h light, 8 h darkness with constant 20°C). For vernalization treatment,
seedlings were grown at 5°C in short-day conditions (8 h light, 16 h darkness with constant 5°C) for
number of weeks indicated in the respective experiment description.
Protein co-immunoprecipitation

4 g of Arabidopsis seedlings were crosslinked and ground in fine powder by liquid N2, following
ChIP protocol[21]. Immunoprecipitation of protein complexes was done as described in[17] using Protein
A beads (Thermo Fisher Scientific, #10008D) coupled with HA antibody (CST, #3724). IP product
was resuspended with 2X NuPAGE sample buffer (Thermo Fisher Scientific, #NP0007) and
denatured at 70°C for 15 min with 100 mM DTT. The final samples were run on NuPAGE Bis-Tris
gels (Thermo Fisher Scientific, # NP0321) and subjected to in-gel digestion.

Protein bands corresponding to heavy and light chains of antibody were excised from NuPAGE gels.
The rest of gels were combined and performed in-gel digestion as described previously[52]. Briefly, gel
pieces were destained and then incubated with 10 ng/µl trypsin (Promega, #V5111) overnight at 37°C.
Peptides were extracted by incubation with 5% trifluoroacetic acid (TFA) for 1 h, followed by
addition of 2.5% TFA/50% acetonitrile (ACN) for 1 h at 37°C. The combined supernatants were dried
in a SpeedVac concentrator (Thermo Fisher Scientific, #SPD120) for mass spectrometry (MS)
analysis.

Mass spectrometry
Samples were analyzed by LC-MS using UltiMate™ 3000 RSLCnano System (Thermo Fisher
Scientific, #ULTIM3000RSLCNANO) interfaced with Q Exactive HF. Digested samples were loaded
on a fused silica trap column Acclaim PepMap 100, 0.075 mm x 2 cm (Thermo Fisher Scientific,
#164569). After washing for 5 min at 5 µl/min with 0.1% TFA, the trap column was brought in-line
with an analytical column (Nanoease MZ peptide BEH C18, 130A, 1.7um, 0.075 mm x250 mm,
Waters, # 186008800) for LC-MS/MS. Peptides were fractionated at 300 nL/min using a segmented
linear gradient from 4 to 90% B (A: 0.1% formic acid, B: 0.08% formic acid, 80% ACN): 5 min, 4-
10% B; 60 min, 10-40% B; 15 min, 40-55% B; 10 min, 55-90% B. Mass spectrometry data was
acquired using a data-dependent acquisition procedure with a cyclic series of a full scan acquired with
a resolution of 120,000 followed by MS/MS scans (30% of collision energy in the HCD cell) with
resolution of 30,000 of 20 most intense ions with dynamic exclusion duration of 20 sec.

The peak list of the LC-MS/MS were generated by Thermo Proteome Discoverer (v. 2.1) into
MASCOT Generic Format (MGF) and searched against TAIR protein database (containing 35,386
sequences) using an in-house version of X! tandem (SLEDGEHAMMER (2015.09.01), the gpm.org)
with carbamidomethylation on cysteine as fixed modification and oxidation of methionine as variable
modification. A +/-7 ppm and 20 ppm were used as tolerance for precursor and product ions
respectively. Trypsin/P was selected as the digestion enzyme and one missed cleavage per peptide
was allowed. The False Discovery Rate was estimated for all samples by using a reverse database
(FDR<0.01). All identified spectra belong to Arabidopsis database and peptide log (E) < = -2 (E
value<=0.01) was reported. Spectral counts of the proteins in each sample were used for relative
quantitation.

Yeast two hybrid analysis
Yeast two hybrid was performed as described elsewhere[53]. Yeast cultures were grown at 28°C on
Yeast Peptone Dextrose (YPD) or on selective Synthetic Dextrose (SD) media. The AH109 strain was
used for transformation, following protocol described in the Yeast Protocols Handbook (version no.
PR973283 21; Clontech), with both Gal4-BD and Gal4-AD constructs added. Transformants were
selected on SD medium lacking Trp and Leu (SD-LW). Transformed yeast were grown to OD600 and
spotted onto selective SD medium additionally lacking His (SD-LWH) which permitted the
identification of weak protein interactions. Yeast dilutions during spotting were: 1/1, 1/5, 1/25, 1/25.

Expression analysis
Total RNA extraction was performed using the hot phenol method, as described elsewhere[54].
Genomic DNA contamination was removed using TURBO DNA-free kit (Invitrogen, #AM1907)
following the manufacturer's guidelines. cDNA was synthesised using SuperScript IV First-strand
Synthesis System (Invitrogen, #18091050), using gene-specific primers. cDNA was diluted 10 times

before qPCR. All primers are listed in the Table S2. cDNA was amplified using SYBR Green I
Master (Roche, 04887352001) and run on LightCycler 480 machine (Roche, 05015243001). Ct values
were normalized to geometric mean of reference genes: *UBIQUITIN CARRIER PROTEIN 1 (UBC)*
and/or *SERINE/THREONINE PROTEIN PHOSPHATASE 2A (PP2A)*. Chromatin-associated RNA
has been isolated using urea as published before[55].

Nucleosome stability analysis

The protocol has been adapted from[56]. 2g Arabidopsis seedlings were ground in liquid nitrogen,
resuspended in 35 ml NIB buffer (10 mM MES-KOH pH=5.4, 10 mM NaCl, 10 mM KCl, 2.5 mM
EDTA, 250 mM Sucrose, 0.1 mM Spermine, 0.5 mM Spermidine, 1mM DTT, 1x Protease inhibitor
cocktail EDTA-free (Thermo Fisher Scientific, # A32965)) and filtered through one layer of
miracloth. 20% Triton X-100 was added to final concentration of 0.35%, the extract was rotated at
4°C for 10 min and spun at 1000 g for 10 min. The pellet was transferred to Low Protein Binding
tubes (Thermo Fisher Scientific, #90410) and washed twice with MNase buffer (10 mM Tris-HCl
pH=8, 5 mM NaCl, 2.5 mM CaCl2, 2 mM MgCl2, 1x Protease inhibitor cocktail EDTA-free (Thermo
Fisher Scientific, # A32965))) at 1000g for 3 min. The pellet was weighted and MNase buffer was
added in ratio 5 µL buffer:1 µg pellet. The mixture was pre-warmed at 37°C for 5 min. 150 µL of the
mixture was put aside as an uncut sample and other 150 µL was subjected to 5U/ml (final
concentration) MNase (Takara Bio, #2910A) enzymatic digestion at 37°C for 10 min. The
concentration of MNase was experimentally tested to obtain nucleosome ladder as in[47]. The reaction
was stopped by adding EGTA to final concentration of 2 mM. The mixture was spun at 1000 g for 3
min, supernatant was removed, the pellet resuspened in 80 mM salt buffer (80 mM NaCl) and rotated
at 4°C for 15 min. The samples were spun at 1000 g for 3 min and the supernatant was collected as
mobile nucleosome fraction. Samples were adjusted to 500 µL and subjected to DNA recovery by
adding 1/50 volume 0.5 M EDTA, 1/50 vol. 5M NaCl and 2 µL RNAse A (Thermo Fisher Scientific,
#EN0531). The samples were incubated at 37°C for 30 min. Proteins were removed by adding 1/16
vol. 10% SDS to final concentration 0.63% and 2.5 µL Proteinase K (Ambion, #AM2546), followed
by incubation at 50°C for 1 h. The samples were then subjected to standard phenol-chloroform-
isoamyl alcohol and DNA precipitation with ethanol. The enrichment was calculated by qPCR with
primers indicated in Table S2. Ct values in mobile nucleosome fraction were normalized to uncut
DNA and Gypsy-like transposon (At4g07700) as an internal control, following normalizations for
published in[57].

Micrococcal nuclease assay (MNase-qPCR)

2g Arabidopsis seedlings were ground in liquid nitrogen and the pellet was resuspended in Honda
buffer[21], filtered through miracloth and spun at 1000 g, 4°C for 10 min, similarly to ChIP protocol.
Subsequently, a pellet was washed twice in MNase buffer (10 mM Tris-HCl pH=8, 5 mM NaCl, 2.5
mM CaCl2, 2 mM MgCl2, 1x Protease inhibitor cocktail EDTA-free (Thermo Fisher Scientific, #
A32965))) and finally resuspended in 1 mL MNase buffer. The samples were divided into 300 µL
uncut control and 300 µL MNase-digested sample. The mixtures were prewarmed at 37°C for 5 min
and digested samples were treated with 66.6 U/mL (final conc.) MNase enzyme (Takara Bio,
#2910A) at 37°C for 30 min. The reactions were stopped by adding 1/100 volume 0.5 M EDTA and
subjected to protein extraction by adding 0.05% (final conc.) SDS, followed by standard phenol-
chloroform-isoamyl alcohol and DNA precipitation with isopropanol. The enrichment was calculated
by qPCR with primers indicated in Table S2. Ct values in mobile nucleosome fraction were
normalized to uncut DNA.

Formaldehyde-assisted isolation of regulatory elements (FAIRE)

380 FAIRE was performed as shown in[57]. The enrichment was assessed by qPCR using primers indicated
381 in Table S2. Calculations were performed using ΔΔCt method as described in[57] with normalization of

crosslinked sample (FAIRE) to non-crosslinked sample (UN-FAIRE) and then to UBC9
(AT4G27960) as an internal control.

Chromatin immunoprecipitation

Histone modification and PolII ChIP was performed as previously described[17,55], respectively.
Immunoprecipitation was done with antibodies: anti-H3 (Abcam, #ab1791), anti-H3K27me3
(Millipore, #07-449), H2AK119ub (Cell Signalling Technology, #8240), anti-H3K36me3 (Abcam,
#ab9050). PolII antibodies were as in[58]. All ChIP experiments were followed by qPCR with indicated
primer pairs (Table S2).

Statistical analysis

Statistical analyses were performed in Microsoft Excel and R Studio (R v4.0.2). P-value, p-value
levels and sample number are included in figures and figure legends. Significance calculations with
Student T-test were precluded by F-test to determine the usage of heteroscedastic or homoscedastic
test. Unless otherwise stated, Student T-tests were two-tail. P-value levels were marked by asterisks
on the figures as follows: * ($p \leq 0.05$), ** ($p \leq 0.01$), *** ($p \leq 0.001$).

References

1. Schuettengruber, B., Bourbon, H. M., Di Croce, L. & Cavalli, G. Genome Regulation by Polycomb and Trithorax: 70 Years and Counting. *Cell* **171**, 34–57 (2017).
2. Di Croce, L., Helin, K. Transcriptional regulation by Polycomb group proteins. Nat Struct Mol Biol **20**, 1147–1155 (2013).
3. Fursova, N. A. *et al.* Synergy between Variant PRC1 Complexes Defines Polycomb-Mediated Gene Repression. *Mol. Cell* **74**, 1020-1036.e8 (2019).
4. Scelfo, A. *et al.* Functional Landscape of PCGF Proteins Reveals Both RING1A/B-Dependent-and RING1A/B-Independent-Specific Activities. *Mol. Cell* **74**, 1037-1052.e7 (2019).
5. Kahn, T. G. *et al.* Interdependence of PRC1 and PRC2 for recruitment to Polycomb Response Elements. *Nucleic Acids Res.* **44**, 10132–10149 (2016).
6. Kahn, T. G., Stenberg, P., Pirrotta, V. & Schwartz, Y. B. Combinatorial Interactions Are Required for the Efficient Recruitment of Pho Repressive Complex (PhoRC) to Polycomb Response Elements. *PLoS Genet.* **10**, e1004495 (2014).
7. Alfieri, C. *et al.* Structural basis for targeting the chromatin repressor Sfmbt to Polycomb response elements. *Genes Dev.* **27**, 2367–2379 (2013).
8. Frey, F. *et al.* Molecular basis of PRC1 targeting to polycomb response elements by PhoRC. *Genes Dev.* **30**, 1116–1127 (2016).
9. Klose, R. J., Cooper, S., Farcas, A. M., Blackledge, N. P. & Brockdorff, N. Chromatin Sampling - An Emerging Perspective on Targeting Polycomb Repressor Proteins. *PLoS Genet.* **9**, e1003717 (2013).
10. Berry, S., Dean, C. & Howard, M. Slow Chromatin Dynamics Allow Polycomb Target Genes to Filter Fluctuations in Transcription Factor Activity. *Cell Syst.* **4**, 445-457.e8 (2017).
11. Michaels, S. D. & Amasino, R. M. *FLOWERING LOCUS C* encodes a novel MADS domain protein that acts as a repressor of flowering. *Plant Cell* **11**, 949-956 (1999).
12. Sheldon, C. C. The molecular basis of vernalization: The central role of *FLOWERING LOCUS C* (*FLC*). *Proc. Natl. Acad. Sci.* **97**, 3753-3758 (2000).
13. Shindo, C. *et al.* Role of *FRIGIDA* and *FLOWERING LOCUS C* in determining variation in flowering time of Arabidopsis. *Plant Physiol.* **138**, 1163–1173 (2005).
14. Johanson, U. *et al.* Molecular analysis of *FRIGIDA*, a major determinant of natural variation in Arabidopsis flowering time. *Science* **290**, 344–347 (2000).
15. Swiezewski, S., Liu, F., Magusin, A. & Dean, C. Cold-induced silencing by long antisense transcripts of an Arabidopsis Polycomb target. *Nature* **462**, 799-802 (2009).
16. Berry S., Hartley M., Olssen TS, Dean C. Howard M. Local chromatin environment of a Polycomb target instructs its own epigenetic inheritance. *eLife* **4**, e07205 (2015).
17. Yang, H. *et al.* Distinct phases of Polycomb silencing to hold epigenetic memory of cold in

434   Arabidopsis. *Science.* **357**, 1142–1145 (2017).

435   18.   De Lucia, F., Crevillen, P., Jones, A. M. E., Greb, T. & Dean, C. A PHD-Polycomb
436   Repressive Complex 2 triggers the epigenetic silencing of *FLC* during vernalization. *Proc.*
437   *Natl. Acad. Sci. U. S. A.* **105**, 16831–16836 (2008).

438   19.   Finnegan, JE. Time-dependent stabilization of the +1 nucleosome is an early step in the
439   transition to stable cold-induced repression of *FLC. Plant J.* **84**, 875–885 (2015).

440   20.   Wood, C. C. *et al.* The *Arabidopsis thaliana* vernalization response requires a Polycomb-like
441   protein complex that also includes VERNALIZATION INSENSITIVE 3. *Proc. Natl. Acad.*
442   *Sci. U. S. A.* **103**, 14631-14636 (2006).

443   21.   Qüesta, J. I. *et al.* Arabidopsis transcriptional repressor VAL1 triggers Polycomb silencing at
444   *FLC* during vernalization. *Science* **353**, 485–8 (2016).

445   22.   Marquardt, S. *et al.* Functional Consequences of Splicing of the Antisense Transcript
446   *COOLAIR* on *FLC* Transcription. *Mol. Cell* **54**, 156–165 (2014).

447   23.   Baile, F., Merini, W., Hidalgo, I. & Calonje, M. EAR domain-containing transcription factors
448   trigger PRC2-mediated chromatin marking in Arabidopsis. *Plant Cell,* koab139 (2021).

449   24.   Deka, B. & Singh, K. K. Multifaceted regulation of gene expression by the Apoptosis- and
450   Splicing-Associated Protein complex and its components. *Int. J. Biol. Sci.* **13**, 545–560 (2017).

451   25.   Henikoff, S., Henikoff, J. G., Sakai, A., Loeb, G. B. & Ahmad, K. Genome-wide profiling of
452   salt fractions maps physical properties of chromatin. *Genome Res.* **19**, 460–469 (2009).

453   26.   Yang, H., Howard, M. & Dean, C. Antagonistic roles for H3K36me3 and H3K27me3 in the
454   cold-induced epigenetic switch at Arabidopsis *FLC. Curr. Biol.* **24**, 1793–1797 (2014).

455   27.   Xu, L. & Shen, W. H. Polycomb Silencing of KNOX Genes Confines Shoot Stem Cell Niches
456   in Arabidopsis. *Curr. Biol.* **18**, 1966–1971 (2008).

457   28.   Chen, D., Molitor, A., Liu, C. & Shen, W. H. The Arabidopsis PRC1-like RING-finger
458   proteins are necessary for repression of embryonic traits during vegetative growth. *Cell Res.*
459   **20**, 1332–1344 (2010).

460   29.   Merini, W. *et al.* The Arabidopsis Polycomb Repressive Complex 1 (PRC1) components
461   AtBMI1A, B, and C impact gene networks throughout all stages of plant development. *Plant*
462   *Physiol.* **173**, 627–641 (2017).

463   30.   Bratzel, F., López-Torrejón, G., Koch, M., Del Pozo, J. C. & Calonje, M. Keeping cell identity
464   in arabidopsis requires PRC1 RING-finger homologs that catalyze H2A monoubiquitination.
465   *Curr. Biol.* **20**, 1853–1859 (2010).

466   31.   Yang, C. *et al.* VAL-and AtBMI1-Mediated H2Aub initiate the switch from embryonic to
467   postgerminative growth in Arabidopsis. *Curr. Biol.* **23**, 1324–1329 (2013).

468   32.   Yuan, L. *et al.* The transcriptional repressors VAL1 and VAL2 recruit PRC2 for genome-wide
469   Polycomb silencing in Arabidopsis. *Nucleic Acids Res.* **49**, 98–113 (2021).

470   33.   Sun, Q., Csorba, T., Skourti-Stathaki, K., Proudfoot, N. J. & Dean, C. R-loop stabilization
471   represses antisense transcription at the Arabidopsis *FLC* locus. *Science* **340**, 619–621 (2013).

472   34.   Zhu, Y. *et al.* The Arabidopsis Nodulin Homeobox Factor AtNDX interacts with
473   AtRING1A/B and negatively regulates abscisic acid signaling. *Plant Cell* **32**, 703–721 (2020).

474   35.   Xu, C. *et al.* R-loop resolution promotes co-transcriptional chromatin silencing. *Nat. Commun.*
475   **12**, 1790 (2021).

476   36.   Crevillén, P., Sonmez, C., Wu, Z. & Dean, C. A gene loop containing the floral repressor *FLC*
477   is disrupted in the early phase of vernalization. *EMBO J.* **32**, 140–148 (2013).

478   37.   Zhou, Y., Romero-Campero, F. J., Gómez-Zambrano, Á., Turck, F. & Calonje, M. H2A
479   monoubiquitination in *Arabidopsis thaliana* is generally independent of LHP1 and PRC2
480   activity. *Genome Biol.* **18**, 69 (2017).

481   38.   De, S., Cheng, Y., Sun, M. A., Gehred, N. D. & Kassis, J. A. Structure and function of an
482   ectopic Polycomb chromatin domain. *Sci Adv.* **5**, eaau9739 (2019)

483   39.   Sengupta, A. K., Kuhrs, A. & Müller, J. General transcriptional silencing by a Polycomb
484   response element in Drosophila. *Development* **131**, 1959–1965 (2004).

485   40.   Christen, B. & Bienz, M. Imaginal disc silencers from *Ultrabithorax*: evidence for Polycomb
486   response elements. *Mech. Dev.* **48**, 255–266 (1994).

487   41.   Schwerk, C. *et al.* ASAP, a Novel Protein Complex Involved in RNA Processing and
488   Apoptosis. *Mol. Cell. Biol.* **23**, 2981–2990 (2003).

489   42.   Barbour, H., Daou, S., Hendzel, M. & Affar, E. B. Polycomb group-mediated histone H2A
490   monoubiquitination in epigenome regulation and nuclear processes. *Nature Communications*
491   **11**, 5947 (2020).
492   43.   Zhang, X. N. *et al.* Transcriptome analyses reveal SR45 to be a neutral splicing regulator and a
493   suppressor of innate immunity in *Arabidopsis thaliana*. *BMC Genomics* **18**, 772 (2017).
494   44.   Albaqami, M., Laluk, K. & Reddy, A. S. N. The Arabidopsis splicing regulator SR45 confers
495   salt tolerance in a splice isoform-dependent manner. *Plant Mol. Biol.* **100**, 379-390 (2019).
496   45.   Chen, S. L. *et al.* Quantitative Proteomics Reveals a Role for SERINE/ARGININE-Rich 45 in
497   Regulating RNA Metabolism and Modulating Transcriptional Suppression via the ASAP
498   Complex in *Arabidopsis thaliana*. *Front. Plant Sci.* **10**, 1116 (2019).
499   46.   Bi, Y. *et al.* Arabidopsis ACINUS is O-glycosylated and regulates transcription and alternative
500   splicing of regulators of reproductive transitions. *Nat. Commun.* **12**, 945 (2021).
501   47.   Chhun, T. *et al.* HSI2 repressor recruits MED13 and HDA6 to down-regulate seed maturation
502   gene expression directly during Arabidopsis early seedling growth. *Plant Cell Physiol.* **57**,
503   1689-1706 (2016).
504   48.   Zeng, X. *et al.* HISTONE DEACETYLASE 9 functions with Polycomb silencing to repress
505   *FLOWERING LOCUS C* expression. *Plant Physiol.* **182**, 555-565 (2020).
506   49.   Csorba, T., Questa, J. I., Sun, Q. & Dean, C. Antisense *COOLAIR* mediates the coordinated
507   switching of chromatin states at *FLC* during vernalization. *Proc. Natl. Acad. Sci. U. S. A.* **111**,
508   16160–16165 (2014).
509   50.   Li, P., Tao, Z. & Dean, C. Phenotypic evolution through variation in splicing of the noncoding
510   RNA *COOLAIR*. *Genes Dev.* **29**, 696–701 (2015).
511   51.   Qin, F. *et al.* Arabidopsis DREB2A-interacting proteins function as RING E3 ligases and
512   negatively regulate plant drought stress-responsive gene expression. *Plant Cell* **20**, 1693–1707
513   (2008).
514   52.   Rappsilber, J., Mann, M. & Ishihama, Y. Protocol for micro-purification, enrichment, pre-
515   fractionation and storage of peptides for proteomics using StageTips. *Nat. Protoc.* **2**, 1896–
516   1906 (2007).
517   53.   Mikulski, P. *et al.* The chromatin-associated protein PWO1 interacts with plant nuclear lamin-
518   like components to regulate nuclear size. *Plant Cell* **31**, 1141–1154 (2019).
519   54.   Box, M. S., Coustham, V., Dean, C. & Mylne, J. S. Protocol: A simple phenol-based method
520   for 96-well extraction of high quality RNA from Arabidopsis. *Plant Methods* **7**, 7 (2011).
521   55.   Wu, Z. *et al.* Quantitative regulation of *FLC* via coordinated transcriptional initiation and
522   elongation. *Proc. Natl. Acad. Sci. U. S. A.* **113**, 218–223 (2016).
523   56.   Teves, S. S. & Henikoff, S. Salt fractionation of nucleosomes for genome-wide profiling.
524   *Methods Mol. Biol.* **833**, 421–432 (2012).
525   57.   Omidbakhshfard, M. A., Winck, F. V., Arvidsson, S., Riaño-Pachón, D. M. & Mueller-
526   Roeber, B. A step-by-step protocol for formaldehyde-assisted isolation of regulatory elements
527   from *Arabidopsis thaliana*. *J. Integr. Plant Biol.* **56**, 527–538 (2014).
528   58.   Nojima, T. *et al.* Mammalian NET-seq reveals genome-wide nascent transcription coupled to
529   RNA processing. *Cell* **161**, 526–540 (2015).

Acknowledgements

This work was funded by the BBSRC grant BB/P006590/1, Wellcome Trust grant 210654/Z/18/Z and supported by the BBSRC Institute Strategic Programmes GRO (BB/J004588/1) and GEN (BB/P013511/1) and a Royal Society Professorship to CD. The authors would like to thank all members of the Dean lab for discussions.

Author information

Affiliations

Cell and Developmental Biology, John Innes Centre, Norwich, UK

Pawel Mikulski, Philip Wolff, Tiancong Lu, Danling Zhu & Caroline Dean

Present address: SUSTech-PKU Institute of Plant and Food Science, Department of Biology, Southern University of Science and Technology, Shenzhen, China

Danling Zhu

Present address: State Key Laboratory of Plant Genomics and National Center for Plant Gene

Research, Institute of Genetics and Developmental Biology, Chinese Academy of Sciences,

Beijing, China

Tiancong Lu

Present address: Department of Biochemistry, University of Oxford, Oxford, UK

Pawel Mikulski

Contributions

PM, PW and CD conceived the study. PM performed: yeast-two-hybrid, FAIRE, nucleosome

mobility, ChIPs and expression analyses in *val1* and PRC1 mutants. PW performed: expression

analyses, histone marks and PolII ChIPs for ASAP mutants. DZ performed nucleosome position

experiment. TL performed protein co-immunoprecipitation with mass spectrometry. PM, PW and CD

designed the experiments and analyzed the data. CD lead the general concept of research. PW wrote

the paper together with CD with all authors contributing to different relevant sections of the

manuscript.

Ethics declarations

Competing interest

The authors declare no competing interest.

Additional information

Figure legends

Fig.1. *asap* mutants affect pre-cold *FLC* expression only. a. Yeast-two-hybrid of the VAL1-SAP18

interaction. Yeast growth on selective medium SD-LWH is shown. Top panel corresponds to

decreasing yeast culture concentration with dilutions: 1/1, 1/5. 1/25 from OD600 = 0.8. AD = pGAD

vector backbone with Gal4 activating domain; BD = pGBKT vector backbone with Gal4 binding

domain; "empty" = empty vector without insertion as negative control. Red frame depicts protein

pairs showing yeast growth over negative controls. **b.** Schematic summary of interaction from a. **c.**

FLC downregulation slope during vernalization (fold change to NV). Values were normalized as

relative expression to UBC. N = 3 biological replicates, error bars = propagated SEM. **d.** *FLC*

expression in ASAP single mutants and different double mutant combinations within ASAP or with

VAL1. Values correspond to mean relative expression to UBC. Statistics are calculated with two-tail

Student T-test in comparison to wildtype (* = p≤0.05; ** = p≤0.01; *** = p≤0.001). N = 3 biological

replicates; error bars = SEM. **e.** Chromatin-bound *FLC* RNA enrichment over the locus. Shown as a

fold change to wildtype. X-axis labels indicate midpoint of the amplicon relative to TSS and its

position in *FLC* locus. N = 3 biological replicates, error bars = propagated SEM.

Fig.2. SR45 co-transcriptionally represses *FLC*. a. RNA Polymerase II ChIP enrichment. X-axis

depicts the midpoint position of amplicon at the *FLC* locus, 0 =TSS. Y-axis corresponds to %

recovery to input values normalized to housekeeping gene *ACT7*. NV = non-vernalized; n = 3

biological replicates; error bars = SEM. **b.** SR45-dependent change of active-repressive histone

marks. X-axis depicts midpoint position of amplicon at the *FLC* locus. Y-axis corresponds to %

recovery to input values normalized to H3 enrichment. NV = non-vernalized; n = 3 biological

replicates; error bars = SEM.

Fig.3. *FLC* chromatin regulation by VAL1. a. Chromatin-bound *FLC* RNA enrichment over *FLC*

locus. Shown as a fold change to wildtype. X-axis labels indicate midpoint of the amplicon relative to

TSS and its position in *FLC* locus. N = 3 biological replicates, error bars = propagated SEM. **b.** Low

salt-extractable nucleosome fraction over *FLC*. Shown as % recovery to uncut DNA and fold to
AT4G07700 (Gypsy-like transposon). X-axis labels indicate beginning of amplicon relative to TSS,
lines below axis' label indicate regions of increased low-salt nucleosome extractability in *val1* shown
in b. N = 3 biological replicates; error bars = SEM; asterisk = p<0.05 from two-tail Student T-test in
comparison wildtype vs *val1*; n.s. = non-significant; NV = non-vernalized. **c.** Nucleosome position at
FLC in wildtype. Results from MNase-qPCR assay shown as % to uncut DNA. X-axis indicate
beginning of amplicon relative to TSS. N = 2 biological replicates, error bars = SEM. **d.** FAIRE
enrichment at *FLC*. Shown as % recovery to non-crosslinked sample (UN-FAIRE) and fold to UBC9.
X-axis labels indicate beginning of amplicon relative to TSS. N = 2 biological replicates; error bars =
SEM; NV = non-vernalized.
Fig.4. H2Aub dynamics at *FLC*. a. H2Aub chromatin -IP showing strong enrichment of H2AUb at
the 5' end of *FLC* in ColFRI correlating with increased transcription in non-vernalized (NV)
conditions. X-axis show midpoint amplicon at *FLC*. N = 3 biological replicates; error bars = SEM;
NoAb = no antibody negative control. *FLC* expression in wildtype backgrounds in NV is depicted in
graph on the right. N = 3 biological replicates, error bars = SEM. **b.** H2Aub chromatin -IP correlating
cold-induced decrease in *FLC* expression with decreased H2AUb at the 5' end of *FLC*. X-axis show
midpoint amplicon at *FLC*. Graph on the right shows unspliced *FLC* expression during vernalization
timecourse. NV = non-vernalized; Nominal-W = number of weeks of cold; N = 3 biological
replicates; error bars = SEM; NoAb = no antibody negative control. **c.** Histone mark behaviour post-
cold. H3K27me3 and H2Aub enrichment over *FLC*. X-axis show midpoint amplicon at *FLC*. N = 3
biological replicates; error bars = SEM; 6W = 6 weeks of cold treatment; T-nominal = number of days
post-cold; NoAb = no antibody negative control.
Fig.5. VAL1, BMI1B and NDX effect on *FLC* expression. a. *FLC* expression in mutants of VAL1,
PRC1 and NDX for *FLC* RNA. Values correspond to mean relative expression to geometric mean of
UBC and PP2A. Statistics were calculated with two-tail Student t-test in comparison to wildtype. N =
3 biological replicates; error bars = SEM. **b.** *FLC* expression level. Represented as relative expression
to housekeeping genes (see methods). Statistics represent two-tail Student t-test in comparison to the
wildtype, N = 3 biological replicates, error bars = SEM, NV = non-vernalized, W = number of weeks
of cold. **c.** The same data as in **b.** presented as *FLC* downregulation slope during vernalization (fold
change to NV). **d.** H2Aub enrichment at *FLC* in wildtype and *val1 bmi1B*. N = 3 biological replicates;
error bars = SEM; NoAb = no antibody negative control.
Fig.6. VAL1, PRC1 and NDX physically interact and are required for H2Aub accumulation at
the *FLC* nucleation region. a. IP-MS/MS results for PRC1 and NDX from crosslinked VAL1-HA
co-IP. Numbers corresponds to unique peptides. IP-numeral = biological replicate; NV = non-
vernalized; Numeral-W = number of weeks of cold; T-numeral = number of days post-cold; (-) =
negative control. **b.** Yeast-2-hybrid results for PRC1-NDX interaction. Yeast growth on selective
medium (-LWH) is shown. Top panel corresponds to decreasing yeast culture concentration. AD =
pGAD vector backbone with Gal4 activating domain; BD = pGBKT vector backbone with Gal4
binding domain; "empty" = empty vector without insertion as negative control. Red frame depicts
protein pairs showing yeast growth over negative controls (pairs with empty vector). precipitated with
anti-HA antibody. **c.** Schematic summary of interactions from **a.** and **b**. **e.** and **f.** H2Aub ChIP results.
X-axis represents midpoint of the amplicons over *FLC*. NoAb = no antibody negative control; N = 3
biological replicates for H2Aub IPs and 3 qPCR technical replicates for NoAb controls; error bars =
SEM.
Fig.7 Functional relationship of ASAP, PRC1 and PRC2 in *FLC* regulation. a. ASAP and PRC1
act as co-repressors that regulate *FLC* chromatin features and downregulate *FLC* expression in warm.
b. Upon shift to cold, PRC1-mediated H2Aub accumulates at *FLC* PRE along with H3K27me3.
H2Aub and H3K27me3 enrichment gradually increases along with cold duration. In post-cold,

H3K27me3 spreads to *FLC* gene body, whereas H2Aub remains localized to the *FLC* PRE. **c.** At the
later post-cold timepoints, H3K27me3 is stably maintained over the *FLC* gene body whereas H2Aub
is removed from the chromatin. H2K27me3 thus provides long-term epigenetic silencing, whereas
H2Aub is a transient repression signal.
642
643

Fig.1. *asap* mutants affect pre-cold *FLC* expression only. a. Yeast-two-hybrid of the VAL1-SAP18 interaction. Yeast growth on selective medium SD-LWH is shown. Top panel corresponds to decreasing yeast culture concentration with dilutions: 1/1, 1/5. 1/25 from OD600 = 0.8. AD = pGAD vector backbone with Gal4 activating domain; BD = pGBKT vector backbone with Gal4 binding domain; "empty" = empty vector without insertion as negative control. Red frame depicts protein pairs showing yeast growth over negative controls. **b.** Schematic summary of interaction from a. **c.** *FLC* downregulation slope during vernalization (fold change to NV). Values were normalized as relative expression to UBC. N = 3 biological replicates, error bars = propagated SEM. **d.** *FLC* expression in ASAP single mutants and different double mutant combinations within ASAP or with VAL1. Values correspond to mean relative expression to UBC. Statistics are calculated with two-tail Student T-test in comparison to wildtype (* = p≤0.05; ** = p≤0.01; *** = p≤0.001). N = 3 biological replicates; error bars = SEM. **e.** Chromatin-bound *FLC* RNA enrichment over the locus. Shown as a fold change to wildtype. X-axis labels indicate midpoint of the amplicon relative to TSS and its position in *FLC* locus. N = 3 biological replicates, error bars = propagated SEM.

Fig.2. SR45 co-transcriptionally represses *FLC*. a. RNA Polymerase II ChIP enrichment. X-axis depicts the midpoint position of amplicon at the *FLC* locus, 0 =TSS. Y-axis corresponds to % recovery to input values normalized to housekeeping gene *ACT7*. NV = non-vernalized; n = 3 biological replicates; error bars = SEM. **b.** SR45-dependent change of active-repressive histone marks. X-axis depicts midpoint position of amplicon at the *FLC* locus. Y-axis corresponds to % recovery to input values normalized to H3 enrichment. NV = non-vernalized; n = 3 biological replicates; error bars = SEM.

Fig.3. *FLC* chromatin regulation by VAL1. a. Chromatin-bound *FLC* RNA enrichment over *FLC* locus. Shown as a fold change to wildtype. X-axis labels indicate midpoint of the amplicon relative to TSS and its position in *FLC* locus. N = 3 biological replicates, error bars = propagated SEM. **b.** Low salt-extractable nucleosome fraction over *FLC*. Shown as % recovery to uncut DNA and fold to AT4G07700 (Gypsy-like transposon). X-axis labels indicate beginning of amplicon relative to TSS, lines below axis' label indicate regions of increased low-salt nucleosome extractability in *val1* shown in b. N = 3 biological replicates; error bars = SEM; asterisk = p<0.05 from two-tail Student T-test in comparison wildtype vs *val1*; n.s. = non-significant; NV = non-vernalized. **c.** Nucleosome position at *FLC* in wildtype. Results from MNase-qPCR assay shown as % to uncut DNA. X-axis indicate beginning of amplicon relative to TSS. N = 2 biological replicates, error bars = SEM. **d.** FAIRE enrichment at *FLC*. Shown as % recovery to non-crosslinked sample (UN-FAIRE) and fold to UBC9. X-axis labels indicate beginning of amplicon relative to TSS. N = 2 biological replicates; error bars = SEM; NV = non-vernalized.

Fig.4. H2Aub dynamics at *FLC*. a. H2Aub chromatin -IP showing strong enrichment of H2AUb at the 5' end of *FLC* in ColFRI correlating with increased transcription in non-vernalized (NV) conditions. X-axis show midpoint amplicon at *FLC*. N = 3 biological replicates; error bars = SEM; NoAb = no antibody negative control. *FLC* expression in wildtype backgrounds in NV is depicted in graph on the right. N = 3 biological replicates, error bars = SEM. **b.** H2Aub chromatin -IP correlating cold-induced decrease in *FLC* expression with decreased H2AUb at the 5' end of *FLC*. X-axis show midpoint amplicon at *FLC*. Graph on the right shows unspliced *FLC* expression during vernalization timecourse. NV = non-vernalized; Nominal-W = number of weeks of cold; N = 3 biological replicates; error bars = SEM; NoAb = no antibody negative control. **c.** Histone mark behaviour post-cold. H3K27me3 and H2Aub enrichment over *FLC*. X-axis show midpoint amplicon at *FLC*. N = 3 biological replicates; error bars = SEM; 6W = 6 weeks of cold treatment; T-nominal = number of days post-cold; NoAb = no antibody negative control.

Fig.5. VAL1, BMI1B and NDX effect on *FLC* expression. a. *FLC* expression in mutants of VAL1, PRC1 and NDX for *FLC* RNA. Values correspond to mean relative expression to geometric mean of UBC and PP2A. Statistics were calculated with two-tail Student t-test in comparison to wildtype. N = 3 biological replicates; error bars = SEM. **b.** *FLC* expression level. Represented as relative expression to housekeeping genes (see methods). Statistics represent two-tail Student t-test in comparison to the wildtype, N = 3 biological replicates, error bars = SEM, NV = non-vernalized, W = number of weeks of cold. **c.** The same data as in **b.** presented as *FLC* downregulation slope during vernalization (fold change to NV). **d.** H2Aub enrichment at *FLC* in wildtype and *val1 bmi1B*. N = 3 biological replicates; error bars = SEM; NoAb = no antibody negative control.

Fig.6. VAL1, PRC1 and NDX physically interact and are required for H2Aub accumulation at the *FLC* nucleation region. a. IP-MS/MS results for PRC1 and NDX from crosslinked VAL1-HA co-IP. Numbers corresponds to unique peptides. IP-numeral = biological replicate; NV = non-vernalized; Numeral-W = number of weeks of cold; T-numeral = number of days post-cold; (-) = negative control. **b.** Yeast-2-hybrid results for PRC1-NDX interaction. Yeast growth on selective medium (-LWH) is shown. Top panel corresponds to decreasing yeast culture concentration. AD = pGAD vector backbone with Gal4 activating domain; BD = pGBKT vector backbone with Gal4 binding domain; "empty" = empty vector without insertion as negative control. Red frame depicts protein pairs showing yeast growth over negative controls (pairs with empty vector). precipitated with anti-HA antibody. **c.** Schematic summary of interactions from **a.** and **b. e.** and **f.** H2Aub ChIP results. X-axis represents midpoint of the amplicons over *FLC*. NoAb = no antibody negative control; N = 3 biological replicates for H2Aub IPs and 3 qPCR technical replicates for NoAb controls; error bars = SEM.

Fig.7 Functional relationship of ASAP, PRC1 and PRC2 in *FLC* regulation. a. ASAP and PRC1 act as co-repressors that regulate *FLC* chromatin features and downregulate *FLC* expression in warm. **b.** Upon shift to cold, PRC1-mediated H2Aub accumulates at *FLC* PRE along with H3K27me3. H2Aub and H3K27me3 enrichment gradually increases along with cold duration. In post-cold, H3K27me3 spreads to *FLC* gene body, whereas H2Aub remains localized to the *FLC* PRE. **c.** At the later post-cold timepoints, H3K27me3 is stably maintained over the *FLC* gene body whereas H2Aub is removed from the chromatin. H2K27me3 thus provides long-term epigenetic silencing, whereas H2Aub is a transient repression signal.