Multiple stressor null models frequently fail to detect most interactions due to low statistical

power

Benjamin J. Burgess[1]

Michelle C. Jackson[2]

David J. Murrell[1]*

[1] **Centre for Biodiversity and Environment Research, Department of Genetics, Evolution and**

Environment, University College London, Gower Street, London, WC1E 6BT, United Kingdom

[2] **Department of Zoology, University of Oxford, Oxford, OX1 3SZ, United Kingdom**

*corresponding author (d.murrell@ucl.ac.uk)

Temporary Running Head: Limited power in multiple stressor null models

Word count: 6933

20 *Abstract*

1. Most ecosystems are subject to co-occurring, anthropogenically driven changes and

understanding how these multiple stressors interact is a pressing concern. Stressor

interactions are typically studied using null models, with the additive and multiplicative null

expectation being those most widely applied. Such approaches classify interactions as being

synergistic, antagonistic, reversal, or indistinguishable from the null expectation. Despite

their wide-spread use, there has been no thorough analysis of these null models, nor a

systematic test of the robustness of their results to sample size or sampling error in the

estimates of the responses to stressors.

2. We use data simulated from food web models where the true stressor interactions are

known, and analytical results based on the null model equations to uncover how (i) sample

size, (ii) variation in biological responses to the stressors and (iii) statistical significance,

affect the ability to detect non-null interactions.

3. Our analyses lead to three main results. Firstly, it is clear the additive and multiplicative null

models are not directly comparable, and over one third of all simulated interactions had

classifications that were model dependent. Secondly, both null models have weak power to

correctly classify interactions at commonly implemented sample sizes (i.e., ≤6 replicates),

unless data uncertainty is unrealistically low. This means all but the most extreme

interactions are indistinguishable from the null model expectation. Thirdly, we show that

increasing sample size increases the power to detect the true interactions but only very

slowly. However, the biggest gains come from increasing replicates from 3 up to 25 and we

provide an R function for users to determine sample sizes required to detect a critical effect

size of biological interest for the additive model.

4. Our results will aid researchers in the design of their experiments and the subsequent

interpretation of results. We find no clear statistical advantage of using one null model over

the other and argue null model choice should be based on biological relevance rather than

statistical properties. However, there is a pressing need to increase experiment sample sizes

otherwise many biologically important synergistic and antagonistic stressor interactions will

continue to be missed.

Keywords

Ecosystem drivers; Experimental design; Sample size; Critical effect size; Synergism; Antagonism;

Power analysis.

Introduction

Most, if not all ecosystems are currently being impacted by multiple co-occurring, anthropogenically-driven, stressors including land-use change, invasive species, climate change, and pollution (**Brook et al., 2008**; **Dirzo et al., 2014; Hillebrand et al., 2020; Jackson et al., 2020**), and these stressors (also known as disturbances, drivers, factors or pressures, **Orr et al. 2020**) can have a wide range of effects on biodiversity and ecosystem services (**Newbold et al., 2015; Sala et al., 2021; Tittensor et al., 2014; Vinebrooke et al., 2004**). A key goal of ecology is to determine whether the combined effect of multiple stressors is more than (synergistic) or less than (antagonistic) that expected by their individual effects (**Orr et al. 2020**). Such an understanding is crucial to designing management strategies to mitigate against the effects of the stressors (e.g. **Lindenmayer et al. 2020**), and indeed may prevent these strategies from making the situation worse (**Brown et al. 2013**).

The effects of multiple interacting stressors are predominately determined through the implementation of null models where the observed response is compared to an expectation that the stressors are non-interacting (**De Laender, 2018**). Of the range of available null models, the additive null model (**Gurevitch et al., 2000**) is the most widely applied (**e.g., Crain et al., 2008; Jackson et al., 2016**) and has the expectation that the overall effect of the multiple interacting stressors is simply equal to the sum of the effects of the stressors acting individually. The statistical test is therefore whether this null model can be rejected in favour of an alternative hypothesis that interactions are: i) greater than anticipated by the additive null model (*Synergistic interactions*); ii) less than the sum of the individual stressor effects (*Antagonistic interactions*); or iii) opposite to that suggested by the additive null model (*Reversal interactions*) (see **Figure 1**).

Figure 1: Comparison of the additive and multiplicative null models, illustrating the conditions under

which the different interaction classifications are observed. Examples are for both Stressor 1 and

Stressor 2 act upon the response in (a) the same direction, and (b) contrasting directions. Individual

Stressors: Grey bars represent the change in a measured response due to a given stressor (Stressor 1

or Stressor 2).   Null Model Expectations: Yellow bar represents the expected response under an

additive null model; Purple bar represents the expected response under a multiplicative null model.

Interaction Class: Arrows denote how an interaction would be classified for a given observed

response. Blue arrows denote synergistic interactions; Pink arrows denote antagonistic interactions;

Orange arrows denote reversal interactions. For the additive and multiplicative null models, a null

interaction is shown by the purple and yellow lines respectively. When comparing the additive and

multiplicative null models, the black line denotes the range of observed responses where the additive

and multiplicative null models classify interactions differently (i.e., the interaction class is null model

dependent).

While the majority of studies currently apply the additive null model there is a growing body of

literature suggesting alternative null models (**e.g., Tekin et al., 2020; Thompson et al., 2018**) that

may potentially be more appropriate when investigating the impacts of multiple stressors (**Dey &**

Koops, 2021; Schäfer & Piggott, 2018). The greatest issue with the additive null model is that it can

lead to biologically meaningless expectations. For example, if stressor A (perhaps a pollutant)

reduces biomass of our target species by 60%; and stressors B (perhaps an invasive species) reduces

it by 50%, then the expected joint effects under the additive model would be a reduction in biomass

of 110%. Similar problems arise for other responses that may have bounded values (e.g. survival

and/or mortality).

The most common alternative is the multiplicative null model (**Hawkes & Sullivan, 2001; Lajeunesse,**

2011), where the expected interaction is simply the product of the responses to the two individual

stressors acting alone. Under our example above the null expectation for the stressor interaction

would be for a reduction in biomass of 80% (i.e. (0.6+0.5) − (0.5*0.6)). As such, it has been suggested

that a multiplicative null model may better reflect biological and ecological systems because it can

prevent biologically implausible expectations (**Stephens et al. 2013**). Deviations away from the null

expectation can follow the same nomenclature as for the additive model, but the same interaction

can result in a different classification under the two null models (**Figure 1**). Very few analyses have

compared interactions in the same data using both null models (see **Stephens et al., 2013**), but

model dependent interaction types means (i) management decisions can vary greatly depending on

which null model is used, with the potential for remedial action to be ineffective or even detrimental

if the 'wrong' null model is chosen, and (ii) comparisons of experiments or reviews that use different

null models are very difficult to interpret as we do not yet know the circumstances where the model

inferences will differ. It is therefore imperative to understand how the additive and multiplicative

null models inherently relate to one another.

An arguably greater problem is that there is very little understanding of null model robustness to

data uncertainty (variation), or consequently, the sample sizes required to detect non-null

interactions. This means there is no guidance of how many replicates are required to guarantee the

detection of a non-null interaction of a given strength should it exist in the data, and researchers

have been left to base the crucial decisions on experimental design solely on heuristics or economic

constraints (**Lakens, 2021**). Without this guidance there is a risk that many experiments will be

statistically underpowered and lack the ability to reject the null model for all but the most extreme

cases. Indeed, several reviews have reported >70% incidences of null interactions (**Stephens et al.,**

2013; Lange et al., 2018; Burgess et al., 2021); but the failure to reject the null model prediction is

not proof it is true, and we cannot be sure how many biologically important ecological surprises

remain hidden from view due to data constraints. As a result, discussions regarding the relative

commonness or rarity of ecological surprises (**e.g., Darling and Côté, 2008; Côté et al., 2016**) cannot

reach any firm conclusions until the ability of the null models to detect the interactions has been

quantified.

Recently, **Burgess et al. (2021)** simulated data using food web models where parameters were

altered under pairs of 'ecological stressors' to investigate the ability of the additive null model to

uncover the true classification in the face of various levels of measurement/observation error. They

found that it can be very difficult to correctly reject the additive null model at even very low levels of

observation error, and this means many strong synergies and antagonisms may be left classified as

indistinguishable from additive. However, the analyses of **Burgess et al. (2021)** fall short of a power

analysis because they did not investigate the relationship between detection of true stressor-pair

interactions and the sample size of the data for the treatment means. The number of replicates in

experiments to investigate stressor interactions rarely reach double figures. As examples, two recent

meta-analyses (**Gomez Isaza et al., 2020; Seifert et al., 2020**) included no experiments with more

than six replicates per treatment, while a third (**Burgess et al., 2021**) found <1% of experiments used

more than eight replicates per treatment (**Supplementary Information**). It remains unknown

whether increasing current sample sizes would greatly improve the statistical power of the additive

null model and potentially lead to fundamental changes in our understanding of how pair of

ecosystem stressors interact. A further open question is to how general the results of **Burgess et al.**

(2021) are to other null models, or whether the additive model is statistically a poor choice to detect

stressor interactions.

Here, we focus on closing these important gaps in knowledge by focussing on the additive and

multiplicative null models and answering the following questions: (1) What are the conditions under

which the additive and multiplicative null models agree, and disagree? (2) To what degree does

treatment uncertainty (in the form of observation error) reduce the ability of both the additive and

multiplicative null models to correctly infer the 'true' interaction classification? (3) What is the

relationship between the power to correctly reject the null models and sample size, and are current

ranges of sample sizes adequate to detect most non-null interactions?

Methods

The null model approach requires a factorial experiment design with four treatments that each

measure the same metric of interest (e.g., individual survival; population density or biomass; species

richness). Each measure X_x, is the mean value of this metric taken over N_x replicates, where

$x \in \{C, A, B, I\}$. The first treatment, C, is the control which is simply the system (i.e., population,

community) of interest with no stressor. There are two treatments (A, B) that account for the

individual effects of each of the stressors of interest in isolation. The final treatment, I, is the

estimate of the effect of both stressors acting simultaneously. Associated with each treatment is an

estimate of the standard deviation of the response to the treatment, and these are denoted by SD_X,

where $x \in \{C, A, B, I\}$. All three elements, X_x, SD_X, and N_x are required for the additive and

multiplicative null models which are defined below, and from this input each null model computes

an effect size, with associated confidence intervals from which the interaction type is inferred.

Additive null model

The additive null model compares the observed interaction effect $X_O = X_I - X_C$, with $X_E = X_A +$

$X_B - 2X_C$, the response expected under the assumption the joint effect is equal to the sum of the

individual effects of stressors A and B. To compute effect sizes we use Hedges' d which is unbiased

by small sample sizes (**Hedges & Olkin, 1985**). The calculation of the additive effect size, (ES_{Add}), is

given by Equation 1.1, and this requires a pooled sampling standard deviation, denoted by s

(Equation 1.2); while $J(m)$ (Equation 1.3) is the small sample bias correction factor (**Borenstein et al.,**

2009). For each additive effect size a corresponding variance, (V_{Add}), standard error, (SE_{Add}), and

confidence intervals, (CI_{Add}), are calculated by Equations 1.4 - 1.6. Here, we set the significance level

to 0.05 ($Z_{\alpha/2}$ = 1.96), hence 95% confidence intervals are calculated.

1.1)
$$ES_{Add} = \frac{X_I - X}{\cdot J(m)}$$

1.2) s

$$= \sqrt{\frac{(N_I - 1) \cdot (SD_I)^2 \; + \; (N_A - 1) \cdot (SD_A)^2 \; + \; (N_B - 1) \cdot (SD_B)^2 \; + \; (N_C - 1)}{N_I + N_A + N_B + N_C - 4}}$$

1.3)
$$J(m) \; = \; 1 - \frac{3}{4 \cdot (N_I + N_A + N_B + N_C - 4) - 1}$$

1.4)
$$V_{Add} \; = \; (J(m))^2 \cdot (\frac{1}{N_I} + \frac{1}{N_A} + \frac{1}{N_B} + \frac{1}{N_C} + \frac{(ES_{Add})^2}{2(N_I + N_A + N_B + N_C)})$$

1.5)
$$SE_{Add} \; = \; \sqrt{V_{Add}}$$

1.6)
$$CI_{Add} \; = \; Z_{\alpha/2} \cdot SE_{Add}$$

Multiplicative null model

The effect size for the multiplicative null model is derived from the response ratio $RR = \ln\left(\frac{X_O}{X_E}\right) =$

$\ln(X_O) - \ln(X_E)$ **Lajeunesse (2011)**, where the observed effect $X_O = \frac{X_I}{X_C}$ (the joint effect of the

stressors), and the expected effect $X_E = \frac{X_A \cdot X_B}{X_C^2}$. The multiplicative effect size, (ES_{Mul}), is then given by

Equation 2.1; while the corresponding variance, (V_{Mul}), standard error, (SE_{Mul}), and 95% confidence

intervals, (CI_{Mul}), are determined using Equations 2.2 − 2.4 (**Lajeunesse 2011**).

2.1)
$$ES_{Mul} \; = \; \ln(X_I) - \ln(X_A) - \ln(X_B) + \ln(X_C)$$

2.2)
$$V_{Mul} \; = \; \frac{(SD_I)^2}{(X_I)^2 \cdot N_I} + \frac{(SD_A)^2}{(X_A)^2 \cdot N_A} + \frac{(SD_B)^2}{(X_B)^2 \cdot N_B} + \frac{(SD_C)^2}{(X_C)^2 \cdot N_C}$$

2.3)
$$SE_{Mul} \; = \; \sqrt{V_{Mul}}$$

2.4)
$$CI_{Mul} \; = \; Z_{\alpha/2} \cdot SE_{Mul}$$

Interaction classifications

Using the naming convention of **Orr et al., (2020**) (see also Figure 1), an interaction is classified as

null if the 95% confidence intervals for that interaction overlapped zero. In this case the

interpretation is not that the null model is true, rather that we have insufficient data to reject the

null hypothesis. Interactions were classed as synergistic if effect size was positive and the 95%

confidence intervals for that interaction did not overlap zero. Antagonistic and reversal interactions

were classified whenever the effect size is negative, and the 95% confidence intervals do not overlap

zero. For an interaction to be classed as antagonistic, the expected and observed interaction effects

had to both act in the same direction (e.g., for an additive interaction: $X_I - X_C > 0$ & $X_A + X_B - 2X_C > 0$).

In contrast, for an interaction to be classed as reversal, the expected and observed interaction

effects had to both act in differing directions (e.g., for an additive interaction: $X_I - X_C < 0$ & $X_A + X_B -$

$2X_C > 0$). The method for classifying interactions is explored more in the **Supplementary**

Information.

Data simulation

We use the same model as **Burgess et al. (2021)** to simulate data from food chains where population

densities are the response metric to 'stressors' that alter key parameters in the models. We refer the

reader to the **Supplementary Information** for full details, including formulae and details of

distributions. The food chains are based on Lotka-Volterra consumer-resource equations enabling us

to establish food chains of three, four, or five trophic levels and determine the densities of

populations at equilibrium. Regardless of the length of food chain considered, the models contain

three distinct types of trophic level. These levels are: i) basal trophic level, equivalent to a pool of a

limiting nutrient such as silica; ii) consumer trophic level (including primary producers that require

the nutrient), which only consumes the trophic level directly below, and is only consumed by the

trophic level directly above; iii) apex consumer trophic level, which only consumes the trophic level

directly below, but is not itself consumed by another level.

The equilibrium densities are governed by five key parameters (α_i, ε_i, δ_i, δ_i, ω) which stay constant

over time. The attack rate, α_i, denotes the rate at which trophic level I feeds upon a resource

(trophic level i-1). The conversion efficiency, ε_i, represents the proportion of the consumed resource

that is converted into new consumers within trophic level i. The density independent mortality rate,

δ_i, represents the background mortality rate of trophic level i. The density dependent mortality rate,

λ_i, increases the mortality rate of a trophic level with increasing population density. Finally, the basal

input rate, ω, represents the constant rate at which the basal trophic level increases in density.

Stressors are simulated by altering the values of the key parameters relative to a control (baseline)

set of values and stressor interactions are computed using a factorial experiment design. Key

parameter values for each trophic level are drawn from pre-determined uniform distributions.

Hence for example, the value for the conversion efficiency parameter at trophic level i is unlikely to

be the same as that for trophic level j. Food chains are subjected to a stressor by altering the value

for a single parameter (α, ε, δ, or ω) chosen at random for a single trophic level, and keeping all

other parameters constant under the control values. All parameters are equally likely to be selected

to be stressed, although a single parameter is only ever impacted by one stressor (i.e., two stressors

cannot act on the same parameter). For the interaction treatment all parameters were under control

conditions, except for those two which were stressed in the individual treatments which were

instead assigned their stressed values. We then calculate the equilibrium densities of each trophic

level within the food chain for all four treatments using Mathematica v10.4 (**Wolfram Research Inc.,**

2016). We retained datasets where all four treatments resulted in biological feasible (i.e. all densities

non-zero) and locally stable equilibria **(Nattrass et al. 2012)**. All retained datasets showed equilibria

that were either stable nodes or stable foci. We then selected one trophic level at random (excluding

the basal level) to be the focal response group and used its equilibrium densities across the four

treatments to classify the stressor interactions using the null models (Equations 1 and 2, *Interaction*

classifications section). Using this approach, we generated a dataset comprising of 100,000

interactions, with approximately equal numbers of equilibrium densities for populations from the

three different lengths of food chain.

Treatment means, treatment uncertainty, and sample sizes

Given that the food chain models are deterministic, there is an absence of data uncertainty at this

stage. Interactions estimated in the absence of any data uncertainty are referred to as the 'true'

stressor interaction. To mirror empirical experiments, we modelled treatment uncertainty, in the

form of observation error for the equilibrium densities recorded at each treatment within each of

the 100,000 interactions. In brief, we follow the method of **Burgess et al., (2021)**, with each

equilibrium density being multiplied by a number drawn from a Gaussian distribution with a mean of

1.00 and a standard deviation σ, with this process being repeated N times per treatment. Treatment

means are then estimated by taking the mean of the values from these replicates. Smaller values of

σ will result in a tighter distribution of observed mean densities around the true population mean

compared to larger values of σ. Our approach allows for an intuitive understanding of σ, as we can

expect 99.7% of all observed treatment densities to fall within 3σ of the true value. We use 250

levels of σ, hence referred to as levels of observation error, ranging from 1.0×10^{-3} to 2.5×10^{-1} (see

Supplementary Information for more details).

By incorporating 'noise' in this way we were able to test the ability of the null models to recover the

true stressor interaction in the face of model uncertainty, and for different sample sizes. We then

compute the probability of assigning the true interaction classification to each interaction under

every level of observation error, thereby investigating the ability of each null model to correctly

classify interactions (i.e., true positives) in the face of uncertainty (see Supplementary Information

for more details).

Results

Stressor-pair classification across the null models

To compare null model relationships, each of the 100,000 simulated interactions in the absence of

uncertainty was assigned an interaction using both models. Overall, both the additive and

multiplicative null models report similar frequencies of the different interaction classifications, with

both reporting that antagonistic interactions are the most prevalent (additive 0.493; multiplicative

0.519), followed by a slightly reduced number of synergistic interactions (0.485; 0.462), minimal

frequencies of reversal interactions (0.023; 0.020), and a complete absence of null interactions

(0.000; 0.000). However, this overview masks the important fact that one third of individual

interactions change classification when moving from one null model to the other (Table 1). Null

model dependent classifications are predominately interactions assigned as synergistic by one null

model and antagonistic by the other (Table 1, see also Figure 1).

Table 1: Proportions of interactions classed differently, or the same, across additive and
multiplicative null models. Shaded boxes indicated the proportion of interactions where both null
models agree.

		Additive Null Model			
		Synergistic	Antagonistic	Reversal	Null
Multiplicative Null Model	Synergistic	0.299	0.158	0.005	0.000
	Antagonistic	0.181	0.328	0.010	0.000
	Reversal	0.005	0.007	0.007	0.000
	Null	0.000	0.000	0.000	0.000

Robust of the null models to data uncertainty (observation error)

Both null models are sensitive to data uncertainty in the form of observation error, with the correct

classification of synergistic and antagonistic interactions rapidly declining as error increases (Figure

2). In contrast, classifications of reversal interactions, although quite rare, are much less affected by

observation error. The frequency of null interactions rapidly increases with error, accounting for

over 50% of all interactions for both null models by σ~0.005 (i.e., when nearly all sampled treatment

means are within 1.5% of the true values). This shows how even low levels of observation error lead

to the failure to correctly reject the null model (i.e., low statistical power).

Figure 2: The effect of observation error on the frequency of interaction classifications, showing how

the proportions differed from those in the absence of observation error, using an additive null model

(a) and multiplicative null model (b). Frequencies of the different interaction classes for the 100,000

simulated interactions at each level of observation error using either null model. Pink line indicates

null interactions. Orange line denotes reversal interactions. Blue line indicates synergistic

interactions. Red line denotes antagonistic interactions.

Sample size and true positive rate

Increasing sample size naturally leads to a higher rate of correctly rejecting both the additive and

multiplicative null models (i.e., an increase in the true positive rate correctly detecting an

interaction) across all observation error levels tested (Figure 3). However, after an initial increase in

true positive rates by moving from N = 3 to approximately N = 15 replications, further improvement

is slow, and even for large sample sizes, it can be difficult to detect the 'true' interaction if

observation error is anything other than minimal. Moreover, there is no clear superiority in

performance for either model with the multiplicative model showing higher statistical power for all

sample sizes at the lowest observation errors (i.e., $\sigma = 0.001$ or $\sigma = 0.01$), whereas the additive null

model shows higher power for the highest level ($\sigma = 0.1$) of observation error.

Figure 3: Frequency of the 100,000 simulated interactions that are correctly classified (i.e., true

positive rate) for a range of different treatment replicates. Three different levels of observation error

are considered (0.001, 0.01, 0.1), with these being indicated by longdashed, dashed, and dotted lines

respectively. Yellow lines denote use of an additive null model, purple lines denote use of a

multiplicative null model.

Critical effect sizes

By combining and rearranging Equations 1.1 – 1.6, we can show that for a given sample size (N),

there is a minimum additive effect size value, also referred to as the critical effect size Mudge et al.

(2012), which must be exceeded before the interaction can be correctly classified as being non-null:

$$|ES_{Add}| > \sqrt{\left(\frac{1}{N_C} + \frac{1}{N_A} + \frac{1}{N_B} + \frac{1}{N_I}\right) \cdot \left(\frac{2 \cdot (Z_{\alpha/2})^2 \cdot (N_C + N_A + N_B + N_I) \cdot (J(m))^2}{2 \cdot (N_C + N_A + N_B + N_I) - (Z_{\alpha/2})^2 \cdot (J(m))^2}\right)}$$

326   *(Inequality 1)*

Unfortunately, the multiplicative null model is too complex to yield a solution. Figure 4 illustrates the

relationship between critical effect sizes and samples sizes for both the additive (Figure 4a) and

multiplicative (Figure 4b) null models using our simulated data. For the additive null model,

Inequality 1 is confirmed as the exact criterion for whether an effect size leads to a rejection of the

null model or not (Figure 4a), whereas the broken nature of the threshold for the multiplicative

model (Figure 4b) confirms its complex relationship with sample size. Importantly, both null models

show a rapid decline in critical effect size as sample sizes increases from N = 3 to approximately N =

25, and in both models the critical effect size more than halves in this range (albeit on different

scales).

As an aid to experimental design we provide an R code function (Supplementary Information) that

takes the choice of significance level (alpha) and the potential sample size (N) for all treatments and

returns the critical effect size ($|ES_{Add}|$) for the additive model. For example, setting alpha = 0.05 and

N = 3, leads to $|ES_{Add}|$ = 2.192, whereas increasing sample size to N = 16 leads to $|ES_{Add}|$= 0.982.

Only observed effect sizes greater than $|ES_{Add}|$ will be classified as non-additive. To judge how

useful an increase in N might be, we can see that in the freshwater stressor interactions analysed in

Burgess et al. (2021) approximately 20% of interactions are classified as non-null for N = 3, and this

rises to approximately 60% if N = 15 (Figure S4, Supplementary Information).

Figure 4: The relationship between treatment sample size and effect size for the additive (a) and

multiplicative (b) null models. For each treatment sample size (3-100), points are plotted from

simulations where an observation error of 0.01 was used. Pink points illustrate interactions that are

classified as null, while black points denote interactions classified as antagonistic, reversal, or

synergistic. The yellow line (Figure 4a) denotes the critical effect size threshold, computed from

Inequality 1, that interaction effect sizes must exceed in order for an interaction to be significantly

different to zero (and hence classified as being non-null) under the additive null model. As in the

methods, 95% confidence intervals were calculated (i.e., for Inequality 1, $Z_{\alpha/2}$ = 1.96).

Discussion

Our explicit aims were to better understand the statistical properties of the two most frequently used null models for multiple stressor interactions. Our analyses lead to three main conclusions that have important consequences for both experimental design and interpretation of results. Firstly, the additive and multiplicative null models make different assumptions that often lead to different interpretations about the stressor interactions, rendering direct comparisons between the model meaningless. Secondly, both models are very sensitive to uncertainty in the estimation of the stressor effects (i.e., the biological responses), and it can be very hard to reject the null interaction even with very small observation errors. Consequently, only interactions that very strongly deviate from the null expectation can be uncovered, meaning it is likely that many non-null interactions may have been missed in previous experiments. Finally, our results show how sample sizes need to be unrealistically large to uncover these 'hidden' synergies and antagonisms, but that increasing sample sizes from N = 3 to N = 25 yields the biggest increase in statistical power. Overall, we conclude that from an analytical perspective there is not an obviously *better* null model, and that model choice can therefore be based upon biological arguments without undue concerns over loss of statistical power.

The two null models we have focussed on are based on different assumptions, hence it should be no surprise that they lead to different interaction classifications, but it does highlight the importance of model choice. Differences in the results of null models are well-documented within fields such as pharmacology (**Tang et al., 2015**), where the class of observed interaction may be prefixed by the null model implemented (e.g., "Loewe antagonism", **Greco et al., 1992**), but it has been rarely addressed in multiple stressor literature (but see **Stephens et al. 2013** for a direct comparison on a single dataset). Currently, there is no equivalent framework for interpreting or reporting results for multiple stressor interactions. This discrepancy should be remedied given that the ambiguous usage of terms (such as synergistic or antagonistic interactions) are potentially hindering interpretation of

381 multiple stressor interactions both within, and across, disciplines (**Dey & Koops, 2021; Orr et al.,**

382 **2020**). Ultimately, we echo previous calls (**Schäfer & Piggott, 2018**) to thoroughly consider the

383 choice of null model to be implemented, and we argue that this decision needs to be based largely

384 on biological grounds since our results show there is little to choose between the null models when

385 it comes to statistical power. The decision should however be made very carefully, since different

386 choices may well lead to different types of interactions being assigned with the consequences of

387 very different management strategies being required to mitigate the stressor-pair effects. Choosing

388 the null model without reference to the relevance of the biological assumptions behind it could

389 therefore lead to remedial action that makes the situation worse.

391 The additive and multiplicative null models have many similarities regarding their robustness to data

392 uncertainty and their responses to sample size. In particular we show that, for likely realistic levels of

393 observation error, both models have poor statistical power (i.e., ability to correctly reject the null

394 model) at commonly used sample sizes of $3 \leq N \leq 8$. Several meta-analyses have highlighted a

395 dominance for null interactions (e.g., **Burgess et al., 2021; Lange et al., 2018; Stephens et al., 2013**),

396 and our results suggest this should be expected on purely statistical grounds. The important

397 corollary is that many non-null interactions are expected to be hidden from view, masked by data

398 uncertainty in the form of not only observation error as we model here, but also other forms such as

399 spatial, environmental, or genetic variation (**Higdon, 2013**).

401 'Will my study answer my research question?' is the most fundamental question a researcher can

402 ask when designing a study (**Johnson et al., 2015**), and its answer requires an understanding of the

403 statistical power of the tools at hand. Unfortunately, our results show that large sample sizes are

404 required to uncover even modest effect sizes (Figure 4). For example, a sample size of around N =

405 100 is required to detect an effect size of 0.5 as being non-additive (Inequality 1; Figure 4). As is the

case for interaction detection in epidemiological studies (**Greenland, 1983**), we find a slow

convergence of statistical power with sample size (Figure 4). Despite these unwelcome findings we

do observe a rapid initial increase in the minimum detectable effect size for both models (Inequality

1, Figure 4). This means increasing sample sizes from N = 3 up to around N = 25 can have important

consequences for uncovering non-null interactions. Indeed, we estimate that increasing sample sizes

to N = 15 would likely result in an increase of around 40% of non-null interactions in the dataset of

Burgess et al. (2021) (Supplementary Information). However, increasing sample sizes is not a trivial

consideration, and we recommend researchers consider what they believe constitutes a biologically

important effect size *before* the experiment is started. This so-called critical effect size (**Lakens,**

2021) is likely to change from study to study as the question, response metric, and taxonomic

group(s) change. Having determined the critical effect size means our results (Inequality 1, Figure 4,

and our R function *AddCritES* in the SI) can then be used to determine what sample size is required

to uncover it, whether this is even possible given budget and or practical constraints, or whether a

different approach might be necessary. However, we add that there is still high value in collecting

the data if it is then made available for future meta-analytical studies (**e.g., Jackson et al., 2016;**

Burgess et al., 2021) where the pooling of data overcomes the low power of small individual study

sample sizes.

A final consideration should be given to the choice of significance level. There has been much

discussion (e.g. **Mudge et al., 2012**) about how the significance level should be a function of the

sample size of the study, and not a fixed entity (e.g., the canonical level of 5%). For cases of low

power, Inequality 1 shows that a smaller alpha level (i.e., smaller value for Z) will lead to a smaller

critical effect size and therefore higher power to detect non-null stressor interactions. However,

raising the significance level comes at the cost of increased Type-1 errors (i.e., false rejection of the

null model). We argue that in multiple stressor research the cost of Type-1 errors is outweighed by

the increase in power because (i) we expect most interactions to be truly non-null even if only

weakly so (e.g. Table 1); (ii) most studies will not be able to have the large sample sizes required to

uncover even relatively strong effect sizes; and (iii) the ecological cost of failing to uncover the true

stressor interaction could be high. As shown by **Figure S2** (Supplementary Information), increasing

the significance level results in a modest increase in statistical power, across all levels of observation

error and number of replicates. Crucially, the altered level of significance must be justified, and not

arbitrarily chosen (**Lakens, 2021**). Such justification may be provided by either a cost-benefit analysis

(**see Field et al., 2004, Mudge et al., 2012**) or a compromise power analysis (**see Lakens, 2021**), and

extensions to our work could consider building a framework to encourage an informed choice of

significance level.

Finally, a word on the generality of our findings. The results based upon algebraic relationships (i.e.,

regarding critical effect size) are general and can be applied to any dataset, whereas results that

depend upon data simulation might be dependent upon the distribution of the effect sizes

generated by the underlying food chain model (e.g., Table 1; Figures 2, 3). Whilst the food chain

model used here has been shown to generate ecologically plausible results (**Burgess et al., 2021**),

the relationships revealed here may differ with other datasets (for instance, where a response other

than density is considered) which contain differing distributions of effect sizes. Ultimately, we

anticipate that many trends will remain across different datasets (e.g., the dominance of null

interactions at commonly implemented sample sizes), though others (e.g., the frequency of

antagonistic, reversal, and synergistic interactions in the absence of uncertainty) may vary between

datasets.

Conclusions

Robust conclusions about the prevalence of non-null interactions can only be reached once there is a

strong understanding of the abilities of the statistical tools to uncover them in the face of small

sample sizes and the underlying variation in the estimates of the biological responses. Our analyses

have confirmed that both additive and multiplicative null models have low power to detect non-null

interactions at commonly used sample sizes. However, we have provided guidance on the sample

sizes required to uncover specific effect sizes and we hope these results will be used to guide future

experimental designs, as well as lead to an increase in sample sizes beyond the typically small

numbers (<10). An important next step will be to investigate other null models (e.g. **Schäfer &**

Piggott, 2018; Tekin et al., 2020) and assess their abilities to uncover non-null interactions. Finally,

our work highlights the importance of null model choice and its effect on interaction classification,

and although we do not see any statistical advantage to either the additive or multiplicative null

model, deciding which model is biologically appropriate for the study question at hand is an

important choice that has consequences for potential downstream management decisions.

References

Borenstein, M., Cooper, H., Hedges, L. and Valentine, J., 2009. Effect sizes for continuous data.

The Handbook of Research Synthesis and Meta-Analysis, 2, pp.221-235.

Brook, B.W., Sodhi, N.S. and Bradshaw, C.J., 2008. Synergies among extinction drivers under

global change. *Trends in Ecology & Evolution*, 23(8), pp.453-460.

Brown, C.J., Saunders, M.I., Possingham, H.P. and Richardson, A.J., 2013. Managing for

interactions between local and global stressors of ecosystems. *PLoS One*, 8(6), p.e65765.

Burgess, B.J., Purves, D., Mace, G. and Murrell, D.J. (2021), Classifying ecosystem stressor

interactions: Theory highlights the data limitations of the additive null model and the

difficulty in revealing ecological surprises. *Global Change Biology*, 27, pp. 3052-3065.

Christensen, M.R., Graham, M.D., Vinebrooke, R.D., Findlay, D.L., Paterson, M.J. and Turner, M.A.,

2006. Multiple anthropogenic stressors cause ecological surprises in boreal lakes. *Global*

Change Biology, 12(12), pp.2316-2322.

Côté, I.M., Darling, E.S. and Brown, C.J., 2016. Interactions among ecosystem stressors and their

importance in conservation. *Proceedings of the Royal Society B: Biological Sciences*,

283(1824), p.20152592.

Crain, C.M., Kroeker, K. and Halpern, B.S., 2008. Interactive and cumulative effects of multiple

human stressors in marine systems. *Ecology Letters*, 11(12), pp.1304-1315.

Darling, E.S. and Côté, I.M., 2008. Quantifying the evidence for ecological synergies. *Ecology*

Letters, 11(12), pp.1278-1286.

De Laender, F., 2018. Community-and ecosystem-level effects of multiple environmental change

drivers: Beyond null model testing. *Global Change Biology*, 24(11), pp.5021-5030.

Dey, C.J. and Koops, M.A., 2021. The consequences of null model selection for predicting mortality from multiple stressors. *Proceedings of the Royal Society B*, 288(1948), p.20203126.

Dirzo, R., Young, H.S., Galetti, M., Ceballos, G., Isaac, N.J. and Collen, B., 2014. Defaunation in the Anthropocene. *Science*, 345(6195), pp.401-406.

Field, S.A., Tyre, A.J., Jonzén, N., Rhodes, J.R. and Possingham, H.P., 2004. Minimizing the cost of environmental management decisions by optimizing statistical thresholds. *Ecology Letters*, 7(8), pp.669-675.

Gomez Isaza, D.F., Cramp, R.L. and Franklin, C.E., 2020. Living in polluted waters: a meta-analysis of the effects of nitrate and interactions with other environmental stressors on freshwater taxa. *Environmental Pollution*, p.114091.

Greco, W., Unkelbach, H.D., Pöch, G., Sühnel, J., Kundi, M. and Bödeker, W., 1992. Consensus on concepts and terminology for combined-action assessment: the Saariselkä agreement. *Arch Complex Environ Stud*, 4(3), pp.65-9.

Greenland, S., 1983. Tests for interaction in epidemiologic studies: a review and a study of power. *Statistics in Medicine*, 2(2), pp.243-251.

Gurevitch, J., Morrison, J.A. and Hedges, L.V., 2000. The interaction between competition and predation: a meta-analysis of field experiments. *The American Naturalist*, 155(4), pp.435-453.

Hawkes, C.V. and Sullivan, J.J., 2001. The impact of herbivory on plants in different resource conditions: a meta-analysis. *Ecology*, 82(7), pp.2045-2058.

Hedges, L. V., and I. Olkin., 1985. Statistical methods for meta-analysis. *Academic Press*, New York.

Higdon R., 2013. Experimental Design, Variability. In: Dubitzky W., Wolkenhauer O., Cho KH.,

516   Yokota H. (eds) Encyclopaedia of Systems Biology. *Springer*, New York.

517   Hillebrand, H., Donohue, I., Harpole, W.S., Hodapp, D., Kucera, M., Lewandowska, A.M., Merder,

518   J., Montoya, J.M. and Freund, J.A., 2020. Thresholds for ecological responses to global

519   change do not emerge from empirical data. *Nature Ecology & Evolution*, pp.1-8.

520   Jackson, M.C., Loewen, C.J., Vinebrooke, R.D. and Chimimba, C.T., 2016. Net effects of multiple

521   stressors in freshwater ecosystems: a meta-analysis. *Global Change Biology*, 22(1),

522   pp.180-189.

523   Jackson, M.C., Fourie, H.E., Dalu, T., Woodford, D.J., Wasserman, R.J., Zengeya, T.A., Ellender,

524   B.R., Kimberg, P.K., Jordaan, M.S., Chimimba, C.T. and Weyl, O.L., 2020. Food web

525   properties vary with climate and land use in South African streams. *Functional Ecology*,

526   34(8), pp.1653-1665.

527   Johnson, P.C., Barry, S.J., Ferguson, H.M. and Müller, P., 2015. Power analysis for generalized

528   linear mixed models in ecology and evolution. *Methods in Ecology and Evolution*, 6(2),

529   pp.133-142.

530   Lajeunesse, M.J., 2011. On the meta-analysis of response ratios for studies with correlated and

531   multi-group designs. *Ecology*, 92(11), pp.2049-2055.

532   Lakens, D., 2021. Sample Size Justification. *PsyArXiv*. https://doi.org/10.31234/osf.io/9d3yf.

533   Lange, K., Bruder, A., Matthaei, C.D., Brodersen, J. and Paterson, R.A., 2018. Multiple-stressor

534   effects on freshwater fish: Importance of taxonomy and life stage. *Fish and Fisheries*,

535   19(6), pp.974-983.

536   Lindenmayer, D.B., Foster, C.N., Westgate, M.J., Scheele, B.C. and Blanchard, W., Managing

537   interacting disturbances: lessons from a case study in Australian forests. *Journal of*

538   *Applied Ecology*, 57, pp1711– 1716.

Mudge, J.F., Baker, L.F., Edge, C.B. and Houlahan, J.E., 2012. Setting an optimal α that minimizes
errors in null hypothesis significance tests. *PLoS One*, 7(2), p.e32734.

Nattrass, S., Baigent, S. & Murrell, D.J. (2012) Quantifying the Likelihood of Co-existence for
Communities with Asymmetric Competition. *Bulletin of Mathematical Biol*ogy 74,
pp2315–2338.

Newbold, T., Hudson, L.N., Hill, S.L., Contu, S., Lysenko, I., Senior, R.A., Börger, L., Bennett, D.J.,
Choimes, A., Collen, B. and Day, J., 2015. Global effects of land use on local terrestrial
biodiversity. *Nature*, 520(7545), pp45-50.

Orr, J.A., Vinebrooke, R.D., Jackson, M.C., Kroeker, K.J., Kordas, R.L., Mantyka-Pringle, C., Van den
Brink, P.J., De Laender, F., Stoks, R., Holmstrup, M. and Matthaei, C.D., 2020. Towards a
unified study of multiple stressors: divisions and common goals across research
disciplines. *Proceedings of the Royal Society B*, 287(1926), p.20200421.

Sala, E., Mayorga, J., Bradley, D., Cabral, R.B., Atwood, T.B., Auber, A., Cheung, W., Costello, C.,
Ferretti, F., Friedlander, A.M. and Gaines, S.D., 2021. Protecting the global ocean for
biodiversity, food and climate. *Nature*, pp.1-6.

Schäfer, R.B. and Piggott, J.J., 2018. Advancing understanding and prediction in multiple stressor
research through a mechanistic basis for null models. *Global Change Biology*, 24(5),
pp.1817-1826.

Seifert, M., Rost, B., Trimborn, S., Hauck, J., 2020. Meta-analysis of multiple driver effects on
marine phytoplankton highlights modulating role of pCO2. *Global Change Biology*, 26, pp.
6787– 6804.

Stephens, A.E., Srivastava, D.S. and Myers, J.H., 2013. Strength in numbers? Effects of multiple
natural enemy species on plant performance. *Proceedings of the Royal Society B:*
Biological Sciences, 280(1760), p.20122756.

Tang, J., Wennerberg, K. and Aittokallio, T., 2015. What is synergy? The Saariselkä agreement

revisited. *Frontiers in Pharmacology*, 6, p.181.

Tekin, E., Diamant, E.S., Cruz-Loya, M., Enriquez, V., Singh, N., Savage, V.M. and Yeh, P.J., 2020.

Using a newly introduced framework to measure ecological stressor interactions. *Ecology*

Letters, 23, pp1391-1403.

Tittensor, D.P., Walpole, M., Hill, S.L., Boyce, D.G., Britten, G.L., Burgess, N.D., Butchart, S.H.,

Leadley, P.W., Regan, E.C., Alkemade, R. and Baumung, R., 2014. A mid-term analysis of

progress toward international biodiversity targets. *Science*, 346(6206), pp.241-244.

Thompson, P.L., MacLennan, M.M. and Vinebrooke, R.D., 2018. An improved null model for

assessing the net effects of multiple stressors on communities. *Global Change Biology*,

24(1), pp.517-525.

Vinebrooke, R.D., Cottingham, K.L., Norberg, M., Scheffer, J., Dodson, S.I., Maberly, S.C. and

Sommer, U., 2004. Impacts of multiple stressors on biodiversity and ecosystem

functioning: The role of species co-tolerance. *Oikos*, 104(3), pp.451-457.

Wolfram Research. Inc., 2016. Mathematica (Version 10.4). *Wolfram Research, Inc., Champaign,*

Illinois.

Acknowledgments

This research was supported by the United Kingdom Natural Environment Research Council

studentship grant NE/M010481/1. The authors declare no conflicts of interest.

Author contributions

BB and DM conceived the ideas and designed methodology; BB generated the data; BB and DM

analysed the data; BB and DM led the writing of the manuscript. All authors contributed critically to

the drafts and gave final approval for publication.

Data availability

All data used in this work is publicly available via the supporting information of Burgess et al. (2021)

https://doi.org/10.1111/gcb.15630. Code for all results will be made available on Github.

590