Lowland plant migrations into alpine ecosystems amplify soil carbon

loss under climate warming

Tom W. N. Walker[1] [*], Konstantin Gavazov[2,3,4], Thomas Guillaume[3,4,5], Thibault

Lambert[6], Pierre Mariotte[3,4,7], Devin Routh[1], Constant Signarbieux[3,4],

Sebastián Block[1,8], Tamara Münkemüller[9], Hanna Nomoto[1], Thomas W.

Crowther[1], Andreas Richter[10], Alexandre Buttler[3,4] & Jake M. Alexander[1]

[1] Institute of Integrative Biology, ETH Zürich, 8092 Zürich, Switzerland.

[2] Climate Impacts Research Centre, Department of Ecology and Environmental Sciences, Umeå Universitet, 98107 Abisko,

Sweden

[3] Swiss Federal Institute for Forest, Snow and Landscape Research (WSL), 1015 Lausanne, Switzerland.

[4] École Polytechnique Fédérale de Lausanne EPFL, School of Architecture, Civil and Environmental Engineering ENAC,

Laboratory of Ecological Systems ECOS and Plant Ecology Research Laboratory PERL, 1015 Lausanne, Switzerland.

[5] Field-Crop Systems & Plant Nutrition, Agroscope, 1260 Nyon, Switzerland.

[6] Faculty of Geosciences & the Environment, Université de Lausanne, 1015 Lausanne, Switzerland.

[7] Grazing Systems, Agroscope, 1260 Nyon, Switzerland.

[8] Department of Ecology & Evolutionary Biology, Princeton University, 08544 Princeton, USA.

[9] Uni. Grenoble Alpes, CNRS, Uni. Savoie Mont Blanc, LECA, Laboratoire d'Ecologie Alpine, F-38000, Grenoble, France.

[10] Centre of Microbiology & Environmental Systems, Division of Terrestrial Ecosystem Research, University of Vienna,

1090 Vienna, Austria.

[11] International Institute for Applied Systems Analysis, Ecosystem Services and Management Program, 2361 Laxenburg,

Austria

* Correspondence to: thomas.walker@usys.ethz.ch.

Abstract

Climate warming is releasing carbon from soils around the world[1–3], constituting a positive climate

feedback. Warming is also causing species to expand their ranges into new ecosystems[4–9]. Yet, in

most ecosystems, whether range expanding species will amplify or buffer expected soil carbon loss

is unknown[10]. Here we used alpine grasslands as a model system to determine whether the

establishment of herbaceous lowland plants in alpine ecosystems influences short-term soil carbon

storage under warming. We found that warming (< 1 year) led to negligible alpine soil carbon loss,

but its effects became significant and 52% ± 31% (mean ± 95% CIs) larger after lowland plants were

introduced at low density into the ecosystem. We present evidence that soil carbon loss likely

occurred *via* lowland plants increasing rates of root exudation, soil microbial respiration and CO_2

release. Our findings suggest that warming-induced range expansions of herbaceous plants may

yield a rapid positive climate feedback in this system, and that plant range expansions among

herbaceous communities may be an overlooked mediator of warming effects on carbon dynamics.

Introduction

Climate warming is accelerating the release of soil carbon to the atmosphere[1–3], creating a positive climate feedback. At the same time, warming is driving uphill and poleward range expansions of native plant species around the globe[4–9]. Yet while warming and plant range expansions are now co-occurring in many terrestrial ecosystems, the impact of range expanding plants on soil carbon storage under warming remains poorly understood[10]. This is important because plant species adapted to different temperatures can have contrasting effects on carbon cycle processes[11], raising the potential for migrating species to have qualitatively different impacts on carbon dynamics from resident plants. For example, studies of non-native species invasions have revealed that the arrival of novel species in a community can alter carbon cycling in recipient ecosystems[12,13], while woody plant expansions into treeless Arctic ecosystems can reverse the direction of carbon cycle feedbacks to climate warming[14,15]. However, the majority of warming-induced range expansions do not involve non-native species or novel growth forms, but occur among native communities of similar species (e.g. ref [9]). In particular, we know remarkably little about how migrations of herbaceous plants into predominantly herbaceous ecosystems will affect soil carbon stocks, and so the magnitude and direction of resulting climate feedbacks are unknown.

Alpine ecosystems occur in the uppermost elevations of the planet as cool islands within warmer landscapes that are bounded by steep gradients in temperature[16]. They are experiencing both rapid temperature change[17] and faster-than-average species immigration[18], the latter of which mainly involves an influx of herbaceous warm-adapted species from lowlands into herbaceous cold-adapted communities at higher elevations[7–9]. Coupled to this, alpine ecosystems are globally relevant reservoirs of soil carbon and hold more than 90% of their total ecosystem carbon in the soil[1–3,19,20]. As such, alpine ecosystems are not only themselves important for climate feedbacks, but are also sentinels for climate warming impacts on other terrestrial ecosystems. Here, we used alpine ecosystems as a model system to determine how the establishment of range expanding native plant species affects soil carbon storage in warming ecosystems. We first used two whole-community transplant experiments in different alpine regions (western/central Swiss Alps) to establish the effects of simulated warming and lowland plant arrival on alpine soil carbon loss. Second, we used

67   these experiments plus glasshouse and laboratory studies to elucidate potential mechanisms

68   through which lowland plants influence the alpine soil carbon cycle, testing the hypothesis that they

69   act *via* changes to root exudation[21,22].

71
72
Figure 1 | Warming and lowland plant effects on alpine soil carbon loss. Mean ± SE soil carbon content (mg C g[-1] dry mass) in alpine

turfs exposed to warming (W; light grey), warming plus lowland plants (WL; dark grey) or an ambient control (C; white). Data are displayed

for two experiments in the western (left) and central (right) Alps, with letters indicating treatment differences (LMEs; N = 58).

Results & Discussion

<u>Lowland plants amplify warming-induced soil carbon loss</u>

We exploited two independent field experiments in which alpine plant communities plus rooting zone

soil were transplanted from high to low elevation to simulate warming, or back into a high elevation

site as a negative control (Supplementary Fig. S1). Half of the alpine communities at low elevation

were planted with seedlings of local lowland plant species to simulate the arrival of lowland plants in

the ecosystem. Remaining communities at low elevation and all control communities at high

elevation were instead subjected to a planting disturbance control (see Methods for a full account of

control treatments and validation exercises). We found that warming of less than one year caused a

small decline in alpine soil carbon content, but in both regions this effect was stronger and only

became statistically significant when lowland plants were also present (Fig. 1; Table S1; $P < 0.001$).

Indeed, considering field data from both experiments together, we estimate that the arrival of lowland

plants in the ecosystem amplified warming-induced soil carbon loss by 52% ± 31% (mean ± 95%

confidence intervals) per °C relative to warming alone (Supplementary Fig. S3; P = 0.001). These

findings collectively show that once lowland plants establish in alpine communities they can amplify

warming-induced soil carbon loss. We thus suggest that uphill migrations of lowland plants, which

are already globally common[8,18], have a hitherto unrecognised bearing over alpine soil carbon

storage and could ultimately yield a positive climate feedback.

<u>Lowland and alpine plants have distinct effects on soil microbial respiration</u>

Soil carbon loss occurs due to the activity of soil microorganisms, which break down soil organic

matter and release a portion as CO_2 through respiration[23]. We therefore hypothesised that the

observed impact of lowland plants on alpine soil carbon loss was mediated *via* lowland plant effects

on soil microbial respiration. To test this, we performed a glasshouse experiment in which we grew

lowland plants (*Bromus erectus*, *Salvia pratensis*, *Medicago lupulina*) or alpine plants (*Phleum*

alpinum, *Anthyllis vulneraria* subsp. *alpestris*, *Plantago alpina*) in alpine soil collected from the

western Alps region. We found that lowland plants accelerated soil microbial respiration relative to

alpine plants (Fig. 2a; treatment × time: LR = 147.98, P < 0.001). Accelerated microbial activity was

also reflected in spectral analyses of soil pore water, which showed that soil under lowland plants

contained more dissolved organic matter (Fig. 2d,e; Table S2; a_{350}: P < 0.001, F_{tot}: P < 0.001) with a

higher proportion of microbially-derived products (Fig. 2f; Table S2; P = 0.002) than soil under alpine

plants. Importantly, microbial respiration was enhanced despite no changes to microbial biomass

between treatments (Fig. 2g; Table S2; P = 0.863), showing that microbes respired intrinsically faster

(that is, per unit of biomass) in the presence of lowland plants. These findings together imply that

lowland plants accelerate alpine soil carbon loss by stimulating the activity of alpine soil microbes

and the associated release of CO_2 through respiration.

We posit that lowland plants accelerated microbial activity by increasing the quantity of root exudates

released into the soil, for four reasons. First, two-pool models of microbial respiration revealed that

soils supporting any plant contained a soil carbon pool that was effectively absent from a bare soil

control, and which was 1.5 ± 0.7 (mean ± SE) times larger under lowland plants than alpine plants

(Fig. 2b; Table S2; P = 0.046). This carbon pool decayed at a similar rate between treatments (Fig.

2c; Table S2; P = 0.602), indicating that it differed in size, but likely not composition, between lowland

and alpine treatments. Second, the same carbon pool was consumed rapidly by soil microbes (half-

life = 5.1 ± 0.4 days), and we also found that lowland plant effects on microbial respiration were lost

after nine days of incubation without plants (Fig. 2a). Plants thus supplied compounds to microbes

that were depleted within this timeframe. A nine-day decay period corresponds to the time expected

for newly photosynthesised CO_2 to be released through root exudation and respired by soil

microbes[21,22], suggesting that this carbon pool was mostly root exudates. Third, lowland plant effects

on the alpine soil system were rapid, occurring within a period of weeks in the glasshouse experiment

and months in the field experiments. It is unlikely that lowland plants influenced microbial activity *via*

litter inputs within this timeframe[24,25]. Finally, dissolved organic matter contained a higher relative

abundance of protein-like substances than humic-like and fulvic acid-like substances

(Supplementary Fig. S5; Table S2; P < 0.001 in all cases) under lowland plants than alpine plants,

also suggesting that lowland plants caused an enrichment in exudate-derived compounds[26].

Together, these findings suggest that lowland plants induce soil carbon loss from alpine ecosystems

by increasing root exudation relative to resident alpine plants, which accelerates soil microbial

activity and the associated degradation and release of soil carbon as CO_2[27,28].

Plants adapted to different temperatures can possess distinct traits that have opposing effects on

carbon cycle processes[11]. We thus investigated whether lowland plants acted on the alpine soil

system by introducing novel traits into the ecosystem. Publicly available trait data[29] relating to all

species in the field experiments revealed that lowland plants on average possessed more productive

traits than alpine plants (Supplementary Fig. S2a; P = 0.003). This was also the case in the

glasshouse experiment, whereby trait measurements from all plants showed that lowland plants

displayed more productive traits overall (Supplementary Fig. S2b; P = 0.001). Coupled to this, while

lowland plants were planted at low abundance in the field experiments (mean ± SE relative cover:

4.7% ± 0.7%), the magnitude of soil carbon loss caused by lowland plant presence was greater in

plots where lowland plants were more abundant overall (Supplementary Fig. S2e; $F_{1,15}$ = 5.64, P =

0.031). By contrast, we found no evidence that soil carbon loss in the field was alternatively related

to individual lowland species identity (Supplementary Fig. S2c,d) or the composition of the recipient

alpine plant community (Supplementary Fig. S2,g).

Taken together, these findings confirm that lowland plants are functionally distinct from alpine

plants. As a result, our results suggest that their establishment leads to a shift in plant community

functioning that alters carbon cycle processes, with effects that intensify as lowland plants become

more abundant in the community[24,25]. However, it is also possible that lowland plants acted on the

alpine soil system simply by introducing more biomass into the ecosystem. We investigated this

possibility by summarising traits from the glasshouse experiment at the pot scale using a principal

components analysis and testing for relationships between resulting axes of trait variation and

alpine soil microbial respiration (Methods). We found that descriptors of plant biomass represented

the most important axis of trait variation (Supplementary Fig. 4c) and were positively related to

microbial respiration (Supplementary Fig. 4a; PC1: LR = 4.03, N = 20, P = 0.045). However, this

relationship did not differ between lowland and alpine treatments (treatment × PC1: LR = 0.26, N =

20, P = 0.608), ruling it out as the mechanism through which lowland plants uniquely alter the

alpine soil system. Instead, we found that trait variation represented by the second principal axis

(for example, specific leaf area; Supplementary Fig. 4d) had opposing effects on microbial

respiration between lowland and alpine treatments (Supplementary Fig. 4b; treatment × PC2: LR =

5.54, N = 20, P = 0.019). Lowland plants thus displayed relationships with the alpine soil system

that were qualitatively distinct from resident alpine plants. While focussed work is now needed to

pinpoint the specific physiological mechanism behind these differences, the existence of such a

distinction reveals that the arrival of lowland plants in alpine ecosystems has the capacity to initiate

a step-change in plant community functioning, with potential consequences for carbon cycle

processes[24,25].

Fig. 2 | Lowland plants and alpine soil carbon cycling. (a-g) Effects of lowland (L) or alpine (A) plants on alpine soil in a glasshouse experiment (mean ± SE; grey points/ribbons: SEs for bare soil). (a) Total microbial respiration (R; µg C g^{-1} h^{-1}) during follow-up incubations without plants (N = 260). Vertical line indicates time before which A-L comparison differed (LME, Tukey). (b) Size (µg C g^{-1}) and (c) decay rate (µg C g^{-1} d^{-1}) of a carbon pool identified using two-pool models (N = 16). (d) Absorbance (a$_{350}$; m^{-1}), (e) total fluorescence (F$_{tot}$; Raman Units) and (f) fluorescence index (FI) of soil pore water (N = 20). (g) Microbial biomass carbon (C$_{mic}$; mg C g^{-1} dry mass) (N = 18). (h-m) Effects of field warming (W), warming plus lowland plants (WL) or an ambient control (C) on alpine soil (means ± SEs) after one season. (h) Total ecosystem respiration (R$_e$; g CO_2 m^{-2} h^{-1}) averaged over multiple dates in the western and central Alps field experiments (N = 196, N = 90). (i) Total microbial respiration (µg C g^{-1} h^{-1}), (j) total microbial growth (µg C g^{-1} h^{-1}), (k) C$_{mic}$ (µg C g^{-1}), (l) microbial carbon use efficiency (CUE; %), (m) biomass-specific microbial respiration (R$_{mass}$; ng C µg^{-1} C$_{mic}$ h^{-1}) and (n) biomass-specific microbial growth (G$_{mass}$; ng C µg^{-1} C$_{mic}$ h^{-1}) in the western Alps experiment (N = 28). Treatment effects (P < 0.05) are indicated by asterisks (b-g; LMEs) or letters (h-n; LMEs, Tukey). Fills correspond to pools in a conceptual model (o) proposing a timeline for how lowland (L) versus alpine (A) plants impact alpine soil carbon loss.

 <u>Lowland plant-induced carbon loss is pulsed</u>

Soil microbial processes are constrained by multiple feedbacks within the soil system, such as

substrate availability[30,31] and microbial acclimation[32–34], that over time can slow, and eventually

arrest, amplified soil carbon loss[31,33,34]. We thus interrogated the state of the soil system in the field

experiments after one season to explore the possible duration of lowland plant effects on alpine soil

carbon loss. We focussed particularly on intrinsic rates (that is, per unit of biomass) of microbial

respiration and growth as processes that promote soil carbon loss and gain, respectively[28,35], as well

as microbial biomass as the scalar through which these processes affect soil carbon[30,31]. We found

that biomass-specific rates of microbial respiration in the field increased under warming in the

presence, but not absence, of lowland plants (Fig. 2m; Table S1; P = 0.038). By contrast, biomass-

specific rates of microbial growth were slower under warming but remained unaffected by lowland

plant presence (Fig. 2n; Table S1; P = 0.016). These findings support observations from the

glasshouse experiment that lowland plants accelerate intrinsic rates of microbial respiration (Fig.

2a,g), while also confirming that this persists for at least one season of field exposure. Moreover, by

stimulating respiration, but not growth, lowland plants increased the capacity for alpine soils to lose

carbon, but not to gain carbon *via* accumulation into microbial biomass[28,35].

Faster microbial respiration can only yield a loss of soil carbon if it occurs on a per unit of soil

basis[31,34,36], which was the case in the glasshouse experiment (Fig. 2g). However, in the field we

found that lowland plants caused a reduction of microbial biomass carbon in warmed plots (Fig. 2k;

Table S1; P < 0.001). As such, although microbes always respired intrinsically faster in the presence

of lowland plants, in the field they had become less abundant and so we observed no change to

microbial respiration when expressed per unit of soil (Fig. 2i; Table S1; P = 0.200). Depressed

microbial respiration per unit of soil was also evident at the ecosystem scale, in that warming

accelerated total ecosystem respiration (that is, plant plus soil respiration) but its effect was

dampened in plots containing lowland plants (Fig. 2h; Table S1; P < 0.001). The decline in microbial

biomass likely took place because soil microbes are chiefly carbon limited, making their abundance

tightly coupled to soil carbon content (here: r_{26} = 0.61, P = 0.001)[31]. Moreover, since intrinsic rates

of microbial growth were unaffected by lowland plants, a reduction in microbial biomass translated

into a decline in microbial growth per unit of soil (Fig. 2j; Table S1; P = 0.002) – preventing any

recovery of microbial biomass *via* growth[35]. We thus suggest that as soil carbon was lost, microbial

biomass also declined, and at the time of measurement this had entirely eliminated the influence of

accelerated biomass-specific microbial respiration over the soil system. In short, the soil carbon loss

observed in warmed field plots containing lowland plants must have both started and stopped within

one season.

Our findings collectively suggest that the establishment of lowland plants in warming alpine

ecosystems accelerates intrinsic rates of microbial respiration (Fig. 2a), which leads to a pulse of

soil carbon release at baseline levels of microbial biomass (Fig. 1, Fig. 2g). However, resulting soil

carbon loss leads to a decline in microbial biomass (Fig. 2k), a decrease in microbial respiration per

unit of soil (Fig. 2i) and a cessation of further carbon loss from the ecosystem within a one-year

timeframe (Fig. 2o)[31]. Here, we simulated a single event of lowland plant establishment, and at

relatively low abundance in the community (mean ± SE relative cover: 4.7% ± 0.7%). Future work

should thus address whether increases in lowland plant cover or repeated establishment events

could induce additional pulses of alpine soil carbon loss. It is important to note that this interpretation

does not consider the possibility that soil microbes acclimate to the presence of lowland plants, which

could also yield a deceleration in microbial processes over time[32–34]. Nevertheless, it remains true

that biomass-specific microbial growth and respiration affect the soil system as a function of their

biomass, and here changes to microbial biomass were sufficient to explain the behaviour of the soil

system in both glasshouse and field experiments. We also found no evidence that soil microbes

instead changed their resource use strategy in the presence versus absence of lowland plants (that

is, carbon use efficiency; Fig. 2l; Table S1; P < 0.001; Tukey: P = 0.251), pointing away from

microbial acclimation as an alternative mechanism.

Conclusions

In summary, we show that uphill migrations of native lowland plants can rapidly amplify the direct

effects of climate warming on soil carbon loss from alpine ecosystems. This is important because it

demonstrates that plant range expansions between herbaceous plant communities have the

potential to yield a positive feedback to climate warming, at least in the short term. These results

contrast with expectations from woody species expansions in the Arctic, where shrubs with

recalcitrant litter can slow carbon turnover in herbaceous communities and potentially yield a

negative climate feedback[14,15]. We thus suggest that the influence of range expanding species on

ecosystem carbon dynamics depends on the relative trait composition of migrating species versus

the recipient community[37], as well as subsequent effects on environmental processes[14]. While our

findings demonstrate that lowland plants affect the rate of soil carbon loss in the short term, short-

term experiments, such as ours, cannot resolve whether lowland plants will also affect the total

amount of soil carbon lost in the long term. However, here soil carbon loss occurred despite a low

abundance of lowland plants, and increased with total lowland plant cover. Given that both warming

and associated plant migrations will intensify over time[8,9], our estimates of lowland plant impacts

may be conservative – particularly if increases in lowland plant cover or repeated establishment

events cause recurrent pulses of soil carbon loss. Moreover, while alpine ecosystems store only a

small fraction of total global soil carbon[1,2], redistributions among herbaceous plant communities are

occurring globally[4–9], including in similarly cool ecosystems in the Arctic that contain vast soil carbon

stocks[38]. This raises the potential for resulting climate feedbacks to also be widespread. Future work

should focus on testing the conditions under which this feedback could occur in other ecosystems

experiencing influxes of range expanding plant species, on quantifying how deeply it occurs in

shallow alpine soils, and on estimating the magnitude of the climate feedback given both ongoing

warming and variation in rates of species range shifts.

Materials & Methods

Field site description, experimental design. We made use of two independent alpine grassland transplant experiments that were established in Switzerland in the western Alps (46°12′40″N, 7°03′13″E)[39] and central Alps (46°52′20″N, 9°28′58″E) in autumn 2016. The experiments examine effects of climate warming on alpine ecosystems, including the arrival of lowland plants due to warming-induced uphill plant migrations (Supplementary Fig. S1). In the western Alps experiment the plant community is dominated by *Sesleria caerulea*, *Carex sempervirens* and *Vaccinium vitis-idea*. In the central Alps experiment the dominant species are *Nardus stricta*, *Alchemilla xanthochlora*, *Trifolium pratense* and *Anthyllis vulneraria*. Broadly, the experiments share the same design. Turfs (area = 1 m^2, depth = 20 cm) of intact alpine vegetation plus rhizosphere soil were removed from a high elevation control site (western Alps: 2200 m, MAT = 0.7 ºC, TAP = 1448 mm; central Alps: 2000 m, MAT = 1.6 ºC, TAP = 1151 mm; ref. [40]) and replanted either at a low elevation site to simulate climate warming (western Alps: 1400 m, MAT = 6.0 ºC, TAP = 1236 mm, N = 10; central Alps: 1400 m, MAT = 4.4 ºC, TAP = 1010 mm, N = 20; ref. [40]) or back into a high elevation site as a transplantation control (see below). At the low elevation site, we created an additional treatment to simulate the arrival of lowland plants under warming by planting a low abundance of lowland plant species in a grid design (13 cm between individuals), creating a lowland plant density of approximately 56 individuals per m^2 (4.7% ± 0.7% total cover) in a background alpine plant community. Lowland plants were positioned randomly in the grid, but with the same number of individuals per species per plot, and we controlled for planting disturbance in all remaining plots using alpine plants in place of lowland plants (see below). In the western Alps region, lowland plant species were *Achillea millefolium, Bellis perennis, Bromus erectus, Dactylis glomerata, Medicago lupulina, Plantago media* and *Salvia pratensis*, and individuals were planted into one side of each plot in a split-plot design (n = 10). In the central Alps experiment, lowland plant species were *Brachypodium pinnatum, Carex flacca, Carum carvi, Dactylis glomerata, Hypericum perforatum, Plantago lanceolata*, *Primula veris*, *Ranunculus bulbosus*, *Salvia pratensis*, *Silene vulgaris*, *Trifolium montanum* and *Viola hirta*, and individuals were planted into half of the transplanted turfs (n = 10). In the western Alps experiment, lowland individuals were pre-grown in a greenhouse in steam-sterilized field soil and planted into the plots in May 2017. In the central Alps experiment, lowland

individuals were taken from the surrounding vegetation and planted into the plots in autumn 2016.

In both regions, soil was partially removed from lowland individuals prior to planting, and we

performed an additional experiment to eliminate the possibility that small contaminations of lowland

soil biota during planting could have modified alpine soil microbial community composition (see

Supplementary Methods). Overall, the experimental design in both regions yielded three treatments

(n = 10 for each) arranged in a block design: warming only (W); warming plus low elevation plants

(WL); and a high elevation control (C). One turf at the low elevation site in the western Alps

experiment failed, leaving a total sample size of N = 28 and N = 30 in the western Alps and central

Alps, respectively.

Field experiment controls and validation. We controlled for transplantation disturbance in both

experiments by moving alpine control turfs back into a high elevation control site. In the western Alps

experiment, alpine turfs were planted into different positions at their origin high elevation site. In the

central Alps experiment, alpine turfs were planted into a new high elevation site at the same

elevation. As such, we eliminated the possibility that alpine turfs changed simply because they were

disturbed or were moved to a different location (that is, irrespective of elevation). The transplantation

process had no effect on alpine soil carbon content, in that we found no difference in soil carbon

content between replanted alpine control turfs at high elevation sites and undisturbed alpine soil in

the surrounding area (LR = 2.27, N = 16, P = 0.132). We also controlled for potential effects of

planting disturbance in all turfs not planted with lowland plants (that is, remaining turfs at low

elevation and all control turfs at high elevation). In the western Alps experiment, we planted alpine

species from the high elevation control site into the other side of alpine turfs at the low elevation site,

as well as into all control turfs at the high elevation site. This was done in autumn 2016 using the

same approach and randomized hexagonal grid design. In the central Alps experiment, we mimicked

planting disturbance by digging up and replanting existing individuals in alpine turfs at the low

elevation site not containing lowland plants, as well as all control turfs at the high elevation site. This

was done in spring 2017 using the same grid design. It is important to note that we did not perform

a reverse transplantation (that is, from low to high elevation), so we cannot entirely rule out the

possibility that transplantation of any community to any new environment could yield a loss of soil

carbon. Nevertheless, it is well established that climate warming promotes soil carbon loss[27,41,42], and we found that soil from low elevation sites (that is, the potential end-member for warming alpine ecosystems) contained less carbon than soil from the alpine sites (70.1 ± 3.05 mg C g^{-1} soil dry mass compared to 145.0 ± 5.29 mg C g^{-1} soil dry mass; LR = 37.2, N = 27, P < 0.001), validating our expectation that alpine soil would lose carbon under warming. Elevation-based transplant experiments are a powerful approach for assessing warming effects on ecosystems[11,43], especially when they include rigorous disturbance controls (see above) and are performed in multiple locations where the common change from high to low elevation is temperature – here warming of between 2.8 ºC (central alps) and 5.3 ºC (western alps). While small variations in experimental design between regions, such in the order of planting in W and WL treatments, likely introduced some error when comparing experiments, we used both regions together so that we could identify treatment effects that were consistent among them and were thus not artefacts of experimental design.

Field plant community composition. We surveyed the composition of alpine communities in summer 2017. This was done by dividing each plot into 5 × 5 cm cells and visually estimating the canopy cover of all species rooted in every grid cell. Species cover was approximated with an ordinal scale in which the first three cover categories correspond to 1/16, 1/8, and 1/4 of the grid cell area (25 cm^2), and subsequent cover categories were constant increases of 1/4 of the grid cell area. Plants were identified to species level whenever possible, although some taxa were identified only to genus or subfamily levels when species identity was unclear based on vegetative traits. Species cover (cm^2) was then calculated as the species-wise sum of cover estimates across all grid cells in the quarter of the plot used for soil and ecosystem respiration measurements (see below).

Field soil carbon content, microbial physiology, microbial biomass. We collected soil samples from all treatments in both field experiments (N = 58) in August 2017. In each plot, we created a composite sample from three cores (ø = 1 cm, approx. d = 7 cm) no closer than 7 cm from a planted individual and from the same quarter of the plot used for ecosystem respiration measurements (see below; Supplementary Fig. S1). We focused on rhizosphere soil for two reasons. First, alpine soils are shallow and alpine plants are shallow-rooting, making the rhizosphere the most logical target for

a first assessment of warming and lowland plant effects on alpine soil carbon cycling. Second, we

wanted to avoid any potential contamination from the base of the turf (depth = 20 cm). Soils were

kept cool, transported immediately to the laboratory, sieved (2 mm mesh size), adjusted to 60% of

water holding capacity (WHC) and pre-incubated for three days at field temperature prior to

measurements. We measured microbial biomass carbon (mg per g dry mass) *via* chloroform

fumigation extraction followed by analysis on a TOC-VCPH/CPNTNM-1 analyzer (Shimadzu; 48h

incubation period; 1 M KCl extraction for fumigated and non-fumigated samples; conversion factor

0.45)[44]. While this approach also yielded measurements of dissolved organic carbon (DOC; mg C

per g dry mass), treatment effects on this variable echoed those of microbial biomass carbon (Table

S1) so we did not interpret them separately. Soil organic carbon content (%) was measured on dry

soil (60 ºC for 48h) with a Carlo Erba 1110 elemental analyzer (EA, CE Instruments) coupled to a

Delta Plus IRMS via a Conflo III (Thermo Fisher). We additionally quantified microbial growth,

respiration and carbon use efficiency in soils from the western Alps experiment (N = 28) using the

incorporation of ^{18}O into DNA[31]. Briefly, we incubated 500 mg of soil for 24h at field temperature

following addition of either ^{18}O-H_2O to 20 atom% enrichment and 80% of WHC or the same volume

of molecular grade non-labelled H_2O as a natural abundance control. Gas samples were taken at

the start and end of the 24h incubation and analyzed for CO_2 concentration with a Trace GC Ultra

(Thermo Fisher) to determine microbial respiration (µg C per g dry mass per h). At the end of the

incubation, soil was snap-frozen in liquid nitrogen, and DNA was extracted (FastDNA SPIN Kit for

Soil, MP Biomedical), quantified (Quant-iT PicoGreen dsDNA Assay Kit, Thermo Fisher) and

analyzed for ^{18}O abundance and total oxygen content using a thermochemical EA coupled to a Delta

V Advantage IRMS via a Conflo III (Thermo Fisher). Following ref. [31], we calculated total DNA

production during the 24h incubation (µg DNA per gram dry mass) and coupled it with measures of

microbial biomass carbon content to derive microbial growth (µg C per gram dry mass per h) and

carbon use efficiency (%).

Field ecosystem respiration. We measured rates of ecosystem respiration (g CO_2 per m^2 per h)

on multiple dates (western Alps: 7 dates; central Alps: 3 dates) between July and August 2017. One

month prior to the first measurement, we installed permanent gas sampling collars (*h* = 10 cm, ø =

30 cm) in each plot to a depth of 5 cm to create an airtight seal at the soil surface[11]. Measurements

were made by enclosing plants and soil within the collars with an opaque chamber (h = 50 cm, ø =

30 cm) and, following a 30s equilibration period, monitoring the increase in CO_2 concentration at 5s

intervals over a 60s period (Vaisala GMP343 CO_2 probe attached to an HM70 meter, Vaisala).

Chambers contained a fan to mix the headspace during measurements and a small sealable hole

on the top that was opened when placing the chamber on sampling collars to avoid pressurization

inside the chamber. Soil temperature and moisture were recorded for each measurement and used

alongside measurements of air temperature, air pressure, enclosure volume and soil surface area

to calculate the CO_2 flux[11]. Where necessary, we repeated measurements in the field to attain a

visually consistent CO_2 change over time, and during data processing we discarded fluxes with an

r-squared of less than 0.9 for this relationship.

Glasshouse experiment design. We established a six-week pot experiment to test for effects of

lowland plants on the alpine soil system. Specifically, we filled pots (ø = 10 cm, depth = 9 cm) with

sieved soil (mesh size 4 mm) from the high elevation site of the western Alps field experiment, and

planted them with lowland species (*Bromus erectus*, *Salvia pratensis*, *Medicago lupulina*) or alpine

species (*Phleum alpinum*, *Anthyllis vulneraria* ssp. *alpestris*, *Plantago alpina*) to create a lowland

only treatment, an alpine only treatment and a bare soil control (N = 30). We selected a grass, herb

and legume species for each treatment, all of which were found naturally at the low (lowland

treatment) or high (alpine treatment) elevation sites in the western Alps field experiment. Vegetated

treatments contained two individuals of each species planted in a circular pattern such they were

never adjacent to conspecifics or plants of the same functional type (i.e. grass, herb, legume). The

original design also included a mixed community treatment containing one individual of each lowland

and alpine species. However, the mixed treatment had double the species richness of other

vegetated treatments and contained an equal cover of lowland and alpine plants (compared to < 5%

lowland plant cover in field plots), so we focused our analyses on the comparison between alpine

and lowland plant treatments to understand how lowland plants affect the soil differently from alpine

plants. All plants were pre-grown from seed provided by commercial suppliers (www.hauenstein.ch,

www.schutzfilisur.ch, www.saatbau.at) in steam-sterilized potting soil for two to three weeks to

approximately the same biomass and root-washed immediately prior to planting. Pots were arranged

in a randomized block design (n = 10) in a greenhouse under standardized growing conditions (22

ºC; 50% humidity; no artificial light), and were watered every two to three days as needed. After six

weeks, we destructively harvested the plants and soil to characterize treatment effects on plant

physiology and the soil system.

Glasshouse plant physiology. We measured maximum leaf-level photosynthesis (photosynthetic

capacity; A_{max}; µmol CO_2 per m^2 per s) and stomatal conductance (g_s; µmol H_2O per m^2 per s) under

optimum conditions on one individual per species per pot using an open path infrared gas analyzer

connected to a 2.5 cm chamber (CIRAS-2 and PLC-6, PP Systems)[11]. Readings were taken once

for each individual and corrected using measurements of leaf area (cm^2) on the same leaf, which we

also dried and weighed to determine specific leaf area (SLA; cm^2 per gram). Following soil pore

water sampling (below), we quantified total, aboveground and belowground biomass of all individuals

in the experiment by destructively sampling, washing, drying (60 ºC for 48h) and weighing all plant

tissue.

Glasshouse soil pore water analysis. We collected soil pore water from all pots prior to soil

sampling (see below) to determine treatment effects on dissolved organic matter (DOM) quality. Soil

water was sampled by leaching pots twice with 50 mL sterile tap water, combining and filtering

leached water (filter pore size 0.2 µm) and measuring absorbance (200-900 nm with a 1nm

increment; 1 cm quartz cuvette; UV-vis spectrophotometer, Shimadzu) and fluorescence intensity

(excitation: 240-450 nm with a 5 nm increment; emission: 290-550 nm with a 2.5 nm increment;

Fluorolog-3 spectrometer, Horiba). We corrected absorbance spectra against a milli-Q water blank,

calculated Napierian absorption coefficients and used these data to derive absorbance at 350 nm

(a_{350}; a proxy for DOM content). We expressed fluorescence data as total fluorescence (F_{tot}; Raman

Units) and the fluorescence index (FI), i.e. the ratio of emission intensity at 470 nm versus 520 nm

at an excitation wavelength of 370 nm (a proxy for microbially-derived DOM[45]). We also derived

excitation-emission matrices (EEMs) to quantify the relative abundance of different components of

DOM through PARAFAC algorithms[46]. Following preprocessing (i.e. Raman scattering removal and

standardization to Raman Units) and normalization[47], we used the drEEM toolbox[48] in MATLAB (Mathworks) to obtain a six-component PARAFAC model. The model was validated with split-half analysis, random initialization and visualization of EEM residuals, after which we calculated the relative abundance of each component as the maximum fluorescence signal of the component against the total maximum fluorescence signal of all components. Finally, we classified components against classical EEM nomenclature (Table S4; refs [49,50]). While PARAFAC modelling provides a means to discriminate between DOM sources (e.g. components C3 and C4; Table S4), we did not physically separate plant- and microbe-derived DOM when sampling soil pore water. As such, observed changes to DOM composition were likely a result of changes to both plant and microbial processes (see Main Text).

Glasshouse soil carbon pools, microbial respiration. We took soil cores (ø = 1 cm, d = 7 cm) from the center of each pot (i.e. equidistant from all individuals) to quantify soil carbon pools and microbial respiration. Soils were sieved (2 mm mesh size), adjusted to 60% of their WHC and pre-incubated at 22 ºC for three days prior to measurements. We quantified concentrations of microbial biomass carbon in all soils as for field soil (see above; mg C per g dry mass). We determined microbial respiration by incubating 8 g sieved soil in the dark at 22 ºC for a further six-week period without plants. We measured microbial respiration daily for the first four days, then approximately biweekly until the end of the six-week period. We measured CO_2 concentrations (EGM-4, PP Systems) of gas samples taken approximately 24h after flushing enclosed vials (v = 126 mL) with CO_2-free air, and calculated microbial respiration (µg C per g dry mass per h) using accompanying measurements of enclosure volume, air pressure and soil mass[11]. After each measurement, soils were weighed and, when necessary, re-adjusted to 60% of their WHC.

Two-pool mixed models of soil carbon mineralization. We described the kinetics of soil carbon mineralization during the follow-up incubation of glasshouse soil by employing a two-pool mixed model to microbial respiration data[51]. Specifically, we constructed a model that considered a first pool that followed first-order decomposition kinetics (i.e. a small soil carbon pool with a fast turnover)

and a second pool that followed zero-order kinetics (i.e. decay rate was constant over the incubation

period, representing a large soil carbon pool with a slow turnover):

$$C_{min} = C_l(1 - e^{-tk_l}) + B \cdot t$$

where C_{min} is the cumulative CO_2 respired (µg C g^{-1}) over the incubation time t (d), C_l and k_l are the

size (µg C g^{-1}) and decomposition constant (d^{-1}) of the first-order kinetics carbon pool, respectively,

and B is the basal respiration rate (µg C g^{-1} d^{-1}) of the zero-order kinetics carbon pool. Given that

C_{min} and t were known, we ran the two-pool mixed model with the nlsList function (maxiter = 100, tol

= 1 × 10^{-5}, minFactor = 1/1024) in R[52] using the package "nlme"[53] for each pot separately to estimate

values for C_l, k_l and B. Models failed to converge for one sample, and we excluded four additional

samples for which models made negative pool estimates (N = 25).

Statistical analysis. The majority of statistical test outputs appear as *P*-values in the Main Text,

with full outputs being provided in Tables S1 to S3. See text below for exceptions. Data analyses

and visualization were undertaken in R[52] using the packages "cowplot"[54], "drake"[55], "emmeans"[56],

"nlme"[53], "tidyverse"[57] and "vegan"[58]. Where necessary, region was accounted for in statistical

models as a fixed effect, since two levels was too few to be included as a random intercept term. As

such, region effects are presented (Tables S1, S3), but not interpreted. Statistical methods are

described as they appear in the Main Text.

Warming and warming plus lowland plant effects on soil carbon content: we used field data to test

for treatment (C, W, WL) effects on soil carbon content using an LME model including region as a

fixed effect and block as a random intercept term (see Table S1 for test output). We also calculated

how much more soil carbon loss occurred in the presence versus absence of lowland plants by fitting

a single LME testing for effects of soil temperature (plot means of all sampling dates), the presence

or absence of lowland plants and their interaction on soil carbon content (see Supplementary Fig.

S2 for test outputs). In order to compare slopes between C-W and C-WL treatment combinations,

we duplicated C treatment values, assigned one set to each C-W and C-WL pair and accounted for

this using a random intercept term. We also included region as a random slope term. We then tested

how much more soil carbon loss would occur per ºC of warming when lowland plants are present by

calculating the relative difference (% ± 95% confidence intervals) between the C-WL and C-W

slopes.

Plant mechanisms for soil carbon loss: we expressed soil carbon content as the block-wise percent

changes between WL and W treatments to derive a metric for lowland plant induced soil carbon loss

(i.e. % soil carbon remaining). We tested for lowland species identity effects on soil carbon remaining

using separate LMEs for each region (due to different lowland species pools) that tested for individual

species cover (cm^2) effects on soil carbon remaining in WL plots, including species as a random

intercept term (see Supplementary Fig. S3 for test outputs). We also performed separate linear

models including each species as a fixed effect, none of which were statistically significant (Table

S3). We tested for effects of total lowland species cover (cm^2) on soil carbon remaining in WL plots

using a linear model also including region as a fixed effect (treatment × region: $F_{1,15} = 0.48$, $P =$

0.498; region: 0.60, $P = 0.500$). We summarized alpine community composition using an NMDS

(Hellinger distance matrix, two axes), following which we expressed NMDS scores as the block-wise

percent changes between W and WL treatments to derive metrics for lowland plant induced alpine

plant community change, and used a linear model that tested for effects of changes to NMDS scores

on soil carbon remaining in WL plots including region as a fixed effect (see Supplementary Fig. S3

for test outputs). For reference, raw NMDS scores did not significantly differ between W and WL

treatments (LME: NMDS #1: LR = 1.54, $P = 0.214$; NMDS #2: LR = 2.78, $P = 0.095$). Finally, we

tested for differences between alpine and lowland species traits in the field experiment using a

PERMANOVA on a PCA of species mean trait values (log_{10} plant height, log_{10} leaf area, log_{10} seed

mass, SLA, leaf carbon, leaf nitrogen) from the TRY database[29] for all species present (where

necessary, missing values were gap-filled following[59]), and in the glasshouse experiment using a

PERMANOVA on a PCA of physiological measurements (total biomass, leaf biomass, root biomass,

SLA, A_{max}, g_s) for all plants (see Supplementary Fig. S3 for test outputs). Traits were selected to

explore broad life history differences between alpine and lowland plant species[60], but were not used

to identify the specific trait(s) responsible for lowland plant effects on alpine soil carbon loss. We

used TRY traits for field species due to the large number of species present in the communities (N

= 242), but used measured traits in the glasshouse experiment to provide direct evidence of

physiological differences between alpine and lowland species.

<u>Lowland versus alpine plant effects on the soil system:</u> we used the glasshouse experiment to test

for lowland versus alpine plant effects on soil microbial biomass carbon, the size (C_l) and decay rate

(k_l) of the first modelled soil carbon pool and all variables relating to soil DOM quality using LMEs

including block as a random intercept term. We tested for effects of treatment, time and their

interaction on soil microbial respiration using the same LME structure but including the bare soil

treatment. In all cases, including/excluding the bare soil treatment had no effect over model

outcomes, but mean ± SE values for this treatment are included in figures for reference. Finally, we

explored how plant physiology in lowland and alpine treatments affected initial rates of soil microbial

respiration. We defined initial microbial respiration as the pot-wise mean respiration rates of the first

nine days of soil incubation (that is, while treatment effects persisted; see Fig. 2a), and expressed

pot-level plant physiology using scores from the first two axes of a principal components analysis

performed on sums of total, root and shoot biomass and biomass-weighted means of SLA, A_{max} and

g_s. We then tested for effects of treatment, PC1 and PC2 scores and their interaction on initial

microbial respiration using an LME including block as a random intercept term. Statistical test outputs

for LMEs are provided in Table S2, except for treatment and plant physiology effects on microbial

respiration, which appear in the Main Text.

<u>Sequence of warming and lowland plant effects on alpine soil carbon loss:</u> we used field data to test

for treatment (C, W, WL) effects on soil microbial biomass carbon and soil DOC using LME models

including region as a fixed effect and block as a random intercept term. We tested for treatment

effects on ecosystem respiration using an LME that included region as a fixed effect and sampling

date nested in block as a random intercept term. We tested for treatment effects on soil microbial

growth and respiration (per unit of soil and per unit of microbial biomass) and carbon use efficiency

in the western Alps experiment using LMEs including block as a random intercept term. Statistical

test outputs for these variables are provided in Table S1. We also used Pearson Product Moment

correlations to test for associations between soil microbial biomass and soil carbon content,

microbial respiration and growth in the western Alps experiment, irrespective of experimental

treatment.

Acknowledgements

JMA and HN are funded by the European Research Council (grant no. 678841). JMA and TWNW

are funded by the Swiss National Science Foundation (grant no. 31003A-176044). KG is funded by

the Swiss National Science Foundation (grant no. PZ00P2-174047). SB and the Central Alps

experiment are funded by the Swiss National Science Foundation (grant no. 31003A-173210).

Author contributions

TWNW, AB and JMA conceived the study. AR, AB and JMA provided resources for field and

laboratory experiments. SB, TM and HN established and maintained field experiments. TWNW,

KG and TG undertook the field work. TWNW, KG, TG, PM, CS, AB and JMA undertook the

glasshouse experiment. TWNW, KG, TG, PM, CS and AR did the laboratory work. TL performed

the soil pore water sampling and associated analyses. SB and TM performed the soil inoculation

experiment and associated analyses. TWNW did the data analysis, with support from DR, TWC

and JMA. TWNW wrote the manuscript in close collaboration with JMA, and with support from all

co-authors.

Competing interests

The authors declare no competing financial interests.

References

1. Streit, K. *et al.* Soil warming alters microbial substrate use in alpine soils. *Glob. Change Biol.* **20,** 1327–1338

(2014).

2. Parker, T. C., Subke, J.-A. & Wookey, P. A. Rapid carbon turnover beneath shrub and tree vegetation is associated

with low soil carbon stocks at a subarctic treeline. *Glob. Change Biol.* **21,** 2070–2081 (2015).

3. Knowles, J. F., Blanken, P. D., Lawrence, C. R. & Williams, M. W. Evidence for non-steady-state carbon emissions

from snow-scoured alpine tundra. *Nat. Commun.* **10,** 1306 (2019).

4.   Parmesan, C. Ecological and evolutionary responses to recent climate change. *Annu. Rev. Ecol. Evol. Syst.* **37,**

637–669 (2006).

5.   Lenoir, J. & Svenning, J. C. Climate-related range shifts - a global multidimensional synthesis and new research

directions. *Ecography* **38,** 15–28 (2015).

6.   Pecl, G. T. *et al.* Biodiversity redistribution under climate change: Impacts on ecosystems and human well-being.

Science **355,** (2017).

7.   Dolezal, J. *et al.* Vegetation dynamics at the upper elevational limit of vascular plants in Himalaya. *Sci. Rep.* **6,**

24881 (2016).

8.   Rumpf, S. B. *et al.* Range dynamics of mountain plants decrease with elevation. *Proc. Natl. Acad. Sci. USA* **115,**

1848–1853 (2018).

9.   Steinbauer, M. J. *et al.* Accelerated increase in plant species richness on mountain summits is linked to warming.

Nature **556,** 231–234 (2018).

10.   Wallingford, P. D. *et al.* Adjusting the lens of invasion biology to focus on the impacts of climate-driven range shifts.

Nat. Clim. Chang. (2020). doi:10.1038/s41558-020-0768-2

11.   Walker, T. W. N. *et al.* Plastic and genetic responses of a common sedge to warming have contrasting effects on

carbon cycle processes. *Ecol. Lett.* **22,** 159–169 (2019).

12.   Liao, C. *et al.* Altered ecosystem carbon and nitrogen cycles by plant invasion: a meta-analysis. *New Phytol.* **177,**

706–714 (2008).

13.   Waller, L. P. *et al.* Biotic interactions drive ecosystem responses to exotic plant invaders. *Science* **368,** 967–972

(2020).

14.   Pearson, R. G. *et al.* Shifts in Arctic vegetation and associated feedbacks under climate change. *Nat. Clim. Chang.*

3, 673–677 (2013).

15.   Cornelissen, J. H. C. *et al.* Global negative vegetation feedback to climate warming responses of leaf litter

decomposition rates in cold biomes. *Ecol. Lett.* **10,** 619–627 (2007).

16.   Körner, C. *Alpine Plant Life: Functional Ecology of High Mountain Ecosystems*. (Springer, 2003).

17.   Pepin, N. *et al.* Elevation-dependent warming in mountain regions of the world. *Nat. Clim. Chang.* **5,** 424–430

(2015).

18.   Lenoir, J., Gégout, J. C., Marquet, P. A., de Ruffray, P. & Brisse, H. A significant upward shift in plant species

optimum elevation during the 20th century. *Science* **320,** 1768–1771 (2008).

19.   Hagedorn, F., Gavazov, K. & Alexander, J. M. Above- and belowground linkages shape responses of mountain

vegetation to climate change. *Science* **365,** 1119–1123 (2019).

20.   Budge, K., Leifeld, J., Hiltbrunner, E. & Fuhrer, J. Alpine grassland soils contain large proportion of labile carbon

but indicate long turnover times. *Biogeosciences* **8,** 1911–1923 (2011).

21.   Hafner, S. *et al.* Effect of grazing on carbon stocks and assimilate partitioning in a Tibetan montane pasture

revealed by 13CO2 pulse labeling. *Glob. Change Biol.* **18,** 528–538 (2012).

22. Studer, M. S., Siegwolf, R. T. W. & Abiven, S. Carbon transfer, partitioning and residence time in the plant-soil

system: a comparison of two[13] CO_2 labelling techniques. *Biogeosciences* **11,** 1637–1648 (2014).

23. Plante, A. F., Stone, M. M. & McGill, W. B. in *Soil microbiology, ecology and biochemistry* 245–272 (Elsevier,

2015). doi:10.1016/B978-0-12-415955-6.00009-8

24. De Deyn, G. B., Cornelissen, J. H. C. & Bardgett, R. D. Plant functional traits and soil carbon sequestration in

contrasting biomes. *Ecol. Lett.* **11,** 516–531 (2008).

25. Cornwell, W. K. *et al.* Plant species traits are the predominant control on litter decomposition rates within biomes

worldwide. *Ecol. Lett.* **11,** 1065–1071 (2008).

26. Jones, D. L., Nguyen, C. & Finlay, R. D. Carbon flow in the rhizosphere: carbon trading at the soil–root interface.

Plant Soil **321,** 5–33 (2009).

27. Melillo, J. M. *et al.* Soil warming and carbon-cycle feedbacks to the climate system. *Science* **298,** 2173–2176

(2002).

28. Allison, S. D., Wallenstein, M. D. & Bradford, M. A. Soil-carbon response to warming dependent on microbial

physiology. *Nat. Geosci.* **3,** 336–340 (2010).

29. Kattge, J. *et al.* TRY plant trait database - enhanced coverage and open access. *Glob. Change Biol.* **26,** 119–188

(2020).

30. Hartley, I. P., Hopkins, D. W., Garnett, M. H., Sommerkorn, M. & Wookey, P. A. Soil microbial respiration in arctic

soil does not acclimate to temperature. *Ecol. Lett.* **11,** 1092–1100 (2008).

31. Walker, T. W. N. *et al.* Microbial temperature sensitivity and biomass change explain soil carbon loss with warming.

Nat. Clim. Chang. **8,** 885–889 (2018).

32. Crowther, T. W. & Bradford, M. A. Thermal acclimation in widespread heterotrophic soil microbes. *Ecol. Lett.* **16,**

469–477 (2013).

33. Bradford, M. A. *et al.* Cross-biome patterns in soil microbial respiration predictable from evolutionary theory on

thermal adaptation. *Nat. Ecol. Evol.* **3,** 223–231 (2019).

34. Melillo, J. M. *et al.* Long-term pattern and magnitude of soil carbon feedback to the climate system in a warming

world. *Science* **358,** 101–105 (2017).

35. Liang, C., Schimel, J. P. & Jastrow, J. D. The importance of anabolism in microbial control over soil carbon storage.

Nat. Microbiol. **2,** 17105 (2017).

36. Prommer, J. *et al.* Increased microbial growth, biomass, and turnover drive soil organic carbon accumulation at

higher plant diversity. *Glob. Change Biol.* **26,** 669–681 (2020).

37. Bjorkman, A. D. *et al.* Plant functional trait change across a warming tundra biome. *Nature* **562,** 57–62 (2018).

38. Tarnocai, C. *et al.* Soil organic carbon pools in the northern circumpolar permafrost region. *Global Biogeochem.*

Cycles **23,** (2009).

39. Nomoto, H. A. & Alexander, J. M. Drivers of local extinction risk in alpine plants under warming climate. *Ecol. Lett.*

24, 1157–1166 (2021).

40. Karger, D. N. *et al.* Climatologies at high resolution for the earth's land surface areas. *Sci. Data* **4,** 170122 (2017).

41. Crowther, T. W. *et al.* Quantifying global soil carbon losses in response to warming. *Nature* **540,** 104–108 (2016).

42. Davidson, E. A. & Janssens, I. A. Temperature sensitivity of soil carbon decomposition and feedbacks to climate

change. *Nature* **440,** 165–173 (2006).

43. Alexander, J. M., Diez, J. M. & Levine, J. M. Novel competitors shape species' responses to climate change.

Nature **525,** 515–518 (2015).

44. Vance, E. D., Brookes, P. C. & Jenkinson, D. S. An extraction method for measuring soil microbial biomass C. *Soil*

Biol. Biochem. **19,** 703–707 (1987).

45. McKnight, D. M. *et al.* Spectrofluorometric characterization of dissolved organic matter for indication of precursor

organic material and aromaticity. *Limnol. Oceanogr.* **46,** 38–48 (2001).

46. Stedmon, C. A., Markager, S. & Bro, R. Tracing dissolved organic matter in aquatic environments using a new

approach to fluorescence spectroscopy. *Mar Chem* **82,** 239–254 (2003).

47. Zepp, R. G., Sheldon, W. M. & Moran, M. A. Dissolved organic fluorophores in southeastern US coastal waters:

correction method for eliminating Rayleigh and Raman scattering peaks in excitation–emission matrices. *Mar Chem*

89, 15–36 (2004).

48. Murphy, K. R., Stedmon, C. A., Graeber, D. & Bro, R. Fluorescence spectroscopy and multi-way techniques.

PARAFAC. *Anal. Methods* **5,** 6557 (2013).

49. Fellman, J. B., Hood, E. & Spencer, R. G. M. Fluorescence spectroscopy opens new windows into dissolved

organic matter dynamics in freshwater ecosystems: A review. *Limnol. Oceanogr.* **55,** 2452–2462 (2010).

50. *Aquatic organic matter fluorescence.* (Cambridge University Press, 2014). doi:10.1017/CBO9781139045452

51. Bonde, T. A. & Rosswall, T. Seasonal variation of potentially mineralizable nitrogen in four cropping systems. *Soil*

Sci. Soc. Am. J. **51,** 1508–1514 (1987).

52. R Core Team. *R: A language and environment for statistical computing.* (R Foundation for Statistical Computing,

2014).

53. Pinheiro, J., Bates, D., DebRoy, S. & Sarkar, D. *nlme: Linear and Nonlinear Mixed Effects Models.* (R Project,

2019).

54. Wilke, C. O. *cowplot: Streamlined Plot Theme and Plot Annotations for "ggplot2."* (R Project, 2019).

55. Michael Landau, W. The drake R package: a pipeline toolkit for reproducibility and high-performance computing.

JOSS **3,** 550 (2018).

56. Lenth, R. *Estimated Marginal Means, aka Least-Squares Means.* (R Project, 2020).

57. Wickham, H. *et al.* Welcome to the tidyverse. *JOSS* (2017). doi:10.21105/joss.01686

58. Oksanen, J. *et al. vegan: Community Ecology Package.* (R Project, 2019).

59. Schrodt, F. *et al.* BHPMF - a hierarchical Bayesian approach to gap-filling and trait prediction for macroecology and

functional biogeography. *Glob. Ecol. Biogeogr.* **24,** 1510–1521 (2015).

60. Díaz, S. *et al.* The global spectrum of plant form and function. *Nature* **529,** 167–171 (2016).