Lowland plant arrival in alpine ecosystems facilitates a decrease in soil

carbon content under experimental climate warming

Tom W. N. Walker[1] *, Konstantin Gavazov[2,3,4], Thomas Guillaume[3,4,5], Thibault

Lambert[6], Pierre Mariotte[3,4,7], Devin Routh[1], Constant Signarbieux[3,4],

Sebastián Block[1,8], Tamara Münkemüller[9], Hanna Nomoto[1], Thomas W.

Crowther[1], Andreas Richter[10], Alexandre Buttler[3,4] & Jake M. Alexander[1]

[1] Institute of Integrative Biology, ETH Zürich, 8092 Zürich, Switzerland.

[2] Climate Impacts Research Centre, Department of Ecology and Environmental Sciences, Umeå Universitet, 98107

Abisko, Sweden

[3] Swiss Federal Institute for Forest, Snow and Landscape Research (WSL), 1015 Lausanne, Switzerland.

[4] École Polytechnique Fédérale de Lausanne EPFL, School of Architecture, Civil and Environmental Engineering ENAC,

Laboratory of Ecological Systems ECOS and Plant Ecology Research Laboratory PERL, 1015 Lausanne, Switzerland.

[5] Field-Crop Systems & Plant Nutrition, Agroscope, 1260 Nyon, Switzerland.

[6] Faculty of Geosciences & the Environment, Université de Lausanne, 1015 Lausanne, Switzerland.

[7] Grazing Systems, Agroscope, 1725 Posieux, Switzerland.

[8] Department of Ecology & Evolutionary Biology, Princeton University, 08544 Princeton, USA.

[9] Uni. Grenoble Alpes, CNRS, Uni. Savoie Mont Blanc, LECA, Laboratoire d'Ecologie Alpine, F-38000, Grenoble,

France.

[10] Centre of Microbiology & Environmental Systems, Division of Terrestrial Ecosystem Research, University of Vienna,

1090 Vienna, Austria.

[11] International Institute for Applied Systems Analysis, Ecosystem Services and Management Program, 2361 Laxenburg,

Austria

* Correspondence to: thomas.walker@usys.ethz.ch.

Abstract

Climate warming is releasing carbon from soils around the world[1–3], constituting a positive climate feedback. Warming is also causing species to expand their ranges into new ecosystems[4–9]. Yet, in most ecosystems, whether range expanding species will amplify or buffer expected soil carbon loss is unknown[10]. Here we used two whole-community transplant experiments and a follow-up glasshouse experiment to determine whether the establishment of herbaceous lowland plants in alpine ecosystems influences soil carbon content under warming. We found that warming (transplantation to low elevation) led to a negligible decrease in alpine soil carbon content, but its effects became significant and 52% ± 31% (mean ± 95% CIs) larger after lowland plants were introduced at low density into the ecosystem. We present evidence that decreases in soil carbon content likely occurred *via* lowland plants increasing rates of root exudation, soil microbial respiration and CO_2 release under warming. Our findings suggest that warming-induced range expansions of herbaceous plants have the potential to alter climate feedbacks from this system, and that plant range expansions among herbaceous communities may be an overlooked mediator of warming effects on carbon dynamics.

Introduction

Climate warming is expected to accelerate the release of soil carbon to the atmosphere[11]. At the same time, warming is driving uphill and poleward range expansions of native plant species around the globe[4–9]. Yet while warming and plant range expansions are now co-occurring in many terrestrial ecosystems, the impact of range expanding plants on soil carbon storage under warming remains poorly understood[10]. This is important because plant species adapted to different temperatures can have contrasting effects on carbon cycle processes[12], raising the potential for migrating species to have qualitatively different impacts on carbon dynamics from resident plants. For example, studies of non-native species invasions have revealed that the arrival of novel species in a community can alter carbon cycling in recipient ecosystems[13,14], while woody plant expansions into treeless Arctic ecosystems can reverse the direction of carbon cycle feedbacks to climate warming[15,16]. However, the majority of warming-induced range expansions do not involve non-native species or novel growth forms, but occur among native communities of similar species (e.g. ref [9]). In particular, we know remarkably little about how migrations of herbaceous plants into predominantly herbaceous ecosystems will affect carbon cycling, and so the magnitude and direction of resulting climate feedbacks are unknown.

Alpine ecosystems occur in the uppermost elevations of the planet as cool islands within warmer landscapes that are bounded by steep gradients in temperature[17]. They are experiencing both rapid temperature change (0.4 to 0.6 °C per decade)[18,19] and rapid species immigration[20], the latter of which involves an influx of herbaceous warm-adapted species from lowlands into herbaceous cold-adapted communities at higher elevations[7–9]. Coupled to this, alpine ecosystems are globally relevant reservoirs of soil carbon and hold more than 90% of their total ecosystem carbon in the soil[1–3,21,22]. As such, alpine ecosystems are not only themselves important for climate feedbacks, but are also sentinels for climate warming impacts on other terrestrial ecosystems. Here, we used alpine ecosystems as a model system to determine how the establishment of range expanding native plant species could affect soil carbon storage in warming ecosystems. We first used two whole-community transplant experiments in different alpine regions (western/central Swiss Alps) to establish the effects of warming plus lowland plant arrival on alpine soil carbon content. Second,

we used these experiments plus glasshouse and laboratory studies to explore potential

mechanisms through which lowland plants influence the alpine soil carbon cycle, testing the

hypothesis that they act *via* changes to root exudation[23,24].

Figure 1 | Warming and lowland plant effects on alpine soil carbon content in the field experiment. Mean ± SE soil carbon

content (mg C g^{-1} dry mass; i.e. mass-based per-mil) in alpine turfs transplanted to low elevation (warming, W; light grey), transplanted

plus planted with lowland plants (warming plus lowland plant arrival, WL; dark grey) or replanted at high elevation (control, C; white).

Data are displayed for two experiments in the western (left) and central (right) Alps, with letters indicating treatment differences (LMEs;

N = 58).

Results & Discussion

Lowland plants facilitate warming-induced soil carbon loss

We exploited two independent field-based transplant experiments in which alpine plant

communities plus rooting zone soil were transplanted from high to low elevation to simulate a

climate warming scenario, or back into a high elevation site as a negative control (Supplementary

Fig. S1; Methods). A portion of the alpine turfs at low elevation were planted with local lowland

plant species to simulate the arrival of warm-adapted lowland plants in the warmed ecosystem,

with the remaining portion being subjected to a planting disturbance control. Elevation-based

transplant experiments are powerful tools for assessing climate warming effects on ecosystems

because they expose plots to a real-world future temperature regime with natural diurnal and

seasonal cycles while also warming both aboveground and belowground subsystems[25–27]. This is

especially true if they include rigorous disturbance controls (here, see Methods) and are performed in multiple locations where the common change from high to low elevation is temperature (here, warming of 2.8 °C in the central Alps and 5.3 °C in the western Alps). While factors other than temperature can co-vary with elevation[17], such factors either do not vary consistently with elevation among experiments (e.g. precipitation, wind), are not expected to strongly influence plant performance (e.g. UV radiation) or in any case form part of a realistic climate warming scenario (e.g. growing-season length, snow cover)[17,27,28]. In short, the experiments used here examined how the arrival of warm-adapted lowland plants influences alpine ecosystems in a warmed climate matching lowland site conditions (i.e. turf transplantation to low elevation plus lowland plant addition) relative to warming-only (i.e. turf transplantation to low elevation) or control (i.e. turf transplantation within high elevation) scenarios.

We found that transplantation from high to low elevation (i.e. warming-only) caused a small decline in alpine soil carbon content, but in both regions this effect was stronger and only became statistically significant when lowland plants were also planted-in (Fig. 1; Table S1; $P < 0.001$). Specifically, considering field data from both experiments together, the addition of lowland plants to transplanted alpine turfs increased warming-induced soil carbon loss from 4.1 ± 0.23 mg g^{-1} per °C to 6.3 mg g^{-1} per °C, which is a $52\% \pm 31\%$ (mean ± 95% confidence intervals) increase relative to warming alone (Supplementary Fig. S3; $P = 0.001$). We caution against equating changes to soil carbon content with changes to soil carbon stock in the absence of coupled measurements of soil bulk density (Methods). Nevertheless, these findings show that once warm-adapted lowland plants establish in warming alpine communities, they facilitate warming effects on soil carbon loss on a per gram basis. We thus suggest that uphill migrations of lowland plants, which are already globally common[8,20], may have a hitherto unrecognised bearing over alpine soil carbon storage.

<u>Lowland and alpine plants have distinct effects on soil microbial respiration</u>

Soil carbon loss occurs due to a shifting balance between soil carbon inputs *via* net primary production and soil carbon outputs *via* soil microbial respiration[29,30]. We therefore hypothesised that lowland plants impacted alpine soil carbon loss by either decreasing net primary production or

increasing soil microbial respiration relative to alpine plants. To test this, we performed a

glasshouse experiment in which we grew lowland plants (*Bromus erectus*, *Salvia pratensis*,

Medicago lupulina) or alpine plants (*Phleum alpinum*, *Anthyllis vulneraria* subsp. *alpestris*,

Plantago alpina) in alpine soil collected from the western Alps region. The glasshouse experiment

was not a direct simulation of field conditions, in that plants from lowland and alpine communities

were grown separately at a constant temperature and humidity (Methods). Instead, the glasshouse

experiment allowed us to isolate how plants adapted to lowland versus alpine climates differentially

affect the alpine soil system.

134

Fig. 2 | Differences between lowland and alpine plant traits in the glasshouse experiment. Mean ± SE pot-level traits for alpine (A)

and lowland (L) plant treatments in a glasshouse experiment. (a) Aboveground biomass (AGB; g; N = 19). (b) Belowground biomass

(BGB; g; N = 10). (c) Root-to-shoot ratio (N = 19). (d-f) Biomass-weighted (d) specific leaf area (SLA; cm^2 g^{-1}; N = 18), (e) maximum

photosynthetic capacity (A_{max}; μmol m^2 s^{-1}; N = 19) and (f) maximum stomatal conductance (g_s; μmol m^2 s^{-1}; N = 20). Treatment effects

($P < 0.05$) are indicated by asterisks (LMEs; see Table S2).

We first compared lowland and alpine plant traits to test whether lowland plants decreased carbon

inputs into alpine soil. We found that lowland plant treatments produced more aboveground (Fig.

2a; Table S2; P = 0.028) and belowground (Fig. 2b; Table S2; P < 0.001) biomass than alpine

plants, while also containing a higher proportion of root tissue (root-to-shoot ratio: Fig. 2c; Table

S2; P = 0.005). While biomass is an imperfect proxy for net primary production, these findings

point away from the hypothesis that lowland plants cause alpine soil carbon loss by reducing

carbon inputs into soil. On the contrary, the higher biomass and proportionately more root tissue

present in lowland plant treatments suggests that lowland plants may in fact increase soil carbon

inputs. This occurred despite lowland plant treatments possessing similar specific leaf area (SLA:

Fig. 2d; Table S2; $P = 0.861$) to alpine plant treatments, as well as lower maximal photosynthetic

capacity (A_{max}: Fig. 2e; Table S2; $P = 0.026$) and stomatal conductance (g_s: Fig. 2f; Table S2; $P <$

0.001). While the reason for decreased photosynthetic capacity and stomatal conductance in the

lowland plant treatment is unclear, the fact that both variables were lower suggests that

photosynthetic capacity was limited by stomata pore size rather than by photosynthetic

machinery[31]. Indeed, lowland plants accumulated more biomass despite having a lower maximal

photosynthetic capacity, ruling it out as a limiting factor of lowland plant net primary production.

Fig. 3 | Lowland and alpine plant effects on alpine soil microbes in the glasshouse experiment. Effects of alpine (A) versus

lowland (L) plant treatments on alpine soil microbes in a glasshouse experiment (mean ± SE; grey points/ribbons: SEs for bare soil). (a)

Total microbial respiration (R; µg C g^{-1} dry mass h^{-1}) during follow-up incubations without plants (N = 260). (b) Size (µg C g^{-1} dry mass)

and (c) decay rate (µg C g^{-1} dry mass d^{-1}) of a carbon pool identified using two-pool models (N = 16). (d) Microbial biomass carbon (C_{mic};

mg C g^{-1} dry mass; N = 18). For (a), vertical line indicates time before which A-L comparison differed (LME, Tukey; see Main Text). For

(b-d), treatment effects ($P < 0.05$) are indicated by asterisks (LMEs; see Table S2).

We next examined the soil system under lowland versus alpine plants to determine whether lowland plants alternatively increased carbon outputs from alpine soil. We found that lowland plants accelerated soil microbial respiration relative to alpine plants (Fig. 3a; treatment × time: likelihood ratio (LR) = 147.99, $P < 0.001$). Accelerated microbial activity was also reflected in spectral analyses of soil pore water, which showed that soil under lowland plants contained more dissolved organic matter (Fig. 4a,b; Table S2; absorbance (a_{350}): $P < 0.001$, total fluorescence (F_{tot}): $P < 0.001$) with a higher proportion of microbially-derived products (Fig. 4c; Table S2; FI; $P = 0.002$) than soil under alpine plants. Importantly, microbial respiration was enhanced despite no changes to microbial biomass between treatments (Fig. 3d; Table S2; $P = 0.693$), showing that microbes respired intrinsically faster (i.e., per unit of biomass) in the presence of lowland plants. These findings support the hypothesis that lowland plants have the capacity to increase soil carbon outputs relative to alpine plants by stimulating soil microbial respiration and associated CO_2 release. While accelerated microbial respiration can alternatively be a signal of soil carbon accumulation *via* greater microbial growth[30], such a mechanism is unlikely to have been responsible here because it would have led to an increase in microbial biomass carbon under lowland plants, which we did not observe.

Fig. 4 | Lowland and alpine plants effects on alpine soil dissolved organic matter in the glasshouse experiment. Effects of alpine (A) versus lowland (L) plant treatments on alpine soil dissolved organic matter extracted from soil pore water in a glasshouse experiment (mean ± SE; grey ribbons: SEs for bare soil). (a) Absorbance (a_{350}; m^{-1}), (b) total fluorescence (F_{tot}; Raman Units) and (c) fluorescence index (FI) (N = 20). (d-i) Relative abundance (%) of six dissolved organic matter components (C1-C6), where components and associated compound families were derived from PARAFAC modelling (Methods; Table S4). Treatment effects (P < 0.05) are indicated by asterisks (LMEs; see Table S2).

We speculate that lowland plants accelerated microbial activity in the glasshouse experiment by increasing the quantity of root exudates released into the soil, for four reasons. First, two-pool models of microbial respiration revealed that soils supporting any plant contained a soil carbon pool that was effectively absent from a bare soil control, and which was a factor of 1.5 ± 0.7 (mean ± SE) times larger under lowland plants than alpine plants (Fig. 3b; Table S2; *P* = 0.046). This carbon pool decayed at a similar rate between treatments (Fig. 3c; Table S2; *P* = 0.631), indicating that it differed in size, but likely not composition, between lowland and alpine treatments. Second, the same carbon pool was consumed rapidly by soil microbes (half-life = 5.1 ± 0.4 days), and we

also found that lowland plant effects on microbial respiration were lost after nine days of incubation without plants (Fig. 3a). Plants thus supplied compounds to microbes that were depleted within this timeframe. While not directly measured here, a nine-day decay period corresponds to the time expected for newly photosynthesised CO_2 to be released through root exudation and respired by soil microbes[23,24], suggesting that this carbon pool was mostly root exudates. Third, lowland plant effects on the alpine soil system were rapid, occurring within a period of weeks in the glasshouse experiment and months in the field experiments. It is unlikely that lowland plants influenced microbial activity *via* litter inputs within this timeframe[32,33]. Finally, dissolved organic matter contained a higher relative abundance of protein-like substances than humic-like and fulvic acid-like substances (Fig. 4d-i; Table S2; $P < 0.01$ in all cases) under lowland plants than alpine plants, also suggesting that lowland plants caused an enrichment in exudate-derived compounds[34]. While further directed studies are required to resolve whether root exudates are truly involved, our findings collectively suggest that lowland plants have the capacity to increase total root exudation into alpine soil relative to resident alpine plants. In turn, root exudation is a well-established regulator of soil microbial decomposition and the associated release of soil carbon as CO_2[29,35].

One explanation for increased soil carbon outputs under lowland plants is that lowland plants have more biomass than alpine plants and therefore stimulate microbial activity to a greater extent (Fig. 2a-c). We investigated this possibility by summarising pot-level traits from the glasshouse experiment using a principal components analysis and testing for relationships between resulting axes of trait variation and alpine soil microbial respiration (Methods). We found that descriptors of plant biomass represented the most important axis of trait variation (Supplementary Fig. S4c) and were positively related to microbial respiration (Supplementary Fig. S4a; PC1: LR = 4.43, N = 20, P = 0.035). Thus, greater biomass of lowland plants may have contributed to amplified microbial activity in soils with lowland plants in the glasshouse experiment. However, it is unlikely that increased biomass was solely responsible for lowland plant effects in the field experiments because lowland plant cover was low (mean ± SE relative cover: 4.7% ± 0.7%). Furthermore, in the glasshouse experiment we found that trait variation represented by the second principal axis had opposing effects on microbial respiration between lowland and alpine treatments

(Supplementary Fig. S4b; treatment × PC2: LR = 5.98, N = 20, P = 0.015). Lowland plants thus displayed relationships with the alpine soil system that were qualitatively distinct from resident alpine plants. While focussed work is now needed to pinpoint the physiological mechanism behind these differences, the existence of such a distinction raises the possibility that the arrival of warm-adapted lowland plants in warming alpine ecosystems initiates a change in plant community functioning *via* changes to traits.

Plants adapted to different temperatures can possess distinct traits that have opposing effects on carbon cycle processes[12]. We therefore tested whether lowland plants in both the glasshouse and field experiments indeed differed from alpine plants in traits that could affect the alpine soil system. Publicly available trait data[36] (plant height, leaf area, seed mass, SLA, leaf carbon, leaf nitrogen; Methods) relating to all species in the field experiments revealed that lowland plants on average possessed distinct traits from alpine plants (Supplementary Fig. S2a; P = 0.025). This was also the case for measured traits of lowland versus alpine plants under standardised conditions in the glasshouse experiment (multidimensional analyses: Supplementary Fig. S2b; P = 0.001). Coupled to this, while lowland plants were present at low abundance in the field experiments (mean ± SE relative cover: 4.7% ± 0.7%), the magnitude of soil carbon loss caused by lowland plant presence was greater in plots where lowland plants were more abundant overall (Supplementary Fig. S2e; $F_{1,14}$ = 5.79, P = 0.030). By contrast, we found no evidence that soil carbon loss in the field was alternatively related to individual lowland species identity (Supplementary Fig. S2c,d) or the composition of the recipient alpine plant community (Supplementary Fig. S2,g). Based on these observations, we hypothesise that, at least in our experiments, the establishment of warm-adapted lowland plants in warmed alpine ecosystems introduces novel traits into the community that alter plant community functioning and thus carbon cycle processes, with effects that intensify as lowland plants become more abundant in the community[32,33]. An alternative explanation is that, trait differences aside, the ability of alpine plants to facilitate soil carbon loss is suppressed when growing in warmer climates to which they are not adapted. While either explanation could ultimately accelerate soil carbon loss from warming alpine ecosystems following the arrival of

lowland plants, future studies are needed to fully unravel the environmental-dependence of plant

trait effects on soil ecosystem processes.

<u>Lowland plant-induced carbon loss is temporally dynamic</u>

We have shown using two field experiments that lowland plants facilitate a decrease in soil carbon

content from warming alpine ecosystems (Fig. 1) and, with a glasshouse experiment, that they

potentially act by accelerating soil carbon outputs *via* microbial respiration relative to alpine plants

(Fig. 3, Fig. 4), as opposed to decreasing soil carbon inputs. We therefore hypothesised that

lowland plant arrival in transplanted alpine plots in the field experiments would drive a net release

of carbon dioxide from the ecosystem. As for the glasshouse experiment, we initially examined

how lowland plants affected the balance between carbon inputs and outputs of the ecosystem

using carbon dioxide flux measurements taken on multiple dates during the growing season

(Methods). We then used soil samples collected on one representative date to focus on intrinsic

rates (that is, per unit of biomass) of microbial respiration and growth as processes that promote

soil carbon loss and gain, respectively[29,30], as well as microbial biomass as the scalar through

which these processes affect soil carbon[37,38].

We found that transplantation from high to low elevation (i.e. warming-only) affected alpine

ecosystem net carbon dioxide exchange differently between regions (Fig. 5a; Table S1; $P = 0.040$).

In the western Alps, transplantation changed the ecosystem from a net sink to a net source of

carbon dioxide, whereas in the central Alps it made the ecosystem a stronger net carbon dioxide

sink. This effect occurred because although transplantation accelerated rates of both gross primary

production (Fig. 5b; Table S1; $P < 0.001$) and ecosystem respiration (Fig. 5c; Table S1; $P < 0.001$),

in the western Alps it had stronger effects on ecosystem respiration and in the central Alps

stronger effects on gross primary production. Importantly, lowland plants had no significant bearing

over net ecosystem exchange (Fig. 5a), implying that although lowland plants were associated with

soil carbon loss from warmed alpine plots (Fig. 1), this must have occurred prior to carbon dioxide

measurements being taken and was no longer actively occurring. Gross primary production in

warmed alpine plots was also unaffected by lowland plant presence (Fig. 5b), which is consistent

with the low abundance of lowland plants in alpine communities (mean ± SE relative cover: 4.7% ± 0.7%). As such, lowland plants likely did not facilitate alpine soil carbon loss under warming by decreasing carbon inputs into the ecosystem. By contrast, ecosystem respiration in warmed alpine plots was depressed in the presence versus absence of lowland plants (Fig. 5c). These findings generally support the hypothesis that lowland plants affect the alpine soil system by changing carbon outputs. However, they contrast with expectations that lowland plants perpetually increase carbon outputs from the ecosystem and thus raise questions about how soil carbon was lost from warmed plots containing lowland plants (Fig. 1).

Fig. 5 | Warming and lowland plant effects on alpine ecosystem carbon dioxide fluxes in the field experiment. Effects of transplantation to low elevation (warming, W), transplantation plus lowland plants (warming plus lowlands, WL) or replantation at high elevation (control, C) on alpine ecosystem carbon dioxide fluxes (mean ± SE) during one season. (a) Net ecosystem CO_2 exchange (NEE; g CO_2 m^{-2} h^{-1}). (b) Gross primary production (GPP; g CO_2 m^{-2} h^{-1}). (c) Ecosystem respiration (ER; g CO_2 m^{-2} h^{-1}). Measurements were taken *in situ* at both experiments on multiple dates (western Alps: N = 196, central Alps: N = 90), with negative values indicating carbon uptake into the ecosystem. Treatment effects ($P < 0.05$) are indicated by letters (LMEs, Tukey; see Table S1).

Fig. 6 | Warming and lowland plant effects on the alpine soil system in the field experiment. Effects of transplantation to low elevation (warming, W), transplantation plus lowland plants (warming plus lowlands, WL) or replantation at high elevation (control, C) on alpine soil pools and processes (means ± SEs) after one season. (a) Biomass-specific microbial respiration (R_{mass}; ng C µg^{-1} C_{mic} h^{-1}). (b) Biomass-specific microbial growth (G_{mass}; ng C µg^{-1} C_{mic} h^{-1}). (c) C_{mic} (µg C g^{-1}). (d) Total microbial respiration (µg C g^{-1} h^{-1}). (e) Total microbial growth (µg C g^{-1} h^{-1}). (f) Microbial carbon use efficiency (CUE; %). Measurements were taken on soils sampled on one date, for (f) at both experiments (western Alps: N = 28; central Alps: N = 30) and for all other variables at the western Alps experiment (N = 28). Treatment effects ($P < 0.05$) are indicated by letters (LMEs, Tukey; see Table S1).

Carbon cycle processes are constrained by multiple feedbacks within the soil system, such as substrate availability[37,38] and microbial acclimation[39–41], that over time can slow, or even arrest, soil carbon loss[38,40,41]. We thus interrogated the state of the soil system in the field experiments in the western Alps experiment to explore whether such a feedback may be operating here, in particular to limit ecosystem respiration once soil carbon content had decreased in warmed alpine plots containing lowland plants. We found that biomass-specific rates of microbial respiration in the field were greater following transplantation to the warmer site in the presence, but not absence, of

lowland plants (Fig. 6a; Table S1; $P = 0.038$). By contrast, biomass-specific rates of microbial

growth were slower following transplantation but remained unaffected by lowland plant presence

(Fig. 6b; Table S1; $P = 0.016$). Thus, despite lower rates of ecosystem respiration overall, alpine

soil microbes still respired intrinsically faster in warmed plots containing lowland plants. Moreover,

accelerated microbial respiration, but not growth, implies that alpine soils had a higher capacity to

lose carbon under warming, but not to gain carbon *via* accumulation into microbial biomass, when

lowland plants were present[29,30]. These findings align with observations from the glasshouse

experiment that lowland plants generally accelerated intrinsic rates of microbial respiration (Fig. 3),

although in field conditions this effect occurred in tandem with warming.

Faster microbial respiration can only yield a loss of soil carbon if it occurs on a per unit of soil

basis[38,41,42], which was the case in the glasshouse experiment (Fig. 3). However, in the field we

found that lowland plants caused a reduction of microbial biomass carbon in alpine communities

transplanted to the warmer site (Fig. 6c; Table S1; $P < 0.001$). As such, although microbes

generally respired intrinsically faster in the presence of lowland plants, in the field they had

become less abundant and so we observed no change to microbial respiration when expressed

per unit of soil (Fig. 6d; Table S1; $P = 0.550$). The decline in microbial biomass likely took place

because soil microbes are chiefly carbon limited[43,] making their abundance tightly coupled to soil

carbon content (here: $r_{26} = 0.61$, $P = 0.001$)[38]. Moreover, since intrinsic rates of microbial growth

were unaffected by lowland plant presence, a reduction in microbial biomass translated into a

decline in microbial growth per unit of soil (Fig. 6e; Table S1; Table S1; $P = 0.009$) – preventing

any recovery of microbial biomass *via* growth[30]. We thus suggest that as soil carbon was lost,

microbial biomass also declined, and at the time of measurement this had eliminated the influence

of accelerated biomass-specific microbial respiration over the soil system. In short, we propose

based on these relationships that the decrease in soil carbon content observed in warmer

transplanted plots containing lowland plants had both started and stopped within one season.

Taken together, one interpretation of our findings is that the establishment of lowland plants in

warming alpine ecosystems accelerates intrinsic rates of microbial respiration (Fig. 3, Fig. 6a),

leading to soil carbon release at baseline levels of microbial biomass (Fig. 1, Fig. 3c), a coupled decline in microbial biomass (Fig. 6c) and a cessation of further carbon loss from the ecosystem (Fig. 5a, Fig. 6d). Although such a mechanism has been reported in other ecosystems[38,41], applying it here is speculative without additional timepoints because field soil measurements came from a single sampling event after soil carbon had already been lost from the ecosystem. For instance, an alternative mechanism could be that soil microbes acclimate to the presence of lowland plants and this decelerated microbial processes over time[39–41]. Nevertheless, it remains true that biomass-specific microbial growth and respiration affect the soil system as a function of microbial biomass, and here changes to microbial biomass could explain the behaviour of the soil system in both glasshouse and field experiments. We also found no evidence that soil microbes instead changed their resource use strategy in the presence versus absence of lowland plants (that is, carbon use efficiency; Fig. 6f; Table S1; $P < 0.001$; Tukey: $P = 0.996$), pointing away from microbial acclimation as an alternative mechanism. Beyond the mechanism for lowland plant effects on alpine soil carbon loss, it is conceivable that soil carbon loss is not isolated to a single season, but will reoccur in the future even without further warming or lowland plant arrival. This is especially true in the western Alps experiment where warming yielded a net output of carbon dioxide from the ecosystem (Fig. 5a). Moreover, in our field experiments we simulated a single event of lowland plant establishment and at relatively low abundance in the community (mean ± SE relative cover: 4.7% ± 0.7%), raising the possibility that increases in lowland plant cover or repeated establishment events in the future could facilitate further decreases in alpine soil carbon content under warming.

General conclusions

In summary, we show that uphill migrations of native lowland plants have the potential to facilitate warming effects on soil carbon loss from alpine ecosystems. This is important because it demonstrates that plant range expansions between herbaceous plant communities have the potential to contribute significantly to climate feedbacks. These results contrast with expectations from woody species expansions in the Arctic, where shrubs with recalcitrant litter can slow carbon turnover in herbaceous communities and potentially yield a negative climate feedback[15,16]. We thus

suggest that the influence of range expanding species on ecosystem carbon dynamics depends on

the relative trait composition of migrating species versus the recipient community[44], as well as

subsequent effects on environmental processes[15]. While our findings demonstrate that lowland

plants affect the rate of soil carbon release in the short term, short-term experiments, such as ours,

cannot resolve whether lowland plants will also affect the total amount of soil carbon lost in the

long term. This includes whether processes such as genetic adaptation (in both alpine and lowland

plants)[12] or community change[45,46] will moderate soil carbon responses to gradual or sustained

warming. Nevertheless, here decreases in soil carbon content occurred rapidly despite a low

abundance of lowland plants, and increased with total lowland plant cover. Given that both

warming and associated plant migrations will intensify over time[8,9], our estimates of lowland plant

impacts may be conservative – particularly if increases in lowland plant cover or repeated

establishment events cause recurrent pulses of soil carbon release. Moreover, while alpine

ecosystems store only a small fraction of total global soil carbon[1,2], redistributions among

herbaceous plant communities are occurring globally[4–9], including in similarly cool ecosystems in

the Arctic that contain vast soil carbon stocks[47]. This raises the potential for resulting climate

feedbacks to also be widespread. Future work should focus on testing the conditions under which

this feedback could occur in different mountain regions, as well as other ecosystems, experiencing

influxes of range expanding plant species, on quantifying how deeply it occurs in shallow alpine

soils, and on estimating the magnitude of the climate feedback given both ongoing warming and

variation in rates of species range shifts.

Materials & Methods

Field site description, experimental design. We made use of two independent alpine grassland transplant experiments that were established in Switzerland in the western Alps (46°12′40″N, 7°03′13″E)[28] and central Alps (46°52′20″N, 9°28′58″E) in autumn 2016. The experiments simulate effects of climate warming on alpine ecosystems *via* transplantation from high to low elevation, including the arrival of lowland plants due to warming-induced uphill plant migrations (Supplementary Fig. S1). In the western Alps experiment the plant community is dominated by *Sesleria caerulea*, *Carex sempervirens* and *Vaccinium vitis-idea*. In the central Alps experiment the dominant species are *Nardus stricta*, *Alchemilla xanthochlora*, *Trifolium pratense* and *Anthyllis vulneraria*. Broadly, the experiments share the same design. Turfs (area = 1 m^2, depth = 20 cm) of intact alpine vegetation plus rhizosphere soil were removed from a high elevation control site (western Alps: 2200 m, MAT = 0.7 ºC, TAP = 1448 mm; central Alps: 2000 m, MAT = 1.6 ºC, TAP = 1151 mm; ref. [48]) and replanted either at a low elevation site to simulate climate warming (western Alps: 1400 m, MAT = 6.0 ºC, TAP = 1236 mm, N = 10; central Alps: 1400 m, MAT = 4.4 ºC, TAP = 1010 mm, N = 20; ref. [48]) or back into a high elevation site as a transplantation control (see below). At the low elevation site, we created an additional treatment to simulate the arrival of lowland plants under warming by planting a low abundance of lowland plant species in a grid design (13 cm between individuals), creating a lowland plant density of approximately 56 individuals per m^2 (4.7% ± 0.7% total cover) in a background alpine plant community. Lowland plants were positioned randomly in the grid, but with the same number of individuals per species per plot, and we controlled for planting disturbance in all remaining plots using alpine plants in place of lowland plants (see below). In the western Alps region, lowland plant species were *Achillea millefolium, Bellis perennis, Bromus erectus, Dactylis glomerata, Medicago lupulina, Plantago media* and *Salvia pratensis*, and individuals were planted into one side of each plot in a split-plot design (n = 10). In the central Alps experiment, lowland plant species were *Brachypodium pinnatum, Carex flacca, Carum carvi, Dactylis glomerata, Hypericum perforatum, Plantago lanceolata, Primula veris, Ranunculus bulbosus, Salvia pratensis*, *Silene vulgaris*, *Trifolium montanum* and *Viola hirta*, and individuals were planted into half of the transplanted turfs (n = 10). While exact dispersal distances for selected lowland species are unknown, all species are

widespread and are expected to migrate uphill under warming[8] and the horizontal distance between high and low sites in the field experiments was always less than 2 km. In the western Alps experiment, lowland individuals were pre-grown in a greenhouse in steam-sterilized field soil and planted into the plots in May 2017. In the central Alps experiment, lowland individuals were taken from the surrounding vegetation and planted into the plots in autumn 2016. In both regions, soil was partially removed from lowland individuals prior to planting, and we performed an additional experiment to eliminate the possibility that small contaminations of lowland soil biota during planting could have modified alpine soil microbial community composition (see *Field Experiment Soil Biota Validation*, below). Overall, the experimental design in both regions yielded three treatments (n = 10 for each) arranged in a block design: transplantation from high to low elevation (i.e. warming-only, W); transplantation plus low elevation plants (i.e. warming plus lowland plant establishment, WL); and replantation at the high elevation origin site (i.e. control, C). One turf at the low elevation site in the western Alps experiment failed, leaving a total sample size of N = 28 and N = 30 in the western Alps and central Alps, respectively.

Field experiment controls and validation. We controlled for transplantation disturbance in both experiments by moving alpine control turfs back into a high elevation control site. In the western Alps experiment, alpine turfs were planted into different positions at their origin high elevation site. In the central Alps experiment, alpine turfs were planted into a new high elevation site at the same elevation. As such, we eliminated the possibility that alpine turfs changed simply because they were disturbed or were moved to a different location (that is, irrespective of elevation). The transplantation process had no effect on alpine soil carbon content, in that we found no difference in soil carbon content between replanted alpine control turfs at high elevation sites and undisturbed alpine soil in the surrounding area (LR = 2.27, N = 16, *P* = 0.132). We also controlled for potential effects of planting disturbance in all turfs not planted with lowland plants (that is, remaining turfs at low elevation and all control turfs at high elevation). In the western Alps experiment, we planted alpine species from the high elevation control site into the other side of alpine turfs at the low elevation site, as well as into all control turfs at the high elevation site. This was done in autumn 2016 using the same approach and randomized hexagonal grid design. In the central Alps

experiment, we mimicked planting disturbance by digging up and replanting existing individuals in

alpine turfs at the low elevation site not containing lowland plants, as well as all control turfs at the

high elevation site. This was done in spring 2017 using the same grid design. It is important to note

that we did not perform a reverse transplantation (that is, from low to high elevation), so we cannot

entirely rule out the possibility that transplantation of any community to any new environment could

yield a loss of soil carbon. Nevertheless, it is well established that climate warming promotes soil

carbon loss[35,49,50], and we found that soil from low elevation sites (that is, the potential end-member

for warming alpine ecosystems) contained less carbon than soil from the alpine sites (70.1 ± 3.05

mg C g^{-1} soil dry mass compared to 145.0 ± 5.29 mg C g^{-1} soil dry mass; LR = 37.2, N = 27, P <

0.001), validating our expectation that alpine soil would lose carbon under warming. Finally, while

small variations in experimental design between regions, such in the order of planting in W and WL

treatments, likely introduced some error when comparing experiments, we used both regions

together so that we could identify treatment effects that were consistent among them and were

thus not artefacts of experimental design.

Field experiment soil biota validation. We tested for the possibility that soil carbon loss was

alternatively caused by low elevation soil microbes hitchhiking on lowland plants during planting.

Specifically, we performed an experiment at the central Alps site to determine whether inoculating

alpine plots with low elevation soil biota affected soil microbial community composition. We first

generated a low elevation soil biota inoculum from rhizosphere soil from the same area where

lowland plants were sourced and gently mixing and sieving it with tap water (following refs [51,52]).

We then added 3 L of inoculum to one half of each of 10 x 1 m^2 alpine turfs equivalent to those in

the main experiment, followed by 3 L of water to rinse the inoculum from the vegetation and

facilitate its infiltration into the soil. The other half of the plot (separated by an impermeable rubber

barrier buried 15 cm deep into the soil around the perimeter of each subplot) received 6 L of water

as a control. Soil inoculation was performed in mid-May and repeated in mid-June 2017. In

October 2017, we sampled soils by pooling three soil cores (ø = 1 cm, d = 5 cm) from each

subplot, following which we determined soil microbial community composition by sequencing the

16S rRNA gene for bacteria/archaea and the ITS1 region for fungi[53]. DNA was extracted

(NucleoSpin Soil kit, Macherey-Nagel, Düren, Germany) and amplified using quadruplicate PCRs,

alongside extraction and PCR blank controls. PCR products were purified (MinEluteTM PCR

purification kit, Qiagen, Hilden, Germany) prior to sequencing on a MiSeq sequencing platform

(Illumina, San Diego, USA). Sequence data were processed using the OBITools software

package[54] and following[55]. Paired-end reads were merged, assigned to their respective samples

and dereplicated, and sequences with too short read lengths and with only one read were

excluded. Remaining sequences were clustered using the SUMACLUST[56] algorithm and

summarized into operational taxonomic units (OTUs) using the Infomap Community Detection

Algorithm[57], following which we calculated per-sample relative OTU abundances for

bacteria/archaea and fungi separately. We found that the alpine soil microbial community was

resistant to the introduction of low elevation soil biota, in that inoculation had no effect on soil

bacterial (Supplementary Fig. S5a; PERMANOVA; $F_{1,9} = 0.75$, $P = 0.573$) or fungal

(Supplementary Fig. S5b; $F_{1,9} = 0.75$, $P = 0.774$) community composition. The inoculation

represented a strong soil biota treatment relative to contamination by hitchhiking microbes. As

such, even if some contamination by low elevation soil biota did occur, it was unlikely to have

resulted in a change to the extant soil community, ruling out the possibility that contamination was

alternatively responsible for the soil carbon loss observed when lowland plants were present in

transplanted alpine plots.

Field plant community composition. We surveyed the composition of alpine communities in

summer 2017. This was done by dividing each plot into 5 × 5 cm cells and visually estimating the

canopy cover of all species rooted in every grid cell. Species cover was approximated with an

ordinal scale in which the first three cover categories correspond to 1/16, 1/8, and 1/4 of the grid

cell area (25 cm^2), and subsequent cover categories were constant increases of 1/4 of the grid cell

area. Plants were identified to species level whenever possible, although some taxa were identified

only to genus or subfamily levels when species identity was unclear based on vegetative traits.

Species cover (cm^2) was then calculated as the species-wise sum of cover estimates across all

grid cells in the quarter of the plot used for soil and ecosystem respiration measurements (see

below).

Field soil carbon content, microbial physiology, microbial biomass. We collected soil

samples from all treatments in both field experiments (N = 58) in August 2017. In each plot, we

created a composite sample from three cores (ø = 1 cm, approx. d = 7 cm) no closer than 7 cm

from a planted individual and from the same quarter of the plot used for ecosystem respiration

measurements (see below; Supplementary Fig. S1). We focused on rhizosphere soil for two

reasons. First, alpine soils are shallow and alpine plants are shallow-rooting, making the

rhizosphere the most logical target for a first assessment of warming and lowland plant effects on

alpine soil carbon cycling. Second, we wanted to avoid any potential contamination from the base

of the turf (depth = 20 cm). Soils were kept cool, transported immediately to the laboratory, sieved

(2 mm mesh size), adjusted to 60% of water holding capacity (WHC) and pre-incubated for three

days at field temperature prior to measurements. We measured microbial biomass carbon (mg per

g dry mass) *via* chloroform fumigation extraction followed by analysis on a TOC-VCPH/CPNTNM-1

analyzer (Shimadzu; 48h incubation period; 1 M KCl extraction for fumigated and non-fumigated

samples; conversion factor 0.45)[58]. While this approach also yielded measurements of dissolved

organic carbon (DOC; mg C per g dry mass), treatment effects on this variable echoed those of

microbial biomass carbon (Table S1) so we did not interpret them separately. Soil organic carbon

content (%) was measured on dry soil (60 ºC for 48h) with a Carlo Erba 1110 elemental analyzer

(EA, CE Instruments) coupled to a Delta Plus IRMS via a Conflo III (Thermo Fisher). It was not

possible to take widespread measurements of soil bulk density due to the destructive sampling

required while other studies were underway (e.g. ref [28]). Instead, we took additional soil cores (ø =

5 cm, d = 5 cm) from the central Alps experiment in 2021 once other studies were complete to

indirectly explore whether lowland plant effects on soil carbon content in warmed alpine plots could

have occurred due to changes in soil bulk density. We found that although transplantation to the

warmer site increased alpine soil bulk density (LR = 7.18, P = 0.028, Tukey: P < 0.05), lowland

plants had no effect (Tukey: P = 0.999). It is not possible to make direct inferences about the soil

carbon stock using measurements made on different soil cores four years apart. Nevertheless,

these results make it unlikely that lowland plant effects on soil carbon content in warmed alpine

plots occurred simply due to a change in soil bulk density. We additionally quantified microbial

growth, respiration and carbon use efficiency in soils from the western Alps experiment (N = 28)

using the incorporation of ^{18}O into DNA[38]. Briefly, we incubated 500 mg of soil for 24h at field

temperature following addition of either ^{18}O-H_2O to 20 atom% enrichment and 80% of WHC or the

same volume of molecular grade non-labelled H_2O as a natural abundance control. Gas samples

were taken at the start and end of the 24h incubation and analyzed for CO_2 concentration with a

Trace GC Ultra (Thermo Fisher) to determine microbial respiration (μg C per g dry mass per h). At

the end of the incubation, soil was snap-frozen in liquid nitrogen, and DNA was extracted

(FastDNA SPIN Kit for Soil, MP Biomedical), quantified (Quant-iT PicoGreen dsDNA Assay Kit,

Thermo Fisher) and analyzed for ^{18}O abundance and total oxygen content using a thermochemical

EA coupled to a Delta V Advantage IRMS via a Conflo III (Thermo Fisher). Following ref. [38], we

calculated total DNA production during the 24h incubation (μg DNA per gram dry mass) and

coupled it with measures of microbial biomass carbon content to derive microbial growth (μg C per

gram dry mass per h) and carbon use efficiency (%).

Field ecosystem respiration. We measured rates of ecosystem respiration (g CO_2 per m^2 per h)

on multiple dates (western Alps: 7 dates; central Alps: 3 dates) between July and August 2017.

One month prior to the first measurement, we installed permanent gas sampling collars (h = 10 cm,

$ø$ = 30 cm) in each plot to a depth of 5 cm to create an airtight seal at the soil surface[12].

Measurements were made by enclosing plants and soil within the collars with an opaque chamber

(h = 50 cm, $ø$ = 30 cm) and, following a 30s equilibration period, monitoring the increase in CO_2

concentration at 5s intervals over a 60s period (Vaisala GMP343 CO_2 probe attached to an HM70

meter, Vaisala). Chambers contained a fan to mix the headspace during measurements and a

small sealable hole on the top that was opened when placing the chamber on sampling collars to

avoid pressurization inside the chamber. Soil temperature and moisture were recorded for each

measurement and used alongside measurements of air temperature, air pressure, enclosure

volume and soil surface area to calculate the CO_2 flux[12]. Where necessary, we repeated

measurements in the field to attain a visually consistent CO_2 change over time, and during data

processing we discarded fluxes with an *r*-squared of less than 0.9 for this relationship.

Glasshouse experiment design. We established a six-week pot experiment to test for effects of

lowland versus alpine plants on the alpine soil system. Specifically, we filled pots (ø = 10 cm, depth

= 9 cm) with sieved soil (mesh size 4 mm) from the high elevation site of the western Alps field

experiment, and planted them with six lowland species (*Bromus erectus*, *Salvia pratensis*,

Medicago lupulina) or six alpine species (*Phleum alpinum*, *Anthyllis vulneraria* ssp. *alpestris*,

Plantago alpina) to create a lowland only treatment, an alpine only treatment and a bare soil

control (N = 30). We selected a grass, herb and legume species for each treatment, all of which

were found naturally at the low (lowland treatment) or high (alpine treatment) elevation sites in the

western Alps field experiment. Vegetated treatments contained two individuals of all lowland or

alpine species planted in a circular pattern such they were never adjacent to conspecifics or plants

of the same functional type (i.e. grass, herb, legume), making replicates within treatments identical

in species composition and neighbourhoods. The original design also included a mixed community

treatment containing one individual of each lowland and alpine species. However, the mixed

treatment had double the species richness of other vegetated treatments and contained an equal

cover of lowland and alpine plants (compared to < 5% lowland plant cover in field plots), so we

focussed our analyses on the comparison between alpine and lowland plant treatments to

understand how lowland plants affect the soil differently from alpine plants. All plants were pre-

grown from seed provided by commercial suppliers (www.hauenstein.ch, www.schutzfilisur.ch,

www.saatbau.at) in steam-sterilized potting soil for two to three weeks to approximately the same

biomass and root-washed immediately prior to planting. Pots were arranged in a randomized block

design (n = 10) in a greenhouse under standardized growing conditions (22 ºC; 50% humidity; no

artificial light), and were watered every two to three days as needed. After six weeks, we

destructively harvested the plants and soil to characterize treatment effects on plant physiology

and the soil system.

Glasshouse plant physiology. We measured maximum leaf-level photosynthesis (photosynthetic

capacity; A_{max}; µmol CO_2 per m^2 per s) and stomatal conductance (g_s; µmol H_2O per m^2 per s)

under optimum conditions on one individual per species per pot using an open path infrared gas

analyzer connected to a 2.5 cm chamber (CIRAS-2 and PLC-6, PP Systems)[12]. Readings were

taken once for each individual and corrected using measurements of leaf area (cm^2) on the same

leaf, which we also dried and weighed to determine specific leaf area (SLA; cm^2 per gram).

Following soil pore water sampling (below), we quantified total, aboveground and belowground

biomass of all individuals in the experiment by destructively sampling, washing, drying (60 ºC for

48h) and weighing all plant tissue.

Glasshouse soil pore water analysis. We collected soil pore water from all pots prior to soil

sampling (see below) to determine treatment effects on dissolved organic matter (DOM) quality.

Soil water was sampled by leaching pots twice with 50 mL sterile tap water, combining and filtering

leached water (filter pore size 0.2 μm) and measuring absorbance (200-900 nm with a 1nm

increment; 1 cm quartz cuvette; UV-vis spectrophotometer, Shimadzu) and fluorescence intensity

(excitation: 240-450 nm with a 5 nm increment; emission: 290-550 nm with a 2.5 nm increment;

Fluorolog-3 spectrometer, Horiba). We corrected absorbance spectra against a milli-Q water blank,

calculated Napierian absorption coefficients and used these data to derive absorbance at 350 nm

(a_{350}; a proxy for DOM content). We expressed fluorescence data as total fluorescence (F_{tot};

Raman Units) and the fluorescence index (FI), i.e. the ratio of emission intensity at 470 nm versus

520 nm at an excitation wavelength of 370 nm (a proxy for microbially-derived DOM[59]). We also

derived excitation-emission matrices (EEMs) to quantify the relative abundance of different

components of DOM through PARAFAC algorithms[60]. Following preprocessing (i.e. Raman

scattering removal and standardization to Raman Units) and normalization[61], we used the drEEM

toolbox[62] in MATLAB (Mathworks) to obtain a six-component PARAFAC model. The model was

validated with split-half analysis, random initialization and visualization of EEM residuals, after

which we calculated the relative abundance of each component as the maximum fluorescence

signal of the component against the total maximum fluorescence signal of all components. Finally,

we classified components against classical EEM nomenclature (Table S4; refs [63,64]). While

PARAFAC modelling provides a means to discriminate between DOM sources (e.g. components

C3 and C4; Table S4), we did not physically separate plant- and microbe-derived DOM when

sampling soil pore water. As such, observed changes to DOM composition were likely a result of

changes to both plant and microbial processes (see Main Text).

Glasshouse soil carbon pools, microbial respiration. We took soil cores ($\emptyset$ = 1 cm, d = 7 cm)

from the center of each pot (i.e. equidistant from all individuals) to quantify soil carbon pools and

microbial respiration. Soils were sieved (2 mm mesh size), adjusted to 60% of their WHC and pre-

incubated at 22 ºC for three days prior to measurements. We quantified concentrations of microbial

biomass carbon in all soils as for field soil (see above; mg C per g dry mass). We determined

microbial respiration by incubating 8 g sieved soil in the dark at 22 ºC for a further six-week period

without plants. We measured microbial respiration daily for the first four days, then approximately

biweekly until the end of the six-week period. We measured CO_2 concentrations (EGM-4, PP

Systems) of gas samples taken approximately 24h after flushing enclosed vials (v = 126 mL) with

CO_2-free air, and calculated microbial respiration (µg C per g dry mass per h) using accompanying

measurements of enclosure volume, air pressure and soil mass[12]. After each measurement, soils

were weighed and, when necessary, re-adjusted to 60% of their WHC.

Two-pool mixed models of soil carbon mineralization. We described the kinetics of soil carbon

mineralization during the follow-up incubation of glasshouse soil by employing a two-pool mixed

model to microbial respiration data[65]. Specifically, we constructed a model that considered a first

pool that followed first-order decomposition kinetics (i.e. a small soil carbon pool with a fast

turnover) and a second pool that followed zero-order kinetics (i.e. decay rate was constant over the

incubation period, representing a large soil carbon pool with a slow turnover):

$$C_{min} = C_l\left(1 - e^{-tk_l}\right) + B \cdot t$$

where C_{min} is the cumulative CO_2 respired (µg C g^{-1}) over the incubation time t (d), C_l and k_l are the

size (µg C g^{-1}) and decomposition constant (d^{-1}) of the first-order kinetics carbon pool, respectively,

and B is the basal respiration rate (µg C $g^{-1}\,d^{-1}$) of the zero-order kinetics carbon pool. Given that

C_{min} and t were known, we ran the two-pool mixed model with the nlsList function (maxiter = 100,

tol = 1 × 10^{-5}, minFactor = 1/1024) in R[66] using the package "nlme"[67] for each pot separately to

estimate values for C_l, k_l and B. Models failed to converge for one sample, and we excluded four

additional samples for which models made negative pool estimates (N = 25).

Statistical analysis. The majority of statistical test outputs appear as *P*-values in the Main Text,

with full outputs being provided in Tables S1 to S3. See text below for exceptions. Data analyses

and visualization were undertaken in R[66] using the packages "cowplot"[68], "drake"[69], "emmeans"[70],

"nlme"[67], "tidyverse"[71] and "vegan"[72]. Where necessary, region was accounted for in statistical

models as a fixed effect, since two levels was too few to be included as a random intercept term.

As such, region effects are presented (Tables S1, S3), but not interpreted. Statistical methods are

described as they appear in the Main Text.

<u>Warming and warming plus lowland plant effects on soil carbon content:</u> we used field data to test

for treatment (C, W, WL) effects on soil carbon content using an LME model including region as a

fixed effect and block as a random intercept term (see Table S1 for test output). We also

calculated how much more soil carbon loss occurred in the presence versus absence of lowland

plants by fitting a single LME testing for effects of soil temperature (plot means of all sampling

dates), the presence or absence of lowland plants and their interaction on soil carbon content (see

Supplementary Fig. S2 for test outputs). In order to compare slopes between C-W and C-WL

treatment combinations, we duplicated C treatment values, assigned one set to each C-W and C-

WL pair and accounted for this using a random intercept term. We also included region as a

random slope term. We then tested how much more soil carbon loss would occur per ºC of

warming when lowland plants are present by calculating the relative difference (% ± 95%

confidence intervals) between the C-WL and C-W slopes.

<u>Plant mechanisms for soil carbon loss:</u> we expressed soil carbon content as the block-wise percent

changes between WL and W treatments to derive a metric for lowland plant induced soil carbon

loss (i.e. % soil carbon remaining). We tested for lowland species identity effects on soil carbon

remaining using separate LMEs for each region (due to different lowland species pools) that tested

for individual species cover (cm^2) effects on soil carbon remaining in WL plots, including species as

a random intercept term (see Supplementary Fig. S2 for test outputs). We also performed separate

linear models including species as fixed effects, none of which were statistically significant (Table

S3). We tested for effects of total lowland species cover (cm^2) on soil carbon remaining in WL plots

using a linear model also including region as a fixed effect. We summarized alpine community

composition in W and WL treatments using an NMDS (Hellinger distance matrix, two axes),

following which we expressed NMDS scores as the block-wise percent changes between W and

WL treatments to derive metrics for lowland plant induced alpine plant community change, and

used a linear model that tested for effects of changes to NMDS scores on soil carbon remaining in

WL plots including region as a fixed effect (see Supplementary Fig. S2 for test outputs). For

reference, raw NMDS scores did not significantly differ between W and WL treatments (LME:

NMDS #1: LR = 1.57, P = 0.210; NMDS #2: LR = 2.67, P = 0.102). Finally, we tested for

differences between alpine and lowland species traits in the field experiment using a

PERMANOVA on a principal components analysis (PCA) of species mean trait values (log_{10} plant

height, log_{10} leaf area, log_{10} seed mass, SLA, leaf carbon, leaf nitrogen) from the TRY database[36]

for all species present (where necessary, missing values were gap-filled following[73]), and in the

glasshouse experiment using a PERMANOVA on a PCA of physiological measurements (total

biomass, leaf biomass, root biomass, SLA, A_{max}, g_s) for all plants (see Supplementary Fig. S2 for

test outputs). Traits were selected to explore broad life history differences between alpine and

lowland plant species[74], but were not used to identify the specific trait(s) responsible for lowland

plant effects on alpine soil carbon loss. We used TRY traits for field species due to the large

number of species present in the communities (N = 118), but used measured traits in the

glasshouse experiment to provide direct evidence of physiological differences between alpine and

lowland species.

<u>Lowland versus alpine plant effects on the soil system:</u> we used the glasshouse experiment to test

for lowland versus alpine plant treatment effects on soil microbial biomass carbon, the size (C_l) and

decay rate (k_l) of the first modelled soil carbon pool and all variables relating to soil DOM quality

using LMEs including block as a random intercept term. We tested for effects of treatment, time

and their interaction on soil microbial respiration using the same LME structure but including the

bare soil treatment. In all cases, including/excluding the bare soil treatment had no effect over

model outcomes, but mean ± SE values for this treatment are included in figures for reference.

Finally, we explored how plant physiology in lowland and alpine treatments at the pot level affected

initial rates of soil microbial respiration. We defined initial microbial respiration as the pot-wise

mean respiration rates of the first nine days of soil incubation (that is, while treatment effects

persisted; see Fig. 3a), and expressed pot-level plant physiology using scores from the first two

axes of a PCA performed on sums of total, root and shoot biomass and biomass-weighted means

of SLA, A_{max} and g_s. We then tested for effects of treatment, PC1 and PC2 scores and their

interaction on initial microbial respiration using an LME including block as a random intercept term.

Statistical test outputs for LMEs are provided in Table S2, except for treatment and plant

physiology effects on microbial respiration, which appear in the Main Text.

<u>Sequence of warming and lowland plant effects on alpine soil carbon loss:</u> we used field data to

test for treatment (C, W, WL) effects on soil microbial biomass carbon and soil DOC using LME

models including region as a fixed effect and block as a random intercept term. We tested for

treatment effects on ecosystem respiration using an LME that included region as a fixed effect and

sampling date nested in block as a random intercept term. We tested for treatment effects on soil

microbial growth and respiration (per unit of soil and per unit of microbial biomass) and carbon use

efficiency in the western Alps experiment using LMEs including block as a random intercept term.

Statistical test outputs for these variables are provided in Table S1. We also used Pearson Product

Moment correlations to test for associations between soil microbial biomass and soil carbon

content, microbial respiration and growth in the western Alps experiment, irrespective of

experimental treatment.

Acknowledgements

JMA received funding from the European Union's Horizon 2020 research and innovation program

(grant no. 678841). JMA and TWNW received funding from the Swiss National Science Foundation

(grant no. 31003A-176044). KG is funded by the Swiss National Science Foundation (grant no.

PZ00P2-174047). SB and the central Alps experiment were funded by the Swiss National Science

Foundation (grant no. 31003A-173210). TM received funding from the French National Research Agency (ANR-20-CE02-0021).

Author contributions

TWNW, AB and JMA conceived the study. AR, AB and JMA provided resources for field and laboratory experiments. SB, TM and HN established and maintained field experiments. TWNW, KG and TG undertook the field work. TWNW, KG, TG, PM, CS, AB and JMA undertook the glasshouse experiment. TWNW, KG, TG, PM, CS and AR did the laboratory work. TL performed the soil pore water sampling and associated analyses. SB and TM performed the soil inoculation experiment and associated analyses. TWNW did the data analysis, with support from DR, TWC and JMA. TWNW wrote the manuscript in close collaboration with JMA, and with support from all co-authors.

Competing interests

The authors declare no competing financial interests.

References

1. Streit, K. *et al.* Soil warming alters microbial substrate use in alpine soils. *Glob. Change Biol.* **20,** 1327–1338 (2014).

2. Parker, T. C., Subke, J.-A. & Wookey, P. A. Rapid carbon turnover beneath shrub and tree vegetation is associated with low soil carbon stocks at a subarctic treeline. *Glob. Change Biol.* **21,** 2070–2081 (2015).

3. Knowles, J. F., Blanken, P. D., Lawrence, C. R. & Williams, M. W. Evidence for non-steady-state carbon emissions from snow-scoured alpine tundra. *Nat. Commun.* **10,** 1306 (2019).

4. Parmesan, C. Ecological and evolutionary responses to recent climate change. *Annu. Rev. Ecol. Evol. Syst.* **37,** 637–669 (2006).

5. Lenoir, J. & Svenning, J. C. Climate-related range shifts - a global multidimensional synthesis and new research directions. *Ecography* **38,** 15–28 (2015).

6. Pecl, G. T. *et al.* Biodiversity redistribution under climate change: Impacts on ecosystems and human well-being. *Science* **355,** (2017).

7. Dolezal, J. *et al.* Vegetation dynamics at the upper elevational limit of vascular plants in Himalaya. *Sci. Rep.* **6,** 24881 (2016).

8.   Rumpf, S. B. *et al.* Range dynamics of mountain plants decrease with elevation. *Proc. Natl. Acad. Sci. USA* **115,**

1848–1853 (2018).

9.   Steinbauer, M. J. *et al.* Accelerated increase in plant species richness on mountain summits is linked to warming.

Nature **556,** 231–234 (2018).

10.   Wallingford, P. D. *et al.* Adjusting the lens of invasion biology to focus on the impacts of climate-driven range shifts.

Nat. Clim. Chang. (2020). doi:10.1038/s41558-020-0768-2

11.   Canadell, J. G. *et al.* in *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the*

Sixth Assessment Report of the Intergovernmental Panel on Climate Change (eds. Masson-Delmotte, V. et al.)

(Cambridge University Press, 2021).

12.   Walker, T. W. N. *et al.* Plastic and genetic responses of a common sedge to warming have contrasting effects on

carbon cycle processes. *Ecol. Lett.* **22,** 159–169 (2019).

13.   Liao, C. *et al.* Altered ecosystem carbon and nitrogen cycles by plant invasion: a meta-analysis. *New Phytol.* **177,**

706–714 (2008).

14.   Waller, L. P. *et al.* Biotic interactions drive ecosystem responses to exotic plant invaders. *Science* **368,** 967–972

(2020).

15.   Pearson, R. G. *et al.* Shifts in Arctic vegetation and associated feedbacks under climate change. *Nat. Clim. Chang.*

3, 673–677 (2013).

16.   Cornelissen, J. H. C. *et al.* Global negative vegetation feedback to climate warming responses of leaf litter

decomposition rates in cold biomes. *Ecol. Lett.* **10,** 619–627 (2007).

17.   Körner, C. *Alpine Plant Life: Functional Ecology of High Mountain Ecosystems*. (Springer, 2003).

18.   Pepin, N. *et al.* Elevation-dependent warming in mountain regions of the world. *Nat. Clim. Chang.* **5,** 424–430

(2015).

19.   Vitasse, Y. *et al.* Phenological and elevational shifts of plants, animals and fungi under climate change in the

European Alps. *Biol. Rev. Camb. Philos. Soc.* **96,** 1816–1835 (2021).

20.   Lenoir, J., Gégout, J. C., Marquet, P. A., de Ruffray, P. & Brisse, H. A significant upward shift in plant species

optimum elevation during the 20th century. *Science* **320,** 1768–1771 (2008).

21.   Hagedorn, F., Gavazov, K. & Alexander, J. M. Above- and belowground linkages shape responses of mountain

vegetation to climate change. *Science* **365,** 1119–1123 (2019).

22.   Budge, K., Leifeld, J., Hiltbrunner, E. & Fuhrer, J. Alpine grassland soils contain large proportion of labile carbon

but indicate long turnover times. *Biogeosciences* **8,** 1911–1923 (2011).

23.   Hafner, S. *et al.* Effect of grazing on carbon stocks and assimilate partitioning in a Tibetan montane pasture

revealed by 13CO2 pulse labeling. *Glob. Change Biol.* **18,** 528–538 (2012).

24.   Studer, M. S., Siegwolf, R. T. W. & Abiven, S. Carbon transfer, partitioning and residence time in the plant-soil

system: a comparison of two[13] CO_2 labelling techniques. *Biogeosciences* **11,** 1637–1648 (2014).

25.   Hannah, L. in *Climate Change Biology* 273–295 (Elsevier, 2022). doi:10.1016/B978-0-08-102975-6.00013-3

26.   Shaver, G. R. *et al.* Global Warming and Terrestrial Ecosystems: A Conceptual Framework for Analysis. *Bioscience*
50, 871 (2000).

27.   Yang, Y. *et al.* Transplants, open top chambers (otcs) and gradient studies ask different questions in climate
change effects studies. *Front. Plant Sci.* **9,** 1574 (2018).

28.   Nomoto, H. A. & Alexander, J. M. Drivers of local extinction risk in alpine plants under warming climate. *Ecol. Lett.*
24, 1157–1166 (2021).

29.   Allison, S. D., Wallenstein, M. D. & Bradford, M. A. Soil-carbon response to warming dependent on microbial
physiology. *Nat. Geosci.* **3,** 336–340 (2010).

30.   Liang, C., Schimel, J. P. & Jastrow, J. D. The importance of anabolism in microbial control over soil carbon storage.
Nat. Microbiol. **2,** 17105 (2017).

31.   Wong, S. C., Cowan, I. R. & Farquhar, G. D. Stomatal conductance correlates with photosynthetic capacity. *Nature*
282, 424–426 (1979).

32.   De Deyn, G. B., Cornelissen, J. H. C. & Bardgett, R. D. Plant functional traits and soil carbon sequestration in
contrasting biomes. *Ecol. Lett.* **11,** 516–531 (2008).

33.   Cornwell, W. K. *et al.* Plant species traits are the predominant control on litter decomposition rates within biomes
worldwide. *Ecol. Lett.* **11,** 1065–1071 (2008).

34.   Jones, D. L., Nguyen, C. & Finlay, R. D. Carbon flow in the rhizosphere: carbon trading at the soil–root interface.
Plant Soil **321,** 5–33 (2009).

35.   Melillo, J. M. *et al.* Soil warming and carbon-cycle feedbacks to the climate system. *Science* **298,** 2173–2176
(2002).

36.   Kattge, J. *et al.* TRY plant trait database - enhanced coverage and open access. *Glob. Change Biol.* **26,** 119–188
(2020).

37.   Hartley, I. P., Hopkins, D. W., Garnett, M. H., Sommerkorn, M. & Wookey, P. A. Soil microbial respiration in arctic
soil does not acclimate to temperature. *Ecol. Lett.* **11,** 1092–1100 (2008).

38.   Walker, T. W. N. *et al.* Microbial temperature sensitivity and biomass change explain soil carbon loss with warming.
Nat. Clim. Chang. **8,** 885–889 (2018).

39.   Crowther, T. W. & Bradford, M. A. Thermal acclimation in widespread heterotrophic soil microbes. *Ecol. Lett.* **16,**
469–477 (2013).

40.   Bradford, M. A. *et al.* Cross-biome patterns in soil microbial respiration predictable from evolutionary theory on
thermal adaptation. *Nat. Ecol. Evol.* **3,** 223–231 (2019).

41.   Melillo, J. M. *et al.* Long-term pattern and magnitude of soil carbon feedback to the climate system in a warming
world. *Science* **358,** 101–105 (2017).

42.   Prommer, J. *et al.* Increased microbial growth, biomass, and turnover drive soil organic carbon accumulation at
higher plant diversity. *Glob. Change Biol.* **26,** 669–681 (2020).

43.   Hobbie, J. E. & Hobbie, E. A. Microbes in nature are limited by carbon and energy: the starving-survival lifestyle in

soil and consequences for estimating microbial rates. *Front. Microbiol.* **4,** 324 (2013).

44. Bjorkman, A. D. *et al.* Plant functional trait change across a warming tundra biome. *Nature* **562,** 57–62 (2018).

45. Walker, T. N. *et al.* Vascular plants promote ancient peatland carbon loss with climate warming. *Glob. Change Biol.*

22, 1880–1889 (2016).

46. Bardgett, R. D., Manning, P., Morriën, E. & De Vries, F. T. Hierarchical responses of plant-soil interactions to

climate change: consequences for the global carbon cycle. *J Ecol* **101,** 334–343 (2013).

47. Tarnocai, C. *et al.* Soil organic carbon pools in the northern circumpolar permafrost region. *Global Biogeochem.*

Cycles **23,** (2009).

48. Karger, D. N. *et al.* Climatologies at high resolution for the earth's land surface areas. *Sci. Data* **4,** 170122 (2017).

49. Crowther, T. W. *et al.* Quantifying global soil carbon losses in response to warming. *Nature* **540,** 104–108 (2016).

50. Davidson, E. A. & Janssens, I. A. Temperature sensitivity of soil carbon decomposition and feedbacks to climate

change. *Nature* **440,** 165–173 (2006).

51. van de Voorde, T. F. J., van der Putten, W. H. & Bezemer, T. M. Soil inoculation method determines the strength of

plant–soil interactions. *Soil Biol. Biochem.* **55,** 1–6 (2012).

52. De Vries, F. T., Bracht Jørgensen, H., Hedlund, K. & Bardgett, R. D. Disentangling plant and soil microbial controls

on carbon and nitrogen loss in grassland mesocosms. *J Ecol* **103,** 629–640 (2015).

53. Herbold, C. W. *et al.* A flexible and economical barcoding approach for highly multiplexed amplicon sequencing of

diverse target genes. *Front. Microbiol.* **6,** 731 (2015).

54. Boyer, F. *et al.* obitools: a unix-inspired software package for DNA metabarcoding. *Mol. Ecol. Resour.* **16,** 176–182

(2016).

55. Zinger, L. *et al.* Soil community assembly varies across body sizes in a tropical forest. *BioRxiv* (2017).

doi:10.1101/154278

56. Mercer, C., Boyer, F., Bonin, A. & Coissac, E. *SUMATRA and SUMACLUST: fast and exact comparison and*

clustering of sequences. (LECA, 2013).

57. Rosvall, M. & Bergstrom, C. T. Maps of random walks on complex networks reveal community structure. *Proc. Natl.*

Acad. Sci. USA **105,** 1118–1123 (2008).

58. Vance, E. D., Brookes, P. C. & Jenkinson, D. S. An extraction method for measuring soil microbial biomass C. *Soil*

Biol. Biochem. **19,** 703–707 (1987).

59. McKnight, D. M. *et al.* Spectrofluorometric characterization of dissolved organic matter for indication of precursor

organic material and aromaticity. *Limnol. Oceanogr.* **46,** 38–48 (2001).

60. Stedmon, C. A., Markager, S. & Bro, R. Tracing dissolved organic matter in aquatic environments using a new

approach to fluorescence spectroscopy. *Mar Chem* **82,** 239–254 (2003).

61. Zepp, R. G., Sheldon, W. M. & Moran, M. A. Dissolved organic fluorophores in southeastern US coastal waters:

correction method for eliminating Rayleigh and Raman scattering peaks in excitation–emission matrices. *Mar Chem*

89, 15–36 (2004).

62. Murphy, K. R., Stedmon, C. A., Graeber, D. & Bro, R. Fluorescence spectroscopy and multi-way techniques.

PARAFAC. *Anal. Methods* **5,** 6557 (2013).

63. Fellman, J. B., Hood, E. & Spencer, R. G. M. Fluorescence spectroscopy opens new windows into dissolved

organic matter dynamics in freshwater ecosystems: A review. *Limnol. Oceanogr.* **55,** 2452–2462 (2010).

64. *Aquatic organic matter fluorescence.* (Cambridge University Press, 2014). doi:10.1017/CBO9781139045452

65. Bonde, T. A. & Rosswall, T. Seasonal variation of potentially mineralizable nitrogen in four cropping systems. *Soil*

Sci. Soc. Am. J. **51,** 1508–1514 (1987).

66. R Core Team. *R: A language and environment for statistical   computing.* (R Foundation for Statistical Computing,

2014).

67. Pinheiro, J., Bates, D., DebRoy, S. & Sarkar, D. *nlme: Linear and Nonlinear Mixed Effects Models.* (R Project,

2019).

68. Wilke, C. O. *cowplot: Streamlined Plot Theme and Plot Annotations for "ggplot2."* (R Project, 2019).

69. Michael Landau, W. The drake R package: a pipeline toolkit for reproducibility and high-performance computing.

JOSS **3,** 550 (2018).

70. Lenth, R. *Estimated Marginal Means, aka Least-Squares Means.* (R Project, 2020).

71. Wickham, H. *et al.* Welcome to the tidyverse. *JOSS* **4,** 1686 (2019).

72. Oksanen, J. *et al. vegan: Community Ecology Package.* (R Project, 2019).

73. Schrodt, F. *et al.* BHPMF - a hierarchical Bayesian approach to gap-filling and trait prediction for macroecology and

functional biogeography. *Glob. Ecol. Biogeogr.* **24,** 1510–1521 (2015).

74. Díaz, S. *et al.* The global spectrum of plant form and function. *Nature* **529,** 167–171 (2016).

 Supplementary Figure Legends

Fig. S1 | Experimental design. (a) In both field experiments, intact alpine turfs containing alpine plant communities plus

rhizosphere soil were removed from their high elevation home sites and either replanted at the home site (negative

control; C) or transplanted to a low elevation recipient site to simulate climate warming (W, WL). At low elevation, half of

the turfs were planted with a low abundance of lowland plant species to additionally simulate the arrival of migrating plant

species in the ecosystem (WL). Both other treatments (i.e. C, W) were planted with the same abundance of alpine plant

species as a disturbance control. (b) The layout of W and WL treatments at the low elevation site differed between

western and central alps experiments. In the central alps, W and WL treatments were established in distinct turfs planted

adjacently in a block design (N = 10). In the western alps, W and WL treatments were established in different halves of

the same turf in a split-plot design (N = 10).

Fig. S2 | Alpine and lowland plant effects on soil carbon content. (a,b) Trait separation of lowland (yellow) and

alpine (green) plant species, displayed as PC1 and PC2 scores from PCAs containing (a) TRY data for species in field

experiments (N = 242; $F_{1,240}$ = 3.98, P = 0.003) or (b) measured data from a glasshouse experiment on selected species

(N = 109; $F_{1,107}$ = 17.69, P = 0.001). Arrows show loadings and P-values refer to alpine-lowland comparison

(PERMANOVAs). (c-g) Relationships between soil carbon loss (% initial soil carbon content remaining) and cover (cm^2)

of each lowland plant species in the (c) western Alps (N = 36; LME: LR = 0.62, P = 0.4302) and (d) central Alps (N = 80;

LME: LR = 0.33, P = 0.5666) experiments, (e) total lowland plant cover (cm^2; N = 19; see Main Text) and (f,g) alpine

plant community composition (% change in NMDS axes; N = 18; linear models: NMDS #1: $F_{1,14}$ = 1.19, P = 0.2934,

NMDS #2: $F_{1,14}$ = 3.17, P = 0.0966; region: $F_{1,14}$ = 1.29, P = 0.2755). For (c,d), non-significant species-wise fit lines

(yellow) are also displayed.

Fig. S3 | Soil temperature and lowland plant effects on alpine soil carbon content. Relationship between soil

temperature (ºC) and soil carbon content (C; mg g^{-1} dry mass) in alpine turfs exposed to warming (W; light grey),

warming plus lowland plants (WL; dark grey) or an ambient control (C; white). Statistics describe effects of warming (W;

LR = 17.65, P < 0.0001), lowland plants (L; LR = 11.77, P = 0.0006) and their interaction (W × L; LR = 11.72, P = 0.0006)

from one linear mixed effects model (N = 78), but are visualized with separate fit lines (including slopes) for C-W (dashed

lines) and C-WL (solid lines) comparisons in the western Alps (triangles) and central Alps (circles) field experiments.

Fig. S4 | Alpine and lowland plant effects on alpine soil microbial respiration. (a,b) Relationships between plant

physiology, expressed as (a) PC1 and (b) PC2 scores from PCAs of pot-level measurements, and rates of soil microbial

respiration during the first nine days of incubation (initial R; µg C g^{-1}- h^{-1}) in alpine (green) versus lowland (yellow)

treatments of the glasshouse experiment. Fit lines describe relationships as determined by a single linear mixed effects

model including treatment, PC1, PC2 and treatment × PC# interactions (N = 20; Main Text), with different colours in (b)

indicating a significant treatment × PC2 interaction. (c,d) Loadings of plant physiological measurements on (c) PC1 and

(d) PC2 axes. Total, belowground and aboveground biomass are sums of all plants per pot and specific leaf area (SLA),

maximum photosynthetic capacity (A_{max}) and stomatal conductance (g_s) are pot-wise community weighted mean (CWM)

values.

Fig. S5 | Soil inoculation effects on alpine soil community composition. Effects of low elevation soil biota

inoculation on (a) bacterial and (b) fungal community composition in alpine plots (N = 10 in both cases). Axes show the

first two components of principal coordinates analyses (PCoA; Bray Curtis distance) performed on the relative

abundances of bacterial and fungal operational taxonomic units, respectively. *P*-values are from PERMANOVAs testing

for inoculation effects on the same distance matrices.

Supplementary Tables

Table S1. Linear mixed effects model outputs for effects of field treatment (control, warming, warming plus lowland plants), region and their interaction on soil variables. LR: likelihood ratio.

Response	Treatment		Region		Treatment × Region		
	LR	P	LR	P	LR	P	N
Soil carbon content	23.16	< 0.0001	34.53	< 0.0001	4.64	0.0984	58
Net ecosystem CO_2 exchange	23.26	< 0.0001	40.76	< 0.0001	6.43	0.0401	266
Gross primary production	58.99	< 0.0001	50.64	< 0.0001	2.78	0.2491	264
Ecosystem respiration	49.61	< 0.0001	13.08	0.0045	12.98	0.0015	268
Microbial biomass C	33.52	< 0.0001	13.64	0.0034	5.35	0.0688	58
Microbial growth (per gram soil)	9.33	0.0094	-	-	-	-	27
Microbial respiration (per gram soil)	1.20	0.5500	-	-	-	-	28
Microbial growth (biomass-specific)	8.32	0.0156	-	-	-	-	27
Microbial Respiration (biomass-specific)	6.54	0.0381	-	-	-	-	28
Microbial carbon use efficiency	15.68	0.0004	-	-	-	-	58

958 **Table S2. Linear mixed effects model outputs for treatment (alpine plants, lowland plants) effects on plant traits**

959 **and soil variables in the glasshouse experiment. LR: likelihood ratio.**

Response variable	LR	N	*P*
Pot-level plant traits			
Aboveground biomass	4.85	19	0.0277
Belowground biomass	16.31	19	0.0001
Root-to-shoot ratio	7.92	19	0.0049
Specific leaf area (SLA)	0.03	18	0.8605
Maximal photosynthetic capacity (A_{max})	4.98	19	0.0257
Maximal stomatal conductance (g_s)	14.37	20	0.0002
Soil variables			
Soil pore water absorbance (a_{350})	24.84	19	< 0.0001
Soil pore water total fluorescence (F_{tot})	27.72	19	< 0.0001
Soil pore water fluorescence index (FI)	9.55	19	0.0020
Microbial biomass carbon	0.16	19	0.6932
Fast-decaying soil carbon pool size	3.99	16	0.0458
Fast-decaying soil carbon pool decay rate	0.23	16	0.6306
Soil DOM C1 (protein-like)	21.39	19	< 0.0001
Soil DOM C2 (protein-like)	10.42	19	0.0012
Soil DOM C3 (humic-like)	15.65	19	0.0001
Soil DOM C4 (humic-like)	9.60	19	0.0019
Soil DOM C5 (humic-like)	17.39	19	< 0.0001
Soil DOM C6 (fulvic acid-like)	16.50	19	< 0.0001

962   **Table S3. Statistical test outputs for lowland species identity effects on soil carbon loss.**

Explanatory variable	Linear model test statistics		
	F	d.f.	*P*
Western Alps experiment			
Achillea millefolium	0.17	1,4	0.7007
Bellis perennis	0.40	1,4	0.5634
Bromus erectus[*]	-	-	-
Dactylis glomerata[*]	-	-	-
Medicago lupulina[*]	-	-	-
Plantago media	1.19	1,4	0.3364
Salvia pratensis	0.10	1,4	0.7684
Central Alps experiment			
Brachypodium pinnatum	3.70	1,8	0.3053
Carex flacca	28.24	1,8	0.1184
Carum carvi	5.94	1,8	0.2479
Dactylis glomerata[*]	-	-	-
Hypericum perforatum	10.11	1,8	0.1939
Plantago lanceolata[*]	-	-	-
Primula veris	6.41	1,8	0.2395
Ranunculus bulbosus	40.38	1,8	0.0994
Salvia pratensis	0.08	1,8	0.8281
Silene vulgaris	36.12	1,8	0.1050
Trifolium montanum[*]	-	-	-
Viola hirta[*]	-	-	-

964   [*] *NB: species not tested, present in < 3 plots*

Table S4. DOM components identified through PARAFAC modelling.

Name	Excitation λ (nm)	Emission Λ (nm)	General description[†]	Peak name[*]
C1	275	332	Protein-like (tryptophan)	T
C2	270	310	Protein-like (tyrosine)	B
C3	340	434	Humic-like	A/C
C4	305	424	Microbial humic-like	M
C5	375	452	Humic-like	C
C6	< 250 (400)	506	Soil fluvic acid-like	C+

[*] Ref 57; [†] Ref 56