Evolutionary conservation of centriole rotational asymmetry in the human

centrosome

Noémie Gaudin [1], Paula Martin Gil [1], Meriem Boumendjel [1], Dmitry Ershov [2,3], Catherine

Pioche-Durieu [1], Manon Bouix [1], Quentin Delobelle [1], Lucia Maniscalco [1], Thanh Bich Ngan

Phan [1], Vincent Heyer [4,5,6,7], Bernardo Reina-San-Martin [4,5,6,7] and Juliette Azimzadeh [1,8].

[1] Université de Paris, CNRS, Institut Jacques Monod, 75006, Paris, France.

[2] Image Analysis Hub, C2RT, Institut Pasteur, Paris, France.

[3] Hub de Bioinformatique et Biostatistique – Département Biologie Computationnelle, Institut

Pasteur, USR 3756 CNRS, Paris, France.

[4] Institut de Génétique et de Biologie Moléculaire et Cellulaire (IGBMC), Illkirch, France.

[5] Institut National de la Santé et de la Recherche Médicale (INSERM), U1258, Illkirch, France.

[6] Centre National de la Recherche Scientifique (CNRS), UMR7104, Illkirch, France.

[7] Université de Strasbourg, Illkirch, France.

[8] Author for correspondence: juliette.azimzadeh@ijm.fr

Keywords: centriole, centrosome, LRRCC1, VFL1, C2CD3, Joubert syndrome, asymmetry.

Abbreviations:

DA: distal appendage; JBTS: Joubert syndrome; MCC: multiciliated cell; PCM: pericentriolar

material; SAG: SMOOTHENED agonist; SMO: SMOOTHENED; TZ: transition zone.

25   **Abstract**

Centrioles are formed by microtubule triplets in a nine-fold symmetric arrangement. In

flagellated protists and in multiciliated cells, accessory structures tethered to specific triplets

render the centrioles rotationally asymmetric, a property that is key to cytoskeletal and cellular

organization in these contexts. In contrast, centrioles within the centrosome of animal cells

display no conspicuous rotational asymmetry. Here, we uncover rotationally asymmetric

molecular features in human centrioles. Using ultrastructure expansion microscopy, we show

that LRRCC1, the ortholog of a protein originally characterized in flagellate green algae,

associates preferentially to two consecutive triplets in the distal lumen of human centrioles.

LRRCC1 partially co-localizes and affects the recruitment of another distal component,

C2CD3, which also has an asymmetric localization pattern in the centriole lumen. Together,

LRRCC1 and C2CD3 delineate a structure reminiscent of a filamentous density observed by

electron microscopy in flagellates, termed the 'acorn'. Functionally, the depletion of LRRCC1

in human cells induced defects in centriole structure, ciliary assembly and ciliary signaling,

supporting that LRRCC1 cooperates with C2CD3 to organizing the distal region of centrioles.

Since a mutation in the *LRRCC1* gene has been identified in Joubert syndrome patients, this

finding is relevant in the context of human ciliopathies. Taken together, our results demonstrate

that rotational asymmetry is a conserved ancient property of centrioles. Our work also reveals

that asymmetrically localized proteins are key for primary ciliogenesis and ciliary signaling in

human cells.

Introduction

Centrioles are cylindrical structures with a characteristic ninefold symmetry, which results

from the arrangement of their constituent microtubule triplets (LeGuennec et al., 2021). In

animal cells, centrioles are essential for the assembly of centrosomes and cilia. The centrosome,

composed of two centrioles embedded in a pericentriolar material (PCM), is the main organizer

of the microtubule cytoskeleton. In addition, most vertebrate cells possess a primary cilium, a

sensory organelle that assembles from the oldest centriole within the centrosome, called mother

centriole (Kumar and Reiter, 2021).

Centrioles within the centrosome show no apparent rotational asymmetry, *i.e.,* no

structural asymmetry of the microtubule triplets. In vertebrates, the mother centriole carries

distal appendages (DAs) and subdistal appendages arranged in a symmetric manner around the

centriole cylinder (Kumar and Reiter, 2021). In contrast, the centriole/basal body complex of

flagellates, to which the animal centrosome is evolutionary related, is characterized by marked

rotational asymmetries (Azimzadeh, 2021; Yubuki and Leander, 2013). In flagellates, an array

of fibers and microtubules anchored asymmetrically at centrioles controls the spatial

organization of the cell (Feldman et al., 2007; Yubuki and Leander, 2013). The asymmetric

attachment of cytoskeletal elements appears to rely on molecular differences between

microtubule triplets. In the green alga *Chlamydomonas reinhardtii,* Vfl1p (Variable Flagella

number 1 protein) localizes principally at two triplets near the attachment site of a striated fiber

connecting the centrioles (Silflow et al., 2001). This fiber is absent or mispositioned in the *vfl1*

mutant, leading to defects in centriole position and number, and overall cytoskeleton

disorganization (Adams et al., 1985; Feldman et al., 2007). In the same region, a rotationally

asymmetric structure termed the 'acorn' was observed in the centriole lumen by transmission

electron microscopy. The acorn appears as a filament connecting five successive triplets and is

in part colocalized with Vfl1p (Geimer and Melkonian, 2005, 2004).

We recently established that Vfl1p function is conserved in the multiciliated cells (MCCs)

of planarian flatworms, which was later confirmed in xenopus (Basquin et al., 2019; Nommick

et al., 2021). MCCs assemble large numbers of centriole that are polarized in the plane of the

plasma membrane to enable the directional beating of the cilia (Meunier and Azimzadeh,

2016), like in *C. reinhardtii*. The planarian ortholog of Vfl1p is required for the assembly of

two appendages that decorate MCC centrioles asymmetrically, the basal foot and the ciliary

rootlet (Basquin et al., 2019). Depleting Vfl1p orthologs in planarian or xenopus MCCs alters

centriole rotational polarity, reminiscent of the *vfl1* phenotype in *C. reinhardtii* (Adams et al.,

1985; Basquin et al., 2019; Nommick et al., 2021). Intriguingly, the human ortholog of Vfl1p,

called LRRCC1 (Leucine Rich Repeat and Coiled Coil containing 1) localizes at the

centrosome despite the lack of rotationally asymmetric appendage in this organelle (Andersen

et al., 2003; Muto et al., 2008). Furthermore, a homozygous mutation in the *LRRCC1* gene was

identified in two siblings affected by a ciliopathy called Joubert syndrome (JBTS), suggesting

that LRRCC1 might somehow affect the function of non-motile cilia (Shaheen et al., 2016).

Here, we show that LRRCC1 localizes in a rotationally asymmetric manner in the centrioles

of the human centrosome. We further establish that LRRCC1 is required for proper ciliary

assembly and signaling, which likely explains its implication in JBTS. LRRCC1 affects the

recruitment at centrioles of another ciliopathy protein called C2CD3 (C2 domain containing

3), which we found to also localize in a rotationally asymmetric manner, forming a pattern

reminiscent of the acorn described in flagellates. Our findings uncover the unanticipated

rotational asymmetry of centrioles in the human centrosome and show that this property is

connected to the assembly and function of primary cilia.

Results

LRRCC1 localizes asymmetrically at the distal end of centrioles

To investigate a potential role of LRRCC1 at the centrosome, we first sought to determine its

precise localization at the centrosome. We raised antibodies against two different fragments

within the long C-terminal coiled-coil domain of LRRCC1 (Ab1, 2), which both stained the

centrosome region in human Retinal Pigmented Epithelial (RPE1) cells (Fig. 1a; Supplemental

Fig. S1b), as previously reported (Muto et al., 2008). Labeling intensity was decreased in

LRRCC1-depleted cells for both antibodies, supporting their specificity (Fig. 4a; Supplemental

Fig. S1c, e, f, h). LRRCC1 punctate labelling in the centrosomal region indicated that it is

present within centriolar satellites, confirming a previous finding that LRRCC1 interacts with

the satellite component PCM1 (Gupta et al., 2015). Importantly, following nocodazole

treatment to depolymerize microtubules, the satellites were dispersed but a fraction of LRRCC1

was retained at the centrioles (Fig. 1a; Supplemental Fig. S1b, e), supporting that LRRCC1 is

a core centriole component.

To determine LRRCC1 localization more precisely within the centriolar structure, we used

ultrastructure expansion microscopy (U-ExM) (Gambarotto et al., 2019) combined with

imaging on a Zeiss Airyscan 2 confocal microscope, thereby increasing the resolution by a

factor of ~ 8 compared to conventional confocal microscopy. We found that LRRCC1 localizes

at the distal end of centrioles as well as of procentrioles (Fig. 1b). Strikingly, and unlike any

other known centriole components, LRRCC1 decorated the distal end of centrioles in a

rotationally asymmetric manner. Indeed, LRRCC1 was detected close to the triplet blades and

towards the lumen of the centriole (Fig. 1c). The staining was often associated with two or

more consecutive triplets, one of them being usually more brightly labelled than the others. In

addition, a fainter staining was consistently detected along the entire length of all triplets (Fig.

1b, see brighter exposure). This pattern was observed in both RPE1 and HEK 293 cells and

was obtained with both anti-LRRCC1 antibodies (Supplemental Fig. S1g), supporting its

specificity. There was some variability in the distal pattern, particularly in the distance between

LRRCC1 and the centriole wall (Fig. 1c), which possibly resulted from the fact that the

centrioles were not perfectly orthogonal to the imaging plan. To obtain a more accurate picture

of LRRCC1 localization, we generated 3D reconstructions that we realigned, first along the

vertical axis, then with respect to one another using the most intense region of the LRRCC1

labelling as a reference point (Fig. 1d; Supplemental Fig. S2a-c). An average 3D reconstruction

was then generated (Fig. 1e) and revealed that in transverse view, LRRCC1 was mainly

associated to one triplet, and to a lesser extent to its direct neighbor counterclockwise, on their

luminal side. A longitudinal view confirmed that LRRCC1 is principally located at the distal

end of centrioles.

Together, our results show that LRRCC1 is localized asymmetrically within the distal

centriole lumen, establishing that centrioles within the human centrosome are rotationally

asymmetric.

The localization pattern of LRRCC1 is similar at the centrosome and in mouse MCCs

LRRCC1 orthologs are required for establishing centriole rotational polarity in planarian and

xenopus MCCs, like in *C. reinhardtii* (Basquin et al., 2019; Nommick et al., 2021; Silflow et

al., 2001). It is therefore plausible that LRRCC1-related proteins localize asymmetrically in

MCC centrioles, and indeed, Lrrcc1 was recently found associated to the ciliary rootlet in

xenopus MCCs (Nommick et al., 2021). To determine whether LRRCC1 also localizes at the

distal end of MCC centrioles in addition to its rootlet localization, and if so, whether LRRCC1

localization pattern resembles that observed at the centrosome, we analyzed mouse ependymal

and tracheal cells by U-ExM. In *in vitro* differentiated ependymal cells, the labelling generated

by the anti-LRRCC1 antibody was consistent with our observations in human culture cells.

Mouse Lrrcc1 localized asymmetrically at the distal end of centrioles, opposite to the side

where the basal foot is attached (Fig. 2a), as determined by staining the basal foot marker γ-

tubulin (Clare et al., 2014). Lrrcc1 was also present at the distal end of procentrioles forming

via either the centriolar or acentriolar pathways (*i.e.,* around parent centrioles or deuterosomes,

respectively) (Fig. 2b). We also examined tracheal explants, in which centrioles were docked

and polarized at the apical membrane in higher proportions (Fig. 2c). We obtained an average

image of Lrrcc1 labelling from 35 individual centrioles aligned using the position of the basal

foot as a reference point. This revealed that Lrrcc1 is principally located in the vicinity of 3

triplets located opposite the basal foot, to the right of basal foot main axis (triplet number 9, 1

and 2 on the diagram in Fig. 2d). Lrrcc1 was located farther away from the triplet wall than in

centrioles of the centrosome, but this was likely an effect of a deformation of the centrioles

(Fig. 2c, d) caused by the incomplete expansion of the underlying cartilage layer in tracheal

explants. In agreement, Lrrcc1 was located close to the triplets in ependymal cell monolayers,

which expand isometrically. Besides the distal centriole staining, we found no evidence that

Lrrcc1 is associated to the ciliary rootlet in mouse MCCs, unlike in xenopus. The Lrrcc1 pattern

in mouse MCCs was thus similar to the pattern observed at the human centrosome.

Together, these results show that LRRCC1 asymmetric localization is a conserved feature

of mammalian centrioles, presumably linked to the control of centriole rotational polarity and

ciliary beat direction in MCCs.

Centriole rotational asymmetry is not linked to stereotypical patterns of procentriole

position and orientation during centriole duplication

In *C. reinhardtii,* centriole rotational polarity and flagellar beat direction depend on the position

and orientation at which new centrioles arise during cell division. Reflecting the stereotypical

organization of centrioles and procentrioles in this species, Vfl1p is recruited early and at a

fixed position at the distal end of procentrioles (Fig.3a) (Geimer and Melkonian, 2004; Silflow

et al., 2001). We therefore wondered whether this mechanism might be somewhat conserved

in human cells, which could then explain the persistence of LRRCC1 asymmetric localization

despite the lack of asymmetric appendages or ciliary motility in these cells. We first analyzed

the timing of LRRCC1 incorporation into procentrioles. LRRCC1 was already present at an

early stage of centriole assembly, when the procentrioles were only about 100 nm in length and

70 nm in diameter (Fig. 3b, c). LRRCC1 was then detected during successive stages of

procentriole elongation, always localizing asymmetrically and distally (Fig. 3c), like in *C.*

reinhardtii. We then examined LRRCC1 localization in duplicating centrosomes by generating

3D-reconstructions of diplosomes (*i.e.,* orthogonal centriole pairs) from RPE1 and HEK 293

cells processed by U-ExM (Fig. 3d). We analyzed two parameters: the angle between LRRCC1

in the procentriole and the long axis of the parent centriole used as reference (Fig. 3d, LRRCC1

localization in procentrioles), and the angle between procentriole position and LRRCC1 in the

parent centriole (Fig. 3d, Procentriole position with respect to centriolar LRRCC1). We found

that LRRCC1 localization in procentrioles was more often aligned with the long axis of the

parent centriole in RPE1 cells (Fig. 3d, top left panel, quadrants Q1 and Q3, respectively), but

less so in HEK 293 cells (top right panel), in which the distribution was closer to a random

distribution. Thus, human procentrioles do not arise in a fixed orientation, unlike in flagellates.

Next, we analyzed the position of procentrioles with respect to centriolar LRRCC1 (bottom

panels). Based on current models, procentriole assembly is expected to occur at a random

position around parent centrioles in animal cells (Takao et al., 2019). Identification of LRRCC1

provided the first opportunity to directly test this model. In diplosomes from both RPE1 and

HEK 293 cells, the position of procentrioles with respect to LRRCC1 location in the parent

centriole was variable, confirming that the position at which procentrioles assemble is not

precisely controlled in human cells. Interestingly, however, the procentrioles were not

distributed in a completely random fashion either. Procentrioles were found in quadrant Q2

(45-135 degrees clockwise from LRRCC1 centroid) on average 4 times less often than in the

other quadrants, both in RPE1 and HEK 293 cells, suggesting that rotational polarity of the

parent centriole somehow impacts procentriole assembly.

Overall, these results nevertheless suggest that the conservation of centriole rotational

asymmetry in human cells is likely not linked to a role in centriole duplication.

LRRCC1 is required for primary cilium assembly and ciliary signaling

A previous report identified a homozygous mutation in a splice acceptor site of the *LRRCC1*

gene in two siblings diagnosed with JBTS (Shaheen et al., 2016), but how disruption of

LRRCC1 expression affects ciliary assembly and signaling has never been investigated. To

address this, we generated RPE1 cell lines deficient in LRRCC1 using two different

CRISPR/Cas9 strategies and targeting two different regions of the *LRRCC1* locus. We could

not recover null clones despite repeated attempts in RPE1 - both wild type and p53[-/-] (Izquierdo

et al., 2014), HEK 293 and U2-OS cells, suggesting that a complete lack of LRRCC1 is possibly

deleterious. Nevertheless, we obtained partially depleted mutant clones, including three RPE1

clones targeted in either exons 11-12 (clone 1.1) or exons 8-9 (clones 1.2 and 1.9). We could

not evaluate the overall decrease in LRRCC1 levels since the endogenous LRRCC1 protein

was not detected by Western blot (Supplemental Fig. S1a). However, we confirmed the

decrease in centrosomal LRRCC1 levels by immunofluorescence using the two different anti-

LRRCC1 antibodies, as well as by quantitative RT-PCR (Fig. 4a; Supplemental Fig. S1c-e).

Following induction of ciliogenesis, the proportion of ciliated cells was decreased in all

three mutant clones compared to control cells (Fig. 4b). We were unable to obtain stable RPE1

cell lines expressing myc- or GFP-tagged versions of LRRCC1, and transient overexpression

of LRRCC1 in wild-type cells led to a decrease in the proportion of ciliated cells, making

phenotype rescue experiments difficult to interpret. We therefore used RNAi as an independent

method to verify the specificity of ciliary defects observed in CRISPR clones. Targeting

LRRCC1 expression using two different siRNAs, we obtained a comparable decrease in

LRRCC1 levels and in the proportion of ciliated cells as in the CRISPR clones (Fig. 4c;

Supplemental Fig. S1f). Sensory ciliopathies like JBTS result to a large extent from defective

Hedgehog signaling (Romani et al., 2013). We determined the effect of LRRCC1-depletion on

Hedgehog signaling by measuring the ciliary accumulation of the activator SMOOTHENED

(SMO) upon induction of the pathway (Rohatgi et al., 2007). Depletion of LRRCC1 by either

CRISPR editing or RNAi led to a drastic decrease in SMO accumulation at the primary cilium

following induction of the Hedgehog pathway by SMO-agonist (SAG) (Fig. 4d, e), and reduced

expression of the target gene *PTCH1* (Supplemental Fig. S1g) (Goodrich et al., 1996).

Taken together, our results demonstrate that LRRCC1 is required for proper ciliary

assembly and signaling in human cells, further establishing its implication in JBTS.

Depletion of LRRCC1 induces defects in centriole structure

Mutations in distal centriole components can alter centriole length regulation or the assembly

of DAs, which both result in defective ciliogenesis (Reiter and Leroux, 2017; Sharma et al.,

2021). We used U-ExM to search for possible defects in centriole structure in LRRCC1-

depleted RPE1 cells. We measured centriole length in the two RPE1 CRISPR clones that

express the lowest levels of LRRCC1 (1.1 and 1.9). Centrioles were co-stained with anti-

acetylated tubulin and an antibody against the DA component CEP164 to differentiate mother

and daughter centrioles. We observed an increase in centriole length in clone 1.9 (Fig. 5a),

which expresses the lowest levels of LRRCC1, compared to control cells (483 ± 53 nm for

mother and 372 ± 55 nm for daughter centrioles in clone 1.9; 427 ± 56 nm for mother and 320

± 46 nm for daughter centrioles in control cells; mean ± SD). Although on a limited sample

size, we also observed abnormally long centrioles by transmission electron microscopy in this

clone (494 ± 73 nm in clone 1.9, N = 9; 429 ± 52 nm in control cells, N = 3; mean ± SD) (Fig.

5b). However, clone 1.1 was not significantly different from control cells by U-ExM,

suggesting that this phenotype is not fully penetrant at the remaining levels of LRRCC1

expression observed in the CRISPR clones. We next used U-ExM to analyze the structure of

DAs labelled with anti-CEP164 antibody (Fig. 5c-e). In RPE1 control cells, 80 ± 14 % of

mother centriole had 9 properly shaped DAs, but this proportion fell to 57 ± 16 % and 44 ± 17

% (mean ± SD) in clones 1.1 and 1.9, respectively (Fig. 5c, d). Mutant clones exhibited an

increased proportion of centrioles with one or more abnormally shaped DAs (29 ± 17 % and

42 ± 18 % in clones 1.1 and 1.9, respectively, compared to 11 ± 11 % in control cells; mean ±

SD). We obtained similar results in a HEK 293 CRISPR clone expressing half the control levels

of LRRCC1 (Fig. 5e; Supplemental Fig. S1h). LRRCC1-depletion did not affect overall

CEP164 levels at mother centrioles in the CRISPR clones (Supplemental Fig. S3a, d),

consistent with the relatively mild defect in DA morphology observed by U-ExM.

Together, these results show that a partial loss of LRRCC1 leads to abnormal DA

morphology.

LRRCC1 and C2CD3 delineate a rotationally asymmetric structure in human centrioles

We next wanted to determine whether LRRCC1 cooperates with other distal centriole

components. Proteins shown to be recruited early at procentriole distal end include CEP290

(Kim et al., 2008), OFD1 (Singla et al., 2010) and C2CD3 (Thauvin-Robinet et al., 2014). Of

particular interest, OFD1 and C2CD3 are required for DA assembly and centriole length

control, and mutations in these proteins have been implicated in sensory ciliopathies (Singla et

al., 2010; Thauvin-Robinet et al., 2014; Tsai et al., 2019; Lei Wang et al., 2018). We first

determined whether depletion of LRRCC1 either by CRISPR editing or by RNAi led to

modifications in the recruitment of these proteins within centrioles. We found no major

differences in the centrosomal levels of OFD1 and CEP290 compared to control cells

(Supplemental Fig. S3b, c, e, f). In contrast, C2CD3 levels were moderately increased in cells

depleted from LRRCC1 either by CRISPR editing (clones 1.1 and 1.9) or by RNAi (Fig. 6a,

b). We thus analyzed C2CD3 further by U-ExM. As described previously, C2CD3 localized

principally at the distal extremity of centrioles (Fig. 6c) (Tsai et al., 2019; Yang et al., 2018).

Strikingly, the C2CD3 labelling was also asymmetric, both in terms of shape, often a C-shape,

and position within the centriole lumen (Fig. 5d). We next generated an average 3D

reconstruction of the C2CD3 pattern using the region where the C2CD3 signal is closest to the

centriole wall (after realignment along the vertical axis) as a reference point. This confirmed

that C2CD3 forms a C-shaped pattern – or possibly a ring-shaped pattern of uneven intensity,

positioned asymmetrically in the centriole lumen (Fig. 6e). To determine whether this

localization pattern is affected by LRRCC1-depletion, we analyzed C2CD3 in LRRCC1

CRISPR clones 1.1 and 1.9. The C2CD3 pattern was more variable than in control RPE1 cells,

and was often abnormal in shape, position, or both (Fig. 6f). As a result, averaging the signal

from 47 (1.1) or 24 (1.9) individual LRRCC1-depleted centrioles as previously produced

aberrant patterns, most strikingly in the clone expressing lower levels of LRRCC1 (clone 1.9;

Fig. 6g). Thus, LRRCC1 is required for the proper shape and asymmetric positioning of the

C2CD3-containing distal structure.

To determine whether LRRCC1 and C2CD3 might belong to a common structure, we next

examined the respective positions of C2CD3 and LRRCC1 within the centriole. We co-stained

centrioles with our anti-LRRCC1 antibody and a second anti-C2CD3 antibody produced in

sheep (Table 1). We confirmed that LRRCC1 and C2CD3 are present in the same distal region

of the centriole (Fig. 7a). In transverse views, the two proteins were usually not perfectly

colocalized but found in close vicinity of one another near the microtubule wall (Fig. 6e).

However, C2CD3 distal staining was consistently fainter than with the previous antibody, and

we either could not observe a full C-shaped pattern, or we could not image it due to

fluorescence bleaching. Neither anti-C2CD3 antibodies worked in mouse, so we were not able

to compare C2cd3 and Lrrcc1 localization in MCCs. Nevertheless, the results obtained by

individually labelling LRRCC1 and C2CD3 at the centrosome (Fig. 1e, 6e, g) together with the

co-localization data (Fig. 7a) are consistent with the hypothesis that LRRCC1 is located where

the C2CD3-containing structure is closer to the centriole wall (Fig. 7b). C2CD3 was not co-

immunoprecipitated with a GFP-LRRCC1 fusion protein however, suggesting that LRRCC1

and C2CD3 do not directly interact (Supplemental Fig. S4).

Taken together, our results support that C2CD3 localizes asymmetrically in the distal

lumen of human centrioles, a pattern that depends in part on LRRCC1.

Discussion

Here, we show that centrioles within the human centrosome are rotationally asymmetric despite

the apparent nine-fold symmetry of their ultrastructure. This asymmetry is manifested by a

specific enrichment in LRRCC1 near two consecutive triplets, and the asymmetric localization

of C2CD3 within the centriole lumen. Depletion of LRRCC1 perturbed the asymmetric

recruitment of C2CD3 and induced defects in centriole structure, ciliogenesis and ciliary

signaling, supporting that LRRCC1 contributes to organizing the distal centriole region

together with C2CD3. LRRCC1 localizes like its *C. reinhardtii* ortholog Vfl1p*,* and C2CD3

delineates a filamentous structure reminiscent of the acorn described in this species (Geimer

and Melkonian, 2005, 2004) and in *Trypanosoma brucei* (Vaughan and Gull, 2016).

Collectively, our results support that rotational asymmetry is a conserved property of centrioles

linked to ciliary assembly and signaling in humans.

LRRCC1 and C2CD3 belong to a conserved rotationally asymmetric structure

Our work identifies two proteins located asymmetrically in the distal centriole lumen of the

human centrosome, each with a specific pattern. LRRCC1 localizes principally near two

consecutive triplets, with the first triplet counterclockwise labelled approximately 50 % more

than the next one. This pattern is highly reminiscent of the LRRCC1 ortholog Vfl1p, which

localizes predominantly to the triplet facing the second centriole (referred to as triplet 1), and

to a lesser extent to its immediate neighbor counterclockwise (triplet 2; Fig. 7b) (Silflow et al.,

2001). In *C. reinhardtii*, triplets 1 and 2 are positioned directly opposite to the direction of

flagellar beat, which is directed towards triplet 6 (Fig. 7b) (Lin et al., 2012). In mouse MCCs,

Lrrcc1 is associated to triplets located not exactly opposite to the basal foot but with a clockwise

shift of at least 20° from the basal foot axis. However, the beating direction was shown to be

also shifted approximately 20° clockwise relative to the position of the basal foot in bovine

tracheal MCCs (Schneiter et al., 2021) (Fig. 2d). The position of Lrrcc1/Vfl1p-labelled triplets

with respect to ciliary beat direction might thus be similar in *C. reinhardtii* and in animal

MCCs. Overall, the specific localization pattern of Vfl1p-related proteins at the distal end of

centrioles, and their requirement for centriole positioning and ciliary beat orientation when

motile cilia are present, appear to be conserved between flagellates and animals.

The second protein conferring rotational asymmetry to human centrioles, C2CD3, delineates

a C-shape positioned asymmetrically in the lumen. The localization of C2CD3 had been studied

before, including by super-resolution microscopy approaches, without this asymmetry being

noted (Tsai et al., 2019; Yang et al., 2018). This difference might result from the fact that we

corrected the vertical alignment of the centrioles prior to averaging multiple images. Several

lines of evidence support the hypothesis that the C2CD3-containing structure is homologous

to the acorn, a 10 nm-filament connecting five consecutive triplets along the centriole wall and

across the lumen in *C. reinhardtii* (Fig. 7b) (Geimer and Melkonian, 2004; Vaughan and Gull,

2016). First, the C2CD3 labelling is consistent with a circular filament. Second, the C2CD3-

containing structure is localized asymmetrically in the distal lumen in close vicinity of at least

three consecutive triplets. Third, C2CD3 is partially co-localized with LRRCC1 near the

microtubule wall. And last, C2CD3 orthologs are found in a variety of flagellated unicellular

eukaryotes, including the green alga *Micromonas pusilla* (Zhang and Aravind, 2012),

suggesting an ancestral association to centrioles and cilia. The partial co-localization of Vfl1p

and the acorn in *C. reinhardtii*, and the observation that both are already present at the distal

end of procentrioles, led to propose that Vfl1p might also be a component of the acorn (Geimer

and Melkonian, 2004). Consistent with this idea, both LRRCC1 and C2CD3 are recruited early

at the distal end of human procentrioles, and LRRCC1 is required for the proper recruitment

and/or positioning of the C2CD3-containing structure. C2CD3 recruitment at the centrioles

also depends on the proteins CEP120 and Talpid3 (Tsai et al., 2019), which possibly explains

why C2CD3 recruitment is not diminished, and is even increased in LRRCC1-depleted

centrioles. Future work will help deciphering the relationships between these different proteins

and characterize in more details the architecture of the rotationally asymmetric structure at the

distal end of mammalian centrioles.

Rotationally asymmetric centriole components are required for ciliogenesis

Our results uncover a link between centriole rotational asymmetry and primary ciliogenesis in

human cells. Mutations in C2CD3 have been involved in several sensory ciliopathies, including

JBTS (Boczek et al., 2018; Cortés et al., 2016; Ooi, 2015; Thauvin-Robinet et al., 2014). The

associated ciliary defects are likely caused by anomalies in the structure of centrioles, since

depleting C2CD3 inhibits centriole elongation and DA assembly, whereas C2CD3

overexpression leads to centriole hyper-elongation (Thauvin-Robinet et al., 2014; Wang et al.,

2018; Ye et al., 2014). We observed similar defects in LRRCC1-depleted cells, but of

comparatively lesser extent. Defects in DA morphology were observed in all CRISPR clones

analyzed, and centriole length was increased in the clone with the most severe phenotype. The

fact that LRRCC1-depletion has a more limited impact on centriole assembly than perturbation

of C2CD3 levels suggests that LRRCC1 might not be directly involved in centriole length

control or DA formation, however. The defects observed in LRRCC1-depleted cells could

instead result indirectly from the abnormal localization of C2CD3. Besides the defects in

centriole structure, it is likely that LRRCC1-depletion also perturbs the organization of the

ciliary gate, as LRRCC1-depleted cells exhibited a drastic reduction in Hedgehog signaling.

Loss of ciliary gate integrity interferes with the accumulation of SMO in the cilium upon

activation of the Hedgehog pathway and is a frequent consequence of ciliopathic mutations

(Garcia-Gonzalo and Reiter, 2017). The ciliary gate consists of the TZ and the DA region,

which both contribute to regulating protein trafficking in and out of the cilium (Garcia-Gonzalo

and Reiter, 2017; Nachury, 2018). The anomalies in DA morphology observed in LRRCC1-

depleted cells could disrupt the organization of the so-called DA matrix (Yang et al., 2018),

thus preventing SMO accumulation in the cilium. Another, non-mutually exclusive possibility

is that the architecture of the TZ, which forms directly in contact with the distal end of the

centriole, is altered by LRRCC1-depletion. In either case, our observations in RPE1 cells are

consistent with the JBTS diagnosis in two siblings carrying a mutation in the *LRRCC1* gene

(Shaheen et al., 2016), further establishing that *LRRCC1* is a novel ciliopathy gene. Besides

signaling, ciliary gate integrity is required for axoneme extension and indeed, LRRCC1-

depleted cells formed cilia at lower frequency than control cells – a defect that might also result

from perturbed DA architecture. In the *vfl1* mutant of *C. reinhardtii*, both unanchored

centrioles and centriole docked at the plasma membrane but lacking a flagellum were observed

(Adams et al., 1985). This would support that LRRCC1/Vfl1p requirement for properly

assembling the ciliary gate is a conserved functional aspect of this family of proteins (Fig. 7c).

Why is there a rotationally asymmetric structure at the base of primary cilia, and how does

this structure form and contribute to the assembly of the DAs and the cilium remain open

questions. In *C. reinhardtii* and in MCCs, LRRCC1 function is linked to the assembly of

asymmetric appendages, which must be correctly positioned in relation to ciliary beat direction

(Fig. 7c). An asymmetric structure present early during centriole assembly and ultimately

located near the cilium appears well suited for this task. The conservation of such a structure

at the base of the primary cilium could perhaps indicate that primary cilia also possess

rotationally asymmetric features, which would open interesting new perspectives on ciliary

roles in heath and disease.

Other roles for centriole rotational assembly in human cells

Our finding that procentrioles do not form completely at random with respect to LRRCC1 location in the parent centriole suggests that centriole rotational polarity can influence centriole duplication in human cells. In *C. reinhartdtii*, procentrioles are formed at fixed positions with respect to the parent centrioles, to which they are bound by a complex array of fibrous and microtubular roots (Fig. 7c) (Geimer and Melkonian, 2004; Yubuki and Leander, 2013). The process is likely different at the centrosome since the roots typical of flagellates are not conserved in animal cells (Azimzadeh, 2021; Yubuki and Leander, 2013). In mammalian cells, procentrioles form near the wall of the parent centriole following the recruitment of early centriole proteins directly to the PCM components CEP152 and CEP192 (Yamamoto and Kitagawa, 2021). It is nonetheless conceivable that an asymmetry in triplet composition could result in local changes in PCM composition, which in turn could negatively impact PLK4 activation in this region. For instance, our analyses in planarian MCCs led us to postulate that linkers might be tethered to one side of the centrioles in a VFL1-dependent manner and independently of centriole appendages (Basquin et al., 2019). Future work will allow deciphering how centriole rotational asymmetry influences centriole duplication, and whether it affects other aspects of centriole positioning and cellular organization.

Acknowledgements

The authors are deeply grateful to Marine Laporte, Virginie Hamel, Paul Guichard and Davide Gambarotto for teaching them the U-ExM procedure and for sharing antibodies; Arnaud Echard and Takashi Ochi for critical reading of the manuscript; Amélie-Rose Boudjema and Alice Meunier for providing mouse ependymal cells and Isabelle Le Parco for the tracheal tissue; Juliane Da Graça and Simon Herman for technical help; Rémi Le Borgne for help with transmission electron microscopy and for critical reading of the manuscript. We acknowledge

the core imaging facility of Institut Jacques Monod (ImagoSeine facility, member of the France

BioImaging infrastructure supported by grant ANR-10-INBS-04 from the French National

Research Agency). This work was supported by La Ligue Contre le Cancer and Fondation ARC

pour la recherche sur le cancer funding to J.A. N.G. was recipient of a MESRI PhD fellowship

from the French Government and a 4th year PhD fellowship from the Fondation pour la

Recherche Médicale.

Declaration of interests

The authors declare no competing interests.

Material and Methods

Cell culture

RPE1 cells (hTERT-RPE1, RRID:CVCL_4388) were cultured in DMEM/F-12 medium (ThermoFisher Scientific) supplemented with 10 % fetal calf serum (ThermoFisher Scientific), 100 U/mL penicillin and 100 μg/mL streptomycin (ThermoFisher Scientific). Ciliogenesis was induced by culturing RPE1 cells in medium without serum during 48 hours. HEK 293 cells (kind gift from F. Causeret, Institut Imagine, Paris) were cultured in DMEM medium (ThermoFisher Scientific) supplemented with 10 % fetal calf serum and antibiotics as previously. All cells were kept at 37°C in the presence of 5 % CO_2.

Mouse ependymal cells and tracheal tissue

All experiments were performed in accordance with French Agricultural Ministry and European guidelines for the care and use of laboratory animals. *In vitro* differentiated ependymal cells were a kind gift from A.R. Boudjema and A. Meunier (IBENS, Paris). They were prepared as described previously (Delgehyr et al., 2015; Mercey et al., 2019) from Cen2GFP mice (CB6-Tg(CAG-EGFP/CETN2)3-4Jgg/J, The Jackson Laboratory). The fragment of trachea was obtained from a wild-type mouse of the Swiss background (kindly provided by I. Le Parco, Institut Jacques Monod).

CRISPR/Cas9 editing

LRRCC1 mutant clones were obtained by two different CRISPR/Cas9 strategies. First, RPE1 cells were co-transfected with plasmid px154-1 (U6p-gRNA#1_U6p-gRNA#2_CMVpnCas9-EGFP_SV40p-PuroR-pA with gRNA#1: 5'- AGA ATT CTA CCC TAC CTG CGG - 3' and gRNA#2: 5'- TAA GGT AGT GCT TCC TAC CGG -3') targeting the *LRRCC1* locus in exon

8, and px155-24 (U6p-gRNA#3_U6p-gRNA#4_CMVpnCas9-mCherry_SV40p-PuroR-pA;

gRNA#1: 5'- ATC TAC TCG GAA AGC TGA CGG -3' and 5'- GCT TGA GGG CTC AAA

TAC CGG - 3') targeting exon 9. Both constructs express the nickase mutant of Cas9 fused to

either EGFP or mCherry. Two days after transfection, EGFP- and mCherry-positive cells were

sorted by flow cytometry and grown at low concentration. Individual clones were picked after

2 weeks and analyzed by PCR to detect short insertions/deletions. A single clone was obtained

(clone 1.1), which was further characterized by genomic sequencing. Both alleles of *LRRCC1*

contained deletions (~ 0.6 kb deletion of exon 9 and a ~1.5 kb deletion of exon 8) leading to

frameshifts. In a second approach, cells were co-transfected using a mix of 3 CRISPR/Cas9

Knockout Plasmids (sc-413781; Santa Cruz Biotechnology) targeting exons 11 (5'- CTT GTT

CTC TTT CTC GAT GAA GG – 3' and 5' - ACT TCT TGC ATT GAA AGA ACA GG - 3')

or 12 (5' - CGT GTT AAG CCA GCA GTA TAT GG – 3') of *LRRCC1,* together with the

corresponding Homology Directed Repair plasmids carrying a puromycine-resistance cassette

(sc-413781-HDR; Santa Cruz Biotechnology), following the recommendations of the

manufacturer. Mutant clones were selected by addition of 2 µg/mL puromycine in the culture

medium and further screened by immunofluorescence.

487   Inducible HEK 293 cell lines

LRRCC1 full-length coding sequence was amplified from cDNA clone IMAGE:5272572

(Genbank accession: BC070092.1), corresponding to the longest isoform of LRRCC1

(NM_033402.5), after correction of a frameshift error by PCR mutagenesis. As N- and C-

terminal GFP fusions were not targeted to the centrosome, we inserted the GFP tag within the

LRRCC1 sequence in disordered regions present between the leucine rich repeat and coiled-

coil domains, either after amino acid 251 or 402. The fusions were cloned into the pCDNA-

5FRT (ThermoFischer Scientific) vector using the Gibson assembly method (Gibson et al.,

2009) and then integrated into the Flp-In-293 cell line using the Flp-In system (ThermoFischer

Scientific). Expression of the GFP-LRRCC1 fusions was induced by culturing the Flp-In-293

cell lines overnight in medium supplemented with 1 µ/mL doxycycline (ThermoFischer

Scientific).

RNAi

Ready to use double-stranded siRNA LRRCC1-si1 (target sequence: 5'- AAG GAG AAA

GAT GGA GAC GAT - 3') (Muto et al., 2008), LRRCC1-si2 (target sequence: 5'- TTA GAT

GAC CAA ATT CTA CAA - 3') and control siRNA (AllStars Negative Control) were

purchased from Qiagen. siRNAs were delivered into cells using Lipofectamine RNAiMAX

diluted in OptiMEM medium (ThermoFisher Scientific). Cells were fixed after 48 hours and

processed for immunofluorescence. For RNAi-depletion of ciliated cells, RPE1 cells grown in

complete culture medium were treated by RNAi, incubated for 2 days, then submitted to a

second round of RNAi. After 8 hours, cells were washed 3 x in PBS then cultured during 24

hours in serum-free medium to induce ciliogenesis.

qRT-PCR

Total RNA extracts were obtained using the Nucleospin RNA kit (Macherey-Nagel) and

cDNAs were synthetized using SuperScript III reverse transcriptase (Thermofisher Scientific).

qPCR was performed with the GoTaq qPCR Master Mix (Promega) in a LightCycler 480

instrument (Roche) using the primers listed in Table 2. Quantification of relative mRNA levels

was performed using CHMP2A and EMC7 as reference genes following the MIQE guidelines

(Bustin et al., 2009).

Antibodies

Fragments encoding either aa 671-805 (Ab1) or aa 961-1032 (Ab2) of LRRCC1 (NP_208325.3) were cloned in pGST-Parallel1 and expressed in *Escherichia coli.* The GST-fusion proteins were purified under native conditions using glutathione agarose (ThermoFisher Scientific) and the LRRCC1 fragments were recovered by Tev protease cleavage and dialyzed before rabbit immunization (Covalab). Antibodies were affinity-purified over the corresponding GST-LRRCC1 fusion bound to Affi-Gel 10 resin (Bio-Rad). Other primary and secondary antibodies used in this study are listed in Table 1.

Western blot

For whole-cell extracts, Flp-In-293 cell lines expressing the GFP-LRRCC1 fusions were induced overnight with doxycycline, collected by centrifugation, and resuspended in Western blot sample buffer prior to incubation at 95 °C for 5 minutes. For immunoprecipitation experiments, doxycycline-induced cells expressing LRRCC1 with a GFP inserted after aa 402 were resuspended in lysis buffer (50 mM Tris pH 8, 150 mM NaCl, 1 % NP-40, 0.5 % sodium deoxycholate, 0.1 % SDS) supplemented with 1 mM $MgCl_2$, 20 µg/mL DNAse I (Roche) and a protease inhibitor cocktail (Complete mini, EDTA-free, Roche). After 30 minutes on ice, the lysates were centrifuged at 15 000 g for 10 minutes at 4°C. The supernatants were then incubated with Dynabeads M-280 sheep-anti rabbit magnetic beads (ThermoFischer Scientific) previously incubated with rabbit anti-IgGs, either anti-GFP or anti-HA tag for the control IP (Table 1), and rotated for 3 hours at 4°C. After 3 washes with lysis buffer, immunoprecipitated proteins were recovered by resuspending the beads in sample buffer and heating at 95 °C for 5 minutes. The samples were then run on 4-20% Mini-Protean TGX precast protein gels (Bio-Rad) and transferred onto PVDF membrane using the iBlot 2 blot system (ThermoFischer Scientific). The membranes were blocked and incubated with antibodies following standard

procedures, then visualized using Pierce ECL plus chemiluminescence reagents

(ThermoFischer Scientific) on an ImageQuant LAS 4000 camera system (GE Healthcare Life

Sciences).

Immunofluorescence

Cells were fixed in cold methanol for 5 minutes at – 20 °C, blocked 10 minutes with 3 % BSA

(Sigma Aldrich) in PBS containing 0.05 % Tween-20 (PBST-0.05%), then incubated with

primary antibodies diluted in PBST-0.05% containing 3 % BSA for 1 hour. After washing 3 x

1 minute in PBST-0.05%, cells were incubated 2 hours with secondary antibodies in PBST-

0.05% containing 3 % BSA and 5 µg/mL Hoechst 33342 (ThermoFischer Scientific), washed

in PBST-0.05% as previously, and mounted using Fluorescence Mounting Medium (Agilent).

For staining of primary cilia with anti-acetylated tubulin, cells were incubated 2 hours on ice

prior to methanol fixation. For quantification of SMO accumulation within cilia, confluent cells

cultured during 24 hours in serum-free medium were supplemented with 200 nM SAG (Sigma)

diluted in DMSO, or DMSO alone for 24 hours. Cells were then co-stained for SMO and

ARL13B to determine the position of the primary cilium. For all experiments involving

induction of ciliogenesis by serum deprivation, we verified that cells were arrested in G0 by

immunofluorescence staining of Ki67. To visualize centriolar LRRCC1, and to quantify

CEP290 centrosomal levels, cells were treated during 1 hour with 5 µM nocodazole prior to

fixation. Images were acquired using an Axio Observer Z.1 microscope (Zeiss) equipped with

a sCMOS Orca Flash4 LT camera (Hamamatsu) and a 63x objective (Plan Apo, N.A. 1.4). The

structured illumination microscopy (SIM) image was acquired on an ELYRA PS.1 (Zeiss)

equipped with an EMCCD iXon 885 camera (Andor) and a 63x objective (Plan Apo, N.A. 1.4).

Ultrastructure expansion microscopy

We used the U-ExM protocol described in (Gambarotto et al., 2019) with slight modifications.

Cells grown on glass coverslips were incubated in a fresh solution of 1 % acrylamide and 0.7

% formaldehyde diluted in PBS. After incubating 5 hours to overnight at 37 °C, the coverslips

were washed with PBS and placed cells down on a drop of 35 μL monomer solution (19.3 %

sodium acrylate, 10 % acrylamide, 0.1 % bis-acrylamide in PBS) to which 0.5 % TEMED and

0.1 % ammonium persulfate were added just before use. The coverslips were incubated 5

minutes on ice then 1 hour at 37°C, then transferred to denaturation buffer (200 mM SDS, 200

mM NaCl, 50 mM Tris pH9) for 15 minutes with agitation to detach the gels from the

coverslips. The gels were then incubated in denaturation buffer 1.5 hours at 95 °C, washed 2 x

30 minutes in deionized water then incubated overnight in water at room temperature to allow

expansion of the gel. The gels were measured at this step to determine the coefficient of

expansion. After 2 x 10 minutes in PBS, the gels were cut into smaller pieces then incubated 3

hours at 37 °C with primary antibodies diluted in saturation buffer (3 % BSA, 0.05 % Tween-

20 in PBS). The gel fragments were then washed 3 x 10 minutes in PBST-0.1%, incubated 3 h

with secondary antibodies and washed in PBST-0.1% as previously. Finally, the gels were

incubated 2 x 30 minutes in de-ionized water then left to expand overnight in de-ionized water

to regain their maximum size. For U-ExM of mouse tracheal cells, a fragment of WT mouse

trachea (kind gift from I. Le Parco, IJM, Paris) was adhered on a poly-lysine coated coverslip

then processed as described above with the following modifications: for the first step, the

fragment of trachea was incubated overnight to 48 hours in 1 % acrylamide and 0.7 %

formaldehyde in PBS; they were placed 15 min on ice prior to the 1-hour incubation at 37 °C

and the transfer to denaturation buffer. Note that GFP fluorescence was quenched during U-

ExM processing, so the GFP-Cen2 construct expressed in ependymal cells was not detectable

in final samples. Gels were imaged on Lab-Tek chamber slides (0.15 mm) coated with poly-

lysine (ThermoFisher Scientific). Images were acquired at room temperature using either a

LSM780 confocal microscope (Zeiss) equipped with an oil 63x objective (Plan Apo, N.A. 1.4),

or an LSM980 confocal microscope with Airyscan 2 (Zeiss) equipped with an oil 63x objective

(Plan Apo, N.A. 1.4).

Image analysis

Protein levels were determined using ImageJ software (Schneider et al., 2012) by measuring

the fluorescence intensity in the centrosome or cilium area and subtracting the cytoplasmic

background in z-series taken at 0.5-μm interval.

To generate average images of LRRCC1 and C2CD3, U-ExM samples were systematically

imaged using Airyscan 2. Only centrioles that were nearly perpendicular to the imaging plane

were acquired in order to maximize the resolution in transverse views. Calculating the average

image consisted of several steps: cropping out individual centrioles, aligning them, providing

reference points, standardizing centrioles using the reference points, and averaging

(Supplemental Fig. S2). The cropping was done in ImageJ, and for aligning and providing the

reference points a graphical user interface was developed based on Napari (Sofroniew et al.,

2020). Centriole alignment: the direction of centriole long axis was selected manually and used

to position the centriole vertically. Providing the reference points: 9 points were manually

selected to outline the circle of microtubules triplets and the location of the protein of interest.

The centriole was also framed in Z dimension with a rectangle. Standardization: the reference

points were used to calculate all necessary transformations (rotation, scaling and translation)

to map the original image of a centriole to the standard image. Averaging: an average image

was calculated for all the successive XY planes of the standardized image stacks. For alignment

of tracheal cell centrioles, since the current version of the graphical user interface can only

accommodate two channels, the position of the basal foot provided by the γ-tubulin channel

was reported manually in the acetylated tubulin channel using Image J. The images were then

processed as before using the manual annotation as a reference point for the basal foot.

For analysis of procentriole position and LRRCC1 location in procentrioles, 3D-

reconstructions of diplosomes processed for U-ExM were obtained using IMARIS software

(Oxford Instruments).

Electron microscopy

RPE1 cells were grown at confluence before induction of ciliogenesis for 72 hours by serum

deprivation. Cells were fixed 30 minutes in 2.5 % glutaraldehyde (Electron Microscopy

Sciences), 2 % paraformaldehyde (Electron Microscopy Sciences), 1 mM $CaCl_2$ in PBS, then

washed 3 x 5 minutes in PBS. Samples were then post-fixed during 30 minutes in 1 % Osmium

tetroxide (Electron Microscopy Sciences), then washed 3 x 5 minutes in water. Dehydration

was performed using graded series of ethanol in water for 5 minutes 30%, 50%, 70%, 90%,

100%, 100%. Resin infiltration was performed by incubating 30 minutes in an Agar low

viscosity resin (Agar Scientific Ltd) and EtOH (1:2) mix, then 30 minutes in a resin and EtOH

(2:1) mix followed by overnight incubation in pure resin. The resin was then changed and the

samples further incubated during 1.5 hours prior to inclusion in gelatin capsules and overnight

polymerization at 60 °C. 70 nm sections were obtained using an EM UC6 ultramicrotome

(Leica), post-stained in 4 % aqueous uranyl acetate and lead citrate, and observed at 80 kV

with a Tecnai12 transmission electron microscope (ThermoFischer Scientific) equipped with a

1K×1K Keen View camera (OSIS).

Statistical analysis

All statistical analyses were performed using the Prism7 for Mac OS X software (GraphPad

Software, Inc.). All values are provided as mean ± SD. The number of experimental replicates

and the statistical test used are indicated in the figure legends, and the p values are included

when statistically different.

References

Adams GM, Wright RL, Jarvik JW. 1985. Defective temporal and spatial control of flagellar assembly in a mutant of Chlamydomonas reinhardtii with variable flagellar number. *J Cell Biol* **100**:955–964.

Azimzadeh J. 2021. Evolution of the centrosome, from the periphery to the center. *Current Opinion in Structural Biology* **66**:96–103. doi:10.1016/j.sbi.2020.10.020

Basquin C, Ershov D, Gaudin N, Vu HT, Louis B, Papon JF, Orfila AM, Mansour S, Rink JC, Azimzadeh J. 2019. Emergence of a Bilaterally Symmetric Pattern from Chiral Components in the Planarian Epidermis. *Dev Cell* **51**:516-525 e5. doi:10.1016/j.devcel.2019.10.021

Boczek NJ, Hopp K, Benoit L, Kraft D, Cousin MA, Blackburn PR, Madsen CD, Oliver GR, Nair AA, Na J, Bianchi DW, Beek G, Harris PC, Pichurin P, Klee EW. 2018. Characterization of three ciliopathy pedigrees expands the phenotype associated with biallelic C2CD3 variants. *European Journal of Human Genetics* **26**:1797–1809. doi:10.1038/s41431-018-0222-3

Bustin SA, Benes V, Garson JA, Hellemans J, Huggett J, Kubista M, Mueller R, Nolan T, Pfaffl MW, Shipley GL, Vandesompele J, Wittwer CT. 2009. The MIQE guidelines: minimum information for publication of quantitative real-time PCR experiments. *Clin Chem* **55**:611–622. doi:10.1373/clinchem.2008.112797

Clare DK, Magescas J, Piolot T, Dumoux M, Vesque C, Pichard E, Dang T, Duvauchelle B, Poirier F, Delacour D. 2014. Basal foot MTOC organizes pillar MTs required for coordination of beating cilia. *Nature communications* **5**:4888. doi:10.1038/ncomms5888

Cortés CR, McInerney-Leo AM, Vogel I, Rondón Galeano MC, Leo PJ, Harris JE, Anderson LK, Keith PA, Brown MA, Ramsing M, Duncan EL, Zankl A, Wicking C. 2016.

Mutations in human C2CD3 cause skeletal dysplasia and provide new insights into
phenotypic and cellular consequences of altered C2CD3 function. *Scientific Reports* **6**.
doi:10.1038/srep24083

Delgehyr N, Meunier A, Faucourt M, Grau MB, Strehl L, Janke C, Spassky N. 2015.
Ependymal cell differentiation, from monociliated to multiciliated cells. *Methods in Cell*
Biology **127**:19–35. doi:10.1016/bs.mcb.2015.01.004

Feldman JL, Geimer S, Marshall WF. 2007. The mother centriole plays an instructive role in
defining cell geometry. *PLoS Biol* **5**:e149.

Gambarotto D, Zwettler FU, le Guennec M, Schmidt-Cernohorska M, Fortun D, Borgers S,
Heine J, Schloetel JG, Reuss M, Unser M, Boyden ES, Sauer M, Hamel V, Guichard P.
2019. Imaging cellular ultrastructures using expansion microscopy (U-ExM). *Nature*
Methods **16**:71–74. doi:10.1038/s41592-018-0238-1

Garcia-Gonzalo FR, Reiter JF. 2017. Open Sesame: How Transition Fibers and the Transition
Zone Control Ciliary Composition. *Cold Spring Harb Perspect Biol* **9**.
doi:10.1101/cshperspect.a028134

Geimer S, Melkonian M. 2005. Centrin scaffold in Chlamydomonas reinhardtii revealed by
immunoelectron microscopy. *Eukaryot Cell* **4**:1253–1263.

Geimer S, Melkonian M. 2004. The ultrastructure of the Chlamydomonas reinhardtii basal
apparatus: identification of an early marker of radial asymmetry inherent in the basal
body. *J Cell Sci* **117**:2663–2674.

Gibson DG, Young L, Chuang RY, Venter JC, Hutchison CA, Smith HO. 2009. Enzymatic
assembly of DNA molecules up to several hundred kilobases. *Nature Methods* **6**:343–345.
doi:10.1038/nmeth.1318

Goodrich L v., Johnson RL, Milenkovic L, McMahon JA, Scott MP. 1996. Conservation of the

hedgehog/patched signaling pathway from flies to mice: Induction of a mouse patched

gene by Hedgehog. *Genes and Development* **10**:301–312. doi:10.1101/gad.10.3.301

Gupta GD, Coyaud É, Gonçalves J, Mojarad BA, Liu Y, Wu Q, Gheiratmand L, Comartin D,

Tkach JM, Cheung SWT, Bashkurov M, Hasegan M, Knight JD, Lin ZY, Schueler M,

Hildebrandt F, Moffat J, Gingras AC, Raught B, Pelletier L. 2015. A Dynamic Protein

Interaction Landscape of the Human Centrosome-Cilium Interface. *Cell* **163**:1484–1499.

doi:10.1016/j.cell.2015.10.065

Izquierdo D, Wang WJ, Uryu K, Tsou MF. 2014. Stabilization of cartwheel-less centrioles for

duplication requires CEP295-mediated centriole-to-centrosome conversion. *Cell Rep*

8:957–965. doi:10.1016/j.celrep.2014.07.022

Kim J, Krishnaswami SR, Gleeson JG. 2008. CEP290 interacts with the centriolar satellite

component PCM-1 and is required for Rab8 localization to the primary cilium. *Hum Mol*

Genet **17**:3796–3805. doi:10.1093/hmg/ddn277

Kumar D, Reiter J. 2021. How the centriole builds its cilium: of mothers, daughters, and the

acquisition of appendages. *Current Opinion in Structural Biology*.

doi:10.1016/j.sbi.2020.09.006

LeGuennec M, Klena N, Aeschlimann G, Hamel V, Guichard P. 2021. Overview of the

centriole architecture. *Current Opinion in Structural Biology* **66**:58–65.

doi:10.1016/j.sbi.2020.09.015

Lin J, Heuser T, Song K, Fu X, Nicastro D. 2012. One of the Nine Doublet Microtubules of

Eukaryotic Flagella Exhibits Unique and Partially Conserved Structures. *PLoS ONE* **7**.

doi:10.1371/journal.pone.0046494

Mercey O, al Jord A, Rostaing P, Mahuzier A, Fortoul A, Boudjema AR, Faucourt M, Spassky
N, Meunier A. 2019. Dynamics of centriole amplification in centrosome-depleted brain
multiciliated progenitors. *Scientific Reports* **9**:1–11. doi:10.1038/s41598-019-49416-2

Meunier A, Azimzadeh J. 2016. Multiciliated cells in animals. *Cold Spring Harbor*
Perspectives in Biology **8**. doi:10.1101/cshperspect.a028233

Muto Y, Yoshioka T, Kimura M, Matsunami M, Saya H, Okano Y. 2008. An evolutionarily
conserved leucine-rich repeat protein CLERC is a centrosomal protein required for spindle
pole integrity. *Cell Cycle* **7**:2738–2748.

Nachury M v. 2018. The molecular machines that traffic signaling receptors into and out of
cilia. *Current Opinion in Cell Biology*. doi:10.1016/j.ceb.2018.03.004

Nommick A, Boutin C, Rosnet O, Bazellières E, Thomé V, Loiseau E, Viallat A, Kodjabachian
L. 2021. Lrrcc1 and Ccdc61 are conserved effectors of multiciliated cell function. *bioRxiv*
2021.01.30.428946. doi:10.1101/2021.01.30.428946

Ooi J. 2015. Mutations in C2CD3 cause oral-facial-digital syndrome through deregulation of
centriole length. *Clinical Genetics* **87**:328–329. doi:10.1111/cge.12545

Reiter JF, Leroux MR. 2017. Genes and molecular pathways underpinning ciliopathies. *Nat*
Rev Mol Cell Biol **18**:533–547. doi:10.1038/nrm.2017.60

Rohatgi R, Milenkovic L, Scott MP. 2007. Patched1 regulates hedgehog signaling at the
primary cilium. *Science* **317**:372–376. doi:10.1126/science.1139740

Romani M, Micalizzi A, Valente EM. 2013. Joubert syndrome: congenital cerebellar ataxia
with the molar tooth. *Lancet neurology* **12**:894–905. doi:10.1016/S1474-4422(13)70136-
4

Schneiter M, Halm S, Odriozola A, Mogel H, Rička J, Stoffel MH, Zuber B, Frenz M, Tschanz
SA. 2021. Multi-scale alignment of respiratory cilia and its relation to mucociliary
function. *Journal of Structural Biology* **213**:107680. doi:10.1016/j.jsb.2020.107680

Shaheen R, Szymanska K, Basu B, Patel N, Ewida N, Faqeih E, al Hashem A, Derar N, Alsharif

H, Aldahmesh MA, Alazami AM, Hashem M, Ibrahim N, Abdulwahab FM, Sonbul R,

Alkuraya H, Alnemer M, al Tala S, Al-Husain M, Morsy H, Seidahmed MZ, Meriki N,

Al-Owain M, AlShahwan S, Tabarki B, Salih MA, Ciliopathy W, Faquih T, El-Kalioby

M, Ueffing M, Boldt K, Logan C v, Parry DA, al Tassan N, Monies D, Megarbane A,

Abouelhoda M, Halees A, Johnson CA, Alkuraya FS. 2016. Characterizing the morbid

genome of ciliopathies. *Genome biology* **17**:242. doi:10.1186/s13059-016-1099-5

Sharma A, Olieric N, Steinmetz MO. 2021. Centriole length control. *Current Opinion in*

Structural Biology **66**:89–95. doi:10.1016/j.sbi.2020.10.011

Silflow CD, LaVoie M, Tam LW, Tousey S, Sanders M, Wu W, Borodovsky M, Lefebvre PA.

2001. The Vfl1 Protein in Chlamydomonas localizes in a rotationally asymmetric pattern

at the distal ends of the basal bodies. *J Cell Biol* **153**:63–74.

Singla V, Romaguera-Ros M, Garcia-Verdugo JM, Reiter JF. 2010. Ofd1, a human disease

gene, regulates the length and distal structure of centrioles. *Dev Cell* **18**:410–424.

Sofroniew N, Lambert T, Evans K, Nunez-Iglesias J, Yamauchi K, Solak AC, Bokota G,

ziyangczi, Buckley G, Winston P, Tung T, Pop DD, Hector, Freeman J, Bussonnier M,

Boone P, Royer L, Har-Gil H, Axelrod S, Rokem A, Bryant, Kiggins J, Huang M, Vemuri

P, Dunham R, Manz T, jakirkham, Wood C, Siqueira A de, Chopra B. 2020. napari/napari:

0.3.8rc1. doi:10.5281/ZENODO.4046812

Takao D, Yamamoto S, Kitagawa D. 2019. A theory of centriole duplication based on self-

organized spatial pattern formation. *Journal of Cell Biology* **218**:3537–3547.

doi:10.1083/JCB.201904156

Thauvin-Robinet C, Lee JS, Lopez E, Herranz-Pérez V, Shida T, Franco B, Jego L, Ye F,

Pasquier L, Loget P, Gigot N, Aral B, Lopes CAM, St-Onge J, Bruel AL, Thevenon J,

González-Granero S, Alby C, Munnich A, Vekemans M, Huet F, Fry AM, Saunier S,

Rivière JB, Attié-Bitach T, Garcia-Verdugo JM, Faivre L, Mégarbané A, Nachury M v.

2014. The oral-facial-digital syndrome gene C2CD3 encodes a positive regulator of

centriole elongation. *Nature Genetics* **46**:905–911. doi:10.1038/ng.3031

Tsai JJ, Hsu W bin, Liu JH, Chang CW, Tang TK. 2019. CEP120 interacts with C2CD3 and

Talpid3 and is required for centriole appendage assembly and ciliogenesis. *Scientific*

Reports **9**. doi:10.1038/s41598-019-42577-0

Vaughan S, Gull K. 2016. Basal body structure and cell cycle-dependent biogenesis in

Trypanosoma brucei. *Cilia*. doi:10.1186/s13630-016-0023-7

Wang Lei, Failler M, Fu W, Dynlacht BD. 2018. A distal centriolar protein network controls

organelle maturation and asymmetry. *Nature Communications* **9**. doi:10.1038/s41467-

018-06286-y

Wang L, Failler M, Fu W, Dynlacht BD. 2018. A distal centriolar protein network controls

organelle   maturation   and   asymmetry.   *Nature   communications*   **9**:3938.

doi:10.1038/s41467-018-06286-y

Yamamoto S, Kitagawa D. 2021. Emerging insights into symmetry breaking in centriole

duplication: updated view on centriole duplication theory. *Current Opinion in Structural*

Biology. doi:10.1016/j.sbi.2020.08.005

Yang TT, Chong WM, Wang WJ, Mazo G, Tanos B, Chen Z, Tran TMN, Chen Y de, Weng

RR, Huang CE, Jane WN, Tsou MFB, Liao JC. 2018. Super-resolution architecture of

mammalian centriole distal appendages reveals distinct blade and matrix functional

components. *Nature Communications* **9**:1–11. doi:10.1038/s41467-018-04469-1

Ye X, Zeng H, Ning G, Reiter JF, Liu A. 2014. C2cd3 is critical for centriolar distal appendage

assembly and ciliary vesicle docking in mammals. *Proceedings of the National Academy*

of   Sciences   of   the   United   States   of   America   **111**:2164–2169.

doi:10.1073/pnas.1318737111

Yubuki N, Leander BS. 2013. Evolution of microtubule organizing centers across the tree of

eukaryotes. *The Plant journal : for cell and molecular biology* **75**:230–244.

doi:10.1111/tpj.12145

Zhang D, Aravind L. 2012. Novel transglutaminase-like peptidase and C2 domains elucidate

the structure, biogenesis and evolution of the ciliary compartment. *Cell Cycle* **11**:3861–

3875. doi:10.4161/cc.22068

Figure legends

Figure 1. LRRCC1 is localized in a rotationally asymmetric manner at the distal end of

centrioles in the human centrosome. a) LRRCC1 localization in non-treated RPE1 cells

(left) or in cells treated with nocodazole to disperse the pericentriolar satellites (right).

LRRCC1 (Ab2, yellow), γ-tubulin (PCM, magenta) and DNA (cyan). Bar, 5 μm (insets, 2 μm).

b) Longitudinal view of centrioles and procentrioles in the duplicating centrosome of an RPE1

cell analyzed by U-ExM. LRRCC1 (Ab2, yellow), acetylated tubulin (magenta). Bar, 0.5 μm.

c) Centrioles from WT RPE1 cells as seen from the distal end. LRRCC1 (Ab2, yellow),

acetylated tubulin (magenta). Bar, 0.2 μm. **d)** Workflow for calculating the average staining

from 3D-reconstructed individual centrioles generated from confocal z-stacks. The brightest

part of LRRCC1 signal was used as a reference point to align the centrioles. **e)** Average

LRRCC1 staining obtained from 34 individual centrioles viewed from the distal end, in

transverse and longitudinal views. A diagram representing the average pattern in transverse

view is also included.

Figure 2. The LRRCC1 rotationally asymmetric pattern is conserved in mouse MCCs.

a) Centrioles in the cytoplasm of mouse ependymal cells differentiating *in vitro* analyzed by

U-ExM, in longitudinal and transverse view. Lrrcc1 (Ab2, yellow), γ-tubulin (basal foot cap,

cyan) and acetylated tubulin (magenta). Of note, γ-tubulin was also detected in the proximal

lumen of centrioles. Bar, 0.2 μm. **b)** Procentrioles assembling via the centriolar (right) or the

deuterosome pathway (left and center) in ependymal cells. Lrrcc1 (Ab2, yellow), acetylated

tubulin (magenta). Bar, 0.2 μm. **c)** Transverse view of centrioles docked at the apical membrane

in fully differentiated mouse tracheal cells, viewed from the distal end. Lrrcc1 (Ab2, yellow),

γ-tubulin (cyan) and acetylated tubulin (magenta). Bar, 0.2 μm. **d)** Average image generated

from 35 individual centrioles from mouse trachea, viewed from the distal end, shown in

transverse and longitudinal views. The position of the basal foot (cyan dotted line) stained with

γ-tubulin was used as a reference point to align the centrioles. A diagram of the average pattern

in transverse view is shown, in which the direction of ciliary beat (Schneiter et al., 2021) is

represented by a dotted arrow and the basal foot axis by a green line. Triplets are numbered

counterclockwise from the LRRCC1 signal.

Figure 3. The conservation of centriole rotational asymmetry in humans is not linked to

stereotypical patterns of procentriole position and orientation. a) Early stage of

procentriole assembly stained for LRRCC1 (Ab2, yellow), SAS-6 (cyan) and acetylated tubulin

(magenta) in a HEK 293 cell. The brightness of the acetylated tubulin labelling was increased

in the inserts. Bar, 0.1 μm. **b)** Successive stages of centriole elongation in HEK 293 cells

stained for LRRCC1 (Ab2, yellow) and acetylated tubulin (magenta). Bar, 0.1 μm. **c)** Location

of LRRCC1 in the procentrioles (top panels) and position of the procentriole relative to its

parent centriole polarity (bottom panels), in RPE1 and HEK 293 centrioles analyzed by U-

ExM. For each diplosome, the angle between LRRCC1 in the procentriole and the centriole

long axis (top panels), or between the procentriole and LRRCC1 in the centriole (bottom

panels) was measured. The number of diplosomes analyzed is indicated. p values are indicated

when statistically different from a random distribution (χ^2-test).

Figure 4. LRRCC1 is required for ciliary assembly and signaling. a) Left: LRRCC1

staining (Ab2) of WT or LRRCC1-defficient RPE1 cells obtained by CRISPR/Cas9 editing

(clones 1.1, 1.2 and 1.9). Bar, 2 μm. Right: quantification of fluorescence intensity. Bars, mean

± SD, 3 independent experiments. p values are provided when statistically significant from the

corresponding control (One-way ANOVA). **b)** Percentage of ciliated cells in WT or LRRCC1-

deficient serum-deprived cells. Bars, mean ± SD, ≥ 331 cells from 4 independent experiments

for each condition. p values are provided when statistically significant from the corresponding

control (One-way ANOVA). **c)** Percentage of ciliated cells in serum deprived RPE1 cells

treated with control or LRRCC1 siRNAs. Bars, mean ± SD, ≥ 83 cells from 3 independent

experiments for each condition. p values are provided when statistically significant from the

corresponding control (One-way ANOVA). **d)** Left: SMO (yellow) accumulation at primary

cilia (ARL13B, magenta) following SAG-induction of the Hedgehog pathway, in WT or

CRISPR RPE1 cells. Bar, 2 μm. Right: quantification of ciliary SMO expressed as a percentage

of the SAG-treated WT mean. Bars, mean ± SD, 3 independent experiments. p values are

provided when statistically significant from the corresponding control (One-way ANOVA). **e)**

Ciliary SMO expressed as a percentage of the SAG-induced WT mean in RPE1 cells treated

with control or LRRCC1 siRNAs. Bars, mean ± SD, 3 independent experiments. p values are

provided when statistically significant from the corresponding control (One-way ANOVA).

Figure 5. Depleting LRRCC1 induces defects in centriole structure. a) Centriole length in

mother (MC) and daughter (DC) centrioles analyzed by U-ExM in WT or LRRCC1-deficient

clones (1.1 and 1.9). Left: Centrioles were stained for acetylated tubulin (magenta) and CEP164

(yellow) to measure centriole length (arrows). Bar, 0.5 μm. Right: Quantification from 3

independent experiments. p values are provided when statistically significant from the

corresponding control (One-way ANOVA). **b)** Transmission electron microscopy view of

centrioles in WT and CRISPR (clone 1.9) RPE1 cells. Note that the two 1.9 centrioles are from

the same cell. Bar, 0.5 μm. **c)** Examples of normal DAs, DAs with abnormal morphology

(white arrowhead: abnormal spacing between consecutive DAs; cyan arrowhead: abnormal DA

shape) or missing DAs (grey arrowhead) analyzed by U-ExM. Bar, 1 μm. **d)** Percentages of

centrioles presenting DA defects in WT or CRISPR RPE1 cells. ≥ 87 centrioles from 8

independent experiments for each condition. p values are provided when statistically

significant from the corresponding control (Two-way ANOVA). **e)** Percentages of centrioles

presenting DA defects in WT or CRISPR HEK 293 (clone 21) cells. $\geq$ 40 centrioles from 4

independent experiments for each condition. p values are provided when statistically

significant from the corresponding control (Two-way ANOVA).

Figure 6. C2CD3 localizes asymmetrically at the distal end of centrioles and is affected

by LRRCC1-depletion. a) C2CD3 levels at the centrosome of WT or LRRCC1 CRISPR

RPE1 clones. Bars, mean $\pm$ SD, 3 independent experiments. p values are provided when

statistically significant from the corresponding control (One-way ANOVA). **b)** C2CD3 levels

at the centrosome in RPE1 cells treated with control or LRRCC1 siRNAs. Bars, mean $\pm$ SD, 3

independent experiments. p values are provided when statistically significant from the

corresponding control (One-way ANOVA). **c)** Longitudinal view of a centriole analyzed by U-

ExM and stained for C2CD3 (yellow) and acetylated tubulin (magenta). Bar, 0.2 μm. **d)**

Centrioles from WT RPE1 cells as viewed from the distal end. C2CD3 (yellow), acetylated

tubulin (magenta). Bar, 0.2 μm. **e)** Average C2CD3 staining obtained from 40 individual

centrioles from WT RPE1 cells viewed from the distal end, in transverse and longitudinal

views. The region where the C2CD3 signal is closest to the triplet wall after correction of

centriole orientation was used as a reference point to align the individual centrioles. A diagram

of the average pattern in transverse view is included. **f)** Centrioles from LRRCC1-deficient

cells (CRISPR clones 1.1 and 1.9) in transverse section as viewed from the distal end. C2CD3

(yellow), acetylated tubulin (magenta). Bar, 0.2 μm. **g)** Average C2CD3 staining obtained from

47 (clone 1.1) or 24 (clone 1.9) individual centrioles from LRRCC1-deficient RPE1 cells

viewed from the distal end, in transverse and longitudinal views.

Figure 7. C2CD3 and LRRCC1 partially colocalize at the distal end of centrioles. a) RPE1

centrioles processed for U-ExM and stained for LRRCC1 (Ab2, yellow), C2CD3 (cyan) and

acetylated tubulin (magenta). Bar, 0.1 μm. **b)** Model showing the possible location of LRRCC1

and C2CD3 relative to each other within human centrioles. Right panel: diagram showing the

respective positions of the acorn (Geimer and Melkonian, 2004) and Vfl1p (Silflow et al., 2001)

in *C. reinhardtii*. The direction of the flagellar beat is indicated by a dotted arrow, and the distal

striated fiber is in grey. **c)** Evolution of the roles played by Vfl1p/LRRCC1 proteins and

associated rotationally asymmetric centriolar substructures. In *C. reinhardtii*, Vfl1p is required

for proper ciliary assembly (1), as well as for the formation of fibers and microtubular roots (2)

that control the position of centrioles and procentrioles (3), and overall cellular organization

(Adams et al., 1985; Silflow et al., 2001). In human cells, LRRCC1 and C2CD3 are required

for primary cilium assembly (1) - this study and (Thauvin-Robinet et al., 2014; Ye et al., 2014)

- and a role in asymmetric anchoring of centrosome or cytoskeletal components might also be

conserved (2).

Supplemental material

Fig. S1 provides additional data on LRRCC1 expression in CRISPR clones and RNAi-treated

cells. **Fig. S2** presents the image analysis pipeline for generating the average images of

LRRCC1 and C2CD3 staining. **Fig. S3** shows the quantification of the DA component

CEP164, and the distal centriole components CEP290 and OFD1, at the centrosome of RPE1

cells depleted from LRRCC1 by CRISPR or RNAi. **Fig. S4** shows that C2CD3 is not co-

immunoprecipitated with GFP-LRRCC1.

Table 1. Antibodies used in this study.

Antibody	Dilution IF	Dilution U-ExM	Dilution WB	RRID Identifier	Source	Reference
Primary antibodies						
Goat anti-ARL13B	1:100	/	/	RRID:AB_2058502	Santa Cruz Biotechnology	sc-102318
Guinea pig anti-alpha tubulin AA344 monobody	/	1:500	/		Geneva antibody facility	scFv-S11B
Guinea pig anti-beta tubulin AA345 monobody	/	1:500	/		Geneva antibody facility	scFv-F2C
Mouse anti-acetylated tubulin (6-11B-1)	1:1000	1:500	/	RRID:AB_628409	Santa Cruz Biotechnology	sc-23950
Mouse anti-CEP290 (B-7)	1:500	/	/	RRID:AB_2890036	Santa Cruz Biotechnology	sc-390462
Mouse anti-CBY (8-2)	1:500	/	/	RRID:AB_1561972	Santa Cruz Biotechnology	sc-101551
Mouse anti-gamma tubulin (GTU88)	1:2000	1:200	/	RRID:AB_532292	Sigma-Aldrich	T5326
Mouse anti-SAS-6	/	1:100	/	RRID:AB_1128357	Santa Cruz Biotechnology	sc-81431
Mouse anti-Smoothened	1:200	/	/	RRID:AB_1270802	Abcam	ab72130
Rabbit anti-ARL13B	1:500	/	/	RRID:AB_2060867	Proteintech	17711-1-AP
Rabbit anti-C2CD3	1:500	1:500	/	RRID:AB_10669542	Sigma-Aldrich	HPA038552
Rabbit anti-C2CD3	/	/	1:1000	RRID:AB_2718714	ThermoFisher Scientific	PA5-72860
Rabbit anti-CEP164	1:500	1:300	/	RRID:AB_2651175	Proteintech	22227-1-AP
Rabbit anti-GFP	/	/	1:1000	RRID:AB_591816	MBL International	598
Rabbit anti-HA	/	/	1:1000	RRID:AB_631618	Santa Cruz Biotechnology	sc-805
Rabbit anti-KI67	1:1000	/	/	RRID:AB_443209	Abcam	ab15580
Rabbit anti-LRRCC1 Ab1	1:500	1:200	/		This study	
Rabbit anti-LRRCC1 Ab2	1:500	1:300	1:1000		This study	
Rabbit anti-OFD1	1:500	/	/	RRID:AB_2890033.	Sigma-Aldrich	ABC961
Sheep anti-C2CD3	1:200	1:100	/	RRID:AB_10997138	R&D systems	AF7348
Secondary antibodies						
Donkey Anti-Goat IgG H&L (Alexa Fluor® 488)	1:500	1:500	/	RRID:AB_2687506	Abcam	ab150129
Donkey Anti-Goat IgG H&L (Alexa Fluor® 568)	1:500	1:500	/	RRID:AB_2636995	Abcam	ab175474
Donkey Anti-Goat IgG H&L (Alexa Fluor® 647)	1:500	1:100	/	RRID:AB_2732857	Abcam	ab150131
Donkey Anti-Mouse IgG H&L (Alexa Fluor® 488)	1:500	1:500	/	RRID:AB_2732856	Abcam	ab150105
Donkey Anti-Mouse IgG H&L (Alexa Fluor® 568)	1:500	1:500	/	RRID:AB_2636996	Abcam	ab175472
Donkey Anti-Mouse IgG H&L (Alexa Fluor® 647)	1:500	1:100	/	RRID:AB_2890037	Abcam	ab150107
Donkey Anti-Rabbit IgG H&L (Alexa Fluor® 488)	1:500	1:500	/	RRID:AB_2636877	Abcam	ab150073
Donkey Anti-Rabbit IgG H&L (Alexa Fluor® 647)	1:500	1:500	/	RRID:AB_2752244	Abcam	ab150075
Donkey Anti-Sheep IgG H&L (Alexa Fluor® 647)	/	1:100	/	RRID:AB_2884038	Abcam	ab150179
Goat Anti-Guinea Pig IgG (H+L) (Alexa Fluor 568)	/	1:100	/	RRID:AB_141954	ThermoFisher Scientific	A-11075
Goat Anti-Rabbit IgG (H+L) Horseradish Peroxidase Conjugate	/	/	1:1000	RRID:AB_2536530	ThermoFisher Scientific	G-21234

Table 2. PCR primers used in this study.

Primer	Sequence
LRRCC1-1_Fw	CAA CAA GGA TCT TCT CTA GCC CA
LRRCC1-1_Rv	AGT TTG GTC GTC TAT GAT TTT GCA
LRRCC1-2_Fw	GCA CAA CAA GGA TCT TCT CTA GC
LRRCC1-2_Rv	TCG CAG ACA TTC ATT CTC TCT AGA
PTCH1_Fw	CCC CTG TAC GAA GTG GAC ACT CTC
PTCH1_Rv	AAG GAA GAT CAC CAC TAC CTT GGC T
CHMP2A_Fw	ATG GGC ACC ATG AAC AGA CAG
CHMP2A_Rv	TCT CCT CTT CAT CTT CCT CAT CAC
EMC7_Fw	GTC AGA CTG CCC TAT CCT CTC C
EMC7_Rv	CAT GTC AGG ATC ACT TGT GTT GAC

Figure 1

Figure 2

Figure 3

Figure 4

Figure 6

Figure 7

a

b

c