Title: Patterns of pan-genome occupancy and gene co-expression under water-deficit in *Brachypodium*

distachyon.

Authors: Rubén Sancho[1,2], Pilar Catalán[,1,2], Bruno Contreras-Moreira[2,3,4], Thomas E. Juenger[5], David L.

Des Marais*[6]

Affiliations:

[1] Department of Agricultural and Environmental Sciences, High Polytechnic School of Huesca,

University of Zaragoza, Huesca, Spain

[2] Grupo de Bioquímica, Biofísica y Biología Computacional (BIFI, UNIZAR), Unidad Asociada al CSIC,

Spain

[3] Estación Experimental de Aula Dei-Consejo Superior de Investigaciones Científicas, Zaragoza, Spain

[4] Fundación ARAID, Zaragoza, Spain

[5] Department of Integrative Biology, The University of Texas at Austin, Austin, TX. USA

[6] Department of Civil and Environmental Engineering, Massachusetts Institute of Technology,

Cambridge, MA. USA

* Corresponding author:

David L. Des Marais. 15 Vassar Street Room 48-325, Cambridge, MA, 02139 (USA). Phone:

617.258.6482. Email: dldesmar@mit.edu

Running Title: Transcriptome variation in *Brachypodium*

ABSTRACT

Natural populations are characterized by abundant genetic diversity driven by a range of different types of mutation. The tractability of sequencing complete genomes has allowed new insights into the variable composition of genomes, summarized as a species pan-genome. These analyses demonstrate that many genes are absent from the first reference genomes, whose analysis dominated the initial years of the genomic era. Our field now turns towards understanding the functional consequence of these highly variable genomes. Here, we analyzed weighted gene co-expression networks from leaf transcriptome data for drought response in the purple false brome *Brachypodium distachyon* and the differential expression of genes putatively involved in adaptation to this stressor. We specifically asked whether genes with variable "occupancy" in the pan-genome – genes which are either present in all studied genotypes or missing in some genotypes – show different distributions among co-expression modules. Co-expression analysis united genes expressed in drought-stressed plants into nine modules covering 72 hub genes (87 hub isoforms), and genes expressed under controlled water conditions into 13 modules, covering 190 hub genes (251 hub isoforms). We find that low occupancy pan-genes are under-represented among several modules, while other modules are over-enriched for low-occupancy pan-genes. We also provide new insight into the regulation of drought response in *B. distachyon*, specifically identifying one module with an apparent role in primary metabolism that is strongly responsive to drought. Our work shows the power of integrating pan-genomic analysis with transcriptomic data using factorial experiments to understand the functional genomics of environmental response.

INTRODUCTION

Soil water availability is a critical factor determining plant growth, development, and reproduction

(Bohnert, Nelson, & Jensen, 1995). Plants are able to cope with and acclimate to a range of soil water

contents through the reprogramming of their physiology, growth, and development over time scales

ranging from hours to seasons (Chaves, Maroco, & Pereira, 2003). Many of these acclimation strategies

arise from altered transcriptional profiles (Fisher et al., 2016; Miao, Han et al., 2017). Drought-responsive

gene regulatory pathways have been investigated extensively in model plant systems such as *Arabidopsis*

thaliana, maize, and rice (Borah et al., 2017; Hayano-Kanashiro et al., 2009; Janiak, Kwa, & Szarejko,

2015; Nakashima, Ito, & Yamaguchi-Shinozaki, 2009; Nakashima, Yamaguchi-Shinozaki, & Shinozaki,

2014). A clear emerging theme, however, is that diverse species and varieties of plants exhibit diverse

stress response mechanisms (Des Marais et al., 2012; Juenger, 2013; Pinheiro & Chaves, 2011), often

controlled by complex regulatory networks. Understanding the genetic control of this phenotypic

diversity is a priority for understanding the response of natural populations to climate change, and for

designing resilient crop species (Benfey & Mitchell-Olds, 2008).

Recent studies have brought attention to the remarkable variation in gene content among plant

populations (Alonge et al., 2020; Gao et al., 2019; Gordon et al., 2017; Haberer et al., 2020), reflected in a

species' pan-genome. A pan-genome refers to the genomic content of a species as a whole, rather than the

composition of a single, reference, individual genotype (Koonin & Wolf, 2008). In practice, pan-genomes

are estimated by deeply resequencing the genomes of a diversity panel of genotypes, often using a

reference genome to aid in final assembly and annotation (Lei et al., 2021). In the diploid model grass

Brachypodium distachyon, genomic analysis of 56 inbred natural "accessions" revealed that the total pan-

genome of the species comprised nearly twice the number of genes in any single accession (Gordon et al.,

2017). Remarkably, only 73% of genes in a given accession are found in at least 95% of the other

accessions (Gordon et al., 2017) – so-called "core genes" (universal or nearly universal genes; Koonin &

Wolf, 2008) and "soft-core genes" (found in at least 95% of accessions; Kaas et al. 2012) – suggesting

that a large number of genes are unique to subsets of accessions or even to individual accessions. The list

of core genes in *B. distachyon* is enriched for annotations associated with essential cellular processes such

as primary metabolism. Lower-occupancy genes, or "shell genes," are found in 5-94% of accessions and

their annotations are enriched for many processes related to environmental response, including disease

resistance. Similar patterns have been observed in the pan-genomes of *Arabidopsis thaliana*, barley,

sunflower, and an ever-growing number of additional plant species (Bayer et al., 2020; Contreras-Moreira

et al., 2017; Hübner et al., 2019). The DNA sequences of core genes bear the hallmark of strong purifying

selection and are typically expressed at a higher level and in more tissues as compared to shell genes.

Shell genes may be the result of gene duplications or deletions in the ancestor of a subset of studied

genotypes and, indeed, the vast majority of shell genes in *B. distachyon* appear to be functional, as

homologs are found in other species' genomes (Gordon et al., 2017).

The preceding observations raise the intriguing possibility that shell genes may represent segregating

variation that could be shaped by natural selection and thereby facilitate local adaptation or adaptive

responses to a variable environment. Multiple studies in *Arabidopsis thaliana* demonstrate the role of

segregating functional gene copies – effectively large-effect mutations – in shaping whole-plant response

to the abiotic environment (Monroe et al., 2016, 2018). The phenotypic effect size of a mutation can

determine the likelihood that the mutant will become fixed in a population, with large-effect mutations

more likely than not to confer deleterious phenotypes that may be removed from populations by natural

selection (Fisher, 1930). The observation that any two accessions of *B. distachyon* likely differ in the

presence or absence of hundreds of functional gene copies begs the question as to how potentially

function-changing gene deletions escape the purging effects of purifying selection. Pan-genomics requires

that we reconceptualize how we interpret "gene loss" as we move beyond a reference-genome view of

genome function. Genes identified as "missing" from a subset of accessions might represent deletions of

genes whose function was no longer constrained by purifying selection in some novel environment,

duplicated genes that originated in a subset of accessions and thus are absent in other accessions, or

paralogs that were both present in a common ancestor and then reciprocally lost in subsets of accessions.

The pleiotropic effect of a mutation can be affected by the number of genes with which it interacts

(Jeong et al., 2001); if a gene has relatively few interacting partners then its presence or absence in a

particular accession may have a small fitness effect and thus be maintained in populations. Similarly, the

efficacy of selection to purge deleterious alleles may be reduced if a gene is only expressed in a subset of

environments experienced by a species (Paaby & Rockman, 2014). In the context of pan-genomes, we

consider gene presence/absence polymorphisms as mutations, and we explore functional gene turnover by

testing the hypothesis that shell genes and core genes differ in their topological positions in

environmentally responsive gene co-expression networks.

Gene co-expression networks are widely used to interpret functional genomic data by assessing

patterns of correlation among genes via a threshold that assigns a connection weight to each gene pair

(Langfelder & Horvath, 2008; Zhang & Horvath, 2005). Sets of genes with similar expression profiles are

assigned to modules by applying graph clustering algorithms (Mao et al., 2009). Genes, or "nodes," in

such networks show considerable variation in the extent to which their expression co-varies with other

genes. As such, co-expression networks are generally considered "scale-free" in the sense that few nodes

have many neighboring nodes while many nodes have few neighboring nodes (Guelzim et al., 2002).

Modules are often comprised of genes with similar functions (Stuart et al., 2003; Wolfe, Kohane, &

Butte, 2005). High connectivity "hub" genes that show a large number of interactions with other genes

within a weighted co-expression network are candidates for key players in regulating cellular processes

(Albert, Jeong, & Barabási, 2000; Carlson et al., 2006; Dong & Horvath, 2007). As such, hub genes might

be expected to encode essential cellular functions and thus show pleiotropic effects when mutated or

deleted. By contrast, genes with fewer close co-expression relationships are often situated on the

periphery of networks and might, therefore, exhibit fewer pleiotropic effects when missing or mutated

(Des Marais et al., 2017a; Masalia, Bewick, & Burke, 2017; Porth et al., 2014). In this context, we

hypothesize that pan-genome core genes may be over-represented among co-expression network "hub-

genes," as both appear to be involved in core cellular processes and may therefore show deleterious

effects when deleted. Conversely, we hypothesize that pan-genome "shell genes" – whose patterns of

expression and thereby phenotypic effects are more restricted and condition-specific – will be enriched

among lowly connected (non-hub) genes in gene co-expression networks.

Here, we study the relationship between a plant's pan-genome and its gene co-expression networks

using *Brachypodium distachyon. Brachypodium* is a small genus of the subfamily Pooideae (Poaceae)

that contains ~20 species distributed worldwide (Catalán et al., 2016; Scholthof et al., 2018; Hasterok et

al. 2022). The annual diploid species *B. distachyon* is a model for temperate cereals and biofuel grasses

(Vogel et al. 2010; Mur et al. 2011; Catalán et al. 2014; Scholthof et al. 2018); a reference genome for

one *B. distachyon* accession, Bd21 (IBI, 2010) is now complemented by 54 deeply resequenced natural

accessions (Gordon et al., 2017). Recent studies demonstrate the utility of *B. distachyon* and its close

congeners for elucidating the evolution and ecology of plant-abiotic interactions, focusing especially on

responses to soil drying, aridity, and water use strategy (Des Marais & Juenger, 2016; Des Marais et al.,

2017b; Monroe et al. 2021; Fisher et al., 2016; Handakumbura et al., 2019; Manzaneda et al., 2015, 2012;

Martínez et al., 2018; Skalska et al., 2020; Verelst et al., 2013). In the present study, we first identify and

characterize gene co-expression modules associated with response to soil drying. We then test the

hypothesis that the occupancy of pan-genes – whether they are part of the shell or core gene sets of the

pan-genome – is associated with their connectivity in the *B. distachyon* gene co-expression network. Our

work demonstrates the dynamic nature of plant genomes and sets up future work on the functional

consequences of diversity on the evolution of gene regulatory networks.

MATERIALS AND METHODS

Plant material, experimental design, total-RNA extraction and 3' cDNA tag libraries preparation

Sampling herein follows our earlier work documenting physiological and developmental response of

33 diploid natural accessions of *Brachypodium distachyon* (L.) P. Beauv. to soil drying (Figure 1; Table

S1) (Des Marais et al., 2017b). The sampled accessions were inbred for more than five generations (Filiz

et al., 2009; Vogel et al., 2006, 2009) and represent the geographic and ecological diversity of *B.*

distachyon across the Mediterranean region. Whole genome resequencing data is available for all studied

accessions (Gordon et al., 2017).

A total of 264 individual plants from the 33 accessions were grown under two greenhouse conditions,

restriction of water (drought, D) and well-watered (water, W). We sampled four biological replicates per

treatment and accession combinations [33 accessions x 4 replicates x 2 treatments (D and W)]. For a full

description of the growth and treatment conditions, please see Des Marais, Lasky, et al. (2017b). In short,

seeds were stratified at 6°C for 14d and then greenhouse-grown in 600 mL of Profile porous ceramic

rooting media (Profile Products) in Deepot D40H pots (650 mL; Stuewe & Sons). For the first 21d of

growth, all plants were watered to field capacity every other day. Daytime high temperatures ranged from

23°C to 28°C and night-time lows from 14°C to 18°C. On day 21 four treatment regimes were

implemented: Cool Wet, Cool Drought, Hot Wet, and Hot Drought, with Drought and Wet plants

spatially randomized within single blocks of Hot or Cool conditions. The hot treatment raised air

temperatures in the plant canopies by ~10°C. Because temperature was confounded with experimental

block in the design used for the current study, we did not include a temperature effect in any of the

statistical models used herein. Well-watered plants (hereafter "Water") were watered to field-capacity

every second day with fresh water, whereas drought plants (hereafter "Drought") were hand-watered daily

by pipette such that the soil water was reduced by 5% each day. The final soil water content of Drought

plants on day 11 of treatment – 32d post-germination – was 45% field capacity, which corresponds to a

decrease in water potential of 1.2 MPa as compared to field capacity in this growth media (Des Marais et

al., 2012).

For each plant, the two youngest, fully expanded leaves of the tallest tiller were excised with a razor

blade at the base of the lamina and flash-frozen on liquid nitrogen. Tissue was ground to a fine powder

under liquid nitrogen using a Mixer Mill MM 300 (Retsch GmbH). RNA was extracted using the Sigma

Spectrum Total Plant RNA kit, including on-column DNase treatment, following the manufacturer's

protocol, and quantified using a NanoDrop spectrophotometer (Thermo Scientific). We used a RNA-Seq

library protocol (3' cDNA tag libraries with fragment of 300-500 bp) for sequencing on the Illumina

HiSeq platform adapted from Meyer et al. (2011). This Tag-Seq method yields only one sequence per

expressed transcript in the RNA pool, allowing for higher sequencing coverage per gene as a function of

total sequencing effort (Tandonnet & Torres, 2017).

Pre-processing of sequences, quantifying transcript abundances, normalizing procedures, and

controlling batch effects

Sequencing was carried out using an Illumina HiSeq2500 platform (100 bp Single-End, SE,

sequencing). Quality control of SE reads was performed with FastQC software. Adapters and low quality

reads were removed and filtered with Trimmomatic-0.32 (Bolger, Lohse, & Usadel, 2014). Total numbers

of raw and filtered SE reads for each accession and treatment are shown in Table S2. Quantifying the

abundances of transcripts from RNA-Seq data was done with Kallisto v0.43.1 (Bray et al., 2016). To

accommodate the library preparation and sequencing protocols (3' tag from fragments of 300-500 bp),

pseudoalignments of RNA-Seq data were carried out using as references 500 bp from the 3' tails of the *B.*

*distachyon*_314_v3.1 transcriptome (IBI 2010;

https://phytozome.jgi.doe.gov/pz/portal.html#!info?alias=Org_Bdistachyon). We applied estimated

average fragment lengths of 100 bp and standard deviations of fragment length of 20. Resulting numbers

of transcripts per million (TPM) were recorded.

Exploratory analysis of the data set and the subsequent filtering and normalization of transcripts

abundance between samples, and the *in silico* technical replicate steps (bootstrap values computed with

Kallisto), were conducted with the Sleuth package (Pimentel et al., 2017). A total of 16,386 targets

(transcripts/isoforms) were recovered after the normalizing and filtering step using Sleuth. This program

was also used for batch-correction of data and of differentially expressed (DE) genes. To account for

library preparation batch effects, date of library preparation was included as a covariate with condition

variable in the full model (Table S2).

Co-expression network analysis of normalized transcripts abundance

Co-expression networks for the Drought and Water (control) data sets were carried out using the

transcripts per million (TPM) estimates and the R package WGCNA (Langfelder and Horvath 2008). We

analyzed 16,386 transcripts that were filtered and normalized for 127 Drought and 124 Water individual

samples (individual plants). After the removal of putative outliers, we retained 121 Drought and 108

Water samples that were used for network construction.

Identical parameters were used for the Drought and the Water data sets to construct their respective

co-expression networks. The *BlockwiseModules* function was used to perform automatic network

construction and module detection on the large expression data set of transcripts. Parameters for co-

expression network construction were fitted checking different values. We chose the Pearson correlation

and unsigned network type, soft thresholding power 6 (high scale free, $R^2 > 0.85$), a minimum module size

of 30, and a medium sensitivity (deepSplit = 2) for the cluster splitting. The topological overlap matrix

(TOM) was generated using the TOMtype unsigned approach. Module clustering was performed with

function *cutreeDynamic* and the Partitioning Around Medoids (PAM) option activated. Module merging

was conducted with mergeCutHeight set to 0.30. K_{diff} was calculated using WGCNA to estimate the

relationship between connectivity among genes within *vs* between co-expression modules.

Both isoform and gene counts were calculated. Isoform counts included all transcripts identified (e.g.,

Bradi1g1234.1; Bradi1g1234.2; Bradi1g1234.3) and gene counts only included different genes expressed,

thus different isoforms from the same gene were reduced to a single gene count in each case (e.g.,

Bradi1g12345.1 and Bradi1g12345.2 are two isoforms of one gene, Bradi1g12345).

Detection of highly connected nodes (hub genes/ isoforms) within co-expression networks

Three representative descriptors of modules, module eigengene (ME), intramodular connectivity

(k_{IM}), and eigengene-based connectivity (k_{ME}; or its equivalent module membership, MM) were

calculated using the WGCNA package. Briefly, ME is defined as the first principal component of a given

module and is often considered to represent the gene expression profiles within the module. k_{IM} measures

how connected, or co-expressed, a given gene is with respect to the genes of a particular module. Thus,

intra-modular connectivity is also the connectivity in the subnetwork defined by the module. MM is the

correlation of gene expression profile with the module eigengene (ME) of a given module. MM values

close to 1 or -1 indicate genes highly connected to the module. The sign of MM indicates a positive or a

negative relationship between a gene and the eigengene of the module (Langfelder & Horvath, 2010).

Genes with absolute MM value over 0.9 were considered "hub genes." Correlations between MM values

transformed by a power of $\beta = 6$ and k_{IM} values were also calculated.

Pan-genome occupancy of clustered, hub, and differentially expressed genes across accessions

Because the *B. distachyon* accessions studied herein comprise a subset of those included in the

original pan-genome (Gordon et al. 2017), we re-ran the clustering procedures used in our earlier analysis

with only the 33 accessions used here. We clustered CDS sequences from the annotated genomes of each

of the studied accessions to define core, soft-core, and shell genes with the software

GET_HOMOLOGUES-EST v03012018 (Contreras-Moreira et al., 2017) using the OMCL algorithm (-

M) and a stringent percent-sequence identity threshold (–S 98). The resulting pan-genome matrix was

interrogated to identify "core" genes observed in all 33 accessions, "soft-core" genes observed in 32 and

31, and "shell" genes observed in 30 or fewer accessions. Occupancy was defined as the number of

accessions that contain a particular gene model. We tested whether each module showed an excess or

deficit of shell genes as compared to genome averages of pan-gene occupancy using a Fisher's Exact

Test, as implemented in the R programming language (R Core Team, 2022).

Enrichment and GO/KEGG annotation of clustered genes

Gene ontology (GO) and the Kyoto Encyclopedia of Genes and Genomes (KEGG) annotations for

the *B. distachyon* 314 (Bd21 acccession) v.3.1 reference genome were retrieved

(http://phytozome.jgi.doe.gov/; IBI 2010). Gene lists were tested for functional enrichments with the

PANTHER (Protein ANalysis THrough Evolutionary Relationships) overrepresentation test

(http://www.pantherdb.org). The original *B. distachyon* 314 (Bd21 accession) v.3.1 gene ids were

converted to v.3.0 with help from Ensembl Plants (Howe et al., 2020) to match those in PANTHER16.0

(Mi et al., 2021). Tests were conducted on all genes and on both conditions -- Drought and Water --

applying the Fisher's Exact test with False Discovery Rate (FDR) multiple test correction. This analysis

was applied on different data sets: all genes, and pan-genome core, soft-core, and shell genes for each co-

expressed module.

Drought versus Watered modular structure preservation and comparison between consensus and
set-specific modules

Permutation tests were performed to check for preservation of the module topology in the Drought

(discovery data) and the Water (test data) networks using both the approach of Langfelder et al.

(Langfelder, Luo, Oldham, & Horvath, 2011) as well as the modulePreservation function of the NetRep

(Ritchie et al. 2016) R package with null="all" (include all nodes) option for RNA-Seq data. All NetRep

test statistics (Module coherence, Average node contribution, Concordance of node contributions, Density

of correlation structure, Concordance of correlation structure, Average edge weight, and Concordance of

weighted degree) were evaluated through permutation analysis; therefore, a module was considered

preserved if all the statistics had a permutation test P-value < 0.01. Searching for modules that could play

a role in drought response, we focused on Drought modules that were unpreserved in the Water network

(P-value > 0.01 in at least 1 of the seven statistics presented in Ritchie et al. (2016). The Consensus

modules (Cons) and the respective relationships between Consensus and Drought (D) and Water (W)

modules were performed as described in Langfelder and Horvath (2008).

Annotations of upstream DNA motifs in the co-expression modules

Genes assigned to modules in the Drought and Water networks were further analyzed with the

objective of discovering DNA motifs putatively involved in their expression in each module. Motif

analysis was carried out using a protocol based on RSAT::Plants (Ksouri et al., 2021). This approach

allowed us to discover DNA motifs enriched in the promoter regions of co-expressed genes and to match

them to curated signatures of experimentally described transcription factors. First, -500 bp (upstream) to

+200 bp (downstream) sequences around the transcription start sites of the genes detected in each module

and 50 random negative controls from other gene promoter regions of equal size were extracted from the

B. distachyon Bd21 v3.0 (Ensembl Plants version 46) reference genome. Then, the peak-motifs (Thomas-

Chollier et al., 2012) option was used to discover enriched motifs applying a 2^{nd} order Markov genomic

model, and GO enrichment was computed for them. The analyses generated a report with links to similar

curated motifs in the database footprintDB as scored with normalized correlation (Ncor) (Sebastian and

Contreras-Moreira 2014). For each module a highly supported DNA motif was selected according to

Ncor, e-value, and the number of sites (i.e., putative cis-regulatory elements, CREs) used to compile the

motif. The matrix-scan tool (Turatsinze et al., 2008), with a weight threshold set to 70% of the motif

length (60% for the small module D8), was used to scan the discovered motifs and to identify individual

genes within each module harboring putative CREs.

All the protein sequences predicted for each co-expression module were analyzed using iTAK (Plant

Transcription factor & Protein Kinase Identifier and Classifier) online (v1.6) (Zheng et al., 2016) to

annotate their respective transcription factors, transcriptional regulators and related protein kinases

(classification system by Lehti-Shiu and Shiu (2012)).

Differentially expressed (DE) isoforms/genes

In order to determine how many isoforms and genes were differentially expressed between the two

treatments (D vs W), the two data sets were analyzed through the sleuth_result function (Pimentel et al.,

2017). This function computes likelihood ratio tests (LRT) for null (no treatment effect) and alternative

(treatment effect) hypotheses, attending to the full and reduced fitted models. A significant q-value ≤ 1E-

6 threshold was fixed to detect DE isoforms. The 50 most differentially expressed genes (25 up-regulated

and 25 down-regulated) were classified based on fold-change of the average TPM values between the

drought and water treatments.

Data availability

291   The *Brachypodium distachyon* filtered RNA-Seq data were deposited in the ENA (European

292   Nucleotide Archive; https://www.ebi.ac.uk/ena) with the consecutive accession numbers from

293   ERR6133302 to ERR6133575. All the R scripts, supplementary Excel files, TPM counts and adjacency

294   matrices are available in the Github repository identified by the following doi: xxxxxxxxxx created using

295   Zenodo.

RESULTS

Our RNA-Seq dataset comprised 121 Drought and 108 Water samples, obtained from 33 inbred

accessions of *B. distachyon* (Figure 1). After the filtering and quality control steps, we identified 16,386

transcripts in our analyses comprising 4,941 isoforms of 3,789 differentially expressed (DE) genes

between the water and drought treatments (Supplementary files S1; S2). Of these genes, 2,808 were

upregulated (3,591 isoforms) and 980 downregulated (1,350 isoforms) in D vs W conditions. One gene

(Bradi1g09950) showed both up-regulated (Bradi1g09950.2) and down-regulated (Bradi1g09950.1)

isoforms under drought conditions (Supplementary file S2).

Modular distribution of genes in the gene co-expression networks

The 33 genetically diverse accessions, along with random experimental variance, provided a wealth

of expression variance that was leveraged to estimate major axes of variation – co-expression modules –

from the RNA-Seq data. The modular distribution of genes in the Drought and Water gene co-expression

networks showed differences both in the number and the size of the modules (Figures S1a,b and S2a,b;

Tables 1a,b and S3a,b). The Drought co-expression network comprised 9 modules (D1-D9) containing a

total of 5,020 isoforms (min = 38, max = 2,477 isoforms per module), corresponding to 4,006 genes (min

= 27, max = 1,986 genes per module). The largest D1 module contained 15.1% of the isoforms (16.1% of

the genes) whereas two modules (D2, D3) clustered 4-6% of the isoforms and genes each and the

remaining modules clustered ≤ 2% of isoforms and genes each (Fig. S1a; Table 1a). 11,366 isoforms

(69.4%; 8,313 genes, 67.4%) were not clustered in any Drought module using our criteria (gray or "zero"

D0 module) (Figs. S1a and S2a; Table 1a). The Water co-expression network showed 13 modules (W1-

W13) containing a total of 6,711 isoforms (min = 48, max = 1,866 isoforms per module), corresponding

to 5,407 genes (min = 40, max = 1,439 genes per module). The largest W1 contained 11.4% of the

isoforms (11.6% of the genes) whereas six modules (W2-W7) clustered over >2-6% of the isoforms and

genes and the remaining six modules ≤ 2% each (Fig. S1b; Table 1b). 9,675 isoforms (59.0%; 6,934

genes, 56.0%) were not clustered in any Water module (gray or "zero" W0 module; Figs. S1b and S2b;

Table 1b). Hereafter, co-expression modules identified in the Drought RNA-Seq dataset will be labeled

with the prefix "D" while those modules identified in the Water (control) dataset will have the "W"

prefix.

While the genes comprising co-expression modules likely represent functionally related genes,

modules are not discrete entities and many genes within a module are likely co-expressed with genes in

other modules. The largest Drought modules showed a positive (D1, D2, D3, D4, and D5) or slightly

negative (D6 and D7) mean k_{diff}, while the smallest modules had more negative values (D8 and D9)

(Table 1a). Negative k_{diff} values for a node indicate that connectivity out of the module is higher than

intra-modular connectivity. Similarly, the Water modules showed positive mean k_{diff} values in the largest

modules (W1, W3, W5, W6, W7, and W9) or slightly negative (W4), with the exception of one large and

one intermediate modules (W2 and W8) both having negative values similar to those of the smallest

modules (Table 1b). High positive linear correlations between MM values and k_{IM} values were recovered

in both Drought (Fig. S3a) and Water (Fig. S3b) networks, thus validating the criterion of high MM

(>0.9) for selecting hub genes. Collectively, these results suggest that modules of both Drought and Water

networks were statistically well-supported but also that many of their genes likely share transcriptional

activity with genes included in other modules.

Preservation and correspondence of Drought and Water networks

We tested the hypothesis that some co-expression modules are only observed under one of our two

treatment conditions using two approaches. First, a permutation test was performed using NetRep (Ritchie

et al., 2016) to test for the preservation of module topology in the Drought versus the Water networks,

defined as discovery and test data sets, respectively. This test comprises seven topological statistics on

each module and condition (drought vs water), quantifying the replication of the structural relationship

between nodes composing each module under the null hypothesis that the module of interest is not

preserved. All Drought and Water modules were topologically preserved according to the seven NetRep

statistics (permutation test P-values < 0.01). We further tested for correspondence between Drought (D)

and Water (W) set-specific and Drought-Water Consensus (Cons) co-expression modules using WGCNA

(Fig. 2a,b). Consensus modules are those shared by two or more networks (Langfelder & Horvath, 2007).

We found that several Drought specific modules were comprised of no or very few isoforms included in

the Consensus modules (Fig. 2a). Thus, the Drought modules D5 (green), D7 (black) and D9 (magenta)

did not show a clear correspondence with consensus modules. The seemingly conflicting results between

NetRep – which detected module preservation between Drought and Water modules – and WGCNA –

which identified three modules that were not preserved – reflected the different sensitivities of these two

approaches to changing co-expression relationships within modules. This suggested that all modules were

preserved in a broad sense between treatments, but that nodes within modules D5, D7, and D9 could have

slightly different co-expression relationships than those observed in the corresponding modules in the

Water treatment. We also found Water specific modules (Fig. 2b), W1 (turquoise), W6 (red), W8 (pink),

and W12 (tan), that did not overlap with the Consensus modules or overlapped only with the non-co-

expressed gray module. W1 and W6 overlapped with the gray consensus module, whereas the smaller

modules W8 and W12 did not overlap with any consensus module.

Seven modules in the Drought network and eleven in the Water network showed a significant GO

term enrichment (Table 2a,b; Supplementary file S3). Both networks shared modules enriched for

different biological processes. Potentially equivalent modules between the D and W networks were

inferred attending to the common consensus module with which they overlapped (Figure 2a,b) and their

GO enrichments (Fig. 2c). Thus, the D1 and W2 modules matched Cons1 and were GO-enriched in

nitrogen, amide, and peptide metabolic and biosynthetic processes, D2 and W10 matched Cons4 and were

enriched in the photosynthesis, D3 and W3 matched Cons3 and were enriched in processes of transport

and locations of compounds, D4 and W9 were enriched in nucleic acid metabolic processes, and D8 and

W13 in nitrogen, amide, and peptide biosynthetic and metabolic processes. However, some modules

showed enrichment in biological process unique to one or the other co-expression network. For example,

D5 is enriched in genes involved in protein folding, response to heat, temperature, and abiotic stimulus,

and W6 in genes predicted to be involved in cell wall organization or biogenesis. Collectively, these co-

expression analyses suggest that the detected modules represent groups of functionally related genes, and

that many functional relationships among genes were conserved in the Water and Drought networks.

Regulatory motifs of genes in the Drought and Water modules

We detected statistically over-represented sequence motifs upstream of the gene sequences in several

modules. These motifs represent putative *cis*-regulatory elements (CREs) located in proximal promoter

regions of co-expressed genes in the Drought (Table 3a; Fig. S4a) and Water (Table 3b; Fig. S4b)

networks. A generally low but variable proportion of genes harboring putative CREs in their proximal

promoters was detected in each module. The drought (Table 3a) and water (Table 3b) modules showed

between 3.5-54.2% and 0.1-23% of genes with the predicted CREs, respectively. Genes in the same

module shared a conserved regulatory architecture. For example, calmodulin-binding CREs were

enriched in the D4 (associated with nucleobase-containing compound, heterocycle and nucleic acid

metabolic processes GO terms) and W9 modules (also enriched for nucleobase-containing compound GO

processes, among others). We also observed enriched motifs in treatment-specific modules, like the

proximal promoters of 27.8% of the genes in module D5 contained CREs similar to those bound by

transcription factor B-3 (Table 3a) (Scharf et al., 2012), known to regulate heat shock responses in

Arabidopsis thaliana (Bechtold et al., 2013; Guo et al., 2016; Nover et al., 2001).

Additionally, all the predicted encoded proteins in each module were analysed to annotate the

transcription factors (TF), transcriptional regulators (TR), and kinases (Table S4a,b). For example, TRs

annotated in the D5 module – identified above as putatively drought-specific -- are involved in responses

to abscisic acid, heat stress, water deprivation and defense, and in zinc, chromatin, and metal ion binding

(Table S4a).

Hub nodes of the Drought and Water networks

Hub nodes were detected in both the Drought (Table 1a) and the Water (Table 1b) networks. A total

of 87 (0.53%; 1.73% excluding grey module) hub node isoforms from 72 (0.58%; 1.8% excluding grey

module) hub genes were identified in the Drought network, and 251 (1.53%; 3.74% excluding grey

module) hub node isoforms from 190 (1.53%; 3.51% excluding grey module) hub genes in the Water

network. Roughly, more than twice per-module fraction of hubs was detected in the Water network

(1.53% hub nodes/ 1.53% hub genes; Table 1b) compared to the Drought network (0.53% hub nodes/

0.58% hub genes; Table 1a).

Pan-genome analyses: occupancy of all clustered and hub genes

The studied pan-genome subset contained 34,310 pan-genome clusters (hereafter "pan-genes") which

were classified by the number of accessions with a gene model represented in a given cluster to determine

their occupancy. We found 16,057 (46.8%) core gene clusters with at least one member in every

accession. We analyzed these core genes along with the 5,642 (16.4%) clusters from the soft-core pan-

genome (occupancy in 31 or 32 accessions) to account for gene annotation errors and uncertainty with

orthology assignments. In contrast, there were 12,611 (36.8%) shell genes (i.e., observed in fewer than 31

accessions). Of the 34,310 pan-genes, 15,848 were represented by sequenced RNA tags, after our filtering

and normalizing steps. The occupancy of the expressed pan-genes was 12,137 (76.6%) core, 1,869

(11.8%) soft-core and 1,842 (11.6%) shell genes. The discrepancy between the total number of pan-genes

and that of genes represented by RNA-Seq tags may have been caused by the filtering of lowly expressed

genes. Additionally, many genes in the genome or pan-genome were likely not expressed in leaf tissue at

the developmental stage assessed herein. However, our results were consistent with prior studies which

found that pan-genome core genes are frequently more highly expressed (Gao et al., 2019; Gordon et al.,

2017; Tao et al., 2021) and therefore more likely to be sampled in RNA-Sequencing libraries (Figure S5).

The distribution of pan-gene occupancy varied considerably among modules in both Drought and

Water networks (Table 4a,b; Fig. 3a,b). Most modules in both co-expression networks showed ratios of

shell genes consistent with the genome averages (Fisher's Test; Table 4a,b), with notable exceptions.

Among Drought modules, D8 showed a considerable excess of shell genes (63.2%). Conversely, D2

(10.0%) and D3 (11.2%) each show a deficit of shell genes. Water co-expression modules showed

considerably more variation in pan-gene occupancy. Most large modules in the water network showed

significant deficits of shell genes, though W4 (27.8%), W9 (21.8%), W11 (25.5%), W12 (27.5%), and

W13 (60.0%) had an excess of shell genes relative to the genome averages (note that W11, W12, and

W13 were very small modules with fewer than 60 genes each).

To investigate the proportion of putative hub genes in each module that were members of the shell

gene sets, greater departures from genome averages were required to reach statistical significance given

the comparatively small number of these genes (Table 4c,d; Fig. 3c,d). Among the Drought modules, D8

and D9 did not show hub genes whereas other modules showed a predominance (>80%) of core or soft-

core hub genes with the exception of D7, whose four hub genes were shell genes (Table 4c). Among the

Water modules, W9 had a high proportion of shell hub genes (33.3%) as well as W4 and W12 though

these were poorly enriched with hub genes, whereas W7 showed a low percentage of hub shell genes.

Only the shell genes of module W7 are significantly under-represented among the hub genes of the Water

modules, according to the Fisher test (Table 4d).

As expected, core and soft-core genes were mostly enriched in the same GO terms as those in their

respective complete modules (Table 2a,b). Similarly, shell genes corresponded to genes involved in the

same biological process as those included in their complete modules. However, the most significant GO

term of the shell genes of D1 was photosynthesis, which was not found to be significant among core

genes. Similarly, the shell genes of D8 were enriched in photosynthesis, nitrogen, amide, and peptide

biosynthetic and metabolic processes terms, while core gene sets did not show any significant

enrichments (Table 2a,b and supplementary file S3).

We analyzed the pan-genome occupancy of genes with putative CREs in the modules with at least 10

detected genes. These corresponded predominantly to core genes (>50%) in both networks, whereas 7-

22% of the genes with predicted motifs in the Drought network and 0-27.3% in the Water network

corresponded to shell genes (Table 3a,b).

Enrichments of differentially expressed genes with a pan-genomic perspective

72.2%, 17.8% and 10% of the differentially expressed genes with upregulated isoforms in the drought

condition were core, soft-core, and shell genes, respectively. On the other hand, 67.5%, 20.9% and 11.6%

of the genes with downregulated isoforms in the drought condition were core, soft-core, and shell genes,

respectively (Table 5; supplementary file S2). 59.3% of DE isoforms (58.7% of DE genes) and 47.4% DE

isoforms (44.6% DE genes) of Drought (Table S5a) and Water (Table S5b) networks, respectively, were

not assigned to any modules (i.e., they are members of the gray or zero module). Among the hub nodes

(isoforms), 64.4% (56/87) and 34.3% (86/251) of them correspond to DE isoforms of the Drought (Table

S6a) and Water (Table S6b) networks, respectively.

Five of the nine Drought co-expression modules had a predominance (>50%) of upregulated DE

isoforms except for the large modules D2 and D3 and the small modules D8 and D9 (Table S5a).

Similarly, among Water modules, only one large module (W4) and three small modules (W10, W11 and

W13) had a predominance of downregulated DE isoforms in the drought condition compared to the water

condition (Table S5b).

21 out of the 25 most strongly upregulated genes did not cluster with other genes in the drought co-

expression network (i.e., they were members of the grey D0 module), while the majority of these strongly

upregulated genes did cluster in a module in the water network (Table S7). The 25 most strongly

upregulated genes by drought showed a range of predicted functions which included two predicted

dehydrins (Decena et al. 2021), two ABA-associated proteins, and two lipid transfer proteins (LTPs)

(Table S7). LTPs were among the most highly induced transcripts in an *Arabidopsis thaliana* experiment

that imposed very similar soil drying conditions to those imposed in the present study (Des Marais et al.,

2012). The 25 most strongly downregulated genes showed markedly different patterns than the most

strongly upregulated genes. Most of the genes cluster in D2 and W10, and four of these genes (three

clustered in D2 and one in W10) were hub genes (Table S7). Both of these modules were enriched for

genes involved in photosynthesis (Table 2a,b) and the annotations for many of these genes suggested

associations with the light reactions of photosynthesis.

DISCUSSION

Large scale transcriptome data sets have been used to construct co-expression networks for gene and

gene regulation discovery in model plant systems and crops (Aoki et al, 2007, 2016; Masalia et al., 2017;

Miao et al., 2017). The co-expression network approach allows testing hypotheses on gene functions from

their putative regulatory interactions with other functionally known genes classified in the same modules

(Mochida et al., 2011), and on links between signalling pathways and phenotypic response to

environmental stress (Des Marais et al., 2012). However, the mechanisms of and consequences for

genetic diversity in environmentally responsive gene regulatory networks are less often considered (Sun

& Dinneny, 2018). Our system-level approach allowed us to construct a drought-responsive gene co-

expression network from leaf tissue transcriptome profiles of *B. distachyon* accessions and to identify

modules of putatively co-regulated genes within it. We integrated these network hypotheses with

information about gene presence/absence variation as represented in the *B. distachyon* pan-genome.

Regulatory control of *Brachypodium* response to soil drying

B. distachyon is an annual species native to seasonally dry environments in the Mediterranean, where

it has likely evolved mechanisms to tolerate short-term soil drying during the growing season as well as

unpredictably timed end-of-season drought (López-Álvarez et al., 2015). Several past studies have

identified mechanisms of response to soil drying comprising transcriptomic, metabolic, physiological, and

developmental plasticity (Bertolini et al., 2013; Chen et al., 2016; Decena et al., 2021; Des Marais, et al.,

2016, 2017b; Fisher et al., 2016; Gordon et al., 2014; Handakumbura et al., 2019; Luo et al., 2011;

Manzaneda et al., 2015; Priest et al., 2014; Ruíz et al., 2016; Verelst et al., 2013), as well as considerable

genetic diversity of response (GxE; Des Marais et al., 2017b; Handakumbura et al., 2019). Priest et al.

(2014) provided the first transcriptomic assessment of response to drying, exposing the Bd21 accession to

a simulated severe drying stress by removing plants from soil to desiccate on a lab benchtop. These

authors observed a strong transcriptional signature of down-regulated photosynthesis, cell division, and

cell growth. Subsequent work imposing a more gradual soil drying stress in Bd21 found the opposite

pattern, directly observing sustained cell division and transcriptomic patterns of altered primary

metabolism, rather than outright downregulation (Verelst et al., 2013). Indeed, studies imposing moderate

drying on diverse *B. distachyon* accessions revealed increased leaf mass per area and greater root biomass

in several accessions in response to drying (Des Marais et al., 2017b; Handakumbura et al., 2019), both of

which require considerable investment of carbohydrates. In the present study we do, however, observe

several strongly downregulated genes with annotated functions related to the light reactions of

photosynthesis as well as RuBisCO assembly and function (Table S7).

How can we reconcile these transcriptional signatures of reduced photosynthesis with the observation

that carbohydrate-intensive processes like root growth continue under drying? The effects of soil drying

on photosynthesis are complex, and the reduction of internal leaf CO_2 (c_i) caused by stomatal closure can

affect the redox status of cells (Pinheiro & Chaves, 2011). As the Calvin Cycle reduces available CO_2, it

is also a strong sink for energy captured by the photosystems. As this sink is lowered by decreased c_i,

continued high irradiance can lead to increased expression of photoprotective mechanisms and decreased

expression of photochemistry as cells try to protect themselves from excess energy (Demmig-Adams &

Adams, 1996). As a result, studies of soil drying responses often observe decreased activity of the

photosystems and increased expression of, for example, photorespiration or other energy sinks (Wingler

et al., 1999). While we have no direct measurements of photorespiration or the quantum yield of

photosystem II in the current study, our observation of decreased expression of transcripts associated with

photosystem proteins is consistent with these mechanisms.

In light of this past evidence for an important role of photosynthesis and primary metabolism in

drying response, we focus here on Drought module 5 (D5). D5 showed a low correlation with the

Consensus modules (Fig. 2a), consistent with the hypothesis that the genes in this module are involved in

regulating plant response to drought stress. Its co-expressed genes, both core and soft-core, are involved

in protein folding, response to heat, temperature, and abiotic stimulus (Table 2a). The four DE hub nodes

(three genes) of D5 were upregulated in the drought condition compared to the water condition (Table S8)

and each of these four hubs is a core gene. Molecular chaperones, especially heat shock proteins (HSPs),

were predominantly annotated in both co-expressed and upregulated DE genes (Table S8). Related to the

presence of chaperones, the annotated DNA motifs concurred with the GO enrichment of the protein

folding and the Heat stress transcription factor B-3 (Table 3a).

Topological position of pan-genes

One key conclusion with respect to the evolution of pan-genomes and gene regulation arise from our

analysis. First, pan-genes are non-randomly assorted among co-expression modules in *B. distachyon.* This

observation suggests that the functions conferred by some co-expression modules are likely under

stronger purifying selection than those conferred by other modules. These latter modules represent

possible regulatory variation on which natural selection may act. Such a prospect was anticipated by

Wagner and Altenberg when they argued that modularity allows for evolutionary tinkering (Wagner &

Altenberg, 1996). Our results point towards a role for segregating gene copies in generating this modular

variation. This conclusion extends earlier work indicated that low-occupancy genes tend to be enriched

for functional classes of genes putatively involved in local adaptation such as disease resistance and gene

regulation (Gordon et al., 2017).

Pleiotropy can have a strong effect on the rate of molecular evolution and on the roles that functional

gene variants might play in evolutionary change. Pleiotropy is often correlated with the position of a gene

or protein in biochemical and gene regulatory networks (Erwin & Davidson, 2009; Jeong et al., 2001),

which are now readily inferred from high dimensional datasets such as the genome-wide gene co-

expression networks studies herein. Here, we considered the case of potentially large-effect mutations –

segregating gene copies identified from a grass pan-genome – and ask whether such "pan-genes" are

unevenly distributed in gene co-expression networks. Focusing on the well-Watered (control)

environment, we find that shell genes – pan-genes found in fewer than 31 of our studied accessions -- are

statistically under-represented among the genes in five of the six largest (in terms of total number of

genes) co-expression modules (Table 4b). These large modules are enriched for GO terms comprising

essential processes such as protein synthesis, primary metabolism, various processes related to

phosphorus metabolism and signalling, and cell wall organization (Table 2b). Moreover, shell genes are

generally under-represented among module hub genes (diagnosed as those whose expression most highly

correlated with the module as a whole, and thus possibly the most topologically connected among genes

in a module) in these five Water modules (Table 4d). Collectively, these results support the hypothesis

that core pan-genome genes are centrally located in gene co-expression networks and involved in

biological processes likely to be under strong purifying selection.

Water module 4 (W4), comprising 590 genes, is the only large water module that is statistically

enriched for shell genes (Table 4b). Shell genes in W4 are associated with a range of GO terms including

processes related to photosynthesis (Table 2b). In general, the lists of shell genes in modules do not tend

to have strong GO enrichments, perhaps owing to the relatively small numbers of genes in these lists.

Interestingly, among the Drought modules, the only module for which shell genes do have an enrichment

(D8; Table 4a) includes GO terms associated with photosynthesis (Table 2a). Shell genes represent genes

found in some sampled accessions but missing in others, suggesting that *B. distachyon* may harbor

genetic diversity in molecular pathways related to photosynthesis. Whether these segregating variants

represent adaptive genetic diversity reflecting the broad geographical coverage of our sampling, or simply

mildly deleterious copy number variants on their way to being lost will require a larger population

sample. Previously, we demonstrated significant genetic variation among the same *Brachypodium*

accessions used herein for leaf carbon content, leaf C:N ratios, and water use efficiency (WUE; Des

Marais, et al.; 2017b). Among these, WUE was significantly associated with principal components

summarizing climate diversity; it is possible that some of the segregating variation in photosynthesis gene

presence/absence is involved in local adaptation to climate.

The co-expression network approach we employed here, in concert with a modern pan-genomic

perspective on segregating genetic variation, allows us to identify subsets of genes involved in core

metabolic processes such as photosynthesis, suggesting possible candidate genes regulating natural

genetic variation in resource assimilation and growth. These genes make attractive targets for further

hypothesis testing via genome editing. Collectively, our work demonstrates the importance of accounting

for both gene copy number variation and regulatory interactions in studying genome function and

evolution.

ACKNOWLEDGEMENTS

We thank R. Hopkins, E. Sukamtoh, J. Bonnette, and B. Whitaker for assistance with data collection. This

work was supported by the USDA (NIFA-2011-67012-30663) to D.L.D., NSF (IOS-0922457) to T.E.J.,

the Spanish Ministry of Science and Innovation CGL2016-79790-P and PID2019-108195GB-I00,

University of Zaragoza UZ2016_TEC02 grant projects to P.C., and the Joint Genome Institute

FP00006675 contract to P.C. and FP00006746 to D.L.D.. R.S. was funded by a Spanish Ministry of

Economy and Competitiveness (Mineco) FPI PhD fellowship, Mineco and Ibercaja-CAI mobility grants

and Instituto de Estudios Altoaragoneses grant. B.C.M. was funded by Fundación ARAID. P.C. and R.S.

were partially funded by a European Social Fund/Spanish Aragón Government Bioflora grant.

REFERENCES

Albert, R., Jeong, H., & Barabási, A.-L. (2000). Error and Attack Tolerance of Complex Networks.

Nature, *406*, 378–382. doi: 10.1038/35019019

Alonge, M., Wang, X., Benoit, M., Soyk, S., Pereira, L., Zhang, L., … Lippman, Z. B. (2020). Major

Impacts of Widespread Structural Variation on Gene Expression and Crop Improvement in Tomato.

Cell, *182*(1), 145-161.e23. doi: 10.1016/j.cell.2020.05.021

Aoki, K., Ogata, Y., & Shibata, D. (2007). Approaches for Extracting Practical Information from Gene

Co-expression Networks in Plant Biology. *Plant Cell Physiology*, *48*(3), 381–390. doi:

10.1093/pcp/pcm013

Aoki, Y., Okamura, Y., Tadaka, S., Kinoshita, K., & Obayashi, T. (2016). ATTED-II in 2016: A Plant

Coexpression Database Towards Lineage-Specific Coexpression. *Plants and Cell Physiology*,

57(1), e5(1-9). doi: 10.1093/pcp/pcv165

Bayer, P. E., Golicz, A. A., Scheben, A., Batley, J., & Edwards, D. (2020). Plant pan-genomes are the

new reference. *Nature Plants*, *6*(8), 914–920. doi: 10.1038/s41477-020-0733-0

Bechtold, U., Albihlal, W. S., Lawson, T., Fryer, M. J., Sparrow, P. A. C., Richard, F., … Mullineaux, P.

M. (2013). Arabidopsis HEAT SHOCK TRANSCRIPTION FACTORA1b overexpression

enhances water productivity, resistance to drought, and infection. *Journal of Experimental Botany*,

64(11), 3467–3481. doi: 10.1093/jxb/ert185

Benfey, P. N., & Mitchell-Olds, T. (2008). From Genotype to Phenotype: Systems Biology Meets Natural

Variation. *Science*, *320*, 495–497. doi: 10.1161/circgen.117.001946

Bertolini, E., Verelst, W., Horner, D. S., Gianfranceschi, L., Piccolo, V., Inzé, D., … Mica, E. (2013).

Addressing the role of microRNAs in reprogramming leaf growth during drought stress in

Brachypodium distachyon. *Molecular Plant*, *6*(2), 423–443. doi: 10.1093/mp/sss160

Bohnert, H. J., Nelson, D. E., & Jensenayb, R. G. (1995). Adaptations to Environmental Stresses. *The

Plant Cell*, *7*, 1099–1111.

Bolger, A. M., Lohse, M., & Usadel, B. (2014). Trimmomatic: A flexible trimmer for Illumina sequence

data. *Bioinformatics*, *30*(15), 2114–2120.

Borah, P., Sharma, E., Kaur, A., Chandel, G., Mohapatra, T., Kapoor, S., & Khurana, J. (2017). Analysis

of drought-responsive signalling network in two contrasting rice cultivars using transcriptome-

based approach. *Scientific Report*, *42131*, 1–21. doi: 10.1038/srep42131

Bray, N. L., Pimentel, H., Melsted, P., & Pachter, L. (2016). Near-optimal probabilistic RNA-seq

quantification. *Nature Biotechnology*, *34*(5), 525–527. doi: 10.1038/nbt.3519

Carlson, M. R. J., Zhang, B., Fang, Z., Mischel, P. S., Horvath, S., & Nelson, S. F. (2006). Gene

connectivity, function, and sequence conservation: predictions from modular yeast co-expression

networks. *BMC Genomics*, *7*(40), 1–15. doi: 10.1186/1471-2164-7-40

Catalán, P., Chalhoub, B., Chochois, V., Garvin, D. F., Hasterok, R., Manzaneda, A. J., … Voxeur, A.
(2014). Update on the genomics and basic biology of Brachypodium. *Trends in Plant Science*,
19(7), 414–418. doi: 10.1016/j.tplants.2014.05.002

Catalán, P., López-Alvarez, D., Díaz-Pérez, A., Sancho, R., & López-Herranz, M. L. (2016). Phylogeny
and Evolution of the Genus Brachypodium. In J P Vogel (Ed.), *Genetics and genomics of*
Brachypodium. Plant Genetics and Genomics: Crops Models (pp. 9–38). Springer. doi:
10.1007/7397

Chaves, M. M., Maroco, J. P., & Pereira, J. S. (2003). Understanding plant responses to drought — from
genes to whole plant. *Functional Plant Biology*, *30*, 239–264. doi: 10.1071/FP02076

Chen, L., Han, J., Deng, X., Tan, S., Li, L., Li, L., … Zhang, W. (2016). Expansion and stress responses
of AP2/EREBP superfamily in Brachypodium Distachyon. *Scientific Reports*, *6*(January), 1–14.
doi: 10.1038/srep21623

Contreras-Moreira, B., Cantalapiedra, C. P., García-Pereira, M. J., Gordon, S. P., Vogel, J. P., Igartua, E.,
… Vinuesa, P. (2017). Analysis of Plant Pan-Genomes and Transcriptomes with GET _
HOMOLOGUES-EST, a Clustering Solution for Sequences of the Same Species. *Frontiers in*
Genetics, *8*(184), 1–16. doi: 10.3389/fpls.2017.00184

Decena, M. A., Gálvez-Rojas, S., Agostini, F., Sancho, R., Contreras-Moreira, B., Des Marais, D. L., …
Catalán, P. (2021). Comparative genomics, evolution, and drought-induced expression of dehydrin
genes in model Brachypodium grasses. *Plants*, *10*(2664), 1–29. doi: 10.3390/plants10122664

Demmig-Adams, B., & Adams, W. W. (1996). The role of xanthophyll cycle carotenoids in the protection
of photosynthesis. *Trends in Plant Science*, *1*(1), 21–26. doi: 10.1016/S1360-1385(96)80019-7

Des Marais, D. L., Guerrero, R. F., Lasky, J. R., & Scarpino, S. V. (2017a). Topological features of gene
regulatory networks predict patterns of natural diversity in environmental response. *Proceedings of*
the Royal Society B: Biological Sciences, *284*(1856). doi: 10.1098/rspb.2017.0914

Des Marais, D. L., & Juenger, T. E. (2016). Brachypodium and the Abiotic Environment. In John P Vogel
(Ed.), *Genetics and genomics of Brachypodium. Plant Genetics and Genomics: Crops Models,*
volume 18 (pp. 291–311). Switzerland: Springer.

Des Marais, D. L., Lasky, J. R., Verslues, P. E., Chang, T. Z., & Juenger, T. E. (2017b). Interactive
effects of water limitation and elevated temperature on the physiology, development and fitness of
diverse accessions of Brachypodium distachyon. *New Phytologist*, *214*, 132–144. doi:
10.1111/nph.14316

Des Marais, D. L., Mckay, J. K., Richards, J. H., Sen, S., Wayne, T., & Juenger, T. E. (2012).

Physiological Genomics of Response to Soil Drying in Diverse Arabidopsis Accessions. *The Plant*

Cell, *24*, 893–914. doi: 10.1105/tpc.112.096180

Des Marais, D. L., Razzaque, S., Hernandez, K. M., Garvin, D. F., & Juenger, T. E. (2016). Quantitative

trait loci associated with natural diversity in water-use efficiency and response to soil drying in

Brachypodium distachyon. *Plant Science*, *251*, 2–11. doi: 10.1016/j.plantsci.2016.03.010

Dong, J., & Horvath, S. (2007). Understanding network concepts in modules. *BMC Systems Biology*,

1(24), 1–20. doi: 10.1186/1752-0509-1-24

Erwin, D. H., & Davidson, E. H. (2009). The evolution of hierarchical gene regulatory networks. *Nature*

Reviews Genetics, *10*(2), 141–148. doi: 10.1038/nrg2499

Filiz, E., Ozdemir, B. S., Budak, F., Vogel, J. P., Tuna, M., & Budak, H. (2009). Molecular,

morphological, and cytological analysis of diverse Brachypodium distachyon inbred lines. *Genome*,

52(10), 876–890. doi: 10.1139/g09-062

Fisher, L. H., Han, J., Corke, F. M., Akinyemi, A., Didion, T., Nielsen, K. K., … Bosch, M. (2016).

Linking Dynamic Phenotyping with Metabolite Analysis to Study Natural Variation in Drought

Responses of Brachypodium distachyon. *Frontiers in Plant Science*, *7*, 1–15. doi:

10.3389/fpls.2016.01751

Fisher, R. A. (1930). *The genetical theory of natural selection*. Oxford Clarendon Press. doi:

10.5962/bhl.title.27468

Gao, L., Gonda, I., Sun, H., Ma, Q., Bao, K., Tieman, D. M., … Fei, Z. (2019). The tomato pan-genome

uncovers new genes and a rare allele regulating fruit flavor. *Nature Genetics*, *51*(6), 1044–1051.

doi: 10.1038/s41588-019-0410-2

Gibson, G. (2016). On the Evaluation of Module Preservation. *Cell Systems*, *3*, 17–19. doi:

10.1016/j.cels.2016.07.009

Gordon, S. P., Contreras-Moreira, B., Woods, D. P., Des Marais, D. L., Burgess, D., Shu, S., … Vogel, J.

P. (2017). Extensive gene content variation in the Brachypodium distachyon pan-genome correlates

with population structure. *Nature Communications*, *8*(2184). doi: 10.1038/s41467-017-02292-8

Gordon, S. P., Priest, H., Des Marais, D. L., Schackwitz, W., Figueroa, M., Martin, J., … Vogel, J. P.

(2014). Genome diversity in Brachypodium distachyon: Deep sequencing of highly diverse inbred

lines. *The Plant Journal*, *79*, 361–374. doi: 10.1111/tpj.12569

Guelzim, N., Bottani, S., Bourgine, P., & Képès, F. (2002). Topological and causal structure of the yeast

transcriptional regulatory network. *Nature Genetics*, *31*(1), 60–63. doi: 10.1038/ng873

Guo, M., Liu, J. H., Ma, X., Luo, D. X., Gong, Z. H., & Lu, M. H. (2016). The plant heat stress

transcription factors (HSFS): Structure, regulation, and function in response to abiotic stresses.

Frontiers in Plant Science, *7*(FEB2016). doi: 10.3389/fpls.2016.00114

Haberer, G., Kamal, N., Bauer, E., Gundlach, H., Fischer, I., Seidel, M. A., … Mayer, K. F. X. (2020).

European maize genomes highlight intraspecies variation in repeat and gene content. *Nature*

Genetics, *52*(9), 950–957. doi: 10.1038/s41588-020-0671-9

Handakumbura, P. P., Stanfill, B., Rivas-Ubach, A., Fortin, D., Vogel, J. P., & Jansson, C. (2019).

Metabotyping as a Stopover in Genome-to-Phenome Mapping. *Scientific Reports*, *9*(1), 1–12. doi:

10.1038/s41598-019-38483-0

Hasterok, R., P. Catalan, S. P. Hazen, A. C. Roulin, J. P. Vogel, K. Wang and L. A. J. Mur (2022).

"*Brachypodium*: Twenty years as a grass biology model system; the way forward?" <u>Trends in Plant Science</u>.

Hayano-Kanashiro, C., Calderón-Vázquez, C., Ibarra-Laclette, E., Herrera-Estrella, L., & Simpson, J.

(2009). Analysis of Gene Expression and Physiological Responses in Three Mexican Maize

Landraces under Drought Stress and Recovery Irrigation. *PLoS ONE*, *4*(10), e7531. doi:

10.1371/journal.pone.0007531

Howe, K. L., Contreras-Moreira, B., De Silva, N., Maslen, G., Akanni, W., Allen, J., … Flicek, P. (2020).

Ensembl Genomes 2020-enabling non-vertebrate genomic research. *Nucleic Acids Research*,

48(D1), D689–D695. doi: 10.1093/nar/gkz890

Hübner, S., Bercovich, N., Todesco, M., Mandel, J. R., Odenheimer, J., Ziegler, E., … Rieseberg, L. H.

(2019). Sunflower pan-genome analysis shows that hybridization altered gene content and disease

resistance. *Nature Plants*, *5*(1), 54–62. doi: 10.1038/s41477-018-0329-0

IBI. (2010). Genome sequencing and analysis of the model grass Brachypodium distachyon. *Nature*, *463*,

763–768. doi: 10.1038/nature08747

Janiak, A., Kwa, M., & Szarejko, I. (2015). Gene expression regulation in roots under drought. *Journal of*

Experimental Botany, *67*(4), 1003–1014. doi: 10.1093/jxb/erv512

Jeong, H., Mason, S. P., Barabási, A. L., & Oltvai, Z. N. (2001). Lethality and centrality in protein

networks. *Nature*, *411*(6833), 41–42. doi: 10.1038/35075138

Juenger, T. E. (2013). Natural variation and genetic constraints on drought tolerance. *Current Opinion in*

Plant Biology, *16*(3), 274–281. doi: 10.1016/j.pbi.2013.02.001

Kaas, R. S., C. Friis, D. W. Ussery and F. M. Aarestrup (2012). "Estimating variation within the genes and
inferring the phylogeny of 186 sequenced diverse Escherichia coli genomes." BMC Genomics **13**(577):
10.1186/1471-2164-1113-1577.

Koonin, E. V., & Wolf, Y. I. (2008). Genomics of bacteria and archaea: The emerging dynamic view of
the prokaryotic world. *Nucleic Acids Research*, *36*(21), 6688–6719. doi: 10.1093/nar/gkn668

Ksouri, N., Castro-Mondragón, J. A., Montardit-Tarda, F., van Helden, J., Contreras-Moreira, B., &
Gogorcena, Y. (2021). Tuning promoter boundaries improves regulatory motif discovery in
nonmodel plants: the peach example. *Plant Physiology*, *185*(3), 1242–1258. doi:
10.1093/plphys/kiaa091

Langfelder, P., & Horvath, S. (2007). Eigengene networks for studying the relationships between co-
expression modules. *BMC Systems Biology*, *1*, 54. doi: 10.1186/1752-0509-1-54

Langfelder, P., & Horvath, S. (2008). WGCNA: an R package for weighted correlation network analysis.
BMC Bioinformatics, *9*(559), 1–13. doi: 10.1186/1471-2105-9-559

Langfelder, P., & Horvath, S. (2010). Overview of network terminology. Retrieved April 16, 2016, from
https://labs.genetics.ucla.edu/horvath/CoexpressionNetwork/ModulePreservation/Tutorials/glossary
Table.pdf

Langfelder, P., Luo, R., Oldham, M. C., & Horvath, S. (2011). Is my network module preserved and
reproducible? *PLoS Computational Biology*, *7*(1). doi: 10.1371/journal.pcbi.1001057

Lehti-Shiu, M. D., & Shiu, S. H. (2012). Diversity, classification and function of the plant protein kinase
superfamily. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *367*(1602),
2619–2639. doi: 10.1098/rstb.2012.0003

Lei, L., Goltsman, E., Goodstein, D., Wu, G. A., Rokhsar, D. S., & Vogel, J. P. (2021). Plant Pan-
Genomics Comes of Age. *Annual Review of Plant Biology*, *72*(1), 411–435. doi: 10.1146/annurev-
arplant-080720-105454

López-Álvarez, D., Manzaneda, A. J., Rey, P. J., Giraldo, P., Benavente, E., Allainguillaume, J., …
Catalan, P. (2015). Environmental niche variation and evolutionary diversification of the
Brachypodium distachyon grass complex species in their native circum-Mediterranean range.
American Journal of Botany, *102*(7), 1073–1088. doi: 10.3732/ajb.1500128

Luo, N., Liu, J., Yu, X., & Jiang, Y. (2011). Natural variation of drought response in Brachypodium
distachyon. *Physiologia Plantarum*, *141*(1), 19–29. doi: 10.1111/j.1399-3054.2010.01413.x

Manzaneda, A. J., Rey, P. J., Anderson, J. T., Raskin, E., Weiss-Lehman, C., & Mitchell-Olds, T. (2015).

Natural variation, differentiation, and genetic trade-offs of ecophysiological traits in response to

water limitation in Brachypodium distachyon and its descendent allotetraploid B. hybridum

(Poaceae). *Evolution*, *69*(10), 2689–2704. doi: 10.1111/evo.12776

Manzaneda, A. J., Rey, P. J., Bastida, J. M., Weiss-Lehman, C., Raskin, E., & Mitchell-Olds, T. (2012).

Environmental aridity is associated with cytotype segregation and polyploidy occurrence in

Brachypodium distachyon (Poaceae). *New Phytologist*, *193*(3), 797–805. doi: 10.1111/j.1469-

8137.2011.03988.x

Mao, L., Hemert, J. L. Van, Dash, S., & Dickerson, J. A. (2009). Arabidopsis gene co-expression network

and its functional modules. *BMC Bioinformatics*, *10*(346), 1–24. doi: 10.1186/1471-2105-10-346

Martínez, L. M., Fernández-ocaña, A., Rey, P. J., Salido, T., Amil-ruiz, F., & Manzaneda, A. J. (2018).

Variation in functional responses to water stress and differentiation between natural allopolyploid

populations in the Brachypodium distachyon species complex. *Annals of Botany*, *00*, 1–14. doi:

10.1093/aob/mcy037

Masalia, R. R., Bewick, A. J., & Burke, J. M. (2017). Connectivity in gene coexpression networks

negatively correlates with rates of molecular evolution in flowering plants. *PLoS ONE*, *12*(7),

e0182289. doi: https://doi.org/10.1371/journal.pone.0182289

Meyer, E., Aglyamova, G. V., & Matz, M. V. (2011). Profiling gene expression responses of coral larvae

(Acropora millepora) to elevated temperature and settlement inducers using a novel RNA-Seq

procedure. *Molecular Ecology*, *20*(17), 3599–3616. doi: 10.1111/j.1365-294X.2011.05205.x

Mi, H., Ebert, D., Muruganujan, A., Mills, C., Albou, L. P., Mushayamaha, T., & Thomas, P. D. (2021).

PANTHER version 16: A revised family classification, tree-based classification tool, enhancer

regions and extensive API. *Nucleic Acids Research*, *49*(D1), D394–D403. doi:

10.1093/nar/gkaa1106

Miao, Z., Han, Z., Zhang, T., Chen, S., & Ma, C. (2017). A systems approach to a spatio- temporal

understanding of the drought stress response in maize. *Scientific Reports*, *7*(6590), 1–14. doi:

10.1038/s41598-017-06929-y

Mochida, K., Uehara-Yamaguchi, Y., Yoshida, T., Sakurai, T., & Shinozaki, K. (2011). Global

Landscape of a Co-Expressed Gene Network in Barley and its Application to Gene Discovery in

Triticeae Crops. *Plants and Cell Physiology*, *52*(5), 785–803. doi: 10.1093/pcp/pcr035

Monroe, J. G., H. Cai and D. L. Des Marais (2021). "Diversity in non-linear responses to soil moisture shapes

evolutionary constraints in *Brachypodium*." G3 **11**(12): jkab334.

Monroe, J. G., McGovern, C., Lasky, J. R., Grogan, K., Beck, J., & McKay, J. K. (2016). Adaptation to
warmer climates by parallel functional evolution of CBF genes in Arabidopsis thaliana. *Molecular*
Ecology, *25*(15), 3632–3644. doi: 10.1111/mec.13711

Monroe, J. G., Powell, T., Price, N., Mullen, J. L., Howard, A., Evans, K., … McKay, J. K. (2018).
Drought adaptation in arabidopsis thaliana by extensive genetic loss-of-function. *ELife*, *7*, 1–18.
doi: 10.7554/eLife.41038

Mur, L. A. J., Allainguillaume, J., Catalán, P., Hasterok, R., Jenkins, G., Lesniewska, K., … Vogel, J.
(2011). Exploiting the Brachypodium tool box in cereal and grass research. *New Phytologist*,
191(2), 334–347. doi: 10.1111/j.1469-8137.2011.03748.x

Nakashima, K., Ito, Y., & Yamaguchi-Shinozaki, K. (2009). Transcriptional Regulatory Networks in
Response to Abiotic Stresses in Arabidopsis and Grasses. *Plant Physiology*, *149*, 88–95. doi:
10.1104/pp.108.129791

Nakashima, K., Yamaguchi-Shinozaki, K., & Shinozaki, K. (2014). The transcriptional regulatory
network in the drought response and its crosstalk in abiotic stress responses including drought, cold,
and heat. *Frontiers in Plant Science*, *5*(170), 1–7. doi: 10.3389/fpls.2014.00170

Nover, L., Bharti, K., Döring, P., Mishra, S. K., Ganguli, A., & Scharf, K. D. (2001). Arabidopsis and the
heat stress transcription factor world: How many heat stress transcription factors do we need? *Cell*
Stress and Chaperones, *6*(3), 177–189. doi: 10.1379/1466-1268(2001)006<0177:aathst>2.0.co;2

Paaby, A. B., & Rockman, M. V. (2014). Cryptic genetic variation: Evolution's hidden substrate. *Nature*
Reviews Genetics, *15*(4), 247–258. doi: 10.1038/nrg3688

Pimentel, H., Bray, N. L., Puente, S., Melsted, P., & Pachter, L. (2017). Differential analysis of RNA-seq
incorporating quantification uncertainty. *Nature Methods*, *14*(7), 687–690. doi:
10.1038/nmeth.4324

Pinheiro, C., & Chaves, M. M. (2011). Photosynthesis and drought: Can we make metabolic connections
from available data? *Journal of Experimental Botany*, *62*(3), 869–882. doi: 10.1093/jxb/erq340

Porth, I., Klápště, J., McKown, A. D., La Mantia, J., Hamelin, R. C., Skyba, O., … Douglas, C. J. (2014).
Extensive functional pleiotropy of REVOLUTA substantiated through forward genetics. *Plant*
Physiology, *164*(2), 548–554. doi: 10.1104/pp.113.228783

Priest, H. D., Fox, S. E., Rowley, E. R., Murray, J. R., Michael, T. P., & Mockler, T. C. (2014). Analysis
of Global Gene Expression in Brachypodium distachyon Reveals Extensive Network Plasticity in
Response to Abiotic Stress. *PLoS ONE*, *9*(1), e87499. doi: 10.1371/journal.pone.0087499

R Core Team. (2022). *R: A language and environment for statistical computing. R Foundation for*
Statistical Computing. Vienna, Austria.

Ritchie, S. C., Watts, S., Fearnley, L. G., Holt, K. E., Abraham, G., & Inouye, M. (2016). A Scalable
Permutation Approach Reveals Replication and Preservation Patterns of Network Modules in Large
Datasets. *Cell Systems*, *3*, 71–82. doi: 10.1016/j.cels.2016.06.012

Ruíz, M., Quemada, M., García, R. M., Carrillo, J. M., & Benavente, E. (2016). Use of thermographic
imaging to screen for drought-tolerant genotypes in Brachypodium distachyon. *Crop and Pasture*
Science, *67*(1), 99–108. doi: 10.1071/CP15134

Scharf, K. D., Berberich, T., Ebersberger, I., & Nover, L. (2012). The plant heat stress transcription factor
(Hsf) family: Structure, function and evolution. *Biochimica et Biophysica Acta - Gene Regulatory*
Mechanisms, *1819*(2), 104–119. doi: 10.1016/j.bbagrm.2011.10.002

Scholthof, K.-B. G., Irigoyen, S., Catalán, P., & Mandadi, K. K. (2018). Brachypodium: A monocot grass
model system for plant biology. *Plant Cell*, *30*, 1673–1694.

Sebastian, A., & Contreras-Moreira, B. (2014). footprintDB: a database of transcription factors with
annotated cis elements and binding interfaces. *Bioinformatics*, *30*(2), 258–265. doi:
10.1093/bioinformatics/btt663

Skalska, A., Stritt, C., Wyler, M., Williams, H. W., Vickers, M., Han, J., … Mur, L. A. J. (2020). Genetic
and methylome variation in Turkish Brachypodium distachyon accessions differentiate two
geographically distinct subpopulations. *International Journal of Molecular Sciences*, *21*(18), 1–17.
doi: 10.3390/ijms21186700

Stuart, J. M., Segal, E., Koller, D., & Kim, S. K. (2003). A Gene Coexpression Network for Global
Discovery of Conserved Genetic Modules. *Science*, *302*, 249–255.

Sun, Y., & Dinneny, J. R. (2018). Q&A: How do gene regulatory networks control environmental
responses in plants? *BMC Biology*, *16*(38), 18–21. doi: 10.1186/s12915-018-0506-7

Tandonnet, S., & Torres, T. T. (2017). Traditional versus 3′ RNA-seq in a non-model species. *Genomics*
Data, *11*, 9–16. doi: 10.1016/j.gdata.2016.11.002

Tao, Y., Luo, H., Xu, J., Cruickshank, A., Zhao, X., Teng, F., … Mace, E. (2021). Extensive variation
within the pan-genome of cultivated and wild sorghum. *Nature Plants*, *7*(6), 766–773. doi:
10.1038/s41477-021-00925-x

Thomas-Chollier, M., Herrmann, C., Defrance, M., Sand, O., Thieffry, D., & Van Helden, J. (2012).
RSAT peak-motifs: Motif analysis in full-size ChIP-seq datasets. *Nucleic Acids Research*, *40*(4).

doi: 10.1093/nar/gkr1104

Turatsinze, J. V., Thomas-Chollier, M., Defrance, M., & van Helden, J. (2008). Using RSAT to scan
genome sequences for transcription factor binding sites and cis-regulatory modules. *Nature*
Protocols, *3*(10), 1578–1588. doi: 10.1038/nprot.2008.97

Verelst, W., Bertolini, E., De Bodt, S., Vandepoele, K., Demeulenaere, M., Pè, M. E., & Inzé, D. (2013).
Molecular and physiological analysis of growth-limiting drought stress in Brachypodium
distachyon leaves. *Molecular Plant*, *6*(2), 311–322. doi: 10.1093/mp/sss098

Vogel, John P., Garvin, D. F., Leong, O. M., & Hayden, D. M. (2006). Agrobacterium-mediated
transformation and inbred line development in the model grass Brachypodium distachyon. *Plant*
Cell, Tissue and Organ Culture, *84*(2), 199–211. doi: 10.1007/s11240-005-9023-9

Vogel, John P., Garvin, D. F., Mockler, T. C., Schmutz, J., Rokhsar, D., Bevan, M. W., … Baxter, I.
(2010). Genome sequencing and analysis of the model grass Brachypodium distachyon. *Nature*,
463(7282), 763–768. doi: 10.1038/nature08747

Vogel, John P, Tuna, M., Budak, H., Huo, N., Gu, Y. Q., & Steinwand, M. A. (2009). Development of
SSR markers and analysis of diversity in Turkish populations of Brachypodium distachyon. *BMC*
Plant Biology, *9*, 88. doi: 10.1186/1471-2229-9-88

Wagner, G. P., & Altenberg, L. (1996). PERSPECTIVE: COMPLEX ADAPTATIONS AND THE
EVOLUTION OF EVOLVABILITY. *Evolution*, *50*(3), 967–976. doi: 10.1111/j.1558-
5646.1996.tb02339.x

Wingler, A., Quick, W. P., Bungard, R. A., Bailey, K. J., Lea, P. J., & Leegood, R. C. (1999). The role of
photorespiration during drought stress: An analysis utilizing barley mutants with reduced activities
of photorespiratory enzymes. *Plant, Cell and Environment*, *22*(4), 361–373. doi: 10.1046/j.1365-
3040.1999.00410.x

Wolfe, C. J., Kohane, I. S., & Butte, A. J. (2005). Systematic survery reveals general applicability of
"guilt-by-association" within gene coexpression networks. *BMC Bioinformatics*, *6*(227), 1–10. doi:
10.1186/1471-2105-6-227

Zhang, B., & Horvath, S. (2005). A General Framework for Weighted Gene Co-expression Network
Analysis. *Statistical Applications in Genetics and Molecular Biology*, *4*(1), Article 17.

Zheng, Y., Jiao, C., Sun, H., Rosli, H. G., Pombo, M. A., Zhang, P., … Fei, Z. (2016). iTAK: A Program
for Genome-wide Prediction and Classification of Plant Transcription Factors, Transcriptional
Regulators, and Protein Kinases. *Molecular Plant*, *9*(12), 1667–1670. doi:

10.1016/j.molp.2016.09.014

DATA ACCESSIBILITY AND BENEFIT-SHARING

Data Accessibility

RNA Sequence data are available at the ENA (European Nucleotide Archive; https://www.ebi.ac.uk/ena)

with the consecutive accession numbers from ERR6133302 to ERR6133575.

All the R scripts, supplementary Excel files, TPM counts and adjacency matrices are available in the

Github repository identified by the following doi: xxxxxxxxx created using Zenodo.

Experimental details and metadata for samples studied herein were reported previously (Des Marais et al.

2017b).

Benefit Sharing

All accessions used in the current study originated from collections generated by the *Brachypodium*

research community over several decades. Benefits from this research accrue from the sharing of our data

and results on public databases as described above.

AUTHOR CONTRIBUTION

D.L.D. and T.E.J. designed the research. D.L.D. performed the experimental manipulations and generated

the data. D.L.D., R.S., and B.C.-M. performed the analyses. D.L.D. and R.S. wrote the primary draft of

the manuscript, and all authors contributed to the final draft of the manuscript.

TABLES

Table 1. Statistics of the number and percentage of isoforms (all and hub isoforms) and genes (all and

hub genes), and mean k_{diff} (the difference between intra- and inter-modular connectivity) per module for

each Drought **(a)** and Water **(b)** co-expression networks. The quantity of total counts of genes in drought

and water networks is different because we counted the genes for each module independently. When

several isoforms of the same gene were clustered in different modules, the gene was counted multiple

times; however, it was counted only once if it clustered in the same module. For example, if D1 has three

isoforms (Bradi1g10000.1, Bradi1g10000.2 and Bradi1g10000.3) of the same gene, it was counted only

once (Bradi1g10000); when isoforms were found in different modules (e.g., Bradi1g10000.1 in D1,

Bradi1g10000.2 in D2, Bradi1g10000.3 in D3), the gene was counted three times.

(a)

Drought modules		isoforms (nodes) per module		genes per module		mean Kdiff
		all	hub (KME>0.9)	all	hub (KME>0.9)	
D0	Gray	11366 (69.4%)	nd	8313 (67.4%)	nd	nd
D1	Turquoise	2477 (15.1%)	4 (0.16%)	1986 (16.1%)	3 (0.15%)	7.9
D2	Blue	979 (6.0%)	20 (2.04%)	750 (6.1%)	13 (1.73%)	3.7
D3	Brown	750 (4.6%)	41 (5.47%)	627 (5.1%)	38 (6.06%)	10.1
D4	Yellow	318 (1.9%)	4 (1.26%)	258 (2.1%)	4 (1.55%)	3.7
D5	Green	214 (1.3%)	4 (1.87%)	169 (1.4%)	3 (1.78%)	0.8
D6	Red	111 (0.7%)	10 (9.01%)	95 (0.8%)	7 (7.37%)	-1.5
D7	Black	68 (0.4%)	4 (5.88%)	48 (0.4%)	4 (8.3%)	-0.2
D8	Pink	65 (0.4%)	0 (0%)	57 (0.5%)	0 (0%)	-5.9
D9	Magenta	38 (0.2%)	0 (0%)	27 (0.2%)	0 (0%)	-8.8
Total counts		**16386**	**87 (0.53%)**	**12330**	**72 (0.58%)**	
Total unique counts		**16386**	**87 (0.53%)**	**12137**	**72 (0.59%)**	
Total unique counts (excluding gray module)		**5020**	**87 (1.73%)**	**4006**	**72 (1.80%)**	

909   **(b)**

Water modules		isoforms (nodes) per module		genes per module		mean Kdiff
		all	hub (KME>0.9)	all	hub (KME>0.9)	
W0	**Gray**	9675 (59.0%)	nd	6934 (56.0%)	nd	nd
W1	**Turquoise**	1866 (11.4%)	149 (7.98%)	1439 (11.6%)	115 (7.99%)	26.8
W2	**Blue**	870 (5.3%)	4 (0.46%)	727 (5.9%)	3 (0.41%)	-3.7
W3	**Brown**	827 (5.0%)	2 (0.24%)	696 (5.6%)	1 (0.14%)	0.3
W4	**Yellow**	701 (4.3%)	2 (0.29%)	590 (4.8%)	2 (0.34%)	-0.7
W5	**Green**	607 (3.7%)	12 (1.98%)	509 (4.1%)	10 (1.96%)	3.1
W6	**Red**	435 (2.7%)	9 (2.07%)	358 (2.9%)	8 (2.08%)	2.1
W7	**Black**	390 (2.4%)	26 (6.67%)	313 (2.5%)	23 (7.35%)	4.6
W8	**Pink**	288 (1.7%)	2 (0.69%)	245 (2.0%)	2 (0.82%)	-2.2
W9	**Magenta**	282 (1.7%)	22 (7.80)	239 (1.9%)	21 (8.79%)	5.8
W10	**Purple**	246 (1.5%)	5 (2.03%)	185 (1.5%)	2 (1.08%)	-1.4
W11	**Greenyellow**	89 (0.5%)	8 (8.99%)	55 (0.4%)	1 (1.81%)	-3.1
W12	**Tan**	62 (0.4%)	1 (1.61%)	51 (0.4%)	1 (1.96%)	-0.9
W13	**Salmon**	48 (0.3%)	9 (18.75%)	40 (0.3%)	1 (2.5%)	-1.6
Total counts		**16386**	**251 (1.53%)**	**12381**	**190 (1.53%)**	
Total unique counts		**16386**	**251 (1.53%)**	**12137**	**190 (1.57%)**	
Total unique counts (excluding gray module)		**6711**	**251 (3.74%)**	**5407**	**190 (3.51%)**	

Table 2. Summary of the enrichment analysis according to the statistically significant GO biological process for the genes (all, core, soft-core and shell genes) clustered in the Drought (D) (**a**) and Water (W) (**b**) modules applying the statistical overrepresentation test of Panther (http://pantherdb.org/) tool. The biological processes are summarized according to the lowest False Discovery Rate (FDR) values (see Supplementary file S3).

(a)

Modules		All genes	core genes	soft-core genes	Shell genes
D1	turquoise	nitrogen, amide and peptide metabolic and biosynthetic processes	nitrogen, amide and peptide metabolic and biosynthetic processes	nitrogen, amide and peptide metabolic and biosynthetic processes	photosynthesis, nitrogen, amide and peptide metabolic and biosynthetic processes
D2	blue	photosynthesis	photosynthesis	photosynthesis	*No statistically significant terms*
D3	brown	organonitrogen compound metabolic process, location/transport and phosphorylation	organonitrogen compound metabolic process, location/transport and phosphorylation	*No statistically significant terms*	*No statistically significant terms*
D4	yellow	nucleobase-containing compound, heterocycle and nucleic acid metabolic processes	*No statistically significant terms*	*No statistically significant terms*	*No statistically significant terms*
D5	green	protein folding, response to heat, temperature and abiotic stimulus	protein folding, response to heat, temperature and abiotic stimulus	response to heat, temperature, and protein folding	*No statistically significant terms*
D6	red	lipid and small molecule metabolic process	lipid and small molecule metabolic process	*No statistically significant terms*	*No statistically significant terms*
D7	black	*No statistically significant terms*	*No statistically significant terms*	*No statistically significant terms*	*No statistically significant terms*
D8	pink	photosynthesis, nitrogen, amide and peptide biosynthetic and metabolic processes	*No statistically significant terms*	*No statistically significant terms*	photosynthesis, nitrogen, amide and peptide biosynthetic and metabolic processes
D9	magenta	*No statistically significant terms*	*No statistically significant terms*	*No statistically significant terms*	*No statistically significant terms*

(b)

	Modules	All genes	core genes	soft-core genes	Shell genes
W1	turquoise	nitrogen compound, organic substance, primary metabolic processes	nitrogen compound, organic substance, primary metabolic processes	protein ubiquitination, protein modification by small protein conjugation	*No statistically significant terms*
W2	blue	peptide, amide and nitrogen biosynthetic and metabolic processes	peptide, amide and nitrogen biosynthetic and metabolic processes	peptide, amide and nitrogen biosynthetic and metabolic processes	peptide, amide and nitrogen biosynthetic and metabolic processes
W3	brown	protein, peptide, amide and ion transport/location	protein, peptide, amide and ion transport/location	protein and ion transport/location	*No statistically significant terms*
W4	yellow	photosynthesis, generation of precursor metabolites and energy, cellular nitrogen compound and organic substance biosynthetic processes, and electron transport chain	organic substance, primary, nitrogen compound, nucleobase-containing compound metabolic processes	*No statistically significant terms*	photosynthesis, generation of precursor metabolites and energy, cellular nitrogen compound and organic substance biosynthetic processes, and electron transport chain
W5	green	organonitrogen and phosphate-containing compounds, and phosphorus metabolic processes and protein phosphorylation	organonitrogen and phosphate-containing compounds, small molecule, protein and phosphorus metabolic processes and protein phosphorylation	*No statistically significant terms*	*No statistically significant terms*
W6	red	plant-type cell wall organization or biogenesis	plant-type cell wall organization or biogenesis	*No statistically significant terms*	*No statistically significant terms*
W7	black	glutathione, cellular amino acid, oxoacid, organic acid and sulfur compound metabolic processes	glutathione, cellular amino acid, oxoacid, organic acid and sulfur compound metabolic processes	*No statistically significant terms*	*No statistically significant terms*

W8	pink	small molecule, starch, proline and carbohydrate metabolic processes	small molecule, starch, proline and carbohydrate metabolic processes	*No statistically significant terms*	*No statistically significant terms*
W9	magenta	regulation of nucleic acid-templated transcription, and RNA, nucleobase-containing compound, macromolecule biosynthesis and metabolic processes	cellular protein modification process, protein dephosphorylation, protein modification process, macromolecule modification and dephosphorylation	regulation of nucleic acid-templated transcription, and RNA, nucleobase-containing compound, macromolecule biosynthesis and metabolic processes	*No statistically significant terms*
W10	purple	photosynthesis	photosynthesis	photosynthesis	*No statistically significant terms*
W11	greenyellow	*No statistically significant terms*	*No statistically significant terms*	*No statistically significant terms*	*No statistically significant terms*
W12	tan	*No statistically significant terms*	*No statistically significant terms*	*No statistically significant terms*	*No statistically significant terms*
W13	salmon	macromolecule, cellular nitrogen compound, peptide and amide biosynthetic processes	*No statistically significant terms*	*No statistically significant terms*	macromolecule, cellular nitrogen compound, peptide and amide biosynthetic processes

Table 3. DNA motifs and cis-regulatory elements (CREs) of co-expressed genes assigned to Drought (D) (**a**) and Water (W) (**b**) modules. Modules (numeric and color codes of the modules); Accession (footprintDB), Names (gene name); Consensus sequence of motif; Protein name (Swissprot); Ncor (normalized correlation score); e-value; sites (number of sites used to compile the DNA motif); Proportion of genes with CREs per module as detected by matrix-scan in -500 to +200 bp windows; Total genes with CREs [occupancy of core, soft-core and shell genes]. The dashes indicate that no results were retrieved.

(a)

Drought modules		Accession (footprintDB)	Names	consensus	Protein (Swissprot)	Ncor	evalue	sites	Proportion of genes with CREs per module	Total genes with CREs [occupancy of core, soft-core and shell genes]
D1	turquoise	MA1353.1 (JASPAR 2020), M0609 (AthalianaCistrome v4_May2016)	AT1G72740, AT1G72740. DAP, T04493	ssmaAAACCCTAGcy	Telomere repeat-binding factor 5 (MYB transcription factor)	0.741	1.40E-56	225	9.0%	178 [139 (78.1%); 26 (14.6%); 13 (7.3%)]
D2	blue	ABI4 (Athamap 20091028)	ABI4	scCACCaCCCrc	Ethylene-responsive transcription factor ABI4 (Protein ABSCISIC ACID INSENSITIVE 4)	0.697	5.80E-07	178	17.1%	128 [102 (79.7%); 17 (13.3%); 9 (7.0%)]

D3	brown	M1684_1.02 (CISBP 1.02)	T153999_1.02, WRKY60	gtCGGTCAACgk	Probable WRKY transcription factor 60 (WRKY DNA-binding protein 60)	0.877	1.30E-09	165	11.6%	73 [54 (74.0%); 13 (17.8%); 6 (8.2%)]
D4	yellow	MA1197.1 (JASPAR 2020), M0350 (AthalianaCistrome v4_May2016)	CAMTA1, CAMTA1.DAP, T27097	bgCGACGCGCTks	Calmodulin-binding transcription activator 1, AtCAMTA1 (Ethylene-induced calmodulin-binding protein b, EICBP.b) (Signal-responsive protein 2, AtSR2)	0.717	7.80E-18	69	19.4%	50 [26 (52.0%); 13 (26.0%); 11 (22.0%)]
D5	green	M0455 (AthalianaCistrome v4_May2016)	HSFB3.ampDAP, T08630	kkCTCTrGAAGgk	Heat stress transcription factor B-3, AtHsfB3 (AtHsf-05)	0.691	9.10E-13	89	27.8%	47 [36 (76.6%); 5 (10.6%); 6 (12.8%)]
D6	red	Not found	Not found	Not found	Not found	Not found	Not found	Not found	-	-
D7	black	AHL25_3 (Arabidopsis PBM 20140210)	AHL25	wcAAAATAAATak	AT-hook motif nuclear-localized protein 25 (AT-hook protein of GA feedback 1)	0.643	0.46	35	54.2%	26 [16 (61.6%); 5 (19.2%); 5 (19.2%)]

					Probable plastid-lipid-associated protein 14, chloroplastic, AtPap14 (Fibrillin-11) (OBP3-responsive protein 1)					
D8	pink	MA1274.1 (JASPAR 2020), M0256 (AthalianaCistrome v4_May2016)	OBP3, OBP3.DAP, T16682	mrArGAATArrrAARrdh	Probable plastid-lipid-associated protein 14, chloroplastic, AtPap14 (Fibrillin-11) (OBP3-responsive protein 1)	0.623	1.50E-12	14	3.5%	2
D9	magenta	Not found	Not found	Not found	Not found	Not found	Not found	Not found	-	-

(b)

Water modules		Accession (footprintDB)	Names	consensus	Protein (Swissprot)	Ncor	evalue	sites	Genes with putative CREs in promoter	
									Proportion of genes with CREs per module	Total genes with CREs [occupancy of core, soft-core and shell genes]
W1	turquoise	M0587_1.02 (CISBP 1.02)	PK22848.1, T073801_1.02	wkwcTAAAATTTTAGwww	No results	0.59	6.80E-29	32	0.1%	2
W2	blue	At2g20350:M05132:TRANSFAC	At2g20350/M05132/TRANSFAC	mmAGGCCCATCwv	Ethylene-responsive transcription factor ERF120	0.59	1.90E-97	244	22.3%	162 [115 (71.0%); 25 (15.4%); 22 (13.6%)]

						Not found	Not found	Not found	-	-
W3	brown	Not found	Not found	Not found	Not found					
W4	yellow	MA1379.1 (JASPAR 2020), M0357 (AthalianaCistrome v4_May2016)	SOL1, SOL1.DAP, T15201	wwTTAAwwwddAAAwr	Carboxypeptidase SOL1, EC 3.4.17.- (Protein SUPPRESSOR OF LLP1 1)	0.662	1.10E-09	458	0.3%	2
W5	green	UP00581A_2 (UniPROBE 20160601)	vascular plant one zinc finger protein 2, VOZ2	sgAGTCAACGtcgv	Transcription factor VOZ2 (Protein VASCULAR PLANT ONE-ZINC FINGER 2, AtVOZ2)	0.88	1.90E-10	126	5.7%	29 [20 (69.0%); 4 (13.8%); 5 (17.2%)]
W6	red	M0523 (AthalianaCistrome v4_May2016)	MYB58.DAP, T03869	ytgCyACCAACCAvm	Transcription factor MYB58	0.837	1.10E-11	45	2.5%	9
W7	black	UP00582A_2 (UniPROBE 20160601)	ARABIDOPSIS THALIANA WRKY DNA-BINDING PROTEIN 18, At4g31800, AtWRKY18, F11C18.16, F28M20.10, WRKY transcription factor 1, WRKY transcription	mmsGGTCAAACGyr	WRKY transcription factor 18	0.698	7.40E-07	27	6.1%	19 [17 (89.5%); 2 (10.5%); 0 (0.0%)]

				factor 18, WRKY1						
W8	pink	Not found	Not found	Not found	Not found	Not found	Not found	Not found	-	-
W9	magenta	MA1197.1 (JASPAR 2020), M0350 (AthalianaCistrome v4_May2016)	CAMTA1, CAMTA1.DAP, T27097	ssaCCGCGTcss	Calmodulin-binding transcription activator 1, AtCAMTA1 (Ethylene-induced calmodulin-binding protein b, EICBP.b) (Signal-responsive protein 2, AtSR2)	0.755	4.50E-19	95	23.0%	55 [28 (50.9%); 12 (21.8%); 15 (27.3%)]
W10	purple	Not found	Not found	Not found	Not found	Not found	Not found	Not found	-	-
W11	green-yellow	Not found	Not found	Not found	Not found	Not found	Not found	Not found	-	-
W12	tan	Not found	Not found	Not found	Not found	Not found	Not found	Not found	-	-
W13	salmon	Not identified	Not identified	trAAATymwwwTTCAht	Not identified	0.504	0.00017	35	20.0%	8

Table 4. Occupancies [core (33 accessions); soft-core (31-32 accessions) and shell (≤30 accessions)] of the co-expressed genes **(a; b)** and hub genes **(c; d)** for each module of the Drought (D) and Water (W) co-expression networks. Asterisks indicate significant differences comparing shell and total genes between modules and network by Fisher test (*p ≤ 0.05; **p ≤ 0.01; ***p ≤ 0.001).

(a)

Drought modules		Genes	Drought genes occupancy		
			Core (33)	Soft-core (32 or 31)	Shell (≤30)
D0	gray	8313	5764 (69.3%)	1269 (15.3%)	1280 (15.4%)
D1	turquoise	1986	1422 (71.6%)	265 (13.3%)	299 (15.1%)
D2	blue	750	549 (73.2%)	126 (16.8%)	75 (10.0%)***
D3	brown	627	440 (70.2%)	117 (18.7%)	70 (11.2%)*
D4	yellow	258	160 (62.0%)	49 (19.0%)	49 (19.0%)
D5	green	169	123 (72.8%)	27 (16.0%)	19 (11.2%)
D6	red	95	62 (65.3%)	18 (19.0%)	15 (15.8%)
D7	black	48	28 (58.3%)	9 (18.8%)	11 (22.9%)
D8	pink	57	17 (29.8%)	4 (7.0%)	36 (63.2%)***
D9	magenta	27	20 (74.1%)	6 (22.2%)	1 (3.7%)
Total counts		12330	8585 (69.6%)	1890 (15.3%)	1855 (15.0%)
Total unique counts		12137	8426 (69.4)	1869 (15.4%)	1842 (15.2%)
Total unique counts (excluding gray module)		4006	2813 (70.2%)	618 (15.4%)	575 (14.4%)

(b)

Water modules		Genes	Water genes occupancy		
			Core (33)	Soft-core (32 or 31)	Shell (≤30)
W0	gray	6934	4788 (69.1%)	1006 (14.5%)	1140 (16.4%)*
W1	turquoise	1439	1094 (76.0%)	188 (13.1%)	157 (10.9%)***
W2	blue	727	544 (74.8%)	111 (15.3%)	72 (9.9%)***
W3	brown	696	500 (71.8%)	128 (18.4%)	68 (9.8%)***
W4	yellow	590	353 (59.8%)	73 (12.4%)	164 (27.8%)***
W5	green	509	375 (73.7%)	101 (19.8%)	33 (6.5%)***
W6	red	358	258 (72.1%)	67 (18.7%)	33 (9.2%)**
W7	black	313	216 (69.0%)	59 (18.8%)	38 (12.1%)
W8	pink	245	162 (66.1%)	50 (20.4%)	33 (13.5%)
W9	magenta	239	124 (51.9%)	63 (26.4%)	52 (21.8%)*
W10	purple	185	127 (68.6%)	36 (19.5%)	22 (11.9%)
W11	greenyellow	55	33 (60.0%)	8 (14.5%)	14 (25.5%)
W12	tan	51	31 (60.8%)	6 (11.8%)	14 (27.5%)
W13	salmon	40	13 (32.5%)	3 (7.5%)	24 (60.0%)***
Total counts		12381	8618 (69.6%)	1899 (15.3%)	1864 (15.1%)
Total unique counts		12137	8426 (69.4)	1869 (15.4%)	1842 (15.2%)
Total unique counts (excluding gray module)		5407	3795 (70.2%)	890 (16.5%)	722 (13.3%)

(c)

Drought modules		Hub Genes	Drought hub genes (KME>0.9) occupancy		
			Core (33)	Soft-core (32 or 31)	Shell (≤30)
D1	turquoise	3	3 (100%)	0 (0%)	0 (0%)
D2	blue	13	10 (76.9%)	1 (7.7%)	2 (15.4%)
D3	brown	38	20 (52.6%)	18 (47.4%)	0 (0%)
D4	yellow	4	3 (75%)	1 (25%)	0 (0%)
D5	green	3	3 (100%)	0 (0%)	0 (0%)
D6	red	7	4 (57.1%)	3 (42.9%)	0 (0%)
D7	black	4	0 (0%)	0 (0%)	4 (100%)
D8	pink	0	0 (0%)	0 (0%)	0 (0%)
D9	magenta	0	0 (0%)	0 (0%)	0 (0%)
Total counts		72	43 (59.7%)	23 (31.9%)	6 (8.3%)
Total unique counts		72	43 (59.7%)	23 (31.9%)	6 (8.3%)

(d)

Water modules		Hub Genes	Water hub genes (KME>0.9) occupancy		
			Core (33)	Soft-core (32 or 31)	Shell (≤30)
W1	turquoise	115	82 (71.3%)	18 (15.7%)	15 (13.0%)
W2	blue	3	2 (66.7%)	1 (33.3%)	0 (0%)
W3	brown	1	1 (100%)	0 (0%)	0 (0%)
W4	yellow	2	1 (50%)	0 (0%)	1 (50%)
W5	green	10	6 (60%)	3 (30%)	1 (10%)
W6	red	8	7 (87.5%)	1 (12.5%)	0 (0%)
W7	black	23	16 (69.6%)	7 (30.4%)	0 (0%)***
W8	pink	2	2 (100%)	0 (0%)	0 (0%)
W9	magenta	21	13 (61.9%)	1 (4.8%)	7 (33.3%)
W10	purple	2	2 (100%)	0 (0%)	0 (0%)
W11	greenyellow	1	1 (100%)	0 (0%)	0 (0%)
W12	tan	1	0 (0%)	0 (0%)	1 (100%)
W13	salmon	1	1 (100%)	0 (0%)	0 (0%)
Total counts		190	134 (70.5%)	31 (16.3%)	25 (13.2%)

Total unique counts	190	134 (70.5%)	31 (16.3%)	25 (13.2%)

Table 5. Pan-genome analysis of the differentially expressed (DE) isoforms according to their gene occupancy: core (33 accessions), soft-core (31-32 accessions) and shell (≤30 accessions).* One gene was represented by one up-regulated and one down-regulated isoform.

	Gene occupancy			
	core	**soft-core**	**shell**	**TOTAL**
upregulated	2027 (72.2%)	500 (17.8%)	281 (10%)	**2808**
downregulated	661 (67.5%)	205 (20.9%)	114 (11.6%)	**980**
up/down-regulated	1*	0	0	**1**
TOTAL	**2689**	**705**	**395**	**3789**

FIGURES

Figure 1. Summary of the experimental design and analyses performed in the 33 accessions of the model grass *Brachypodium distachyon* under drought (D) and water (control) (W) conditions.

Figure 2 Correspondence (number of nodes) of Drought (D) (**a**) and Water (W) (**b**) set-specific and Drought-Water consensus (Cons) modules. Each row of the table corresponds to one Drought/Water set-specific module, and each column corresponds to one consensus module. Numbers in the table indicate node counts in the intersection of the corresponding modules. Coloring of the table encodes − log(p), with p being the Fisher's exact test p-value for the overlap of the two modules. The stronger the red color, the more significant the overlap is. (**c**) Comparison of the GO enrichments for each D and W module, according to the correlation with the common consensus module.

Figure 3. Proportion (%) of occupancies [core (33); soft-core (31-32) and shell (≤30)] of the co-expressed genes (**a; b**) and hub genes (**c; d**) for each Drought (D) and Water (W) network.