Intra-Articular Delivery of an Indoleamine 2,3-Dioxygenase Galectin-3 Fusion Protein for Osteoarthritis Treatment in Male Lewis Rats

Brittany D. Partain Ph.D.[1], Evelyn Bracho-Sanchez Ph.D.[1], Shaheen A. Farhadi Ph.D.[1], Elena G. Yarmola Ph.D.[1], Benjamin G. Keselowsky Ph.D.[1], Gregory A. Hudalla Ph.D.[1], and Kyle D. Allen Ph.D.[1,2]

[1] J. Crayton Pruitt Family Department of Biomedical Engineering, University of Florida, Gainesville, FL, USA
[2] Department of Orthopedics and Sports Medicine, University of Florida, Gainesville, FL, USA

Corresponding Author:
Kyle D. Allen, Ph.D.
Associate Professor
J. Crayton Pruitt Family Department of Biomedical Engineering
University of Florida
1275 Center Drive
Biomedical Sciences Building
Gainesville, Florida 32611
T: (352) 273-9337
F: (352) 273 9221
kyle.allen@bme.ufl.edu

Key Words: Knee Osteoarthritis, Indoleamine 2,3-Dioxygenase, Galectin-3, Metabolic Reprogramming, Intra-articular Drug Delivery, Inflammation, Pain and Disability

Running Head: IDO-Gal3 for OA Treatment

Grants and other financial supporters of the study: This work was supported by the NIH / National Institute of Arthritis and Musculoskeletal and Skin Diseases (Grant R01AR068424, R01AR071431) and NIH/National Institute of Dental and Craniofacial Research (Grant R01DE027301). KDA is part of a patent on the magnetic capture technology (US Patent No. 10,695,036). BGK and GAH are founders, hold stock, and serve on the scientific advisory board of Anchor Biologics, Inc. The University of Florida has pending patents related to molecules and their uses reported in this paper.

Abstract (250 words)

Objective: Controlling joint inflammation can improve osteoarthritis (OA) symptoms; however,

current treatments often fail to provide long-term effects. We have developed an indoleamine 2,3-

dioxygenase and galectin-3 fusion protein (IDO-Gal3). IDO converts tryptophan to kynurenines,

directing the local environment toward an anti-inflammatory state; Gal3 binds carbohydrates and

extends IDO's joint residence time. In this study, we evaluated IDO-Gal3's ability to alter OA-

associated inflammation and pain-related behaviors in a rat model of established knee OA.

Methods: Joint residence was first evaluated with an analog Gal3 fusion protein (NanoLuc™

and Gal3, NL-Gal3) that produces luminescence from furimazine. OA was induced in male Lewis

rats via a medial collateral ligament and medial meniscus transection (MCLT+MMT). At 8 weeks,

NL or NL-Gal3 were injected intra-articularly (n=8 per group), and bioluminescence was tracked

for 4 weeks. Next, IDO-Gal3's ability to modulate OA pain and inflammation was assessed. Again,

OA was induced via MCLT+MMT in male Lewis rats, with IDO-Gal3 or saline injected into OA-

affected knees at 8 weeks post-surgery (n=7 per group). Gait and tactile sensitivity were then

assessed weekly. At 12 weeks, intra-articular levels of IL6, CCL2, and CTXII were assessed.

Results: The Gal3 fusion increased joint residence in OA and contralateral knees ($p<0.0001$).

In OA-affected animals, IDO-Gal3 improved tactile sensitivity ($p=0.002$), increased walking

velocities ($p≤0.033$), and improved vertical ground reaction forces ($p≤0.04$). Finally, IDO-Gal3

decreased intra-articular IL6 levels within the OA-affected joint ($p=0.0025$).

Conclusion: Intra-articular IDO-Gal3 delivery provided long-term modulation of joint

inflammation and pain-related behaviors in rats with established OA.

Introduction

Osteoarthritis (OA) is associated with chronic local, low grade inflammation (1). Controlling joint inflammation is important, especially if control of inflammation leads to improvement of OA symptoms. Current treatments for OA include oral non-steroidal anti-inflammatory drugs (NSAIDs) and intra-articular injections of corticosteroids or hyaluronic acid (2). Although intra-articular injections increase the local concentration of an anti-inflammatory drug and decrease off target effects, the efficacy of this approach is often hindered by rapid joint clearance (3). Short joint residence times typically necessitate repeated injections, which can lead to increased risk of infection and chondrocyte death (4–6).

Multiple intra-articular delivery strategies aim to extend joint residence times (7). For example, therapeutics are commonly encapsulated in carriers that extend joint residence; however, harsh encapsulation conditions can reduce a drug's bioactivity (8,9). Alternatively, conjugating protein-based drugs to synthetic particles can extend joint residence, improve bioactivity, and limit degradation (10); however, determining an effective dose to circumvent joint clearance and maintain efficacy has proven difficult (11). In this regard, enzyme-based therapeutics may offer an advantage (12). With the catalytic capabilities of enzymes, target molecules can be continually and locally produced, as long as the enzyme's substrate is available. This decreases the necessary dose of the enzyme, and conjugating enzymes to anchoring molecules can further enhance bioactivity and extend joint residence.

To provide long-lasting local control of inflammation, we developed an indoleamine 2,3-dioxygenase galectin-3 fusion protein (IDO-Gal3). IDO is an immunomodulatory enzyme that converts tryptophan into kynurenines (13–16). By locally degrading tryptophan, IDO induces effector T cell apoptosis, promotes regulatory T cell proliferation, modulates macrophage differentiation toward an anti-inflammatory phenotype, and maintains an immature phenotype on dendritic cells that results in suppression of antigen-specific T cell proliferation (15,17–19). Gal3

binds to β-galactoside glycoconjugates and glycosaminoglycans, and these sites are prevalent in

multiple joint tissues. Via this binding, Gal3 can extend an enzyme's local residence time (20–

22). In this work, IDO-Gal3's ability to alter OA-associated inflammation and pain-related

behaviors was evaluated in a rat model of established knee OA.

Methods

Experimental Designs

Animal procedures were approved by the University of Florida's Institutional Animal Care and

Use Committee. In our first experiment, joint retention of a Gal3 fusion protein was assessed

(Supplemental Fig. 1). Here, 16 male Lewis rats (250 g) were acquired from Charles River

Laboratories (Wilmington, MA, USA) and acclimated to the University of Florida for 1 week. Rats

then underwent a medial collateral ligament and medial meniscus transection surgery

(MCLT+MMT). At 8 weeks post-surgery, background luminescence was measured for each rat

(1 sec exposure, field of view D, PerkinElmer IVIS, Waltham, MA, USA). Both knees were then

injected with either 50 μL NanoLuc™ Luciferase (NL, n=8, 3.27 μM) or a NanoLuc™ Luciferase

galectin-3 fusion protein (NL-Gal3, n=8, 3.27 μM). For luminescence imaging, knees were injected

with 50 μL furimazine (1:50 dilution in PBS, Nano-Glo™, PRN1120, Promega, Madison, WI,

USA), then assessed with the above IVIS settings. Luminescence imaging was repeated 1, 2, 4,

8, 12, 16, 20, 24, and 28 days after NL or NL-Gal 3 injection. On day 28, rats were euthanized via

exsanguination under deep anesthesia, and the patella, tibia, femur, and meniscus were

dissected, preserving the synovium and fat pad connected to each bone. Tissue samples were

placed in a 24 well plate and incubated with diluted furimazine for 1 min at 25°C. After incubation,

tissues were removed, and luminescence was measured using the autoexposure setting in field

of view C.

In our second experiment, we evaluated IDO's ability to modulate OA-related inflammation and

symptoms (Supplemental Fig. 2). Here, 16 additional male Lewis rats (250 g, 3 months) were

acquired from Charles River Laboratories and acclimated to the University of Florida for 1 week.

Rats then underwent baseline behavioral testing followed by MCLT+MMT surgery; one rat was

euthanized due to surgical complications. Post-surgery, rodent gait was assessed at 3, 5, and 7

weeks and tactile sensitivity was assessed weekly (except for week 6 due to university closure).

At 8 weeks post-surgery, rats received 30 µL injections of either saline (n=7) or IDO-Gal3 (n=8,

33 pg/µL) in the MCLT+MMT knee. Tactile sensitivity testing was conducted the day after

injection, then weekly out to 11 weeks post-surgery. Gait testing was performed 2 days after

injection, then weekly out to 11 weeks post-surgery. Rats were then euthanized via

exsanguination under deep anesthesia. Immediately post-mortem, IL6, CCL2, and CTXII were

assessed in both knees using magnetic capture. Here, saline samples from a contralateral and

operated limb were lost due to insufficient collection of magnetic beads. Following magnetic

capture, knees were processed for histology. In histology, one IDO-Gal3 treated rat showed an

intact medial meniscus (failure to fully transect the meniscus); this animal was removed from the

analysis. Thus, the final datasets are n=6-7 for saline treatment and n=7 for IDO-Gal3 treatment.

Statistical analyses are described for each method below, with analyses performed at α=0.05 in

R Studio.

Detailed Methods

Protein Production

Genes encoding NL-Gal3 (IDT, Coralville, IA, USA) and recombinant human Gal3 (OriGene,

Rockville, MD, USA) were inserted into pET-21d(+) vectors between the Ncol and Xhol sites, then

transformed into One Shot™ TOP10 Chemically Competent E. coli (ThermoFisher, Waltham, MA,

USA). NL was amplified from the NL-Gal3 gene using PCR and mutated to have Ncol and Xhol

flank the genes using the following primers: 5'-CGC CTC GAG CGC CAG AAT GCG TT-3' and

5'-GCT TAG CCA TGG CGG TCT TCA CAC TCG AAG-3'. PCR products were digested using

NcoI and XhoI and re-inserted into pET-21d(+) vectors. Genes encoding IDO were generously

provided by Dr. Carlos Gonzalez's Laboratory at the University of Florida. Amplification and

insertion of the BamHI restriction site was performed using the following primers: 5'-CAG CTA

CCA TGG CAC ACG CTA TGG AAA-3' and GAG AAC 5'-GGA TCC ACC TTC CTT CAA AAG-

3'. The NL-Gal3 gene was digested with NcoI and BamHI to remove the NL gene and insert the

IDO gene. After overnight treatment on ampicillin (100 μg/mL) doped LB/agar plates (37°C),

positive clones were used to inoculate 5 mL of LB broth containing ampicillin (100 μg/mL).

Cultures were grown overnight on an orbital shaker (37°C, 220 rpm), with plasmids isolated using

a Plasmid Miniprep Kit (Qiagen, Hilden, Germany).

Origami™ B (DE3) E. coli (Novagen, Madison, WI, USA) were transformed with pET-21d-Gal3,

pET-21d-NL, pET-21d-NL-Gal3, or pET-21d-IDO-Gal3 vectors and selected on ampicillin (100

μg/mL) and kanamycin A (50 μg/mL) doped LB/agar plates overnight (37°C). Positive clones were

used to inoculate 5 mL of LB broth containing ampicillin (100 μg/mL) and kanamycin A (50 μg/mL)

and grown overnight (37°C, 220 rpm). Cultures were expanded in 1 L 2xTY broth containing

ampicillin (100 μg/mL) and kanamycin A (50 μg/mL) and grown on an orbital shaker (37°C, 225

rpm) until the optical density reached 0.6-0.8 (λ=600). Protein expression was induced via 0.5

mM isopropyl β-D-1-thiogalactopyranoside (ThermoFisher, Waltham, MA, USA) for 18 hrs (18°C,

225 rpm). Bacteria were pelleted via centrifugation, washed with PBS, and incubated for 20 mins

with B-PER™ bacterial protein extraction reagent (ThermoFisher, Waltham, MA, USA), 1 Pierce

protease inhibitor tablet (ThermoFisher, Waltham, MA, USA), 300 units DNAse I from bovine

pancreas (MilliporeSigma, Burlington, MA, USA), and 100 μg lysozyme (MilliporeSigma,

Burlington, MA, USA). Lysed bacteria were centrifuged, and supernatant was loaded into columns

containing HisPur™ Cobalt Superflow Agarose (ThermoFisher, Waltham, MA, USA). Proteins

were eluted using an imidazole gradient (0-250 mM), then centrifuged on Amicon Ultra Centrifugal

Filters with a 10 kDa cutoff (MilliporeSigma, Burlington, MA, USA). A second purification step was

performed for IDO-Gal3 using size exclusion chromatography in an ÄKTA Pure Protein

Purification System (GE Life Sciences, Marlborough, MA, USA). Molecular weight and purity were

determined by sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE).

Endotoxin contaminants were removed via Detoxi-Gel Endotoxin Removing Columns

(ThermoFisher, Waltham, MA, USA). Final endotoxin content was determined to be below 0.1

EU/mL using Pierce™ LAL Chromogenic Endotoxin Quantitation Kit (ThermoFisher, Waltham,

MA, USA).

Rat OA Model

Rats were anesthetized using 2.5% isoflurane (Patterson Veterinary, Greeley, CO, USA). Right

knees were aseptically prepared with betadine surgical scrub (Purdue Products, Stamford, CT,

USA) and 70% ethanol in triplicate, ending with a fourth betadine scrub. Via a 1-2 cm skin incision

and blunt muscle dissection, the medial collateral ligament was exposed and then transected.

Knee abduction was then used to expose the medial meniscus, with the meniscus transected

radially in its central portion. Joint capsule and muscle were closed with absorbable 5-0 vicryl

braided sutures (Ethicon, Somerville, NJ, USA). Skin was closed with 4-0 ethilon nylon

monofilament (Ethicon, Somerville, NJ, USA). For post-surgical pain, rats received subcutaneous

buprenorphine (0.05 mg/kg, Patterson Veterinary, Greeley, CO, USA) peri-operatively then every

12 hrs for 48 hrs.

Intra-articular Injections

Rats were anesthetized and aseptically prepared as described above. A 1 mL 27G x 3/8

syringe (Becton Dickinson, Franklin Lakes, NJ, USA) was inserted through the patellar ligament

into the joint space. After injection, the needle was removed, gauze wetted with 70% ethanol was

pressed against the injection site, and the knee was flexed.

IVIS Image Analysis

A region of interest (ROI) was drawn around the largest signal then copied to all other images. After subtracting background radiance, average radiance was calculated in the ROI. Average radiance for each animal was $\log_{10}$ normalized, plotted against time, then area under the curve (AUC) was calculated. Using AUCs, unpaired Student's *t*-tests were conducted to evaluate clearance differences between NL and NL-Gal3 treated MCLT+MMT knees or NL and NL-Gal3 treated contralateral knees.

Gait Analysis

Rodent gait was analyzed using our GAITOR system (23). Briefly, high-speed videography (IDT M3, Pasadena, CA, 500 fps) and Kistler 3-axis load cells (±2 kN, Sindelfingen, Germany) were used to collect gait data during walking. Videos were analyzed to determine stride length, percentage stance time, temporal symmetry, and spatial symmetry (24). Force recordings were analyzed for peak vertical force at weeks 7-11 (25).

After IDO-Gal3 or saline treatment, gait differences were assessed with linear mixed effects models treating the animal identifier as a random factor. If indicated, comparison of least square group means was conducted using Tukey's HSD corrections for multiple comparisons. For stride length, percentage stance time, and peak vertical force, data were first visualized relative to velocity, then statistically analyzed as both means and velocity-corrected residuals. For residualization, pre-treatment data (week 7) was used as the control line (24). For stance time imbalance and gait symmetries, data were also compared to mathematical definitions for balanced and symmetric gait using Bonferroni-correct Student's *t*-tests.

Tactile Sensitivity

50% paw withdrawal thresholds were determined using Chaplan's up-down method (26). Here, von Frey filaments (0.6, 1.4, 2.0, 4.0, 6.0, 8.0, 15.0, and 26.0 g, Stoelting Co., Wood Dale, IL,

USA) were applied to the hind foot's plantar region, starting with the 4.0 g filament. If paw

withdrawal occurred, a less stiff filament was applied; if paw withdrawal did not occur, a stiffer

filament was applied. Using these data, the force where paw withdrawal is equally likely to

stimulus tolerance can be approximated (26). Tactile sensitivity was evaluated using a linear

mixed effects model treating week and group as fixed effects and animal ID as a random effect.

When indicated, comparisons of least squared means were conducted, correcting for

compounding type 1 errors using Tukey's HSD adjustment.

Magnetic Capture

Magnetic capture was used to assess intra-articular levels of IL6, CCL2, and CTXII (27). In

magnetic capture, particles containing superparamagnetic iron oxide nanoparticles are used to

magnetically precipitate target molecules from synovial fluid (28). In this work, biotin-streptavidin

binding was used to functionalize magnetic particles (Dynabeads MyOne™ Streptavidin C1, Cat.

#65001, Life Technologies, Carlsbad, CA, USA) with antibodies for CTXII (cat. #AC-08F1,

ImmunoDiagnostic Systems, Copenhagen, Denmark), IL6, and CCL2 (cat. #517703 and 505908,

Biolegend, San Diego, CA, USA). Antibody-functionalized particles were injected into rat knees,

incubated for 2 hrs, then washed from the joint using five 50 µL PBS washes. Particles in wash

solutions were magnetically separated then washed again. CTXII, IL6, and CCL2 were released

from particles via a 15 mins treatment of 100 mM Glycine-Tris buffer (pH 3.1, containing 2% BSA

and 2 mM EDTA). Following release, particles were magnetically separated, and supernatant was

adjusted to pH 8.3.

In supernatants, CTXII was evaluated using the Cartilaps CTXII ELISA (cat. #AC-08F1,

ImmunoDiagnostic Systems Cartilaps kit, Copenhagen, Denmark) according to manufacturer's

instructions. CCL2 was quantified using a rat CCL2 ELISA (Cat. # KRC1012 Life Technologies,

Carlsbad, CA, USA) according to manufacturer's instructions and buffer modifications described

in Yarmola et al (29). IL6 was quantified using an ELISA developed from an antibody pair (biotin

anti-rat IL6 detection antibody, cat. #517703, and purified anti-rat IL6 coating antibody, cat.

#517701, Biolegend, San Diego, CA, USA). For this ELISA, coating antibodies were diluted in

buffer containing 100 mM $NaHCO_3$ and 34 mM Na_2CO_3 (pH 9.5) and incubated on Nunc

MaxiSorp™ microwells (Cat. #434797, ThermoFisher Scientific Inc., Waltham, MA, USA) for 30

mins on a plate shaker then overnight (4°C). Plates were washed 5 times (PBS with 0.5% Tween-

20), blocked for 1 hr in PBS containing 2% BSA and 10% heat-treated bovine serum (blocking

solution), then washed again. Samples and standards were pre-incubated with detection

antibodies for 30 mins at room temperature then overnight at 4°C, then added to the plate,

incubated for 3 hrs, and washed. Avidin-HRP (100 µl, cat. #405103, Biolegend, San Diego, CA,

USA) was diluted 500 times in blocking solution, incubated in the wells for 30 mins, and then

washed. Finally, tetramethylbenzidine substrate (100 µL) was added and incubated for 15 mins,

with the reaction stopped using diluted sulfuric acid (100 µL). Absorbance was then read at 450

and 650 nm.

To quantify particles, particles were re-suspended in 60 µL of PBS, sonicated (Model M1800H,

Branson Ultrasonic Corporation, Danbury, CT, USA), and read for absorbance at 450 nm. Known

particle concentrations were used as standards.

Following ELISA and particle quantification, biomarker levels were estimated as described in

(28) with assay validation provided in (27,28). IL6, CCL2, and CTXII levels in rat knees were

compared with one-way ANOVAs with *post hoc* Tukey's HSD tests when indicated.

Histology

Knees were fixed in 10% neutral buffered formalin (Fisher Scientific, Pittsburgh, PA, USA) for

48 hrs, decalcified in Cal-Ex (Fisher Scientific, Pittsburgh, PA, USA) for 5 days, dehydrated using

an ethanol ladder, and infiltrated with paraffin wax. Then, 10 µm frontal sections were acquired

with at least one section taken at every 100 µm between the anterior to posterior meniscal horns.

Slides were stained with toluidine blue; 3 sections per knee were assessed by 3 blinded graders

using our histological grading software (GEKO) (30,31). Histological differences between IDO-

Gal3 treated and saline treated knees were assessed using Student's *t*-tests.

<u>Results</u>

Galectin-3 (Gal3) increased local retention of NanoLuc™ Luciferase (NL, $p<0.0001$, Fig. 1A).

At 4 weeks post-injection, NL-Gal3 was present on the femur, patella/fat pad, meniscus, and tibia

of both knees, while NL alone was not present (Fig. 1B).

The paw of MCLT+MMT-operated limbs became sensitized to touch after surgery, indicated

by a decreasing 50% paw withdrawal threshold ($p=0.002$, Fig. 2A). After treatment with either

saline or IDO-Gal3, tactile sensitivity improved relative to pre-treatment levels ($p=0.0008$),

reaching the maximum of our testing range in 6 of 7 IDO-Gal3-treated animals. While the average

tactile sensitivity for IDO-Gal3 and saline treated animals did not vary, increases in tactile

sensitivity thresholds relative to baseline values were larger in IDO-Gal3 treated animals relative

to saline treated animals at weeks 9-11 ($p≤0.041$, Fig. 2B).

Prior to treatment, saline and IDO-Gal3 treated rats used similar walking velocities ($p≥0.89$,

Fig. 3A). Following treatment, IDO-Gal3 treated rats achieved higher velocities than saline

controls on week 10 and week 11 ($p<0.001$, $p=0.033$, respectively). Before treatment, saline and

IDO-Gal3 treated rats also used balanced, symmetric gaits. After treatment, the gait of saline

treated rats became temporally asymmetric (asynchronous foot-strike sequence) at weeks 8-11,

differing from IDO treatment animals at weeks 8-10 ($p≤0.035$, Fig. 3B). While neither saline nor

IDO-Gal3 treated animals had imbalanced stance times, saline treated animals tended to spend

more time on the contralateral limb (imbalance > 0), while IDO-Gal3 treated animals tended to

spend more time on the affected limb (imbalance < 0). At weeks 8 and 9, stance time imbalance

differed between saline and IDO-Gal3 treated animals ($p≤0.025$, Fig. 3C). Spatially asymmetric

gait patterns were not observed in either group (Fig. 3D).

Many gait parameters correlate to velocity (left column, Fig. 4). As such, velocity-dependent

gait parameters were analyzed both as averages and velocity-corrected residuals, allowing for

the evaluation of shifts due to and independent of velocity. In conjunction with velocity changes,

differences between IDO-Gal3 and saline treated rats were found for average left (contralateral)

limb percent stance time (p=0.04, treatment main effect), right (operated) limb percent stance

time (p=0.005, week-treatment interaction), stride length (p=0.02, week-treatment interaction),

and right (operated) limb peak vertical force (p=0.03, treatment main effect). Specifically,

differences between IDO-Gal3 and saline treated rats were found for average left limb percent

stance time at weeks 9, 10, and 11 (p<0.05), average stride length at weeks 9, 10, and 11

(p<0.05), and average right limb peak vertical force at weeks 9, 10, and 11 (p<0.04). Despite a

significant interaction, differences at specific weeks were not detected for average right limb

percent stance time. Peak vertical forces were similar in the left and right hind limbs of IDO-Gal3

treated rats, indicating balanced weight bearing. Saline treated rats reduced loads on the

operated limb relative to their contralateral limb at weeks 9 and 11 (Supplemental Fig. 3). Again,

to correct for velocity effects, residuals were calculated using week 7 data as the control line.

Following velocity corrections, shifts in ground reaction forces were only significant in saline

treated animals at week 9.

At 4 weeks after intra-articular injections, interleukin-6 (IL6) levels in saline treated knees were

elevated relative to contralateral controls (p=0.0025); moreover, IDO-Gal3 treated knees had

lower IL6 levels relative to saline treated knees (p=0.0021, Fig. 5A). A similar trend occurred for

intra-articular levels of CCL2 and CTXII, but neither CCL2 nor CTXII were statistically elevated in

saline treated knees compared to contralateral controls (p=0.073, p=0.090, respectively). As

such, shifts due to IDO-Gal3 treatment were also not statistically significant (Fig. 5B and 5C).

Please note, at the time of injection (8 weeks after MCLT+MMT surgery), full thickness cartilage

lesions are expected (24). As such, all MCLT+MMT operated knees had marked cartilage

damage. Treatment with IDO-Gal3 did not markedly alter cartilage histological measures,

osteophyte size, or medial capsule thickness. However, IDO-Gal3 treated knees had lower

epiphyseal trabecular bone area relative to saline controls (p=0.031, Fig. 6).

Discussion

Inflammation is recognized as a key mediator of both joint destruction and OA pain (32,33).

Thus, developing strategies for long-term control of inflammation remains a promising path for

OA drug development. Many preclinical OA studies deliver therapeutics prior to or shortly after

joint injury (34–37), when joint remodeling and pain-related behaviors are minimal. However, pain

is the primary reason OA patients seek treatment (38). To this end, we developed a novel

enzyme-based strategy for sustained control of joint inflammation and evaluated this therapeutic

after behaviors indicative of OA pain had developed.

Our first goal was to demonstrate sustained intra-articular residence time of an enzyme due to

Gal3 conjugation. Here, joint residence of nanoluciferase (a reporter enzyme) was extended by

Gal3 conjugation relative to no conjugation. Gal3 has an affinity for β-galactosides found on

extracellular matrix proteins, cell surface receptors, and lubricin (39–41). Gal3 also binds sulfated

glycosaminoglycans, like chondroitin sulfate, which is abundantly found in healthy cartilage (21).

However, in our data, operated and contralateral joints had similar nanoluciferase retention,

suggesting Gal3 is not primarily binding to chondroitin sulfate in cartilage (which is lost as OA

develops). Nonetheless, Gal3 conjugation extended joint residence in healthy and OA-affected

joints, showing luminescence signal across multiple tissues at 4 weeks after injection in both

healthy and OA-affected joints.

Use of Gal3 to extend joint residence requires careful consideration. Unconjugated Gal3 can

have pro-inflammatory actions; however, these actions require that Gal3 can cluster cell surface

receptors via an oligomerization that initiates cell adhesion, differentiation, or apoptotic signals

(42). As shown in our prior work, conjugation of Gal3 to another protein prevents this action and

limits these pro-inflammatory effects (22). Moreover, screens of synoviocyte viability in

preparation for this study confirmed that IDO-Gal3 was not cytotoxic for the doses used in this

study (Supplemental Fig. 4).

Our second goal was to evaluate IDO's ability to modulate OA-associated inflammation. Here,

IDO-Gal3 was injected into knees with established OA. At 4 weeks post-injection, IDO-Gal3

significantly decreased intra-articular IL6 levels. IDO-Gal3 tended to have similar effects on CCL2

and CTXII; however, in saline treated animals, neither CCL2 nor CTXII were significantly elevated

in MCLT+MMT operated knees relative to contralateral controls. Thus, any effects related to IDO-

Gal3 delivery were not detectable in our statistical analyses. While we have previously detected

elevated CCL2 and CTXII levels in rodent OA models (27,29), this was in untreated animals and

saline injection may have reduced effect sizes to some degree. Nonetheless, IDO-Gal3 did affect

IL6 levels at 4 weeks after injection, demonstrating potential to provide sustained effects on OA-

associated inflammation.

Our third goal was to evaluate IDO's ability to modulate OA-related symptoms. Importantly,

compensatory gaits can develop due to both pain avoidance and mechanical joint dysfunction,

and in rodent OA models, these compensations can be unilateral (limping) or bilateral (shuffle-

stepping), even for unilateral injury (24). Because both unilateral and bilateral compensations

reduce loading on the injured limb (43), peak vertical force often provides the most sensitive gait

variable for the evaluation of OA effects in rodents. Here, IDO-Gal3 treated rats had increased

loading on the operated limb relative to saline treated animals at weeks 9, 10, and 11. Shifts in

dynamic limb weight bearing were confirmed by spatial and temporal shifts in the gait cycle. For

example, IDO-Gal3 treated rats also used symmetric and balanced gaits; saline treated animals

were asymmetric and imbalanced at weeks 8 and 9 and asymmetric at week 10. IDO-Gal3 treated

rats used higher walking velocities than saline treated animals at weeks 10 and 11. Finally, IDO-

Gal3 treated animals used longer stride lengths than saline treated animals at weeks 9, 10, and

11. Combined, these data demonstrate improvement in limb use (as detected via gait parameters)

in IDO-Gal3 treated animals relative to saline controls.

Gait parameters that are correlated to walking velocity were first shown as raw values, then

analyzed as both average values and differences that were residualized to pre-treatment data.

After accounting for velocity differences, the gait profiles of IDO-Gal3 and saline treated animals

were no longer statistically significant. This indicates IDO-Gal3 treated animals did not use

fundamentally different gaits, but instead, IDO-Gal3 treated animals achieved higher functional

limb use (higher limb forces, longer stride lengths, improved stance times) via the use of faster

walking velocities. Combined with the use of symmetric and balanced gaits, these data

demonstrate a strong improvement in joint use for rats injected with IDO-Gal3.

Testing of tactile sensitivity evaluates withdrawal reflexes related to limb hypersensitivity. Here,

IDO-Gal3 treatment substantially increased paw withdrawal thresholds relative to pre-treatment

levels, with all but one IDO-Gal3 treated rat markedly increasing their paw withdrawal threshold

after injection. Withdrawal thresholds also tended to increase with saline treatment, but to a lesser

extent.

Histologically, MCLT+MMT surgery causes significant cartilage damage by 8 weeks post-

surgery. Due to cartilage's limited repair capacity, treatment at 8 weeks post-surgery was not

anticipated to improve cartilage parameters. However, IDO-Gal3 treatment did alter IL6 levels,

reduce tactile sensitivity, and improve gait parameters. Moreover, subchondral bone grades

followed these changes in inflammation and behavior, where epiphyseal trabecular bone area

indicated less area dedicated to trabecular bone and more area to marrow voids, trending toward

percentages seen in healthy knees (approximately 60%, (31)). As such, it is plausible that

treatment with IDO-Gal3 shortly after joint trauma could arrest OA progression; however, further

studies are needed to address this question.

Several metabolic pathways are altered in OA joints, including tryptophan metabolism (44).

IDO modulates the immune system primarily through local depletion of L-tryptophan and

production of kynurenines (45). Recent work suggests IDO-driven shifts in local metabolites can

cause immunomodulatory effects on T cells and antigen presenting cells (46). Consistent with

these findings, the transfection of IDO plasmids into PMA-differentiated THP-1 (dTHP-1) cells

results in increased expression of anti-inflammatory markers and decreased expression of pro-

inflammatory markers (17). Due to inflammation, IDO is often upregulated in antigen presenting

cells (45), and these antigen presenting cells can then activate regulatory T cells, inhibit effector

T cell responses, and attenuate pro-inflammatory responses (47). Taken together, these qualities

support IDO's ability to restrain pro-inflammatory cascades. However, specific immunomodulatory

effects of IDO in OA remain unanswered. Future work will also need to assess shifts in joint-level

metabolic pathways and immune cell phenotypes after intra-articular IDO-Gal3 injection.

IDO has been explored in other forms of arthritis, primarily rheumatoid arthritis. Here, studies

have focused on assessment of downstream metabolites, rather than behavioral assays. For

example, IDO inhibition in collagen-induced arthritis (CIA) increased disease severity and

elevated Th1 and Th17 responses in the joint (14,48). Related to this, intra-articular injection of

dendritic cells modified to overexpress IDO reduced joint damage in the CIA model (49). However,

inability to sustain IDO's effects has limited progress (50). Our approach, directly injecting IDO

conjugated to Gal3, may overcome these issues.

The limitations of the study include the loss of magnetic nanoparticles in a subset of our

animals, lowering statistical power of our intra-articular inflammation analysis. Additionally,

because intra-articular inflammation was prioritized in this study, the direct assessment of

metabolites in synovial fluid was not possible. Furthermore, therapeutic efficacy was not

evaluated as a function of dose or time after administration. We intend to evaluate these

parameters in follow-up studies. Nonetheless, these data demonstrate proof-of-principle that

intra-articular delivery of IDO-Gal3 affects the inflammatory environment in the OA joint for

sustained periods of time and these effects correspond to changes in OA-related behaviors in
animals with established knee OA.

Conclusions

In this study, a novel enzyme-based therapeutic was tested in a rat model of established knee
OA. First, we demonstrated that Gal3 conjugation can extend the joint residence of a reporter
enzyme out to 4 weeks. Moreover, intra-articular injection of IDO-Gal3 in the OA-affected limb
produced sustained effects on rodent walking, improved tactile sensitivity, and decreased IL6
levels. These data demonstrate the potential of IDO-Gal3 to serve as a metabolic reprogramming
strategy for long-term, control of intra-articular inflammation and OA-related pain. Future studies
will evaluate how intra-articular delivery of IDO-Gal3 modifies joint metabolism, immune cell
polarization, and progression of joint damage in additional rat OA models.

Author Contributions

BDP conducted and analyzed *in vitro* viability experiments, clearance and biodistribution
experiments, tactile sensitivity experiments, gait analysis, and histological experiments. EBS
expressed and purified IDO-Gal3. SAF expressed and purified NL-Gal3. EGY performed
magnetic capture experiments and analyzed data with BDP. KDA performed MMT surgeries.
BDP, EBS, BGK, GAH, and KDA contributed to the study design and data interpretation. The
manuscript was prepared by BDP, with critical feedback from all authors. All authors agree with
the content and interpretation of the data as presented.

Acknowledgements

The authors would like to thank Natalie Thurlow and Mallory Box for assisting with gait
processing. The authors would also like to acknowledge Nicole Sieling for sectioning and staining
samples for histology, and Kiara Chan, Jacob Griffin, and Jessica Aldrich for grading histological

images. We would also like to thank Sabrina Freeman for providing scientific and technical

support regarding IDO.

References

1. Neogi T, Guermazi A, Roemer F, Nevitt MC, Scholz J, Arendt-Nielsen L, et al. Association of Joint Inflammation With Pain Sensitization in Knee Osteoarthritis: The Multicenter Osteoarthritis Study HHS Public Access. *Arthritis Rheumatol* 2016;68:654–661.

2. Zhang W, Moskowitz RW, Nuki G, Abramson S, Altman RD, Arden N, et al. OARSI recommendations for the management of hip and knee osteoarthritis, Part II: OARSI evidence-based, expert consensus guidelines. *Osteoarthritis and cartilage* 2008;16:137–162.

3. Evans CH, Kraus VB, Setton LA. Progress in intra-articular therapy. *Nature reviews Rheumatology* 2014;10:11–22.

4. Holland C, Jaeger L, Smentkowski U, Weber B, Otto C. Septic and Aseptic Complications of Corticosteroid Injections An Assessment of 278 Cases Reviewed by Expert Commissions and Mediation Boards From. 2012:425–430.

5. Dragoo JL, Danial CM, Braun HJ, Pouliot MA, Kim HJ. The chondrotoxicity of single-dose corticosteroids. *Knee Surgery, Sports Traumatology, Arthroscopy* 2012;20:1809–1814.

6. Huang Y, Cai G quan, Peng JP, Shen C. Glucocorticoids induce apoptosis and matrix metalloproteinase-13 expression in chondrocytes through the NOX4/ROS/p38 MAPK pathway. *Journal of Steroid Biochemistry and Molecular Biology* 2018;181:52–62.

7. Maudens P, Jordan O, Allémann E. Recent advances in intra-articular drug delivery systems for osteoarthritis therapy. *Drug Discovery Today* 2018;23:1761–1775.

8. Giteau A, Venier-Julienne MC, Aubert-Pouëssel A, Benoit JP. How to achieve sustained and complete protein release from PLGA-based microparticles? *International Journal of Pharmaceutics* 2008;350:14–26.

9. Whitaker MJ, Hao J, Davies OR, Serhatkulu G, Stolnik-Trenkic S, Howdle SM, et al. The production of protein-loaded microparticles by supercritical fluid enhanced mixing and spraying. In: *Journal of Controlled Release.*Vol 101. Elsevier; 2005:85–92.

10. Whitmire RE, Wilson DS, Singh A, Levenston ME, Murthy N, García AJ. Self-assembling nanoparticles for intra-articular delivery of anti-inflammatory proteins. *Biomaterials* 2012;33:7665–75.

11. Edwards SHR. Intra-articular drug delivery: The challenge to extend drug residence time within the joint. *The Veterinary Journal* 2011;190:15–21.

12. Dean SN, Turner KB, Medintz IL, Walper SA. Targeting and delivery of therapeutic enzymes. *Therapeutic Delivery* 2017;8:577–595.

13. Soliman H, Mediavilla-Varela M, Antonia S. Indoleamine 2,3-dioxygenase: is it an immune suppressor? *Cancer journal (Sudbury, Mass)* 2010;16:354–9.

14. Criado G, Simelyte E, Inglis JJ, Essex D, Williams RO. Indoleamine 2,3 Dioxygenase– Mediated Tryptophan Catabolism Regulates Accumulation of Th1/Th17 Cells in the Joint in Collagen-Induced Arthritis. *ARTHRITIS & RHEUMATISM* 2009;60:1342–1351.

15. Fallarino F, Grohmann U, You S, McGrath BC, Cavener DR, Vacca C, et al. The combined effects of tryptophan starvation and tryptophan catabolites down-regulate T cell receptor zeta-chain and induce a regulatory phenotype in naive T cells. *Journal of immunology (Baltimore, Md : 1950)* 2006;176:6752–61.

16. Mellor AL, Munn DH. Tryptophan catabolism and T-cell tolerance: immunosuppression by starvation? *Immunology Today* 1999;20:469–473.

17. Wang XF, Wang HS, Wang H, Zhang F, Wang KF, Guo Q, et al. The role of indoleamine 2,3-dioxygenase (IDO) in immune tolerance: Focus on macrophage polarization of THP-1 cells. *Cellular Immunology* 2014;289:42–48.

18. Frumento G, Rotondo R, Tonetti M, Damonte G, Benatti U, Ferrara GB. Tryptophan-derived
catabolites are responsible for inhibition of T and natural killer cell proliferation induced by
indoleamine 2,3-dioxygenase. *Journal of Experimental Medicine* 2002;196:459–468.
19. Bracho-Sanchez E, Hassanzadeh A, Brusko MA, Wallet MA, Keselowsky BG. Dendritic
cells treated with exogenous indoleamine 2,3-dioxygenase maintain an immature phenotype
and suppress antigen-specific T cell proliferation. *Journal of Immunology and Regenerative*
Medicine 2019;5:100015.
20. Ochieng J, Furtak V, Lukyanov P. Extracellular functions of galectin-3. *Glycoconjugate*
Journal 2002;19:527–535.
21. Talaga ML, Fan N, Fueri AL, Brown RK, Bandyopadhyay P, Dam TK. Multitasking Human
Lectin Galectin-3 Interacts with Sulfated Glycosaminoglycans and Chondroitin Sulfate
Proteoglycans. *Biochemistry* 2016;55:4541–4551.
22. Farhadi SA, Bracho-Sanchez E, Fettis MM, Seroski DT, Freeman SL, Restuccia A, et al.
Locally anchoring enzymes to tissues via extracellular glycan recognition. *Nature*
Communications 2018;9:1–14.
23. Jacobs BY, Lakes EH, Reiter AJ, Lake SP, Ham TR, Leipzig ND, et al. The Open Source
GAITOR Suite for Rodent Gait Analysis. *Scientific Reports* 2018;8:9797.
24. Kloefkorn HE, Jacobs BY, Loye AM, Allen KD. Spatiotemporal gait compensations
following medial collateral ligament and medial meniscus injury in the rat: correlating gait
patterns to joint damage. *Arthritis research & therapy* 2015;17:287.
25. Jacobs BY, Dunnigan K, Pires-Fernandes M, Allen KD. Unique spatiotemporal and dynamic
gait compensations in the rat monoiodoacetate injection and medial meniscus transection
models of knee osteoarthritis. *Osteoarthritis and Cartilage* 2017;25:750–758.
26. Chaplan SR, Bach FW, Pogrel JW, Chung JM, Yaksh TL. Quantitative assessment of tactile
allodynia in the rat paw. *Journal of Neuroscience Methods* 1994;53:55–63.
27. Allen KD, Chan KM, Yarmola EG, Shah YY, Partain BD. The effects of age on the severity
of joint damage and intra-articular inflammation following a simulated medial meniscus
injury in 3, 6, and 9 month old male rats. *Connective Tissue Research* 2019:1–13.
28. Yarmola EG, Shah Y, Arnold DP, Dobson J, Allen KD. Magnetic Capture of a Molecular
Biomarker from Synovial Fluid in a Rat Model of Knee Osteoarthritis. *Annals of Biomedical*
Engineering 2016;44:1159–1169.
29. Yarmola EG, Shah YY, Lakes EH, Pacheco YC, Xie DF, Dobson J, et al. Use of magnetic
capture to identify elevated levels of CCL2 following intra-articular injection of
monoiodoacetate in rats. *Connective Tissue Research* 2019.
30. Kloefkorn HE, Jacobs BY, Xie DF, Allen KD. A graphic user interface for the evaluation of
knee osteoarthritis (GEKO): an open-source tool for histological grading. *Osteoarthritis and*
Cartilage 2019;27:114–117.
31. Kloefkorn HE, Allen KD. Quantitative histological grading methods to assess subchondral
bone and synovium changes subsequent to medial meniscus transection in the rat. *Connective*
Tissue Research 2017;58.
32. Chen D, Shen J, Zhao W, Wang T, Han L, Hamilton JL, et al. Osteoarthritis: Toward a
comprehensive understanding of pathological mechanism. *Bone Research* 2017;5:1–13.
33. Loeser RF, Goldring SR, Scanzello CR, Goldring MB. Osteoarthritis: a disease of the joint as
an organ. *Arthritis and rheumatism* 2012;64:1697–707.

34. Geiger BC, Wang S, Padera RF, Grodzinsky AJ, Hammond PT. Cartilage-penetrating
nanocarriers improve delivery and efficacy of growth factor treatment of osteoarthritis.
Science Translational Medicine 2018;10:eaat8800.
35. Zhang Z, Leong DJ, Xu L, He Z, Wang A, Navati M, et al. Curcumin slows osteoarthritis
progression and relieves osteoarthritis-associated pain symptoms in a post-traumatic
osteoarthritis mouse model. 2016.
36. Horisawa E, Hirota T, Kawazoe S, Yamada J, Yamamoto H, Takeuchi H, et al. Prolonged
anti-inflammatory action of DL-lactide/glycolide copolymer nanospheres containing
betamethasone sodium phosphate for an intra-articular delivery system in antigen-induced
arthritic rabbit. *Pharmaceutical Research* 2002;19:403–410.
37. Shi D, Xu X, Ye Y, Song K, Cheng Y, Di J, et al. Photo-cross-linked scaffold with
kartogenin-encapsulated nanoparticles for cartilage regeneration. *ACS Nano* 2016;10:1292–
1299.
38. Creamer P. Osteoarthritis pain and its treatment. *Current Opinion in Rheumatology*
2000;12:450–455.
39. Elola MT, Wolfenstein-Todel C, Troncoso MF, Vasta GR, Rabinovich GA. Review
Galectins: matricellular glycan-binding proteins linking cell adhesion, migration, and
survival.
40. Restuccia A, Fettis MM, Hudalla GA. Glycomaterials for immunomodulation,
immunotherapy, and infection prophylaxis. *Journal of Materials Chemistry B* 2016;4:1569–
1585.
41. Reesink HL, Bonnevie ED, Liu S, Shurer CR, Hollander MJ, Bonassar LJ, et al. Galectin-3
Binds to Lubricin and Reinforces the Lubricating Boundary Layer of Articular Cartilage.
Scientific Reports 2016;6:25463.
42. Hernandez JD, Baum LG. Ah, sweet mystery of death! Galectins and control of cell fate.
Glycobiology 2002;12:127–136.
43. Jacobs BY, Kloefkorn HE, Allen KD. Gait Analysis Methods for Rodent Models of
Osteoarthritis. *Current Pain and Headache Reports* 2014;18.
44. Carlson AK, Rawle RA, Adams E, Greenwood MC, Bothner B, June RK. Application of
global metabolomic profiling of synovial fluid for osteoarthritis biomarkers. *Biochemical and
Biophysical Research Communications* 2018;499:182–188.
45. Yeung AWS, Terentis AC, King NJC, Thomas SR. Role of indoleamine 2,3-dioxygenase in
health and disease. *Clinical Science* 2015;129:601–672.
46. Pallotta MT, Orabona C, Volpi C, Vacca C, Belladonna ML, Bianchi R, et al. Indoleamine
2,3-dioxygenase is a signaling protein in long-term tolerance by dendritic cells. *Nature
Immunology* 2011;12:870–878.
47. Huang L, Baban B, Johnson BA, Mellor AL, Mellor AL. Dendritic cells, indoleamine 2,3
dioxygenase and acquired immune privilege. *International reviews of immunology*
2010;29:133–55.
48. Szántó S, Koreny T, Mikecz K, Glant TT, Szekanecz Z, Varga J. Inhibition of indoleamine
2,3-dioxygenase-mediated tryptophan catabolism accelerates collagen-induced arthritis in
mice. *Arthritis Research and Therapy* 2007;9:1–7.
49. Bianco NR, Kim SH, Ruffner MA, Robbins PD. Therapeutic effect of exosomes from
indoleamine 2,3-dioxygenase-positive dendritic cells in collagen-induced arthritis and
delayed-type hypersensitivity disease models. *Arthritis & Rheumatism* 2009;60:380–389.

50. Williams RO. Exploitation of the IDO pathway in the therapy of rheumatoid arthritis.
International Journal of Tryptophan Research 2013;6:67–73.
570
571

Figures

Fig. 1: Joint retention and 4-week post-injection joint distribution for knees injected with

NL and NL-Gal3. A) The joint residence of NL-Gal3 was significantly longer than the joint

residence of unconjugated NL in both MCLT+MMT operated and contralateral joints ($p<0.0001$,

linear mixed effects model). Data are presented as mean + 95% confidence interval. B) Post-

mortem analysis of luminescence at 4 weeks after injection demonstrated the presence of NL-

Gal3 on the femur, patella/fat pad, meniscus, and tibia of MCLT+MMT operated and contralateral

joints. Unconjugated NL was not found.

Fig. 2: Tactile sensitivity before and after intra-articular injection of IDO-Gal3 or saline in MCLT+MMT operated knees. The 50% paw withdrawal threshold in MCLT+MMT operated limbs decreased at week 5 and week 7 relative to week 0 (^, $p<0.05$), indicating a heightened sensitivity to touch caused by MCLT+MMT surgery. Following intra-articular injections, 50% paw withdrawal thresholds increased in both saline and IDO-Gal3 treated animals relative to pretreatment (#, $p<0.05$), with 6 of 7 rats returning to baseline sensitivity levels in the IDO-Gal3 group by week 9. The average tactile sensitivity for IDO-Gal3 and saline treated animals did not vary after treatment; however, when change in tactile sensitivity relative to each animal's week 7 value was assessed (paired analysis), increases in tactile sensitivity in IDO-Gal3 treated animals were larger than saline treated animals at weeks 9-11 ($p \leq 0.041$, Fig. 2B). As Chaplan's up-down protocol provides a discrete set of data, results are shown as box plots with median, 25-75% interquartile range, and the lower and upper fence. Data outside the fence are plotted as individual data points.

Fig. 3: Walking velocity and spatiotemporal gait pattern analysis before and after intra-articular injection of IDO-Gal3 or saline in MCLT+MMT operated knees. Prior to treatment (weeks 3-7), saline and IDO-Gal3 treated rats used similar walking velocities (p≥0.89, Fig. 3A).

However, following treatment, rats treated with IDO-Gal3 used faster walking velocities than

saline controls on week 10 and week 11 (p<0.001, p=0.033, respectively). Balanced gaits are

defined as stance time being equal on the left and right limb, or a difference of stance time

between limbs being near zero. Temporally symmetric gaits have foot strike sequences that are

equally spaced in time, where a right foot strike occurs halfway between two left foot strikes (i.e.

temporal symmetry ≈ 0.5). Prior to treatment, saline and IDO-Gal3 treated rats used balanced,

symmetric gaits. After treatment, the gait of saline treated rats became temporally asymmetric at

weeks 8-11 (^symmetry > 0.5, p<0.05, Bonferroni-corrected t-test) and differed from IDO-Gal3

treatment at weeks 8-10 (p≤0.035, Tukey's HSD pairwise test, Fig. 3B). Neither saline nor IDO-

Gal3 treated animals had imbalanced stance times, though saline treated animals tended to

spend more time on the contralateral limb (imbalance > 0) while IDO-Gal3 treated animals tended

to spend more time on the affected limb (imbalance < 0). Here, at weeks 8 and 9, stance time

imbalance differed between saline and IDO-Gal3 treated animals (p≤0.025, Tukey's HSD pairwise

test, Fig. 3C). Spatial symmetry measures the symmetry of the foot placement, rather than the

timing of the foot strike. Again, a spatially symmetric gait has a right foot placement about halfway

between two left foot placements (spatial symmetry ≈ 0.5). Here, both IDO-Gal3 and saline treated

animals had spatially symmetric gait patterns throughout the experiment (Fig. 3D). This

experiment includes over 1500 gait trials, with 65-131 trials collected at each treatment-timepoint.

As such, data are plotted using density plots, with bars indicating the 95% confidence interval of

each treatment-timepoint mean as predicted by our linear mixed effects statistical model.

Fig. 4: Gait measurements before and after intra-articular injection of IDO-Gal3 or saline

in MCLT+MMT operated knees. Many gait parameters correlate to velocity. As such, velocity-

dependent gait parameters are first shown relative to velocity (left column), with each data point

shown. Then, shifts in gait parameters were analyzed both as averages and velocity-corrected

residuals (using week 7 data as the control line). This allowed for the evaluation of gait shifts

related to velocity changes (middle column) and independent of velocity changes (right column).

In conjunction with velocity changes, differences between IDO-Gal3 and saline treated rats were

found for average left contralateral limb percent stance time (p=0.04, treatment main effect), right

operated limb percent stance time (p=0.005, week-treatment interaction), stride length (p=0.02,

week-treatment interaction), and right operated limb peak vertical force (p=0.03, treatment main

effect). Specifically, differences between IDO-Gal3 and saline treated rats were found for average

left limb percent stance time at weeks 9, 10, and 11 (p≤0.048), average stride length at weeks 9,

10, and 11 (p≤0.046), and average right limb peak vertical force at weeks 9, 10, and 11 (p≤0.033).

Despite a significant interaction, differences at specific weeks were not detected for average right

limb percent stance time. Residualizing data to week 7 control lines eliminated these statistical

differences, indicating differences in the gait patterns of IDO-Gal3 and saline treated rats were

primarily related to increased walking speed. At week 11, stride lengths were longer in both IDO-

Gal3 and saline treated rats relative to the week 7 control line (^, p<0.05); this is typical for

experiments lasting longer than a month, as rats continue to have significant skeletal growth into

adulthood. Again, data are plotted using density plots, with bars indicating the 95% confidence

interval of each treatment-timepoint mean as predicted by our linear mixed effects statistical

model.

Fig. 5: Intra-articular levels of IL6, CCL2, and CTXII 4 weeks after intra-articular injection of IDO-Gal3 or saline in MCLT+MMT operated knees. At 4 weeks after intra-articular injection, joint levels of IL6 were significantly decreased in the IDO-Gal3 treated operated knees compared to saline treated operated knees (A, p=0.03). Intra-articular levels of CCL2 (B) and CTXII (C) were not elevated in saline treated operated knees compared to contralateral controls, and as such, shifts due to IDO-Gal3 treatment could not be detected (p=0.073 for CCL2 and p=0.090 for CTXII). Bars represent mean +/- 95% confidence interval.

Fig. 6: Histological scores of joint remodeling. At 12 weeks after surgery and 4 weeks after intra-articular injection, similar degrees of cartilage damage are seen in IDO-Gal3 and saline treated knees (A). In scoring the histological images, tibial plateau width (B), total affected cartilage width (C), medial joint capsule width (D), synovial cell density (E), and osteophyte area (F) were similar for IDO-Gal3 and saline treated groups. Epiphyseal trabecular bone area was lower in IDO-treated animals relative to saline controls, indicating marrow voids in the trabecular region were larger in IDO-Gal3 treated animals (G, p=0.031). Bars represent mean +/- 95% confidence intervals.

Supplemental Fig. 1: Flowchart describing the experimental design used to assess the joint residence and joint distribution of an enzyme after conjugation to galectin-3. Sixteen 3-month old, male Lewis Rats underwent medial collateral ligament transection plus medial meniscus transection (MCLT+MMT) surgery. At 8 weeks after MCLT+MMT surgery, rats were injected with either NanoLuc™ (NL, n=8) or NanoLuc™ galectin-3 (NL-Gal3, n=8) in both the operated and contralateral knees. Following NL or NL-Gal3 injection, rats were injected with the NL substrate, furimazine, and immediately imaged. Furimazine injections and IVIS imaging were repeated 1, 2, 4, 8, 12, 16, 20, 24, and 28 days after injection. After imaging on day 28, rats were euthanized and knees were dissected to isolate the patellar tissue, tibial tissue, femoral tissue, and meniscus for assessment of joint distribution.

Supplemental Fig. 2: Flowchart describing the experimental design used to assess the

ability of IDO-Gal3 to modulate OA-related pain and inflammation. Sixteen 3-month old, male

Lewis rats underwent baseline gait and tactile sensitivity testing, then MCLT+MMT surgery.

Following surgery, one rat was euthanized due to surgical complications, and another was

excluded after histological evaluation showed no evidence of meniscus transection (OA did not

develop). Remaining rats underwent gait assessments on post-surgery weeks 3, 5, 7, 8, 9, 10,

and 11, and tactile sensitivity testing on post-surgery weeks 1, 2, 3, 4, 5, 7, 8, 9, 10, and 11. At 8

weeks after MCLT+MMT surgery, rats received either an injection of saline (n=7) or IDO-Gal3

(n=7) in the MCLT+MMT operated knee. At 12 weeks post-surgery, all rats were euthanized, then

assessed for IL6, CCL2, and CTXII via magnetic capture and for joint damage using histological

evaluations.

Supplemental Fig. 3: Peak vertical force measurements before and after intra-articular injection of IDO-Gal3 or saline into MCLT+MMT operated knees. This figure provides an alternate visualization of data presented in Fig. 4D and 4E, placing the left and right limbs of saline and IDO-Gal3 animals next to each other. Here, saline animals showed lower peak vertical forces in their affected limbs at week 9 and week 11 (p≤0.003). After residualizing data to the week 7 control line, differences at week 9 remained (p=0.009), while differences at week 11 were no longer significant.

Supplemental Fig. 4: Synoviocyte viability after exposure to Gal3, IDO, and IDO-Gal3, as visualized with a LIVE/DEAD™ Cell Imaging Kit. In a preliminary experiment, rat synoviocyte viability after exposure to IDO-Gal3 was assessed via LIVE/DEAD™ Cell Imaging Kit (R37601, Invitrogen, Carlsbad, CA, USA) following manufacturer's instructions. Cells were grown to 70%-80% confluence. Then, synoviocytes were incubated with Triton X-100, tryptophan free media, kynurenine-supplemented media (2080 µg/mL), tryptophan free media with a kynurenine supplement (2080 µg/mL), Gal3-supplemented media (8.3 µg/mL), IDO-supplemented media (15 µg/mL, and IDO-Gal3-supplemented media (2.37 µg/mL) (n=5/group) for 24 hrs. Concentrations of tryptophan and kynurenine were selected to be 100x greater than levels found in the OA joint (12.67 mM/mL and 8.74 µg/mL, respectively [1]); IDO concentrations were based on necessary levels to affect 100x concentrations of tryptophan in a 24 hrs period. Synoviocyte death was not observed in tryptophan free media, kynurenine-supplemented media, tryptophan free media with a kynurenine supplement, Gal3-supplemented media, IDO-supplemented media, and IDO-Gal3-supplemented media. Treatment with Triton X-100 (positive control) elicited complete cell death.

[1] T. Igari, M. Tsuchizawa, T. Shimamura, Alteration of Tryptophan Metabolism in the Synovial Fluid of Patients with Rheumatoid Arthritis and Osteoarthritis, Tohoku J. Exp. Med. 153 (1987) 79–86. https://www.jstage.jst.go.jp/article/tjem1920/153/2/153_2_79/_pdf