Competent immune responses to SARS-CoV-2 variants in older adults following mRNA vaccination

Mladen Jergovič*[1,2], Jennifer L. Uhrlaub*[1,2], Makiko Watanabe[1,2], Christine M. Bradshaw[1,2], Lisa M. White[3], Bonnie J. LaFleur[3], Taylor Edwards[4], Ryan Sprissler[4,5], Michael Worobey[6], Deepta Bhattacharya[1,3], Janko Nikolich-Žugich[1,2,3,#]

1- Department of Immunobiology, University of Arizona College of Medicine, Tucson, AZ, USA

2- University of Arizona Center on Aging, University of Arizona, College of Medicine, Tucson, Tucson, AZ, USA;

3- BIO5 Institute, University of Arizona, Tucson, USA

4- University of Arizona Genetics Core, University of Arizona, Tucson, AZ, USA

5- Center for Applied Genetics and Genomic Medicine, University of Arizona, Tucson, AZ, USA

6-Department of Ecology and Evolutionary Biology, University of Arizona, Tucson, AZ 85721, USA.

- To whom correspondence should be addressed - nikolich@arizona.edu or by mail at P.O. Box 245221, Department of Immunobiology, University of Arizona College of Medicine-Tucson, 1501 N. Campbell Ave., Tucson, AZ 85724.

Graphical Abstract

Created with BioRender.com

Abstract

Aging is associated with a reduced magnitude of primary immune responses to

vaccination and constriction of immune receptor repertoire diversity. Clinical trials

demonstrate high efficacy of mRNA based SARS-CoV-2 vaccines in older adults but

concerns about virus variant escape have not been well addressed. We have

conducted an in-depth analysis of humoral and cellular immunity against an early-

pandemic viral isolate and compared that to the P.1. (Gamma) and B.1.617.2 (Delta)

variants, as well to a B.1.595 SARS-CoV-2 isolate bearing Spike mutation E484Q, in

<55 and >65 age cohorts of mRNA vaccine recipients. As reported, robust immunity

required the second dose of vaccine. Older vaccine recipients exhibited an expected 3-

5x reduction (but not a complete loss) in neutralizing antibody titers against both P.1.

(Gamma) and the B.1.595 virus at the peak of the boosted response. However, older

vaccinees manifest robust cellular immunity against early-pandemic SARS-CoV-2 and

more recent variants, which remained statistically comparable to the adult group. While

the duration of these immune responses remains to be determined over longer periods

of time, these results provide reasons for optimism regarding vaccine protection of older

adults against SARS-CoV-2 variants and inform thinking about boost vaccination with

variant vaccines.

eTOC summary: Vaccine responses are often diminished with aging, but we found

strong responses to SARS-CoV-2 in older adults following mRNA vaccination. T cell

responses were not diminished when confronted by SARS-CoV-2 variants. Neutralizing

Ab were reduced but not more than those in adults.

Introduction

RNA viruses have a high mutation rate resulting in diverse viral populations (Robson *et al.*, 2020). Since its emergence in the human population in late 2019, SARS-CoV-2 has infected more than 180 million people globally leading to the appearance of multiple new variants. Several variants of concern (VOC) have quickly become dominant in their countries of identification, spreading across the globe. B.1.1.7 (Alpha) was identified in the UK in December 2020 and spread to more than 90 other countries (Karim, 2021). Other identified VOC with broad spread include B.1.351 (Beta), P.1. (Gamma) and B.1.617.2 (Delta). These lineages are transmitted more efficiently, conferring evolutionary advantage over wild-type virus (Jewell, 2021). Two mRNA vaccines based on the first published genome sequence of SARS-CoV-2 (Wuhan/Hu-1/2019), BNT162b2 (Pfizer) and mRNA-1273 (Moderna), are being deployed to reduce COVID-19 disease severity and transmission, raising the obvious question of whether they will afford similar protection against variants of concern. Reduced plasma antibody (Ab) neutralization titers against pseudovirus in mRNA vaccine recipients has been reported for multiple variants (Garcia-Beltran *et al.*, 2021). Another report showed that B.1.351 (Beta) variant might be more effective in escaping humoral immunity than B.1.1.7 (Alpha) variant (Planas *et al.*, 2021), as one may expect from the location of its mutations to Ab-binding regions. Despite reduced humoral immunity, T cell responses to Alpha and Beta variants seem to be preserved in adult mRNA vaccine recipients (Geers *et al.*, 2021). Similar results were obtained with adenovirus vaccine recipients, which showed 3-5 fold lower neutralizing antibody titers against Beta and Gamma variants but uncompromised T cell responses (Alter *et al.*, 2021).

Efficacy and immunogenicity of many vaccines are known to be decreased in advanced age (Schmader *et al.*, 2012; Young *et al.*, 2017; Crooke *et al.*, 2019). Fortunately, SARS-CoV-2 mRNA based vaccines were shown to be both well tolerated and highly effective in older adults (Anderson *et al.*, 2020; Spector *et al.*, 2020; Li *et al.*, 2021). It remains unknown whether the breadth of the immune response to these mRNA vaccines will remain sufficient for protection against new viral variants.

To answer this, we have performed comprehensive analysis of antigen specific humoral and cellular immune responses in adult mRNA vaccinees of different ages. For humoral responses we compared a viral isolate representative of the early pandemic (USA/WA1/2020, hereafter 'WA1') to the P.1. (Gamma) and a local Arizona virus variant within lineage B.1.595 but, uniquely for that lineage, bearing Spike mutation E484Q (USA/AZ-WL-CVG00126, hereafter 'AZ-E484Q'). For T cell responses, we compared WA1 with both the P.1 (Gamma) and B.1.617.2 (Delta) variants. We show that the magnitude and neutralization capacity of the overall antibody response to these vaccines is not reduced in our older adult cohort (>65 years) as might have been expected based on historic vaccine studies. Regarding variants, we confirm lower neutralizing antibody titers against the P.1. (Gamma) variant and AZ-E484Q, with the reduction more pronounced in our older cohort than in younger adults. Importantly, robust cellular immunity against WA1, P.1. (Gamma), and B.1.617.2 (Delta) was preserved in both adult and old mRNA vaccine recipients. Overall, the effect of age on the immune response to the COVID mRNA vaccines was measurable, but smaller than expected for humoral immunity, and equal to that found in adults for cellular immunity, suggesting that these vaccines will provide considerable protection even as variants continue to emerge.

Results and Discussion

A total of 25 participants were enrolled in our study before receiving either the BNT162b2 (Pfizer) or mRNA-1273 (Moderna) COVID vaccine. Blood draws were collected prior to initial dose of vaccine, 7-9, and 18-26 days after first vaccine dose, and 7-14 days and 2-3 months after their second dose. These time points are designated as T0, T1, T2, T3, and T4 on all graphs. One participant was excluded from the study because they had a high neutralizing antibody titer and a strong antigen (Ag) specific T cell response before vaccination, indicating prior infection. Our final cohort included 15 participants aged 53 and younger (<55) and 9 participants over the age of 65 (>65).

Antibody ELISA assays for the receptor-binding domain (RBD) and S2 region of Spike

protein on plasma samples at each time point show similar rates and levels of

seroconversion between cohorts (Figure 1A, B). Area under curve (AUC) values confirm

that participants over age 65 have robust and comparable antibody responses to the

younger cohort with no significant differences measured at any time point (Figure

1C). Neutralizing antibody test titers utilizing WA1 were also comparable between both

cohorts across time points (Figure 2A). These data taken together are important

because they demonstrate that the humoral immune response in advanced age is

robust in response to COVID mRNA vaccines. We found a diminished capacity to

neutralize the P.1. (Gamma) variant and AZ-E484Q, with both being less well

neutralized at the third time point in the older cohort but only AZ-E484Q being less well

neutralized by the adult cohort (Figure 2A). E484K, a mutation in the receptor binding

domain (RBD) of Spike protein, is present in both B.1.351 (Beta) and P.1. (Gamma) and

has already been demonstrated to impact neutralization capacity (Wang *et al.*, 2021).

We chose AZ-E484Q to test against vaccine induced antibodies because it has Spike

E484Q and D614G mutations in common with B.1.617.1 (Kappa) but not L452R. L452R

has already been shown by others to impact neutralization in pseudotyped virus

systems (Li *et al.*, 2020) but it remained an open question how E484Q impacts

neutralization capacity (Verghese *et al.*, 2021). Our use of authentic B.1.595 SARS-

CoV-2, which in Spike bears only E484Q and D614G, demonstrates that E484Q also

affords the virus an opportunity to escape neutralization.

Antibody assays are considered the gold-standard when assessing the quality of

immunity after vaccination because antibodies can provide sterilizing immunity.

However, the establishment of memory B cell populations is also critical to lasting

vaccine efficacy as these are the resources available for response to the virus, or a

variant, upon next encounter. To assess whether there is an impact of age on the

formation of memory B cell populations we measured the frequency and number of

circulating memory B cells specific for SARS-CoV-2 pre-vaccination and one week after

booster dose by dual staining with tetramers specific for RBD and S1 using flow

cytometry (Shroff *et al.*, 2021) (gating strategy in Figure 3A). Both groups of participants

had a small (0.1-0.3% of all B cells) but detectable population of antigen specific cells

double positive for S1 and RBD tetramer (Figure 3A). This population increased as a

percentage of B cells post vaccination in all but two (one from each cohort) participants

with no difference between groups (Figure 3B). Similar results were obtained with S1

single-positive cells (Figure 3D) again with no difference between adult and older

participants post vaccination. By multiplying percentages of tetramer positive B cells

with total B cell percentage and lymphocyte counts we have calculated the absolute

numbers of SARS-CoV-2 specific B cells in circulation and again observed no difference

between cohorts for S1+RBD+ double positive cells (Figure 3C) or S1 single positive

cells (Figure 3E). Next, we examined the phenotype of the SARS-CoV-2 specific B cells

(gating strategy in Figure S1) with respect to class switching, as it has been previously

reported that aging is associated with a decline in the percentage and numbers of

switched memory B cells (Frasca *et al.*, 2008). To investigate this possibility, we

examined differentiation and class switching of total tetramer (S1+) positive B cells by

flow cytometric staining for CD27, IgM, IgD, CD21 and CD11c (representative flow

cytometric gating in Figure S1A). Ag-specific B cells from adult and older participants

expressed CD27 at identical levels (Figure S1B) and equal numbers of both CD27

positive and negative cells were class switched (Figure S1C,D). Adult and older

participants also displayed no difference in classical memory (CD21+) phenotype

among the class switched tetramer-positive cells (Figure S1E). Thus overall, we

conclude that induction and differentiation of SARS-CoV-2 specific B cells through

vaccination was not compromised by aging.

Finally, we measured antigen-specific T cells prior to, and at all time points available

post, vaccination. Given that ELISpot was previously reported to be a highly sensitive

method for detection of rare antigen specific T cells (Chauvat *et al.*, 2014) we

simultaneously measured the number of T cells expressing costimulatory molecules

CD137 and OX-40 by flow cytometry (used in several reports examining SARS-Cov-2

specific T cell immunity - (Grifoni *et al.*, 2020; Rydyznski Moderbacher *et al.*, 2020)

(representative flow cytometry in Figure S2A) and performed IFN-γ ELISpot on PBMCs

from select participants stimulated with peptide pools corresponding to the Spike protein

of WA1. Confirming prior data, ELISpot proved to be a much more sensitive method for

enumeration of Ag-specific T cells. We detected a statistically significant increase in

ELISpots per 10^6 PBMC's after vaccination analyzing just 5 samples, whereas parallel

flow cytometry samples showed no significant differences (Figure S2B-E). The only

limitation of ELISpot, as compared to flow cytometry based enumeration, is that total T

cell responses are measured without separate quantification of CD4 and CD8

responses.

We also analyzed Ag-specific T cell responses to stimulation with Spike protein peptide

pools from WA1 versus two variants of concern by ELISpot. Participant PBMC's were

stimulated with 16-mer overlapping peptide pools corresponding to the spike protein of

WA1, P.1. (Gamma) and B.1.617.2 (Delta). In accordance with previously published

results (Sahin *et al.*, 2020), mRNA vaccines induced a robust T cell response to WA1,

an early 2020 virus, which did not differ for Gamma and Delta variants, as evidenced by

a 10-fold increase in ELISpots from post-vaccination samples stimulated by S peptide

pools compared to unstimulated wells (Figure 4A). Next, we analyzed T cell responses

of both groups at different time points. Given that some samples had high responses in

the unstimulated wells we subtracted the number of ELISpots from the unstimulated

well to calculate the number of Ag-specific ELISpots for each time point. Data from aged

mice (Brien *et al.*, 2009; Fulton *et al.*, 2013; Jergović *et al.*, 2019) showed that induction

of Ag-specific T cell responses becomes delayed and decreased with age. In our data,

there was a slightly lower response in the older cohort at day 7 post first dose with all

three variants but it was not statistically significant (Figure 4B). However, after a second

(booster) dose both groups had a very robust T cell response against all three viral

variants examined (Figures 4C, D, E). Therefore, while we acknowledge the possibility

that the initiation of the Ag-specific T cell response may be delayed in some of our older

participants, we conclude that the outcome after booster dose is competent T cell

mediated immunity against WA1 and variants of concern virus in all mRNA vaccine

recipients in our study.

It is well established that primary immune responses wane with age and contribute to

the increased susceptibility to infection experienced by older adults (reviewed in

Nikolich-Žugich, 2018). There is also evidence that generation of immune memory in

older adults is reduced, but not maintenance of memory. Studies with multiple

pathogens have shown decreased T cell receptor repertoire with age (Yager *et al.*,

2008; Smithey *et al.*, 2018) which could mean easier escape from existing immunity for

pathogen variants. All of the above-mentioned findings warrant an extensive and long

term monitoring of immunity in SARS-CoV-2 vaccinees, especially those over age 65.

The emergence of SARS-CoV-2 effected older adults especially hard with more than

80% of deaths in those over age 65 (Lithander *et al.*, 2020). Vaccination reduces these

rates dramatically (Haas *et al.*, 2021) demonstrating that the principles of immunology

hold true for this virus and that immune memory is what our species is lacking. Well

tolerated and effective, SARS-CoV-2 mRNA vaccines induce potent humoral and

cellular immune responses (Anderson *et al.*, 2020; Sahin *et al.*, 2020). Their deployment

also offers an opportunity to establish, in a truly immune naive population, correlates,

and contours, of protective immunity. Recent elegant studies in rhesus macaques have

shown that even sub-sterilizing neutralizing antibody titers are protective and decrease

SARS-CoV-2 severity (McMahan *et al.*, 2021). The same research study demonstrated

a protective role for T cell memory responses and this is supported by human research

showing that accumulation of oligoclonal CD8 T cells in bronchoalveolar lavage fluid

inversely correlated with disease severity (Liao *et al.*, 2020). Recently, Collier et al.

analyzed 102 partially and 38 fully vaccinated subjects and concluded that subjects >80

years of age produced lower primary, but not secondary (boosted) antibody neutralizing

responses, including those against variants (Collier, Ferreira *et al.* 2021). They did not

analyze T cell immunity against variants and, somewhat curiously, did not observe

increased T cell responses following boosting in their older groups (>80). These authors

argue that older adults – those over 80 – remain vulnerable at least until they receive

the second vaccine dose. While this study with larger cohorts highlights the potential

importance of boosting the antibody response in the oldest adults, the key issue

remains that of biological relevance of the totality of measured differences. Given the

published and above-cited literature, we suggest that a decrease in neutralizing titer to

viral variants, accompanied by robust immune memory responses to both WA1 and the

variants, shown in this study for both B and T cells, gives reasons for optimism relative

to protection of older adults. This is supported by the efficacy of the mRNA vaccines in

protecting this vulnerable group. The decrease in antibody titer when challenged with

more recent SARS-CoV-2 variants does suggest that the breadth of immunity may be

more narrow in advanced age; a challenge that can be met with booster doses of

heterologous sequence virus. Further studies on the durability and breadth of

protection by current and future heterologous vaccines in older adults will be necessary

to answer all the germane questions on their immunity and SARS-CoV-2 protection.

Materials and methods

Study participants

This study was approved by the University of Arizona IRB (Protocol#2102460536). We collected blood from 16 adults <55 years old (mean age 31.3 ± 6.2) and 9 above 65 years old (71 ± 5.2). Demographics are provided in Table I. Samples for all participants were collected prior to vaccination, one week after first dose (mRNA vaccine: Pfizer N=15; Moderna N=10), day before booster dose and 7-10 days after booster. A smaller number of participants was sampled 3 months following first dose. The time points are labeled T0, T1, T2, T3, and T4 in all graphs. Blood for complete blood count was collected in BD vacutainer with EDTA and submitted to Sonora Quest Laboratories (Arizona). Blood for peripheral blood mononuclear cells (PBMCs) and plasma was collected in BD Vacutainer with sodium heparin. Plasma was separated by centrifugation at 1100 rpm for 10 minutes and PBMC was isolated from the buffy coat by Ficoll-Paque PLUS (GE Healthcare) and cryopreserved in fetal calf serum + 10% DMSO.

Virus

SARS-Related Coronavirus 2 (SARS-CoV-2), Isolate USA-WA1/2020, was deposited by Dr. Natalie J. Thornburg at the Centers for Disease Control and Prevention and obtained from the World Reference Center for Emerging Viruses and Arboviruses (WRCEVA). P.1. (Gamma) variant was obtained through BEI Resources, NIAID, NIH: SARS-Related Coronavirus 2, Isolate hCoV-19/Japan/TY7-503/2021, NR-54982, contributed by National Institute of Infectious Diseases. AZ-E484Q, a B.1.595 virus (GISAID: EPI_ISL_765942) was grown from a nasopharyngeal swab on Calu-3 cells for 48 hours. AZ-E484Q has Spike D614G and E484Q mutations but has not emerged as a variant of concern. Stocks of SARS-CoV-2 were generated as a single passage from received stock vial (USA-WA1/2020 and P.1.) or primary culture (B.1.595) on mycoplasma negative Vero cells (ATCC #CCL-81) at a MOI of 0.005 for 48 h. Supernatant and cell lysate were combined, subjected to a single freeze-thaw, and then centrifuged at 3000RPM for 10 min to remove cell debris.

Plaque Reduction Neutralization Test

A plaque reduction neutralization test (PRNT) for WA1 and SARS-CoV-2 variants was performed as described in Ripperger *et al.*, 2020. Briefly, mycoplasma negative Vero cells (ATCC # CCL-81) were plated in flat bottom, 96 well, tissue culture plates and grown overnight. Three-fold serial dilutions (1:20 - 1:43,740) of plasma/serum samples were incubated with 100 plaque forming units of virus for 1 h at 37°C. Plasma/serum dilutions plus virus were transferred to the cell plates and incubated for 2 h at 37°C, 5% CO_2 then overlayed with 1% methylcellulose in media. After 96 hours, plates were fixed with 10% Neutral Buffered Formalin for 30 min and stained with 1% crystal violet. Plaques were imaged using an ImmunoSpot Versa (Cellular Technology Limited) plate reader. The serum/plasma dilution that contained 10 or less plaques was designated as the NT90 titer.

ELISpot assays

T cell specific immunity to peptide pools corresponding to Spike, Nucleocapsid, and Matrix proteins were measured as previously described (Shroff *et al.*, 2021). Briefly, frozen PBMCs were thawed and rested overnight incubated in 24-well plates overnight at 37°C with 5% CO_2. Following day cells were stimulated in X-VIVO 15 media with 5% male human AB serum containing ~1 nmol of peptide pool corresponding of SARS-CoV-2 Prot S of US-WA (wild-type), Gamma (P.1./B1.1.28) and Delta (B.1.617.2) variants or positive control anti-CD3 mAb CD3-2 from or blank media as negative control. Overlapping 16mer peptide pools were purchased from 21st century Biochemicals,Inc. Cell suspensions were transferred to pre-coated Human IFN-γ ELISpot plus kit (Mabtech) and developed after 20h according to manufacturer instructions. Spots were imaged and counted using an ImmunoSpot Versa (Cellular Technology Limited).

Antibody ELISA

Serological assays were performed exactly as previously described in Shroff *et al.*,

2021). Area under the curve (AUC) values were calculated using R version 4.0.3 (R

Core Team, 2020) and the DescTools package (Signorelli, 2021).

Flow cytometry

Cryopreserved PBMC (2-5 x 10^6/sample) were thawed in prewarmed RPMI-1640 with L-

glutamine (Lonza) + 10% FCS. Thawed PBMCS were rested overnight at 37 ^{o}C in X-

VIVO15 Serum-free Hematopoietic Cell Medium (Lonza) supplemented with 5% human

Ab serum. Cells were stained with surface antibodies in PBS (Lonza) + 2% FCS, and

then stained with the live dead fixable blue dye (Thermofisher). Samples were acquired

using a Cytek Aurora cytometer (Cytek) and analyzed by FlowJo software (Tree Star).

Statistical analysis

SPSS and Graph Pad Prism were used for statistical analysis. Upon inspection of data

distribution by group differences were calculated either by Mann Whitney U-test, one-

way ANOVA or Kruskal Wallis test with post hoc correction. For all statistical differences

*$p<0.05$, **$p<0.01$, ***$p<0.001$. ****$p<0.0001$.

Competing Interest Statement

J.N.Ž. is co-chair of the scientific advisory board of and receives research funding from

Young Blood Institute, Inc. Sana Biotechnology has licensed intellectual property of D.B.

and Washington University in St. Louis. D.B. is a co-founder of Clade Therapeutics.

B.J.L. has a financial interest in Cofactor Genomics, Inc. and Iron Horse Dx.

Table I – Demographic characteristics of the cohort

	<55 yrs (N=16)	>65 yrs (N=9)
Gender [N (%)]		
Male	5 (31.3)	3 (33.3)
Female	11 (68.8)	6 (66.7)
Age [mean (range)]	33.3 (24-52)	71 (65-80)
Race [N (%)]		
White	8 (50)	7 (77.8)
Asian	8 (50)	1 (11.1)
Unknown	0	1 (11.1)
Vaccine Type [N (%)]		
Pfizer	9 (56.3)	3 (33.3)
Moderna	7 (43.8)	6 (66.7)

Figure 1. Antibody responses to mRNA vaccines are comparable regardless of age.

Time points denote days 7-9 (T1) and 18-26 (T2) days post first and 7-14 days (T3) and 2-3 months (T4) post second vaccine dose. Semi-quantitative 1:60 serum dilution ELISA results for reactivity to RBD **(A)** and S2 **(B)** of SARS-CoV-2 Spike protein. Individuals are connected across time points. Gray shaded area indicates positivity threshold for each assay. There is no statistical difference (Two-way ANOVA with Šídák's multiple comparisons test) between age groups at any time point measured. **C)**

Quantitative titers for RBD and S2 were calculated for each individual at each time point

and shown as area under the curve (AUC) values. There is no statistical difference

(Two-way ANOVA with Šídák's multiple comparisons test) between age groups at any

time point measured.

Figure 2

mRNA vaccines induce robust neutralization titers in adults under 55 and over 65

that is variably affected, but not abolished, by SARS-CoV-2 variants.

Time points are pre-bleed (T0); days 7-9 (T1) and 18-26 (T2) days post first dose of

vaccine; and 7-14 days (T3) and 2-3 months (T4) after second dose of vaccine. Virus

neutralization assays were performed using the USA-WA1/2020, P.1. (Gamma), and

AZ-E484Q isolates of SARS-CoV-2. Serial 1:3 dilutions of plasma were assayed for

neutralization capacity on Vero cells. The highest dilution capable of preventing >90% of

plaques was considered to be the PRNT90 value. Statistical significance was assessed

by Two-way ANOVA with Tukey's multiple comparisons test and determined significant

only at time point 3 as represented to the right of the legend. Neutralizing titers were

comparable between the <55 and > 65 cohorts to USA-WA1/2020; reduced against P.1.

(Gamma) as compared to WA1 for only the >65 cohort; and reduced in both cohorts

against B.1.595.

Figure 3. mRNA vaccines induce SARS-CoV-2 specific memory B cells in adult and old vaccinee cohorts. A) Representative flow cytometric gating after gating on total lymphocytes. Doublets and dead cells were excluded and percentage of S1 and RBD tetramer positive cells was analyzed on total CD19+ B cells; B) Percentage of tetramer double positive (S1+RBD+) B cells was increased after vaccine booster dose (T3) compared to pre-vaccination (T0) and to an equal extent in adult and old groups C) absolute numbers of S1+RBD+ B cells were also equally increased in <55 and >65 cohorts. D) Percentage of S1+ tetramer single positive B cells was equally increased in

<55 and >65 cohorts; **E)** as was their absolute number per ml of blood. Mann-Whitney U

test, for all statistical differences *p<0.05, **p<0.01, ***p<0.001. ****p<0.0001.

Figure 4. mRNA vaccines induce SARS-CoV-2 specific T cells reactive to Spike

peptide pools from all three variants A) 10^6 PBMCs per well were stimulated with

Spike peptide pools from USA-WA1/2020, P.1 (Gamma), and B.1.617.2 (Delta) cultured

overnight in pre-coated IFN-γ ELISpot plates. All three peptide pools induced an equally

strong IFN-γ response compared to unstimulated wells **B)** Ag-specific ELISpot numbers

were calculated by subtracting the unstimulated wells of each participant from the

peptide stimulated wells. USA-WA1/2020 induced an equal number of ELIspots in <55

and >65 cohorts at each time point. **C)** Similarly, P.1 (Gamma) peptide pool induced a

strong response at all time points post-vaccination which did not differ between <55 and

>65 cohorts. **D)** Identical results were obtained with stimulation with B.1.617.2 (Delta)

pool. Kruskal Wallis test with Dunn's post hoc correction. For all statistical differences

*p<0.05, **p<0.01, ***p<0.001. ****p<0.0001.

S1. SARS-CoV-2 specific memory B cells differentiate equally in <55 and >65

cohorts. A) Representative flow cytometric gating to examine B cell differentiation. **B)**

Percentage of CD27 expressing B cells among tetramer positive cells was equal

between cohorts. **C & D)** Equal percentage of SARS-CoV-2 specific B cells were class

switched among CD27 positive and negative cells between <55 and >65 cohorts. **E)**

Expression of CD21 (Classical memory) was also equal between <55 and >65 cohorts.

Mann-Whitney U test, for all statistical differences *p<0.05, **p<0.01, ***p<0.001.

****p<0.0001.

Figure S2. ELISpot is more specific method for enumeration of SARS-CoV-2

specific T cells. A) Representative flow cytometric gating for identifying CD137+OX-

40+ double positive CD4 and CD8 T cells. **B)** Representative unstimulated and Spike

protein peptide pool stimulated ELISpot wells. PBMCS from 5 participants were

stimulated overnight with USA-WA1/2020 spike peptide pool. SARS-CoV-2 specific T

cells were enumerated by either flow cytometry **(C,D)** or ELISpot **(E)**. ELISpot proved to

be a more sensitive method with statistically significant difference before and after

vaccination with N=5. Mann-Whitney U test, for all statistical differences *$p<0.05$,

**$p<0.01$, ***$p<0.001$. ****$p<0.0001$.

References

Anderson, E. J. *et al.* (2020) 'Safety and Immunogenicity of SARS-CoV-2 mRNA-1273 Vaccine in Older Adults', *New England Journal of Medicine*, 383(25), pp. 2427–2438. doi: 10.1056/nejmoa2028436.

Brien, J. D. *et al.* (2009) 'Key role of T cell defects in age-related vulnerability to West Nile virus.', *The Journal of experimental medicine*, 206(12), pp. 2735–2745. doi: 10.1084/jem.20090222.

Chauvat, A. *et al.* (2014) 'Clinical validation of IFNγ/IL-10 and IFNγ/IL-2 FluoroSpot assays for the detection of Tr1 T cells and influenza vaccine monitoring in humans.', *Human vaccines & immunotherapeutics*, 10(1), pp. 104–113. doi: 10.4161/hv.26593.

Collier DA, Ferreira I, Kotagiri P, et al. (2021) "Age-related immune response heterogeneity to SARS-CoV-2 vaccine BNT162b2." Nature.

Crooke, S. N. *et al.* (2019) 'Immunosenescence and human vaccine immune responses', *Immunity & Ageing*, 16(1), pp. 1–16. doi: 10.1186/s12979-019-0164-9.

Frasca, D. *et al.* (2008) 'Aging Down-Regulates the Transcription Factor E2A, Activation-Induced Cytidine Deaminase, and Ig Class Switch in Human B Cells', *The Journal of Immunology*, 180(8), pp. 5283 LP – 5290. doi: 10.4049/jimmunol.180.8.5283.

Fulton, R. B. *et al.* (2013) 'Aged mice exhibit a severely diminished CD8 T cell response following respiratory syncytial virus infection.', *Journal of virology*, 87(23), pp. 12694–700. doi: 10.1128/JVI.02282-12.

Grifoni, A. *et al.* (2020) 'Targets of T cell responses to SARS-CoV-2 coronavirus in humans with COVID-19 disease and unexposed individuals', *Cell*. doi: 10.1016/j.cell.2020.05.015.

Haas, E. J. *et al.* (2021) 'Impact and effectiveness of mRNA BNT162b2 vaccine against SARS-CoV-2 infections and COVID-19 cases, hospitalisations, and deaths following a nationwide vaccination campaign in Israel: an observational study using national surveillance data', *The Lancet*, 397(10287), pp. 1819–1829. doi: 10.1016/S0140-6736(21)00947-8.

Jergović, M. *et al.* (2019) 'Defective Transcriptional Programming of Effector CD8 T Cells in Aged Mice Is Cell-Extrinsic and Can Be Corrected by Administration of IL-12 and IL-18', *Frontiers in immunology*, 10, p. 2206. doi: 10.3389/fimmu.2019.02206.

Li, J. *et al.* (2021) 'Safety and immunogenicity of the SARS-CoV-2 BNT162b1 mRNA vaccine in younger and older Chinese adults: a randomized, placebo-controlled, double-blind phase 1 study', *Nature Medicine*, 27(6), pp. 1062–1070. doi: 10.1038/s41591-021-01330-9.

Li, Q. *et al.* (2020) 'The Impact of Mutations in SARS-CoV-2 Spike on Viral Infectivity and Antigenicity', *Cell*. 2020/07/17, 182(5), pp. 1284-1294.e9. doi: 10.1016/j.cell.2020.07.012.

Liao, M. *et al.* (2020) 'Single-cell landscape of bronchoalveolar immune cells in patients with COVID-19', *Nature Medicine*, 26(6), pp. 842–844. doi: 10.1038/s41591-020-0901-9.

Lithander, F. E. *et al.* (2020) 'COVID-19 in older people: A rapid clinical review', *Age and Ageing*, 49(4), pp. 501–515. doi: 10.1093/ageing/afaa093.

McMahan, K. *et al.* (2021) 'Correlates of protection against SARS-CoV-2 in rhesus macaques', *Nature*, 590(7847), pp. 630–634. doi: 10.1038/s41586-020-03041-6.

Nikolich-Žugich, J. (2018) 'The twilight of immunity: emerging concepts in aging of the immune system', *Nature Immunology*, 19(1), pp. 10–19. doi: 10.1038/s41590-017-0006-x.

Rydyznski Moderbacher, C. *et al.* (2020) 'Antigen-Specific Adaptive Immunity to SARS-CoV-2 in Acute COVID-19 and Associations with Age and Disease Severity', *Cell*, pp. 1–17. doi: 10.1016/j.cell.2020.09.038.

Sahin, U. *et al.* (2020) 'COVID-19 vaccine BNT162b1 elicits human antibody and TH1 T cell responses', *Nature*, 586(7830), pp. 594–599. doi: 10.1038/s41586-020-2814-7.

Schmader, K. E. *et al.* (2012) 'Efficacy, safety, and tolerability of herpes zoster vaccine in persons aged 50-59 years', *Clinical Infectious Diseases*, 54(7), pp. 922–928. doi: 10.1093/cid/cir970.

Shroff, R. T. *et al.* (2021) 'Immune Responses to COVID-19 mRNA Vaccines in Patients
with Solid Tumors on Active, Immunosuppressive Cancer Therapy', *medRxiv : the*
preprint server for health sciences, p. 2021.05.13.21257129. doi:
10.1101/2021.05.13.21257129.

Smithey, M. J. *et al.* (2018) 'Lifelong CMV infection improves immune defense in old
mice by broadening the mobilized TCR repertoire against third-party infection',
Proceedings of the National Academy of Sciences of the United States of America.
2018/07/02, 115(29), pp. E6817–E6825. doi: 10.1073/pnas.1719451115.

Spector, S. A. *et al.* (2020) 'Efficacy and Safety of the mRNA-1273 SARS-CoV-2
Vaccine', pp. 1–14. doi: 10.1056/NEJMoa2035389.

Verghese M, Jiang B, Iwai N, Mar M, Sahoo MK, Yamamoto F, Mfuh KO, Miller J, Wang
H, Zehnder J, Pinsky BA. A SARS-CoV-2 Variant with L452R and E484Q Neutralization
Resistance Mutations. J Clin Microbiol. 2021 Jun 18;59(7):e0074121. doi:
10.1128/JCM.00741-21. Epub 2021 Jun 18. PMID: 33952596.

Wang, R. *et al.* (2021) 'Analysis of SARS-CoV-2 variant mutations reveals neutralization
escape mechanisms and the ability to use ACE2 receptors from additional species.',
Immunity. doi: 10.1016/j.immuni.2021.06.003.

Yager, E. J. *et al.* (2008) 'Age-associated decline in T cell repertoire diversity leads to
holes in the repertoire and impaired immunity to influenza virus', *The Journal of*
Experimental Medicine, 205(3), pp. 711–723. doi: 10.1084/jem.20071140.

Young, B. *et al.* (2017) 'Do antibody responses to the influenza vaccine persist year-
round in the elderly? A systematic review and meta-analysis.', *Vaccine*, 35(2), pp. 212–
221. doi: 10.1016/j.vaccine.2016.11.013.

509