High-speed imaging of light-induced photoreceptor microsaccades in compound eyes

Joni Kemppainen[1], Neveen Mansour[1], Jouni Takalo[1] & Mikko Juusola[1,2*]

[1]Department of Biomedical Science, University of Sheffield, Sheffield S10 2TN, UK

[2]National Key Laboratory of Cognitive Neuroscience and Learning, Beijing, Beijing Normal

University, Beijing 100875, China.

* Correspondence: Mikko Juusola

m.juusola@sheffield.ac.uk

Keywords: photoreceptor, open rhabdom, fused rhabdom, microsaccade, *Drosophila*, deep pseudopupil, goniometry, hyperacuity, active sampling, active vision

Abstract

Inside compound eyes, photoreceptors contract to light changes, sharpening retinal images of the moving world in time. Current methods to measure these so-called photoreceptor microsaccades in living insects are spatially limited and technically challenging. Here, we present goniometric high-speed deep pseudopupil (GHS-DPP) microscopy to assess how the rhabdomeric insect photoreceptors and their microsaccades are organised across the compound eyes. This method enables non-invasive rhabdomere orientation mapping, whilst their microsaccades can be locally light-activated, revealing the eyes' underlying active sampling motifs. By comparing the microsaccades in wild-type *Drosophila*'s open rhabdom eyes to *spam*-mutant eyes, reverted to an ancestral fused rhabdom state, and honeybee's fused rhabdom eyes, we show how different eye types sample light

information. These results show different ways compound eyes initiate the conversion of spatial light

patterns in the environment into temporal neural signals and highlight how this active sampling can

evolve with insects' visual needs.

Introduction

Because the insect compound eyes extend from the rigid head exoskeleton, appearing stationary to

an outside observer, it was long assumed that their inner workings would also be static. Therefore, as

the eyes' ommatidial faceting sets their photoreceptor spacing, it was deduced that the compound

eyes could only sample a pixelated low-resolution image of the world[1-3].

Fig. 1 | Active sampling by photomechanical photoreceptor microsaccades and the deep pseudopupil phenomena. a, In *the conventional static sampling theory*[1, 2], ommatidial facets set a compound eye's photoreceptor spacing, limiting the finest image details the eye could resolve. However, inside an ommatidium, incoming light intensity changes make its R1-R7/8 photoreceptors rapidly recoil axially and swing laterally. These so-called ultrafast photoreceptor microsaccades enable *Drosophila* to see the world in a finer resolution than its eyes' photoreceptor spacing, explained by *the new active sampling theory*[4, 5]. Left: *Drosophila* eye computer graphics (CG) model highlights the axial microsaccade component; R1-R7/8s first recoil and then slide towards the ommatidium lens. Right: concurrently, the light-activated R1-R7/8 also swing sideways (laterally).

A local (incident) light intensity change evokes microsaccades only in those ommatidia facing the stimulus. If this happens in the frontal left and right eye ommatidia with overlapping receptive fields, their microsaccades are synchronous yet have mirror-symmetric lateral components[4, 5]. Meanwhile, elsewhere across the eyes, the photoreceptors stay still because the eye curvature and the ommatidial screening pigments block them from seeing the stimulus[4, 5]. **b**, The optical principle of the deep pseudopupil (DPP). DPP is a virtual image of several distal R1-R7/8 rhabdomere tips (highlighted in blue, yellow and green for three nearby ommatidia), which align with the angle the eye is observed at while being ~10x-magnified by the ommatidial lens system. These virtual rhabdomere images are optically brought together when the microscopes' focal plane is ~200 μm under the eye surface (as shown in image 2). Because of the optical magnification, the rhabdomere tips, which appear deep inside the eye, are actually positioned at ~20 μm from the inner surface of the ommatidium lens.

However, recent results on the *Drosophila* compound eyes are now replacing this *static* viewpoint with a new concept of morphodynamic *active* sampling[4, 5]. Sophisticated experiments have revealed how photoreceptor microsaccades *locally*[4, 5] (**Fig. 1a**) and intraocular muscle contractions *globally*[4-7] move, stretch and recoil intraommatidial optical structures, improving vision morphodynamically. During the *local ultrafast* (<100 ms) *photomechanical* microsaccades, the photoreceptors of a single ommatidium concurrently recoil axially (**Fig. 1a**, left) and swing laterally (right) to increase sampling resolution in space and sharpen light input in time for super-resolution vision[4]. And, with the left and right eye photoreceptor pairs generating mirror-symmetric microsaccades, this active sampling further expands the flies' hyperacute stereopsis[5]. Conversely, the intraocular muscle contractions shift one eye's entire retina (its sampling matrix) *globally* regarding the other eye[4-7]. In head-immobilised *Drosophila*, these drifts and vergence movements, which also happen underneath the eyes' rigid ommatidial lens cover, hidden from the outside view, are typically 10-to-100-times slower than the local photoreceptor microsaccades[4, 5]. But in freely behaving flies[6], their dynamics may strengthen to combat adaptive perceptual fading[4, 7] and contribute to attentive saccadic viewing and object tracking[6].

Minute photomechanical photoreceptor contractions (<~200 nm) were first measured in *ex vivo*

Drosophila preparations using atomic force microscopy (AMF)[8]. Initially, these movements, caused

by PIP$_2$ cleavage in the microvillar photoreceptor membrane[8, 9], were thought to be too small to alter

the photoreceptors' light input[8]. However, later live-microscopy experiments[4], using the cornea

neutralisation method[10], showed that *ex vivo* AFM underestimates the size of the lateral rhabdomere

movements[4]. Intense light modulation *in vivo* could rapidly swing an R1-R6 rhabdomere about its

width (~1,400 nm) sideways. And with similar microsaccades also occurring in synaptically decoupled

photoreceptors[4], the results demonstrated active sampling inside an ommatidium for the first time[4].

Unfortunately, both these methods are technically demanding and spatially limiting and thus ill-suited

for mapping the microsaccade movement sizes and directions across the left and right compound

eyes.

In the *Drosophila* compound eyes, an optical phenomenon called the deep pseudopupil (DPP) arises

from the regular arrangement of ommatidia (**Fig. 1b**), each containing R1-R7/8 photoreceptors in

which open rhabdomeres act as waveguides[11]. By focussing a microscope's image plane below the

eye's outer surface (**Fig. 1b**, inset), virtual images of several ommatidia's R1-R7/8 rhabdomere

patterns (**Fig. 1b**, highlighted in blue, green and yellow for the nearby ommatidia) become

superimposed, revealing their stereotypical yet ~10-times magnified trapezoid arrangement. And

since these virtual images fuse at the microscope focal plane of ~200 μm, we see the rhabdomere

tips inside the eye, ~20 μm away from the inner surface of their ommatidium lenses. Thus, DPP

microscopy offers a versatile, non-invasive method to observe retinal tissue in living flies and other

insects (**Fig. 2**). First, to observe a well-defined, clear DPP pattern in dipteran eyes requires precisely

organised rhabdomeres across the neighbouring ommatidia[12], and DPP microscopy with epi-

illumination[13, 14] (frontally, through the eye optics) has been used to study retinal degeneration that

breaks this order[15-17]. Second, because the rhabdomeres contributing to the DPP image are those

facing the observer, the DPP microscopy provides the "gold standard" measure for the binocular

77   overlap over the left and right compound eyes[18-20]. Finally, because any lateral retinal tissue

78   movement shifts the DPP similarly, DPP microscopy can be used to investigate how the eye-muscle-

79   induced retinal micromovements shift the photoreceptors' receptive fields[5, 21].

Fig. 2 | Goniometric high-speed deep pseudopupil (GHS-DPP) imaging system. a, Its integral

parts are a sideways mounted stereo microscope with a high-speed (100-1,000 fps) digital camera

and a goniometric rotation stage system. *Drosophila* eyes are imaged under the antidromic dual

infrared illumination, invisible to the flies[4, 22]. Photoreceptor microsaccades are activated by

ultraviolet (UV or green) light stimulation, delivered through the ocular slot system. **b**, The fly's x/y-

rotations are read using a 1,024-step rotary encoder and an Arduino board. **c**, The high-power LEDs and the camera were controlled over the BNC interface. **d**, An infrared (IR) passing but UV blocking optical filter in the front of the camera decouples the UV-stimulus from the imaging pathway. See **Supplementary Video 1**.

Here, we present a novel goniometric high-speed deep pseudopupil (GHS-DPP) microscopy (**Fig. 2**)

with invisible (850 nm) infrared back-illumination, developed to study active sampling in insect

compound eyes. We first use it to measure the photoreceptor microsaccade dynamics and directions

in wild-type *Drosophila melanogaster*, possessing the archetypical open rhabdom dipteran eyes (**Fig.**

3, top), and transgenic *spam* null-mutants, in which rhabdomeres fail to separate (**Fig. 3**, middle),

forming an ancestral, fused rhabdom (apposition) eye[23]. Finally, we measure photoreceptor

microsaccades in the Honeybee (*Apis mellifera*) apposition eyes (Fig. 3, bottom) and compare these

dynamics to wild-type and *spam Drosophila*. Our results show that active sampling by photoreceptor

microsaccades occurs both in the open and fused rhabdom eyes. We analyse their functional

similarities and differences and discuss what these results could mean for the evolution of compound

eyes and insect vision in general[24].

Fig. 3 | Deep pseudopupil (DPP) in the wild-type and *spam Drosophila* and honeybee apposition eyes. a, Left: the *Drosophila* compound eye comprises ~750 ommatidia, each having a light focusing facet lens and eight photoreceptors; with their light-sensitive parts, the rhabdomeres (coloured), protruding centrally. Middle: Computer graphics (CG) models of the open wild-type rhabdomeres; here shown for their left eye's southern hemisphere. Right: The CG-model simulated DPP image appears as a magnified "virtual" image of the open rhabdomere patterns. Here, DPP is

shown at the focal plane of 200 μm into the eye for the right eye ventral section, corresponding to frame #2 in **Fig. 1b** and the bottom-left frame in **Fig 4b**.

b, Left: in the *spam*, R1-R7/8 rhabdomeres reverted back into their ancestral fused rhabdom form. Middle: inside the ommatidia (in the retinal tissue), the fused rhabdom resembles a single rod. Right: their DPP appears as a tiny bright disc. Colouring indicates the contributing R1-R7/8 rhabdomeres' relative locations in the virtual DPP rhabdom image.

c, Left and Middle: In the honeybee apposition eye ommatidia, the photoreceptor rhabdomeres form a single rod-like waveguide, the rhabdom, centred ~100 μm behind the ommatidial lens.

Right: In the DPP image, the honeybee fused rhabdom appears as a single disk, similar to *spam*.

Results

Left and right eye DPP patterning rotate systematically and mirror-symmetrically

We first inspected the wild-type and *spam* flies' DPP patterns in still images taken under antidromic infrared illumination in perceptual darkness (**Fig. 4**). Characteristically, the wild-type DPP image merged the neighbouring ommatidia's R1-R7/8 rhabdomere images into a classic trapezoid pattern of seven small bright discs (**Fig. 4a**). In contrast, the *spam* DPP appeared as a tiny bright disc (**Fig. 4d**). These different DPP patterns were entirely expected, as they directly follow the deep pseudopupil theory[12] (**Fig. 1b**) and are reproduced by our CG-modelling (**Fig. 3**).

Fig. 4 | Wild-type and *spam* DPP imaged by the 850 nm GHS-DPP microscopy. a, Forward-facing wild-type eyes (horizontal rotation = 0°) with the red and blue square boxes indicating their left and right DPPs, respectively. **b**, Wild-type eyes, viewed on the dorsal side (1st row), at the midline (2nd row), and on the ventral side (3rd row), reveal their DPPS ventral-dorsal symmetry. **c**, The wild-type rhabdomere orientation across the left and right eyes as mapped with the GHS-DPP

microscopy. The green lines indicate the R1-R2-R3 rhabdomere axes, and the yellow lines the R3-R4-R5 axes. R1-R7/8 rhabdomeres rotate systematically with changing eye location, producing concentrically-expanding diamonds-patterns[5], highlighting the characteristic right-left eye and north-south (dorsal-ventral) hemisphere mirror-symmetricity. **d**, The left and right eye *spam* DPPs appear circular-symmetric discs and are more oversaturated due to their eyes' lighter pigmentation. (**e**) *spam* eyes with their DPPs are viewed on the dorsal side, midline, and ventral side.

Because *Drosophila* has a small head and fairly transparent (not densely pigmented) cuticle, the

infrared GHS-DPP microscopy makes it straightforward to record, measure and map the DPP pattern

changes across the eyes. Specifically, for *Drosophila*, we could use relatively low-power infrared

illumination - propagating through its intact head - and still obtain high signal-to-noise DPP imaging at

high frame rates (≥100 fps). The real benefit was that since the tested flies required no surgical cuticle

removal (to improve infrared throughput for better DPP image quality) and suffered very little or no

heat damage, individual *Drosophila* regularly provided consistent, repeatable results throughout

hours-long experiments.

In the wild-type, the left and right eyes' DPPs are mirror-symmetric: shown in the binocular upper-

frontal view (**Fig. 4a**; the blue and red boxes highlight their right and left DPPs, respectively).

Interestingly, the north and south hemispheres of the eye also have mirror-symmetric DPP patterns[25]

but fuse at the equator (mid-line) to form larger elongated triangle shapes (**Fig. 4b**). The DPP

orientation, as the angular rotation between R3-R2-R1 rhabdomeres (yellow line) and R3-R4-R5

rhabdomeres (green), shifts between nearby eye locations in regular small steps, generating the left

and right eyes' mirror-symmetric global map (**Fig. 4c**). In this global map, local DPP alignments follow

a concentrically expanding diamond-shaped pattern. Therefore, the underlying developmentally

rotated R1-R7/8 rhabdomere orientations at each eye position align with the frontally expanding optic

flow field[5] (**Supplementary Video 3**). On the contrary, in the *spam* mutant, because their DPPs

appear as homogeneous circular or oval discs (**Fig. 4d**), neither such gradual rotations nor their left-

right and north-south divisions were readily observable (**Fig. 4e**).

Photoreceptor microsaccades' lateral and axial components

After imaging the eyes' DPPs, we tested whether the *spam* eyes can generate ultrafast (time-to-peak

<100 ms) photomechanical microsaccades, akin to the open rhabdom wild-type[4, 5] eyes. In the first

instance (**Fig. 5a** and **b**), these experiments were performed at the fixed left and right eye locations

(±28° horizontal, -37° vertical) using a bright 200-ms-long ultraviolet (365 nm) flash stimulus

(**Supplementary Video 2**). In each fly, the UV flash was delivered locally at the centre of the observed

DPP photoreceptors' receptive fields (**Supplementary Video 4**) and repeated 25 times to obtain

robust estimates and statistics of the resulting response dynamics.

A flash of any wavelength R1-R8 photoreceptors are sensitive to (~300 to ~650 nm)[22, 26] evokes a

photoreceptor microsaccade[5]. Inside an ommatidium, the number of light-activated photoreceptors

and their combined contraction strength set its photoreceptor microsaccade amplitude[5]. Because

these photoreceptors are mechanically coupled and likely pivoted[5], it only takes one (say R1) to be

light-activated, and its contraction alone can move its neighbours (R2-R8) too[5] (**Supplementary**

Video 4). As *Drosophila* R1-R6 possesses the sensitising UV-pigment and R7s are UV-sensitive[22, 26],

a UV-flash evokes larger photoreceptor microsaccades than, say, an amber-flash, which only

activates R8y cells[5]. Therefore, in *Drosophila* experiments, UV-flash is a good choice of stimulus.

In wild-type, a UV flash always evoked a local photomechanical photoreceptor microsaccade, making

the observed DPP rapidly jump laterally in the front-to-back (north-west) direction (**Fig. 5c**) before

swiftly returning in darkness, as expected for normal eye function[4, 5]. However, remarkably, we found

that the fused rhabdom *spam* eyes also generate robust ultrafast DPP microsaccades (**Fig. 5d**) with

broadly similar temporal dynamics. In both phenotypes, besides moving laterally (**Fig. 5d**, above), the

photoreceptors moved simultaneously also axially (below). This axial component, resulting from the

rhabdomeres contracting away and elongating towards the ommatidium lens (and the camera)[5], can

be directly measured as a proportional DPP darkening and lightening[5]. To eliminate any motion

artefacts, we measured the axial component from the DPP image pixels' dynamic intensity change,

tracking frame-by-frame only the pixels within the rhabdomere tips. As expected, the fast axial DPP

brightness changes systematically time-locked with the corresponding lateral DPP movements (**Fig.**

5d-f), consistent with both phenotypes having the same photomechanical phototransduction origin.

Qualitatively, both the wild-type and *spam* had similar looking microsaccade kinematics and

probabilities (**Fig. 5d-f**), but the overall displacement amplitudes appeared much smaller in *spam* (**Fig.**

5d, below). Maximum amplitude measurements confirmed that the *spam* DPP microsaccades (both

their lateral and axial components) were indeed smaller than the wild-type (**Fig. 5g**). Similarly, the

spam flies' calculated maximum microsaccade activation speed was significantly slower (**Fig. 5h**).

However, these phenotypes' logarithmic growth factors during the activation phase did not differ

statistically (**Fig. 5i**), meaning that the *spam* DPP microsaccades' slower speed resulted from their

shorter travelling distance but not from changes in their duration. Accordingly, there was no significant

change in the half-rise times (**Fig. 5j**). The similar logarithmic growth factors and half-rise times

suggest that the likely photomechanical cause of the DPP microsaccades, the PIP_2 cleavage from the

microvillar membrane during phototransduction[8], is unaffected by the *spam* mutation. Overall, these

results show that the *spam* flies have similar microsaccade kinematics as the wild-type but are only

smaller.

We further inspected individual flies' DPP microsaccade variations to the 25 times repeated light

flashes. A qualitative comparison between the selected wild-type and *spam* flies showed that most

variation occurs in the total microsaccade size; that is, how far the DPP and the rhabdomeres travel

during the activation phase (**Fig. 5k**). Furthermore, we found no difference between their DPP

displacements' standard deviations (**Fig. 5l**), indicating similar intrinsic amplitude jitter, or

photomechanical stochasticity, affecting wild-type and *spam* microsaccades. These results highlight

that DPP microsaccades vary considerably between the flies and between individual flies' repeated

responses, suggesting that their kinematics may further reflect top-down regulation from the brain[27,]

[28], i.e. the flies internal state (circadian rhythm, attention or activity state)[27, 29, 30]. Such variability could

result, for instance, from the slow eye-muscle-induced whole retina drifting[5], structurally fluctuating

the local rhabdomere tension (their anchoring and pivoting dynamics) from one trial to another[5].

Notably, however, the real benefit of the local photoreceptor microsaccades' stochastic variability,

irrespective of its cause, is that it effectively removes aliasing from the retinal sampling matrix[4, 5, 31].

Fig. 5 | DPP microsaccade kinematics in the wild-type and *spam*. a, In the local recordings, the UV flash was repeated 25 times following a regular inter-stimulus interval (ISI). **b**, These experiments were done at one fixed location in each eye. **c**, Image time series from the stimulus onset (time zero) to 180 ms later, showing how the same DPP pattern, plotted against fixed red x-y-coordinate axes, moves back-to-front (North-West) about one rhabdomere width (~1.5 mm inside

ommatidia). **d**, Wild-type and spam DDP patents and their tracked movement directions (orange arrows). Mean rhabdomere lateral (continuous traces) and axial (fast DPP darkening/brightening, dotted) displacement components in the wild-type (1st row) and *spam* (2nd row) appear similar in shape but are smaller in *spam*. The thick lines show the corresponding population means. **e-f**, The probability plots, containing the separate (single) responses, indicate that the photoreceptor microsaccades occur over a predictable but variable range. **g-j**, The microsaccade amplitude, the top speed, the logistic growth factor and the half-rise time, respectively, in the wild-type and *spam* (Welch's t-test; p = 0.027, p = 0.031, p = 0.86, p = 0.62). **k**, The DPP microsaccade responses of one selected wild-type and one *spam* show variability in repeated responses that may indicate purposeful modulation. **l**, Quantified displacement standard deviations (std) for all the wild-type and *spam* flies do not differ statistically (Welch's t-test; p = 0.16).

Mapping photoreceptor microsaccades' movement directions across the eyes

Next, we mapped DPP microsaccades and their directions from about 200 distinct locations over the

left and right eyes (**Fig. 6a** and **b; Supplementary Video 5**). In the wild-type, the DPP microsaccades

moved approximately from back-to-front or south-to-north, depending on the eye region (**Fig. 6c**). But

we did not image the DPP microsaccade relaxation phase, which occurs during light decrements

(darkening), moving slower and in opposite directions to the activation phase[4, 5] (returning

rhabdomeres to their initial starting positions).

Fig. 6 | DPP microsaccades across the left and right wild-type and *spam* eyes. a, During the 200 ms UV flash, the camera recorded the DPP microsaccades 100 frames-per-second. **b**, After each flash, the fly was rotated, recording approximately 200 locations on the left and right eyes. Blue arrows indicate DPP microsaccade directions. **c**, Mean microsaccade vector map of wild-type flies (N = 5; the arrows indicate eye-location-specific normalised microsaccade directions). The small insets show the dorsal, the anterior and the ventral vector map views from the top to bottom. **d**, Mean microsaccade vector map of spam files (N=5) appears similar to the wild-type but only slightly noisier. **e**, Angular difference plot of the wild-type and *spam* vector maps show that the wild-type and *spam* vector maps are mostly collinear. **f**, On the anterior eye regions, the angular

difference graph suggests that the *spam* right eye microsaccades are clockwise-rotated and the left eye microsaccades counterclockwise-rotated compared to the wild-type (as viewed from outside). See **Supplementary Videos 2** and **5**.

Interestingly, we discovered that the fused rhabdom *spam* DPP microsaccades generally moved along

similar directions to those of wild-type (**Fig. 6d**). We calculated the absolute angular difference in the

rotation of the microsaccades between the wild-type and *spam* flies and found their resulting global

maps broadly similar (**Fig. 6e**; **Supplementary Video 5**). However, in a more detailed inspection, the

spam left eye DPP microsaccades on the anterior parts showed slightly more counterclockwise

rotation and the right eye DPP microsaccades slightly more clockwise rotation than the wild-type (**Fig.**

6f).

These results demonstrate active sampling - by photomechanical photoreceptor microsaccades -

occurring in a spatially-coordinated manner across the *spam* fused rhabdom eyes. Moreover, since

the wild-type microsaccade directions (their forward-and-back movement axes) align with their R1-

R2-R3 rhabdomere axis (**Fig. 3c**, yellow arrows)[5], it seems highly likely that the *spam* R1-R8

photoreceptors inside the ommatidia would also rotate in an eye-position-dependent manner, resulting

in the ventral-dorsal hemispheric DPP mirroring. However, we could not directly confirm this from the

round and oval *spam* DPP images (**Fig. 3e**). Nevertheless, in both phenotypes, the photoreceptors'

photomechanical active sampling makes their receptive fields scan the world in their eye-location-

specific directions, broadly matching the flies' concentrically expanding optic flow field during the

forward locomotion[5] (**Supplementary Video 6**).

Fig. 7 | DPP microsaccades in honeybee fused rhabdom apposition eyes.

a, Head-immobilised living honeybee prepared for the high-speed DPP imaging experiments.

b, Honeybee compound eye image series under antidromic infrared illumination, when the microscope gradually focusses inside the eye. First, the facet lenses appear (frames #3 and #4) before the DPP emerges as a small bright disk (frames #11 and #12).

c, High-speed imaging of photomechanical photoreceptor microsaccades. A characteristic slightly elongated DPP image of superimposed rhabdoms of neighbouring ommatidia. The DPP elongation is caused by the "ellipsoid" (non-spherical) honeybee eye shape. The scale bars give the actual

rhabdom dimensions in ommatidia. The tested UV and green flashes evoked photomechanical photoreceptor microsaccades, seen as ultrafast lateral DPP "jump" and axial DPP darkening synchronised in time. The average DPP microsaccade to the green flash (green traces) was 1.54-times larger than to the UV flash (purple traces), consistent with the ommatidia having more 1.6-times more green photoreceptors (4) than UV (2.5) photoreceptors (inset).

d, Spontaneous slow eye-muscle-movement-induced (whole retina) DPP drifts were 10-to-100-times slower than light flash triggered photomechanical photoreceptor microsaccades. However, unlike in *Drosophila*[5], the sluggish eye-muscle-activity shifted the Honeybee DPP positions over seconds and minutes more than the ultrafast (<200 ms) DPP microsaccade displacements superimposed on them.

Photoreceptor microsaccades in honeybee fused rhabdom apposition eyes

Our current results, together with those from the previous studies[4, 5, 8, 9], indicate that photomechanical

transduction reactions within *Drosophila* photoreceptors are responsible for their ultrafast light-

induced microsaccades. As these PIP_2-bound reaction steps[8] are thought to be evolutionarily

conserved in rhabdomeric photoreceptors[32, 33], we next tested whether the Honeybee (**Fig. 7**) fused

rhabdom apposition eyes also generate photomechanical photoreceptor microsaccades.

The honeybee eyes and head are about 10-times larger than *Drosophila*'s and more densely

pigmented, thicker, and sturdier. With long hairs, mouthparts and antennae sticking out, the bee head

looks striking, like a 16th-century knight's helmet (**Fig. 7a**). Unfortunately, these structural features

complicate GHS-DPP microscopy. The head's thick pigmented cuticle filters out infrared light and the

antidromic light path (from the back of the head to the rhabdomere tips) is ~10-times longer than in

Drosophila. Therefore, we had to surgically remove parts of the rear head cuticle and use a more

powerful and condensed infrared beam to achieve sufficiently-high signal-to-noise ratio DPP images

of the back-illuminated rhabdoms (**Fig. 7b**). Once a honeybee was aligned correctly, a point-like DPP

image of one superimposed rhabdom (**Fig. 7b**, frame #12), collected from the neighbouring

ommatidia, emerged with the microscope focussing through the ommatidial surface into the eye. These DPP patterns matched our CG-model prediction (cf. **Fig. 3**) and appeared similar to those in the *spam* fused rhabdom eye (cf. **Fig. 4e**).

We learned through trial and error not to strive for the best spatial resolution in high-speed imaging experiments. Instead, we optimised the setup to enable high-speed *in vivo* imaging by balancing the infrared power and exposure time with the DPP image contrast and temporal resolution (**Fig. 7c**, left). This optimization process was complex as the temporal resolution was essential to avoid blurring caused by the photoreceptor contractions. Too low infrared power or frame rate, and the small and fast photoreceptor microsaccades were undetectable from the noisy DPP images. Too high power, and the honeybee was near-instantly killed by the heat cooking its brain. However, with appropriate settings, we could repeatedly record bee photoreceptor microsaccades *in vivo* (**Fig. 7c**).

Honeybee fused rhabdom photoreceptors generated photomechanical microsaccades to both tested UV (**Fig. 7c**, middle) and green light flashes (right); the given examples were recorded after prolonged dark-adaptation. Unsurprisingly, the DPP microsaccades were relatively small, with their maximum displacement range (≤ 1 µm) equating to ≤ 1° receptive field jumps in space, being close to our earlier prediction[5]. This prediction was based on the bee rhabdom's envisaged "rod-like" rigidity, ~1.9° receptive field half-width[34] and the ~1° interommatidial angle[1] (see Discussion). Notably, the green flash evoked on average 1.54-times larger microsaccades than the UV flash. The greater green-sensitivity is consistent with each honeybee ommatidium (**Fig. 7c**, inset) having 4 large green-sensitive photoreceptors and 2.5 UV-sensitive ones (2 large photoreceptors at the opposite walls of each ommatidium + 1 small photoreceptor underneath at the base). In *Drosophila*, the number of light-activated photoreceptors and their combined contraction strength set the DPP microsaccade amplitude[5]. Thus, it seems probable that similar additive photomechanics would govern the Honeybee DPP microsaccades too. Also expectedly, the microsaccades showed synchronised lateral and axial components, comparable to *Drosophila* (**Fig. 5d-l**).

Nevertheless, whilst indicative of their phototransduction origin, these recorded dynamics were somewhat slower than expected[5], with the photomechanical rhabdom movements reaching their maxima in about 80-150 ms (**Fig. 7c**), akin to *Drosophila* (**Fig. 5**). The probable explanation for this speed range is the prolonged dark-adaptation, short 100 ms flashes (note, *Drosophila* was tested with 200 ms flashes, **Fig. 5a**) and 10 s inter-flash-intervals used in these experiments. After all, dark-adaption is well-known to decelerate honeybee phototransduction dramatically. Interestingly, however, the DPP microsaccades showed a consistent photomechanical transient (a nudge) ~30-50 ms from the light onset (black arrows). Such a nudge could, for example, signal a fast and large axial (inward) component, which DPP imaging with suboptimal resolution (of a relatively low signal-to-noise ratio) cannot resolve. Moreover, the microsaccade dynamics varied greatly from trial to trial, even more so than in *Drosophila* (cf. **Fig. 5k**), suggesting that they could be modulated or influenced by intrinsic processes, such as intraocular muscles extruding a force on the ommatidial structures.

To test whether the photoreceptor microsaccade variability (**Fig. 7c**) could, in part, reflect spontaneous intraocular muscle activity drifting the entire retina (and thus potentially inflicting variable tension to the rhabdoms[5]), we next monitored honeybee DPP continuously in darkness (**Fig. 7d**). These long-term recordings lasted up to 16 minutes. The recordings showed slow wave-like lateral retina movements, occurring about 2-3-times in a minute, and gradual axial creep, almost certainly[5] pulling the observed rhabdoms inwards (DPP darkening). These spontaneous, presumably muscle-activity-induced, components (**Fig. 7d**) differed clearly from the ultrafast photomechanical photoreceptor microsaccades (**Fig. 7c**). They were largely unsynchronised in time, and most crucially, showed 10-100-times slower dynamics, broadly comparable to our earlier findings of intraocular muscle activity in *Drosophila*[4, 5].

Discussion

We recorded photoreceptor microsaccades across the wild-type *Drosophila*, *spam* mutant and

honeybee compound eyes using a novel infrared GHS-DPP microscopy and analysed their active

sampling kinematics. Remarkably, we found the *spam* mutants and honeybee generating ultrafast

light-induced microsaccades akin to the wild-type *Drosophila*. Furthermore, in *spam*, the lateral

microsaccade movements oriented locally, forming mirror-symmetric left and right eye sampling maps,

largely similar to the wild-type flies. These results demonstrate that photoreceptor microsaccades are

not limited to the open rhabdom eye design but also occur in fused rhabdom eyes. Most insects,

including honeybees, possess fused rhabdom eyes[23, 24], in which phototransduction reagents,

including PIP$_2$, are thought to function alike in *Drosophila*[32, 35]. Therefore, it seems probable that all

insect compound eye photoreceptors would generate active sampling.

Photomechanical photoreceptor microsaccade dynamics

Photomechanical photoreceptor microsaccades are not reflex-like uniform contractions[4, 5]. Instead, at

each moment, they actively and continuously auto-regulate photon sampling dynamics by moving and

narrowing the photoreceptors' receptive fields in respect to environmental light contrast changes to

maximise information capture[4]. These dynamics rapidly adapt to immediate light history and are

different at dim and bright conditions and to positive (light increments) and negative contrasts (light

decrements)[4, 5]. So, from the viewpoint of sampling theory, photoreceptor microsaccades are not

passive but constitute a form of ultrafast *morphodynamic* active sampling[4, 5]. However, in this study,

we only examined photoreceptor microsaccades in one stimulus condition; to bright light flashes after

prolonged dark adaptation.

The spam and honeybee DPP microsaccade displacements were generally smaller than the wild-type

Drosophila's. This finding is consistent with the inter-rhabdomeric coupling hypothesis[5]. The fused

rhabdom rhabdomeres embrace each other and rigidify, and therefore, during microsaccades, they

would have less flexibility to move sideways than the open wild-type *Drosophila* rhabdomeres.

Conversely, in the much larger honeybee eye, the rhabdoms are further away from the ommatidium

lenses, reducing their receptive field sizes and interommatidial angles[5, 36]. The photoreceptor

microsaccades then seem scaled down in proportion to the interommatidial angle, presumably for

scanning the best image resolution (**Fig. 8**). These active sampling (or micro-scanning) strategies are

not mutually exclusive. Both structure-function relationships could be evolutionarily tuned to scale the

insect photoreceptors' active sampling dynamics to each species' unique visual needs. For example,

we would predict for fast-flying flies, such as houseflies (*Musca domestica*) and blowflies (*Calliphora*

vicina), having more ommatidia tiling their eyes more densely, that their photoreceptor microsaccades

be smaller and faster than those of slow-flying *Drosophila* of fewer less-densely-packed ommatidia.

This way, a fast-eye's photoreceptor receptive fields would sample the world in higher velocity and

resolution for higher visual information capture[37] - but these high-rate processes would make them

metabolically more expensive[37] - than those of a slow-eye. The fast eyes should also have more

frontal ommatidia with the fastest microsaccades to accentuate acuity[4] and stereoscopic range[5] than

the slow-eyes.

Fig. 8 | Fruit fly and Honeybee photoreceptor microsaccades scale with their receptive field half-width and interommatidial angle, presumably maximising retinal image acuity.

a, *Drosophila* photomechanical photoreceptor microsaccades typically shift their rhabdomeres 1-1.5 μm laterally (max < 2 μm), equating to ~3-4.5° receptive field movements in the visual space. R1-R6 photoreceptors' receptive field half-widths (Δρ, retinal pixels) are between 4.5-6°, over-completely tiling up the visual space[4, 5]. The average interommatidial angle[38] (φ) is 4.5-5°. Now consider this as image sampling by a digital camera. The spatial image information doubles when its sensor is moved, and two consecutive images are taken a 1/2-pixel apart and time-integrated for enhanced resolution. But if a pixel (receptive field) moves more, it eventually fuses with its

neighbour (if that neighbour pixel was still – i.e. not detecting light changes). Because of this complete pixel fusion, acuity would decrease as the resulting neural image would contain fewer pixels. Therefore, by limiting micro-scanning to interommatidial angle, *Drosophila* can time-integrate a neural image, which greatly surpasses its compound eyes' optical limits[4, 5].

b, Honeybee photoreceptor microsaccades shift their receptive field maximally <1°, approaching the eye's average interommatidial angle[1]. Equally, such displacement is less than their average receptive field half-width (~1.8°) in the front of the eye[34]. As this active sampling strategy is broadly comparable to that of *Drosophila*, we predict that honeybee vision surpasses its compound eyes' static pixelation limit, similar to what we have shown for *Drosophila*[4, 5].

Ultrafast microsaccades of dissociated *ex vivo Drosophila* photoreceptors show both *lateral* and *axial*

components[4], implying that underneath the ommatidial lenses, light changes make photoreceptors

bounce inwards and outwards and sideways in a complex piston motion[4, 5]. GHS-DPP microscopy

can reveal both of these components (**Fig. 5d** and **7c**), enabling us to estimate how they shape the

way photoreceptors encode visual space in neural time through modelling[4, 5]. The axial component

pulls the rhabdomere away from the ommatidium lens to collect light from a narrower angle. The lateral

component makes the resulting receptive field scan the visual space. GHS-DPP microscopy produces

2D image sequences ideal for measuring the *lateral* photoreceptor microsaccade dynamics. However,

the method detects less well their *axial* components, moving rhabdomeres to and from the ommatidial

lens. The estimation becomes less reliable the further away the rhabdom/rhabdomeres are from the

ommatidium lens. Following the laws of physics, the proportional DPP intensity change

(brightening/darkening), indicating the axial rhabdomere movement, diminishes with distance[5].

Therefore, GHS-DPP imaging can underestimate the overall microsaccade dynamics if these show

little sideways movement but have a larger (concealed) axial component. We suspect this would be

the case with the honeybee photoreceptor microsaccades. Honeybee rhabdoms are 4.5-times longer

than *Drosophila* rhabdomeres, thus having greater potential for axial contraction, but positioned about

100 μm from the ommatidium lens[39]; 5-times the distance in *Drosophila*[36].

Photoreceptor microsaccade movement direction maps

The microsaccades in the *spam* mutants were slightly more rotated than the wild-type. Nevertheless, with such minor differences, their global maps should broadly match the forward flight optic flow field, similar to what we have previously shown for the wild-type[5]. Theoretically, this correspondence between active sampling and optic flow should improve the visual resolution of the moving world[5]. We further conjectured that since the microsaccades follow the R1-R2-R3 rhabdomere orientation axis[5] (**Supplementary Video 5**), their movement directions are set during development, perhaps guided by some lowest resistance (minimum energy) anchoring[5]. Thus, the observed wild-type and *spam* microsaccade direction differences could reflect slight differences in their R1-R7/8 rhabdomere orientations. Unfortunately, in the circular-symmetric *spam* DPP, the separate R1-R7/8 rhabdomeres are not directly identifiable under infrared illumination. In the future, this hypothesis could be tested by expressing GFP in selected R1-R7/8 photoreceptors or rhabdomeres. Moreover, because the high-speed imaging in the larger and densely pigmented honeybee eyes was experimentally challenging, we also left mapping their microsaccade movement directions for future studies.

Photoreceptor microsaccades and eye-muscle-induced whole retina movements mix

Some sporadic interference in the measured microsaccades can originate from the intraocular muscles' activity, which exerts force on the retina. But since such muscle activity typically occurs over much longer time intervals[21] or during active viewing[6], being practically absent in firmly restrained flies[5] and sluggish in head-immobilised honeybees, the eye muscle induced movements had little influence on the ultrafast photomechanical microsaccades shown in this report (i.e. influencing them perhaps only through variable ommatidial tension).

However, during normal behaviours (in non-restricted, free-moving conditions), the local photoreceptor microsaccades and the global eye-muscle-induced whole retina movements must interact in active sampling. On the top of any eye-muscle-induced whole retina movements, the

photomechanical photoreceptor microsaccades will ensue, leading to complex superimposed

spatiotemporal ("super-saccadic") sampling dynamics. This sophistication arises because each retina

movement will change its photoreceptors' light input, evoking their photomechanical microsaccades.

Notably, active sampling can be even more elaborate if the eye-muscle-induced whole retina

movements were partly voluntary and depended upon the attentive state. In those circumstances, an

insect could use eye-muscle-induced retina movements together with other directional senses, such

as antennal casting[40], to get a better idea of what an encountered object might be. After all, integration

of multisensory information reduces uncertainty, increasing fitness.

Finally, we note that there are uncontrolled genetic differences between the wild-type and *spam*

phenotypes, which could potentially contribute to their observed DPP microsaccade differences.

Nevertheless, such differences would not alter the general demonstration of the active sampling of

the fused rhabdom eye photoreceptors as it occurs with somewhat comparable dynamics to the

honeybee rhabdom.

Conclusions

Goniometric high-speed deep pseudopupil (GHS-DPP) microscopy provides an innovative non-

invasive way to image photoreceptor microsaccades globally across the left and right eyes, or locally,

in great detail at specific eye locations. We explained how to use it and gave out free open-source

(GPLv3) software tools to quantify and compare active sampling in different insect eyes. Our results

demonstrate active sampling both in open and fused rhabdom eyes. Thus, the GHS-DPP microscopy

shows real potential as a powerful tool to study how the insect eyes actively sample the visual world.

Authors declare no competing interests.

Author Contributions

Conceptualization (MJ, JK, JT), Data curation (JK, NM), Formal Analysis (JK), Funding acquisition

(MJ), Investigation (JK, NM, JT, MJ), Methodology (JK, JT, MJ), Project administration (MJ),

Resources (MJ), Software (JK, JT), Supervision (MJ), Validation (JK, JT, MJ), Visualization (JK, MJ),

Writing – main paper original draft (JK), Writing – review & editing (MJ, JK, JT).

Funding

This work was supported by Jane and Aatos Erkko Foundation Fellowships (MJ and JT), The

Leverhulme Trust (RPG-2012–567: MJ), the BBSRC (BB/F012071/1, BB/D001900/1 and

BB/H013849/1: MJ), the EPSRC (EP/P006094/1: MJ), the Open Research Fund of the State Key

Laboratory of Cognitive Neuroscience and Learning (MJ), High-End Foreign Expert Grant by Chinese

Government (GDT20151100004: MJ).

Acknowledgements

We kindly thank Andrew Zelhof for the spam fly line and Gonzalo de Polavieja, Roger C Hardie, Merlin

Juusola, Gregor Belušič and the Juusola laboratory members for discussions and comments.

1   Data Availability Statement

The software code and datasets for this study can be found in the https://github.com/JuusolaLab/GHS-

DPP_paper

1.   Land, M.F. Visual acuity in insects. *Annu Rev Entomol* **42**, 147-177 (1997).

2.   Laughlin, S.B. The role of sensory adaptation in the retina. *J Exp Biol* **146**, 39-62 (1989).

3.   Barlow, H.B. The size of ommatidia in apposition eyes. *J Exp Biol* **29**, 667-674 (1952).

4.   Juusola, M. et al. Microsaccadic sampling of moving image information provides *Drosophila*
hyperacute vision. *Elife* **6** (2017).

5.   Kemppainen, J. et al. Binocular Mirror-Symmetric Microsaccadic Sampling Enables *Drosophila*
Hyperacute 3D-Vision. *bioRxiv* (2021).

6.   Franceschini, N., Chagneux, R., Kirschfeld, K. & Mucke, A. in Gottingen Neurobiology Report: Synapse
- Transmission Modulation. (eds. N. Elsner & H. Penzlin) 1 (Georg Thieme Verlag, Stuttgard, New
York; 1991).

7.   Hengstenberg, R. Eye muscle system of housefly *Musca-Domestica* .1. Analysis of clock spikes and
their source. *Kybernetik* **9**, 56-77 (1971).

8.   Hardie, R.C. & Franze, K. Photomechanical responses in *Drosophila* photoreceptors. *Science* **338**, 260-
263 (2012).

9.   Hardie, R.C. & Juusola, M. Phototransduction in *Drosophila*. *Curr Opin Neurobiol* **34**, 37-45 (2015).

10. Franceschini, N. & Kirschfeld, K. Optical study *in vivo* of photoreceptor elements in compound eye of
Drosophila. *Kybernetik* **8**, 1-13 (1971).

11. Franceschini, N. & Kirschfeld, K. Phenomena of pseudopupil in compound eye of *Drosophila*.
Kybernetik **9**, 159-182 (1971).

12. Franceschini, N. in Information processing in the visual systems of *Anthropods* (ed. R. Wehner) 75-82
(Springer-Verlag, Berlin, Heidelberg, New York; 1972).

13. Douglass, J.K. & Wehling, M.F. Rapid mapping of compound eye visual sampling parameters with
FACETS, a highly automated wide-field goniometer. *J Comp Physiol A* **202**, 839-851 (2016).

14. Muñoz Arias, M., Douglass, J.K., Wehling, M.F. & Stavenga, D.G. Automated charting of the visual
space of insect compound eyes. *arxiv* (2021).

15. Kurada, P. & O'Tousa, J.E. Retinal degeneration caused by dominant rhodopsin mutations in
Drosophila. *Neuron* **14**, 571-579 (1995).

16. Zelhof, A.C., Koundakjian, E., Scully, A.L., Hardy, R.W. & Pounds, L. Mutation of the photoreceptor
specific homeodomain gene Pph13 results in defects in phototransduction and rhabdomere
morphogenesis. *Development* **130**, 4383-4392 (2003).

17. Huang, Y., Xie, J. & Wang, T. A fluorescence-based genetic screen to study retinal degeneration in
Drosophila. *Plos One* **10**, e0144925 (2015).

18. Seidl, R. & Kaiser, W. Visual field size, binocular domain and the ommatidial array of the compound
eyes in worker honey bees. *J Comp Physiol A* **143**, 17-26 (1981).

19. Taylor, G.J. et al. Bumblebee visual allometry results in locally improved resolution and globally
improved sensitivity. *Elife* **8**, e40613 (2019).

20. Aptekar, J.W. et al. Method and software for using m-sequences to characterize parallel components
of higher-order visual tracking behavior in *Drosophila*. *Front Neural Circuits* **8**, 130 (2014).

21. Hengstenberg, R. in Information Processing in the Visual Systems of Anthropods 93-96 (Springer,
1972).

22. Wardill, T.J. et al. Multiple spectral inputs improve motion discrimination in the *Drosophila* visual
system. *Science* **336**, 925-931 (2012).

23. Zelhof, A.C., Hardy, R.W., Becker, A. & Zuker, C.S. Transforming the architecture of compound eyes.
Nature **443**, 696-699 (2006).

24. Osorio, D. Spam and the evolution of the fly's eye. *Bioessays* **29**, 111-115 (2007).

25. Ready, D.F., Hanson, T.E. & Benzer, S. Development of the *Drosophila* retina, a neurocrystalline
lattice. *Dev Biol* **53**, 217-240 (1976).

26. Hardie, R.C. in Progress in Sensory Physiology. (eds. H. Autrum et al.) (Springer, Berlin; 1985).

27. Tang, S. & Juusola, M. Intrinsic activity in the fly brain gates visual information during behavioral
choices. *Plos One* **5** (2010).

28.   Zheng, L. et al. Feedback network controls photoreceptor output at the layer of first visual synapses
in *Drosophila*. *J Gen Physiol* **127**, 495-510 (2006).

29.   van Swinderen, B. Attention-like processes in *Drosophila* require short-term memory genes. *Science*
315, 1590-1593 (2007).

30.   Fujiwara, T., Cruz, T.L., Bohnslav, J.P. & Chiappe, M.E. A faithful internal representation of walking
movements in the *Drosophila* visual system. *Nat Neurosci* **20**, 72-81 (2017).

31.   Juusola, M. & Song, Z. How a fly photoreceptor samples light information in time. *J Physiol-London*
595, 5427-5437 (2017).

32.   Fain, G.L., Hardie, R. & Laughlin, S.B. Phototransduction and the evolution of photoreceptors. *Curr*
Biol **20**, R114-124 (2010).

33.   Juusola, M., Song, Z. & Hardie, R.C. in Encyclopedia of Computational Neuroscience. (eds. D. Jaeger &
R. Jung) 2359-2376 (Springer New York, New York, NY; 2015).

34.   Rigosi, E., Wiederman, S.D. & O'Carroll, D.C. Visual acuity of the honey bee retina and the limits for
feature detection. *Sci Rep-Uk* **7** (2017).

35.   Yau, K.W. & Hardie, R.C. Phototransduction motifs and variations. *Cell* **139**, 246-264 (2009).

36.   Stavenga, D.G. Angular and spectral sensitivity of fly photoreceptors. II. Dependence on facet lens F-
number and rhabdomere type in *Drosophila*. *J Comp Physiol A* **189**, 189-202 (2003).

37.   Song, Z. & Juusola, M. Refractory sampling links efficiency and costs of sensory encoding to stimulus
statistics. *J Neurosci* **34**, 7216-7237 (2014).

38.   Gonzalez-Bellido, P.T., Wardill, T.J. & Juusola, M. Compound eyes and retinal information processing
in miniature *dipteran* species match their specific ecological demands. *P Natl Acad Sci USA* **108**,
4224-4229 (2011).

39.   Varela, F.G. & Wiitanen, W. The optics of the compound eye of the honeybee (Apis mellifera). *J Gen*
Physiol **55**, 336-358 (1970).

40.   Baker, K.L. et al. Algorithms for Olfactory Search across Species. *J Neurosci* **38**, 9383-9389 (2018).

41.   Juusola, M., Dau, A., Zheng, L. & Rien, D.N. Electrophysiological method for recording intracellular
voltage responses of *Drosophila* photoreceptors and interneurons to light stimuli *in vivo*. *Jove-J Vis*
Exp (2016).

42.   Juusola, M. & Hardie, R.C. Light adaptation in *Drosophila* photoreceptors: I. Response dynamics and
signaling efficiency at 25 degrees C. *J Gen Physiol* **117**, 3-25 (2001).

43.   Bradski, G. & Kaehler, A. OpenCV. *Dr. Dobb's J Software Tools* **3** (2000).

44.   Virtanen, P. et al. SciPy 1.0: fundamental algorithms for scientific computing in Python. *Nat Methods*
17, 261-272 (2020).

Methods

We describe the GHS-DPP imaging system hardware configuration, how to prepare the flies,

experimental protocols, and data analysis tools and principles. All software - from recording to analysis

- are available under a free and open (GPLv3) software license in a GitHub repository:

https://github.com/JuusolaLab/GHS-DPP_paper.

Supplementary Videos 1-2 show how the GHS-DPP microscopy was put together and works.

Supplementary Video 3 shows how ommatidial rhabdomere orientation across the *Drosophila* eyes

matches the optic flow field during forward translation.

Supplementary Video 4 shows the relationship between the number of locally activated

photoreceptor microsaccades, the angular size of the light stimulus and the number of ommatidial

photoreceptors pooled in the deep pseudopupil imaging (given the microscope system's numerical

aperture, NA).

Supplementary Videos 2 and **5** show how active sampling - by photomechanical photoreceptor

microsaccades - looks like in slow motion and how the microsaccades were analysed across the

compound eyes.

Supplementary Video 6 shows how photoreceptor microsaccade movement directions match the

ommatidial rhabdomere orientation across the *Drosophila* eyes.

Experimental setup

The GHS-DPP imaging system's two primary components are a rotation stage system and a

stereomicroscope (**Fig. 2a**). The rotation system allows precise control over the fly eyes' yaw and

pitch using two perpendicularly mounted rotation stages (Thorlabs PR01/M, USA), mounted

horizontally and vertically. An additional small 3-axis micromanipulator, connected to the vertical

rotation stage, was used to control the fly's initial position. Crucially, the vertical rotation stage further

rested on a 2-axis micromanipulator so that the intersection point of the two rotation stage axes could

be centred at the microscope's field of view. The rotation stage positions were acquired digitally, using

two 1024-step rotation encoders (YUMO E6B2-CW23E) and an Arduino board (Arduino Uno, Italy)

running a custom C++ program (**Fig. 2b**). In addition, the rotation stages were fitted with stepper

motors for fully automated experiments (**Fig. 2a**). Still, in this study, the experiments were performed

manually, accurately focussing on DPP at all tested eye locations. **Supplementary Video 1** shows

how the GHS-DPP microscope system was put together. Note that the stereo microscope's horizontal mounting is not critical but resulted from our earlier design choices. By having an upright rotation stage, this configuration works well for the binocular compound eye measurements.

The stereomicroscope was mounted sideways to function with the rotation stages. A high-intensity ultraviolet LED (UV-OptoLED, Cairn Research, UK) was inserted in the microscope's ocular slot, enabling direct light stimulation of the observed DPP rhabdomeres. This 365 nm UV-LED and the two infrared 850 nm LEDs (IR-OptoLED, Cairn Research, UK), which provided antidromic non-stimulating illumination of the DPP rhabdomeres (without activating their phototransduction) through the fly head capsule, were connected to their separate driver units (Dual OptoLED Power Supply, Cairn Research, UK). The LEDs were controlled over the BNC interface using a computer connected data acquisition system (PCI-6221 with BNC-2090A and PCI-6733 with BNC-100, National Instruments, USA) (**Fig. 2c**). The two infrared LEDs were mounted apart from each other at different angles to prevent the pipette tip or the fly body from blocking the illumination, which would otherwise frequently happen with a single point source. Both the infrared LEDs were in bespoke holders, having a convex lens with adjustable lens-to-LED distance enabled beam focusing. The second unit mounted to the microscope was the high-speed optical camera (Orca Flash 4.0 C13440, Hamamatsu, Japan), which also sent a trigger signal over the BNC interface to the data acquisition system to time the stimulus delivery precisely. We typically acquired 2,048 x 2,048 pixel full-frame images at 100 fps, and occasionally - by cropping the sensor to 2,000 x 200 pixel - collected images at 1,000 fps. The camera had a transparent infrared and opaque UV filter on its pathway, ideally stopping the UV stimulation light from reaching and polluting the image sensor (**Fig. 2d**). Besides these filters, the microscope was configured with a beam-splitter (SZX-BS, Olympus, Japan), a photo adapter piece (SZ-PHA, Olympus, Japan) and a magnification changer (U-TVCAC, Olympus, Japan).

The setup was mounted on a vibration isolation table (air table), which uncoupled any building vibrations that could affect the motion analysis results. The rotation stage system was connected to

the table by magnetic clamps (Magnetic Stand, World Precision Instruments), whilst the microscope

system - attached to a thick steel pole of a steel base plate - was heavy enough to ensure its fixed

position. The whole setup was enclosed inside a black-painted metal cubicle, with its only open side

supporting black curtains for performing the experiments in controlled, dark conditions. Accordingly,

the Arduino board's few surface-mounted LEDs were covered with electrical insulation tape to

minimise any light noise.

Flies and preparation

We used wild-type *Drosophila* flies (Berlin) and a fused rhabdom *spam* null-mutant line (w;

spam1/spam1 Frt; sqh-GFP/Tm6B, a gift from Andrew Zelhof) in the experiments. The flies were

maintained in an incubator at 25 ℃, under a 12:12h light-dark cycle. Only healthy 3- to 20-days-old

male and female flies, climbing up the vial, were selected for the experiments. We avoided using very

young flies (< 3 days) because their soft heads could bulge during the imaging, presumably due to

spontaneous eye muscle activity. The flies were prepared for the experiments using the plastic pipette

tip immobilisation technique[41] (**Supplementary Fig.1**). Previously, we tested a copper hook tethering

technique[5], allowing simultaneous behavioural experiments (**Supplementary Fig.1a**). But because

both these immobilisations gave broadly similar results[5], the more laborious and time-consuming

tethering was not used here.

Supplementary Fig.1 | The pipette tip preparation technique. **a**, Flies can be either tethered on a small copper hook with UV-cured glue or immobilised on a plastic pipette tip with beeswax. **b**, With the help of a funnel piece, a fly was captured in a 1,000 µl plastic pipette tip. **c**, Air puffs generated by a hand-held syringe pushed the fly towards the pipette tip's small end. **d**, The fly was immobilised from its dorsal thorax. **e**, the head was secured near the proboscis using melted beeswax. **f**, The pipette excess was cut off with a razor blade, and then the preparation was inserted into the GHS-DPP imaging system.

In the pipette-tip-fixation method, a 1,000 µl plastic pipette tip was first linked to a funnel piece, and the funnel was connected to a vial full of flies. One fly at a time was lured into the pipette tip through the funnel driven by *Drosophila*'s innate geotaxis behaviour (**Supplementary Fig.1b**). This way, there was no need to immobilise the flies by CO_2 or ice-cooling that could potentially affect the microsaccades. Next, the pipette tip was viewed under a stereo preparation microscope (Olympus SZX-9, Japan). At the same time, the fly was gently pushed towards the tip opening by puffing air from a hand-held syringe (**Supplementary Fig.1c**). When needed, the pipette tip's opening was adjusted using a razor blade to ensure that the fly head passed through without any deformation, minimising structural damage to the eyes. The fly was air-puffed until its head and upper thorax protruded the tip end. When the fly was ideally positioned, it was quickly immobilised by applying melted beeswax on its thorax (**Supplementary Fig.1d**) and ventrally to its head and proboscis (**Supplementary Fig.1e**). In this correct position, the pipette would not shadow the dual-IR illumination during the experiments.

Additionally, the proboscis could have been pulled out with forceps and waxed on the pipette's outside, or the antennae waxed. However, these manoeuvres, which we often use in preparing *Drosophila* for

intracellular electrophysiology[41, 42], were omitted because they made no difference in the observed

microsaccades. Furthermore, these procedures would have prolonged the preparation making,

potentially increasing structural damage. Conversely, some more wax was routinely applied on the

dorsal side of the head, under the ocelli, to secure the head position further. Finally, the pipette was

cut from its large end using a razor blade (**Supplementary Fig.1f**), and the preparation was carefully

placed in the setup (**Supplementary Video 2**).

Honeybee

Honeybees (*Apis   mellifera*) were kept indoors   in   a hive   that provided the   bees an outside

(nature) access   through   a plastic portal. Worker bees   were   captured   from   the   portal and

prepared using the pipette tip technique similar to the *Drosophila* with few adjustments. First, the

pipette tip's small opening was cut larger to fit the bee head. Second, the bee was ice-cooled before

the waxing to prevent it from escaping. During the cooling, the bee inside a plastic vial was placed in

ice until its leg movements temporarily halted. Although the bees quickly recovered, as judged by their

antennal movements, the ice-cooling may have generally affected their physiology. Finally, a cuticle

section (between the antennae and ocelli and the dorsal part of the eyes) was removed by a razor

blade after the waxing. This microsurgery greatly increased the DPP image's brightness, allowing us

to image photoreceptor microsaccades in high frame rates (100Hz and 500 Hz).

Data acquisition and software

To enable non-specialist users to operate the GHS-DPP microsaccade imaging experiments, we

created a free and open (GPLv3) recording software called *Gonio Imsoft. Gonio Imsoft* interfaced with

the open-source microscopy software *MicroManager* to control the high-speed camera. In addition, it

used   the   *NI-DAQmx*   module   (controlling   the   data   acquisition)   and   the   *PySerial*   module

(communicating with the Arduino microcontroller), reading out the rotary encoders. *Gonio Imsoft* ran

on a Windows 10 platform.

We used the Olympus DF-PLAPO 1X objective for typical experiments with the microscope's

continuous zoom maxed out and the additional magnification changer set to the 2x-position. In this

configuration, a small-to-average-size fly head falls entirely within the field of view. The microscope's

light path selector was set to its mid-state, resulting in an 80%/20% light intensity split between the

camera and the eyepieces, respectively. The two infrared illumination LEDs were focused on the back

of the fly head. Using the 2-axis micrometre, mounted on the horizontal rotation stage, the vertical

rotation stage was centred so that the fly head, when brought to the field-of-view by the 3-axis

micromanipulator (mounted on the vertical rotation stage), remained in the centre of the field of view

in all possible horizontal and vertical rotation combinations (**Supplementary Video 2**). However, the

microscope's focus remained only approximately correct since the fly head is not perfectly round and

hence refocusing was needed to maintain a sharp DPP image during the experiments.

The microscope was first focused so that the ommatidial lenses appeared crisp and clear. Then, the

focus was brought deeper into the eye until the DPP became clear and visible. In this ommatidial axes'

converge point, images formed by individual ommatidia superimpose, creating a magnified, virtual

image of the rhabdomere tips[11] (**Supplementary Video 4**). Using our computer graphics (CG)

simulations on wild-type and *spam* eyes, we further confirmed that the retinal patterning (**Fig. 1a** and

b) indeed resulted in a one-spot DDP image in the *spam* (**Fig. 1c**).

Fig. 9 | The amount of DPP forming ommatidia depends on the microscope numerical aperture. a, A spherical compound eye illustrated with seven centred hexagonal rings. **b**, The nth centred hexagonal number quantifies the number of ommatidia, N, within the nth ring. **c**, Light rays emerging from the retina midpoints can be collected by the microscope only if their incident angle is smaller than the microscope's half-collection angle, Θ. **d**, Some light from the uncollected ommatidia may reach the microscope if it originates off-centre (blue). **e**, light originating 1 μm off-centre is approximately 3° diverted from the optic axis. **f**, The r parameters were estimated from the wild-type and *spam* DPPs, and one wild-type DPP rhabdomere. The scale bar corresponds to 50 μm in the DPP images and 5 μm physically in the retina. **g**, The amount of DPP forming ommatidia (here c=0) rises sharply with the increasing microscope numerical aperture, NA.

The DPP is purely an optical phenomenon, informing us more about how the observed rhabdomere
summation image is affected by the used microscope system[11] than how the fly compound eye
converges visual inputs from the world[28]. Therefore, the DPP cannot tell us too much about the eyes'
neural superposition design, in which R1-R6 photoreceptor outputs from six neighbouring ommatidia,
sampling light over the same small visual area, are synaptically pooled. Nevertheless, it is helpful to
estimate the number of ommatidia forming in the optical DPP image as this improves our
understanding of any blurring and irregularities in it. On a hexagonal tiling, the number of hexagons
within an nth concentric hexagonal circle (**Fig. 9a** and **b**) can be calculated using the formula

$$648 \quad \begin{cases} N = 3n(n-1)+1 & \text{if} \quad n \geq 1 \\ N = n & \text{if} \quad n < 1 \\ N = 0 & \text{if} \quad n \leq 0 \end{cases} \tag{1}$$

650 The values of N are known as the centred hexagonal numbers. The piecewise formulation also

651 accounts for the $n < 1$ cases, usually left undefined. By finding the right n, we use it to denote the

652 number of ommatidia participating in the DPP image formation. From the eye-microscope geometry

653 (**Fig. 9c** and **d**), we can estimate the value of n, after which the illumination light rays no longer can

654 enter the microscope's entrance pupil, and write

$$656 \quad n = \frac{\theta + 2(0.5-c)\alpha}{\Delta\varphi} + 1 \tag{2}$$

θ is the half-angle subtended by the microscope lens to the eye, $\Delta\varphi$ is the interommatidial angle, c is

the minimum counted contribution term and α is the mean deviation angle between the outermost

rhabdomere receptive axes and the ommatidial axis. If we were to estimate the number of ommatidia

that project all their rhabdomeric light to the microscope, $c = 1$, and conversely, if we were interested

in ommatidia casting *any* rhabdomeric light to the microscope, $c = 0$. For counting ommatidia with

other contributions, the correct c depends on the microscope NA. But for a 50% minimum contribution

estimate, we can use the former estimates. Then next, ideally, if the rhabdomeres formed a single

point on the ommatidial axis (and the eye was perfectly spherical), α would be zero. And, if the

rhabdomeres were organised into a hexagonal shape so that the neighbouring rhabdomeres'

receptive fields (RFs) perfectly overlap at infinity, α would nearly equal the interommatidial angle $\Delta\varphi$.

By replacing the half-angle with the numerical aperture (NA) and considering that in *Drosophila* 1 µm

displacement results in a 3° angular change[5], we can write

$$672 \quad n = \frac{\sin^{-1}\left(\frac{NA}{m}\right) + 2(0.5-c)\left(3°/\text{µm } r\right)}{\Delta\varphi} + 1 \tag{3}$$

NA is the numerical aperture of the used microscope, m is the refractive index of microscope objective

immersion medium (m = 1 for air), and r is the radius of a circle in the ommatidial retinal plane that

representatively contains all the rhabdomeres. The numerical aperture in our microscope system was

0.11, and the *Drosophila* interommatidial angle is approximately 5° [36, 38]. Since the DPP image is a

~10x magnified image of the rhabdomere tips, the r parameters for the wild-type and *spam* can be

estimated directly from the DPP still images as 2.8 µm and 1.5 µm (**Fig. 9f**). Inserting these values in

(3) and then calculating (1), we approximate that in our microscope system, 18 ommatidia would

contribute at least 50% to the deep pseudopupil image formation in the wild-type and 12 ommatidia in

the *spam* eye. For one rhabdomere in the wild-type DPP pattern, we can similarly estimate that its r

is 0.78 µm (**Fig. 9f**), suggesting that 10 ommatidia form it. Notice, however, that because of the

structural asymmetricities, these ommatidium counts are likely slightly overestimated. For example,

the *Drosophila* ommatidial R1-R6 rhabdomeres are not hexagonally arranged around the lens centre.

Instead, their sizes and distances vary[4], forming an asymmetric (slanted) trapezoid arrangement (**Fig.**

3c), and the eye is not perfectly spherical. Moreover, infrared illumination, by passing through the

ommatidial screening pigments that block non-axial green and UV transmission, could potentially

merge more ommatidia into the DPP image than visible light. Nevertheless, since these upper-bound

estimates vary with the used microscope systems (**Fig. 9g**), we calculated them for a range of air-

objective NAs (**Supplementary Table 1**).

Supplementary Table 1. The estimated number of ommatidia forming the DPP image.

Microscope NA (in air)	N (wild-type ommatidia) (r = 2.8 µm)			N (*Spam* ommatidia) (r = 1.5 µm)			One wild-type rhabdomere (r = 0.78 µm)		
	contributing at least								
	100%	50%	>0%	100%	50%	0%	100%	50%	>0%

0.001	0	7.3	15	0.11	3.2	6.2	0.54	1.8	3.1
0.01	0	8.0	16	0.21	3.7	7.1	0.65	2.2	3.8
0.05	0	11	23	0.67	6.3	12	1.3	4.4	7.4
0.1	0.47	17	33	1.9	11	20	4.4	9.1	14
0.2	4.1	32	61	11	26	41	17	25	32
0.4	38	90	143	56	84	112	68	83	97
0.6	115	195	274	146	189	231	165	187	209
0.8	268	380	492	314	374	434	341	372	403
0.95	521	671	821	585	665	745	622	663	705

In this study, we performed two kinds of experiments: *local recordings* at a fixed position on the left

and right eye and *global recordings* across the eyes. In the local recordings, we imaged one location

on the left eye (+28° horizontal and -37° vertical) and another on the right eye (-28° horizontal and -

37° vertical) because, at these locations, the upper infrared LED illuminated the eyes ideally, forming

the crispest DPP. The stimulus UV-LED was flashed for 200 ms while simultaneously the high-speed

camera acquired images 100 frames per second, yielding 20 image frames per flash. As a standard

procedure, this recording process was repeated 25 times to obtain the mean photoreceptor

microsaccade estimates and inspect their variability in an individual fly. Between the repeats, we

initially used 10 s inter-stimulus-interval (ISI) for both the wild-type and spam flies. But ISI was later

shortened ISI to 2 s, as there was no significant reduction in the response amplitude. All images were

saved as 16-bit unsigned-integer, grayscale TIFF images.

The global imaging procedure was similar, but it was performed only once at each location. Instead,

after each flash, the fly was rotated in 10° steps from -40° to +50° (limited by the vertical rotation stage,

covering the microscope objective or the two illumination LEDs) using the horizontal rotation stage.

After completing the horizontal "line scan", the vertical rotation stage was advanced in a 10° step,

covering a range from -110° to +110°. By rotating the fly and imaging in between, we scanned

approximately 200 distinct locations on the left and right eyes. Overall, it took about 10 s to reorient

the fly, refocus at the deep pseudopupil and start the image acquisition protocol again.

Data analysis

To quantify the 3-dimensional rhabdomere movement fields, we created a free and open (GPLv3)

data analysis software called *Gonio Analysis*. It allows drawing regions of interest (ROIs) around the

deep pseudopupil, performing motion analysis, and finally, translating the motion results from the

camera image coordinates to the fly's 3D frame-of-reference, using the digitally read rotation stage

values. Here, we shortly describe its data analysis principles.

Rectangular ROIs were drawn by hand for the first frame of each location only by selecting the deep

pseudopupil. Next, these pseudopupil images were used as template images for cross-correlation

based motion analysis (**Fig. 10a**). We used the computer vision library *OpenCV* [43] and its

matchTemplate routine for the following 2D cross-correlation

727                                $$R(x,y) = \frac{\sum_{x',y'}(T'(x',y')\,I'(x+x',y+y'))}{\sqrt{\sum_{x',y'}T'(x',y')^2\,\sum_{x',y'}I'(x+x',y+y')^2}}$$ (4)

728                                $$T'(x',y') = T(x',y') - \frac{1}{w*h} * \sum_{x',y'}T(x'',y'')$$ (5)

729                $$I'(x+x',y+y') = I(x+x',y+y') - \frac{1}{(w*h)} * \sum_{x'',y''}I(x+x'',y+y'')$$ (6)

Fig. 10 | Cross-correlation based motion analysis by template matching. **a**, The first frame of the DPP image, manually cropped, was used as the template image. By 2D cross-correlation analysis, the DPP image locations were searched among the source images. **b**, A cross-correlation result image *R* can be presented as a height-map, giving a better view of the cross-correlation landscape. **c**, Two superimposed R height-maps reveal a small shift Δd in the location of the primary maxima; This is the DDP displacement. **d**, The motion analysis results were transformed from the camera image coordinates to the fly coordinate system using the rotation stage values.

R is the 2-dimensional cross-correlation image, and $R(x,y)$ is its value at the pixel (x,y). x', x" and y', y"

are summation indices within ranges [0, 1, 2, …, *w*-1] and [0, 1, 2, ..., *h*-1], *w* and *h* are the width and

height of the template image. *I* is the source image, and *T* is the template image.

In the cross-correlation results of the images *R* produced by the template matching (**Fig. 10b**), the

higher values were signed for the higher similarity between the template and source images at each

location. Therefore, using two source images acquired at different times, the template displacement

between these images can be calculated by comparing their resulting cross-correlation image's peak

values (**Fig. 10c**). Furthermore, we visually confirmed some motion analysis results by creating videos

in which the rectangular ROI box was moved according to the motion analysis results, readily following

the moving DPP. On the other hand, we note that computing the complete cross-correlation with

uncropped source images (cf. **Fig. 1**) is inefficient and can lead to a false match. Therefore, we instead

performed the cross-correlation only in the near vicinity (30 pixels) of the DPP cropping without

truncating the responses.

In the template matching, the motion analysis results were given in the camera image coordinates. To

translate them from the camera system to the 3D space in the fly's frame of reference (**Fig. 10d**), we

used the digitally read vertical v and horizontal h rotation stage values on the following set of equations

751
$$\begin{cases} y = \cos(h)\cos(v) \\ z = y\tan(v) \\ |x| = \sqrt{1 - y^2 - z^2} \end{cases} \tag{7}$$

753 to calculate the microscope's (x, y, z) location. Using the same equation set, in short, $P(h, v) \rightarrow (x, y, z)$,

754 we then calculated the camera x unit vector as the vector from $P(h, v)$ to a slightly displaced

755 point $P(h + \Delta h, v)$

757
$$\hat{i}_{cam}(h, v) = \overrightarrow{P(h, v)\, P(h + \Delta h, v)} \tag{8}$$

759 where Δh is ideally as small as possible but large enough not to cause errors because of limited

760 floating-point precision. Since the y-camera-unit only depends on the vertical rotation and its x-

761 component in the fly coordinate system is conveniently zero (i.e. the y-unit-vector is always

762 perpendicular to the great circle arc that the microscope travels along from the fly's point of view), it is

763 simply

765
$$\hat{j}_{cam}(v) = -\sin(v)\hat{j} + \cos(v)\hat{k} \tag{9}$$

767 Finally, the (x, y, z) movement vectors can be then calculated using camera unit vectors as

769
$$v(m_x, m_y, h, v) = m_x \hat{\imath}_{cam} + m_y \hat{\jmath}_{cam} \tag{10}$$

where m_x and m_y are the camera image x and y movement values produced by the cross-correlation.

We averaged the results over many flies using simple N-nearest neighbour interpolation for the

microsaccade vector maps acquired in the global recordings. For each interpolation point, from each

of the N imaged flies, the nearest 3D vector was selected, but only if the angular distance of the 3D

vector was not larger than 2-times the angular interpolation step of 5°. And, these equal or less than

N vectors were averaged together only if there were N/2 of them or more. The difference in the wild-

type and *spam* vector maps were calculated point-wise as

779
$$e(v_{wtb}, v_{spam}) = \frac{1}{\pi} \cos^{-1}\left(\frac{v_{wtb} \cdot v_{spam}}{\|v_{wtb}\|\|v_{spam}\|}\right) \tag{11}$$

where v_{wtb} and v_{spam} are the microsaccade vectors and the operators $\cdot$ and $\|\ \|$ denote the inner

product and the vector norm, respectively. Finally, for rotation direction analysis, we rotated the

vectors on the x-axis and calculated whether the spam vectors were rotated clockwise or

counterclockwise compared to the wild-type and then used this result to sign the error in (11).

In the local recordings, we focused solely on the directionless microsaccade magnitude, calculated

using the Pythagorean theorem from the camera coordinate movement values. From these animal-

specific mean magnitude traces, we calculated the microsaccade amplitude, speed, logistic growth

factor and half-rise time quantifications. The probability graphs were calculated as 1D-histograms at

each time point, with 20 stitched together to cover the whole imaging period. The total displacement

values were quantified from the magnitude data by simply taking the mean of the last 7 data points,

the last 70 ms of the imaging. The maximum speed was calculated from the highest value between

frames displacement. For the logistic growth factor and the half-rise time, we fitted the data with the

sigmoidal logistic function

$$f(t) = \frac{L}{1+e^{-k(t-t_0)}}$$ (12)

where L is the maximum value corresponding to the total rhabdomeric displacement, k is the logistic

growth factor that in our case characterises the microsaccade activation phase time-duration, and t_0

is the half-rise time.

Statistics and Reproducibility

The presented results are readily reproducible. Every healthy *Drosophila* with functional vision will

show them. The figures and figure legends give the sample sizes (how many flies were used) and the

number of recorded responses (to repeated stimulation). In contrast, because honeybee has black

"armoured" head cuticle, which makes preparing them significantly more complicated, we show only

exemplary (repeatable and reproducible) results from one bee. These results reveal photoreceptor

microsaccades occurring also in the bee eye, with comparable ultra-fast dynamics to *Drosophila*, and

demonstrate that GHS-DPP microscopy can capture them too. All the quantified local recording

parameters (displacement, speed, growth factor and rise-time) appeared reasonably normally

distributed. However, since the *Drosophila* wild-type sample size was 3-times larger than the *spam*

one, we used Welch's t-test[44] that performs better with uneven samples for all comparisons.