RESEARCH

Context dependency of nucleotide probabilities and variants in human DNA

Yuhu Liang[1,2], Christian Grønbæk[2], Piero Fariselli[3] and Anders Krogh[1,2]*

*Correspondence: akrogh@di.ku.dk
[1]Department of Computer Science, University of Copenhagen, Copenhagen, Denmark
Full list of author information is available at the end of the article

Abstract

Background: Genomic DNA has been shaped by mutational processes through evolution. The cellular machinery for error correction and repair has left its marks in the nucleotide composition along with structural and functional constraints. Therefore, the probability of observing a base in a certain position in the human genome is highly context-dependent. Mutations are also known to depend on the genomic context, but in previous work, the nucleotide distribution and the mutability have not been combined.

Results: Here we use a context-dependent nucleotide model as the basis for a mutability model for the human genome. We first investigate simple models of nucleotides conditioned on sequence context and develop a bidirectional Markov model that depends on up to 14 nucleotides to each side. We show how the genome predictability varies across different types of genomic regions. Surprisingly, this model can predict a base from its context with an average of more than 50% accuracy. Inspired by DNA substitution models, we develop a model of mutability that estimates a mutation matrix (called the alpha matrix) on top of the nucleotide distribution. The advantage of this separation into two terms is that the alpha matrix can be estimated from a much smaller context than the nucleotide model, but the final model will still depend on the full context of the nucleotide model. With the bidirectional Markov model of order 14 and an alpha matrix dependent on just one base to each side, we obtain a model that compares well with a model of mutability that estimates mutation probabilities directly conditioned on three nucleotides to each side. For high-probability population variants, which are mainly CpG sites, the simple model fits better than our hybrid model, but for somatic variants it is opposite. Interestingly, the model is not very sensitive to the size of the context for the alpha matrix.

Conclusions: Our study found strong context dependencies of nucleotides in the human genome. The best model can estimate the nucleotide probabilities depending on contexts up to 14 nucleotides to each side. Based on these models, a substitution model was constructed that separates into the context model and an alpha matrix dependent in a small context. These models fit variants very well.

Keywords: DNA Context; Bidirectional Markov model; Subsitutional model

Background

The evolution of species can be followed in chromosomal DNA, which has undergone mutations and selection, and mutational processes have been essential for the development of life on earth. On the other hand mutations need to be controlled, because if an essential gene is mutated it may result in severe disease or loss of viability. This balance between plasticity and stability is important for sustaining stable life

forms [1]. The question we ask in this study is, how this balance is reflected in the local sequence properties of human DNA and how the sequence context affects mutations. More precisely, we consider models of mutability that depend on the sequence context of e.g. k bases on each side of the position in question.

It is well known that the sequence context influences mutational processes. For instance, the mutation of C to T is much more common in CpG dinucleotides than in other contexts in the human genome[2, 3], and previous studies have reported that the immediate neighbouring bases (up to a 7 base context) influence mutation rates [4, 5, 6]. Another study showed point mutations can be affected by sequence motifs [7]. The cellular machinery includes components for maintaining genome integrity, such as DNA repair mechanisms, which result in mutational biases[8, 9] and other processes may lead to other biases. These mechanisms together govern the intrinsic mutability. Following [10], we use the term mutability rather than mutation rate, because we are not considering the detailed evolutionary process and there is no time in our models, although the same ideas are easily applicable to estimation of context sensitive mutation rates.

Models of mutability can be estimated from observed variants by simply estimating the probability of a mutation given a context. However, such models are estimated from fairly small and biased sets of variants without utilizing the mutability foot-print in the genome. Here we propose to split the context dependent mutability into a nucleotide distribution and a variant part. The nucleotide distribution can be estimated from the whole genome and the variant part from variants, thereby allowing the two parts to have different context sizes. Due to the size of the human genome, the context dependent nucleotide distribution can be estimated from a much larger context than the variant part. The variant part can depend on a *smaller* context and can thus be estimated from a small number of variants.

In the first part of the paper, we focus on estimation of the probability of observing a base in the genome, given a context. One measure to quantify the context sensitivity is predictability. In a random sequence of nucleotides with no context sensitivity, we would only be able to predict a given base with an accuracy of 25% (random guessing), so this is the lower boundary of predictability. However, due to the mutational biasses discussed above and the repetitive nature of genomes, we would expect that a genome is more predictable than a random sequence. We show that a human genomic base can be predicted with an average of 51% using our most sophisticated model.

In the second part of the paper, we estimate a mutability model based on the context dependent nucleotide distribution found. For a fixed context dependent nucleotide distribution model, we show that the mutability is not very sensitive to the context size of the variant part. We compare to a simple mutability model conditioned on a 7 base context as in [5] and show that they differ between different types of mutations.

Results

Context modeling of the human genome

In our first model, we simply estimate the conditional probability of a nucleotide given k bases to each side. For base x_i at a genomic position i these probabilities

are written as

$$P(x_i|x_{i-k}, \ldots, x_{i-1}, x_{i+1}, \ldots, x_{i+k}).$$

They are estimated from the genomic frequencies of the 4 possible $(2k+1)$-mers of the given context. A $k = 3$ model corresponds to a neighbourhood of 7 as used in [5], and we use this model as our baseline. Since we are estimating frequencies from all positions on both strands, they are automatically strand symmetric.

One can of course use other values of k as long as a model can be reliably estimated. As the 4 probabilities sum to one, there are $3 * 4^{2k}$ free parameters in the model, so the $k = 3$ model has around 12,000 free parameters, which can easily be estimated from the 6 billion sites of the two strands of the human genome. A $k = 7$ model has approximately 0.8 billion free parameters, and is thus the upper limit of what we can hope to reliably estimate for a genome like the human. Even with $k = 7$ there are many contexts that occur only once or very rarely. To avoid over-fitting, we have used an interpolated model, in which a model of order k is used to regularize a model of order $k + 1$ and so on (see Methods). Our second model has $k = 7$, but is interpolated from $k = 4$.

A Markov model of order k yields probabilities of the four bases conditional on the k previous bases. A Markov model also can be used to estimate from both strands, as above, which means that for base i, it can give two different probabilities: $P(x_i|x_{i-1}, \ldots, x_{i-k})$ on the direct strand and $P(\hat{x}_i|\hat{x}_{i+1}, \ldots, \hat{x}_{i+k})$ on the opposite strand, where $\hat{x}_i$ means the complementary base to base x_i. Note that these models are estimated from both strands as the central models, which means that a model estimated using a 5' context is identical to the complementary of a model estimated using a 3' context and therefore, without loss of generality, we always assume 5' models.

Our third model is a bidrectional interpolated Markov model of order $k = 14$, interpolated from $k = 8$ (see Methods). It is called bidirectional, because we use the *average* between the probability of x_i from one strand and the probability of $\hat{x}_i$ from the opposite strand as explained above. Note that this model with $k = 14$ has the same number of free parameters $(3 * 10^{14})$ as the central model with $k = 7$ described above, because both use 14 bases as context. However, the bidirectional Markov model actually uses a context of 28 bases for prediction, because of the averaging over the two directions. This model is called BM14 in the following.

We have developed a program written in C that implements these different models. Instead of saving counts for each context, it dynamically calculates the count based on a Burrows-Wheeler encoded genome [11] to save memory. The performance of our models can be evaluated by the accuracy, which is the fraction of positions, where the most probable base given the context equals the actual base in the reference genome. The accuracy is shown in Figure 1 for the different models mentioned above (Additional file 1: Table S1).

For estimating the performance shown in Figure 1 , we have used leave-one-out cross-validation at the nucleotide level. It means that when estimating the probabilities for a given site in the genome, that site is excluded in the counts for model

Figure 1 Prediction accuracy for the three models.

Baseline with $k = 3$, Central model with $k = 7$ and the bidirectional Markov model with $k = 14$ (BM14). The bar-plot shows accuracy for each chromosome and average accuracy on the whole genome. Results using nucleotide-based cross-validation.

estimation. We have also done a chromosome-based cross-validation for comparison and calculated the overall accuracies for each chromosome using a model estimated from the *other* chromosomes. The difference between nucleotide-based and chromosome-based cross validation is only 0.5 percentage points (p.p.) on average, but for the Y chromosome, it is more than 3 p.p. (Additional file 1: Table S2 and Additional file 2: Figure S1).

With interpolation it is in principle possible to go beyond $k = 14$, because for contexts with zero counts, the probabilities are equal to a lower order estimate, so it should adapt without over-fitting. We have not explored higher k so much, but in Figure S2, we have run the bi-directional Markov model from $k = 10$ to $k = 20$ for chromosome 20 and different values of the interpolation constant described in Methods. Up to $k \simeq 14$ the models steeply improve and are insensitive to the interpolation constant. Above $k = 14$ we still see a monotonous improvement that seems to level off at around 52% for the best model. Chromosome 20 was chosen for this experiment, because it is small and has a prediction accuracy similar to the average for the BM14 model.

The predictive performance of BM14 on different regions in the human genome is shown in Figure 2. As expected, the model is very good at predicting in repetitive sequences with an overall accuracy of 64%, but there are quite large differences between different types of repeats. The most common type of repeat in the human genome, the ALU sequences, is 87% correctly predicted, whereas LINE1 for instance is only at 63% (Additional file 1: Table S3).

The density of the reference base probability for different genomic regions is shown in Figure 3. For each feature except for CDS there are two peaks of which one is due to repeats. However, in positions where the reference probability is above 0.4,

Figure 2 Prediction accuracies for BM14 in different regions across all chromosomes.

The accuracy for different features on the chromosome 1 to Y, is indicated by colored dots. The line shows the overall accuracy for each chromosome.

Figure 3 Density profile of reference probabilities in different genomic regions obtained with BM14.

repeats account for a large proportion compared to other features. (Additional file 1: Table S4).

To further elucidate the predictability across different regions, we show in Figure 4 the reference probabilities across human 3' and 5' splice sites that averaged over all introns annotated in Chromosome 1 (Chr1). The probability shows a large jump from a level of almost random prediction (~ 0.28) in the coding region to a fairly high value (~ 0.36) in the intron. The conservation plot in the same figure presents an opposite trend.

To test whether the model can be improved for non-repeat regions, we estimated a restricted model from everything *outside* coding regions and repeats. There is little difference between the restricted model and the full one in terms of prediction

Figure 4 Probabilities (top) and conservation score (bottom) of reference bases across 3' and 5' splice sites.

The probabilities of the reference bases by BM14 were averaged for each position for the first/last 100 nt in coding sequence and 500 nt in introns. The conservation score is phastcons from the UCSC browser.

accuracy or reference probability as seen in (Additional file 2: Figure S3) and we did not analyze this model further.

We briefly examined the performance of a bidirectional Markov model on some other species. Because of the smaller genome sizes, we used an interpolated bidirectional Markov model of order $k = 10$ in this analysis. The density plot of the reference probabilities in (Additional file 2: Figure S4) shows that a main peak occurs for human and E.coli genomes. In positions where the reference probability is above ~ 0.55, the curve of human is higher than that of other species, which could be caused by repeats in human genome. *A. thaliana*, *C. elegans* and *S. cerevisiae* have two peaks, of which the peak towards low probability is associated with coding regions.

Variants

We next evaluated BM14 on variant datasets. We assume that our models are valid for all genomes, and variants found in population studies, such as the 1000 Genome Project (1KGP) [12], should be predicted with the same accuracy as the

Figure 5 The triangle plot for probabilities of Ref-Alt alleles.

It was estimated by $k = 7$ context central model and $k = 14$ bidirectional Markov model, respectively. Only SNPs from 1000 Genome Project dataset were calculated.

corresponding positions in the reference genome. We identified ~ 73 million bi-allelic single nucleotide polymorphisms (SNPs) in the 1KGP. The probability of the reference (Pref) was plotted against the probability of the alternative (Palt) shown in Figure 5 for the $k = 7$ central model and BM14. The latter shows a larger concentration of sites in the middle of the plot. Note the unexpected asymmetry between the corners at Pref$\simeq 1$ and Palt$\simeq 1$ for both models.

This asymmetry is also reflected in the fact that the reference allele had the highest probability in 38.82% of cases and the alternative allele in only 24.20% for BM14. The density plot of Pref-Palt in Figure 6A also shows a peak near 1 when all SNPs are used. However, when rare SNPs are ignored, the right peak decreases in size and a peak in the left side of the plot appears and the density becomes symmetric when only including SNPs with allele frequency above 20%. The far majority of SNPs with a reference probability higher than 0.875 in the 1KGP dataset belong to repeats.

We also compared Pref and Palt for different types of single nucleotide variants (SNVs) in coding (Figure 6B) and non-coding regions (Additional file 2: Figure S5). Clinically relevant mutations from the Clinvar database are almost indistinguishable from 1KGP in coding regions, whereas somatic mutations have a clear tendency to mutate towards a more probable base (Palt > Pref). In non-coding regions, the somatic mutations are also shifted towards a higher probability for the alternative and have the same peak at high reference probability as 1KGP.

To see if there is a difference between damaging and benign SNPs, we show the same densities for Polyphen2 predictions [13] on Chr1 in Figure 6C. On Chr1 there is

Figure 6 Density profiles of Pref - Palt for SNPs on Chromosome 1.

A. SNPs from 1KGP. The different lines represent SNPs with allele frequencies greater than 0, 0.01, 0.1 and 0.2, respectively. SNP counts are shown in the legend after the dash. B. Density profiles show variants of ClinVar, somatic mutations (COSMIC) and 1KGP database in coding regions. C. Densities of damaging and benign variants predicted by Polyphen-2 based on the HumanVar database and annotated on 1KGP database by ANNOVAR software.

a total of 32,841 SNPs classified as benign and 15,299 SNP sclassified as damaging. There is a small shift of the damaging SNPs towards higher probability of the alternative allele. However, when comparing the two classes, we saw that for only 21% of damaging SNPs the reference allele had the highest probability whereas for 29% the alternative allele had the highest probability. For benign SNPs, these numbers are 26.5% and 24%.

Context-dependent models of substitutions

It is possible to estimate context dependent models of single nucleotide substitutions from a set of known variants. Since SNV sampling is very biased and variants are not fully observed, the context size needs to be much smaller than for the nucleotide distribution models described above. In the previously mentioned work [5] a seven nucleotide context is used. Here we want to explore the possibility of using our genome models to obtain models of substitutions. The rationale is that to maintain the context dependent nucleotide probabilities, they must be reflected in the mutability.

We assume the genome has reached approximate equilibrium. To keep this state, the mutability towards a nucleotide should be higher, the higher the probability of that nucleotide is in the given context. Therefore we set the probability of a mutation from a to b to be proportional to the probability of nucleotide b (in that context) with a constant that depends on the nucleotides and which can also depend on the context. This model is inspired by the general time-reversible stationary Markov model [14, 15], in which the off-diagonal rates are $\mu_{ab} = \alpha_{ab}\pi_b$ with symmetric α_{ab} for nucleotides $a \neq b$ and the equilibrium distribution $P(a) = \pi_a$. The mathematical theory does not apply directly here, because reversibility is too restrictive, so we do not require the α matrix to be symmetric, but we can still estimate an α matrix that best fits a set of variants. For lack of a better term, we call α the "alpha matrix".

Whereas the nucleotide distribution can be estimated from the whole genome using large contexts, the αs must be estimated from observed mutations. We hypothesize that the αs are less context dependent, and thus can be estimated from

Figure 7 Substitution model.

Model substitution probabilities shown for the models with context-insensitive α (k=0), the ones with α depending on 1, 2, and 3 bases to each side (k=1, 2, 3), and the simple model conditioned on the 3 bases to each side. The model substitution probability for a site is the sum of the probabilities for the three possible substitutions. A: The cumulative distribution of model substitution probabilities for all sites (solid lines) and for SNPs (dashed) on Chr1 shown for the five models. Note that for all models there are very few sites with substitution probability above 0.3. B: The fraction of sites on Chr1 with an observed variant in the 1000 Genome project (1KG) plotted against p. The y values are SNP counts in small probability intervals (10^{-4}) divided by total counts. The curves are smoothed with splines. Estimates are noisy for larger probabilities due to low counts. C: As B for SNPs in 1KG, Clinvar and COSMIC for the $k = 1$ model and simple only. For latter two, counts are scaled so they sum to the number of SNPs in the 1KG set for Chr1. For high mutability values there are few SNPs, so the curves are very noisy especially for Clinvar.

Table 1 α matrixes for $k = 0$ and $k = 1$ estimated by substitution model.

α matrix, $k = 0$					α matrix, $k = 1$, CG sites only			
	A	C	G	T		A	C	G	T
A	–	0.019	0.066	0.012	ACG	0.041	–	0.041	0.717
C	0.025	–	0.034	0.096	CCG	0.035	–	0.066	0.555
G	0.096	0.033	–	0.025	GCG	0.062	–	0.035	0.566
T	0.012	0.065	0.019	–	TCG	0.043	–	0.048	0.483

Left: The α matrix for $k = 0$ estimated from all chromosomes except Chr1. Right: The part of the α matrix for $k = 1$ corresponding to contexts with CG preceded by one base, so they correspond to mutations of C in these contexts.

a smaller context than the nucleotide distributions. Details of the estimation procedure is described in Methods.

We estimated αs from all chromosomes except Chr1 for symmetrical contexts of size 0, 3, 5, and 7 ($k = 0, 1, 2,$ and 3) using SNPs from the 1KGP and the BM14 model for the nucleotide distribution. The alpha matrix is shown in Table 1 (left) for $k = 0$. Notice that it is essentially strand-symmetric, but not symmetric in normal matrix-sense, so it violates reversibility. Similarly, we estimated a simple conditional model with a 7-mer context ($k = 3$) from the same data, which is called the simple model in the following. The simple model is similar to one of the models in [5], but the variants used for estimation are slightly different. The models were then applied to Chr1 where we calculated the probability of a mutation given the context for all positions with an observed SNP. In Figure 7B the fraction of sites with a certain mutability that has an observed SNP is plotted against mutability for some of the models. Ideally these should be linear, but we see a significant deviation from linear for the simple model and for the α models with $k > 0$. The models with $k = 1$–3 behave almost the same, and up to a substitution probability of ~ 0.25 they are

very close to the simple model. The total fraction of sites with probability above 0.25 is very small for all models, see Figure 7A.

Above a mutability of 0.25, our models with $k > 0$ deviate significantly from the diagonal line. It turns out that these rare reference genome sites with high substitution probability are mainly CpG sites. The alpha matrix for $k = 1$ is shown in Table 1 for the CG contexts, where it is evident that the C to T values are very large, ranging from 0.48 to 0.72, which should be compared to the largest α of 0.22 that is not a CG context, see (Additional file 1: Table S5). For contexts where the T has high probability according to the nucleotide distribution, the substitution probabilities will become large, because it is the product of α and the nucleotide probability. It suggests – as expected – that these substitutions are very likely at unselected positions.

We applied the model also to SNVs from Clinvar and COSMIC as shown in Figure 7C for $k = 1$ and for the simple model. The number of variants is small for large mutability values, so the data are a bit noisy, but it is evident that the somatic SNVs from COSMIC follow the model more closely than germline SNPs in this domain.

Discussion

We developed context dependent models of the nucleotide distribution in the human genome. The most advanced one, a bi-directional Markov model with a context of 14 nucleotides to each side, can predict a nucleotide with 51% accuracy. We use interpolation from lower orders, so it is in principle possible to go above $k = 14$, but we saw that this did not change the model very much, and the predictability of just above 50% is close to an upper limit for this type of model. The high predictability is, to a large extent, due to repeats. It is interesting that approximately half the human genome is said to be repetitive [16], which superficially coincides with the predictability, but an exact definition of repetitive regions is a challenge and some report a higher repetitive fraction (see e.g. [17]). Perhaps predictability is a better measure of the repetitiveness of a genome.

Not surprisingly, the predictability is highly dependent on the type of the genomic region. Coding regions can be predicted with only 36% accuracy, whereas Alu repeat regions are at 87% and simple repeats even higher (Figure 2). When looking more closely at splice sites we see – as expected – a negative correlation between conservation and the probability of the reference base (Figure 4), although such a correlation is weak, when looked at genome wide due to the lack of conservation of repeats. There are also differences between chromosomes, where especially the Y chromosome and Chr19 stand out with higher predictability than others, likely due to their high repeat content.

We compared the probability of the reference allele to the alternative allele on single nucleotide variants from the 1000 Genomes Project. There is a peak with SNPs that have a reference probability close to one, which skews the distribution away from symmetry (Figure 6). Almost all SNPs in this peak (with reference probabilities over 0.875) fall in repeat regions and one possibility is that some of them are mapping artefacts. They also have relatively low allele frequencies, and when considering only SNPs with high allele frequency, the plot becomes symmetric.

Therefore, another factor that may explain the asymmetry is that the reference genome, which is not a genome of a single individual, contains very few rare alleles.

The sequence models presented here estimate distributions of the bases for a given context and reflect inherent properties of the cellular machinery responsible for replication, error correction, and so on, as well as the physical properties of DNA, such as curvature and bendability. A mutation that moves a base closer to this distribution is likely to be more probable than one that moves it away, at least if selection is ignored. To explore this, we have derived a model that takes the context dependent nucleotide distribution into account.

In our model, we are assuming that the variation of a site in the human DNA can be described by a context sensitive continuous Markov model with a rate matrix that is a product between the nucleotide distribution and an "alpha matrix". The alpha matrix can be estimated from known variants and it can depend on a smaller context than the model for the nucleotide distribution and can be estimated from a relatively small number of SNVs. It means that our model for mutability have a very large context due to the context dependent nucleotide distribution even if the alpha matrix uses a smaller context.

The model does not depend strongly on the context size for the alpha matrix for contexts of the two neighbours or larger ($k \geq 1$). Our models behave very similarly to a simple mutability model, which is estimated from SNPs alone and a context of three nucleotides to each side except in a regime of very high mutability (Figure 7b). Our models seem to over-estimate the SNP mutability from 1KGP when the values are larger than about 0.25. However, this is not the case for somatic mutations, and the mutations seem to be well-described by these models (Figure 7C).

The model is inspired by the general time-reversible model from evolutionary theory, which has six free parameters corresponding to a symmetric alpha matrix, and with rates depending on the equilibrium distribution. However, although time-reversibility would be desirable, it is not likely that the context dependent nucleotide distribution we estimate is an equilibrium distribution for the entire genome. In fact, when inspecting the estimated alpha matrix for zero context (Table 1) and a context of one nucleotide to each side (Additional file 1: Table S5), it is evident that it is not symmetric. For the latter there are very large deviations from symmetry for contexts with NCG, where N can be any base. In these contexts, α_{CT} is consistently 10-20 times larger than α_{TC} corresponding to a strong tendency to mutate from CG to TG.

Even if the α matrix depends on a small context, the substitution still depends on the full context of the nucleotide distribution. This construction is very attractive, because substitution models estimated from variants alone need to have small contexts due to the limited number of variants and the strong sampling biases.

Conclusions

There are strong context dependencies of nucleotides in genomes. We have shown how one can estimate a model of the nucleotide probabilities depending on contexts up to 14 nucleotides to each side. Building on these models, it was shown how it is possible to make models of mutations that combine the context dependent nucleotide probabilities with a mutation matrix, called the alpha matrix, to give

mutation probabilities ("mutabilities") that depend on the same large context. It was shown that these models fit observed mutations very well and especially somatic ones. Importantly, the alpha matrix can depend on a much smaller context of just one to three bases to each side and does not depend strongly on this parameter.

These models can form the basis for a better understanding of human mutations and we believe it will be possible to use them in a wide range of applications from GWAS studies to analysis of somatic mutations.

Methods

Conditional probability models for the central base

The base at position i (chromosome, coordinate) in the reference genome is called x_i and the symmetric sequence context around it is called

$$s_i(k) = x_{i-k}, x_{i-k+1}, \ldots, x_{i-1}, x_{i+1}, x_{i+2}, \ldots, x_{i+k}. \tag{1}$$

If it is clear from the context which k, we call it s_i to ease notation. To estimate the conditional probability of base b at position i, we use the counts $n(b|s_i)$ of the occurrences in the same context throughout the reference genome (on both strands):

$$P(b|s_i) = \frac{n(b|s_i) - \delta_{b,x_i}}{N(s_i) - 1}, \tag{2}$$

where

$$N(s_i) = \sum_b n(b|s_i).$$

We use the Kronecker δ_{b,x_i}, which is 1 if $x_i = b$ and otherwise 0, to ensure that we only count *other* contexts, when estimating probabilities at position i. This is leave-one-out cross-validation and is discussed further below.

For large contexts, the counts become small and thus the probabilities cannot be reliably estimated. To interpolate between different orders of the model, we use regularization by pseudo-counts obtained from the $k-1$ model. Specifically, for order k, we define pseudo-counts

$$r(b|s_i(k)) = \gamma P(b|s_i(k-1)),$$

where γ is the strength of pseudo-counts. Now the model of order k is estimated as before, but using the actual counts plus pseudo-counts,

$$P(b|s_i(k)) = \frac{n(b|s_i(k)) - \delta_{b,x_i} + r(b|s_i(k))}{N(s_i(k)) - 1 + \gamma}.$$

The advantage of pseudo-counts is that they have minor influence, when there is plenty of data (actual counts are high), but have strong effect at low counts. With $k = 4$ counts are on average $6 * 10^9 / 4^9 \simeq 23000$, so we assume that psudo-counts are not needed. Therefore, our interpolated model starts with unregularized estimates for $k = 4$, and then use the pseudo-counts iteratively for $k = 5$ to $k = 7$ for the interpolated model. We used a strength of $\gamma = 100$ for the pseudo-counts (a few experiments showed that the model is relatively robust to changes in γ, see below).

Markov models

In a Markov model of order k, the probability of a base is conditioned on the k previous bases. If we redefine the k-context in (1) to be the k previous bases,

$$s_i(k) = x_{i-k}, x_{i-k+1}, \ldots, x_{i-1},$$

we can use exactly the same formulation as above. In this case however, the context size is not $2k$ letters as above, but only k letters. Therefore, one can estimate Markov models up to sizes around $k = 14$ for the human genome, and we used a model interpolated from $k = 8$ to $k = 14$ analogously to the central interpolated model described above.

Due to the interpolation, larger k are possible, and we did a small experiment with k ranging from 10 to 20 and with four different values of the interpolation constant γ resulting in Figure S2. These tests were done only on chromosome 20 with a model estimated from all chromosomes *except* 20. Although small gains can be obtained with larger k values and different γ, we decided to stick to our initial choice of $k = 14$ and $\gamma = 100$.

Estimating a "forward" Markov model from both strands of the human genome will automatically make it strand-symmetric. For a given position in the genome, the model can therefore give two sets of base probabilities: one for the forward strand and one for the reverse strand. Our final Markov probabilities are the average between the two as described in the main text and referred to as bidirectional.

Cross-validation

Our way of estimating the conditional probability of seeing one of the four bases given the surrounding context can be seen as a leave-one-out procedure. In particular, the estimate depends on the reference base at the considered position as well as the context. To obtain an estimate that is independent of the reference base at the position, a natural way to proceed is to consider the average of the four base-dependent estimates over all occurrences of the given context. This average turns out to be equal to the estimate that includes all positions. To see this, average (2) over all sites (skipping the k dependence for clarity)

$$\bar{P}(b'|s) = \frac{1}{N(s)} \sum_b n(b|s) \frac{n(b'|s) - \delta_{b,b'}}{N(s) - 1}.$$

Since $\sum_s n(b|s)\delta_{b,b'} = n(b'|s)$, we get

$$\bar{P}(b'|s) = \frac{1}{N(s)(N(s) - 1)}(N(s)n(b'|s) - n(b'|s)) = \frac{n(b'|s)}{N(s)}.$$

We also assessed our models by cross-validation by chromosomes. One chromosome was used as test data, and the remaining chromosomes as training data. We repeated this step 24 times to calculate the fraction correct predictions for each chromosome.

Substitution models

A simple model estimates mutability as the fraction of all sites with context $\hat{s}$ having a specific mutation. More specifically,

$$P_{\text{Simple}}\,(a \to b|\hat{s}) = \frac{n(a \to b|\hat{s})}{n(a|\hat{s})}.\tag{3}$$

Here $n(a \to b|\hat{s})$ is the number of observed mutations $a \to b$ in context $\hat{s}$ and $n(a|\hat{s})$ is the number of times we see reference base a in context $\hat{s}$ (as above). We use $\hat{s}$ to indicate that the context may be different from the context s for the genome model above. We have used this model with a symmetric context of three bases to each side, which we call the simple model.

We will now derive a continuous time Markov model with context dependent substitution rates $\mu_{ab|s}$ that takes the nucleotide distribution into account. We also assume a constant evolutionary time, which is infinitesimally small compared to the rates, so we can approximate the substitution probability by the first-order term in the Taylor expansion of an exponential

$$P(a \to b|s) \simeq \delta_{a,b} + \mu_{ab|s},$$

where time is set to 1. The diagonal rates are $-\sum_{b \neq a} \mu_{ab|s}$, so in the following we will not write the diagonal terms. For a stationary, reversible Markov model with $P(a|s)$ as equilibrium probabilities the rates can be written as

$$P(a \to b|s) \simeq \mu_{ab|s} = \alpha_{ab|s} P(b|s) \qquad (a \neq b).$$

with a symmetric matrix α_{ab}. This is the general time-reversible six-parameter model (see e.g. [15]). Inspired by this model, we assume that mutability is given by the same equation, but without requiring that the nucleotide distribution is the equilibrium distribution and without requiring that α is symmetric.

The above expression factorizes the rates into the nucleotide distribution and the α-term that encapsulates the mutations. Now we assume the αs depend on a *smaller* context $\hat{s}$ than the context s for the genome model $P(a|s)$, so the above can be written as

$$P(a \to b|s) \simeq \mu_{ab|s} = \alpha_{ab|\hat{s}} P(b|s) \qquad (a \neq b)\tag{4}$$

In analogy with (3), $P(a \to b|s) = n(a \to b|s)/n(a|s)$ with s instead of $\hat{s}$, so combining with the above

$$n(a \to b|s) \simeq n(a|s)\alpha_{ab|\hat{s}} P(b|s) \qquad (a \neq b)$$

To estimate the αs we sum over all contexts that contains $\hat{s}$, which we write as $s|\hat{s} \subseteq s$, so

$$n(a \to b|\hat{s}) = \sum_{s|\hat{s} \subseteq s} n(a \to b|s) \simeq \alpha_{ab|\hat{s}} \sum_{s|\hat{s} \subseteq s} n(a|s) P(b|s)$$

The last sum depends only on the nucleotide distribution. It can be rewritten as a sum over all positions in the genome, where the reference base, r_i, equals a and where the context is $\hat{s}$. We call this term $Z_{ab|\hat{s}}$,

$$Z_{ab|\hat{s}} = \frac{1}{n(a|\hat{s})} \sum_{s|\hat{s} \subseteq s} n(a|s) P(b|s) = \frac{1}{n(a|\hat{s})} \sum_{i|r_i=a \wedge \hat{s} \subseteq s_i} P(b|s_i),$$

For convenience, it is normalized by $n(a|\hat{s})$, so it is the average probability of base b over all positions with reference base a and context $\hat{s}$. As an estimate of α we then have

$$\alpha_{ab|\hat{s}} = \frac{1}{Z_{ab|\hat{s}}} \frac{n(a \to b|\hat{s})}{n(a|\hat{s})} = \frac{P_{\text{Simple}}(a \to b|\hat{s})}{Z_{ab|\hat{s}}}$$

Note that we can rewrite the the original probability (4) in terms of the simple model as

$$P(a \to b|s) \simeq \frac{P(b|s)}{Z_{ab|\hat{s}}} P_{\text{Simple}}(a \to b|\hat{s})$$

for $\hat{s} \subseteq s$. The factor is 1 when $\hat{s} = s$, so the models are identical as they should be when they use the same context. The equation directly shows how the wider context from the genome model can modulate the simpler estimate. If the probability of base b in context s is larger than the mean $Z_{ab|\hat{s}}$, the mutability becomes larger than in the simple model, and if it is smaller, the mutability becomes smaller.

The first order approximation assumes the rates are small. When calculating the total mutability of a site, we therefore use the approximation $1 - P(a \to a|s) \simeq 1 - e^{\mu_{aa|s}}$. For small α's it makes little difference whether it is the exponentiated form or not.

Data
The human reference genome, GRCh38.p13, was downloaded from NCBI (released March 2019 by Genome Reference Consortium). We considered only primary assemblies of chromosomes 1 to 22 and X, Y. Genomic annotation bed files were downloaded from the UCSC Genome Browser. These were 3'-UTR, 5'-UTR, CDS, Introns, Genes, and Repeats. PhastCons scores file was downloaded from the UCSC, as well.

Variants were downloaded from the 1000 Genome Project (release March 2019, phased 20190312_biallelic_SNV_and_INDEL) in VCF format. The Indels were filtered from 1KGP dataset.

ClinVar (clinvar_20200310.vcf) [18, 19] and somatic mutation data (CosmicCodingMuts.vcf and CosmicNonCodingVariants.vcf) [20] were obtained from NCBI and COSMIC, respectively.

Data analysis
Accuracy We calculated the probabilities of the 4 bases for every position in the genome using the models in question. From these we obtained the overall accuracy

per chromosome. By intersection with the bed feature files [21, 22], we calculated feature-specific quantities. Fraction corrects were plotted by a dot-plot (Figure 2) for these features of different chromosomes.

For the plot of probability (Figure 4) around exons, we calculated both probabilities at human 3' and 5' splice sites. A 100nt long protein coding sequence was used after the 3' splice site and in front of the 5' splice site, respectively. In addition, we did a plot by using conservation scores of Phastcons at the same regions [23].

Prediction probability of SNP variants We considered only single nucleotide bi-allelic variants. The model probabilities for the reference allele and alternative allele were extracted for of each SNP site of 1KGP, ClinVar and COSMIC databases.

The triangle plots were made using reference probabilities against alternative probabilities of 1KGP database. Additionally, we used ANNOVAR software [24] to annotate benign and damaging SNPs on 1KGP predicted by PolyPhen2 [13]to be associated with single genetic disease. Densities were plotted by only using these reference probabilities or both reference and alternative probabilities.

Test Markov model on other species The genomes of *Arabidopsis thaliana*, *Escherichia coli*, *Caenorhabditis elegans* and *Saccharomyces cerevisiae* were downloaded from NCBI. We tested our bi-direct markov model on these species and also *Homo sapiens*, then calculated their prediction probability by setting $k = 10$, $\gamma = 100$ and interpolated from $k = 6$.

Software

Counting of k-mers and estimation of probabilities is implemented in the C programming language. The program counts the contexts for each site using a Burrows-Wheeler transform to minimize memory usage, rather than storing the k-mers. The program is called predictDNA and relies on an index built with the program makeabwt. The software is open sourse and available at GiHub: https://github.com/AndersKrogh/abwt/releases/tag/v1.2.1

Acknowledgements
We thank Hanne Munkholm for her big help and support with compute servers. AK acknowledges a visiting scholarship from University of Torino.

Funding
YL acknowledges China Scholarship Council (Grant 201804910693) for Ph.D. financial support.

Availability of data and materials
All data used in this study are publicly available. All data can be downloaded from NCBI, UCSC, 1KGP and COSMIC database as we mentioned in our methods.

Ethics approval and consent to participate
Not applicable.

Competing interests
The authors declare no competing interests.

Consent for publication
Not applicable.

Authors' contributions
AK and PF initiated the project. YL and AK performed most analyses and drafted the paper with assistance from CG. All authors participated in revision and approved the final version.

Author details
[1]Department of Computer Science, University of Copenhagen, Copenhagen, Denmark. [2]Department of Biology, University of Copenhagen, Copenhagen, Denmark. [3]Department of Medical Sciences, University of Torino, Torino, Italy.

References
 1. Schubert, I., Vu, G.T.: Genome stability and evolution: attempting a holistic view. Trends in plant science **21**, 749–757 (2016)
 2. Cooper, D.N., Youssoufian, H.: The CpG dinucleotide and human genetic disease. Human genetics **78**, 151–155 (1988)
 3. Hess, S.T., Blake, J.D., Blake, R.D.: Wide variations in neighbor-dependent substitution rates. Journal of molecular biology **236**, 1022–1033 (1994)
 4. Krawczak, M., Ball, E.V., Cooper, D.N.: Neighboring-nucleotide effects on the rates of germ-line single-base-pair substitution in human genes. American journal of human genetics **63**, 474–488 (1998)
 5. Aggarwala, V., Voight, B.F.: An expanded sequence context model broadly explains variability in polymorphism levels across the human genome. Nature genetics **48**, 349–355 (2016). doi:10.1038/ng.3511
 6. Carlson, J., Locke, A.E., Flickinger, M., Zawistowski, M., Levy, S., Myers, R.M., Boehnke, M., Kang, H.M., Scott, L.J., Li, J.Z., *et al.*: Extremely rare variants reveal patterns of germline mutation rate heterogeneity in humans. Nature communications **1**, 1–13 (2018)
 7. Zhu, Y., Neeman, T., Yap, V.B., Huttley, G.A.: Statistical methods for identifying sequence motifs affecting point mutations. Genetics **205**, 843–856 (2017)
 8. Lind, P.A., Andersson, D.I.: Whole-genome mutational biases in bacteria. Proceedings of the National Academy of Sciences **105**, 17878–17883 (2008)
 9. Pearson, C.E., Edamura, K.N., Cleary, J.D.: Repeat instability: mechanisms of dynamic mutations. Nature Reviews Genetics **6**, 729–742 (2005)
10. Zavolan, M., Kepler, T.B.: Statistical inference of sequence-dependent mutation rates. Current opinion in genetics & development **11**, 612–615 (2001)
11. Li, H., Durbin, R.: Fast and accurate short read alignment with burrows–wheeler transform. bioinformatics **25**, 1754–1760 (2009)
12. Consortium, T..G.P.: A global reference for human genetic variation. Nature **526**, 68–74 (2015). doi:10.1038/nature15393
13. Adzhubei, I.A., Schmidt, S., Peshkin, L., Ramensky, V.E., Gerasimova, A., Bork, P., Kondrashov, A.S., Sunyaev, S.R.: A method and server for predicting damaging missense mutations. Nature methods **7**(4), 248–249 (2010)
14. Tavaré, S.: Some probabilistic and statistical problems in the analysis of dna sequences. Lectures on Mathematics in the Life Sciences **17**, 57–86 (1986)
15. Felsenstein, J., Felenstein, J.: Inferring Phylogenies vol. 2. Sinauer associates Sunderland, MA, ??? (2004)
16. Lander, E.S., Linton, L.M., Birren, B., Nusbaum, C., Zody, M.C., Baldwin, J., Devon, K., Dewar, K., Doyle, M., FitzHugh, W., Funke, R., Gage, D., Harris, K., Heaford, A., Howland, J., Kann, L., Lehoczky, J., LeVine, R., McEwan, P., McKernan, K., Meldrim, J., Mesirov, J.P., Miranda, C., Morris, W., Naylor, J., Raymond, C., Rosetti, M., Santos, R., Sheridan, A., Sougnez, C., Stange-Thomann, Y., Stojanovic, N., Subramanian, A., Wyman, D., Rogers, J., Sulston, J., Ainscough, R., Beck, S., Bentley, D., Burton, J., Clee, C., Carter, N., Coulson, A., Deadman, R., Deloukas, P., Dunham, A., Dunham, I., Durbin, R., French, L., Grafham, D., Gregory, S., Hubbard, T., Humphray, S., Hunt, A., Jones, M., Lloyd, C., McMurray, A., Matthews, L., Mercer, S., Milne, S., Mullikin, J.C., Mungall, A., Plumb, R., Ross, M., Shownkeen, R., Sims, S., Waterston, R.H., Wilson, R.K., Hillier, L.W., McPherson, J.D., Marra, M.A., Mardis, E.R., Fulton, L.A., Chinwalla, A.T., Pepin, K.H., Gish, W.R., Chissoe, S.L., Wendl, M.C., Delehaunty, K.D., Miner, T.L., Delehaunty, A., Kramer, J.B., Cook, L.L., Fulton, R.S., Johnson, D.L., Minx, P.J., Clifton, S.W., Hawkins, T., Branscomb, E., Predki, P., Richardson, P., Wenning, S., Slezak, T., Doggett, N., Cheng, J.-F., Olsen, A., Lucas, S., Elkin, C., Uberbacher, E., Frazier, M., Gibbs, R.A., Muzny, D.M., Scherer, S.E., Bouck, J.B., Sodergren, E.J., Worley, K.C., Rives, C.M., Gorrell, J.H., Metzker, M.L., Naylor, S.L., Kucherlapati, R.S., Nelson, D.L., Weinstock, G.M., Sakaki, Y., Fujiyama, A., Hattori, M., Yada, T., Toyoda, A., Itoh, T., Kawagoe, C., Watanabe, H., Totoki, Y., Taylor, T., Weissenbach, J., Heilig, R., Saurin, W., Artiguenave, F., Brottier, P., Bruls, T., Pelletier, E., Robert, C., Wincker, P., Smith, D.R., Doucette-Stamm, L., Rubenfield, M., Weinstock, K., Lee, H.M., Dubois, J., Rosenthal, A., Platzer, M., Nyakatura, G., Taudien, S., Rump, A., Yang, H., Yu, J., Wang, J., Huang, G., Gu, J., Hood, L., Rowen, L., Madan, A., Qin, S., Davis, R.W., Federspiel, N.A., Abola, A.P., Proctor, M.J., Myers, R.M., Schmutz, J., Dickson, M., Grimwood, J., Cox, D.R., Olson, M.V., Kaul, R., Raymond, C., Shimizu, N., Kawasaki, K., Minoshima, S., Evans, G.A., Athanasiou, M., Schultz, R., Roe, B.A., Chen, F., Pan, H., Ramser, J., Lehrach, H., Reinhardt, R., McCombie, W.R., de la Bastide, M., Dedhia, N., Blöcker, H., Hornischer, K., Nordsiek, G., Agarwala, R., Aravind, L., Bailey, J.A., Bateman, A., Batzoglou, S., Birney, E., Bork, P., Brown, D.G., Burge, C.B., Cerutti, L., Chen, H.-C., Church, D., Clamp, M., Copley, R.R., Doerks, T., Eddy, S.R., Eichler, E.E., Furey, T.S., Galagan, J., Gilbert, J.G., Harmon, C., Hayashizaki, Y., Haussler, D., Hermjakob, H., Hokamp, K., Jang, W., Johnson, L.S., Jones, T.A., Kasif, S., Kaspryzk, A., Kennedy, S., Kent, W.J., Kitts, P., Koonin, E.V., Korf, I., Kulp, D., Lancet, D., Lowe, T.M., McLysaght, A., Mikkelsen, T., Moran, J.V., Mulder, N., Pollara, V.J., Ponting, C.P., Schuler, G., Schultz, J., Slater, G., Smit, A.F., Stupka, E., Szustakowski, J., Thierry-Mieg, D., Thierry-Mieg, J., Wagner, L., Wallis, J., Wheeler, R., Williams, A., Wolf, Y.I., Wolfe, K.H., Yang, S.P., Yeh, R.F., Collins, F., Guyer, M.S., Peterson, J., Felsenfeld, A., Wetterstrand, K.A., Patrinos, A., Morgan, M.J., de Jong, P., Catanese, J.J., Osoegawa, K., Shizuya, H., Choi, S., Chen, Y.J., Szustakowski, J., Consortium, I.H.G.S.: Initial sequencing and analysis of the human genome. Nature **409**, 860–921 (2001). doi:10.1038/35057062
17. de Koning, A.P.J., Gu, W., Castoe, T.A., Batzer, M.A., Pollock, D.D.: Repetitive elements may comprise over two-thirds of the human genome. PLoS genetics **7**, 1002384 (2011). doi:10.1371/journal.pgen.1002384
18. Landrum, M.J., Lee, J.M., Benson, M., Brown, G.R., Chao, C., Chitipiralla, S., Gu, B., Hart, J., Hoffman, D.,

Jang, W., *et al.*: ClinVar: improving access to variant interpretations and supporting evidence. Nucleic acids research **46**, 1062–1067 (2018)

19. Landrum, M.J., Lee, J.M., Riley, G.R., Jang, W., Rubinstein, W.S., Church, D.M., Maglott, D.R.: ClinVar: public archive of relationships among sequence variation and human phenotype. Nucleic acids research **4**, 980–985 (2014)

20. Tate, J.G., Bamford, S., Jubb, H.C., Sondka, Z., Beare, D.M., Bindal, N., Boutselakis, H., Cole, C.G., Creatore, C., Dawson, E., *et al.*: Cosmic: the catalogue of somatic mutations in cancer. Nucleic acids research **47**, 941–947 (2019)

21. Quinlan, A.R., Hall, I.M.: BEDTools: a flexible suite of utilities for comparing genomic features. Bioinformatics **26**, 841–842 (2010)

22. Quinlan, A.R.: BEDTools: the Swiss-army tool for genome feature analysis. Current protocols in bioinformatics **47**, 11–12 (2014)

23. Castle, J.C.: SNPs occur in regions with less genomic sequence conservation. PLoS One **6**, 20660 (2011)

24. Wang, K., Li, M., Hakonarson, H.: ANNOVAR: functional annotation of genetic variants from high-throughput sequencing data. Nucleic acids research **38**, 164–164 (2010)

Additional Files

Additional file 1 — Supplementary tables S1-S5
Supplementary tables: Table S1, S2, S3, S4, S5

Additional file 2 — Supplementary Figures
Supplementary figures: Figure S1, S2, S3, S4, S5