Roles of proteins containing immunoglobulin-like domains in the conjugation of bacterial

plasmids.

Mário Hüttener[1], Jon Hergueta[1], Manuel Bernabeu[1], Alejandro Prieto[1], Sonia Aznar[1], Susana

Merino[1], Joan Tomás[1] and Antonio Juárez[1,2*]

[1]Department of Genetics, Microbiology and Statistics, University of Barcelona, Barcelona, 08028,

-. Spain.

[2]Institute for Bioengineering of Catalonia, The Barcelona Institute of Science and Technology,

Barcelona, 08028, - Spain.

Keywords: plasmid conjugation; Big proteins; multiple antibiotic resistance

17 *Corresponding author: Prof. Antonio Juárez, – ajuarez@ub.edu

Abstract

Horizontal transfer of bacterial plasmids generates genetic variability and contributes to the

dissemination of the genes that enable bacterial cells to develop antimicrobial resistance (AMR).

Several aspects of the conjugative process have long been known, namely, those related to the

proteins that participate in the establishment of cell-to-cell contact and to the enzymatic

processes associated with the processing of plasmid DNA and its transfer to the recipient cell. In

this work, we describe the role of newly identified proteins that influence the conjugation of

several plasmids. Genes encoding high-molecular-weight proteins that contain one or several

immunoglobulin-like domains (Big) are located in the transfer regions of several plasmids that

usually harbor AMR determinants. These Big proteins are exported to the external medium and

target two extracellular organelles: the flagella and the conjugative pili. The plasmid-encoded

Big proteins facilitate conjugation by reducing cell motility and facilitating cell-to-cell contact by

binding both to the flagella and to the conjugative pilus. They use the same export machinery as

that used by the conjugative pilus components. In the examples characterized in this paper,

these proteins influence conjugation at environmental temperatures (i.e., 25°C). This suggests

that they may play relevant roles in the dissemination of plasmids in natural environments.

Taking into account that they interact with outer surface organelles, they could be targeted to

control the dissemination of different bacterial plasmids carrying AMR determinants.

Introduction

Bacterial infectious diseases, despite the availability of antibiotics, remain an important public

health issue, representing the second leading cause of death worldwide [1]. The gradual increase

in the resistance rates of several important bacterial pathogens represents a serious threat to

public health [2–4]. Indeed, multidrug-resistant bacteria are the cause of a slow-growing

pandemic. The dissemination of multiple antimicrobial resistance (AMR) genes has been largely

attributed to the acquisition of plasmids by horizontal gene transfer (HGT), especially in Gram-

negative bacteria [5–7], as well as in Gram-positive bacteria [8]. Plasmids can confer resistance

to the major classes of antimicrobials [9].

For decades, plasmid incompatibility [10] has been a useful tool for grouping bacterial plasmids.

In recent years, other approaches have been considered. Based on phylogenetic analysis of

conjugative relaxase, the protein required to initiate plasmid mobilization through conjugation,

plasmids can be grouped into different relaxase families [11]. Plasmids belonging to the

incompatibility group (Inc) HI (MOB$_H$ relaxase family) are widespread in *Enterobacteriaceae* and

most commonly include genetic elements encoding multiple AMR determinants [12]. IncHI plasmids, often >200 kb in size, share a common core of approximately 160 kb. The differences in size are due to the presence of distinct insertion elements, including many AMR determinants [13]. IncHI-encoded AMR can be present in enterobacterial such as *Salmonella, Escherichia coli* [14], *Klebsiella pneumoniae* [15] and *Citrobacter freundii* [16]. Plasmids of the IncHI2 subgroup predominate in antibiotic-resistant *Salmonella* isolates. In *S.* Typhi*,* more than 40% of isolates harbor an IncHI plasmid [17]. In recent years, a novel role of IncHI plasmids in AMR spread has been reported. The emergence of Gram-negative bacteria with AMR, especially those producing carbapenemases, led to reintroduction of colistin as a last resort antibiotic for the treatment of severe infections [18]. In contrast to its limited clinical use, colistin is widely used in veterinary medicine [19]. In the past, colistin resistance was associated with chromosomal mutations only [20]. Nevertheless, plasmid-mediated resistance, conferred by the mobilized colistin resistance gene (*mcr-1*), has emerged recently. Since its discovery in 2016 in China [21], *mcr* genes, have been detected in animals, food, the human microbiota, and clinical samples in over thirty countries [22–26]. IncHI2 plasmids represent 20.5% of the overall plasmids encoding the *mcr-1* gene worldwide but up to 41% in Europe [27]. Of special concern is the presence of the *mcr-1* resistance determinant in *Enterobacteriaceae* carrying carbapenem resistance genes, such as bla_{NDM} and bla_{KPC}. The combination of these AMR determinants seriously compromises the treatment of infections caused by pathogenic strains harboring these plasmids [28,29]. An example of this is the recent report of an AMR clone of the highly virulent *E. coli* ST95 lineage [14]. *E. coli* ST95 clones underlie neonatal meningitis and sepsis. They are usually sensitive to several antibiotics. This clone harbors an IncHI2 plasmid that encodes, among other factors, determinants of resistance to colistin and multiple other antibiotics (including the extended-spectrum beta-lactamase *blaCTX-M-1*). The spread of such an AMR ST95 clone could pose a threat to human health worldwide [14].

IncHI plasmid conjugation has a distinctive feature: while optimal conjugation rates are obtained at temperatures found outside the host (30°C and below), conjugative transfer is repressed at temperatures encountered within the host (37°C) [30,31]. The plasmid R27 is the prototype of IncHI1 plasmids. It harbors the Tn*10* transposon, which confers resistance to Tc, and has been exhaustively studied. The R27 replication and conjugation determinants are well characterized [32,33], and its complete nucleotide sequence is available [34]. Several ORFs from the plasmid R27 (66%) do not show similarity to any known ORFs.

IncA/C plasmids belong to the same MOB$_H$ relaxase family as IncHI plasmids. They were originally identified in the 1970s among multidrug-resistant *Aeromonas hydrophila* and *Vibrio* spp. isolates

that infected cultured fish [35,36]. Since the 1990s, these plasmids have received increasing

interest because of their role in mobilizing AMR in enterobacteria and other Gram-negative

microorganisms [37–40]. They have an extremely broad host range that includes members of

Beta-, Gamma- and Deltaproteobacteria [41] and play a relevant role in the global spread of

AMR [42,43]. They represented 50% of all plasmids isolated from bla_{NDM}-producing *Klebsiella*

pneumoniae of clinical origin characterized in a recent study [44].

Proteins containing an immunoglobulin (Ig)-like domain contain several chains of approximately

70-100 amino acid residues present in antiparallel β-strands and organized in two β-sheets that

are packed against each other in a β-sandwich. The Ig-like domain has been identified in a large

number of proteins with diverse biological functions, is widely distributed in nature, and is

present in vertebrates, invertebrates, plants, fungi, parasites, bacteria, and viruses [45].

Bacterial proteins containing Ig-like domains (Big) exhibit a wide range of functions. They include

fimbrial subunits, adhesins, membrane transporters and several enzymes (as reviewed in [46]).

In a previous report, we studied a high-molecular-weight extracellular protein (the RSP protein)

that contains a Big domain and plays an essential role in IncHI plasmids conjugation. Among

other targets, the RSP protein appears to be associated with flagella, reducing cell motility.

Under specific mating conditions, it could be shown that binding of the RSP protein to the flagella

influences conjugation [47]. In this report, we present novel data about the roles of these

plasmid-encoded Big proteins. We show that two Big proteins bind both flagella and the

conjugative pilus to favor conjugation of the IncHI1 plasmid R27. Furthermore, we also show

that other groups of plasmids such as IncA/C and IncP2 also encode these proteins. We provide

evidence for their role in the conjugation of IncA/C plasmids. The role of plasmid-encoded Big

proteins in plasmid conjugation is discussed.

Results

The RSP protein interacts in vitro both with a new R27-encoded Big protein and with a protein

involved in plasmid conjugation.

To gain further insight into the role of the RSP protein in the conjugation of the R27 plasmid, we

decided to assess whether this protein interacts with other proteins expressed by the *Salmonella*

strain SL1344 (R27). We performed immunoprecipitation of a cellular extract of strains SL1344

(R27 RSP-Flag) and SL1344 (R27 Δ*rsp*) and analyzed the proteins that specifically coprecipitated

with the RSP protein. Two R27-encoded proteins were found to specifically coprecipitate with

the RSP protein (Table S1). The protein showing the highest score (187.77) and coverage (58.56)

was the R27_p055 protein. The *R0055* gene was mapped between transfer regions 2 and 1 of

the R27 plasmid (Fig. 1A). The *R0055* gene product is a 794 AA protein with a molecular mass of

86.75 kDa. As with the RSP protein, the protein encoded by *R0055* also contains bacterial Ig-like

domains (Big_1 and _3): a Big_1 domain spanning amino acid residues 143 to 254, and a Big_3

domain spanning residues 537 to 693. The R27_p055 protein, herein termed RSP2, also contains

a DUF4165 domain of unknown function (amino acids 23 to 142) (Fig. 1B).

The second protein showing a high score (59.46) and coverage (39.92) was the R27-encoded

TrhH protein. This protein shares 26% of identity with the IncF TraH protein, which is involved

in plasmid conjugation [32].

The RSP2 protein shows RSP-dependent expression in the external medium.

To determine whether the RSP2 protein is also present in the cell-free secretome of strain

SL1344 (R27), we analyzed the cell-free secreted protein profile of this strain by SDS-PAGE (Fig.

1C). In addition to the band corresponding to the already characterized RSP protein, a second

band corresponding to a protein of molecular mass equivalent to that of the RSP2 protein, was

observed. The band was isolated and analyzed by LC-MS/MS. It corresponded to the RSP2

protein (Fig. 1C). To identify the RSP2 protein in the different cellular compartments, a Flag-tag

was added to the *rsp2* gene (see Materials and Methods section for details). Cultures of strains

SL1344 wt and SL1344 (R27 RSP2-Flag) were grown in LB medium at 25°C to an OD_{600nm} of 2.0.

Samples were then collected, and the different cellular fractions were obtained. The RSP2

protein was detected in the different fractions by Western blotting, using anti-Flag-specific

antibodies (Fig. 2A-C). The protein was identified in the same cell compartments as the RSP

protein (i.-e., periplasm, inner membrane, cytoplasm, and cell-free secreted proteins).

As the above reported data show that the RSP2 protein can be exported to the external medium,

we decided to study whether this protein is exported by the R27-encoded type IV secretion

system that is also used by RSP [47]. To address this point, we first used the SSPred program

(http://www.bioinformatics.org/sspred/html/sspred.html) for *in silico* prediction of whether

the RSP2 protein, similar to the RSP protein, could be exported through a type IV secretion

system (Fig S1). To provide evidence supporting this hypothesis, we used strains SL1344 (R27

RSP2-Flag) and SL1344 (R27 ΔtrhC RSP2-Flag) and analyzed the presence of RSP2 protein in the

protein profile of the cell-free secreted fractions (secretome). This protein could not be detected

in the secretome of strain SL1344 (R27 ΔtrhC) (Fig. 3A). Immunodetection of RSP2-Flag by using

anti-Flag specific antibodies confirmed the requirement for TrhC expression for RSP2 export (Fig.

3B). We next checked whether RSP2 export in strain SL1344 (R27 ΔtrhC) could be made to occur

by providing the gene encoding the TrhC ATPase *in trans*, cloned in the plasmid pBR322 (plasmid

pBR322-*trhC*). Complementation of RSP2 export was observed (Fig. 3B), suggesting that, as was

the case for the RSP protein, the R27-encoded type IV secretion system mediates export of the

RSP2 protein.

We next addressed the question whether there is co-dependence in protein export between the

RSP and the RSP2 proteins. To that end, a Flag-tag was added to the *rsp2* gene in the strain

SL1344 (R27 Δ*rsp*). Expression of both RSP and RSP2 proteins was assessed in the strains SL1344

(R27 RSP2-Flag), SL1344 (R27 Δ*rsp* RSP2-Flag) and SL1344 (R27 Δ*rsp* RSP2-Flag pLG338-rsp). The

results obtained show that export of the RSP2 protein is dependent on the RSP expression (Fig.

4A and 4B).

Expression of the RSP2 protein influences the motility and conjugation of strain SL1344 (R27).

We previously showed that the expression of the RSP protein is essential for R27 plasmid

conjugation, and that SL1344 cells that express RSP show reduced motility compared to plasmid-

free cells [47]. Considering the observed interaction of both RSP and RSP2 proteins, we studied

whether, as is the case for the RSP protein, expression of the RSP2 protein influences cell motility

and/or conjugation.

After constructing an R27 derivative lacking the *rsp2* gene (plasmid R27 Δ*rsp2*), we performed a

comparative motility assay with the *Salmonella* strain SL1344 and its derivatives harboring the

R27, R27 Δ*rsp2*, and R27 Δ*rsp2* pLG338-*rsp2* plasmids. The results obtained (Fig.5A) showed that

the RSP2 protein influences the motility of strain SL1344 (R27).

We were able to show previously that the RSP protein is associated with the flagella synthesized

by the Salmonella strain SL1344 [47]. We decided next to assess whether RSP2 also targets the

flagella. In an attempt to detect the RSP2 protein by transmission electron microscopy, we used

strain SL1344 (R27 RSP2-Flag) and gold-labelled anti-Flag monoclonal antibodies (Fig 5B and 5C).

As was the case for RSP, RSP2 also exhibited binding to the flagella.

To assess the role of the RSP2 protein on plasmid conjugation, we constructed a R27 derivative

lacking the *rsp2* gene (plasmid R27 Δ*rsp2*) and compared the conjugation frequencies of strains

SL1344 (R27), SL1344 (R27 Δ*rsp2*) and SL1344 (R27 Δ*rsp2* pLG338-*rsp2*) growing at 25°C in liquid

media. We also compared the conjugation frequencies of strains SL1344 (R27) and SL1344 (R27

Δ*rsp2*) growing cells on nitrocellulose filters placed on LB plates. When cells were grown in liquid

medium, transfer of the R27 Δ*rsp2* plasmid was detected at a frequency that was approximately

two logs lower than that of wt R27 (Fig.5D). The presence *in trans* of the RSP2 protein encoded

by the pLG338-*rsp2* plasmid, restored the conjugation frequency of the wt R27 plasmid. When

cells were grown on solid medium, we also detected a significant decrease in the conjugation
frequency of the R27 Δ*rsp2* plasmid compared to that of the wt R27 plasmid (Fig. 5E).

Relationship between the RSP and RSP2 proteins and the conjugative machinery of the R27
plasmid.

As mentioned above, immunoprecipitation of the RSP protein indicated interactions with both
the RSP protein and the R27 *trhH* gene product. The TrhH protein of the plasmid R27 shares
identity with the TraH protein encoded by IncF plasmids [32]. The TraH protein is a component
of the outer membrane complex involved in conjugation [48] and has been shown to be required
for pilus assembly [49]. On the other hand, the *trhA* gene product encoded in IncHI plasmids has
been considered to be the pilin subunit itself [50]. We decided to analyze whether the
expression of the R27 TrhH and/or TrhA proteins encoded by IncHI plasmids influences the
expression of the RSP and RSP2 proteins. After inactivation of the *trhH* or *trhA* genes of strain
SL1344 (R27), the expression of the RSP and RSP2 proteins in the external medium was analyzed.
As both the RSP and RSP2 proteins were copurified with flagella when a conventional flagella
purification protocol was used (see the Material and Methods section), we obtained fractions
containing flagella from the SL1344 wt, SL1344 (R27), SL1344 (R27 Δ*trhA*) and SL1344 (R27
Δ*trhH*) isogenic derivatives. Proteins were analyzed by SDS-PAGE (Fig. 6A). The expression of
both the RSP and RSP2 proteins in the extracellular medium is dependent upon the function of
TrhH or TrhA. We also analyzed the intracellular expression of the RSP and RSP2 proteins in the
trhH and *trhA* genetic backgrounds. Lack of either TrhH (Fig. 6B and 6C) or TrhA (Fig. 6D and 6E)
function resulted in intracellular accumulation of either RSP or RSP2. Considering that
expression of the RSP and RSP2 proteins reduces cell motility in strain SL1344 (R27) and that
inactivation of the *trhH* gene interferes with the presence of these proteins in the extracellular
medium, it could be expected that *trhH* inactivation would also result in increased motility in
SL1344 cells harboring plasmid R27 Δ*trhH*. We then performed a mobility assay of the plasmid-
free SL1344 strain and of its SL1344 (R27) SL1344 (R27 Δ*rsp*)*,* SL1344 (R27 Δ*rsp2*) and SL1344
(R27 Δ*trhH*) derivatives. In accordance with (i) the observed requirement of TrhH function for
RSP and RSP2 export and (ii) the effect of the RSP/RSP2 proteins on cell motility, the motility of
strain SL1344 (R27 Δ*trhH*) was significantly reduced compared to that of the wt strain (Fig. 7).
The RSP and RSP2 proteins bind the R27 conjugative pili.

Taking into account the observed relationship between elements of the conjugation machinery
of the R27 plasmid and the RSP and RSP2 proteins, we decided to assess whether these proteins,
in addition of binding the flagella, they also target the conjugative pilus. To that end, we blocked

flagella expression and used electron microscopy on order to detect either the RSP or RSP2

proteins bound to the conjugative pili. To prevent flagella expression, we constructed Δ*fliC/fljB*

(flagellin subunit) derivatives of strains SL1344 (R27) and SL1344 (R27 RSP2-Flag). Immunogold

transmission electron microscopy imaging of these strains by using either polyclonal anti-RSP

antibodies [47] or monoclonal anti-Flag antibodies showed that in both examples, gold particles

were associated with tubular structures that likely corresponded to the conjugative pilus (Fig.

8A and 8C). When the Δ*trhH* allele was introduced into the mutant derivatives lacking flagella,

the tubular structures were no longer detected, thus supporting the hypothesis that they

corresponded to the conjugative pilus (Fig. 8B and 8D).

The RSP2 protein is specific for IncHI1 plasmids.

The RSP protein is restricted to the IncHI plasmids, from both the IncHI1 and IncHI2 subgroups

[47]. Upon having shown the relevant role of the RSP2 protein in the conjugation of the IncHI1

plasmid R27, we performed a BLAST search to identify the *rsp2* gene in other bacterial plasmids

(Table S2). In contrast to the RSP protein, the RSP2 protein is present in IncHI1 plasmids but not

in IncHI2 plasmids. Interestingly, a group of IncN plasmids also contains an homolog of the R27

rsp2 gene.

Distribution of proteins containing the Big domain among bacterial plasmids.

We next addressed the question of whether proteins containing Big domains are a feature of

only IncHI plasmids or whether they are also encoded by plasmids of other incompatibility

groups. We used for this analysis the genome viewer integrated into the NCBI database,

configuring it to show the features and domains of the annotated proteins. The search was

performed in assembled and sequenced plasmids of all incompatibility groups, selecting those

that presented proteins with annotated Big domains. Among all the plasmids analyzed, proteins

with these domains were also found in the plasmids of the IncA/C and IncP2 incompatibility

groups. Big proteins from IncA/C plasmids show a high degree of similarity (Fig. S2) and can be

mapped to the corresponding *tra* regions (Fig. S3 and Fig. S4). Big proteins from IncP2 plasmids

also show a very high degree of similarity (Fig. S5) and can be mapped close to a pilin gene (Fig.

S4).

The ALG87338.1 gene product of the IncA/C plasmid pKAZ3 is required for plasmid conjugation

at low temperature.

Upon having identified genes encoding proteins containing Big domains harbored on plasmids

different from those of the IncHI group, we also aimed to assess whether a protein containing a

Big domain encoded by a plasmid different from the IncHI group is also found in the secretome and influences conjugation. For this study, we selected the IncA/C plasmid pKAZ3, encoding the ALG87338.1 gene, the product of which contains a Big domain. The plasmid pKAZ3 was isolated from an antibiotic-contaminated lake and confers multiple-antibiotic resistance [51]. The ALG87338.1 gene product is an 1843 AA protein (200.75 kDa) that contains Big (3_2 and 3_3) and DUF4165 domains (Fig. S6). Plasmid pKAZ3 was first transferred to strain SL1344, and the secretomes of strains SL1344 and SL1344 (pKAZ3) were compared. A band corresponding to a protein of a molecular mass corresponding to the ALG87388.1 gene product was identified (Fig. 9A). This was confirmed by LC-MS/MS analysis of the protein excised from the SDS-PAGE gel. We next generated an ALG87388.1 knockout mutant derivative of plasmid pKAZ3. Considering that expression of either the RSP or RSP2 protein reduces SL1344 cell motility, we also compared the motility of strains SL1344, SL1344 (pKAZ3) and SL1344 (pKAZ3 ΔALG87388.1) (Fig. 9B). We observed that, while the acquisition of the pKAZ3 plasmid reduced cellular motility, there were no significant differences in motility between strains SL1344 (pKAZ3) and SL1344 (pKAZ3 ΔALG87388.1). We also compared the conjugation frequencies of strains SL1344 (pKAZ3) and SL1344 (pKAZ3 ΔALG87388.1), at both 37°C and at 25°C (Fig. 9C). ΔALG87388.1 did not influence the conjugation frequency at 37°C, but it had a very strong effect at 25°C.

Discussion

The conjugative process has been studied for decades, and a detailed picture of the molecular mechanism of conjugational DNA transfer is available (as reviewed in [50,52,53]). Plasmids such as the F factor, RP4, R388 or pTi have been studied as reference models to determine the function of several Tra proteins in plasmid conjugation by using genetic and biochemical approaches [50]. Comprehensive information is available on processes such as conjugative pilus biosynthesis, the establishment of donor-recipient cell contact or the assembly and activity of the relaxosome, which are key aspects of bacterial conjugation that must occur prior to the DNA transfer process. Nevertheless, several key questions, such as those regarding the function of several Tra proteins or the pilus´s ability to transport DNA between distant donor and recipient cells, remain to be answered [50]. In this report, we have elaborated on the role of plasmid-encoded high-molecular-weight Big proteins in the conjugation process of different plasmids.

In the IncHI plasmid R27, two genes that were mapped to the Tra2 and Tra1-Tra2 intergenic regions (*rsp* and *rsp2*, respectively) [32,33], encode proteins containing Big domains that participate in the conjugation process. Both the RSP and RSP2 proteins bind flagella, this resulting in a reduced cell motility. Export of the RSP2 protein to the external medium appears

to be dependent upon RSP export. This may explain that, whereas loss of RSP function

completely compensates for the effect of R27 acquisition on bacterial cell motility, loss of RSP2

function only shows a partial effect on cell motility.

The relationship between the RSP and RSP2 proteins, and the conjugative machinery of the R27

plasmid is shown in this work. Expression of key elements required for the synthesis of the

conjugative pilus of incHI1 plasmids (TrhH protein) and of the pilin subunit itself (TrhA protein)

is required for the correct translocation of the RSP and RSP2 proteins to the external surface of

the cells. In addition, imaging of cells lacking flagella showed short filaments that likely

corresponded to the conjugative pilus. These filaments were targeted both by the RSP and RSP2

proteins. Hence, there is an interaction between the RSP and RSP2 proteins and the R27

conjugative pilus.

IncHI plasmids and *Salmonella* participate in regulatory crosstalk that, is based on the expression

of both the tetracycline determinant and other plasmid-encoded genes [54,55]. One of the

effects of the crosstalk is the alteration in cellular motility, which is mediated by both plasmid-

borne regulators [56] and the binding of the RSP and RSP2 proteins to the flagellar structure

([47], this work). Nevertheless, these proteins also play a second and critical role in IncHI plasmid

conjugation. They also bind the conjugative pilus, which is likely required to facilitate

transmission of the conjugative plasmid. Plasmid-encoded Big proteins binding to the flagella

and thus reducing motility may favor cell-to-cell contact and hence conjugation. Binding to the

conjugative pilus may contribute to the stabilization of the pilus structure. The overall effect of

the expression of these proteins is to significantly favor the transfer of the plasmid that encode

them.

Genes encoding proteins containing Big domains are not restricted to IncHI plasmids. The

genomic analysis performed in this work has shown that such proteins are also present in IncA/C

and IncP2 plasmids. Both groups of plasmids have a wide host range. IncA/C plasmids are also

key players in the dissemination of AMR in *Enterobacteriaceae* and other Gram-negative

microorganisms. IncP2 plasmids are high-molecular-weight plasmids that are prevalent in

Pseudomonas and contribute to the dissemination of AMR within this genus [57,58]. As the

search that we performed had some limitations, it cannot be ruled out that other groups of

plasmids also express Big proteins. By using the genome viewer integrated into the NCBI

database, only those plasmids that were well-annotated and characterized were considered. In

addition, proteins that contain Big domains were annotated because of the confirmation that

this domain, was present, not because they shared amino acid sequence similarity. Notably, the

RSP2 protein is specific of a subgroup of IncHI plasmids, incHI1. IncHI2 plasmids that encode the
rsp gene lack the *rsp2* gene. Interestingly, our BLAST analysis also showed that the *rsp2* gene has
jumped to plasmids of a different Inc group, IncN. This may indicate that the *rsp2* genes are
spreading and may also influence conjugation in these plasmids.

The identified high-molecular-weight Big proteins encoded by IncA/C plasmids show a very high
degree of homology, which suggests that they play similar roles in these plasmids. The genes
encoding these proteins were also mapped to the *tra* region *(traV/A/W/F/N genes)* and, in some
instances, were adjacent to the pilin genes. A recent study focused on performing a
comprehensive analysis of IncC plasmid conjugation [59], and the gene encoding the identified
Big protein encoded by the pMS6198A plasmid (the product of the gene MS6198_A094) was
considered not to influence conjugation, despite mapping within the Tra1 region, between the
*dsbC*A and *traL* genes. The likely reason for this was that the conjugation experiments were
performed only at 37°C. We showed here that the ALG87338.1 gene product of the IncA/C
plasmid pKAZ3 positively influenced conjugation at 25°C (i.e., at environmental temperature).
IncHI and IncA/C plasmids belong to the same relaxase family (MOB_H). Interestingly, they also
share the requirement for a high-molecular-weight Big protein for efficient conjugation at
environmental temperatures. The identified Big proteins of IncP2 plasmids also show a high
degree of similarity. While the molecular mass of these proteins is lower than that of the Big
proteins of IncHI and IncA/C plasmids, they map close to a pilin, which also suggests that they
may play a role in the conjugative process.

Out of the IncHI plasmids, which have classically been characterized as being conjugative only
at temperatures below 30°C, plasmids harbored by *Enterobacteriaceae* have usually been
studied in cells growing at the optimal growth temperature for these microorganisms, that is,
37°C. Our study also highlights the importance of studying plasmid conjugation at other
temperatures, closer to those that most microorganisms encounter outside their warm-blooded
hosts. Indeed, AMR transmission occurs mainly in natural environments at temperatures that
rarely reach 37°C.

The relevance of the Big proteins studied in this work is not only their role as elements favoring
plasmid conjugation but also the fact that, as they are associated with extracellular appendages
of the bacterial cells, they can be targeted with specific antibodies either to restrict the
dissemination of the plasmids that encode them or, when expressed within the human body, to
control infections caused by bacteria that express the plasmid that encode these proteins.

Materials and Methods

Bacterial strains, plasmids, and growth conditions.

The bacterial strains (Table S3) were routinely grown in Luria-Bertani (LB) medium (10 g l^{-1} NaCl, 10 g l^{-1} tryptone and 5 g l^{-1} yeast extract) with vigorous shaking at 200 rpm (Innova 3100, New Brunswick Scientific). The antibiotics used were chloramphenicol (Cm) (25 µg ml^{-1}), tetracycline (Tc) (15 µg ml^{-1}), carbenicillin (Cb) (100 µg ml^{-1}) and kanamycin (Km) (50 µg ml^{-1}) (Sigma-Aldrich).

Oligonucleotides.

The oligonucleotides (from 5' to 3') used in this work are listed in Table S4.

Genetic manipulation.

All enzymes used to perform standard molecular and genetic procedures were used according to the manufacturer's recommendations. To introduce plasmids into *E. coli* and *Salmonella*, bacterial cells were grown until an OD$_{600\ nm}$ of 0.6. Cells were then washed several times with 10% glycerol, and the respective plasmids or DNA was electroporated by using an Eppendorf gene pulser (Electroporator 2510).

Deletions of the *rsp2* (ORF *R0055), fliC, fljB, trhH* and *trhA* genes were performed in strain SL1344 (R27) by using the λ Red recombination method, as previously described [60]. The antibiotic resistance determinant of the plasmids pKD3/pKD4 was amplified using the corresponding oligonucleotides (P1/P2 series, see Table S4). The mutants were confirmed by PCR using the corresponding oligonucleotides (P1up/P2down series, see Table S4). We used phage P22 HT for combining mutations by transduction [61]. When necessary, the antibiotic resistance cassette was eliminated using the FLP/FRT-mediated site-specific recombination method as previously described [62].

For deletion of the ALG87338.1 gene from the pKAZ3 plasmid, we first constructed the plasmid pKD46-Km^R following the strategy described in [63]. Briefly, kanamycin resistance from the pKD4 plasmid was amplified using the PkmXmnIFW/PkmXmnIRv oligonucleotides together with Phusion Hot Start II High-Fidelity DNA Polymerase (Thermo Scientific) following the manufacturer's recommendations. The corresponding XmnI-flanked kanamycin resistance fragment was cloned into the vector pKD46 previously digested with the same enzyme. The resulting plasmid was termed pKD46-Km^R. Deletion of ALG87338.1 was performed in strain SL1344 (pKAZ3) pKD46-Km^R by using the λ Red recombination method. The antibiotic resistance determinant of the plasmid pKD3 was amplified using the corresponding oligonucleotides

(P1/P2 series, see Table S4). The mutants were confirmed by PCR using the corresponding

oligonucleotides (P1up/P2down series, see Table S4).

Recombinational transfer of the Flag sequence into the *rsp2* gene was achieved by following a

previously described methodology [64]. The template vector encoding Flag and Km^r used was

the pSUB11 plasmid. The primers used for construction of the Flag-tagged derivative were

R27_p0553XP1 and R27_p0553XP2 (Table S4). The correct insertion of the Flag-tag was

confirmed by PCR using oligonucleotides R27_p0553XP1UP and R27_p0553XP2DOWN (Table

S4).

To construct the plasmid pLG338-*rsp2*, the ORF *R0055* (*rsp2*) (GenBank accession number

AF250878.1, position 56639-59970) was amplified using the oligonucleotides R27_R55 pLG322

ECORI fw/R27_R55 pLG322 Bam rv (see Table S4 for the sequences) together with Phusion Hot

Start II High-Fidelity DNA Polymerase (Thermo Scientific) following the manufacturer's

recommendations. The corresponding EcoRI/BamHI fragment was cloned into the vector

pLG338-30 previously digested with the same enzymes. The resulting plasmids were Sanger

sequenced and termed pLG338-*rsp2*.

Plasmid conjugation.

The R27 and pKAZ3 plasmids were conjugated as described previously [47]. The mating

frequency was calculated as the number of transconjugants per donor cell. Student´s *t*-test was

used to determine statistical significance, and the values were obtained by using GraphPad

Prism 8 software. A *P* value of less than 0.05 was considered significant.

Motility assays.

The motility assay was performed as previously described [47]. The experiments were repeated

three times with three plates of each strain in each experiment. The colony diameter was

measured and plotted, and standard errors were calculated. Student´s t-test was used to

determine statistical significance, and the values were obtained by using the GraphPad Prism 8

software. A P value of less than 0.05 was considered significant.

Flagellum Isolation.

Flagellum isolation was prepared as previously described [47].

Cell-free secreted proteins (secretome).

Cell-free supernatants were prepared as previously described [47].

Cell fractionation.

Cell fractionation was performed as previously described [47].

Immunogold electron microscopy.

Immunogold microscopy experiments were performed as previously described [47].

Electrophoresis and Western blotting analysis of proteins.

Protein samples were analyzed by 10% and 12.5% SDS-PAGE [65]. Proteins were transferred from the gels to PVDF membranes using the Trans-Blot Turbo system (Bio-Rad). Western blot analysis was performed with a monoclonal antibody raised against the Flag-epitope (Sigma) diluted 1:10.000 in a solution containing PBS, 0.2% Triton, and 3% skim milk and incubated for 16 hours at 4°C. The membranes were washed for 20 minutes each with PBS and 0.2% Triton solution. The washing step was repeated three times. Thereafter, the membranes were incubated with horseradish peroxidase-conjugated goat anti-mouse IgG (Invitrogen) diluted 1:5.000 in a solution containing PBS and 0.2% Triton for 1 hour at room temperature. Again, three washing steps of 45 minutes with PBS and 0.2% Triton solution were performed, and detection was performed by enhanced chemiluminescence using ImageQuant LAS 54000 imaging system software (GE Healthcare Lifesciences).

RSP-Flag Immunoprecipitation.

For RSP-Flag protein immunoprecipitation, the strains SL1344 (R27 RSP-Flag) and SL1344 (R27 Δrsp, negative control) were grown in LB medium for 16 hours at 37ºC. One hundred milliliters of fresh LB medium was inoculated 1:100 with both overnight-cultured strains and grown at 25°C until an $OD_{600\,nm}$ of 2.0 was reached. Then, the cells were centrifuged at 9.000 rpm for 30 minutes at 4°C, the pellets were discarded, and the supernatants were filtered through 0.22 µm filters. For each immunoprecipitation protocol, we used 100 µl of Anti-Flag M2 Affinity Gel (Sigma-Aldrich) and 100 ml of each supernatant and incubated the mixture under slow rotation at 4°C for 16 hours. Each supernatant was loaded onto a Poly-Prep chromatography column (Bio-Rad), and the flowthrough was stored at -20°C for further analysis. Each column was washed out with 100 ml of washing buffer (50 mM Tris-HCl pH 7.5, 150 mM NaCl), and elution was performed 3 times with 0.3 ml of elution buffer (0.1M glycine pH 3.5). Elution fractions were concentrated using a trichloroacetic acid (TCA) precipitation protocol, briefly, 1 ml of the samples was mixed with 0.5 ml of a 45% TCA solution (w/v). The samples were kept on ice for 30 minutes and centrifugated for 30 minutes at 13,400 rpm. The supernatants were carefully

discarded, acetone was added to the protein pellets, and the pellets were again centrifugated

for 30 minutes at 13,400 rpm. The supernatants were carefully discarded. The protein pellets

were dried and solubilized with 1x Laemmli Sample Buffer (Bio-Rad). Samples were boiled for 10

minutes and loaded onto a 12.5% SDS-PAGE gel. When the protein samples entered the stacking

phase, the gel run was stopped, and samples were excised from the gel and sent to the

Proteomic Platform (Barcelona Science Park, Barcelona, Spain) for protein identification, as

previously described [55]. Immunoprecipitation and protein identification experiments were

repeated twice.

Acknowledgments and funding sources

451 This work was supported by grants from Fundació "La Marató TV3", Spain (Project ref. 201818

452 10), and PID2019-107479RB-I00 (AEI/FEDER, UE) from the Ministerio de Economía, Industria y

453 Competitividad, and CERCA Programme/Generalitat de Catalunya to A.J. Authors wish to thank

454 Dr. Álvaro San Millán (CNB-CSIC, Madrid) who kindly provided the pKAZ3 plasmid.

References

1. Morens DM, Folkers GK, Fauci AS. The challenge of emerging and re-emerging infectious diseases. Nature. Nature Publishing Group; 2004. pp. 242–249. doi:10.1038/nature02759

2. Meyer E, Schwab F, Schroeren-Boersch B, Gastmeier P. Dramatic increase of third-generation cephalosporin-resistant *E. coli* in German intensive care units: Secular trends in antibiotic drug use and bacterial resistance, 2001 to 2008. Critical Care. 2010;14. doi:10.1186/cc9062

3. Rossolini GM, Mantengoli E, Docquier J-D, Musmanno RA. Epidemiology of infections caused by multiresistant Gram-negatives: ESBLs, MBLs, panresistant strains. NEW MICROBIOLOGICA. 2007.

4. Spellberg B, Guidos R, Gilbert D, Bradley J, Boucher HW, Scheld WM, et al. The epidemic of antibiotic-resistant infections: A call to action for the medical community from the infectious diseases society of America. Clinical Infectious Diseases. Clin Infect Dis; 2008. pp. 155–164. doi:10.1086/524891

5. Carattoli A. Plasmids and the spread of resistance. International Journal of Medical Microbiology. Int J Med Microbiol; 2013. pp. 298–304. doi:10.1016/j.ijmm.2013.02.001

6. Wang J, Stephan R, Zurfluh K, Hächler H, Fanning S. Characterization of the genetic environment of *blaESBL* genes, integrons and toxin-antitoxin systems identified on large transferrable plasmids in multi-drug resistant *Escherichia coli*. Frontiers in Microbiology. 2015;6. doi:10.3389/fmicb.2014.00716

7. Thomas CM, Nielsen KM. Mechanisms of, and barriers to, horizontal gene transfer between bacteria. Nature Reviews Microbiology. Nat Rev Microbiol; 2005. pp. 711–721. doi:10.1038/nrmicro1234

8. Vrancianu CO, Popa LI, Bleotu C, Chifiriuc MC. Targeting Plasmids to Limit Acquisition and Transmission of Antimicrobial Resistance. Frontiers in Microbiology. Frontiers Media S.A.; 2020. doi:10.3389/fmicb.2020.00761

9. Carattoli A. Resistance plasmid families in Enterobacteriaceae. Antimicrobial Agents and Chemotherapy. Antimicrob Agents Chemother; 2009. pp. 2227–2238. doi:10.1128/AAC.01707-08

10. Novick RP. Plasmid incompatibility. Microbiological Reviews. American Society for Microbiology (ASM); 1987. pp. 381–395. doi:10.1128/mmbr.51.4.381-395.1987

11. Garcillán-Barcia MP, Francia MV, de La Cruz F. The diversity of conjugative relaxases and its application in plasmid classification. FEMS Microbiology Reviews. 2009;33: 657–687. doi:10.1111/j.1574-6976.2009.00168.x

491   12.   Phan MD, Wain J. IncHI plasmids, a dynamic link between resistance and pathogenicity.
492   Journal of infection in developing countries. J Infect Dev Ctries; 2008. pp. 272–278.
493   doi:10.3855/jidc.221

494   13.   Gilmour MW, Thomson NR, Sanders M, Parkhill J, Taylor DE. The complete nucleotide
495   sequence of the resistance plasmid R478: Defining the backbone components of
496   incompatibility group H conjugative plasmids through comparative genomics. Plasmid.
497   2004;52: 182–202. doi:10.1016/j.plasmid.2004.06.006

498   14.   Forde BM, Zowawi HM, Harris PNA, Roberts L, Ibrahim E, Shaikh N, et al. Discovery of
499   *mcr-1* -Mediated Colistin Resistance in a Highly Virulent *Escherichia coli* Lineage.
500   mSphere. 2018;3. doi:10.1128/msphere.00486-18

501   15.   Villa L, Poirel L, Nordmann P, Carta C, Carattoli A. Complete sequencing of an IncH
502   plasmid carrying the *bla ndm-1*, *bla ctx-m-15* and *qnrB1* genes. Journal of Antimicrobial
503   Chemotherapy. 2012;67: 1645–1650. doi:10.1093/jac/dks114

504   16.   Dolejska M, Villa L, Poirel L, Nordmann P, Carattoli A. Complete sequencing of an IncHI1
505   plasmid encoding the carbapenemase NDM-1, the ArmA 16S RNA methylase and a
506   resistance-nodulation-cell division/multidrug efflux pump. Journal of Antimicrobial
507   Chemotherapy. 2013;68: 34–39. doi:10.1093/jac/dks357

508   17.   Holt KE, Phan MD, Baker S, Duy PT, Nga TVT, Nair S, et al. Emergence of a Globally
509   Dominant IncHI1 Plasmid Type Associated with Multiple Drug Resistant Typhoid. Ryan
510   ET, editor. PLoS Neglected Tropical Diseases. 2011;5: e1245.
511   doi:10.1371/journal.pntd.0001245

512   18.   Lim LM, Ly N, Anderson D, Yang JC, Macander L, Jarkowski A, et al. Resurgence of colistin:
513   A review of resistance, toxicity, pharmacodynamics, and dosing. Pharmacotherapy.
514   Pharmacotherapy; 2010. pp. 1279–1291. doi:10.1592/phco.30.12.1279

515   19.   Catry B, Cavaleri M, Baptiste K, Grave K, Grein K, Holm A, et al. Use of colistin-containing
516   products within the European Union and European Economic Area (EU/EEA):
517   development of resistance in animals and possible impact on human and animal health.
518   International Journal of Antimicrobial Agents. Elsevier B.V.; 2015. pp. 297–306.
519   doi:10.1016/j.ijantimicag.2015.06.005

520   20.   Olaitan AO, Morand S, Rolain JM. Mechanisms of polymyxin resistance: Acquired and
521   intrinsic resistance in bacteria. Frontiers in Microbiology. Frontiers Media S.A.; 2014.
522   doi:10.3389/fmicb.2014.00643

523   21.   Liu YY, Wang Y, Walsh TR, Yi LX, Zhang R, Spencer J, et al. Emergence of plasmid-mediated
524   colistin resistance mechanism MCR-1 in animals and human beings in China: A
525   microbiological and molecular biological study. The Lancet Infectious Diseases. 2016;16:
526   161–168. doi:10.1016/S1473-3099(15)00424-7

527   22.   Gao R, Hu Y, Li Z, Sun J, Wang Q, Lin J, et al. Dissemination and Mechanism for the MCR-
528   1 Colistin Resistance. PLoS Pathogens. 2016;12. doi:10.1371/journal.ppat.1005957

529   23.   Xavier BB, Lammens C, Ruhal R, Malhotra-Kumar S, Butaye P, Goossens H, et al.
530   Identification of a novel plasmid-mediated colistinresistance gene, *mcr-2*, in *Escherichia*
531   *coli*, Belgium, june 2016. Eurosurveillance. 2016;21. doi:10.2807/1560-
532   7917.ES.2016.21.27.30280

533   24.   Borowiak M, Fischer J, Hammerl JA, Hendriksen RS, Szabo I, Malorny B. Identification of
534   a novel transposon-associated phosphoethanolamine transferase gene, *mcr-5*,
535   conferring colistin resistance in d-tartrate fermenting *Salmonella enterica* subsp. enterica
536   serovar Paratyphi B. Journal of Antimicrobial Chemotherapy. 2017;72: 3317–3324.
537   doi:10.1093/jac/dkx327

538   25.   Carattoli A, Villa L, Feudi C, Curcio L, Orsini S, Luppi A, et al. Novel plasmid-mediated
539   colistin resistance *mcr-4* gene in *Salmonella* and *Escherichia coli*, Italy 2013, Spain and
540   Belgium, 2015 to 2016. Eurosurveillance. 2017;22. doi:10.2807/1560-
541   7917.ES.2017.22.31.30589

542   26.   Yin W, Li H, Shen Y, Liu Z, Wang S, Shen Z, et al. Novel Plasmid-Mediated Colistin
543   Resistance Gene *mcr-3* in *Escherichia coli* Downloaded from. 2021;8: 543–560.
544   doi:10.1128/mBio

545   27.   Matamoros S, Van Hattem JM, Arcilla MS, Willemse N, Melles DC, Penders J, et al. Global
546   phylogenetic analysis of *Escherichia coli* and plasmids carrying the *mcr-1* gene indicates
547   bacterial diversity but plasmid restriction. Scientific Reports. 2017;7: 1–9.
548   doi:10.1038/s41598-017-15539-7

549   28.   Wang Y, Tian GB, Zhang R, Shen Y, Tyrrell JM, Huang X, et al. Prevalence, risk factors,
550   outcomes, and molecular epidemiology of *mcr-1*-positive Enterobacteriaceae in patients
551   and healthy adults from China: an epidemiological and clinical study. The Lancet
552   Infectious Diseases. 2017;17: 390–399. doi:10.1016/S1473-3099(16)30527-8

553   29.   Zheng B, Dong H, Xu H, Lv J, Zhang J, Jiang X, et al. Coexistence of MCR-1 and NDM-1 in
554   Clinical *Escherichia coli* Isolates: Clinical Infectious Diseases. 2016;63: 1393–1395.
555   doi:10.1093/cid/ciw553

556   30.   Maher D, Sherburne R, Taylor DE. H-pilus assembly kinetics determined by electron
557   microscopy. Journal of Bacteriology. 1993;175: 2175–2183. doi:10.1128/JB.175.8.2175-
558   2183.1993

559   31.   Alonso G, Baptista K, Ngo T, Taylor DE. Transcriptional organization of the temperature-
560   sensitive transfer system from the IncHI1 plasmid R27. Microbiology. 2005;151: 3563–
561   3573. doi:10.1099/mic.0.28256-0

562   32.   Lawley TD, Gilmour MW, Gunton JE, Standeven LJ, Taylor DE. Functional and mutational
563   analysis of conjugative transfer region 1 (Tra1) from the IncHI1 plasmid R27. Journal of
564   Bacteriology. 2002;184: 2173–2180. doi:10.1128/JB.184.8.2173-2180.2002

33. Lawley TD, Gilmour MW, Gunton JE, Tracz DM, Taylor DE. Functional and mutational
analysis of conjugative transfer region 2 (Tra2) from the IncHI1 plasmid R27. Journal of
Bacteriology. 2003;185: 581–591. doi:10.1128/JB.185.2.581-591.2003

34. Sherburne CK, Lawley TD, Gilmour MW, Blattner FR, Burland V, Grotbeck E, et al. The
complete DNA sequence and analysis of R27, a large IncHI plasmid from *Salmonella* typhi
that is temperature sensitive for transfer. Nucleic Acids Research. 2000;28: 2177–2186.
doi:10.1093/nar/28.10.2177

35. Aoki T, Egusa S, Kimura T, Watanabe T. Detection of R factors in naturally occurring
Aeromonas salmonicida strains. Applied microbiology. 1971;22: 716–717.
doi:10.1128/aem.22.4.716-717.1971

36. Watanabe T, Aoki T, Ogata Y, Egusa S. R FACTORS RELATED TO FISH CULTURING. Annals
of the New York Academy of Sciences. 1971;182: 383–410. doi:10.1111/j.1749-
6632.1971.tb30674.x

37. Welch TJ, Fricke WF, McDermott PF, White DG, Rosso ML, Rasko DA, et al. Multiple
antimicrobial resistance in plague: An emerging public health risk. PLoS ONE. 2007;2.
doi:10.1371/journal.pone.0000309

38. Fricke WF, Welch TJ, McDermott PF, Mammel MK, LeClerc JE, White DG, et al.
Comparative genomics of the IncA/C multidrug resistance plasmid family. Journal of
Bacteriology. 2009;191: 4750–4757. doi:10.1128/JB.00189-09

39. Call DR, Singer RS, Meng D, Broschat SL, Orfe LH, Anderson JM, et al. *blaCMY-2*-positive
IncA/C plasmids from *Escherichia coli* and *Salmonella enterica* are a distinct component
of a larger lineage of plasmids. Antimicrobial Agents and Chemotherapy. 2010;54: 590–
596. doi:10.1128/AAC.00055-09

40. Fernández-Alarcón C, Singer RS, Johnson TJ. Comparative genomics of multidrug
resistance-encoding incA/C plasmids from commensal and pathogenic *Escherichia coli*
from multiple animal sources. PLoS ONE. 2011;6. doi:10.1371/journal.pone.0023415

41. Suzuki H, Yano H, Brown CJ, Top EM. Predicting plasmid promiscuity based on genomic
signature. Journal of Bacteriology. 2010;192: 6045–6055. doi:10.1128/JB.00277-10

42. Eda R, Nakamura M, Takayama Y, Maehana S, Nakano R, Yano H, et al. Trends and
molecular characteristics of carbapenemase-producing Enterobacteriaceae in Japanese
hospital from 2006 to 2015. Journal of Infection and Chemotherapy. 2020;26: 667–671.
doi:10.1016/j.jiac.2020.02.002

43. Rozwandowicz M, Brouwer MSM, Fischer J, Wagenaar JA, Gonzalez-Zorn B, Guerra B, et
al. Plasmids carrying antimicrobial resistance genes in Enterobacteriaceae. Journal of
Antimicrobial Chemotherapy. 2018;73: 1121–1137. doi:10.1093/jac/dkx488

44. Qamar MU, Ejaz H, Walsh TR, Shah AA, al Farraj DA, Alkufeidy RM, et al. Clonal
relatedness and plasmid profiling of extensively drug-resistant New Delhi metallo-β-

lactamase-producing *Klebsiella pneumoniae* clinical isolates. Future Microbiology.
2021;16: 229–239. doi:10.2217/fmb-2020-0315

45.   Halaby DM, Mornon JPE. The immunoglobulin superfamily: An insight on its tissular,
species, and functional diversity. Journal of Molecular Evolution. 1998;46: 389–400.
doi:10.1007/PL00006318

46.   Bodelón G, Palomino C, Fernández LA. Immunoglobulin domains in *Escherichia coli* and
other enterobacteria: From pathogenesis to applications in antibody technologies. FEMS
Microbiology Reviews. Oxford Academic; 2013. pp. 204–250. doi:10.1111/j.1574-
6976.2012.00347.x

47.   Hüttener M, Prieto A, Aznar S, Bernabeu M, Glaría E, Valledor AF, et al. Expression of a
novel class of bacterial Ig-like proteins is required for IncHI plasmid conjugation. PLoS
Genetics. 2019;15: e1008399. doi:10.1371/journal.pgen.1008399

48.   Arutyunov D, Arenson B, Manchak J, Frost LS. F plasmid TraF and TraH are components
of an outer membrane complex involved in conjugation. Journal of Bacteriology.
2010;192: 1730–1734. doi:10.1128/JB.00726-09

49.   Frost LS, Ippen-Ihler K, Skurray RA. Analysis of the sequence and gene products of the
transfer region of the F sex factor. Microbiology and Molecular Biology Reviews. 1994;58.

50.   Virolle C, Goldlust K, Djermoun S, Bigot S, Lesterlin C. Plasmid transfer by conjugation in
gram-negative bacteria: From the cellular to the community level. Genes. MDPI AG;
2020. pp. 1–33. doi:10.3390/genes11111239

51.   Flach CF, Johnning A, Nilsson I, Smalla K, Kristiansson E, Larsson DGJ. Isolation of novel
IncA/C and IncN fluoroquinolone resistance plasmids from an antibiotic-polluted lake.
Journal of Antimicrobial Chemotherapy. 2015;70: 2709–2717. doi:10.1093/jac/dkv167

52.   Smillie C, Garcillán-Barcia MP, Francia MV, Rocha EPC, de la Cruz F. Mobility of Plasmids.
Microbiology   and   Molecular   Biology   Reviews.   2010;74:   434–452.
doi:10.1128/mmbr.00020-10

53.   Cabezón E, Ripoll-Rozada J, Peña A, de la Cruz F, Arechaga I. Towards an integrated model
of bacterial conjugation. FEMS Microbiology Reviews. Oxford University Press; 2015. pp.
81–95. doi:10.1111/1574-6976.12085

54.   Paytubi S, Aznar S, Madrid C, Balsalobre C, Dillon SC, Dorman CJ, et al. A novel role for
antibiotic resistance plasmids in facilitating *Salmonella* adaptation to non-host
environments. Environmental Microbiology. 2014;16: 950–962. doi:10.1111/1462-
2920.12244

55.   Hüttener M, Prieto A, Aznar S, Dietrich M, Paytubi S, Juárez A. Tetracycline alters gene
expression in *Salmonella* strains that harbor the Tn10 transposon. Environmental
Microbiology Reports. 2018;10: 202–209. doi:10.1111/1758-2229.12621

56.   Luque A, Paytubi S, Sánchez-Montejo J, Gibert M, Balsalobre C, Madrid C. Crosstalk
639   between bacterial conjugation and motility is mediated by plasmid-borne regulators.
640   Environmental Microbiology Reports. 2019;11: 708–717. doi:10.1111/1758-2229.12784

641   57.   Jacoby GA, Sutton L, Knobel L, Mammen P. Properties of IncP-2 plasmids of *Pseudomonas*
642   spp.   Antimicrobial   Agents   and   Chemotherapy.   1983;24:   168–175.
643   doi:10.1128/AAC.24.2.168

644   58.   Cazares A, Moore MP, Hall JPJ, Wright LL, Grimes M, Emond-Rhéault JG, et al. A
645   megaplasmid family driving dissemination of multidrug resistance in *Pseudomonas*.
646   Nature Communications. 2020;11: 1–13. doi:10.1038/s41467-020-15081-7

647   59.   Hancock SJ, Phan MD, Luo Z, Lo AW, Peters KM, Nhu NTK, et al. Comprehensive analysis
648   of IncC plasmid conjugation identifies a crucial role for the transcriptional regulator AcaB.
649   Nature Microbiology. 2020;5: 1340–1348. doi:10.1038/s41564-020-0775-0

650   60.   Datsenko KA, Wanner BL. One-step inactivation of chromosomal genes in *Escherichia coli*
651   K-12 using PCR products. Proceedings of the National Academy of Sciences of the United
652   States of America. 2000;97: 6640–6645. doi:10.1073/pnas.120163297

653   61.   Sternberg NL, Maurer R. Bacteriophage-Mediated Generalized Transduction in
654   *Escherichia coli* and *Salmonella* typhimurium. Methods in Enzymology. 1991;204: 18–43.
655   doi:10.1016/0076-6879(91)04004-8

656   62.   Cherepanov PP, Wackernagel W. Gene disruption in *Escherichia coli*: TcR and KmR
657   cassettes with the option of Flp-catalyzed excision of the antibiotic-resistance
658   determinant. Gene. 1995;158: 9–14. doi:10.1016/0378-1119(95)00193-A

659   63.   Doublet B, Douard G, Targant H, Meunier D, Madec JY, Cloeckaert A. Antibiotic marker
660   modifications of λ Red and FLP helper plasmids, pKD46 and pCP20, for inactivation of
661   chromosomal genes using PCR products in multidrug-resistant strains. Journal of
662   Microbiological Methods. 2008;75: 359–361. doi:10.1016/j.mimet.2008.06.010

663   64.   Uzzau S, Figueroa-Bossi N, Rubino S, Bossi L. Epitope tagging of chromosomal genes in
664   *Salmonella*. Proceedings of the National Academy of Sciences of the United States of
665   America. 2001;98: 15264–15269. doi:10.1073/pnas.261348198

666   65.   Sambrook J and Russell D. Molecular Cloning: A Laboratory Manual. 3rd ed. NYC: Cold
667   Spring Harbor Laboratory Press; 2001.

 Figures

674   **Figure 1.** Identification of the RSP2 protein. (A) Genetic map of the R27 plasmid region where both the *rsp* and *rsp2* (*R0055*) genes were mapped. (B)

675   Comparison of the Big domains of the RSP and RSP2 proteins. (C) Detection of the RSP2 protein in the cell-free secretome of the SL1344 (R27) strain. Arrows

676   point to the bands corresponding to the RSP and RSP2 proteins, the latter of which was confirmed by LC-MS/MS analysis.

Figure 2. Immunodetection of the RSP2-Flag protein in different cellular compartments. (A) Coomassie blue staining of the different cellular fractions obtained from strains SL11344 wt, SL1344 (R27 RSP-Flag) and SL1344 (R27 RSP2-Flag). (B) Immunodetection of the RSP-Flag and RSP2-Flag proteins in the periplasm, cytoplasm, and outer and inner membrane fractions of strains SL1344 (R27 RSP-Flag) and SL1344 (R27 RSP2-Flag), respectively. (C) Immunodetection of the RSP2-Flag protein in the cell-free secretome of strain SL1344 (R27 RSP2-Flag) grown at 25ºC until OD$_{600\,nm}$ of 2.0.

1 – R27 RSP2-Flag
2 – R27 ΔtrhC RSP2-Flag pBR322
3 – R27 ΔtrhC RSP2-Flag pBR322-trhC
4 – R27 ΔtrhC RSP-Flag pBR322
5 – R27 ΔtrhC RSP-Flag pBR322-trhC

Figure 3. RSP2 export requires the type IV secretion system encoded by the R27 plasmid. (A)

Coomassie blue staining for SDS-PAGE analysis of the cell-free secretome of the strains SL1344

(R27 RSP2-Flag), SL1344 (R27 ΔtrhC RSP2-Flag pBR322), SL1344 (R27 ΔtrhC RSP2-Flag pBR322-

trhC), (R27 ΔtrhC RSP-Flag pBR322) and (R27 ΔtrhC RSP-Flag pBR322-trhC) grown at 25ºC until

OD$_{600\ nm}$ of 2.0. (B) Immunodetection of the RSP-Flag and RSP2-Flag proteins with anti-Flag

antibodies. Arrows point to the RSP-Flag and RSP2-Flag proteins. The experiment was repeated

three times. A representative experiment is shown.

– R27 RSP2-Flag
– R27 Δ*rsp* RSP2-Flag
– R27 Δ*rsp* RSP2-Flag pLG338-*rsp*

Figure 4. RSP2 export is dependent on the expression of the RSP protein. (A) Coomassie blue staining for SDS-PAGE analysis of the cell-free secretomes of the strains SL1344 (R27 RSP2-Flag), SL1344 (R27 Δ*rsp* RSP2-Flag) and SL1344 (R27 Δ*rsp* RSP2-Flag pLG338-*rsp*) grown at 25ºC until $OD_{600\ nm}$ of 2.0. (B) Immunodetection of the RSP2-Flag protein with anti-Flag antibodies. RSP2-Flag protein is indicated. The experiment was repeated three times. A representative experiment is shown.

Figure 5. The RSP2 protein influences the motility of strain SL1344 (R27) and the conjugation of

the R27 plasmid. (A) Effect of the *rsp2* allele on the motility of strain SL1344 (R27). The results

are the means of three independent experiments. Standard deviations are shown. Statistical

analysis showed significant differences, analyzed by unpaired two-sided Student´s t-test (*P-

value < 0.0001). (B and C) The RSP2 protein binds to the flagella of strain SL1344 (R27).

Immunogold electron microscopy of cells from strains SL1344 (R27 RSP2-Flag) (B) and SL1344

(R27 Δ*rsp2*) (C) using monoclonal anti-Flag antibodies and goat anti-mouse IgG conjugated to 12

nm gold particles. Arrows point to the RSP protein associated with the flagella in (B) and only to

the flagella in (C). Bars represent 2 μm. (D and E) Effect of the *rsp2* allele on the conjugation

frequency of the R27 plasmid in liquid (D) and solid (E) media, respectively. The data shown are

the means and standard deviations of three independent experiments. Statistical analysis

showed significant differences, analyzed by unpaired two-sided Student´s t-test (D) (*P-value

0.0082, **P-value 0.0095) and (E) (**P-value 0.0428).

Figure 6. Expression of the RSP and RSP2 proteins depends on the R27 TrhA and TrhH functions. (A) SDS-PAGE analysis of the purified flagellar fractions of the plasmid-free strain SL1344 and strain SL1344 harboring (R27) and the (R27 ΔtrhA) and (R27 ΔtrhH) derivatives. (B) SDS-PAGE analysis of the cell extract and cell-free secretome of the strains SL1344 (R27 ΔtrhH), SL1344 (R27 ΔtrhH RSP-Flag) and SL1344 (R27 ΔtrhH RSP2-Flag), respectively. (C) Immunodetection of RSP-Flag and RSP2-Flag in the intracellular compartments of strain SL1344 harboring the corresponding *trhH* derivatives of the R27 plasmid. (D) SDS-PAGE analysis of the cell extract and cell-free secretome of the strains SL1344 (R27 RSP-Flag), SL1344 (R27 RSP2-Flag), SL1344 (R27 ΔtrhA), SL1344 (R27 ΔtrhA RSP-

734   Flag) and SL1344 (R27 Δ*trhA* RSP2-Flag), respectively. (E) Immunodetection of RSP-Flag and RSP2-Flag in the intracellular compartments of strain SL1344

735   harboring the corresponding *trhA* derivatives of the R27 plasmid. The experiments were repeated three times. A representative experiment is shown.

Figure 7. Loss of TrhH function impairs the effect of the RSP and RSP2 proteins on SL1344 cell

motility. The motility of the different strains was measured as the halo diameter of the different

colonies growing on motility agar. The results are the means of three independent experiments.

Standard deviations are shown. Statistical analysis showed a significant difference analyzed by

unpaired two-sided Student´s t-test (*P-value 0.0009, **P-value <0.0001).

Figure 8. The RSP and RSP2 proteins bind to the conjugative pilus encoded by the R27 plasmid. Immunogold electron microscopy of cells from strains SL1344 Δ*fliC* Δ*fljB* (R27) (A) and SL1344 Δ*fliC* Δ*fljB* (R27 Δ*trhH*) (B) using polyclonal anti-RSP antibodies and goat anti-rabbit IgG conjugated to 12 nm gold particles. Arrows point to the RSP protein associated with the conjugative pilus. Immunogold electron microscopy of cells from strains SL1344 Δ*fliC* Δ*fljB* (R27 RSP2-Flag) (C) and SL1344 Δ*fliC* Δ*fljB* (R27 RSP2-Flag Δ*trhH*) (D) using anti-Flag monoclonal antibodies and goat anti-mouse IgG conjugated to 12 nm gold particles. Arrows point to the RSP2 protein associated with the conjugative pilus. Bars represent 1 μm (in A, B and D) and 2 μm (in C).

Figure 9. The ALG87388.1 gene product of the pKAZ3 plasmid influences conjugation at 25°C. (A) SDS-PAGE analysis of the cell-free secretomes of strains

SL1344, SL1344 (R27) and SL1344 (pKAZ3) grown at 25ºC until $OD_{600\,nm}$ of 2.0. Asterisks indicated the detection of the ALG87388.1 protein in the secretome

of strain SL1344 (pKAZ3) confirmed by LC-MS/MS. Arrows indicate the RSP and RSP2 protein. (B) Motility of the SL1344, SL1344 (pKAZ3) and SL1344 (pKAZ3

ΔALG87388.1) strains. The data shown are the means and standard deviations of three independent experiments. Statistical analysis showed significant

differences analyzed by unpaired two-sided Student´s t-test (*P-value 0.0469; n.s. not significant). (C and D). Effect of the ΔALG87388.1 allele on the

conjugation frequency of the pKAZ3 plasmid at 25°C (C) and 37°C (D) in liquid media. The data shown are the means and standard deviations of three

independent experiments. Statistical analysis showed significant differences analyzed by unpaired two-sided Student´s t-test (*P-value 0.042; n.s. not

significant).