A burst of regulatory and protein innovation at the origin of placental mammals drove the

emergence of placenta and underpins divergent early pregnancy strategies in modern

mammals.

Alysha S. Taylor[1,2], Haidee Tinning[1], Vladimir Ovchinnikov[3], William Smith[1,4], Anna L. Pullinger[1],

Ruth A. Sutton[1], Bede Constantinides[2,5], Dapeng Wang[6,7], Niamh Forde[*1], Mary J O'Connell[*2,3]

[1]Discovery and Translational Sciences Department, Leeds Institute of Cardiovascular and

Metabolic Medicine, Faculty of Medicine and Health, University of Leeds, LS2 9JT, United

Kingdom.

[2]School of Biology, Faculty of Biological Sciences, University of Leeds, LS2 9JT, United Kingdom.

[3]School of Life Sciences, Faculty of Medicine and Health Sciences, University of Nottingham,

NG7 2RD, United Kingdom.

[4] Leeds Fertility, Leeds Teaching Hospitals NHS Trust, York Road, Seacroft, Leeds, LS14 6UH,

United Kingdom.

[5]Present address: Modernising Medical Microbiology Consortium, Nuffield Department of

Clinical Medicine, John Radcliffe Hospital, University of Oxford, OX3 9DU, United Kingdom

[6] LeedsOmics, University of Leeds, United Kingdom.

[7] Present address: Wellcome Centre for Human Genetics, University of Oxford, OX3 7BN, United

Kingdom.

[*] Both corresponding authors contributed equally to this manuscript: Correspondence should

be addressed to n.forde@leeds.ac.uk or drmary.oconnell@gmail.com

Summary paragraph: The origin of live birth in mammals ~148 million years ago was a dramatic shift in reproductive strategy, yet the molecular changes that established mammal viviparity are largely unknown. Although progesterone receptor signalling predates the origin of mammals andis highly conserved in, and critical for, successful mammal pregnancy, it alone cannot explain the origin and subsequent diversity of implantation strategies throughout the placental mammal radiation. MiRNAs are known to be flexible and dynamic regulators with a well established role in the pathophysiology of mammal placenta. We propose that a dynamic core microRNA (miRNA) network originated early in placental mammal evolution, responds to conserved mammal pregnancy cues (*e.g.* progesterone), and facilitates species-specific responses. Here we identify 13 miRNAs that arose at the origin of placental mammals and were subsequently retained in all descendent lineages. The expression of these 13 miRNAs in response to early pregnancy molecules is regulated in a species-specific manner in endometrial epithelia of species with extremes of implantation strategies. Furthermore, these 13 miRNAs preferentially target 84 proteins under positive selective pressure on ancestral eutherian. Discovery of this core "live-birth" toolkit and specifically adapted proteins helps explain the origin and evolution of the placenta in mammals.

Successful pregnancy in eutheria is contingent on a developmentally competent embryo, appropriate endometrial function, and on formation of the placenta and on the molecular cross-talk across these components [1,2]. Yet placental organ morphology, and more generally, the underlying regulation of successful pregnancy, varies across mammals. Protein coding alterations along with innovations in regulatory networks drive the origin and evolution of novel traits [3]. Moreover, bursts of evolution of new microRNAs (miRNAs) are known to be associated with morphological innovation [4–6], and miRNAs are known to regulate placental function in both normal and pathophysiological conditions [7,8]. We have uncovered a core placental toolkit in mammals that evolved through a synergy of molecular events involving the emergence of new miRNAs combined with adaptive amino acid changes in key proteins. We propose that this conserved core toolkit contributes to the diversity of implantation strategies and pregnancy morphologies observed in modern eutheria (Figure 1).

A core miRNA toolkit of 13 miRNAs arose at the origin of therian and eutherian mammals and

were never subsequently lost.

Because significant miRNA family expansions have been found to correlate with major transitions in animal evolution[4–6], we interrogated MirGeneDB[9] and identified 112 miRNA families that originated on either the therian (6 miRNA families) or eutherian stem lineage (106 miRNA families). Six of these miRNA families emerged on the older therian stem lineage (mir-340, mir-483, mir-671, mir-675, mir-1251, and mir-3613), of which only mir-340 is phylogenetically conserved in both Theria and Eutheria (Figure 2a). Given that mir-671 is present in tasmanian devil and in all Eutheria sampled, we classified mir-671 as a 'therian stem lineage miRNA'. A total of 106 miRNAs emerged at the origin of eutheria, and 11 of these remained conserved in all extant eutheria sampled (Figure 2a). In total, this yielded 13 miRNAs that originated on stem mammal lineages and were subsequently retained in all descendent lineages, henceforth referred to as "the core miRNA toolkit". Expression and function of some of the core miRNA toolkit has been demonstrated for normal placenta (*e.g.* mir-433[10], mir-28[11], mir-378[12]), and endometrium (*e.g.* mir-505[13] and mir-542[14]). Others have been implicated in pathophysiological pregnancies, *e.g.* mir-185, mir-188, mir-423 and mir-542[15–17] have been implicated in preeclampsia, mir-127[10] has been associated with placentomegaly, mir-324 is associated with LGA pregnancies[18], mir-331 is associated with placenta from intra-amniotic infection[19], and mir-505 can be associated with preterm birth[20]. Collectively all 13 members of the core miRNA toolkit have been implicated in critical roles in placental mammal pregnancies.

Targets were identified in the human genome for the core miRNA toolkit using TargetScan[21]. The predicted functions of the targets of the core miRNA toolkit include reproductive functions (55 target transcripts), metabolic process (1476 target transcripts), and Biological regulation (1269 target transcripts) (Supplementary Figure S1b). More specifically, some of the targets were implicated in pathways associated with INPP5E regulation, Neurophilin interactions, and VEGF interactions with its receptor (VEGR), each involved in the process of angiogenesis. TGF-β signalling and p53 regulation are amongst the predicted targets and are implicated in cell proliferation (Supplementary Figure S3). Both angiogenesis and

proliferation are required for successful pregnancy. The syncytins are a family of endogenous

retrovirus-derived protein coding regions that were domesticated in mammals and are

essential for promoting placental formation[22] and 4/13 of the core miRNA toolkit (miR-185-3p,

miR-188-3p, miR-423-5p and miR-433-3p) are predicted to target the syncytins with at least 7-

mer binding. In the case of miR-423-5p, there are two predicted target sites for *syncytin-1*, one

of which has a site overlapping with that of the miR-185-3p binding site, indicating

dynamic/competitive binding between these miRNAs and their *syncytin-1* target. The predicted

targets of the core miRNA toolkit also include 130 gene families that have been proposed to

have emerged on the eutherian stem lineage[23]. In addition, the targets included genes that

evolved endometrial expression on the stem eutherian lineage, and that are hypothesized to

have assisted in the remodelling of the uterine landscape during the evolution of the mammal

placenta[1].

Evidence of positive selection on the stem eutherian lineage.

Protein coding alterations (such as the birth of new genes and gene families, gene loss,

co-option, and selective pressure variation) along with innovations in regulatory networks drive

the development of novel traits[3]. A number of cases of positively selected amino acids

(indicative of protein functional shift) are known to have had a direct role in endometrial

function, *e.g.* the galectin family of proteins involved in immune modification at the maternal-

fetal interface[24]. We looked for evidence of adaptive evolution in single gene orthologous

families (SGOs) on the stem eutherian lineage. We focussed on SGOs to optimise our ability to

assign function to orthologs and to accurately trace evolutionary histories. We chose a total of

ten Eutheria that demonstrate the greatest range of diversity in placental morphology, plus

four outgroup species, one from each of the *Monotremata, Marsupialia, Aves* and *Teleostei*

(Supplementary Table S1). Annotated gene families were taken from Ensembl 90[25]. Following

our filtering regime we extracted a total of 1,437 SGOs. Applying codon models of evolution to

these SGOs, we identified signatures of positive selection on amino acid residues on the stem

eutherian lineage in 237 SGOs. The functions of these genes are predominantly cellular

processes, metabolic processes, and biological regulation (Supplementary Figure S2). Out of

these 237 positively selected SGOs, 115 contained positively selected amino acid residues that

were subsequently unaltered in all descendent lineages. The 115 SGOs are functionally

enriched for chromosomal maintenance, telomere activity, p53 signalling, cell cycle, and the

inflammatory immune response - activities that have been associated with the formation of the

placental tissue in pregnancy[26–28]. We then studied whether there is a significant association

between the core miRNA toolkit and stem-lineage positively selected proteins.

The core miRNA toolkit preferentially targets genes under positive selection in the stem

eutherian lineage.

Synergy between regulatory and protein coding innovations often drives substantial

phenotypic novelty[2,3,29]. Therefore, to test the hypothesis that innovation both at the level of

regulation and of protein coding change underpinned the origin of placentation in mammals,

we performed a simulation study on the targets of the 13 stem lineage miRNAs. Out of 115

SGOs with evidence of positive selection on the ancestral eutherian lineage, 84 were found to

be significantly enriched for binding sites for the 13 members of the core miRNA toolkit

(p=1.35618e-11), with a mean of 6.66 binding sites per transcript (median=4.0) (Figure 2b). We

determined if the number of binding sites in this subset of 84 positively selected SGOs was

significantly different than one would expect by random chance. We estimated the number of

binding sites per transcript for the core miRNA toolkit in 100 randomly sampled gene sets and

found it is significantly lower, with a mean=1.64 (median=1.0). This suggests that the 13

miRNAs in the core miRNA toolkit preferentially target the positively selected SGOs (6.66

binding sites compared to 1.64) (Figure 2b). Combined, this indicates that a co-evolutionary

process arose in a short time in early mammal evolution that resulted in altered protein

function, as well as a new miRNA-mediated regulatory network.

The functions of the 84 positively selected SGOs targeted by the core miRNA toolkit

broadly fall within the categories of cell cycle, DNA damage & DNA metabolic processes, and

regulation of hair cycle & hair cycle – a uniquely mammal characteristic (Supplementary Figure

S3). 21/84 SGOs are significantly more likely to interact with one another (p<0.05) in

comparison to any other gene in the genome (Supplementary Figure S2b). Of course there is

significant variation found in modern mammals in other facets such as telomere biology, cancer

incidence, body mass and maximum lifespan[30–33], therefore innovation at this node was not

expected to be entirely skewed to pregnancy (Supplementary Table S2)

Species-specific regulation of the core miRNA toolkit by key early pregnancy molecules in

species with different implantation strategies.

Implantation in eutherian mammals displays wide variation in both embryological

morphology and bi-lateral signalling between the embryo and maternal endometrium, and

degree of invasiveness (invasive in human, superficial in bovine). This variation is governed, in

part, by conserved signalling pathways, *e.g.* the sustained actions of the hormone progesterone

(P4)[34], but also diverse molecular cues such as the maternal recognition of pregnancy signal,

e.g. chorionic gonadotrophin (hCG) in human and interferon (IFNT) in bovine[35,36]. We asked the

question what molecules, that are involved in bi-lateral communication between the embryo

and endometrium, regulate the core mammal miRNA toolkit and if they are regulated in a

species-specific manner. We cultured endometrial epithelial cells from human and bovine[37] and

exposed them for 24 hr to recombinant forms of conserved (P4, CAPG, and PDI) and diverse

(IFNT and hCG) molecular cues important for early pregnancy success in placental mammals.

CAPG and PDI have recently been identified as produced by the bovine conceptus during

pregnancy recognition and are highly conserved (in terms of sequence identity and

phylogenetic distribution) across placental mammals[37,38]. We then examined the expression of

the 13 stem lineage miRNAs in these cells using an LNA-based approach. Treatment of bovine

endometrial epithelial cells with 10µg/mL of P4 during the early luteal phase resulted in

increased expression of miR-505-5p (Supplementary Figure S4). In summary, treatment with

the evolutionarily conserved early pregnancy proteins (P4, CAPG and PDI) in the endometrial

epithelial cells of bovine and/or human resulted in a change in expression of 11/13 of the stem

lineage miRNAs (Figure 1).

Intriguingly, addition of recombinant bovine forms of bCAPG and bPDI proteins to

bovine or human endometrial epithelial cells altered expression of selected miRNAs in a

species-specific manner (Figure 3&4 respectively). Treatment of bovine cells with recombinant

bCAPG resulted in increased expression of miR-331-3p, miR-324-5p and miR-505-5p in
endometrial epithelial cells (p<0.05). Whereas in human endometrial epithelial cells treated
with 1000ng/µl bCAPG, the expression of miR-127-3p, miR-151a-3p (a paralog of mir-28
originating on the Eutherian stem lineage), and miR-188-5p showed a significant decrease in
expression compared to vehicle control (p<0.05: Figure 3). Treatment with recombinant bPDI
decreased expression of miR-185-5p in bovine epithelial cells (P<0.05). In human Ishikawa
immortalised endometrial epithelial cells treated with 1000ng/µl bPDI the expression of miR-
151a-5p, miR-185-5p, miR-378a-3p and miR-532-5p (a paralogue of mir-188 originating on the
Eutherian stem lineage) were significantly decreased compared to vehicle control (Figure 4).

In contrast, addition of the species specific pregnancy recognition signals (IFNT in
bovine: hCG in human) to receptive endometrial epithelial cells altered expression of one
member of the core miRNA toolkit (Supplementary Figures S5 and S6 for IFNT and hCG data
respectively). These data demonstrate that the expression of the core miRNA toolkit is not
altered by the species-specific pregnancy recognition signals (IFNT and hCG).

Human and bovine represent two distinct implantation strategies for mammals and
these lineages last shared a common ancestor some ~92 million years ago, representing ~184
million years of independent evolution [39,40]. None of the 13 stem lineage miRNAs are regulated
by the species-specific pregnancy recognition signals (IFNT and hCG), but the expression of the
core miRNA toolkit is modified by proteins that are highly conserved amongst placental
mammals (CAPG and PDI). Combined, our results show that the preferential targeting of the
core mammal miRNA toolkit, and protein functional shift were essential to the establishment of
mammalian implantation, and that subsequent diversification of this network facilitated the
range of implantation strategies observed today.

In summary, this work identifies a core regulatory network that drove a major transition
- the origin of live birth in mammals. With the core now defined, future work can focus on the
accessory elements that drove the subsequent diversification of mammal placenta.

195
196
197
198   **ACKNOWLEDGEMENTS**

We would like to thank Dr Fuller Bazer from Texas A&M for the kind gift of recombinant ovine IFNT. We would like to acknowledge the assistance of Stefania Mountevedi in helping with the bovine cell isolation and culture. Recombinant proteins CAPG and PDI were generated by the Protein and Proteome Analysis core (NUPPA), Newcastle University, Newcastle upon Tyne, UK. This work was funded by University of Leeds under the LARS PhD scholarship (AST) and Maternity leave support scheme to MOC for BC, and by the University of Nottingham postdoctoral fellowship funding to MOC for VO. This work was supported by N8 agri-food pump priming, QR GCRF, as well as BBSRC grant number BB/R017522/1 to NF. This work was undertaken on ARC3, part of the High-Performance Computing facilities at the University of Leeds, UK. Figure 1 was created using Biorender.

CONTRIBUTIONS

NF and MJOC conceived of the study and designed the experiments. AST, BC, VO and MJOC carried out all evolutionary analyses and DW carried out statistical tests. NF, AST, HT, WS, ALP, and RAS undertook all molecular analyses. NF and MJOC drafted the manuscript, all authors contributed to revising, reading, and critiquing the manuscript.

REFERENCES

1. Lynch, V. J. *et al.* Ancient transposable elements transformed the uterine regulatory landscape and transcriptome during the evolution of mammalian pregnancy. *Cell Rep.* **10**, 551–561 (2015).

2. Kin, K. *et al.* The Transcriptomic Evolution of Mammalian Pregnancy: Gene Expression Innovations in Endometrial Stromal Fibroblasts. *Genome Biol. Evol.* **8**, 2459–2473 (2016).

3. Wagner, G. P. Evolutionary innovations and novelties: Let us get down to business! *Zool. Anz.* **256**, 75–81 (2015).

4. Hertel, J. *et al.* The expansion of the metazoan microRNA repertoire. *BMC Genomics* **7**, 25 (2006).

5. Prochnik, S. E., Rokhsar, D. S. & Aboobaker, A. A. Evidence for a microRNA expansion in the bilaterian ancestor. *Dev. Genes Evol.* **217**, 73–77 (2007).

6. Grimson, A. *et al.* The early origins of microRNAs and Piwi-interacting RNAs in animals. *Nature* **455**, 10.1038/nature07415 (2008).

231   7.   Keniry, A. *et al.* The H19 lincRNA is a developmental reservoir of miR-675 which
232   suppresses growth and Igf1r. *Nat. Cell Biol.* **14**, 659–665 (2012).

233   8.   Munaut, C. *et al.* Dysregulated circulating miRNAs in preeclampsia. *Biomed. reports* **5**,
234   686–692 (2016).

235   9.   Fromm, B. *et al.* MirGeneDB 2.0: the metazoan microRNA complement. *Nucleic Acids*
236   *Res.* **48**, D132–D141 (2020).

237   10.   Ito, M. *et al.* A trans-homologue interaction between reciprocally imprinted miR-127 and
238   Rtl1 regulates placenta development. *Development* **142**, 2425–2430 (2015).

239   11.   Farrokhnia, F., Aplin, J. D., Westwood, M. & Forbes, K. MicroRNA regulation of mitogenic
240   signaling networks in the human placenta. *J. Biol. Chem.* **289**, 30404–30416 (2014).

241   12.   Luo, L. *et al.* MicroRNA-378a-5p promotes trophoblast cell survival, migration and
242   invasion by targeting Nodal. *J. Cell Sci.* **125**, 3124 LP – 3132 (2012).

243   13.   Morales-Prieto, D. M. *et al.* MicroRNA expression profiles of trophoblastic cells. *Placenta*
244   **33**, 725–734 (2012).

245   14.   Tochigi, H. *et al.* Loss of miR-542-3p enhances IGFBP-1 expression in decidualizing
246   human endometrial stromal cells. *Sci. Rep.* **7**, 40001 (2017).

247   15.   Fu, G., Brkić, J., Hayder, H. & Peng, C. MicroRNAs in human placental development and
248   pregnancy complications. *Int. J. Mol. Sci.* **14**, 5519–5544 (2013).

249   16.   Harapan, H. & Andalas, M. The role of microRNAs in the proliferation, differentiation,
250   invasion, and apoptosis of trophoblasts during the occurrence of preeclampsia—A
251   systematic review. *Tzu Chi Med. J.* **27**, 54–64 (2015).

252   17.   Hosseini, M. K., Gunel, T., Gumusoglu, E., Benian, A. & Aydinli, K. MicroRNA expression
253   profiling in placenta and maternal plasma in early pregnancy loss. *Mol. Med. Rep.* **17**,
254   4941–4952 (2018).

255   18.   Rahman, M. L. *et al.* Regulation of birthweight by placenta-derived miRNAs: evidence
256   from an arsenic-exposed birth cohort in Bangladesh. *Epigenetics* **13**, 573–590 (2018).

257   19.   do Imperio, G. E. *et al.* Chorioamnionitis Induces a Specific Signature of Placental ABC
258   Transporters Associated with an Increase of miR-331-5p in the Human Preterm Placenta.
259   *Cell. Physiol. Biochem.* **45**, 591–604 (2018).

260   20.   Fallen, S. *et al.* Extracellular vesicle RNAs reflect placenta dysfunction and are a
261   biomarker source for preterm labour. *J. Cell. Mol. Med.* **22**, 2760–2773 (2018).

262   21.   Agarwal, V., Bell, G. W., Nam, J. W. & Bartel, D. P. Predicting effective microRNA target
263   sites in mammalian mRNAs. *Elife* **4**, 1–38 (2015).

22.   Esnault, C., Cornelis, G., Heidmann, O. & Heidmann, T. Differential Evolutionary Fate of

an Ancestral Primate Endogenous Retrovirus Envelope Gene, the EnvV Syncytin,

Captured for a Function in Placentation. *PLOS Genet.* **9**, e1003400 (2013).

23.   Dunwell, T. L., Paps, J. & Holland, P. W. H. Novel and divergent genes in the evolution of

placental mammals. *Proceedings. Biol. Sci.* **284**, 20171357 (2017).

24.   Than, N. G. *et al.* A primate subfamily of galectins expressed at the maternal–fetal

interface that promote immune cell death. *Proc. Natl. Acad. Sci.* **106**, 9731 LP – 9736

(2009).

25.   Yates, A. *et al.* Ensembl 2016. *Nucleic Acids Res.* **44**, D710–D716 (2016).

26.   Hauguel-de Mouzon, S. & Guerre-Millo, M. The Placenta Cytokine Network and

Inflammatory Signals. *Placenta* **27**, 794–798 (2006).

27.   Woods, L., Perez-Garcia, V. & Hemberger, M. Regulation of Placental Development and

Its Impact on Fetal Growth—New Insights From Mouse Models   . *Frontiers in*

Endocrinology   **9**, 570 (2018).

28.   Gal, H. *et al.* Molecular pathways of senescence regulate placental structure and

function. *EMBO J.* **38**, e100849–e100849 (2019).

29.   Lynch, V. J., Leclerc, R. D., May, G. & Wagner, G. P. Transposon-mediated rewiring of

gene regulatory networks contributed to the evolution of pregnancy in mammals. *Nat.*

Genet. **43**, 1154–9 (2011).

30.   Tian, X. *et al.* Evolution of telomere maintenance and tumour suppressor mechanisms

across mammals. *Philos. Trans. R. Soc. B Biol. Sci.* **373**, 20160443 (2018).

31.   McNab, B. K. An analysis of the factors that influence the level and scaling of mammalian

BMR. *Comp. Biochem. Physiol. Part A Mol. Integr. Physiol.* **151**, 5–28 (2008).

32.   Garratt, M., Gaillard, J.-M. J., Brooks, R. C. P., Lemaitre, J.-F. & Lemaître, J.-F.

Diversification of the eutherian placenta is associated with changes in the pace of life.

Proc. Natl. Acad. Sci. USA **110**, 7760–7765 (2013).

33.   Welch, J. J., Bininda-Emonds, O. R. P. & Bromham, L. Correlates of substitution rate

variation in mammalian protein-coding sequences. *BMC Evol. Biol.* **8**, 53 (2008).

34.   Spencer, T. E. & Bazer, F. W. Temporal and Spatial Alterations in Uterine Estrogen

Receptor and Progesterone Receptor Gene Expression During the Estrous Cycle and

Early Pregnancy in the Ewe1. *Biol. Reprod.* **53**, 1527–1543 (1995).

35.   Godkin, J. D., Bazer, F. W., Moffatt, J., Sessions, F. & Roberts, R. M. Purification and

properties of a major, low molecular weight protein released by the trophoblast of sheep

blastocysts at Day 13–21. *Reproduction* **65**, 141–150 (1982).

36.   Morgan, F. J. & Canfield, R. E. Nature of the Subunits of Human Chorionic Gonadotropin.

Endocrinology **88**, 1045–1053 (1971).

37.   Tinning, H. *et al.* The role of CAPG in molecular communication between the embryo and

the uterine endometrium: Is its function conserved in species with different implantation

strategies? *FASEB J.* **34**, 11015–11029 (2020).

38.   Forde, N. *et al.* Conceptus-Induced Changes in the Endometrial Transcriptome: How

Soon Does the Cow Know She Is Pregnant?1. *Biol. Reprod.* **85**, 144–156 (2011).

306
307
308 **Figure 1:**
309
310

311
312

Figure 1. Regulation of core miRNA toolkit by early pregnancy markers. *The endometrial*
epithelium of bovine (superficial implantation strategy) (top panel) or human (invasive
implantation strategy) (bottom panel) are regulated by molecules important for endometrial
function in early pregnancy in eutheria (P4: progesterone; hCG: human chorionic

gonadotrophin; Interferon Tau: IFNT; Macrophage capping protein: CAPG, and protein disulfide

isomerase: PDI).

Figure 2:

Figure 2: Phylogenetic distribution of the miRNA families and their levels of targeting in positively selected genes. (a) *Phylogenetic distribution of miRNA families specific to therian and eutherian mammals. The mammal phylogeny displaying the species sampled in our analysis. The corresponding matrix shows the presence (dark red) or absence (pale grey) of miRNA families across the species sampled.* ***(b)*** *Violin plot comparing the number of target miRNA binding sites from the core miRNA toolkit per transcript in the 84 genes that underwent positive selection on the stem Eutherian lineage (PSGs) (left) compared to sets of randomly sampled targets of the miRNAs that do not have signatures of positive selection on the stem Eutherian lineage (nPSG) (right). For each of the 84 PSGs, the mean number of predicted miRNA binding sites was determined for each target transcript in human (depicted in green). This was compared to the mean number of binding sites for each of the nPSGs (depicted in blue). The mean number of binding sites was determined to be significantly different between the two datasets when p≤0.05, two-sample t-test with unequal variance.*

344
345 **Figure 3:**

346
Figure 3. ***Regulation of stem lineage miRNAs in bovine and human endometrial epithelial cells***
treated with bCAPG. *Expression of stem lineage miRNAs miR-127-3p, miR-151a-3p, miR-188-*
5p, miR-324-3p, miR-331-3p, and miR-505-5p in either bovine (left hand side of each pair) or
human (right hand side of each pair) endometrial epithelial cells. Primary bovine endometrial
epithelial cells were treated with vehicle control (grey circle), or 1000ng/μl bCAPG (purple circle)
for 24 hours. Human Ishikawa immortalized endometrial epithelial cells were treated with
vehicle control (grey circle), or 1000ng/μl bCAPG (purple circle) for 24 hours. Significant
differences in miRNA expression values determined when p≤0.05 are depicted by an asterisk (*).
355

364
Figure 4: *Regulation of stem lineage miRNAs in bovine and human endometrial epithelial cells*
treated with bPDI. *Expression of stem lineage miRNAs miR-127-3p, miR-151a-3p, miR-188-5p,*
miR-324-3p, miR-331-3p, and miR-505-5p in either bovine (left hand side of each pair) or human
(right hand side of each pair) endometrial epithelial cells. Primary bovine endometrial epithelial
cells were treated with vehicle control (grey circle), or 1000ng/µl bPDI (orange circle) for 24
hours. Human Ishikawa immortalized endometrial epithelial cells were treated with vehicle
control (grey circle), or 1000ng/µl bPDI (orange circle) for 24 hours. Significant differences in
miRNA expression values determined when p≤0.05 are depicted by an asterisk (*).

384
385
386

387   **Supplementary Figures and Tables**

Supplementary Figure S1:

Figure S1. Number of predicted targets for each of the 13 stem lineage miRNA family and the

functional annotation of the target genes using Panther DB. (A) *Targets were predicted for*

each of the 13 stem lineage miRNAs using TargetScan70 (Agarwal et al., 2015). TargetScan

output was filtered for targets with8mer-A1, 7mer-m8 and 7mer-A1 complementary binding to

the seed region. ***(B)*** *Functional annotation of predicted targets of the 13 stem lineage miRNAs.*

Filtered stem lineage miRNA target transcripts were analysed for functional enrichment using

PANTHERv.14 (Muruganujan et al., 2018), where PANTHERv.14 found functional annotations to

be significantly enriched when p≤0.05.

Supplementary Figure S2:

405
406
Figure S2. Broad functions of the genes with signatures of positive selection and sample of
their interactions. (A) *Functional annotation using Gene Ontology Biological Process terms of*
115 SGOs that underwent positive selection on the stem eutherian lineage and where the amino
acid substitution was fixed on all extant Eutheria tested. The absolute number (out of 115) of
positively selected genes in a given category are shown in the X axis and the functional
annotations on the Y-axis. ***(B)*** *String interaction network of the same set of 115 genes. Network*
has 106 total nodes and 8 edges (expected edges =4). Background nodes, with no high
confidence interactions from experimental and database sources are faded. Nodes with high
confidence interactions from experimentally determined (pink lines) or curated database (blue
lines) sources are depicted in colour. The network was found to be significantly enriched for
gene-gene interactions (p=0.0468). Average node degree is 0.151, with an average local
clustering coefficient of 0.104. Minimum interaction score is 0.700.
419
420

422   **Supplementary Figure S3:**

424   ***Figure S3: A standard TreeMap from REVIGO displaying the GO biological process terms***
425   ***present in the 84 PSGs that were predicted targets of the 13 stem lineage miRNAs****. Rectangle*
426   *size represents semantic uniqueness of GO term, defined by REVIGO as the negative of average*
427   *similarity to all other terms present in human.*

Figure S4. Expression of stem lineage miRNAs in bovine endometrial epithelial cells treated with P4. *Expression of stem lineage miRNA (i) miR-28-3p, (ii) miR-127-3p, (iii) miR-151a-3p, (iv) miR-151a-5p, (v) miR-185-5p, (vi) miR-188-5p, (vii) miR-324-5p, (viii) miR-331-3p, (ix) miR-340-5p, (x) miR-378a-5p, (xi) miR-423-3p, (xii) miR-423-5p, (xiii) miR-433-3p, (xiv) miR-505-5p, (xv) miR-532-5p, (xvi) miR-542-3p, (xvii) miR-660-5p, (xviii) miR-671-3p, (xix) miR-671-5p and (xx) miR-708-5p in bovine endometrial epithelial cells treated with vehicle control (grey circle), 0.1µg/mL (light blue), 1.0µg/mL (medium blue) or 10µg/mL P4 (dark blue circle) for 24 hours. Significant differences in miRNA expression values determined when p≤0.05 are depicted by an asterisk (*).*

Supplementary Figure S5:

457
458
Figure S5. Expression of stem lineage miRNAs in bovine endometrial explants treated with
recombinant oIFNT. Expression of stem lineage miRNA (i) miR-28-3p, (ii) miR-127-3p, (iii) miR-

151a-3p, (iv) miR-151a-5p, (v) miR-185-5p, (vi) miR-188-5p, (vii) miR-324-5p, (viii) miR-331-3p,
(ix) miR- 340-5p, (x) miR-378a-5p, (xi) miR-423-3p, (xii) miR-423-5p, (xiii) miR-433-3p, (xiv) miR-
505-5p, (xv) miR-532-5p, (xvi) miR-542-3p, (xvii) miR- 660-5p, (xviii) miR-671-3p, (xix) miR-671-
5p and (xx) miR-708-5p in bovine endometrial explants treated with vehicle control (grey
circle), or 1000ng/µl oIFNT (green circle) for 24 hours. Significant differences in miRNA
expression values determined when $p < 0.05$ are depicted by an asterisk (*).
467
468

Supplementary Figure S6:

Figure S6. Expression of stem lineage miRNAs in human endometrial epithelial cells treated with hCG. Expression of stem lineage miRNA (i) miR-28-3p, (ii) miR-127-3p, (iii) miR-151a-3p, (iv) miR-151a-5p, (v) miR-185-5p, (vi) miR-188-5p, (vii) miR-324-5p, (viii) miR-331-3p, (ix) miR-340-5p, (x) miR-378a-5p, (xi) miR-423-3p, (xii) miR-423-5p, (xiii) miR-433-3p, (xiv) miR-505-5p, (xv) miR-532-5p, (xvi) miR-542-3p, (xvii) miR- 660-5p, (xviii) miR-671-3p, (xix) miR-671-5p and

480 (xx) miR-708-5p in human Ishikawa immortalized endometrial epithelial cells were treated with

481 control (dark grey circle), vehicle control (light grey circle), or P4 (light blue circle), or P4+hCG

482 (darker blue circles) for 24 hours. No differences in miRNA expression were determined

483 (p>0.05).

 Supplementary Table S1: Vertebrate Species sampled, genome version, coverage and

 completion level.

Clade	Species	Version	Genome Quality
Fish	Zebrafish	GRCz11	Full Genome, Chromosome Level
	Chicken	Gallus_gallus-5.0	70X Coverage, Chromosome Level
Monotremes	Platypus	OANA5	6X Coverage, Chromosome Level
Metatheria	Opossum	monDom5	7.33X Coverage, Chromosome Level
Eutheria	Elephant	Loxafr3.0	7X Coverage, Scaffold Level
	Armadillo	Dasnov3.0	6X Coverage, Scaffold Level
	Mouse	GRCm38.p6	High Quality Reference Assembly
	Human	GRCh38.p12	High Quality Reference Assembly
	Mouse Lemur	Mmur_3.0	221.6X Coverage, Chromosome Level
	Bushbaby	OtoGar3	137X Coverage, Scaffold Level
	Cat	Felis_catus_9.0	72X Coverage, Chromosome Level
	Microbat	Myoluc2.0	7X Coverage, Scaffold Level
	Horse	Equ Cab 2	6.79X Coverage, Chromosome Level
	Cow	UMD3.1	9X Coverage, Chromosome Level

Table S1: Set of 14 species sampled for the selective pressure analysis. Using Ensembl 92 (Yates *et al.*,
2016)*, a dataset of genomes representative of (i) vertebrate outgroup clades, or (ii) variations in
placental morphology in metatherian and eutherian mammals. For each clade, genomes were chosen
based on genome coverage for downstream homology searching. 'Clade' refers to the taxonomic group
of each included species. 'Species' denotes the included species, by their common name. 'Version'
denotes the genome assembly version included in this analysis. 'Genome Quality' refers to the coverage
and assembly level of each species included in this analysis.*

496 **Supplementary Table S2: Example of three functionally related proteins under positive**

497 **selection on stem eutherian lineage**

Family	Gene Name	lnL	*p2*	ω2	Positions in the alignment predicted to be positively selected.
PTHR13211	WRAP53	-10149.383	0.04821	37.80743	72, 88, 114, 225, 253, 260, 261, 331, 423, 480, 493, 506, 510, 513, 522, 528
PTHR12066	TERT	-36003.667	0.02302	954.86770	121, 122, 140, 166, 197, 508, 878, 1020, 1110
PTHR14487	ACD	-15296.106	0.10315	30.62227	138, 143, 148, 178, 185, 200, 202, 206, 208, 209, 214, 269, 278, 296, 298, 314, 347, 363, 368, 379, 390, 394, 395, 412, 414, 450, 470, 474, 524, 530, 532, 533, 545, 548, 561, 571, 575, 586, 685, 698

498 *Table S2: The panther family ID and common gene names are provided for a set of 3 proteins*
499 *extracted to illustrate the cases of positive selection identified. The LnL value associated with*
500 *the fit of the codeml model (PAML) to the data are provided as are the associated proportion of*
501 *sites (p2) that have the corresponding ω2 (or Dn/Ds ratio). The sites estimated to be positively*
502 *selected are given in the final column, these are numbered as per aligned codon position.*
503