Phenomics data processing: Extracting dose-response curve parameters from high-resolution temperature courses and repeated field-based wheat height measurements

Lukas Roth[a,*], Hans-Peter Piepho[b], Andreas Hund[a]

[a]*ETH Zurich, Institute of Agricultural Sciences, Universitätstrasse 2, 8092 Zurich, Switzerland*
[b]*University of Hohenheim, Institute for Crop Science, Biostatistics Unit, Fruwirthstrasse 23, 70593 Stuttgart, Germany*

Abstract

Temperature is a main driver of plant growth and development. New phenotyping tools enable quantifying the temperature response of hundreds of genotypes. Yet, for field-derived data, temperature response modeling bears flaws and pitfalls concerning the interpretation of derived parameters. In this study, climate data from five growing seasons with differing temperature distributions served as starting point for a growth simulation of wheat stem elongation, based on a four-parametric temperature response function (Wang-Engel) including all cardinal temperatures. In a novel approach, we re-extracted dose-responses from the simulation by combining high-resolution (hours) temperature courses with low-resolution (days) height data. The collection of such data is common in field phenotyping platforms. To take advantage of the lack of supra-optimal temperatures during the stem elongation, simpler (linear and asymptotic) models to predict temperature-response parameters were investigated. The asymptotic model extracted the base temperature of growth and the maximum absolute growth rate with high precision, whereas simpler, linear models failed to do so. Additionally, the asymptotic model provided a proxy estimate for the optimum temperature. However, when including seasonally changing cardinal temperatures, the prediction accuracy of the asymptotic model was strongly reduced. In a field study with three winter wheat varieties, significant differences were found for all three asymptotic dose-response curve parameters. We conclude that the asymptotic model based on high-resolution temperature courses is suitable to extract meaningful parameters from field-based data.

Keywords: growth, high-throughput field phenotyping, stem elongation, wheat (Triticum aestivum).

1. Introduction

As early as 1984, it was noted that the validity and scope of a crop growth model depends on the available data, which are, however, severely limited (Porter and Rayner, 1984). Today, high-throughout field phenotyping

---

*Corresponding author: lukas.roth@usys.ethz.ch

(HTFP) potentially enables to quantify crop growth—and consequently crop response—of hundreds of genotypes

under field conditions (Araus et al., 2018; Tardieu et al., 2020). Understanding crop responses to environmental

factors is essential to secure global food demands: Improvement in wheat yields stagnated on a global scale in

the past three decades (Brisson et al., 2010; Laidig et al., 2017). It was suspected that for Europe, the changing

climate increasingly impacts wheat yields (Brisson et al., 2010). Mitigating these changes in plant breeding to

improve yield requires understanding crop responses to environmental factors (Ramirez-Villegas et al., 2015). A

main driver of plant growth and development is temperature (Porter and Gawith, 1999).

1.1. Approaches to quantify temperature responses under controlled conditions

The effects of changes in temperature on crops are well studied in controlled environments, but the translation

of insights to the field is not straightforward. The temperature response of developmental processes is usually

studied by exposing plants to different temperature regimes—often during rather short time period of their life

cycle. Two different approaches are used: one can either (1) expose different plants of the same genotype to

different temperatures (Slafer and Rawson, 1995; Hund et al., 2012; Reimer et al., 2013), or (2) apply short

phases of different temperatures to the same plant. Examples of the latter approach are the studies on leaf

growth performed in the indoor platform 'Phenoarch', summarized by Parent and Tardieu (2012). This approach

requires that the temperature response is measured at a constant growth phase, during which growth is linear

when the temperature remains constant. Hence, the response-pattern can be observed non-destructively on the

same plant, offering a high throughput. In such experiments, the conditions of all but one covariate may be kept

constant. Furthermore, the reversibility of the process can be tested by repeatedly applying the same conditions

during the course of the experiment.

In general, developmental processes follow some sort of peak function in response to increasing tempera-

tures, i.e., there is a base temperature T_{min} at which growth starts, an optimal temperature T_{opt} at which growth

reaches its maximum absolute growth rate r_{max}, and a maximum temperature T_{max} at which growth stops (Porter

and Gawith, 1999). Depending on the research field, different models are used to approximate such functions—

for an overview see, e.g., Wang et al. (2017); Parent et al. (2019) for plants, and Rebaudo et al. (2018); Re-

baudo and Rabhi (2018) for ectoterms (i.e., animals whose regulation of body temperature depends on external

sources). In their work, Wang et al. (2017) distinguish four types of temperature response functions with in-

creasing complexity: linear functions with no optimum or maximum temperature (Type 1), capped linear or

curvilinear functions with an optimum but no maximum temperature (Type 2), segmented linear functions with

optimum and maximum temperatures (Type 3), and segmented linear functions or curvilinear functions with

three cardinal temperatures (Type 4). Clearly, functions of Type 4 are most appropriate to describe temperature

responses. Among the most widely used Type 4 functions for plants are the Wang-Engel cardinal temperature

function (Wang et al., 2017) and a modified version of the reaction rates equation from Johnson et al. (1942)

applied to different growth processes in a variety of major crops by Parent and Tardieu (2012). We will refer to

these different response functions in detail further down.

1.2. Cardinal temperatures of wheat

In their meta-analyses reporting cardinal temperatures of different developmental processes of wheat, Porter
and Gawith (1999) summarized 65 controlled condition and field studies with regard to the minimum (T_{min}),
optimum (T_{opt}) and maximum (T_{max}) temperature. Mean cardinal temperatures (based on daily averages) for
shoot growth were 3.0 °C (T_{min}), 20.3 °C (T_{opt}) and > 20.9 °C (T_{max}). Notably, T_{opt} and T_{max} were very close
together in many studies.

While Porter and Gawith (1999) were reporting on cardinal temperatures related to the phenology of wheat,
Parent and Tardieu (2012) were relating to those of a single organ, specifically, a wheat leaf. In contrast to the
comparably low T_{opt} summarized by Porter and Gawith (1999), Parent and Tardieu (2012) reported a significantly
higher T_{opt} of 27.7 °C when summarizing eight studies and own data. Since a wheat stem is composed of several
growing internodes, we consider its (long-term) elongation to be closer related to phenology than to (short-term)
organ growth. Consequently, we will use the more conservative, i.e., lower, temperature optimum reported by
Porter and Gawith (1999) in this study.

Yet, in addition to the short-term response of growing organs within the borders of genotype-specific cardinal
temperatures, there is a long-term adjustment of these cardinal temperatures during the growing season. Slafer
and Rawson (1995) examined T_{min} and T_{opt} for leaf appearance rates of vernalized winter wheat plants within
three phases, i) after vernalization up to the terminal spikelet, ii) from the terminal spikelet up to heading, and
iii) from heading to anthesis. The authors found that, averaged across cultivars, T_{min} was -1.9 °C, 1.2 °C and
8.1 °C, while T_{opt} was < 22 °C, 25 °C and > 25 °C for the three different phases, respectively. In addition, the
cultivars differed substantially for the two parameters. Consequently, one may assume that in particular the
base temperature may change throughout the season in a cultivar-specific manner. Nevertheless, the estimates
from Slafer and Rawson (1995) of cardinal temperatures and their changes were derived from a controlled
environment study. The benefits of controlled environment phenotyping lies in the ability to test a wide range of
temperature regimes while keeping other influential factors, such as vapor pressure deficit and light, constant.
However, plants usually do not experience such conditions in the field, as discussed by Slafer and Rawson (1995).

1.3. Approaches to quantify temperature responses with high-throughput in the field

Under field conditions, the temperature regimes are distinctly different from the controlled-environment stud-
ies considered so far. The environmental covariates in the field follow an annual and diurnal pattern to which
plants adjust their life cycle. HTFP offers the possibility to assess crop growth under such more realistic condi-
tions with high temporal and spatial resolution (Rebetzke et al., 2019). Hence, if assuming that temperatures
measured at a local weather station are closely related with effective in-canopy temperatures—i.e., meristem
temperatures (Gallagher et al., 1979)—one can use the temperature course during the season to quantify the
genotype-specific temperature responses. Still, such studies are sparse. Grieder et al. (2015) extracted the base

temperature and response to increasing temperatures at early growth stages before terminal spikelet formation.
The authors were able to extract heritable slopes and intercepts of linear responses. Kronenberg et al. (2020a)
were able to extract heritable response traits for the stem elongation phase based on data from three seasons.
Again, the authors used linear regressions of growth versus temperature as model. Kronenberg et al. (2020a)
argued that the recorded average temperatures in the measuring intervals were well below the optimum temper-
ature of 27.7 °C reported by Parent and Tardieu (2012), which would justify approximating a curvilinear with
a linear function. However, the optimum temperature of 20.3 °C summarized by Porter and Gawith (1999) is
distinctly lower. Following those results, it is likely that growth reached its optimum at several hours of the
day during their measurement period. Therefore, the extracted temperature response (slope) may be greatly
affected by the optimum temperature and maximum abolute growth rate at this temperature. Thus, modelling
plant response to temperature based on HTFP-derived data poses several challenges, e.g., the comparably sparse
measuring frequency, the unpredictable range of temperatures, and the changes of cardinal temperatures during
the measuring period.

1.4. Research aims

HTFP data are characterized by irregular and long trait measurement intervals but regular and short covariate
measurement intervals from a weather station. Our main research question was how to parameterize genotype-
specific temperature response pattern from such HTFP data. Using the stem elongation phase of winter wheat as
an example, we aimed to evaluate a new modelling approach with regard to its suitability. Therefore, we simu-
lated the temperature-response in the stem elongation phase using a Wang-Engel response function. The chosen
parameters were based on existing literature on cardinal temperatures in winter wheat (Porter and Gawith, 1999)
and enhanced with realistic temperature recordings of the stem elongation phase from a local weather station
at the field phenotyping platform site of ETH Zurich (FIP) (Kronenberg et al., 2020a). To evaluate the strengths
and limitations of our approach, we added changing cardinal temperatures, shifts in the start and end of the
elongation phase, measurement noise, and spatial field inhomogeneities to our simulation (Appendix, Figure 6).

2. Background

Modeling the stem elongation phase of winter wheat on a HTFP trait level (i.e., plant heights) requires as-
suming a theoretical whole-plant mechanism that is the result of the combination of a vast number of underlying
physiological mechanisms (Tardieu et al., 2020). Although the number of combinations of such mechanism may
be near-infinite, Tardieu et al. (2020) argued that natural selection and breeding activities have constrained
them to a limited number, resulting in a 'meta-mechanism' that is consistent across modeling scales. In the fol-
lowing, we provide background information on tools that allow extracting parameters of the 'meta-mechanism'
temperature response from HTFP data.

 2.1. Parametric versus semi-parametric approaches

Semi-parametric approaches (e.g., using smoothing techniques) are often favored to extract traits related to
growth processes from HTFP time series (van Eeuwijk et al., 2019; Roth et al., 2021a; Pérez-Valencia et al., 2022),
as they avoid the daunting task of modeling explicitly the—potentially unknown—influence of time-dependent
covariate courses. In a previous research, we have shown that semi-parametric approaches are suitable to extract
timing of key stages and quantities at defined time points or periods from simulated HTFP data. Those data were
generated using a dose-response curve that was unknown to the model being fitted, and corresponding covariate
dependencies masked by a "smoothing" part in the model (Roth et al., 2021b).

In contrast, to extract temperature responses, one cannot avoid modeling a dose-response curve explicitly, as
one is interested in its parameters. Consequently, we hereafter propose a parametric approach. Although our
focus is on HTFP data, the approach is also directly applicable to indoor platform data, but additionally considers
field-specific characteristics.

In comparison to greenhouse or climate chamber data, HTFP data have two major drawbacks: (i) Field-based
measurements are notoriously noisy due to environmental and soil inhomogeneities and measurement errors;
(ii) the suitability of measured trait time series and covariate courses to extract dose-response curves requires
large numbers of years and/or locations or the good luck of the scientist to catch at least one year with a (yet
unknown) close-to-optimal distribution of temperatures along the season. Research in agriculture has already
addressed the issue of spatial heterogeneity with the development of highly specialized experimental designs (e.g.,
by augmenting row-column designs with check cultivars (Piepho and Williams, 2016)) and matching statistical
analysis (e.g., by adding spatial components to the model (Piepho and Williams, 2010)). We could already
show that following these principles is of advantage for HTFP as well (Roth et al., 2021b). For brevity, we
therefore focus on the extraction of response traits from time series but refrain from further processing to adjusted
genotype means. To allow determining and comparing the performance of non-linear and linear models and the
requirements on an 'close-to-optimal' distribution of temperatures along the season, we use a simulation that is
parametrized based on real field data.

2.2. Temperature response functions

The linear temperature response model used in Grieder et al. (2015) is defined as

$$r_{\text{linear}}(T) = a \cdot (T - T_{\text{min}}), \tag{1}$$

155 where T is the temperature, a is the slope of the response, and T_{min} the base temperature. Note that such a model
156 causes negative growth rates r for $T < T_{\text{min}}$ and does not include an optimum temperature T_{opt}. Therefore, it
157 is likely that the extracted temperature response (slope) is affected by the optimum temperature and maximum
158 absolute growth rate at this temperature. Additionally, using such a Type 1 response (a linear function with no
159 optimum or maximum temperature, Wang et al. (2017)) will come to its limitation when data ranges span a

whole growing season where temperatures also extend to supra-optimal ranges (Weikai and Hunt, 1999; Parent et al., 2019). In the crop model community, models describing the response to temperature vary widely, but most of them consider an optimum temperature beyond which the growth rate levels off (Bonhomme, 2000; Parent et al., 2010; Wang et al., 2017).

Arrhenius type functions were shown to adequately describe the dose-response relationship between temperature and growth across a wide range of temperatures and species (Parent et al., 2010; Parent and Tardieu, 2012). However, it was disputed whether this approach allows for a mechanistic interpretation of the different model parameters (Clavijo Michelangeli et al., 2016). Evidently, research in wheat so far mainly focused on determining (non-mechanistic) cardinal temperatures such as the lower and upper temperature of growth T_{min} and T_{max} and the optimum temperature T_{opt} where the growth rate r is maximal (Porter and Gawith, 1999). Wang et al. (2017) could show that using the Wang-Engel temperature function (Wang and Engel, 1998) allows to adequately replace an Arrhenius type functions to describe growth rates using these cardinal temperatures,

$$r_{\text{Wang-Engel}}(T) = r_{\text{max}} \cdot \frac{2(T - T_{\text{min}})^\alpha (T_{\text{opt}} - T_{\text{min}})^\alpha - (T - T_{\text{min}})^{2\alpha}}{(T_{\text{opt}} - T_{\text{min}})^{2\alpha}}$$

$$\alpha = \frac{\ln(2)}{\ln\big((T_{\text{max}} - T_{\text{min}})/(T_{\text{opt}} - T_{\text{min}})\big)},$$

(2)

where T is the temperature, r_{max} the maximum absolute growth rate at the temperature optimum T_{opt}, T_{min} the lower base temperature and T_{max} the upper base temperature of growth.

The Wang-Engel temperature function is based on three cardinal temperatures T_{min}, T_{max} and T_{opt}. If aiming to extract genotype-specific cardinal temperatures as well as the maximum absolute growth rate r_{max} from HTFP data, this would require to fit a four-parametric non-linear model with parameters that have interdependent constrains (e.g., $T_{\text{min}} < T_{\text{opt}}$). In our experience, extracting realistic parameter values by fitting such a model to noisy HTFP data is practically impossible for different reasons, i.e., (i) the optimum temperature T_{opt} and maximum absolute growth rate r_{max} are closely correlated, (ii) for field experiments performed in target environments (without stress treatments) in temperate climates, the maximum temperature T_{max} is typically outside the range of observed data, and (iii) proximity of cardinal temperatures T_{min} and T_{opt} may make it challenging to achieve convergence. Some authors have reacted to these restrictions by fixing one or multiple parameters and normalizing growth rates (Parent and Tardieu, 2012).

As a previously unconsidered alternative, we propose a Type 2 response that integrates an optimum temperature (Wang et al., 2017) to approximate the Arrhenius type functions of Parent and Tardieu (2012) based on an asymptotic model (Figure 1c),

$$r = r_{\text{max}} \cdot (1 - \exp(-\exp(s) \cdot (T - T_{\text{min}})))$$

$$r_{\text{asym}}(T) = \begin{cases} r, & r > 0 \\ 0, & \text{otherwise} \end{cases},$$

(3)

where T is the temperature, r_{max} is the maximum absolute growth rate (and therefore the asymptote of the curve),

T_{min} the base temperature where the growth rate is zero, and s characterizes the steepness of the response (natural

logarithm of the rate constant, thus '*lrc*') (Pinheiro and Bates, 2000). Please note that (1) *lrc* is a parameter

in our model and not—as the name may indicate—a constant, (2) Equation 3 reduces to a monomolecular /

Mitscherlich growth function if replacing the first factor r_{max} with one and substituting s with $\ln(r_{max})$ (Heinen,

1999). The proposed asymptotic model does not consider a maximum temperature and therefore supra-optimal

range, and has no inflection point (Paine et al., 2012). It requires fitting a three-parametric non-linear model.

Monomolecular growth functions such as the asymptotic function were frequently used to describe growth rates in

proportion to a substrate level, e.g., in response to photon flux density (Causton and Dale, 1990). Nevertheless,

we are not aware of previous applications to temperature response problems in plant science. We would like

to note that such an approach is only appropriate if one expects the maximum temperature T_{max} to be outside

the range of observed data, as it is the case for the stem elongation phase of winter wheat—at least under the

temperate climate of Switzerland.

2.3. Combining irregular and long trait measurement intervals (days) with constant and short covariate measure-
ment intervals (hours)

Independent of the specific dose-response model, one can describe a phenotype y at time point t (Figure 1a)

as the result of such a dose-response model $r()$ (Figure 1c) dependent on a temperature course T_t (Figure 1b)

and a genotype-specific set of crop model parameters θ, e.g., $\theta = (T_{min}, T_{opt}, r_{max})$,

$$y_t = \int_{t_0}^{t} r(T_{t'}, \theta) \, dt', \tag{4}$$

205   where t_0 is the time point were growth started and therefore a timing of key stage trait that needs to be determined

206   beforehand. Trait and covariates such as canopy height and temperatures are only measured at certain points in

207   time, dependent on the measurement interval Δm. We therefore may discretize Equation 4 to

$$y_t = \sum_{t'=t_0}^{t} r(T_{t'}, \theta) \cdot \Delta m, \tag{5}$$

where t' are measurement time points with $(t' = t_0, t_0 + \Delta m, t_0 + 2\Delta m, \ldots, t)$, e.g., days.

Note that fitting Equation 5 to data implies the same measurement interval for traits y_t and temperatures $T_{t'}$.

Although the goal of HTFP is to measure at small measurement intervals, unfavorable weather conditions may pre-

vent regular measurements at all times, while registering covariates (e.g., temperature) is weather-independent

and done with constant high frequency. Outdoor phenotyping therefore produces trait measurements with ir-

regular and long measurement intervals Δm_d (e.g., measurements every 3–4 days) but covariate measurements

with constant and short measurement intervals Δm_h (e.g., hourly measurements). Therefore, before applying

Equation 5 to data, an aggregation step is required.

Figure 1: Schematic drawing of trait measurements with irregular and long intervals for two contrasting genotypes (green and blue) in their growth phase (solid) and outside their growth phase (grey) (a), covariate measurements with short and constant intervals (b), and the asymptotic dose-response model based on a maximum absolute growth rate ($r_{\max}$), minimum temperature ($T_{\min}$), and steepness of the response (lrc) (c).

A common technique is to align trait and covariate measurements by aggregating the covariate values to a mean (e.g., Kronenberg et al., 2017, 2020a),

$$y_t = \sum_{d=t_0}^{t} \left(r\left(\overline{T}_d, \theta\right) \cdot \Delta m_d \right), \tag{6}$$

where $\overline{T}_d$ is the mean of all covariate measurements made in the time period between successive trait measurement time points d where $(d = t_0, t_0 + \Delta m_1, t_0 + \Delta m_1 + \Delta m_2, \ldots, t)$. In this study, we propose an alternative approach: One may also apply the dose-response function r to each covariate value T_{dh} where h indexes covariate values between successive trait measurement time points d, $(h = 1, \ldots, n_d)$, and sum up the responses to the trait measurements,

$$y_t = \sum_{d=t_0}^{t} \left(\sum_{h=1}^{n_d} r\left(T_{dh}, \theta\right) \cdot \Delta m_h \right). \tag{7}$$

In the following, the first method is denoted the T_{mean} method and the second method the T_{course} method.

3. Materials and Methods

3.1. Simulation of canopy height data

For phenotyping experiments, one aims extracting (growth) parameter that are genotype-specific. In this work we used a simulation to demonstrate the suitability of dose-response models to extract temperature response parameters. The main advantage of using a simulation instead of real data is that one can report the expected precision and potential modeling artifacts. For real data experiments, judging on the heritability of extracted parameters would be the only choice, which does not allow distinguishing between poor performance caused by modeling artifacts and poor performance caused by lack of genetic effects.

As base for the simulation, we used the Wang-Engel function that has a similar shape as the Arrhenius function but is based on cardinal temperatures where one can rely on corresponding literature for temperature ranges

(see Section Background). This approach allows examining whether simpler functions such as the asymptotic or linear function allow extracting temperature response parameters as well, despite the fact that they simplify the underlying 'biological' function.

The canopy growth of wheat genotypes was simulated using a growth function g_T that was based on the Wang-Engel dose-response model of Equation 2 parameterized with four genotype-specific dose-response crop model parameters $\theta^C = (T_{\min}, T_{\text{opt}}, T_{\max}, r_{\max})$. To mimic the characteristic variation in start and stop of the elongation phase for a diverse genotype set, this model was extended with time points for the start and end of growth, parameterized with two genotype-specific timing traits $\theta^T = (t\text{PH}_{\text{start}}, t\text{PH}_{\text{stop}})$. Due to this genotype-specific start and end of the growth process, values for measurement time points at the beginning and end of the season were unbalanced, potentially leading to modeling artifacts. An environment inhomogeneity factor u and a measurement error e were added, leading to the model

$$y_{it} = g_T(t, \theta_i^C, \theta_i^T; T_{dh}) + u_{it} + e_t, \tag{8}$$

where y_{it} is the measured canopy height (phenotype) for the ith genotype at campaign time points t in intervals of three days ($t = 1, 4, 7, \ldots, T$), θ_i^C and θ_i^T are genotype-specific crop growth model parameters respectively timing traits, T_{dh} are hourly (h) temperature readings nested in measurement days (d), u_{it} a time point and genotype-specific inhomogeneity error (i.e., simulating the influence of other covariates than temperature) and e_t a time point specific measurement error (i.e., simulating random measurement errors). The growth function g_T was specified as an implementation of Equation 7 based on the Wang-Engel function of Equation 2,

$$g_T(t, \theta^C, \theta^T; T_{dh}) = \sum_{d=1}^{t} \begin{cases} \sum_{h=1}^{72} r_{\text{Wang-Engel}}(T_{dh}, \theta^C) & t\text{PH}_{\text{start}} < d < t\text{PH}_{\text{stop}} \\ 0, & \text{otherwise}. \end{cases} \tag{9}$$

The genotype-specific traits θ_i^C and θ_i^T and the error e_t were simulated using independent (genotypic) effects with variance σ^2, and the field and environment imhomogeneities error u_{it} according to a first-order autoregressive (AR1) model, all assuming normality. Specific μ and σ^2 were chosen based on literature if available, and otherwise based on own field data (Roth, 2021). Corresponding sources and selected distributions for all simulation input parameter are summarized in Table 1. Two alternative simulations were performed: One with changing cardinal temperatures with time (indicated with '→' in Table 1), and one with fixed cardinal temperatures (set to the mean of changing cardinal temperatures indicated in Table 1). Changing cardinal temperatures $T_{\min}$ and T_{opt} were implemented as linear interpolation between the start of the growth phase $t\text{PH}_{\text{start}}$ with the corresponding lower cardinal temperate limit (relating to 'terminal spikelet' in Porter and Gawith (1999)), and the end of the growth phase $t\text{PH}_{\text{stop}}$ with the corresponding upper cardinal temperature limit (relating to 'anthesis' in Porter and Gawith (1999)). Changing cardinal temperatures $T_{\max}$ were implemented as 5 °C ramp between the start of the growth phase $t\text{PH}_{\text{start}}$ and the end of the growth phase $t\text{PH}_{\text{stop}}$ from 30 to 35 °C (moving around the $T_{\max}$ of 31 °C reported for 'anthesis' in Porter and Gawith (1999), as precise values for 'terminal spikelet' are

not provided).

Canopy growth was simulated for a measurement interval of three, seven and 14 days for a period between

the 15th of March and 20th of July based on temperature courses of three years (Figure 2). Additionally, an

irregular measurement frequency of two measurements per week with frequent shifts by one ($P(1) = 1/3$), two

($P(2) = 2/9$) or three ($P(3) = 1/9$) days was simulated. A total of 1,000 differing genotypes were assumed,

resulting in 5,000 genotype time series. As temperature data source, real weather data of five consecutive years

at the ETH research station of agricultural sciences in Lindau Eschikon, Switzerland (47.449 N, 8.682 E, 556 m

a.s.l.) were used.

3.2. Extracting dose-response curves

To extract the Wang-Engel dose-response curve parameters $T_{\min}$, T_{opt} and $r_{\max}$, the linear and asymptotic

growth response models (Equation 1 and 3) were fitted to canopy height data between $t\mathrm{PH}_{\mathrm{start}}$ and $t\mathrm{PH}_{\mathrm{stop}}$ for

two covariate options, T_{mean} based on averaged temperatures between measurements (Equation 6), and T_{course}

based on temperature courses with a period of one hour (Equation 7). In addition to the linear and asymptotic

model, Equation 3 was enhanced with a slope parameter for $T_{\min}$ in dependence of time of measurement to test

if changing cardinal temperatures can be integrated in the temperature response function,

$$r = r_{\max} \cdot (1 - \exp(-\exp(s) \cdot (T - (T_{\min} + a \cdot t)))) , \tag{10}$$

where a is the slope and t the day of measurement. Maximum-likelihood optimization was used to fit Equation

6 to a set of t data points, $X_t = (\bar{T}_t, y_t - y_{(t-1)})$. Fitting Equation 7 to data required modifying the definition of

the data set so that $X_t = (\vec{T}_t, y_t - y_{(t-1)})$, where $\vec{T}_t$ is a vector of covariate readings between trait measurement

time points $t-1$ and t.

As an independent random measurement errors was assumed, the parameter σ was fixed to the well-known

measurement error of Structure-from-Motion based height estimations (Roth et al., 2020), herein the simulation

input σ_m (Table 1). Parameters were optimized using the method L-BFGS-B of the base R function *optim*.

To allow examining the effect of noise caused by inhomogeneities and measurement errors, the extraction was

once performed on raw simulated data, once on data contaminated with the measurement error e, and once with

both measurement and inhomogeneities errors e and u. Bias, variance, root-mean squared error (RMSE) and

Pearson's correlation were calculated after extracting the parameters $T_{\min}$, *lrc* and $r_{\max}$ for the asymptotic model

and a (slope) and $T_{\min}$ for the linear model. Bias was defined as $\frac{1}{n}\sum_{i=1}^{n} \hat{\theta}_i - \theta_i$, RMSE as $\sqrt{\frac{1}{n}\sum_{i=1}^{n} (\hat{\theta}_i - \theta_i)^2}$,

where n is the number of simulated time series, $\hat{\theta}_i$ the extracted parameter value for the ith time series and θ_i

the simulation input parameter value.

3.3. Real field data case study

294   To give an outlook on the usefulness of combining high-resolution (hours) covariate data with low-resolution

295   (days) HTFP data, the asymptotic growth response model and the novel T_{course} approach were applied to terres-

296   trial laser scanning height data published in Kronenberg et al. (2020a). Three genotypes were selected (Nara,

Table 1: Model input parameters for the simulation.

		Values	Source
θ_i^T	$\mathcal{N}\left(\mu,\sigma^2\right)$	$\mu_{T_{\min}} = 1.5 \to 9.5$ °C	Porter and Gawith (1999) (Terminal spikelet→Anthesis)
		$\sigma_{T_{\min}} = 2$ °C	Assumption
		$\mu_{T_{\mathrm{opt}}} = 10.6 \to 21$ °C	Porter and Gawith (1999) (Terminal spikelet→Anthesis)
		$\sigma_{T_{\mathrm{opt}}} = 2$ °C	Assumption
		$\mu_{T_{\max}} = 30 \to 35$ °C °C	Porter and Gawith (1999) (Anthesis)
		$\sigma_{T_{\max}} = 2$ °C	Assumption
		$\mu_{r_{\max}} = 1.05$ mm h^{-1}, $\sigma_{r_{\max}} = 0.12$ mm h^{-1}	Roth (2021)
θ_i^{PH}	$\mathcal{N}\left(\mu,\sigma^2\right)$	2016: $\mu_{t\mathrm{PH}_{\mathrm{start}}} = 108$ d, $\sigma_{t\mathrm{PH}_{\mathrm{start}}} = 2.8$ d	Kronenberg et al. (2020a)
		2017: $\mu_{t\mathrm{PH}_{\mathrm{start}}} = 103$ d, $\sigma_{t\mathrm{PH}_{\mathrm{start}}} = 3.0$ d	Kronenberg et al. (2020a)
		2018: $\mu_{t\mathrm{PH}_{\mathrm{start}}} = 101$ d, $\sigma_{t\mathrm{PH}_{\mathrm{start}}} = 3.1$ d	Roth (2021)
		2019: $\mu_{t\mathrm{PH}_{\mathrm{start}}} = 95$ d, $\sigma_{t\mathrm{PH}_{\mathrm{start}}} = 3.0$ d	Estimation based on 2016–2018 data
		2020: $\mu_{t\mathrm{PH}_{\mathrm{start}}} = 100$ d, $\sigma_{t\mathrm{PH}_{\mathrm{start}}} = 2.9.0$ d	Estimation based on 2016–2018 data
		2016: $\mu_{t\mathrm{PH}_{\mathrm{stop}}} = 165$ d, $\sigma_{t\mathrm{PH}_{\mathrm{stop}}} = 2.5$ d	Kronenberg et al. (2020a)
		2017: $\mu_{t\mathrm{PH}_{\mathrm{stop}}} = 162$ d, $\sigma_{t\mathrm{PH}_{\mathrm{stop}}} = 3.5$ d	Kronenberg et al. (2020a)
		2018: $\mu_{t\mathrm{PH}_{\mathrm{stop}}} = 158$ d, $\sigma_{t\mathrm{PH}_{\mathrm{stop}}} = 4.0$ d	Roth (2021)
		2019: $\mu_{t\mathrm{PH}_{\mathrm{stop}}} = 164$ d, $\sigma_{t\mathrm{PH}_{\mathrm{stop}}} = 3.0$ d	Estimation based on 2016–2018 data
		2020: $\mu_{t\mathrm{PH}_{\mathrm{stop}}} = 160$ d, $\sigma_{t\mathrm{PH}_{\mathrm{stop}}} = 3.1$ d	Estimation based on 2016–2018 data
u_{it}	AR(1)	$\rho = 0.9$, $\sigma = 0.003$ m d^{-1}	Assumption
e_t	$\mathcal{N}\left(\mu_m,\sigma_m^2\right)$	$\mu_m = 0$, $\sigma_m = 0.01$ m	Roth et al. (2020)

$\to$ Connects lower and upper limit of cardinal temperature ramps
AR(1): Autoregressive model of order 1.

Figure 2: Measured temperatures in the stem elongation phase of winter wheat at the field phenotyping platform site of ETH Zurich (FIP) on an hourly scale (T_{course}) and aggregated to a 3-day scale (T_{mean}) (a) and Pearson's correlation between simulation input and extracted parameter values in dependence on the measurement interval (b). Grey boxes in (a) indicate cardinal temperature ranges from terminal spikelet to anthesis according to Porter and Gawith (1999), early season corresponds to the first half of the stem elongation phase and late season to the second half. 4 days (irr.) in (b) corresponds to an intended measurement frequency of two measurements per week with frequent shifts by one, two or three days due to averse weather conditions, resulting in an irregular measurement frequency.

297 CH Claro, Suretta; Agroscope (Nyon, Switzerland) / DSP (Delley, Switzerland)). These genotypes were used as
298 spatial checks in 2017 and therefore replicated 18 times, thus resulting in 54 plot based time series.

4. Results

Aggregating measured temperatures at the field phenotyping platform site of ETH Zurich (FIP) to hourly
means (T_{course}) and 3-day means (T_{mean}) reveled severe differences in their distribution (Figure 2a): Temperatures
close to the cardinal temperature T_{min} and between the cardinal temperatures T_{opt} and T_{max} were frequent for
T_{course} in all years, but almost absent for T_{mean} in the years 2018 and 2020, and partly absent in 2016. While 2018
was a year with high temperatures throughout the stem elongation, 2020 was characterized by high temperatures
in the early elongation phase. 2016 and 2017 were years with distinct temperature differences between the early
and late season, while differences for 2019 were more moderate.

Simulating the stem elongation by combining the T_{course} measurements with a Wang-Engel dose-response
function resulted in an average final height slightly higher than observed for elite varieties under field condi-
tions by Kronenberg et al. (2020b) (Appendix, Figure 7), but realistic individual plant height time series with
characteristic starts, stops, and lag phases (Figure 3). Fitting the asymptotic model (Equation 3) based on T_{course}
(Equation 7) to the simulated data converged for 97% of all genotypes, fitting the model to real data of three
genotypes with 18 replications for 91% of all plots. Based on visual inspections of simulated versus fitted curves
(Figure 4a) the asymptotic model was able to model well the Wang-Engel curve increase in growth rate from T_{min}
to T_{opt} while systematically underestimating r_{max} and slightly overestimating T_{min}. Parameters extracted from real
data were in the range of simulation input values with slightly lower r_{max} and higher T_{min} values (Figure 4b).
Even tough for T_{min} the extracted values for the three genotypes were very close, significant differences between
genotypes for all three parameters T_{min}, r_{max} and lrc were found.

Fitting the cardinal temperature shift enhanced model (Equation 10) to data converged for 90% of all geno-
types. All other models converged for all genotypes. In a preliminary run, we also tried to fit the Wang-Engel
model to simulated data, but failed to extract meaningful parameters. This was partly due to the complexity
of defining valid non-overlapping ranges for the cardinal temperatures, but mainly because the model did not
converge in most cases.

Based on the RMSE, the base temperature T_{min} were best estimated with the asymptotic dose-response model
based on T_{course} covariates, while the maximum absolute growth rate r_{max} was best estimated with the asymptotic
dose-response model based on T_{mean} covariates (Table 2). While the variance was drastically lower for both
parameters if using T_{course}, the bias slightly increased for r_{max} but decreased for T_{min}. The RMSE, bias and
variance of the linear model for the parameter T_{min} was much higher than for the asymptotic model.

Introducing a random measurement error e had little effect on the correlation for the slope estimated by the
linear model; extracted values correlated almost 1:1 for the simulation with and without such noise (Table 2).
In contrast, the Pearson's r for T_{min} was close to zero, but the Spearman's rank correlation revealed nevertheless

Figure 3: Simulated growth rates (a) and canopy heights (b) for 200 out of 1000 simulated genotypes and three years. Simulated growth rates are based on a Wang-Engel dose-response curve model.

Figure 4: Simulated Wang-Engel dose-response curves and fitted asymptotic dose-response curves for four selected simulated genotypes (a), and real field data example with asymptotic curves fitted to TLS height data (terrestrial laser scanner) (Kronenberg et al., 2017) of three genotypes in 2017 (b). Lines and bars in (b) denote means of eighteen replications per genotype, error bars standard errors, letters significant differences ($\alpha = 0.05$) based on one-way ANOVAs and Tukey's Honest Significant Difference method.

a high robustness of the extracted ranking to noise. The asymptotic model was more susceptible to noise than the linear model, indicated by increasing RMSE, bias and variance for T_{min} and r_{max} . Using temperature courses instead of mean temperatures was of benefit for the parameters r_{max} and T_{min}.

Correlating simulated versus predicted parameter values revealed further differences between the methods (Table 2, Figure 5): While the asymptotic model based on T_{course} was able to extract r_{max} and T_{min} with very strong correlations to the input values, the same model based on T_{mean} yielded only moderate correlations for r_{max} and T_{min}.

The parameter r_{max} extracted by the asymptotic model based on T_{course} was not only correlated with the simulated r_{max}, but also weakly related to T_{min} and T_{opt} (Figure 5). Later correlations are at odds with the simulation input where T_{min}, T_{opt} and r_{max} were uncorrelated, and is therefore an indicator of a modeling artifact. The asymptotic model additionally allowed estimating T_{opt} using lrc as proxy with a moderate correlation of lrc to T_{opt} for T_{course}, while for T_{mean} the correlation was weak.

For the linear model, both parameter estimates of slope and T_{min} were uncorrelated with the corresponding simulated input parameter (Figure 5b). Instead, the slope was related to r_{max}, T_{min} and T_{opt}, while T_{min} based on the estimated intercept was not correlated to any input parameter. Both the linear and asymptotic model were not affected by supra-optimal temperatures: the input parameter T_{max} that defined growth in the supra-optimal range above T_{opt} was uncorrelated with any extracted parameter.

Adding noise to the simulated growth time series led to increased variances for the parameter estimates of r_{max} and T_{min} based on the asymptotic model (Table 2). While the RMSE increased by a factor of two for T_{min} and the correlation to input values dropped towards 0.5, it remained low for r_{max} while the correlation with input parameter dropped towards 0.3. The bias for r_{max} remained unchanged, while the bias for T_{min} decreased. Adding additional noise caused by highly auto-correlated spatial inhomogeneities did not further decrease the RMSE or increase the variance and bias.

Simulating changing cardinal temperatures for all genotypes alike decreased correlations of extracted parameters for all models (Figure 5c). The ranking for T_{mean} and T_{course} for the asymptotic model switched: While for both models the ability to extract T_{min} was drastically reduced, T_{mean} remained its ability to extract r_{max} but T_{course} lost it. The curvature parameter lrc was now moderately correlated with T_{min} and weakly with T_{opt} for T_{mean}, therefore reducing its suitability as proxy trait for T_{opt}. For T_{course} the correlation of lrc with T_{opt} remained moderate, therefore retaining its suitability as proxy trait for T_{opt}. The slope extracted by the linear model was moderately related to T_{min} and weakly to r_{max} and T_{opt}. Again, T_{max} was uncorrelated with any extracted parameter.

Integrating changing cardinal temperatures in the parameter extraction model turned out to be difficult (Appendix, Table 3): For r_{max} no improvement in parameter correlation and only slight improvements in RMSE, bias and variances were found. For T_{min} the correlation with input values slightly increased but remained weak, and RMSE, bias and variances slightly improved.

Table 2: Bias, variance, root-mean squared error, and correlations for the linear and asymptotic model based on mean temperature and temperature courses for simulated data without noise, with noise due to measurement errors, and with noise due to field and environment inhomogeneities. Reported correlations are Pearson's correlation between parameters extracted on simulated data and simulation inputs, and Pearson's and Spearman's correlation of the extracted parameters with and without noise due to measurement errors.

Parameter	Model	Noise	Bias	Var	RMSE	$\text{cor}_\text{P}(\hat{\theta}, \theta)$	$\text{cor}_\text{P}(\hat{\theta}_e, \hat{\theta})$	$\text{cor}_\text{S}(\hat{\theta}_e, \hat{\theta})$
a (slope)	Linear T_mean	-	0.0617	0.00244	0.0617	0.08		
		e	0.0625	0.00253	0.0625	0.08	0.94	0.94
		$e + u$	0.0625	0.00253	0.0625	0.08		
r_max	Asymptotic T_mean	-	0.0622	0.0412	0.0931	0.52		
		e	0.0965	0.187	0.140	0.30	0.35	0.62
		$e + u$	0.0967	0.189	0.140	0.30		
	Asymptotic T_course	-	0.0988	0.00628	0.101	0.82		
		e	0.0978	0.119	0.116	0.32	0.57	0.86
		$e + u$	0.098	0.115	0.116	0.33		
T_min	Linear T_mean	-	9.23	124e+03	9.67	0.02		
		e	8.78	222000	9.67	0.02	-0.02	0.84
		$e + u$	8.73	272000	9.61	0.00		
	Asymptotic T_mean	-	0.954	2.40	1.26	0.65		
		e	0.893	5.35	1.59	0.36	0.43	0.45
		$e + u$	0.896	5.46	1.57	0.35		
	Asymptotic T_course	-	-0.491	1.36	0.674	0.83		
		e	-0.394	5.28	1.29	0.51	0.71	0.73
		$e + u$	-0.369	5.43	1.32	0.50		
lrc	Asymptotic T_mean	-						
		e					0.48	0.49
	Asymptotic T_course	-						
		e					0.32	0.48

e: Random measurement error noise
u: Field and environment inhomogeneities
$\text{cor}_\text{P}(\hat{\theta}, \theta)$: Pearson's r between simulation input and extracted values
$\text{cor}_\text{P}(\hat{\theta}_e, \hat{\theta})$: Pearson's r between extracted values from simulation without noise (-) and with random measurement error (e)
$\text{cor}_\text{S}(\hat{\theta}_e, \hat{\theta})$: Spearman's r between extracted values from simulation without noise (-) and with random measurement error (e)

The importance of the measurement interval varied with year and parameter (Figure 2b). Overall, 2019 was best suited to extract all parameters, independent of the measurement interval. For T_{min}, in 2019 the measurement interval effect was almost negligible. For all other years, differences between three days intervals and four days irregular intervals were small for T_{min}, while larger measurement intervals (7 and 14 days) severely reduced the correlations with input parameters. The same pattern, but even stronger reductions between four and seven days intervals, were visible for r_{max}, except for 2019 where the correlation dropped only for a 14 days interval. For lrc, all reductions including the one from three days to four days reduced the correlations, and differences between the years were small.

5. Discussion

Although the growth of simulated canopies was based on a Wang-Engel temperature dose-response curve, using a corresponding model to extract parameters turned out to be not suitable, mainly because of failed convergence. Presumably, the low frequency of measurement values for supra optimal temperatures ($\gtrsim$ 30 °C, Figure 2) prevented estimating the required parameter T_{max}. As suspected, the asymptotic model performed better with percentages of converged time series close to 100%. Using temperature courses instead of aggregated mean values allowed estimating independent parameters with higher correlations to input parameters, to the cost of a slightly decreased convergence rate. Especially the parameter T_{min} profited from taking temperature courses into account (Table 2), which indicates that in the early phase where stem elongation starts the distribution of temperatures and therefore temperature courses are essential to describe temperature responses ($\lesssim$ 5 °C, Figure 2a). This thesis is also supported by the finding that years with a rather cold spring (2016, 2017, 2019) were better suited to extract T_{min} (Figure 2). Consequently, the use of the T_{course} method can provide dose-response curve models with valuable information about the occurrence of temperature values at the extremes, while the T_{mean} method leads to a shrinking of these values towards the mean.

Additionally to T_{min}, the asymptotic model will extract the maximum absolute growth rate r_{max} and the parameter lrc which is related to steepness of the curvature of the asymptotic model. Accordingly, the parameter relates to the optimum temperature T_{opt}. Thus, the model can fully describe temperature responses between the lower base temperature T_{min} and optimum temperature T_{opt}. A limitation of the proposed asymptotic approach may be the high optimum temperatures: Although the extracted base temperature T_{min} and maximum growth rate r_{max} were strongly correlated with input parameters, they were also correlated to the optimum temperature T_{opt}. As the parameters were simulated as uncorrelated, this indicates a modeling bias. Consequently, high optimum temperatures may lead to overestimation of T_{min} and r_{max}

Noise and environment inhomogeneities reduced the correlation to input parameters drastically, independent on whether RMSE stayed stable or not (Table 2). It seems reasonable to assume that this reduction in correlation will lead to reduced estimated heritabilities of measured traits in HTFP. The simulation was not including a randomized field experiment set-up, but only independent genotype time series. Nevertheless, we assume based

Figure 5: Pearson's correlations of simulated data versus extracted temperature response parameters. Provided are results for the simulation without noise (a) versus extracted temperature response parameters with fixed cardinal temperatures (b) and changing cardinal temperatures with time (c) for the linear model based on mean temperature (Linear T_{mean}) and the asymptotic growth model based on mean temperature (Asym T_{mean}) and temperature courses (Asym T_{course}). Note that the slope is derived from the relation slope $= r_{\mathrm{max}}/T_{\mathrm{opt}} - T_{\mathrm{min}}$ and therefore not an independent input parameter of the simulation. Black bold boxes indicate correlations between predicted and true values for identical parameters.

on our previous study on two other intermediate traits (Roth et al., 2021b) that also for dose-response curve traits it is essential to follow the three principles of experimental design, i.e., replication, randomization and blocking (Kempton, 1984). Such an experimental design will allow processing the extracted plot-based parameters using, e.g., linear mixed models to compensate for measurement errors and environment inhomogeneities and therefore ensure the overall heritability.

Simulating changing cardinal temperatures drastically reduced the ability of the asymptotic model to extract mean cardinal temperatures (Figure 5c). We suspect that this effect was overamplified by the simulation and that supposed cardinal temperature ramps in Porter and Gawith (1999) may be too extreme for wheat elite cultivar sets. In such an extreme ramp case, the asymptotic model would yield less heritable parameters than in the case of fixed cardinal temperatures, but would not introduce additional biases. Including an additional parameter to allow a shift of $T_{\min}$ with time only slightly improved the estimations (Appendix, Figure 3). In addition to a reduced convergence rate due to the increased number of parameters, introducing such a parameter also implies assuming the form of an underlying 'biological' function, which may further increases uncertainties as the exact form of the function is (yet) unknown. In addition to changes in cardinal temperatures, one may assume that $r_{\max}$ changes with time as well, but we are not aware of publications that would allow us to incorporate such changes in the simulation and extraction model.

A 'close-to-optimum' year 2019 was found in this study that was best suited to extract temperature dose-response parameters. The year was characterized by mild temperatures and clearly distinct temperature distributions for the early and late season of stem elongation. The early season covered well temperatures around $T_{\min}$, while late season temperatures ranged primarily around T_{opt} but also covered temperatures close to $T_{\max}$ (Figure 2a). In less optimal years, the chosen measurement interval plays an essential role (Figure 2b). Considering the fact that a field crop researcher cannot anticipate the suitability of a coming season to extract response parameters, an interval of at maximum 4 days is advisable.

The literature-based cardinal temperatures together with the sampled temperature courses during stem elongation suggest that T_{opt} but not $T_{\max}$ is reached at numerous hours. Thus, the asymptotic model proved to be superior compared to the more complex Wang-Engel model including $T_{\max}$, but also superior compared to the simpler linear model suitable for temperature ranges above $T_{\min}$ but well-below T_{opt}. The former did not converge while the latter introduced substantial bias with regard to the estimation of cardinal temperatures. The linear response to temperature (slope) was similarly related to both cardinal temperatures $T_{\min}$ and T_{opt} but also to the maximum growth rate $r_{\max}$, but unrelated to the simulated slope (Figure 5b). Thus, the apparent temperature response is mainly driven by cardinal temperatures. When fitting a linear model to a wide temperature range, with significant number of hours in temperature ranges beyond the optimum (Figure 2a), it will inevitably compensate for both the maximum growth rate at the optimum and the base temperature of growth. Using a linear model if cardinal temperatures are changing during the season is even more harmful: The relation of the slope may switch from T_{opt} towards $T_{\min}$ (Figure 5c). Depending on the examined genetic material, applying a linear

model may therefore lead to completely different conclusions.

Kronenberg et al. (2020a) applied such linear functions to wheat canopy height data. The authors argued that
based on average temperatures in the measurement intervals, the temperature optimum T_{opt} of 27 °C (Parent
and Tardieu, 2012) may not be reached. Nevertheless, our results show that the lower T_{opt} of 20.3 °C (Porter and
Gawith, 1999) is reached at many hours during stem elongation, indicating that the slope reported by Kronenberg
et al. (2020a) may be partly affected by growth at optimum temperature and the maximum growth rate. In
contrast, the asymptotic model was able to mimic the curvature of a Wang-Engel model for temperature ranges
below the optimum temperature (Figure 4a) and extracted differing values for r_{max} and *lrc* for three genotypes
measured in the field (Figure 4b).

Whether an asymptotic model would also be more appropriate to model early canopy development (Slafer
and Rawson, 1995; Grieder et al., 2015; Nagelmüller et al., 2016) remains questionable. Instead, the issue may
rather be to model zero growth values below T_{min} appropriately. In summary, while a wide range of functions
is available to model temperature-response, it is critical to know which cardinal temperature are reached under
field conditions during the growth phase of interest, and consequently to use a function that is sufficiently flexible
in critical areas of the affected temperature range.

6. Conclusion

Adequate models to quantify the different aspects of growth response to temperature may greatly improve
our understanding of crop adaptation to certain climatic scenarios. Asymptotic models are a valid choice, if
temperature ranges cover T_{min} and T_{opt} and the occurrence of supra-optimal temperatures during the observed
growth phase is negligible. In contrast, linear models should be avoided in such situations. Using temperature
courses with high temporal resolution as covariate input instead of aggregated mean temperatures is strongly
advisable. The presence of changing cardinal temperatures—related to advancing phenology—may reduce the
ability to extract parameters with an asymptotic model but will not severely bias results.

HTFP measurement intervals of about four days are sufficient to reliably extract T_{min} and r_{max} in an average
season. Some irregularity in the frequency due to measuring breaks related to unsuitable weather conditions
does not greatly affect the precision. Yet, *lrc* may profit from a higher measuring frequency. Not every season is
equally suitable for characterizing dose-response curve parameters from repeated measurements, but an unsuit-
able season can be partially compensated for by increasing the measuring frequency. Consequently, we propose
the following rule of thumb: "For dose-response modeling, one shall measure at least twice a week, but rest on
weekends".

Model and Data availability

466 Data and source code that support the findings of this study are openly available in the ETH gitlab reposi-
467 tory at https://gitlab.ethz.ch/crop_phenotyping/htfp_data_processing and archived in the ETH

468 research collection (http://doi.org/10.5905/ethz-1007-385).

Authorship contribution statement

470 **Lukas Roth**: Conceptualization, Methodology, Software, Formal analysis, Visualization, Writing - Original
471 Draft. **Hans-Peter Piepho**: Conceptualization, Methodology, Writing - Review & Editing. **Andreas Hund**: Con-
472 ceptualization, Supervision, Project administration, Funding acquisition, Writing - Review & Editing.

Acknowledgements

474 We like to acknowledge Lukas Kronenberg for feedback on an early version of the manuscript, and Norbert
475 Kirchgessner for feedback on formula notations (both ETH Zurich).
476 LR received funding from Innosuisse (http://www.innosuisse.ch) in the framework for the project "Trait
477 spotting" (grant number: KTI P-Nr 27059.2 PFLS-LS). HPP was supported by DFG grant PI 377/24-1.

Appendix

Table 3: Bias, variance, root-mean squared error, and Pearson's correlation ($\text{cor}_\text{P}(\hat{\theta}, \theta)$) for values extracted from simulated data with noise and changing cardinal temperatures with the asymptotic model based on temperature courses (T_course) and with an asymptotic model that additionally models the slope of changing cardinal temperatures ($T_\text{course\&ramp}$).

Parameter	Model	Noise	Bias	Var	RMSE	$\text{cor}_\text{P}(\hat{\theta}, \theta)$
r_max	Asymptotic T_course	$u + e + \text{ramp}$	0.144	0.179	0.176	0.28
	Asymptotic $T_\text{course\&ramp}$		0.134	0.207	0.168	0.28
T_min	Asymptotic T_course	$u + e + \text{ramp}$	1.91	11.2	2.21	0.16
	Asymptotic $T_\text{course\&ramp}$		1.86	10.6	2.20	0.23

Data generation

Data extraction

Figure 6: Schematic representation of the workflow. Top: Data generation using a simulation based on a Wang-Engel function parametrised with real temperature courses and cardinal temperatures from literature. Bottom: Data extraction using a linear model and aggregated temperature values (Kronenberg et al., 2020a) and the herein proposed new approach using an asymptotic model directly applied to covariate courses with high temporal resolution.

Figure 7: Simulated and measured plant heights for 2017. True measurement values were captured using a terrestrial laser scanner (Kronenberg et al., 2017) on a subset of the GABI wheat planel (350 genotypes). Simulated measurements values are based on 1000 assumed genotypes.

References

J. L. Araus, S. C. Kefauver, M. Zaman-Allah, M. S. Olsen, and J. E. Cairns. Translating High-Throughput Phenotyping into Genetic Gain. *Trends in Plant Science*, 23(5):451–466, 2018. ISSN 13601385. doi: 10.1016/j.tplants.2018.02.001.

R. Bonhomme. Bases and limits to using 'degree.day' units. *European Journal of Agronomy*, 13:1–10, 2000. ISSN 11610301. doi: 10.1016/S1161-0301(00)00058-7.

N. Brisson, P. Gate, D. Gouache, G. Charmet, F. X. Oury, and F. Huard. Why are wheat yields stagnating in Europe? A comprehensive data analysis for France. *Field Crops Research*, 119:201–212, 2010. ISSN 03784290. doi: 10.1016/j.fcr.2010.07.012.

D. R. Causton and M. P. Dale. The monomolecular and rectangular hyperbola as empirical models of the response of photosynthetic rate to photon flux density, with applications to three Veronica species. *Annals of Botany*, 65:389–394, 1990. ISSN 03057364. doi: 10.1093/oxfordjournals.aob.a087949.

J. A. Clavijo Michelangeli, T. R. Sinclair, and N. Bliznyuk. Using an Arrhenius-type function to describe temperature response of plant developmental processes: Inference and cautions. *New Phytologist*, 210(2):377–379, 2016. ISSN 14698137. doi: 10.1111/nph.13812.

J. N. Gallagher, P. V. Biscoe, and J. S. Wallace. Field studies of cereal leaf growth: IV. Winter wheat leaf extension in relation to temperature and leaf water status. *Journal of Experimental Botany*, 30(117):657–668, 1979. ISSN 00220957. doi: 10.1093/jxb/30.4.657.

C. Grieder, A. Hund, and A. Walter. Image based phenotyping during winter: a powerful tool to assess wheat genetic variation in growth response to temperature. *Functional Plant Biology*, 42:387–396, 2015. ISSN 1445-4408. doi: 10.1071/fp14226.

M. Heinen. Analytical growth equations and their Genstat 5 equivalents. *Netherlands Journal of Agricultural Science*, 47:67–89, 1999. ISSN 00282928. doi: 10.18174/njas.v47i1.479.

A. Hund, R. Reimer, P. Stamp, and A. Walter. Can we improve heterosis for root growth of maize by selecting parental inbred lines with different temperature behaviour? *Philosophical Transactions of the Royal Society B: Biological Sciences*, 367:1580–1588, 2012. ISSN 14712970. doi: 10.1098/rstb.2011.0242.

F. H. Johnson, H. Eyring, and R. W. Williams. The nature of enzyme inhibitions in bacterial luminescence: sulfanilamide, urethane, temperature and pressure. *Journal of Cellular and Comparative Physiology*, 20(3):247–268, 1942.

R. Kempton. The design and analysis of unreplicated field trials. *Vorträge für Pflanzenzüchtung*, 7:219–242, 1984.

L. Kronenberg, K. Yu, A. Walter, and A. Hund. Monitoring the dynamics of wheat stem elongation: genotypes differ at critical stages. *Euphytica*, 213(157), 2017. doi: 10.1007/s10681-017-1940-2.

L. Kronenberg, S. Yates, M. P. Boer, N. Kirchgessner, A. Walter, and A. Hund. Temperature response of wheat affects final height and the timing of stem elongation under field conditions. *Journal of Experimental Botany*, 2020a. doi: 10.1093/jxb/eraa471.

L. Kronenberg, S. Yates, S. Ghiasi, L. Roth, M. Friedli, M. E. Ruckle, R. A. Werner, F. Tschurr, M. Binggeli, N. Buchmann, B. Studer, and A. Walter. Rethinking temperature effects on leaf growth, gene expression and metabolism: Diel variation matters. *Plant, Cell and Environment*, pages 1–15, 2020b. ISSN 0140-7791. doi: 10.1111/pce.13958.

F. Laidig, H.-P. Piepho, D. Rentel, T. Drobek, U. Meyer, and A. Huesken. Breeding progress, environmental variation and correlation of winter wheat yield and quality traits in German official variety trials and on-farm during 1983–2014. *Theoretical and Applied Genetics*, 130:223–245, 2017. doi: 10.1007/s00122-016-2810-3.

S. Nagelmüller, N. Kirchgessner, S. Yates, M. Hiltpold, and A. Walter. Leaf Length Tracker: A novel approach to analyse leaf elongation close to the thermal limit of growth in the field. *Journal of Experimental Botany*, 67(6):1897–1906, 2016. ISSN 14602431. doi: 10.1093/jxb/erw003.

C. E. T. Paine, T. R. Marthews, D. R. Vogt, D. Purves, M. Rees, A. Hector, and L. A. Turnbull. How to fit nonlinear plant growth models and calculate growth rates: An update for ecologists. *Methods in Ecology and Evolution*, 3:245–256, 2012. ISSN 2041210X. doi: 10.1111/j.2041-210X.2011.00155.x.

B. Parent and F. Tardieu. Temperature responses of developmental processes have not been affected by breeding in different ecological areas for 17 crop species. *New Phytologist*, 194:760–774, 2012. ISSN 0028646X. doi: 10.1111/j.1469-8137.2012.04086.x.

B. Parent, O. Turc, Y. Gibon, M. Stitt, and F. Tardieu. Modelling temperature-compensated physiological rates, based on the co-ordination of responses to temperature of developmental processes. *Journal of Experimental Botany*, 61(8):2057–2069, 2010. ISSN 00220957. doi: 10.1093/jxb/erq003.

B. Parent, E. J. Millet, and F. Tardieu. The use of thermal time in plant studies has a sound theoretical basis provided that confounding effects are avoided. *Journal of Experimental Botany*, 70(9):2359–2370, 2019. ISSN 14602431. doi: 10.1093/jxb/ery402.

D. M. Pérez-Valencia, M. X. Rodríguez-Álvarez, M. P. Boer, L. Kronenberg, A. Hund, L. C. Bosquet, E. J. Millet, and F. A. van Eeuwijk. A two-stage approach for the spatio-temporal analysis of high-throughput phenotyping data. *Scientific Reports*, 12(3177), 2022. doi: 10.1038/s41598-022-06935-9.

H. P. Piepho and E. R. Williams. Linear variance models for plant breeding trials. *Plant Breeding*, 129:1–8, 2010. ISSN 01799541. doi: 10.1111/j.1439-0523.2009.01654.x.

H.-P. Piepho and E. R. Williams. Augmented row–column designs for a small number of checks. *Agronomy Journal*, 108(6):2256–2262, 2016. ISSN 14350645. doi: 10.2134/agronj2016.06.0325.

J. C. Pinheiro and D. M. Bates. *Mixed-Effects Models in S and S-PLUS*. Springer-Verlag New York, New York, 2000. ISBN 0-387-98957-9.

J. R. Porter and M. Gawith. Temperatures and the growth and development of wheat a review. *European Journal of Agronomy*, 10:23–36, 1999. doi: 10.1016/S1161-0301(98)00047-1.

J. R. Porter and J. H. Rayner. A winter wheat crop simulation model without water or nutrient limitations. *The Journal of Agricultural Science*, 102:371–382, 1984. ISSN 14695146. doi: 10.1017/S0021859600042702.

J. Ramirez-Villegas, J. Watson, and A. J. Challinor. Identifying traits for genotypic adaptation using crop models. *Journal of Experimental Botany*, 66(12):3451–3462, 2015. ISSN 14602431. doi: 10.1093/jxb/erv014.

F. Rebaudo and V. B. Rabhi. Modeling temperature-dependent development rate and phenology in insects: review of major developments, challenges, and future directions. *Entomologia Experimentalis et Applicata*, 166:607–617, 2018. ISSN 15707458. doi: 10.1111/eea.12693.

F. Rebaudo, Q. Struelens, and O. Dangles. Modelling temperature-dependent development rate and phenology in arthropods: The devRate
package for R. *Methods in Ecology and Evolution*, 9:1144–1150, 2018. ISSN 2041210X. doi: 10.1111/2041-210X.12935.
G. J. Rebetzke, J. Jimenez-Berni, R. A. Fischer, D. M. Deery, and D. J. Smith. Review: High-throughput phenotyping to enhance the use of
crop genetic resources. *Plant Science*, 282:40–48, 2019. doi: 10.1016/j.plantsci.2018.06.017.
R. Reimer, B. Stich, A. E. Melchinger, T. A. Schrag, A. P. Sørensen, P. Stamp, and A. Hund. Root response to temperature extremes: Association
mapping of temperate maize (Zea mays L). *Maydica*, 58:156–168, 2013. ISSN 22798013.
L. Roth. *Development of drone-based phenotyping methodologies to support physiological plant breeding of wheat and soybean*. PhD thesis, mar
2021. URL https://www.research-collection.ethz.ch:443/handle/20.500.11850/474271.
L. Roth, M. Camenzind, H. Aasen, L. Kronenberg, C. Barendregt, K.-H. Camp, A. Walter, N. Kirchgessner, and A. Hund. Repeated Multiview
Imaging for Estimating Seedling Tiller Counts of Wheat Genotypes Using Drones. *Plant Phenomics*, 2020(3729715), 2020. doi: 10.34133/
2020/3729715.
L. Roth, M. X. Rodríguez-Álvarez, F. van Eeuwijk, H.-P. Piepho, and A. Hund. Phenomics data processing: A plot-level model for repeated
measurements to extract the timing of key stages and quantities at defined time points. *bioRxiv*, page 2021.05.02.442243, jan 2021a. doi:
10.1101/2021.05.02.442243.
L. Roth, M. X. Rodríguez-Álvarez, F. van Eeuwijk, H.-P. Piepho, and A. Hund. Phenomics data processing: A plot-level model for repeated
measurements to extract the timing of key stages and quantities at defined time points. *Field Crops Research*, 274(108314), 2021b. doi:
10.1016/j.fcr.2021.108314.
G. A. Slafer and H. M. Rawson. Rates and cardinal temperatures for processes of development in wheat: Effects of temperature and thermal
amplitude. *Australian Journal of Plant Physiology*, 22:913–926, 1995. ISSN 03107841. doi: 10.1071/PP9950913.
F. Tardieu, I. S. C. Granato, E. J. Van Oosterom, B. Parent, and G. L. Hammer. Are crop and detailed physiological models equally 'mechanistic'
for predicting the genetic variability of whole-plant behaviour? The nexus between mechanisms and adaptive strategies. *in silico Plants*,
2(1), 2020. doi: 10.1093/insilicoplants/diaa011.
F. A. van Eeuwijk, D. Bustos-Korts, E. J. Millet, M. P. Boer, W. Kruijer, A. Thompson, M. Malosetti, H. Iwata, R. Quiroz, C. Kuppe, O. Muller,
K. N. Blazakis, K. Yu, F. Tardieu, and S. C. Chapman. Modelling strategies for assessing and increasing the effectiveness of new phenotyping
techniques in plant breeding. *Plant Science*, 282:23–39, 2019. ISSN 18732259. doi: 10.1016/j.plantsci.2018.06.018.
E. Wang and T. Engel. Simulation of phenological development of wheat crops. *Agricultural Systems*, 58(1):1–24, 1998. ISSN 0308521X.
doi: 10.1016/S0308-521X(98)00028-6.
E. Wang, P. Martre, Z. Zhao, F. Ewert, A. Maiorano, R. P. Rötter, B. A. Kimball, M. J. Ottman, G. W. Wall, J. W. White, M. P. Reynolds, P. D.
Alderman, P. K. Aggarwal, J. Anothai, B. Basso, C. Biernath, D. Cammarano, A. J. Challinor, G. De Sanctis, J. Doltra, E. Fereres, M. Garcia-
Vila, S. Gayler, G. Hoogenboom, L. A. Hunt, R. C. Izaurralde, M. Jabloun, C. D. Jones, K. C. Kersebaum, A. K. Koehler, L. Liu, C. Müller,
S. Naresh Kumar, C. Nendel, G. O'Leary, J. E. Olesen, T. Palosuo, E. Priesack, E. Eyshi Rezaei, D. Ripoche, A. C. Ruane, M. A. Semenov,
I. Shcherbak, C. Stöckle, P. Stratonovitch, T. Streck, I. Supit, F. Tao, P. Thorburn, K. Waha, D. Wallach, Z. Wang, J. Wolf, Y. Zhu, and
S. Asseng. The uncertainty of crop yield projections is reduced by improved temperature response functions. *Nature Plants*, 3(17102),
2017. ISSN 20550278. doi: 10.1038/nplants.2017.102.
Y. Weikai and L. A. Hunt. An Equation for Modelling the Temperature Response of Plants using only the Cardinal Temperatures. *Annals of
Botany*, 84:607–614, 1999. doi: 10.1006/anbo.1999.0955.

25