1 **Fecal Microbiota Transplantation Drives Colonic Expression of Immune**

2 **Activation Genes in a Mouse Model of Antibiotic Use**

G. Brett Moreau[1], Hale Ozbek[2], Pankaj Kumar[3], Alyse Frisbee[4], Jhansi Leslie[1],

William A. Petri, Jr.[1,2,4]

[1]Department of Medicine, Infectious Diseases and International Health, University

of Virginia School of Medicine, Charlottesville, VA, USA

[2]Department of Pathology, University of Virginia School of Medicine,

Charlottesville, VA, USA

[3]Department of Biochemistry and Molecular Genetics, University of Virginia School

of Medicine, Charlottesville, VA, USA

[4]Department of Microbiology, Immunology, and Cancer Biology, University of

Virginia School of Medicine, Charlottesville, VA, USA

15 **Abstract**

Clostridioides difficile infection (CDI) is the leading hospital acquired infection in

North America. While the standard treatment for CDI remains antibiotics, fecal

microbiota transplantation (FMT) has gained attention as an effective therapy to

prevent relapse. Previous work has focused on colonization resistance mounted

against *C. difficile* by FMT-delivered commensals, but the effects of FMT on the gut

mucosal immune response are poorly understood. Better understanding of the

molecular mechanisms driven by FMT would allow for more targeted therapy

against CDI. To address this important gap in knowledge, microbial community

structure and host gene expression were assessed after FMT in a mouse model of

antibiotic use. Administration of FMT led to a significant increase in microbial

diversity and partial restoration of community structure within 48 hours of

treatment. RNA sequencing of cecal tissue identified large changes in gene

expression between FMT recipient and vehicle control groups. Strikingly, genes

upregulated after FMT treatment were enriched in immune activation pathways,

many of which were associated with pro-inflammatory immune responses. FMT

also upregulated several genes associated with type 2 immunity while repressing

several associated with type 3 immunity, trends that are associated with improved

response to CDI. These results highlight the interplay between the intestinal

microbiota and host transcriptome and identify pathways of interest for exploring

the role of FMT on treatment of recurring CDI.

Introduction

Clostridioides difficile is a Gram-positive spore-forming obligate anaerobe that

causes mild to severe diarrheal disease. In the CDC's 2019 *Antibiotic Resistance*

Threats Report, *C. difficile* was ranked as an urgent threat to human health (1).

The increased prominence of *C. difficile* infection (CDI) is due in part to the

emergence in 2005 of hypervirulent strains (2), which cause more severe disease

and longer hospital stays (3–5). In a single year, *C. difficile* was associated with

nearly half a million infections, 83,000 recurrences and 29,000 deaths in the United

States alone (6). While *C. difficile* is one of the most common hospital-acquired

infections (7), community-acquired infections are also becoming more prevalent

(8). Due to the significant health impact and cost of CDI, development of highly

effective treatments is critical. Risk of CDI is associated with disruption of the

intestinal microbiota through antibiotic exposure (9, 10), which allows *C. difficile* to

bloom and produce toxins, resulting in initiation of diarrhea, damage to the

epithelium, and disruption of the intestinal barrier (7). Thus, maintaining or

restoring microbiota homeostasis is critical for minimizing the risk of CDI.

While the standard treatment for CDI remains antibiotics (11), one in five patients

experience recurrence (12), and antibiotic treatment further disrupts the intestinal

microbiota, increasing susceptibility for relapse. Because of this, fecal microbiota

transplantation (FMT), the administration of the whole microbial community from

the stool of a healthy donor into the patient's gastrointestinal tract, has been

explored as a therapeutic for CDI. While FMT has been implicated as therapeutic

in a wide range of disorders (13), it has proved most successful as treatment for

CDI, with an efficacy of over 90% (14). Despite the efficacy of FMT at treating CDI,

it is still considered an experimental therapy due to concerns of FMT donor

standardization, the evolving regulation of FMT, long-term side effects, and risk of

pathogen transmission, particularly for immunocompromised patients (15). FMT

has been associated with transmission of antibiotic resistant pathogens to

recipients, including one fatality (16). Given the prevalence of recurrent CDI and

the potential risks posed by FMT, more defined therapeutics for CDI are

necessary. Better understanding of the protective mechanisms behind FMT will

allow for more targeted therapeutics such as a defined consortium of probiotic

organisms or a defined cocktail of immunotherapeutic drugs as a replacement for

FMT.

Because *C. difficile* disease is initiated by antibiotic disruption of the intestinal

microbiota (9, 10), much of the focus of FMT has been on its ability to provide

colonization resistance against *C. difficile* (17). Recurrent CDI is characterized by

decreased microbial diversity that is associated with depletion of the *Bacteroidota*

and *Firmicutes* phyla and spikes in *Proteobacteria* (18), and these changes in

diversity can be restored by FMT (19). FMT may also work by influencing the

immune response in the gut to facilitate *C. difficile* clearance and prevent

excessive host damage. It has recently been more greatly appreciated that the

type of host immune response can impact CDI severity (20). While *C. difficile*

toxins cause much of the virulence and epithelial barrier damage seen in CDI, an

especially robust immune response and resulting inflammation has been implicated

in severe disease and worse clinical outcome independently of bacterial burden

(21–23). Specific types of immune responses have also been implicated in

differential CDI outcomes: tissue-regulated type 2 immunity has been associated

with tissue repair and protection from severe disease (24, 25), while a type 3

immune response has been implicated in increased disease severity (26).

Balancing these immune types to produce a response that facilitates *C. difficile*

clearance while minimizing tissue damage is critical for decreased disease

severity. As components from the microbiota and microbially-derived metabolites

can influence immune responses (20), this may be another mechanism through

which FMT drives resistance to CDI.

More mechanistic understanding of FMT's impact on the gut is required to develop

more defined therapeutics. While the ability of FMT to increase microbial diversity

and restore microbial community composition has been well studied (27–30), its

effect on the host transcriptome remains poorly understood. Further limiting our

understanding of FMT is that many studies have investigated FMT in the context of

other intestinal disorders, such as intestinal colitis models (30–32). In addition,

studies often look well after FMT administration (33), making it difficult to

distinguish the effects of FMT from the effects of gradual microbiota recovery in the

absence of antibiotics. This study aims to address these gaps in knowledge by

investigating the immediate impact of FMT in the context of antibiotic depletion of

the host microbiota. The primary findings of this study demonstrate that FMT

elicited significant changes in microbial composition and gene expression within 48

hours of administration and that changes in gene expression were enriched in host

immune activation gene sets. In particular, changes to immune genes suggest an

upregulation of pro-inflammatory and resolving type 2 immune signatures while

suppressing type 3 activation. This work highlights the crosstalk between the

intestinal microbiota and immune responses in the gut, identifying a potential

mechanism by which FMT could protect against CDI.

Results

FMT Partially Restores Microbiome Diversity after Antibiotic Treatment

To assess the effect of FMT on microbiome composition and host gene

expression, mice were given antibiotics to disrupt the gut microbiota, then treated

with either FMT or vehicle control for two days. Forty-eight hours after the initial

FMT administration, FMT recipient, vehicle control, and FMT donor mice were

sacrificed (Figure 1). Cecal contents and cecal tissue samples were collected for

analysis of microbiota community structure and host gene expression, respectively.

First, the V4-16S rRNA gene was sequenced using DNA from cecal contents to

examine microbial community structure. Analysis identified 566 unique amplicon

sequence variants (ASVs) in these samples. Community diversity in each group

was assessed with the hypothesis that antibiotic treatment would decrease

community diversity and that diversity would be restored by FMT.

The alpha diversity of the microbial community was first assessed by looking at

both the overall ASVs observed as a measure of sample richness as well as

Shannon's diversity index, which accounts for both richness and evenness. We

observed significant differences between all groups (Figure 2A) as measured by a

pairwise Wilcoxon Rank test with Bonferroni correction for multiple comparisons.

Treatment with the antibiotic cocktail was sufficient to disrupt diversity in the

vehicle control group relative to the FMT donor group, which was not treated with

antibiotics. While FMT administration partially restored diversity, it was unable to

reach the level observed in the FMT donor group.

Beta diversity between samples was also assessed using Unweighted Unifrac

distance, which takes into account phylogenetic distance between ASVs. Similar to

the results from alpha diversity metrics, FMT donor, vehicle control, and FMT

recipient groups all clustered separately. In addition, the FMT recipient group

clustered more closely with the FMT donor group along Axis 1, which accounted

for most of the variation in the model (Figure 2B). Several FMT recipient samples

clustered with the vehicle control group, and these same samples also had lower

alpha diversity values more similar to those seen in control samples. It is possible

that the FMT failed to establish in these animals, which may account for why they

are more similar to the vehicle control than other FMT recipient samples. Similar

results were observed using Bray-Curtis Dissimilarity index as a different measure

of beta diversity (Supplemental Figure 1), indicating that this outcome is not

dependent on bacterial phylogeny. These data indicate that FMT administration

partially restored community diversity to levels more similar to those seen in mice

that did not receive antibiotics.

FMT Restores Community Composition by Increasing Relative Abundance of

Firmicutes

Next, community composition was assessed to identify relative changes at the

phylum and family levels between groups. The FMT donor group was composed

primarily of members of the *Firmicutes* phylum, with the *Bacteroidota* phylum

making up a minor proportion (Figure 2C) The *Firmicutes* phylum was a smaller

proportion of the antibiotic-treated vehicle control samples relative to the FMT

donor group, with antibiotic-treated mice having relatively higher abundance of

Verrucomicrobiota, *Bacteroidota*, and *Proteobacteria.* FMT administration after

antibiotic treatment restored a community composition more closely resembling the

FMT donor group, with increased *Firmicutes* and decreased *Bacteroidota*,

Verrumicrobiota, and *Proteobacteria* relative to the vehicle control group.

Similar differences in community composition were observed at the Family level,

with the relative abundance of multiple families being intermediate in the FMT

recipient group between the FMT donor and vehicle control groups (Figure 2D).

The most prominent families observed in the FMT donor group were

Lachnospiraceae, *Muribaculaceae*, and *Oscillospiraceae*, whereas these families

made up a minority of the vehicle control samples in favor of relative increases in

the proportion of *Akkermansiaceae*, *Bacteroidaceae*, and *Enterococcaceae*.

Relative to the vehicle control, mice that received FMT had increases in the

proportion of *Lachnospiraceae*, *Muribaculaceae*, and *Oscillospiracea* as well as

decreases in *Akkermansiaceae*, *Bacteroidaceae*, and *Enterococcaceae*, leading

them to a more similar community composition to that seen in the FMT donor

group. Overall, these data indicate that administration of FMT partially restored the

microbial community in antibiotic-treated mice to a similar composition to that of the

FMT donor.

Finally, a random forest model was generated to identify ASVs that best

discriminated between vehicle control and FMT recipient groups. The model

performed well, with a 14% out-of-bag error rate. Of note, the model correctly

classified all samples except for the three FMT recipient samples that exhibited

lower diversity and clustered with Vehicle control samples. Of the top 15 ASVs

selected by the random forest model, 87% (13/15) were more prevalent in the FMT

recipient group relative to controls (Figure 2E), with most of these ASVs from the

Firmicutes phylum. At the genus level, FMT recipient samples had increased

relative abundance of *Lactobacillus ssp.* and *Erysipelatoclostridium ssp.*, while

Akkermansia muciniphilia and *Enterococcus ssp.* had higher relative abundance in

the control samples relative to FMT recipients. The top 15 ASVs were also

enriched in ASVs from the *Lachnospiraceae* family, which were largely absent from

vehicle control samples but present in the FMT recipient group. These data

highlight the significance of restoration of the *Firmicutes* phylum after FMT.

FMT Significantly Changes the Host Transcriptome

The immediate effect of FMT on host gene expression was next assessed. Cecal

contents from a subset of mice (four mice/group) were selected for whole RNA

sequencing. Of note, all four FMT recipient samples came from mice that displayed

increased alpha and beta diversity metrics relative to vehicle controls. Community

diversity and composition analysis in this subset of samples was representative of

diversity and composition observed in the full data set (Supplemental Figure 2).

Principal component analysis (PCA) was first performed on normalized, variance-

stabilized and log_2-transformed count data to further check data quality and assess

similarity between groups. FMT donor samples clustered separately from vehicle

control and FMT recipient samples along PC2, which likely indicates the effect of

antibiotic treatment on transcription (Figure 3A). While there was overlap between

vehicle control and FMT recipient samples, FMT recipient samples appeared to

cluster more closely with FMT donor samples along PC1, which accounted for

most of the total variation in the PCA model. These results suggest that the most

prominent factor distinguishing RNA sequencing results from different samples was

antibiotic treatment. While differences in host gene expression were less

pronounced between FMT recipient and vehicle control samples than differences in

community diversity, there appeared to be some clustering between these

samples, with the FMT recipient group appearing more similar to FMT donors.

Previous work from our lab has demonstrated that type 2 immune responses,

particularly the alarmins IL-25 and IL-33, are protective against CDI, in part by

activating group 2 innate lymphoid cells (ILC2) and promoting an eosinophilic

response at the site of infection (24, 25). Based on these data, a directed approach

was taken to evaluate expression of type 2 genes with the hypothesis that FMT

would promote their expression. While *IL25* expression was not detected in these

samples, expression of *IL33* and its receptor ST2 (*IL1RL1*) was upregulated in

response to FMT (Figure 3B). In addition, expression of the major eosinophil

attractant Eotaxin (*CCL11*) trended higher (FDR-adjusted p = 0.056) in FMT

treated mice, consistent with FMT promoting an immune response that facilitates

eosinophil recruitment. Other markers of type 2 immune responses including YM1

(*CHIL3*) (34, 35) and the IL-13 receptor (*IL13RA1*) were also upregulated in

response to FMT. The ability of FMT to repress IL-17A and related type 3 immune

responses was also of interest, as induction of a type 3 immune response can

promote more severe disease during CDI (26). While *IL17* was not detected,

expression of the IL-17 co-receptor (*IL17RC*) and the transcription factor RORγt

(*RORC*) were inhibited in response to FMT. *SOCS3*, a signaling regulator that

promotes type 1 or 2 immune responses while repressing type 3 differentiation

(36), was also upregulated in response to FMT. Overall, these findings support the

hypothesis that FMT administration may promote shifts towards a type 2-mediated

immune response and away from a type 3 response.

RNA Sequencing Implicates Inflammatory Biomarkers and Altered Lipid

Metabolism in the Immediate Response to FMT.

Next, an unbiased approach was taken to examine transcriptional differences

between vehicle control and FMT recipient samples. A total of 864 genes were

differentially expressed between vehicle control and FMT recipient samples after

false discovery (FDR) correction (Supplemental Table 1). Of these genes, 292 had

at least a 2-fold change ($\log_2$ fold change > 1.0) expression between groups and

50 had at least a 4-fold change (log2 fold change > 2.0). Of these 50 genes, 33

were upregulated in the FMT group relative to the vehicle control and 17 were

downregulated in the FMT condition. Unsupervised hierarchical clustering was

performed on these 50 genes to identify patterns within this gene set (Figure 3C).

Samples clustered according to group, indicating that all three groups had unique

expression patterns.

Many of the upregulated genes in the FMT recipient group encode proteins

associated with the host immune response. These include chemokines *CCL7*,

CCL12, and *IL1B* as well as immune signaling pathway proteins, such as the Toll

adapter protein *MAL* (37) and *PLET1*, which facilitates dendritic cell migration into

the intestine (38). Biomarkers of intestinal inflammation such as the Chitinase 1

(*CHIT1*) gene and inflammasome-dependent antimicrobial peptides *REG3B* and

REG3G (39) were also upregulated in the FMT recipient group compared to

vehicle controls. Immunoglobulin kappa variable genes were identified as both

upregulated and downregulated in response to FMT, but were more abundant in

the FMT group compared to controls. Extracellular matrix remodeling was also

enriched in response to FMT, with upregulated expression of both matrix

metalloproteinases (*MMP3*, *MMP10*, *PRSS27*) and proteinase inhibitors (*TIMP1*,

SERPINE1)*,* pathways that are expressed in response to immune signaling (40).

Downregulated genes made up a minority (17/50, 34%) of differentially expressed

genes and were more heterogeneous in their functions. Genes encoding immune-

associated proteins were also downregulated in response to FMT, including

S100G, Metallothionein 2 (*MT2*), and the tight junction protein Claudin 8 (*CLDN8*).

Several genes encoding bile acid and ion transporters were also downregulated

after FMT, including *SLC34A2*, *SLC10A2*, and *SCNN1G*. Finally, several genes

were involved in host metabolism, such as the Apoliprotein A-IV *APOA4*, the

pancreatic lipase *PLRP2*, and the peptide hormone *CCK*. Overall, genes with the

largest changes in gene expression were consistent with increased immune

activation in response to FMT.

GSEA-Assisted Pathway Analysis Identifies Immune Activation and

Decreased Lipid Processing in Response to FMT.

To better understand the biological processes broadly associated with FMT, Gene

Set Enrichment Analysis (GSEA) (41, 42) was performed utilizing the entire gene

expression data set. Overall, 23 gene sets had an FDR-adjusted p < 0.05 value for

significance (Figure 4), with 74% (17/23) being upregulated in FMT recipient

samples relative to vehicle controls. Upregulated gene sets were dominated by

immune pathways, including "HALLMARK_INFLAMMATORY_RESPONSE",

"HALLMARK_IL6_JAK_STAT3_SIGNALING", , and

"HALLMARK_INTERFERON_GAMMA_RESPONSE", suggesting an immediate

pro-inflammatory response to FMT administration. Many genes were associated

with multiple immune gene sets, with core genes such as *CCL2*, *SOCS3*, and

TIMP1. There was also significant overlap with genes previously identified in

Figure 3C, such as *SERPINE1, CCL7*, and *IL1B*. In addition to these pathways, the

gene set "HALLMARK_EPITHELIAL_MESENCHYMAL_TRANSITION" was

upregulated after FMT, which may suggest genes involved in resolution of

inflammation are being upregulated.

Downregulated gene sets primarily consisted of metabolism pathways, including

"HALLMARK_OXIDATIVE_PHOSPHORYLATION",

"HALLMARK_FATTY_ACID_METABOLISM", and

"HALLMARK_BILE_ACID_METABOLISM". The

"HALLMARK_OXIDATIVE_PHOSPHORYLATION" gene set was distinct from

other downregulated gene sets, with notable genes enriched in this data set

encoding proteins such as Monoamine Oxidase B (*MOAB*), Pyruvate

Dehydrogenase Kinase 4 (*PDK4*), ATPase Na+/K+ transporting subunit beta 1

(*ATP1B1*), and Cytochrome C Oxidase Copper Chaperone 11 (*COX11*). There

was significant overlap between genes in the remaining gene sets. Core genes

associated with these pathways encode proteins involved in fatty acid synthesis,

such as Acyl-CoA Synthetase (*ACSL1*), Malonyl-CoA Decarboxylase (*MLYCD*),

and Aldehyde Dehydrogenase 1 (*ALDH1A1*). Overall, these results suggest that

FMT is associated with broad changes in immune activation as well as

downregulation of metabolic pathways.

Discussion

While the effect of FMT on the intestinal microbiota has been well established, its

impact on host gene expression is more poorly understood. To address this gap in

knowledge, mice were treated with antibiotics before administration of either FMT

or vehicle control. Overall, the primary finding from this work demonstrates that

FMT results in significant changes to the host transcriptome relative to antibiotic

treatment alone, and that these genes are enriched in immune activation

pathways. Immune genes, including both pro-inflammatory pathways and several

genes associated type 2 immunity were upregulated in response to FMT treatment.

In contrast, several pathways were downregulated after FMT, including oxidative

phosphorylation, fatty acid metabolism, and bile acid metabolism. In addition, this

study confirms previous findings that FMT significantly increases microbial diversity

and partially restores community composition to that seen before antibiotic

treatment.

Microbiome diversity and community structure were both partially restored in

antibiotic treated mice within 48 hours of initial FMT administration. This finding is

consistent with previous studies that observed effective restoration of microbiome

diversity and structure after FMT treatment in both animal models (28, 30) and

clinical studies (27, 29, 43, 44). Restoration of *Firmicutes* was a specific hallmark

of FMT treatment, as 13 of the 15 ASVs that best distinguish FMT-treated mice

from vehicle controls were from this phylum. Notable *Firmicutes* restored by FMT

include *Lactabacillus ssp.*, which was the ASV that most strongly predicted FMT

treatment, as well as multiple taxa from the *Lachnospiraceae* family. These

changes suggest a microbial environment post-FMT that is more resistant to CDI,

as FMT success at preventing recurrent *C. difficile* infection is associated with

restoration of the *Firmicutes* and *Bacteroidota* phyla (45). In particular, several

Lactobacillus species have been implicated in protection against recurrent CDI

(46), potentially through production of organic acids that inhibit *C. difficile* growth or

inhibition of *C. difficile* toxin (47). Similarly, *Lachnospiraceae* species provide

colonization resistance against *C. difficile* in a mouse model, leading to decreased

intestinal histopathology and improved survival (48). This may be due in part to

production of short-chain fatty acids from carbohydrates, which inhibit *C. difficile*

growth, mediate epithelial proliferation, promote barrier integrity, and exert anti-

inflammatory effects in the gut (49, 50). In contrast, the three taxa enriched in the

control group relative to FMT-treated samples were *Akkermansia mucinophilia*

from the *Verrucomicrobiota* phylum, *Enterococcus ssp.* from the *Firmicutes*

phylum, and *Sutterellaceae ssp.* from the *Proteobacteria* phylum. Increased

relative abundance of the *Proteobacteria* phylum and *Enterococcus* family have

been observed immediately after antibiotic treatment (51), and these taxa are

reduced after FMT treatment (52). Disruption of the microbiota resulting in

increased relative abundance of *Proteobacteria* and *Enterococcus* also triggered

C. difficile overgrowth and increased spore shedding (53), suggesting that

increased relative abundance of these taxa may promote CDI. Taxa of the phylum

Verrucomicrobiota (including *Akkermansia*) are enriched after broad-spectrum

antibiotic treatment (54), indicating that higher relative abundance of this taxa in

the vehicle control group may be due to antibiotic treatment. Overall, these results

are largely consistent with other findings in the field and suggest that FMT

administration restores a microbial community more similar to before antibiotics

and more resistant to *C. difficile.*

A key finding of this work is the significant impact of FMT administration on host

gene expression. A total of 864 genes were differentially expressed between FMT

treated and vehicle control samples, with a significant proportion of these genes

involved in immune activation pathways. In particular, GSEA analysis identified an

enrichment in immune pathways that are primarily considered to be pro-

inflammatory, such as IL-6, IFNγ, and TNF☐. There is some evidence that

disruption of the intestinal microbiota is associated with disruption of type 1

immune responses and that these responses can be restored through

administration of FMT or microbial associated molecular patterns (MAMPs) (55,

56). Therefore, the introduction of MAMPs through FMT may drive the enrichment

in immune genes observed in this study. Activation of these immune pathways may

facilitate some of the protective effects of FMT independently of the microbiota, as

previous work identified filtered fecal supernatants as sufficient to protect against

recurrent CDI (57). Previous studies have found that FMT diminishes production of

pro-inflammatory cytokines such as IL-6 and TNF☐ (33) while promoting anti-

inflammatory mediators and regulatory T cells (30, 58). However, many of these

studies were in the context of intestinal inflammatory disorders and looked later in

the course of FMT than the current study, so it is unclear how comparable they are

to the acute phase of FMT in a naive setting. Strong inflammatory responses in the

initial response to infection are critical for *C. difficile* clearance and CDI recovery

(59–61), and it is possible that the upregulation of inflammatory genes immediately

after FMT promotes a stronger response upon subsequent *C. difficile*

infection. Additional studies in this context are necessary to elucidate the impact of

these immediate pro-inflammatory responses on CDI.

Signatures of type 2 immune responses and wound repair pathways were also

observed after FMT administration, indicating that the microbiota plays a key role in

maintaining a tissue regulatory and reparative response. Directed analysis of type

2 and type 3 associated genes in the list of differentially expressed genes identified

several type 2 genes (*IL33*, *IL1RL1*, *CHIL3*, *CCL11*) that were upregulated and

several type 3 genes (*RORC, IL17RC*) that were downregulated in response to

FMT. *SOCS3*, a critical signaling molecule that promotes type 2 and inhibits type 3

differentiation, was also identified in multiple gene sets enriched after FMT. These

results are consistent with previous results finding decreased expression of type 2

cytokines in germ-free mice and restoration of their expression after FMT treatment

(62, 63). Specific taxa identified in this study may be critical to these shifts: for

example, *REG3B* and *REG3G*, which were upregulated in response to FMT, were

restored after *Lactobacillus* supplementation in a DSS colitis model (64). Type 2

immune responses are associated with wound repair and eosinophil recruitment in

response to infection or inflammation (65), indicating that upregulation of these

genes may be the next step in restoration of intestinal homeostasis. Indeed, genes

involved in the epithelial to mesenchymal transition, a key process in wound repair

after inflammatory signaling (66), were also identified after FMT by GSEA. Several

genes in this set were also identified in the top 50 differentially regulated genes,

including the metalloprotease *MMP3* and protease inhibitors *TIMP1* and

SERPINE1, which are important for extracellular matrix remodeling (67, 68). In the

context of CDI, type 2 responses and subsequent eosinophilic responses are

critical to reduce disease severity (24, 25). Type 3 responses are slightly more

mixed: early pro-inflammatory responses are required for *C. difficile* clearance, but

prolonged inflammation and neutrophil recruitment is associated with worse

outcomes (26, 69, 70). Together, these gene expression data are consistent with

the paradigm that a type 2 immune response is generally protective against CDI

while a type 3 immune response exacerbates disease, which may partially explain

one of the mechanisms by which FMT protects against CDI.

Several pathways were downregulated in FMT-treated samples compared to

vehicle controls, including oxidative phosphorylation, fatty acid metabolism, and

bile acid metabolism. Downregulation of oxidative phosphorylation genes is

consistent with observations that immune cells are activated in this condition, as

immune activation is accompanied by a shift in immune cell metabolism from

oxidative phosphorylation to glycolysis (71). While immune activation is likely a key

driver of the downregulation of oxidative phosphorylation genes, this may also be

an effect of antibiotic treatment. Oxidative phosphorylation, fatty acid metabolism,

and bile acid metabolism genes were all downregulated in antibiotic treated mice

(44), highlighting the influence of the intestinal microbiota on these pathways. The

microbiota is known to significantly influence lipid absorption and metabolism (72),

indicating that disruption of the microbiota can impact lipid profiles relative to the

healthy gut. *SLC10A2*, which encodes a major primary bile acid importer, was also

downregulated after FMT treatment. Expression of *SLC10A2* is downregulated by

the intestinal microbiota (73), suggesting that restoration of the intestinal microbiota

may prevent bile acid reabsorption and increase the pool of microbially-modified

secondary bile acids, which are protective against CDI (74). Further research is

necessary to determine how these changes in gene expression affect bile acid and

lipid profiles in the gut and their downstream effects on disease susceptibility and

severity.

Together, these data show that FMT quickly restores microbial diversity and

community composition. FMT also significantly alters host gene expression,

particularly immune genes, demonstrating the dynamic and highly coupled

relationship between the host and the microbiota to maintain homeostasis. Many of

these pathways were associated with pro-inflammatory immune responses, which

may indicate an immediate response to the microbial community or microbial

components in the FMT. In addition, FMT upregulated several genes associated

with type 2 immune differentiation and repressed several type 3 biomarkers,

suggesting a shift towards repair and resolution of inflammation. In the context of

CDI, these shifts suggest an environment that is more resistant to subsequent *C.*

difficile infection. Future studies are required to identify how long-lasting these

transcriptional changes are and how they change the response to *C. difficile* during

infection. However, these results highlight the crosstalk between the intestinal

microbiota and host gene expression while identifying modulation of the immune

response as an important potential mechanism through which FMT protects

against intestinal dysbiosis and subsequent infection.

Acknowledgements

This work was funded by the US National Institutes of Health (1R01 AI152477 and

R01 AI124214 to WAP, 1F31AI136421-01A1 and 5T32AI007496-24 to ALF, and

F32DK124048 to JL). The funders played no role in experimental design or data

interpretation.

Methods

Murine FMT Model

All work with animals complied with all relevant ethical regulations for animal

testing and research procedures approved by Institutional Animal Care and Use

Committee at the University of Virginia (IACUC). Murine experiments were carried

out using eight-week-old male C57BL6/6J mice (Jackson Laboratory). Mice were

split into three groups: FMT donor (n=12), FMT recipient (n=10), and vehicle

control (n=10). Mice in vehicle control and FMT recipient groups were given an

antibiotic cocktail in drinking water consisting of 45mg/L Vancomycin (Mylan), 35

mg/L Colistin (Sigma), 35 mg/L Gentamicin (Sigma), and 215 mg/L Metronidazole

(Hospira) for three days. Mice were then switched to normal drinking water and

given a single IP injection (0.016 mg/g) of Clindamycin (Hospira) on Day -1.

On Day 0 and Day 1, FMT recipient mice were given an oral gavage of 100ul FMT,

while vehicle control mice were gavaged with anaerobic PBS as a sham control.

FMT was prepared from fecal pellets from age- and sex- matched donor mice that

were collected and immediately homogenized by vortexing in 1.5ml sterile

anaerobic PBS. Mice were sacrificed 48 hours after the initial FMT treatment on

Day 2. Cecal contents were collected for 16S rRNA sequencing and harvested

cecal tissue were immediately placed in RNA*later* (Sigma) and stored at -80°C.

16S rRNA Gene Sequencing

DNA extraction and 16S rRNA sequencing from murine cecal content was

performed as previously described (25, 28). Briefly, DNA was extracted using the

MagAttract PowerMicrobiome kit (Qiagen) with the EpMotion liquid handling

system (Eppendorf). The V4 region of the 16S rRNA gene was amplified from each

sample using a dual indexing sequencing strategy (75). Samples were sequenced

on the MiSeq platform (Illumina) using the MiSeq Reagent Kit V2 500 cycles

(Illumina #MS102-2003) according to the manufacturer's protocol with

modifications found in the Schloss Wet Lab SOP

(https://github.com/SchlossLab/MiSeq_WetLab_SOP).

16S Sequence Analysis

All analyses were performed in R (version 4.0.2) (76). Raw sequencing reads were

inspected for quality, filtered, paired reads merged, and chimeric sequences

removed using DADA2 (version 1.18.0) (77). Length distribution of non-chimeric

sequences was plotted to ensure lengths matched the expected V4 amplicon size.

Taxonomy was assigned to amplicon sequence variants (ASVs) by aligning with

the Silva reference database (version 138.1) (78). Alpha and beta diversity were

calculated using the phyloseq package (version 1.34.0) (79). The packages

tidyverse (version 1.3.0) (80) and ggpubr (version 0.4.0) were used for data

organization and analysis.

Sample preparation and sequencing quality was assessed using blank control,

water control, and mock community control samples. Blank and water control

samples had significantly fewer reads than other samples, suggesting that

contamination was not a major problem during the sequencing run. A mock

community control sample was also processed to investigate sequencing accuracy

and identified all eight community members included in the mock community

sample at similar relative abundance (Supplemental Table 2). Three additional

ASVs were identified, including a *Bacteroidota* ASV that comprised 0.17% of

sequence abundance, a LD29 ASV that comprised 0.05% of sequence abundance,

and an ASV of unknown genus that comprised 2.1% of sequence abundance. This

unknown ASV is of the *Enterobacteriaceae* Family, which is the same family as

Salmonella ssp., a predicted member of the community. We hypothesize that this

ASV of unknown genus may represent *Salmonella* that was artificially split during

the DADA2 pipeline (81).

Random forest analysis was performed in R using the randomForest package

(version 4.6.14) (82). Samples were first divided into training (70% of samples,

divided equally between FMT recipient and vehicle control samples) and test (30%

of samples) sets. The training set was used for tuning the "mtry" parameter of the

model, while the test set was used to validate model performance. Feature

importance was determined using the Gini index, which measures the total

decrease in node impurity averaged across all trees.

RNA Isolation from Mouse Cecal Tissue

RNA*later*-stabilized cecal tissue was removed using sterile needles and placed in

1ml TRIzol per tissue. Tissues were disrupted and homogenized using TissueLyser

II according to the manufacturer's protocol and then tubes were centrifuged to

separate the aqueous and organic phases. The aqueous phase was precipitated in

ethanol, and RNA was redissolved in 400ul RLT buffer and further purified using

the RNeasy kit (Qiagen) according the manufacturer's protocol. The size,

distribution, and quantity of total RNA in each sample was determined using an

Agilent 2100 bioanalyzer. All samples had an RNA integrity number (RIN) above 8.

RNA Sequencing

RNA Sequencing was performed by Novogene. Briefly, mRNA was isolated using

oligo(dT) beads and randomly fragmented, followed by cDNA synthesis and

second-strand synthesis. After end repair, barcode ligation, and sequencing

adaptor ligation, the double-stranded cDNA library was size selected and PCR

amplified. Sequencing was carried out on an Illumina HiSeq platform PE150 with

paired-end 150 base pair reads.

Gene Expression Profiling

RNA sequencing libraries were checked for their quality using the FastQC program

(https://www.bioinformatics.babraham.ac.uk/projects/fastqc/) and aggregated using

MultiQC (83). Clean reads were mapped to the most updated version of the

transcriptome with the STAR aligner, and counted using HTseq (84). Count tables

were imported into R for differential gene expression analysis. The DESeq2

package (85) was used to exclude genes with low counts, normalize data, estimate

dispersion, and fit counts using a negative binomial model. A heatmap of

differentially expressed genes between FMT recipient and vehicle control groups

was generated from normalized count data in R using the pheatmap package

(version 1.0.12). Differentially expressed genes were also ranked (upregulated to

downregulated) based on $\log_2$ fold change and FDR corrected p-values for

pathway analysis using Gene Set Enrichment Analysis (GSEA) software (41, 42).

Hallmark gene sets from MSigDB were chosen for this analysis.

Data Sharing

Raw sequencing reads have been deposited in the Sequence Read Archive (SRA)

database under Accession number PRJNA521980. Data and code for data

processing are hosted at https://github.com/gbmoreau/FMT_Mouse_16S_RNAseq.

References

1.   2019. Antibiotic resistance threats in the United States, 2019.

http://dx.doi.org/10.15620/cdc:82532, Atlanta, GA.

2.   Bacci S, Mølbak K, Kjeldsen MK, Olsen KEP. 2011. Binary toxin and death

after Clostridium difficile infection. Emerg Infect Dis 17:976–982.

3.   Loo VG, Poirier L, Miller MA, Oughton M, Libman MD, Michaud S, Bourgault

A-M, Nguyen T, Frenette C, Kelly M, Vibien A, Brassard P, Fenn S, Dewar K,

Hudson TJ, Horn R, René P, Monczak Y, Dascal A. 2005. A Predominantly

Clonal Multi-Institutional Outbreak of Clostridium difficile–Associated

Diarrhea with High Morbidity and Mortality. N Engl J Med 353:2442–2449.

4.   Pépin J, Valiquette L, Alary M-E, Villemure P, Pelletier A, Forget K, Pépin K,

Chouinard D. 2004. Clostridium difficile-associated diarrhea in a region of

Quebec from 1991 to 2003: a changing pattern of disease severity. CMAJ

171:466–472.

5.   Kuijper EJ, Barbut F, Brazier JS, Kleinkauf N, Eckmanns T, Lambert ML,

Drudy D, Fitzpatrick F, Wiuff C, Brown DJ, Coia JE, Pituch H, Reichert P,

Even J, Mossong J, Widmer AF, Olsen KE, Allerberger F, Notermans DW,

Delmée M, Coignard B, Wilcox M, Patel B, Frei R, Nagy E, Bouza E, Marin

M, Akerlund T, Virolainen-Julkunen A, Lyytikäinen O, Kotila S, Ingebretsen

A, Smyth B, Rooney P, Poxton IR, Monnet DL. 2008. Update of Clostridium

difficile infection due to PCR ribotype 027 in Europe, 2008. Euro Surveill Bull

Eur sur les Mal Transm = Eur Commun Dis Bull 13.

6.   Lessa FC, Mu Y, Bamberg WM, Beldavs ZG, Dumyati GK, Dunn JR, Farley

MM, Holzbauer SM, Meek JI, Phipps EC, Wilson LE, Winston LG, Cohen JA,

Limbago BM, Fridkin SK, Gerding DN, McDonald LC. 2015. Burden of

Clostridium difficile infection in the United States. N Engl J Med 372:825–

834.

7. Czepiel J, Dróżdż M, Pituch H, Kuijper EJ, Perucki W, Mielimonka A,

Goldman S, Wultańska D, Garlicki A, Biesiada G. 2019. Clostridium difficile

infection: review. Eur J Clin Microbiol Infect Dis 38:1211–1221.

8. Khanna S, Pardi DS, Aronson SL, Kammer PP, Orenstein R, St Sauver JL,

Harmsen WS, Zinsmeister AR. 2012. The epidemiology of community-

acquired Clostridium difficile infection: a population-based study. Am J

Gastroenterol2011/11/22. 107:89–95.

9. Slimings C, Riley T V. 2014. Antibiotics and hospital-acquired Clostridium

difficile infection: update of systematic review and meta-analysis. J

Antimicrob Chemother 69:881–891.

10. Deshpande A, Pasupuleti V, Thota P, Pant C, Rolston DDK, Sferra TJ,

Hernandez A V, Donskey CJ. 2013. Community-associated Clostridium

difficile infection and antibiotics: a meta-analysis. J Antimicrob Chemother

68:1951–1961.

11. Cornely OA, Miller MA, Louie TJ, Crook DW, Gorbach SL. 2012. Treatment

of First Recurrence of Clostridium difficile Infection: Fidaxomicin Versus

Vancomycin. Clin Infect Dis 55:S154–S161.

12. Louie TJ, Miller MA, Mullane KM, Weiss K, Lentnek A, Golan Y, Gorbach S,

Sears P, Shue Y-K. 2011. Fidaxomicin versus vancomycin for Clostridium

difficile infection. N Engl J Med 364:422–431.

13. Cohen NA, Maharshak N. 2017. Novel Indications for Fecal Microbial

Transplantation: Update and Review of the Literature. Dig Dis Sci 62:1131–

1145.

14. Baunwall SMD, Lee MM, Eriksen MK, Mullish BH, Marchesi JR, Dahlerup JF,

Hvas CL. 2020. Faecal microbiota transplantation for recurrent Clostridioides

difficile infection: An updated systematic review and meta-analysis.

EClinicalMedicine 29–30:100642.

15. Gupta S, Mullish BH, Allegretti JR. 2021. Fecal Microbiota Transplantation:

The Evolving Risk Landscape. Off J Am Coll Gastroenterol | ACG 116.

16. DeFilipp Z, Bloom PP, Torres Soto M, Mansour MK, Sater MRA, Huntley

MH, Turbett S, Chung RT, Chen Y-B, Hohmann EL. 2019. Drug-Resistant E.

coli Bacteremia Transmitted by Fecal Microbiota Transplant. N Engl J Med

381:2043–2050.

17. Adamu BO, Lawley TD. 2013. Bacteriotherapy for the treatment of intestinal

dysbiosis caused by Clostridium difficile infection. Curr Opin

Microbiol2013/07/15. 16:596–601.

18. Chang JY, Antonopoulos DA, Kalra A, Tonelli A, Khalife WT, Schmidt TM,

Young VB. 2008. Decreased Diversity of the Fecal Microbiome in Recurrent

Clostridium difficile—Associated Diarrhea. J Infect Dis 197:435–438.

19. Weingarden AR, Chen C, Bobr A, Yao D, Lu Y, Nelson VM, Sadowsky MJ,

Khoruts A. 2014. Microbiota transplantation restores normal fecal bile acid

composition in recurrent Clostridium difficile infection. Am J Physiol

Gastrointest Liver Physiol2013/11/27. 306:G310–G319.

20. Frisbee AL, Petri WA. 2020. Considering the Immune System during Fecal

Microbiota Transplantation for Clostridioides difficile Infection. Trends Mol

Med 26:496–507.

21. Steiner TS, Flores CA, Pizarro TT, Guerrant RL. 1997. Fecal lactoferrin,

interleukin-1beta, and interleukin-8 are elevated in patients with severe

Clostridium difficile colitis. Clin Diagn Lab Immunol 4:719–722.

22. Jiang Z-D, DuPont HL, Garey K, Price M, Graham G, Okhuysen P, Dao-Tran

T, LaRocco M. 2006. A common polymorphism in the interleukin 8 gene

promoter is associated with Clostridium difficile diarrhea. Am J Gastroenterol

101:1112–1116.

23. El Feghaly RE, Stauber JL, Deych E, Gonzalez C, Tarr PI, Haslam DB.

2013. Markers of intestinal inflammation, not bacterial burden, correlate with

clinical outcomes in Clostridium difficile infection. Clin Infect Dis2013/03/13.

56:1713–1721.

24. Buonomo EL, Cowardin CA, Wilson MG, Saleh MM, Pramoonjago P, Petri

WAJ. 2016. Microbiota-Regulated IL-25 Increases Eosinophil Number to

Provide Protection during Clostridium difficile Infection. Cell Rep 16:432–

443.

25. Frisbee AL, Saleh MM, Young MK, Leslie JL, Simpson ME, Abhyankar MM,

Cowardin CA, Ma JZ, Pramoonjago P, Turner SD, Liou AP, Buonomo EL,

Petri WA. 2019. IL-33 drives group 2 innate lymphoid cell-mediated

protection during Clostridium difficile infection. Nat Commun 10:2712.

26.   Saleh MM, Frisbee AL, Leslie JL, Buonomo EL, Cowardin CA, Ma JZ,

Simpson ME, Scully KW, Abhyankar MM, Petri WA. 2019. Colitis-Induced

Th17 Cells Increase the Risk for Severe Subsequent Clostridium difficile

Infection. Cell Host Microbe 25:756-765.e5.

27.   Hamilton MJ, Weingarden AR, Unno T, Khoruts A, Sadowsky MJ. 2013.

High-throughput DNA sequence analysis reveals stable engraftment of gut

microbiota following transplantation of previously frozen fecal bacteria. Gut

Microbes 4:125–135.

28.   Seekatz AM, Theriot CM, Molloy CT, Wozniak KL, Bergin IL, Young VB.

2015. Fecal Microbiota Transplantation Eliminates Clostridium difficile in a

Murine Model of Relapsing Disease. Infect Immun2015/07/13. 83:3838–

3846.

29.   Jacob V, Crawford C, Cohen-Mekelburg S, Viladomiu M, Putzel GG,

Schneider Y, Chabouni F, O'Neil S, Bosworth B, Woo V, Ajami NJ, Petrosino

JF, Gerardin Y, Kassam Z, Smith M, Iliev ID, Sonnenberg GF, Artis D, Scherl

E, Longman RS. 2017. Single Delivery of High-Diversity Fecal Microbiota

Preparation by Colonoscopy Is Safe and Effective in Increasing Microbial

Diversity in Active Ulcerative Colitis. Inflamm Bowel Dis 23:903–911.

30.   Wang Z, Hua W, Li C, Chang H, Liu R, Ni Y, Sun H, Li Y, Wang X, Hou M,

Liu Y, Xu Z, Ji M. 2019. Protective Role of Fecal Microbiota Transplantation

on Colitis and Colitis-Associated Colon Cancer in Mice Is Associated With

Treg Cells. Front Microbiol 10:2498.

31. Burrello C, Garavaglia F, Cribiù FM, Ercoli G, Lopez G, Troisi J, Colucci A,

Guglietta S, Carloni S, Guglielmetti S, Taverniti V, Nizzoli G, Bosari S,

Caprioli F, Rescigno M, Facciotti F. 2018. Therapeutic faecal microbiota

transplantation controls intestinal inflammation through IL10 secretion by

immune cells. Nat Commun 9:5184.

32. Wei Y-L, Chen Y-Q, Gong H, Li N, Wu K-Q, Hu W, Wang B, Liu K-J, Wen L-

Z, Xiao X, Chen D-F. 2018. Fecal Microbiota Transplantation Ameliorates

Experimentally Induced Colitis in Mice by Upregulating AhR. Front Microbiol

9:1921.

33. Konturek PC, Koziel J, Dieterich W, Haziri D, Wirtz S, Glowczyk I, Konturek

K, Neurath MF, Zopf Y. 2016. Successful therapy of Clostridium difficile

infection with fecal microbiota transplantation. J Physiol Pharmacol an Off J

Polish Physiol Soc 67:859–866.

34. Welch JS, Escoubet-Lozach L, Sykes DB, Liddiard K, Greaves DR, Glass

CK. 2002. TH2 cytokines and allergic challenge induce Ym1 expression in

macrophages by a STAT6-dependent mechanism. J Biol Chem 277:42821–

42829.

35. Cai Y, Kumar RK, Zhou J, Foster PS, Webb DC. 2009. Ym1/2 promotes Th2

cytokine expression by inhibiting 12/15(S)-lipoxygenase: identification of a

701 novel pathway for regulating allergic inflammation. J Immunol 182:5393–

702 5399.

703 36. Chen Z, Laurence A, Kanno Y, Pacher-Zavisin M, Zhu B-M, Tato C,

704 Yoshimura A, Hennighausen L, O'Shea JJ. 2006. Selective regulatory

705 function of Socs3 in the formation of IL-17-secreting T cells. Proc Natl Acad

706 Sci 103:8137 LP – 8142.

707 37. Belhaouane I, Hoffmann E, Chamaillard M, Brodin P, Machelart A. 2020.

708 Paradoxical Roles of the MAL/Tirap Adaptor in Pathologies. Front Immunol

709 11:569127.

710 38. Karrich JJ, Romera-Hernández M, Papazian N, Veenbergen S, Cornelissen

711 F, Aparicio-Domingo P, Stenhouse FH, Peddie CD, Hoogenboezem RM, den

712 Hollander CWJ, Gaskell T, Medley T, Boon L, Blackburn CC, Withers DR,

713 Samsom JN, Cupedo T. 2019. Expression of Plet1 controls interstitial

714 migration of murine small intestinal dendritic cells. Eur J Immunol 49:290–

715 301.

716 39. Ratsimandresy RA, Indramohan M, Dorfleutner A, Stehlik C. 2017. The AIM2

717 inflammasome is a central regulator of intestinal homeostasis through the IL-

718 18/IL-22/STAT3 pathway. Cell Mol Immunol 14:127–142.

719 40. Pham TAN, Clare S, Goulding D, Arasteh JM, Stares MD, Browne HP,

720 Keane JA, Page AJ, Kumasaka N, Kane L, Mottram L, Harcourt K, Hale C,

721 Arends MJ, Gaffney DJ, Dougan G, Lawley TD. 2014. Epithelial IL-22RA1-

722 Mediated Fucosylation Promotes Intestinal Colonization Resistance to an

Opportunistic Pathogen. Cell Host Microbe 16:504–516.

41. Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA,

Paulovich A, Pomeroy SL, Golub TR, Lander ES, Mesirov JP. 2005. Gene

set enrichment analysis: a knowledge-based approach for interpreting

genome-wide expression profiles. Proc Natl Acad Sci U S A 102:15545–

15550.

42. Mootha VK, Lindgren CM, Eriksson K-F, Subramanian A, Sihag S, Lehar J,

Puigserver P, Carlsson E, Ridderstråle M, Laurila E, Houstis N, Daly MJ,

Patterson N, Mesirov JP, Golub TR, Tamayo P, Spiegelman B, Lander ES,

Hirschhorn JN, Altshuler D, Groop LC. 2003. PGC-1α-responsive genes

involved in oxidative phosphorylation are coordinately downregulated in

human diabetes. Nat Genet 34:267–273.

43. Nusbaum DJ, Sun F, Ren J, Zhu Z, Ramsy N, Pervolarakis N, Kunde S,

England W, Gao B, Fiehn O, Michail S, Whiteson K. 2018. Gut microbial and

metabolomic profiles after fecal microbiota transplantation in pediatric

ulcerative colitis patients. FEMS Microbiol Ecol 94.

44. Suez J, Zmora N, Zilberman-Schapira G, Mor U, Dori-Bachash M,

Bashiardes S, Zur M, Regev-Lehavi D, Ben-Zeev Brik R, Federici S, Horn M,

Cohen Y, Moor AE, Zeevi D, Korem T, Kotler E, Harmelin A, Itzkovitz S,

Maharshak N, Shibolet O, Pevsner-Fischer M, Shapiro H, Sharon I, Halpern

Z, Segal E, Elinav E. 2018. Post-Antibiotic Gut Mucosal Microbiome

Reconstitution Is Impaired by Probiotics and Improved by Autologous FMT.

Cell 174:1406-1423.e16.

45.   Gupta S, Allen-Vercoe E, Petrof EO. 2016. Fecal microbiota transplantation:

in perspective. Therap Adv Gastroenterol 9:229–239.

46.   Maziade P-J, Pereira P, Goldstein EJC. 2015. A Decade of Experience in

Primary Prevention of Clostridium difficile Infection at a Community Hospital

Using the Probiotic Combination Lactobacillus acidophilus CL1285,

Lactobacillus casei LBC80R, and Lactobacillus rhamnosus CLR2 (Bio-K+).

Clin Infect Dis 60:S144–S147.

47.   Auclair J, Frappier M, Millette M. 2015. Lactobacillus acidophilus CL1285,

Lactobacillus casei LBC80R, and Lactobacillus rhamnosus CLR2 (Bio-K+):

Characterization, Manufacture, Mechanisms of Action, and Quality Control of

a Specific Probiotic Combination for Primary Prevention of Clostridium dif.

Clin Infect Dis 60:S135–S143.

48.   Reeves AE, Koenigsknecht MJ, Bergin IL, Young VB. 2012. Suppression of

Clostridium difficile in the gastrointestinal tracts of germfree mice inoculated

with a murine isolate from the family Lachnospiraceae. Infect

Immun2012/08/13. 80:3786–3794.

49.   Rolfe RD. 1984. Role of volatile fatty acids in colonization resistance to

Clostridium difficile. Infect Immun 45:185–191.

50.   Parada Venegas D, De la Fuente MK, Landskron G, González MJ, Quera R,

Dijkstra G, Harmsen HJM, Faber KN, Hermoso MA. 2019. Short Chain Fatty

Acids (SCFAs)-Mediated Gut Epithelial and Immune Regulation and Its

Relevance for Inflammatory Bowel Diseases . Front Immunol .

51. Schwartz DJ, Langdon AE, Dantas G. 2020. Understanding the impact of

antibiotic perturbation on the human microbiome. Genome Med 12:82.

52. Kellingray L, Gall G Le, Defernez M, Beales ILP, Franslem-Elumogo N,

Narbad A. 2018. Microbial taxonomic and metabolic alterations during faecal

microbiota transplantation to treat Clostridium difficile infection. J Infect

77:107–118.

53. Lawley TD, Clare S, Walker AW, Goulding D, Stabler RA, Croucher N,

Mastroeni P, Scott P, Raisen C, Mottram L, Fairweather NF, Wren BW,

Parkhill J, Dougan G. 2009. Antibiotic treatment of clostridium difficile carrier

mice triggers a supershedder state, spore-mediated transmission, and

severe disease in immunocompromised hosts. Infect Immun 77:3661–3669.

54. Dubourg G, Lagier J-C, Armougom F, Robert C, Audoly G, Papazian L,

Raoult D. 2013. High-level colonisation of the human gut by Verrucomicrobia

following broad-spectrum antibiotic treatment. Int J Antimicrob Agents

41:149–155.

55. Khan N, Vidyarthi A, Nadeem S, Negi S, Nair G, Agrewala JN. 2016.

Alteration in the Gut Microbiota Provokes Susceptibility to Tuberculosis.

Front Immunol 7:529.

56. Xu C, Ruan B, Jiang Y, Xue T, Wang Z, Lu H, Wei M, Wang S, Ye Z, Zhai D,

Wang L, Lu Z. 2017. Antibiotics-induced gut microbiota dysbiosis promotes

tumor initiation via affecting APC-Th1 development in mice. Biochem

Biophys Res Commun 488:418–424.

57. Ott SJ, Waetzig GH, Rehman A, Moltzau-Anderson J, Bharti R, Grasis JA,

Cassidy L, Tholey A, Fickenscher H, Seegert D, Rosenstiel P, Schreiber S.

2017. Efficacy of Sterile Fecal Filtrate Transfer for Treating Patients With

Clostridium difficile Infection. Gastroenterology 152:799-811.e7.

58. Liu Y, Fan L, Cheng Z, Yu L, Cong S, Hu Y, Zhu L, Zhang B, Cheng Y, Zhao

P, Zhao X, Cheng M. 2021. Fecal transplantation alleviates acute liver injury

in mice through regulating Treg/Th17 cytokines balance. Sci Rep 11:1611.

59. Ishida Y, Maegawa T, Kondo T, Kimura A, Iwakura Y, Nakamura S, Mukaida

N. 2004. Essential involvement of IFN-gamma in Clostridium difficile toxin A-

induced enteritis. J Immunol 172:3018–3025.

60. Jafari N V, Kuehne SA, Bryant CE, Elawad M, Wren BW, Minton NP, Allan E,

Bajaj-Elliott M. 2013. Clostridium difficile modulates host innate immunity via

toxin-independent and dependent mechanism(s). PLoS One 8:e69846.

61. Abt MC, Lewis BB, Caballero S, Xiong H, Carter RA, Sušac B, Ling L, Leiner

I, Pamer EG. 2015. Innate Immune Defenses Mediated by Two ILC Subsets

Are Critical for Protection against Acute Clostridium difficile Infection. Cell

Host Microbe 18:27–37.

62. De Salvo C, Wang X-M, Pastorelli L, Mattioli B, Omenetti S, Buela KA,

Chowdhry S, Garg RR, Goodman WA, Rodriguez-Palacios A, Smith DE,

Abbott DW, Cominelli F, Bamias G, Xin W, Lee JJ, Vecchi M, Pizarro TT.

2016. IL-33 Drives Eosinophil Infiltration and Pathogenic Type 2 Helper T-

Cell Immune Responses Leading to Chronic Experimental Ileitis. Am J

Pathol 186:885–898.

63. Zhang L, Ma X, Liu P, Ge W, Hu L, Zuo Z, Xiao H, Liao W. 2021. Treatment

and mechanism of fecal microbiota transplantation in mice with

experimentally induced ulcerative colitis. Exp Biol Med 246:1563–1575.

64. Hou Q, Ye L, Liu H, Huang L, Yang Q, Turner JR, Yu Q. 2018. Lactobacillus

accelerates ISCs regeneration to protect the integrity of intestinal mucosa

through activation of STAT3 signaling pathway induced by LPLs secretion of

IL-22. Cell Death Differ 25:1657–1670.

65. Allen JE, Wynn TA. 2011. Evolution of Th2 immunity: a rapid repair response

to tissue destructive pathogens. PLoS Pathog2011/05/12. 7:e1002003–

e1002003.

66. Kalluri R, Weinberg RA. 2009. The basics of epithelial-mesenchymal

transition. J Clin Invest 119:1420–1428.

67. Manicone AM, McGuire JK. 2008. Matrix metalloproteinases as modulators

of inflammation. Semin Cell Dev Biol 19:34–41.

68. Fingleton B. 2017. Matrix metalloproteinases as regulators of inflammatory

processes. Biochim Biophys Acta - Mol Cell Res 1864:2036–2042.

69. Solomon K, Martin AJ, O'Donoghue C, Chen X, Fenelon L, Fanning S, Kelly

CP, Kyne L. 2013. Mortality in patients with Clostridium difficile infection

correlates with host pro-inflammatory and humoral immune responses. J

Med Microbiol 62:1453–1460.

70.   Jose S, Madan R. 2016. Neutrophil-mediated inflammation in the

834   pathogenesis of Clostridium difficile infections. Anaerobe 41:85–90.

835   71.   Ganeshan K, Chawla A. 2014. Metabolic Regulation of Immune Responses.

836   CircAnn Rev Immunol.

837   72.   Aron-Wisnewsky J, Warmbrunn M V, Nieuwdorp M, Clément K. 2021.

838   Metabolism and Metabolic Disorders and the Microbiome: The Intestinal

839   Microbiota Associated With Obesity, Lipid Metabolism, and Metabolic

840   Health—Pathophysiology and Therapeutic Strategies. Gastroenterology

841   160:573–599.

842   73.   Yang N, Dong Y-Q, Jia G-X, Fan S-M, Li S-Z, Yang S-S, Li Y-B. 2020.

843   ASBT(SLC10A2): A promising target for treatment of diseases and drug

844   discovery. Biomed Pharmacother 132:110835.

845   74.   Buffie CG, Bucci V, Stein RR, McKenney PT, Ling L, Gobourne A, No D, Liu

846   H, Kinnebrew M, Viale A, Littmann E, van den Brink MRM, Jenq RR, Taur Y,

847   Sander C, Cross JR, Toussaint NC, Xavier JB, Pamer EG. 2015. Precision

848   microbiome reconstitution restores bile acid mediated resistance to

849   Clostridium difficile. Nature2014/10/22. 517:205–208.

850   75.   Kozich JJ, Westcott SL, Baxter NT, Highlander SK, Schloss PD. 2013.

851   Development of a dual-index sequencing strategy and curation pipeline for

852   analyzing amplicon sequence data on the MiSeq Illumina sequencing

853   platform. Appl Environ Microbiol2013/06/21. 79:5112–5120.

76.   R Core Team. 2020. R: A language and environment for statistical

computing.

77. Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP.

2016. DADA2: High-resolution sample inference from Illumina amplicon data.

Nat Methods2016/05/23. 13:581–583.

78. Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, Peplies J,

Glöckner FO. 2013. The SILVA ribosomal RNA gene database project:

improved data processing and web-based tools. Nucleic Acids

Res2012/11/28. 41:D590–D596.

79. McMurdie PJ, Holmes S. 2013. phyloseq: An R Package for Reproducible

Interactive Analysis and Graphics of Microbiome Census Data. PLoS One

8:e61217.

80. Wickham H, Averick M, Bryan J, Chang W, McGowan LD, François R,

Grolemund G, Hayes A, Henry L, Hester J. 2019. Welcome to the Tidyverse.

J open source Softw 4:1686.

81. D. SP, Katherine M. 2021. Amplicon Sequence Variants Artificially Split

Bacterial Genomes into Separate Clusters. mSphere 0:e00191-21.

82. Liaw A, Wiener M. 2002. Classification and regression by randomForest. R

news 2:18–22.

83. Ewels P, Magnusson M, Lundin S, Käller M. 2016. MultiQC: summarize

analysis results for multiple tools and samples in a single report.

Bioinformatics 32:3047–3048.

84. Anders S, Pyl PT, Huber W. 2015. HTSeq--a Python framework to work with

high-throughput sequencing data. Bioinformatics 31:166–169.

85.   Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change

and dispersion for RNA-seq data with DESeq2. Genome Biol 15:550.

Figure Legends

Figure 1: *Overview of experimental design.* C57BL6 mice were divided into three

groups: FMT donors (n=12), FMT recipients (n=10), and vehicle controls (n=10).

FMT recipient and vehicle control mice were given an antibiotic cocktail in drinking

water for three days, then switched to regular drinking water and given a single IP

injection of Clindamycin on Day −1. On Day 0 and Day 1, mice were administered

either FMT (a fecal slurry from donor mice) or vehicle (PBS) by oral gavage. On

Day 2, cecal contents and cecal tissue were harvested from mice for downstream

analysis. FMT donor mice were untreated until harvest on Day 2.

Figure 2: *FMT partially restores the microbiome after antibiotic disruption.* A) Alpha

diversity metrics (Total number of observed ASVs and Shannon Index) for FMT

donor (red), FMT recipient (green) and vehicle control (blue) groups. All groups

within each metric are significantly different from each other when analyzed using a

pairwise Wilcoxon Rank test with Bonferroni correction for multiple comparisons. B)

Non-metric multidimensional scaling (NMDS) plot of beta diversity (unweighted

Unifrac) metrics for FMT donor (red), FMT recipient (green) and vehicle control

(blue) groups. The percent of variance explained by each dimension is listed on

each axis. C-D) Relative abundance of taxa within each group at the C) phylum

and D) family levels. Taxa that comprised less than 1% of relative abundance in all

groups are aggregated. E) Ranking of the Top 15 ASVs that distinguish FMT

recipient samples from vehicle controls as determined by a random forest model.

ASVs that are more abundant in the FMT recipient group relative to vehicle

controls are displayed as circles, while ASVs more abundant in the control group

are displayed as triangles.

Figure 3: *FMT significantly changes host gene expression.* A) Principal component

analysis of RNA sequencing reads from FMT Donor (red), FMT recipient (green),

and vehicle Control (blue) samples. B) Transcriptional changes in FMT recipient

group relative the vehicle control group. Select type 2 immunity-associated genes

are colored in blue, while select type 3 immunity-associated genes are colored in

red. *SOCS3* is both a promoter of type 2 responses and an inhibitor of type 17

responses. C) Heatmap of normalized and log2-transformed gene counts for genes

with at least a four-fold change in expression and FDR-adjusted p-value < 0.05

between FMT recipient and vehicle control samples. Samples and genes are both

hierarchically clustered.

Figure 4: *FMT increases expression of host immune pathways.* Ranking of

significantly enriched (FDR adjusted p-value < 0.05) gene sets in either FMT

recipient (red) or vehicle control (blue) samples. Gene sets are ranked according to

Normalized Enrichment Score, which normalizes gene enrichment based on the

size of each gene set.

Enriched Gene Sets (FMT Recipient vs. Vehicle Control)

HALLMARK_INFLAMMATORY_RESPONSE
HALLMARK_TNFA_SIGNALING_VIA_NFKB
HALLMARK_IL6_JAK_STAT3_SIGNALING
HALLMARK_INTERFERON_GAMMA_RESPONSE
HALLMARK_TGFB_BETA_SIGNALING
HALLMARK_TH2_SIGNALING
HALLMARK_COAGULATION
HALLMARK_KRAS_SIGNALING_UP
HALLMARK_ANGIOGENESIS
HALLMARK_INTERFERON_ALPHA_RESPONSE
HALLMARK_APICAL_JUNCTION
HALLMARK_IL2_STAT5_SIGNALING
HALLMARK_APOPTOSIS
HALLMARK_APICAL_SURFACE
HALLMARK_UV_RESPONSE_UP
HALLMARK_MYC_TARGETS_V1
HALLMARK_CHOLESTEROL_HOMEOSTASIS
HALLMARK_BILE_ACID_METABOLISM
HALLMARK_PEROXISOME
HALLMARK_OXIDATIVE_PHOSPHORYLATION

Normalized Enrichment Score

Enrichment
■ FMT Recipient
■ Vehicle Control