1   **Running title:** Leaf senescence and climate warming

2   **Shifts in leaf senescence across the Northern Hemisphere in response to seasonal**

3   **warming**

4   Lei Chen[1,2*], Sergio Rossi[3,4], Nicholas G. Smith[2], Jianquan Liu[1*]

[1]Key Laboratory of Bio-Resource and Eco-Environment of Ministry of Education,

College of Life Sciences, Sichuan University, Chengdu, China

[2]Department of Biological Sciences, Texas Tech University, Lubbock, USA

[3]Département des Sciences Fondamentales, Université du Québec à Chicoutimi,

Chicoutimi (QC), G7H SB1, Canada

[4]Key Laboratory of Vegetation Restoration and Management of Degraded Ecosystems,

Guangdong Provincial Key Laboratory of Applied Botany, South China Botanical

Garden, Chinese Academy of Sciences, Guangzhou, China

* Email of corresponding author:

Lei Chen   Email: chen_lei1029@163.com

Jianquan Liu   Email: liujq@lzu.edu.cn

Abstract

Shifts in plant phenology under ongoing warming affect global vegetation

dynamics and carbon assimilation of the biomes. The response of leaf senescence to

climate is crucial for predicting changes in the physiological processes of trees at

ecosystem scale. We used long-term ground observations, phenological metrics derived

from PhenoCam, and satellite imagery of the Northern Hemisphere to show that the

timings of leaf senescence can advance or delay in case of warming occurring at the

beginning (before June) or during (after June) the main growing season, respectively.

Flux data demonstrated that net photosynthetic carbon assimilation converted from

positive to negative at the end of June. These findings suggest that leaf senescence is

driven by carbon assimilation and nutrient resorption at different growth stages of

leaves. Our results provide new insights into understanding and modelling autumn

phenology and carbon cycling under warming scenarios.

Key words: Climate change, phenology, carbon assimilation, nutrient resorption,

phenocam, flux data

INTRODUCTION

Tree phenology mirrors the timing of budburst, leaf-out, flowering, leaf senescence and other related biological events (Richardson *et al.* 2013; Piao *et al.* 2019). Shifts in tree phenology alter the length of growing season (Cleland *et al.* 2007; Richardson *et al.* 2018) and influence the productivity of terrestrial forest ecosystems (Zhang *et al.* 2020). Phenological changes in trees provide a clear, visible signal of, and an important basis for modelling, how global warming influences terrestrial ecosystems (Chuine & Régnière 2017; Zhang *et al.* 2020). Advances in the dates of spring leaf-out in response to warming have been consistently observed over recent decades (Fu *et al.* 2015; Chen *et al.* 2018). Both advanced and delayed trends in leaf senescence have been reported under warming conditions (Menzel *et al.* 2006; Jeong *et al.* 2011; Gill *et al.* 2015). The ecological mechanisms underlying these contradictory warming responses remain unclear, making it difficult to predict how global warming will impact tree leaf senescence and forest ecosystem functioning in the future (Chen *et al.* 2020; Zhang *et al.* 2020).

The final stage through which tree leaves pass before death is accompanied by the degradation of various macromolecules (Woo *et al.* 2013). The main function of leaves at this late stage of the season is to remobilize nutrients from aging leaves into perennial trunks, twigs, and roots for overwintering and to support growth in the following spring (Vergutz *et al.* 2012). Trees have been shown to delay leaf senescence in order to remobilize more nutrients from old leaves (Estiarte & Penuelas 2015). The progress of leaf senescence therefore depends on whether nutrients have been resorbed to their maximum potential extent. However, the timing of leaf senescence is also determined by the maximum amount of assimilated carbon that can be stored (or sink limitation of photosynthesis) early in the growing season (Paul & Foyer 2001). If warming (or other

factors, e.g., elevated carbon dioxide and light levels) speeds up the rate of

photosynthesis and subsequently the rate at which this maximum storage capacity is

reached, then leaf senescence will be advanced (Fu *et al.* 2014).

Using 500,000 phenological records for 15 temperate trees at 5,000 sites,

phenological metrics derived from PhenoCam and satellite imagery, and carbon flux

measurements from 72 sites in the Northern Hemisphere (Fig. S1), we carried out

detailed analyses of the responses of leaf senescence to warming during the growing

season and aim to disentangle the mechanisms underlying the observed contrasts in the

responses of leaf senescence to warming and provide an ecological basis for predicting

the trajectory of leaf senescence under future warming.

METHODS SUMMARY

Phenology and climate data

Leaf senescence data were obtained from three large phenological datasets in the

Northern Hemisphere: PEP725 phenological database, PhenoCam network and

MODIS phenology product. The FLUXNET2015 dataset was used to examine the

temporal variations in photosynthetic carbon assimilation in growing season. Gridded

daily temperatures, precipitation, radiation, and humidity data with a spatial resolution

of 0.25° in Europe were collected from the database E-OBS. Gridded CLM/ERAi soil

moisture (0-10cm) data between were downloaded from KNMI Climate Explorer.

Global monthly mean temperature data with 0.5° spatial resolution were downloaded

from the Climate Research Unit (CRU TS Version 4.04).

Statistical analysis

Using the PEP725 data, we calculated the daily, weekly, and monthly temperature

sensitivity (S_T, change in days per degree Celsius) of leaf senescence during the

growing season between May and September for each species at each site. From the

calculated daily, weekly and monthly S_T of leaf senescence, we found that responses of

leaf senescence to warming changed from negative in May and June to positive between

July and September. We therefore divided the entire season into two periods: early

season (May-June) and late season (July-September), and further calculated the mean

S_T during the two periods to obtain the S_T during the early ($S_{T-Early}$) and late (S_{T-Late})

season, respectively. The sum of $S_{T-Early}$ and S_{T-Late} of leaf senescence was used to

represent the overall warming responses of leaf senescence during the whole season

while leaves were present. We also calculated the monthly S_T of leaf senescence

between May and September during 2000-2018 based on the dates of leaf senescence

extracted from the PhenoCam network and MODIS phenology product.

Using the PEP725 network, we examined the effects of growing degree days (GDD)

on the leaf senescence dates (change in days GDD^{-1}) in years under low and high

nighttime temperature conditions during the early and late growing season. We

calculated and compared the S_T of leaf senescence during the early, late, and entire

season during the coldest period 1953-1972 and the warmest period 1992-2011 over

the past 60 years. Using partial correlation analysis, we excluded the covariate effects

of soil moisture, precipitation, radiation, humidity and further examined the

relationship between monthly mean temperature from May to September and leaf

senescence dates. Furthermore, we quantified and compared the relative influences of

temperature, soil moisture, precipitation, radiation, humidity on leaf senescence date in

the growing season using boosted regression trees (BRTs).

Using the FLUXNET2015 dataset, we examined the temporal changes in the

photosynthesis carbon assimilation during the growing season based on the Net

Ecosystem Exchange (NEE); we calculated and compared the differences in the

nighttime respiration during the early season (May and June) and the number of frost

days ($T_{min} < 0°C$) in late autumn (October and November) during years with low and

high nighttime temperature.

Full methods and any associated references are available in the Supplementary

Methods.

RESULTS

Using records of leaf senescence for 15 temperature tree species at 5,000 sites from

the PEP725 network, we found the mean S_T of leaf senescence was negative in May

and June, while it gradually converted to positive between July and September (Fig. 1a,

residual degrees of freedom $df=1.02 \times 10^5$). Based on the daily and weekly S_T of leaf

senescence, we also observed a significant increase in S_T throughout the entire season

(Fig. S2a, $df=159$, $P<0.001$; Fig. S2b, $df=20$, $P<0.001$). According to the linear mixed

models, the overall S_T of leaf senescence during the early and late season across all

species and sites were approximately -1.14 and +1.33 days $°C^{-1}$, while S_T of leaf

senescence of the total season was +0.12 days $°C^{-1}$ (Table S2). After excluding the

effects of other climate variables, using partial correlation analysis we also observed a

negative response of leaf senescence to temperature during the early season, but a

positive response during the late season (Fig. S3, $df=4.02 \times 10^4$, $P<0.05$). According to

the calculated relative influence, temperature had the strongest effect on leaf senescence,

followed by soil moisture and radiation (Fig. S4).

Using the PhenoCam network, we again observed a transition in the S_T of leaf

senescence in North America from May to September (Fig. 1b, *df*=558, *P*<0.001).

Based on phenology metrics extracted from MODIS, we consistently observed a

transition in the S_T of leaf senescence from May to September across all biomes except

deserts and xeric shrublands in the Northern Hemisphere (Fig. 1c, *df*=1.30×10^5,

P<0.001). In May and June, the effects of temperature on leaf senescence were negative

across temperate and boreal biomes (Fig. 1c, *df*=1.30×10^5, *P*<0.001). We also

observed negative effects of temperatures in May and June on leaf senescence in tundra,

alpine, Mediterranean regions, deserts and xeric shrublands (Fig. 1c, *df*=1.30×10^5,

P<0.001). The effects of temperature gradually became positive in August or

September in these biomes (Fig. 1c, *df*=1.30×10^5, *P*<0.001). The spatial distribution

of the monthly S_T from May to September in the Northern Hemisphere was shown in

Fig. S5.

Using the FLUXNET2015 data, we detected an obvious changing point at the end

of June (DOY 180) for the net daily carbon assimilation (Fig. 2, n=1.46×10^3, *P*<0.05).

Generally, net carbon assimilation was positive during the early season (before June)

but was negative during the late season (after June) (Fig. 2a, *P*<0.001). This suggested

that carbon assimilation mainly occurs before June. The difference in the net carbon

assimilation between early season (before June) and late season (after June) in forest

was the largest, followed by grassland and savanna (Fig. 2b, *df*=1.46×10^3, *P*<0.05).

Results showed that the timing of leaf senescence was also advanced by greater

GDDs during the early season (May-June), but delayed by greater GDDs during the

late season (July-September) (Fig. S6ab, *df*=1.39×10^4, *P*<0.001). We found that during

the early season, the negative effect of GDDs on leaf senescence was stronger during

years with low nighttime temperatures (Fig. S6a, n=1.39×10^4, $P<0.001$). Furthermore,

during the late season the positive effect of GDDs on leaf senescence was weaker during

years with low nighttime temperature (Fig. S6b, df=1.39×10^4, $P<0.001$). When

nighttime temperature was higher, nighttime ecosystem respiration was significantly

greater during the early season (Fig. S6c, df=840, $P<0.05$), while the number of frost

days was significantly lower during the late season (Fig. S6d, df=844, $P<0.001$).

To assess the effects of climate warming on leaf senescence, we used the PEP725

dataset to calculate and compare the S_T of leaf senescence between the coldest and the

warmest 20-year periods: 1953-1972 and 1992-2011. We found that both the $S_{T\text{-Early}}$

and $S_{T\text{-Late}}$ of leaf senescence were significantly higher during 1992-2011 than those

during 1953-1972 (Fig. 3, df=1.63×10^4, $P<0.05$). However, $S_{T\text{-Late}}$ of leaf senescence

during 1992-2011 increased more compared to $S_{T\text{-Early}}$ of leaf senescence (Fig. 3, n=1.63

$\times 10^4$, $P<0.05$). The S_T of leaf senescence during the whole growing season, i.e., the

sum of $S_{T\text{-Early}}$ and $S_{T\text{-Late}}$ of leaf senescence, also showed a significant increase (Fig. 3,

df=1.63×10^4, $P<0.05$).

DISCUSSION

Earlier leaf senescence reduces photosynthetic carbon assimilation and nutrient

resorption efficiency (i.e., the proportion of nutrients resorbed from old leaves) (Estiarte

& Penuelas 2015). However, trees experiencing late leaf senescence are more at risk

from frost (Schwartz 2003), which may reduce nutrient resorption (Estiarte & Penuelas

2015). The optimal timing of leaf senescence is therefore likely to be a trade-off

between photosynthetic carbon assimilation and autumnal nutrient resorption at

different stages in the seasonal functioning of leaves (Fracheboud *et al.* 2009) (Fig. S7).

When trees reach their maximum carbon storage capacity, they will initialize nutrient

resorption and senescence. Accordingly, more efficient accumulation of carbohydrates

with warmer temperatures in the early season will result in a relatively shorter period

being required to reach the maximum carbon storage capacity (Peng *et al.* 2013). Thus,

early season warming advances leaf senescence. Using the flux data, we further found

net carbon assimilation converted from positive to negative at the end of June (DOY

180). This provides direct physiological evidence that photosynthetic carbon

assimilation mainly occurred during the early season (before June). However, warmer

temperatures later in the season may reduce the risk of late autumn frost (Vitasse *et al.*

2014), enhancing the activities of photosynthetic enzymes (Shi *et al.* 2014) and

reducing the rate of chlorophyll degradation (Fracheboud *et al.* 2009; Estiarte &

Penuelas 2015). This may prolong nutrient remobilization from leaves, reduce

degradation rate of organelle dismantling, increase leaf longevity and eventually delay

the final stage of leaf senescence (Kikuzawa 1995). Therefore, leaf senescence was

advanced by warming during the early season, but was delayed by the warming during

the late season in temperate regions. However, leaf senescence was advanced by

warming throughout both early and late seasons in deserts and xeric shrublands. This

may result from drought stress caused by warmer autumns in arid regions increasing

evaporative demand (Chen *et al.* 2017) and thus initiating leaf senescence early

(Estiarte & Penuelas 2015; Chen *et al.* 2020), supported by the positive correlations

between soil moisture and leaf senescence in May and July.

The parameter growing degree-days (GDD) only takes into account the heat

accumulated above the minimum threshold of temperature that must be exceeded for

tree growth (Miller *et al.* 2001) and therefore provides a more accurate physiological

assessment of leaf senescence in response to warming (Chen *et al.* 2020). Our

observation that leaf senescence was advanced by GDD-based warming during the

early growing season, but was delayed by GDD-based warming during the late

resorption season, are consistent with the contrasting warming responses of leaf

senescence reported previously (Menzel *et al.* 2006; Jeong *et al.* 2011; Gill *et al.* 2015).

These results similarly support trade-off of leaf senescence between carbon assimilation

and nutrient resorption. We further examined this trade-off by comparing the warming

responses of leaf senescence when the nighttime temperature changed, because of the

asymmetric effects of nighttime temperature on carbon assimilation during the early

growing season and frost avoidance during the late resorption season (Peng *et al.* 2013;

Chen *et al.* 2020). As a result, trees will reach their maximum carbon capacity quickly

when nighttime temperatures are lower according to such a trade-off assumption. By

contrast, trees can be expected to accomplish nutrient resorption rapidly during the late

season in order to reduce the risk of frost when nighttime temperatures are lower

(Silvestro *et al.* 2019). Consistent with this, we found that the signals of the effects of

warming on leaf senescence were stronger or weaker when the nighttime temperature

was lower during the early or late season respectively. Our findings indicate that not

only the carbon sink limitation in the early season but also nutrient resorption in the late

season should be considered when modelling autumn phenology of temperate trees

under future warming scenarios. More important, our statistical analyses of leaf

senescence during the warmest 20-year periods across both seasons suggest that global

warming may delay leaf senescence in the future.

In addition to temperature, photoperiod may influence tree phenology (Körner &

Basler 2010). The previous work has evidenced that autumnal phenological events are

commonly more sensitive to photoperiod compared with spring events (Way &

Montgomery 2015). However, such a conservative response to warming may be due to

the counterbalancing effects of warming on leaf senescence at different seasons.

Additionally, despite the photoperiodic control of leaf senescence (Way &

Montgomery 2015), we found that temperature alone had strong predictive power even

when photoperiod was not considered, indicating the dominant role of temperature in

leaf senescence.

Overall, our findings based on three large and complementary datasets illustrate

that the onset of leaf senescence is advanced under early season warming, but delayed

when warming occurs in the late stages of the growing season. Although further

controlled warming experiments in different seasons should be conducted to test the

contrasting seasonal climatic responses of leaf senescence, our study provides new

insights into how to accurately predict whether leaf senescence will be delayed or

advanced in response to climate warming (Menzel *et al.* 2006). If future warming spans

both early and late seasons in temperate regions, as found previously (Menzel *et al.*

2006; Gill *et al.* 2015), leaf senescence could be delayed, rather than advanced.

ACKNOWLEDGEMENTS

We acknowledge all members of the PEP725 network for collecting and providing the

phenological data. This research was supported by the Strategic Priority Research

Program of the Chinese Academy of Sciences (XDB31010300) and the National Key

Research and Development Program of China (2017YFC0505203), and also by the

National Natural Science Foundation of China (grant numbers 31590821, 91731301

and 31561123001), the Fundamental Research Funds for the Central Universities

(2018CDDY-S02-SCU and SCU2019D013), and National High-Level Talents Special

Support Plans.

DATA ACCESSIBILITY

The ground observation phenological data are available at www.pep725.eu. The

phenological metrics derived from digital camera imagery are available at

https://lpdaac.usgs.gov/. The phenology data extracted from satellite images can be downloaded from https://daac.ornl.gov/cgi-bin/dsviewer.pl?ds_id=1674. The climate data used in this study are available at http://ensembles-eu.metoffice.com and https://crudata.uea.ac.uk /cru/data/hrg/ cru_ts_4.04/cruts.2004151855.v4.04/.

Literature Cited

Chen, L., Hänninen, H., Rossi, S., Smith, N.G., Pau, S., Liu, Z. *et al.* (2020). Leaf senescence exhibits stronger climatic responses during warm than during cold autumns. *Nat. Clim. Change*, 10, 777-780.

Chen, L., Huang, J.G., Dawson, A., Zhai, L., Stadt, K.J., Comeau, P.G. *et al.* (2018). Contributions of insects and droughts to growth decline of trembling aspen mixed boreal forest of western Canada. *Glob. Chang. Biol.*, 24, 655-667.

Chen, L., Huang, J.G., Alam, S.A., Zhai, L., Dawson, A., Stadt, K.J. *et al.* (2017). Drought causes reduced growth of trembling aspen in western Canada. *Glob. Chang. Biol.*, 23, 2887-2902.

Chuine, I. & Régnière, J. (2017). Process-based models of phenology for plants and animals. *Annu. Rev. Ecol. Evol. Syst.*, 48, 159-182.

Cleland, E.E., Chuine, I., Menzel, A., Mooney, H.A. & Schwartz, M.D. (2007). Shifting plant phenology in response to global change. *Trends Ecol. Evol.*, 22, 357-365.

Estiarte, M. & Penuelas, J. (2015). Alteration of the phenology of leaf senescence and fall in winter deciduous species by climate change: effects on nutrient proficiency. *Glob. Chang. Biol.*, 21, 1005-1017.

Fracheboud, Y., Luquez, V., Björkén, L., Sjödin, A., Tuominen, H. & Jansson, S. (2009). The control of autumn senescence in European aspen. *Plant Physiol.*, 149, 1982-1991.

Fu, Y., Campioli, M., Vitasse, Y., De Boeck, H.J., Van den Berge, J., AbdElgawad, H.

et al. (2014). Variation in leaf flushing date influences autumnal senescence and

next year's flushing date in two temperate tree species. *Proc. Nat. Acad. Sci.*,

111, 7355-7360.

Fu, Y.H., Zhao, H., Piao, S., Peaucelle, M., Peng, S., Zhou, G. *et al.* (2015). Declining

global warming effects on the phenology of spring leaf unfolding. *Nature*, 526,

104-107.

Gill, A.L., Gallinat, A.S., Sanders-DeMott, R., Rigden, A.J., Short Gianotti, D.J.,

Mantooth, J.A. *et al.* (2015). Changes in autumn senescence in northern

hemisphere deciduous trees: a meta-analysis of autumn phenology studies. *Ann.*

Bot., 116, 875-888.

Jeong, S.J., HO, C.H., GIM, H.J. & Brown, M.E. (2011). Phenology shifts at start vs.

end of growing season in temperate vegetation over the Northern Hemisphere

for the period 1982–2008. *Glob. Chang. Biol.*, 17, 2385-2399.

Kikuzawa, K. (1995). Leaf phenology as an optimal strategy for carbon gain in plants.

Can. J. Bot., 73, 158-163.

Körner, C. & Basler, D. (2010). Phenology under global warming. *Science*, 327, 1461-

1462.

Menzel, A., Sparks, T.H., Estrella, N., Koch, E., Aasa, A., Ahas, R. *et al.* (2006).

European phenological response to climate change matches the warming pattern.

Glob. Chang. Biol., 12, 1969-1976.

Miller, P., Lanier, W. & Brandt, S. (2001). Using growing degree days to predict plant

stages. *Ag/Extension Communications Coordinator, Communications Services*,

Montana State University-Bozeman, Bozeman, MO.

Paul, M.J. & Foyer, C.H. (2001). Sink regulation of photosynthesis. *J. Exp. Bot.*, 52,

1383-1400.

Peng, S., Piao, S., Ciais, P., Myneni, R.B., Chen, A., Chevallier, F. *et al.* (2013). Asymmetric effects of daytime and night-time warming on Northern Hemisphere vegetation. *Nature*, 501, 88-92.

Piao, S., Liu, Q., Chen, A., Janssens, I.A., Fu, Y., Dai, J. *et al.* (2019). Plant phenology and global climate change: Current progresses and challenges. *Glob. Chang. Biol.*, 25, 1922-1940.

Richardson, A.D., Hufkens, K., Milliman, T., Aubrecht, D.M., Furze, M.E., Seyednasrollah, B. *et al.* (2018). Ecosystem warming extends vegetation activity but heightens vulnerability to cold temperatures. *Nature*, 560, 368-371.

Richardson, A.D., Keenan, T.F., Migliavacca, M., Ryu, Y., Sonnentag, O. & Toomey, M. (2013). Climate change, phenology, and phenological control of vegetation feedbacks to the climate system. *Agric For Meteorol*, 169, 156-173.

Schwartz, M.D. (2003). *Phenology: an integrative environmental science*. Springer.

Shi, C., Sun, G., Zhang, H., Xiao, B., Ze, B., Zhang, N. *et al.* (2014). Effects of warming on chlorophyll degradation and carbohydrate accumulation of alpine herbaceous species during plant senescence on the Tibetan Plateau. *PLoS One*, 9, e107874.

Silvestro, R., Rossi, S., Zhang, S., Froment, I., Huang, J.G. & Saracino, A. (2019). From phenology to forest management: Ecotypes selection can avoid early or late frosts, but not both. *For. Ecol. Manag.*, 436, 21-26.

Vergutz, L., Manzoni, S., Porporato, A., Novais, R.F. & Jackson, R.B. (2012). Global resorption efficiencies and concentrations of carbon and nutrients in leaves of terrestrial plants. *Ecol. Monogr.*, 82, 205-220.

Vitasse, Y., Lenz, A. & Körner, C. (2014). The interaction between freezing tolerance and phenology in temperate deciduous trees. *Front. Plant Sci.*, 5, 541.

Way, D.A. & Montgomery, R.A. (2015). Photoperiod constraints on tree phenology,

performance and migration in a warming world. *Plant Cell Environ.*, 38, 1725-

1736.

Woo, H.R., Kim, H.J., Nam, H.G. & Lim, P.O. (2013). Plant leaf senescence and death–

regulation by multiple layers of control and implications for aging in general. *J.*

Cell. Sci., 126, 4823-4833.

Zhang, Y., Commane, R., Zhou, S., Williams, A.P. & Gentine, P. (2020). Light limitation

regulates the response of autumn terrestrial carbon uptake to warming. *Nat.*

Clim. Change, 10, 739-743.

FIGURE CAPTIONS

Fig. 1 Temperature sensitivities (S_T, change in days $℃^{-1}$) of leaf senescence during the

growing season between May and September. The calculated S_T was based on **(a)**

records of leaf senescence for 15 temperate tree species at 5,000 sites in Europe, and

phenological metrics extracted from **(b)** the PhenoCam network and **(c)** the MODIS

phenology product (MCD12Q2 version 6) for different biomes. The length of each box

indicates the interquartile range, the horizontal line inside each box the median, and the

bottom and top of the box the first and third quartiles respectively. The biomes are

Tundra (TUN), Boreal Forests/Taiga (BFT), Temperate Broadleaf & Mixed Forests

(TBM), Temperate Conifer Forests (TCF), Temperate Grasslands, Savannas &

Shrublands (TGS), Montane Grasslands & Shrublands (MGS), Mediterranean Forests,

Woodlands & Scrub (MFW), and Deserts & Xeric Shrublands (DXS). The asterisks

indicate a significant difference in the S_T during the early (May-June) and the late (July-

September) season ($P<0.05$).

Fig. 2 Change in the net daily photosynthetic carbon assimilation (g $Cm^{-2}d^{-1}$). The

estimation of the net carbon assimilation was calculated by multiplying the Net

Ecosystem Exchange (NEE) by -1. The length of each box in **(b)** indicates the

interquartile range, the horizontal line inside each box the median, and the bottom and

top of the box the first and third quartiles respectively. The asterisks indicate a

significant difference in the net photosynthetic carbon assimilation during the early

season (before June) and late season (after June). Different color lines and boxes

represent different vegetation types. The dashed vertical line indicates the change point

of net photosynthetic carbon assimilation (DOY 180).

Fig. 3 Temperature sensitivities (S_T, change in days $℃^{-1}$) of leaf senescence in 15

temperate tree species during 1953-1972 and 1992-2011. Calculations of the S_T values

were based on the temperature in early (May and June), late (July-September) and entire

(May-September) growing seasons. The length of each box indicates the interquartile

range, the horizontal line inside each box the median, and the bottom and top of the box

the first and third quartiles respectively. The asterisks indicate a significant difference

in the S_T between 1953-1972 and 1992-2011 ($P<0.05$).

Fig. 1

Fig. 2

Fig. 3