Trade-offs between cost of ingestion and rate of intake drive defensive toxin use

Tyler E. Douglas[1*], Sofia G. Beskid[1,2,+], Callie E. Gernand[3,+], Brianna E. Nirtaut[3,+], Kristen E.
Tamsil[1,+], Richard W. Fitch[3], and Rebecca D. Tarvin[1*]
[1] Museum of Vertebrate Zoology and Department of Integrative Biology, University of
California Berkeley, Berkeley, CA 94720
[2] Department of Integrative Biology, University of Texas at Austin, Austin, TX, 78712
[3] Department of Chemistry and Physics, Indiana State University, Terre Haute, IN 47809
[+] Authors listed in alphabetical order
9
*Corresponding authors:
Tyler Douglas (tyler.douglas@berkeley.edu)
Rebecca Tarvin (rdtarvin@berkeley.edu)
Phone: 510 642 0308
3101 Valley Life Sciences Building
Berkeley, CA, 94720

16
17   **Keywords:** xenobiotic metabolism, chemical defense, multi-trophic selection, bioaccumulation,
18   enemy-free space

19   **Word count:** 2664

Abstract

Animals that ingest toxins can become unpalatable and even toxic to predators and parasites through toxin sequestration. Because most animals rapidly eliminate toxins to survive their ingestion, it is unclear how populations transition from susceptibility and toxin elimination to tolerance and accumulation as chemical defense emerges. Studies of chemical defense have generally focused on species with active toxin sequestration and target-site insensitivity mutations or toxin-binding proteins that permit survival without necessitating toxin elimination. Here, we investigate whether animals that presumably rely on toxin elimination for survival can utilize ingested toxins for defense. We use the A4 and A3 *Drosophila melanogaster* fly strains from the Drosophila Synthetic Population Resource (DSPR), which respectively possess elevated and reduced metabolic nicotine resistance amongst DSPR fly lines. We find that ingesting nicotine increased A4 but not A3 fly survival against *Leptopilina heterotoma* wasp parasitism. Further, we find that despite possessing genetic variants that enhance toxin elimination, A4 flies accrued more nicotine than A3 individuals likely by consuming more media. Our results suggest that enhanced toxin metabolism can allow for greater toxin intake by offsetting the cost of toxin ingestion. Passive toxin accumulation that accompanies increased toxin intake may underlie the early origins of chemical defense.

Introduction

Most animals survive toxin ingestion by eliminating toxins through metabolic detoxification (1–3). Some chemically defended animals subvert this paradigm by sequestering dietary toxins to deter predators or parasites (4). Because metabolic detoxification serves to prevent toxin accumulation, toxin-sequestering taxa often employ resistance mechanisms that do not degrade toxins (5). For example, target-site insensitivity (TSI), which results from mutations in a protein that prevent toxins from binding, is common in toxin-sequestering insects (6, 7). TSI sometimes co-occurs with toxin-binding proteins that scavenge toxins and prevent them from binding to targets (8–11). Such non-metabolic resistance mechanisms may facilitate the transition from toxin elimination to sequestration by decreasing reliance on toxin breakdown for survival (12).

Although metabolic detoxification degrades toxins, it is unclear whether reliance on this mechanism constrains chemical defense evolution. Metabolic detoxification permits toxin consumption and may ultimately lead to toxin sequestration so long as consumption outpaces degradation. To test this idea, we obtained two isofemale, homozygous strains of *Drosophila melanogaster* from the Drosophila Synthetic Population Resource (DSPR (13)) that possess high and low nicotine resistance (A3 and A4, Bloomington stocks 3852 and 3844), and exposed them to nicotine, a plant allelochemical that targets acetylcholine receptors (14). Although some drosophilids do feed on toxic food sources (15, 16) and the A4 fly strain may have experienced incidental nicotine exposure on tobacco farms that were prevalent at its collection site (17), drosophilids are not known to select nicotine-producing plants as hosts. Nevertheless, the genetic basis of nicotine resistance in *D. melanogaster* is extensively characterized, making this toxin well-suited to modelling the evolutionary origins of chemical defense (18). Compared to A3, A4

flies possess duplicate copies of cytochrome p450s *Cyp28d1* and *Cyp28d2* that are constitutively
expressed at higher levels. A4 flies also overexpress the UDP-glucuronosyltransferase *Ugt86Dd*,
while A3 harbors a mutation in this gene that significantly reduces nicotine resistance (19).
Ugt86Dd is located in a Quantitative Trait Locus (QTL) that contributes 50.3% of the broad-
sense heritability in nicotine resistance of DSPR lines, while a QTL containing *Cyp28d1* and
Cyp28d2 accounts for 5% (18, 20). The contributions of these three genes to nicotine resistance
have been confirmed using gene knockout (21). Previous QTL and expression-QTL studies did
not report evidence for TSI or toxin-binding proteins in A3 or A4 lines. While these mechanisms
could exist, variation in metabolic enzymes appears to underlie the major difference between A3
and A4 nicotine resistance.

Results and Discussion

We first quantified A3 and A4 nicotine resistance by estimating the median lethal
concentration (LC50) of nicotine (Fig. 1). Because A4 flies had low viability in general, to
compare LC50 between strains for this assay we normalized percent survival by the maximum
survival of each line on control food (see supporting data for non-normalized values). The A4
LC50 was nearly twice that of A3 ($LC50_{A4} = 1.9 \pm 0.3$ mM [mean $\pm$ SD], $LC50_{A3} = 1.1 \pm 0.2$
mM; Fig. 1). While A3 survival decreased significantly at 0.5 mM nicotine, A4 survival was not
significantly impacted until 1.75 mM. We proceeded to use an intermediate level of 1.25-mM
nicotine for subsequent experiments.

We next assessed whether ingesting 1.25-mM nicotine after parasitism by the figitid
wasp *Leptopilina heterotoma* increased *D. melanogaster* survival. *Leptopilina heterotoma*
oviposits into the hemocoel of developing fly larvae, and actively suppresses the drosophilid
defensive immune response against endoparasites (22). Thus, developing parasites are exposed
to host hemolymph and, presumably, to circulating toxins consumed by fly larvae. In the control-
fed, unparasitized treatment, $2.8\% \pm 2.7\%$ of A4 larvae survived to adulthood, while in the
nicotine-fed, parasitized treatment, A4 survival increased significantly to $6.8 \pm 4.4\%$ ($p = 0.03$, Z
$= -2.2$; Fig. 2A). Correspondingly, *L. heterotoma* developmental success decreased five-fold
from $37\% \pm 20\%$ to $6.4\% \pm 6.8\%$ ($p < 0.0001$, $Z = 7.0$; Fig. 2B). Thus, nicotine consumption
increased A4 fly survival against parasitism.

In contrast, the survival of parasitized, nicotine-fed A3 larvae ($15 \pm 4.4\%$) was the same
as parasitized, control-fed A3 larvae ($19 \pm 10\%$; $p = 0.36$, $Z = 0.92$; Fig. 2A). However, wasp
developmental success on A3 flies halved from $41 \pm 15\%$ to $21 \pm 9.3\%$ when A3 flies consumed
nicotine ($p = 0.0001$, $Z = 4$; Fig. 2B). This suggests nicotine consumption partially alleviated A3
parasitism-induced mortality. Nicotine consumption decreased unparasitized A3 fly survival by
44% ($p < 0.0001$, $Z = 7.6$), while nicotine consumption decreased parasitized A3 survival by
only a tenth as much: 3.5%. The comparatively insignificant effect of nicotine consumption on
parasitized A3 flies paired with a ~50% decrease in wasp success suggests that nicotine may
have offset parasitism-induced mortality for A3 flies, although to a lesser degree compared to A4
flies.

Next, we quantified nicotine accumulation in whole bodies of nicotine-fed larvae and
adult flies. After 24hr ± 2.5hr on nicotine media, third-instar A4 larvae contained twice as much
nicotine as A3 larvae (9.3 ± 4.6 vs. 4.3 ± 1.0 ng nicotine, $p = 0.016$, $W = 1$; Fig. 2D). Nicotine
continued to accumulate until pupation and persisted through metamorphosis in both strains (Fig.
2D; also observed with ouabain [6]), suggesting that nicotine remained after the meconium was
shed and may provide a defensive advantage into adulthood. The greater amount of nicotine in
A4 could underlie the stronger effect of nicotine on parasite success in A4 versus A3 individuals
(Fig. 2B). Although nicotine-fed A3 adults are ~20% smaller than nicotine-fed A4 adults (Fig.
2C), this difference cannot explain the two-fold difference observed in nicotine accumulation
between strains. The developmental rate of nicotine-fed A3 and A4 flies did not differ
significantly at 1.25 mM nicotine and is also unlikely to underlie differences in nicotine
accumulation (Fig. S1).

Our finding that A4 larvae accumulated more nicotine than A3 defies genotypic
expectations, as A4 flies have genetic variants that are expected to increase nicotine breakdown
(19, 21). To better understand this pattern, we compared relative amounts of cotinine, a
metabolic by-product of nicotine (Fig 2D) between strains. A4 larvae contained significantly
higher levels of cotinine compared to A3 individuals (Fig. 2D). Intriguingly, one-day-old and
three-day-old A3 flies had significantly higher cotinine to nicotine ratios than A4, suggesting that
A4 larvae have a distinct metabolic detoxification pathway compared to A3 ($p_{one-day-old} = 0.031$,
$W_{one-day-old} = 23$, $p_{three-day-old} = 0.008$, $W_{three-day-old} = 25$ (13)). This result matches expectations
based on genotype, as the largest QTL underlying resistance in A4 contains several UGTs, which
convert nicotine to glucuronides instead of cotinine (18).

The higher nicotine levels in A4 flies suggested that A3 flies are unable to survive high
toxin loads, and thus might consume less to avoid nicotine accumulation. To quantify differences
in feeding, we compared A3 and A4 adult body mass when reared on control versus nicotine
food. While nicotine consumption significantly reduced A3 adult body mass, A4 mass remained
unaffected (Fig. 2C), indicating that nicotine sensitivity constrained A3 food intake. The tobacco
hornworm *Manduca sexta* employs a more extreme version of this pattern: nicotine exposure
activates xenobiotic enzymes, which further stimulates feeding (23). Thus, perhaps
unexpectedly, increased metabolic detoxification may promote rather than preclude toxin
accumulation via increased feeding.

Intriguingly, while nicotine consumption increased A4 fly survival against parasitism, A4
flies under all but the nicotine-fed, unparasitized condition had lower viability than A3 flies (Fig.
2A). Thus, in a hypothetical population made only of A3 and A4 flies and exposed to *L.*
heterotoma and nicotine, natural selection may be unlikely to favor A4 individuals. In this
scenario, the evolutionary outcome would depend partly on whether antagonistic pleiotropy
exists among loci determining metabolic resistance and viability. One general viability QTL has
been identified in DSPR strains, but this QTL does not contain detoxification genes. Moreover,
A4 and A3 flies seem to share the same allele at this QTL (15). Furthermore, while A4 survival
was generally lower than A3, A3 (and not A4) female body mass was reduced by nicotine
consumption. Body mass is correlated with fecundity in *D. melanogaster*, and thus nicotine-fed

A4 flies may have greater reproductive success than A3 (24), which would potentially offset the
cost of lower survival.

To our knowledge, *D. melanogaster* does not possess active nicotine sequestration
mechanisms. Some drosophilids, such as *D. sechellia,* are known to acquire chemical defenses
from toxic food sources (25), and *D. melanogaster* self-medicates against parasitoids using
ethanol (26). However, other drosophilids that consume toxins have not been evaluated for
chemical defenses (15, 16). Our finding that flies can utilize nicotine for defense without active
sequestration mechanisms suggests that other organisms that tolerate toxin consumption could
receive a transient defensive advantage, too. The biochemical properties and metabolic context
of each toxin should affect their propensity to bioaccumulate. For example, non-toxic
glucosinolates (GLS) rapidly breakdown into toxic mustard oils; thus, GLS-sequestration
requires adaptations that interrupt this process (16). Many organisms sequester toxic steroids or
alkaloids (4, 27, 28), perhaps because these more readily diffuse or are transported across tissues.
Here we find that in addition to having increased nicotine metabolism, A4 *D. melanogaster* flies
also likely consume much higher quantities of nicotine than A3 flies (Fig. 2C). We hypothesize
that higher intake may allow relatively more nicotine to escape metabolism and permeate into the
hemolymph of A4 flies, affecting *L. heterotoma* development to a greater degree than in A3 flies.
This pattern could be verified with future studies that compare nicotine abundance in different
tissues of A4 and A3 flies.

In conclusion, we find that elevated resistance increases passive toxin accumulation.
Further, this accumulation produces a toxin-mediated fitness advantage against natural enemies,
in animals without identified sequestration mechanisms. Reliance on metabolic detoxification is
likely the ancestral character state for organisms with acquired chemical defenses, and variation
in toxin metabolism is common (29). We therefore propose that one of the first steps in the
evolution of chemical defense may paradoxically be natural selection for increased toxin
metabolism.

Methods

Fly and wasp stocks

Flies were maintained at room temperature on molasses media from the Fly Food Facility at
UCB; survival and parasitism experiments used Ward's Instant Drosophila media to facilitate
toxin dosing.

Wasps were maintained at room temperature on W118 *D. melanogaster* and 70%-honey water.
Experiments used wasps within two weeks of eclosion.

Generation of fly larvae

Approximately one-thousand flies were allowed to lay eggs for three days in three replicate
resealable plastic containers with a layer of molasses-agar smeared with yeast paste. Larvae were

then pooled from each container, and second-instar larvae (L2) were selected based on
morphology under a dissection microscope. Flies were not sorted by sex.

Nicotine-resistance experiment

197   Twenty A4 and A3 L2 larvae were transferred one-by-one from egg-laying chambers into 5
198   replicate vials containing the following nicotine concentrations: 0 mM, 0.5 mM, 1.25 mM, 1.75
199   mM, 2.25 mM, 2.50 mM, 3.00 mM, 4.00 mM, 5.00 mM nicotine-treated media. Vials were
200   checked daily for new pupae and eclosed flies, and daily counts were used to calculate
201   developmental rate across nicotine doses (Fig. S1).

Parasitism experiment

For each fly strain, 400 L2 were transferred into six replicate plastic containers containing
molasses agar. Forty female and twenty male wasps were added to three containers ("wasp"
treatment) while the other three were left unmanipulated ("no-wasp" treatment); all containers
were left for 24hr. One "no-wasp" container contained only 80 L2s. The L2s were then counted
individually (to avoid batch bias) into forty vials containing either control or 1.25-mM nicotine
media. We pooled data on A4 flies from two separate runs of this experiment (average survival
was not significantly different between runs). In run 1 (A4 only), we added 20 larvae to each
vial. In run 2 (A4 and A3), we add 16 larvae to each vial. Vials were checked every 1-2 days for
pupation and emergence. Parasitism was performed prior to nicotine treatment to avoid exposing
L. heterotoma adults to nicotine. Therefore, changes in fly and wasp survival reflect the effects
of nicotine consumption by *D. melanogaster* larvae and not any behavioral change by *L.*
heterotoma.

Nicotine accumulation experiment

One-thousand A4/A3 L2 were distributed one-by-one from egg-laying chambers into five 1.25-
mM nicotine-treated vials. At five developmental stages (3^{rd}-instar larvae, day-1 pupa, day-3
pupae [A4 only], day-1 adult, day-3 adult), we collected five individuals and washed them
individually in glass dissection wells with DI H_2O. Pupae were removed from vials prior to
eclosion to avoid contamination of the adult exoskeleton with nicotine. Individuals from each
stage for each vial were pooled and frozen at -20°C.

Frozen flies were thawed and soaked with methanol (50 μL) at room temperature for 2-3 days to
reach equilibrium. Crude methanolic extracts were transferred to limited volume autosampler
vials and injected directly. Gas chromatographic-mass spectrometric conditions were as
previously described (30); full details are given in the Supplementary Material.

Body Mass Measurement

227   300 A3/A4 L2 were placed one-by-one from egg-laying chambers into twenty vials containing
228   either control or 1.25-mM nicotine media. Upon pupation, individuals were removed and placed
229   onto food-free vials. Adults were starved for 48 hours and then weighed.

Statistical Analysis

Statistical analyses were conducted using Rv3.6.1 (31). LC50s were calculated using adapted version of the 'dose.p' function from the 'MASS' package (32) to a binomial regression model of normalized percent survival versus nicotine dose generated by the 'glmer' function from lme4. Fly survival and wasp success were assessed by applying a least-squared-means test to a binomial regression model of survival as a function of nicotine and (for flies) parasite treatments using the 'glm' function from lme4 (33). Adult fly mass was compared by applying the least-squared-means method described above to a model of average mass per vial as a function of nicotine and sex. Developmental rate and mean nicotine content of flies was compared across strains using Wilcoxon signed-rank tests in base R.

Data accessibility

Raw data files, R script, and detailed metadata are available for download from the Dryad Digital Repository: https://doi.org/10.5061/dryad.w3r2280sc (34).

Acknowledgements
We thank Kirsten Verster, Jessica Aguilar, Mariana Karageorgi, Luis Jazo, Max Lambert for input on the experimental design, Noah Whiteman for helpful discussions, Stuart Macdonald for DSPR lines, Todd Schlenke for wasps. RDT was supported by the Miller Institute for Basic Research in Science, UC Berkeley start-up funds, and the Hellman Fellows Program; RWF was supported by NSF CCLI-DUE-0942345 and DEB-1556982.

Figure 1. A) Nicotine concentration-survival curve for DSPR A3 and A4 *Drosophila melanogaster*. Data are normalized by maximum survival of each strain on control food. Vertical dashed lines represent LC50 of each strain.

Figure 2. A) Nicotine consumption significantly decreases survival in unparasitized A3 and A4 *Drosophila melanogaster* flies. Nicotine consumption increases survival of parasitized A4 but not A3 flies. **B)** Nicotine consumption by A4 and A3 flies significantly decreases *Leptopilina heterotoma* developmental success. **C)** Nicotine consumption reduced A3 but not A4 adult body mass. **D)** Nicotine-fed A3 and A4 flies accumulate nicotine and its metabolic byproduct cotinine across developmental stages. Asterisks indicate significant differences.

1. P. J. Daborn, *et al.*, A single p450 allele associated with insecticide resistance in *Drosophila*. *Science* **297**, 2253–2256 (2002).

2. X. Li, M. R. Berenbaum, M. A. Schuler, Plant allelochemicals differentially regulate *Helicoverpa zea* cytochrome P450 genes. *Insect Mol. Biol.* **11**, 343–351 (2002).

3. P. Kumar, S. S. Pandit, A. Steppuhn, I. T. Baldwin, Natural history-driven, plant-mediated RNAi-based study reveals *CYP6B46*'s role in a nicotine-mediated antipredator herbivore defense. *Proc. Natl. Acad. Sci.* **111**, 1245–1252 (2014).

4. S. S. Duffey, Sequestration of plant natural products by insects. *Ann Rev. Entomol.* **25**,

447-77 (1980).

5.   G. Petschenka, A. A. Agrawal, Milkweed butterfly resistance to plant toxins is linked to
sequestration, not coping with a toxic diet. *Proc. R. Soc. B Biol. Sci.* **282** (2015).

6.   M. Karageorgi, *et al.*, Genome editing retraces the evolution of toxin resistance in the
monarch butterfly. *Nature* **574**, 409–412 (2019).

7.   A. J. Crossthwaite, S. Rendine, M. Stenta, R. Slater, Target-site resistance to
neonicotinoids. *J. Chem. Biol. 2014 74* **7**, 125–128 (2014).

8.   S. N. Caty, *et al.*, Molecular physiology of chemical defenses in a poison frog. *J. Exp.*
Biol. **222** (2019).

9.   F. Abderemane-Ali, *et al.*, Evidence that toxin resistance in poison birds and frogs is not
rooted in sodium channel mutations and may rely on "toxin sponge" proteins. *J. Gen.*
Physiol. **153** (2021).

10.   R. D. Tarvin, *et al.*, Interacting amino acid replacements allow poison frogs to evolve
epibatidine resistance. **357**, 1261-1266 (2017).

11.   B. L. Whitelaw, I. R. Cooke, J. Finn, K. Zenger, J. M. Strugnell, The evolution and origin
of tetrodotoxin acquisition in the blue-ringed octopus (genus *Hapalochlaena*). *Aquat.*
Toxicol. **206**, 114–122 (2019).

12.   S. Dobler, S. Dalla, V. Wagschal, A. A. Agrawal, Community-wide convergent evolution
in insect adaptation to toxic cardenolides by substitutions in the Na,K-ATPase. *Proc. Natl.*
Acad. Sci. U. S. A. **109**, 13040–13045 (2012).

13.   E. G. King, *et al.*, Genetic dissection of a model complex trait using the *Drosophila*
Synthetic Population Resource. *Genome Res.* **22**, 1558–1566 (2012).

14.   J. E. C. I. Yamamoto, *Nicotinoid Insecticides and the Nicotinic Acetylcholine Receptor*
(1999) https:/doi.org/10.1007/978-4-431-67933-2.

15.   J. C. Fogleman, Response of *Drosophila melanogaster* to selection for P450-mediated
resistance to isoquinoline alkaloids. *Chem. Biol. Interact.* **125**, 93–105 (2000).

16.   P. T. Humphrey, *et al.*, Aversion and attraction to harmful plant secondary compounds
jointly shape the foraging ecology of a specialist herbivore. *Ecol. Evol.* **6**, 3256–3268
(2016).

17.   G. E. Addicott, Rhodesian tobacco and world marketts. *South African J. Econ.* **26**, 29–40
(1958).

18.   T. N. Marriage, E. G. King, A. D. Long, S. J. Macdonald, Fine-mapping nicotine
resistance loci in *Drosophila* using a multiparent advanced generation inter-cross
population. *Genetics* **198**, 45–57 (2014).

19.   C. A. Highfill, *et al.*, Naturally segregating variation at *Ugt86Dd* contributes to nicotine
resistance in *Drosophila melanogaster*. *Genetics* **207**, 311–325 (2017).

20.   M. Chakraborty, J. J. Emerson, S. J. Macdonald, A. D. Long, Structural variants exhibit
widespread allelic heterogeneity and shape variation in complex traits. *Nat. Commun.* **10**,
1–11 (2019).

21.   S. J. Macdonald, C. A. Highfill, A naturally-occurring 22-bp coding deletion in *Ugt86Dd*
reduces nicotine resistance in *Drosophila melanogaster*. *BMC Res. Notes* **13**, 1–6 (2020).

22.   R. M. Rizki, T. M. Rizki, Selective destruction of a host blood cell type by a parasitoid
wasp. *Proc. Nati. Acad. Sci. USA* **81**, 6154-6158 (1984).

23.   M. J. Snyder, J. I. Glendinning, Causal connection between detoxification enzyme activity
and consumption of a toxic plant compound. *J. Comp. Physiol. A* **179**, 255-261 (1996).

24.   P. L. Kwang, *et al.*, Lifespan and reproduction in *Drosophila*: New insights from
nutritional geometry. *Proc. Natl. Acad. Sci. U. S. A.* **105**, 2498–2503 (2008).

25.   L. Salazar-Jaramillo., B. Wertheim, Does *Drosophila sechellia* escape parasitoid attack by
feeding on a toxic resource? *PeerJ* **9** (2021).

26.   N. F. Milan, B. Z. Kacsoh, T. A. Schlenke, Alcohol consumption as self-medication
against blood-borne parasites in the fruit fly. *Curr. Biol.* **22**, 488–493 (2012).

27.   S. C. Groen, *et al.*, Multidrug transporters and organic anion transporting polypeptides
protect insects against the toxic effects of cardenolides. *Insect Biochem. Mol. Biol.* **81**, 51–
61 (2017).

28.   C. T. Hanifin, The chemical and evolutionary ccology of tetrodotoxin (TTX) toxicity in
terrestrial vertebrates. *Mar. Drugs* **8**, 577–593 (2010).

29.   M. A. Najarro, *et al.*, Identifying loci contributing to natural variation in xenobiotic
resistance in *Drosophila. PLOS Genet.* **11**, e1005663 (2015).

30.   R. W. Fitch, T. F. Spande, H. M. Garraffo, H. J. C. Yeh, J. W. Daly, Phantasmidine: an
Epibatidine congener from the Ecuadorian poison frog *Epipedobates anthonyi*. *J. Nat.*
Prod. **73**, 331 (2010).

31.   R Core Team. 2021. R: A language and environment for statistical computing. R
Foundation for Statistical Computing, Vienna, Austria

32.   Venables, W. N., & Ripley, B. D. Modern Applied Statistics with S Fourth edition. *World*
53, (2002).

33.   D. Bates, M. Mächler, B. Bolker, S. Walker, Fitting Linear Mixed-Effects Models Using
lme4. *J. Stat. Softw.* **67**, 1–48 (2015).

34.   T.E. Douglas, S.G. Beskid, C.E. Gernand, B.E. Nirtaut, K.E. Tamsil, R.W. Fitch, R.D.
336   Tarvin. Trade-offs between cost of ingestion and rate of intake drive defensive toxin use.
337   Dryad Digital Repository (doi.org/10.5061/dryad.w3r2280sc) (2021).