Trade-offs between cost of ingestion and rate of intake drive defensive toxin use

Tyler E. Douglas[1*], Sofia G. Beskid[1,2,+], Callie E. Gernand[3,+], Brianna E. Nirtaut[3,+], Kristen E.
Tamsil[1,+], Richard W. Fitch[3], and Rebecca D. Tarvin[1*]
[1] Museum of Vertebrate Zoology and Department of Integrative Biology, University of
California Berkeley, Berkeley, CA 94720
[2] Department of Integrative Biology, University of Texas at Austin, Austin, TX, 78712
[3] Department of Chemistry and Physics, Indiana State University, Terre Haute, IN 47809
[+] Authors listed in alphabetical order
9
*Corresponding authors:
Tyler Douglas (tyler.douglas@berkeley.edu)
Rebecca Tarvin (rdtarvin@berkeley.edu)
Phone: 510 642 0308
3101 Valley Life Sciences Building
Berkeley, CA, 94720

Keywords: xenobiotic metabolism, chemical defense, multi-trophic selection, bioaccumulation,
enemy-free space

Word count: 2453

Abstract

Animals that ingest toxins can become unpalatable and even toxic to predators and parasites
through toxin sequestration. Because most animals rapidly eliminate toxins to survive their
ingestion, it is unclear how populations transition from susceptibility and toxin elimination to
tolerance and accumulation as chemical defense emerges. Studies of chemical defense have
generally focused on species with active toxin sequestration and target-site insensitivity
mutations or toxin-binding proteins that permit survival without necessitating toxin elimination.
Here, we investigate whether animals that presumably rely on toxin elimination for survival can
utilize ingested toxins for defense. We use the A4 and A3 *Drosophila melanogaster* fly strains
from the Drosophila Synthetic Population Resource (DSPR), which respectively possess elevated
and reduced metabolic nicotine resistance amongst DSPR fly lines. We find that ingesting
nicotine increased A4 but not A3 fly survival against *Leptopilina heterotoma* wasp parasitism.
Further, we find that despite possessing genetic variants that enhance toxin elimination, A4 flies
accrued more nicotine than A3 individuals likely by consuming more media. Our results suggest
that enhanced toxin metabolism can allow for greater toxin intake by offsetting the cost of toxin
ingestion. Passive toxin accumulation that accompanies increased toxin intake may underlie the
early origins of chemical defense.

Introduction

Most animals survive toxin ingestion by eliminating toxins through metabolic
detoxification (1–3). Some chemically defended animals subvert this paradigm by sequestering
dietary toxins to deter predators or parasites (4). Because metabolic detoxification serves to
prevent toxin accumulation, toxin-sequestering taxa often employ resistance mechanisms that do
not degrade toxins (5). For example, target-site insensitivity (TSI), which results from mutations
in a protein that prevent toxins from binding, is common in toxin-sequestering insects (6, 7). TSI
sometimes co-occurs with toxin-binding proteins that scavenge toxins and prevent them from
binding to targets (8–11). Such non-metabolic resistance mechanisms may facilitate the
transition from toxin elimination to sequestration by decreasing reliance on toxin breakdown for
survival (12).

Although metabolic detoxification degrades toxins, it also permits toxin consumption and
may ultimately lead to toxin sequestration so long as consumption outpaces degradation. To test
this idea, we obtained two isofemale, homozygous strains of *Drosophila melanogaster* from the
Drosophila Synthetic Population Resource (DSPR (13)) that possess high and low nicotine
resistance (A3 and A4, Bloomington stocks 3852 and 3844), as well as the endoparasitoid figitid
wasp, *Leptopilina heterotoma*. Compared to A3, A4 possess duplicate copies of cytochrome
p450s *Cyp28d1* and *Cyp28d2* that are constitutively expressed at higher levels. A4 flies also
harbor a mutation in the UDP-glucuronosyltransferase *Ugt86Dd*. *Ugt86Dd* is located in a
Quantitative Trait Locus (QTL) that contributes 50.3% of the broad-sense heritability in nicotine
resistance of DSPR lines, while a QTL containing *Cyp28d1* and *Cyp28d2* accounts for 5% (14,
15). The contributions of these three genes to nicotine resistance has been confirmed via gene
knockout (16). Previous QTL and expression-QTL studies did not report evidence for TSI or

toxin-binding proteins in A3 or A4 lines. While these mechanisms could exist, variation in
metabolic enzymes appears to underlie the major difference between A3 and A4 nicotine
resistance.

We found that nicotine consumption by both flies killed many developing wasps,
increasing A4 (but not A3) survival. A4 flies, which should detoxify nicotine more rapidly,
accumulated higher levels of nicotine over a 24-hr period than A3 flies. Feeding assays
suggested that this pattern was a result of reduced food consumption by A3 flies when on
nicotine media. Thus, our findings indicate that increased toxin metabolism may facilitate the
evolution of chemical defense by reducing the cost of toxin consumption and allowing for
transitive accumulation of higher toxin loads.

Results and Discussion

We quantified A3 and A4 nicotine resistance by estimating the median lethal
concentration (LC50) of nicotine (Fig. 1). Because A4 flies had low viability in general, to
compare LC50 between strains for this assay we normalized percent survival by the maximum
survival of each line on control food (see supporting data for non-normalized values). The A4
LC50 was nearly twice that of A3 ($LC50_{A4}$ = 1.9 ± 0.3 mM [mean ± SD], $LC50_{A3}$ = 1.1 ± 0.2
mM; Fig. 1). While A3 survival decreased significantly at 0.5 mM nicotine, A4 survival was not
significantly impacted until 1.75 mM. We proceeded to use an intermediate level of 1.25-mM
nicotine for subsequent experiments.

We next assessed whether ingesting 1.25-mM nicotine after parasitism by *Leptopilina*
heterotoma increased *D. melanogaster* survival. *Leptopilina heterotoma* oviposits into the
hemocoel of developing fly larvae, and actively suppresses the drosophilid defensive immune
response against endoparasites (17). Thus, developing parasites are exposed to host hemolymph
and, presumably, to circulating toxins consumed by fly larvae. In the no-nicotine + wasp
treatment, 2.8% ± 2.7% of A4 larvae survived to adulthood, while in the nicotine + wasp
treatment, A4 survival increased significantly to 6.8 ± 4.4% (p = 0.03, Z = -2.2; Fig. 2A).
Correspondingly, *L. heterotoma* survival decreased five-fold from 37% ± 20% to 6.4% ± 6.8% (p
< 0.0001, Z = 7.0; Fig. 2B). Thus, nicotine consumption provided A4 flies with a toxin-mediated
fitness advantage.

In contrast, the survival of parasitized, nicotine-fed A3 larvae (15 ± 4.4%) was the same
as parasitized, control-fed A3 larvae (19 ± 10%; p = 0.36, Z = 0.92; Fig. 2A). However, wasp
survival on A3 flies halved from 41 ± 15% to 21 ± 9.3% when A3 flies consumed nicotine (p =
0.0001, Z = 4; Fig. 2B). This suggests nicotine consumption partially alleviated A3 parasitism-
induced mortality. Nicotine consumption decreased unparasitized A3 fly survival by 44% (p <
0.0001, Z = 7.6), while nicotine consumption decreased parasitized A3 survival by only a tenth
as much – 3.5%. The comparatively insignificant effect of nicotine consumption on parasitized
A3 flies paired with a ~50% decrease in wasp survival suggests that nicotine may have offset
parasitism-induced mortality for A3 flies, although to a lesser degree compared to A4 flies.

Next, we quantified nicotine accumulation in whole bodies of nicotine-fed larvae and
adult flies. After ~24hr on nicotine media, third-instar A4 larvae contained over three times as
much nicotine as A3 larvae (9.3 ± 4.6 ng vs. 3.2 ± 1.4 ng nicotine, p = 0.021, W = 1; Fig. 2D).
Nicotine continued to accumulate until pupation in both strains and persisted through
metamorphosis (Fig. 2DE; also observed with ouabain; 7), suggesting that nicotine remained
after the meconium was shed and may provide a defensive advantage into adulthood. The greater
amount of nicotine in A4 (Fig. 2DE) could underlie the stronger effect of nicotine on parasite
survival in A4 versus A3 individuals (Fig. 2B). Although nicotine-fed A3 adults are smaller than
nicotine-fed A4 adults (Fig. 2C), the difference is not three-fold and thus body size differences
cannot completely explain variation in nicotine accumulation between strains.

Our finding that A4 larvae accumulated more nicotine than A3 contrasted with
expectations, as A4 flies have genetic variants that are expected to increase nicotine breakdown
(16, 18). To better understand this pattern, we compared relative amounts of cotinine, a
metabolic by-product of nicotine (Fig 2E) between strains using integrated areas from GCMS
traces (no standard was available). A4 larvae and pupae contained significantly higher levels of
cotinine compared to A3 individuals (Fig. 2E). Intriguingly, A3 flies have greater relative
amounts of cotinine than nicotine in all stages, yet A4 flies had relatively higher amounts of
cotinine than nicotine in only the larval stage, suggesting that A4 flies have a distinct metabolic
detoxification pathway compared to A3 (Fig. 2E; 13). This result matches expectations based on
genotype, as the largest QTL underlying resistance in A4 contains several UGTs, which convert
nicotine to glucuronides instead of cotinine (15).

The higher nicotine levels in A4 flies suggested that A3 flies are unable to survive high
toxin loads, and thus might consume less to avoid nicotine accumulation. To quantify differences
in feeding, we compared A3 and A4 adult body mass when reared on control versus nicotine
food. While nicotine consumption significantly reduced A3 adult body mass, A4 mass remained
unaffected (Fig. 2C), indicating that nicotine sensitivity constrained A3 food intake. The tobacco
hornworm *Manduca sexta* employs a more extreme version of this pattern: nicotine exposure
activates xenobiotic enzymes, which further stimulates feeding (19). Thus, perhaps
unexpectedly, increased metabolic detoxification may promote rather than preclude toxin
accumulation.

To our knowledge, *D. melanogaster* does not possess active nicotine sequestration
mechanisms. Some drosophilids, such as *D. sechellia,* are known to acquire chemical defenses
from toxic food sources (20), and *D. melanogaster* self-medicates against parasitoids using
ethanol (21). However, other drosophilids that consume toxins have not been evaluated for
chemical defenses (22, 23). Our finding that flies can utilize nicotine for defense without active
sequestration mechanisms suggests that other organisms that tolerate toxin consumption could
receive a transient defensive advantage, too. The biochemical properties and metabolic context
of each toxin should affect their propensity to bioaccumulate. For example, glucosinolates (GLS)
rapidly breakdown into toxic mustard oils; thus, GLS-sequestration requires adaptations that
interrupt this process (23). Many organisms sequester toxic steroids or alkaloids (4, 24, 25),
perhaps because these more readily diffuse or are transported across tissues into the hemolymph.

Here we find that in addition to having increased nicotine metabolism, A4 *D. melanogaster* flies
also likely consume much higher quantities of nicotine than A3 flies (Fig. 2C). We hypothesize
that higher intake may allow relatively more nicotine to escape metabolism and permeate into the
hemolymph, affecting *L. heterotoma* development to a greater degree than A3 flies. This pattern
could be confirmed with future studies that image nicotine abundance in different tissues of A4
and A3 flies.

In conclusion, we find that elevated resistance increases passive toxin accumulation.
Further, this accumulation produces a toxin-mediated fitness advantage against natural enemies,
in animals without identified sequestration mechanisms. Reliance on metabolic detoxification is
likely the ancestral character state for organisms with acquired chemical defenses, and variation
in toxin metabolism is common (27). We therefore propose that one of the first steps in the
evolution of chemical defense may paradoxically be natural selection for increased toxin
metabolism.

Methods

Fly and wasp stocks

172 Flies were maintained at room temperature on molasses media from the Fly Food Facility at
173 UCB; survival and parasitism experiments used Ward's Instant Drosophila media to facilitate
174 toxin dosing.

175 Wasps were maintained at room temperature on W118 *D. melanogaster* and 70%-honey water.
176 Experiments used wasps within two weeks of eclosion.

Generation of fly larvae

178 Approximately one thousand flies were allowed to lay eggs for three days in three replicate
179 resealable plastic containers with a layer of molasses-agar smeared with yeast paste. Larvae were
180 then pooled from each container, and second instar larvae (L2) were selected based on
181 morphology under a dissection microscope. Flies were not sorted by sex.

Nicotine-resistance experiment

183 Twenty L2 were transferred one-by-one from egg-laying chambers into 5 replicate vials
184 containing the following nicotine concentrations: 0 mM, 0.5 mM, 1.25 mM, 1.75 mM, 2.25 mM,
185 2.50 mM, 3.00 mM, 4.00 mM, 5.00 mM nicotine-treated media. Vials were checked daily for
186 eclosed flies.

Parasitism experiment

Four hundred A4/A3 L2 were transferred from egg-laying chambers into five plastic containers
containing molasses agar. Forty female and twenty male wasps were added to three containers
("wasp" treatment) while the other two along with a sixth container of eighty L2s were left
unmanipulated ("no-wasp" treatment); all containers were left for 24hr. The L2s were then
transferred one-by-one (to avoid batch bias) into forty vials containing either control or 1.25-mM

nicotine media. Vials were checked every 1-2 days for pupation and emergence. Parasitism was
performed prior to nicotine treatment to avoid exposing *L. heterotoma* adults to nicotine.
Therefore, changes in fly and wasp survival reflect the effects of nicotine consumption by *D.*
melanogaster larvae and not any behavioral change by *L. heterotoma*. This experiment was run
twice with A4 flies and data were pooled across runs.

Nicotine accumulation experiment

One thousand A4/A3 L2 were distributed one-by-one from egg-laying chambers into five 1.25-
mM nicotine-treated vials. At five developmental stages (3rd-instar larvae, day-1 pupa, day-3
pupae [A4 only], day-1 adult, day-3 adult), we collected five individuals and washed them
individually in glass dissection wells with DI H$_2$O. Pupae were removed from vials prior to
eclosion to avoid contamination of the adult exoskeleton with nicotine. Individuals from each
stage for each vial were pooled and frozen at -20°C.

Frozen flies were thawed and soaked with methanol (50 μL) at room temperature for 2-3 days to
reach equilibrium. Crude methanolic extracts were transferred to limited volume autosampler
vials and injected directly. Gas chromatographic-mass spectrometric conditions were as
previously described (28) and full details are given int the Supplementary Material.

Body Mass Measurement

300 A3/A4 L2 were placed one-by-one from egg-laying chambers into twenty vials containing
either control or 1.25-mM nicotine media. Upon pupation, individuals were removed and placed
onto food-free vials. Adults were starved for 48 hours and then weighed on a Mettler-Toledo
XP6 scale.

Statistical Analysis

Statistical analyses were conducted using Rv3.6.1 (29). LC50s were calculated using adapted
version of the 'dose.p' function from the 'MASS' package (30) to a binomial regression model of
normalized percent survival versus nicotine dose generated by the 'glmer' function from lme4.
Fly and wasp survival were each compared by applying a least-squared-means test to a binomial
regression model of survival as a function of nicotine and (for flies) parasite treatments using the
'glm' function from the 'lme4' package (31). Adult fly mass was compared by applying the least-
squared-means method described above to a model of average mass per vial as a function of
nicotine and sex. Mean nicotine content of flies and was compared across fly strains using a
Wilcoxon signed-rank tests in base R.

Acknowledgements
We thank Kirsten Verster, Jessica Aguilar, Mariana Karageorgi, Luis Jazo, and Max Lambert for
input on the experimental design, Noah Whiteman for helpful discussions, Stuart Macdonald for
DSPR lines, Todd Schlenke for wasps. RDT was supported by the Miller Institute for Basic
Research in Science and UC Berkeley start-up funds; RWF was supported by NSF CCLI-DUE
0942345 and DEB-1556982.

Figure 1. A) Nicotine concentration-survival curve for DSPR A3 and A4 *Drosophila melanogaster*. Data are normalized by maximum survival of each strain on control food. Vertical dashed lines represent estimated LC50 of each strain.

Figure 2. A) Nicotine consumption significantly decreases survival in unparasitized A3 and A4 *Drosophila melanogaster* flies. Nicotine consumption increases survival of parasitized A4 but not A3 flies. **B)** Nicotine consumption by A4 and A3 flies significantly decreases *Leptopilina heterotoma* survival. **C)** Nicotine consumption reduced A3 but not A4 adult body mass. **D)** Nicotine-fed A3 and A4 flies accumulate nicotine across developmental stages. **E)** A4 flies contained higher amounts of cotinine than A3 flies, but A3 flies contained higher quantities of cotinine relative to nicotine than A4. Plot points and bars indicate mean and standard deviation; asterisks indicate significant differences (at $p < 0.05$).

References

1. P. J. Daborn, *et al.*, A single p450 allele associated with insecticide resistance in *Drosophila*. *Science* **297**, 2253–2256 (2002).

2. X. Li, M. R. Berenbaum, M. A. Schuler, Plant allelochemicals differentially regulate *Helicoverpa zea* cytochrome P450 genes. *Insect Mol. Biol.* **11**, 343–351 (2002).

3. P. Kumar, S. S. Pandit, A. Steppuhn, I. T. Baldwin, Natural history-driven, plant-mediated RNAi-based study reveals CYP6B46's role in a nicotine-mediated antipredator herbivore defense. *Proc. Natl. Acad. Sci.* **111**, 1245–1252 (2014).

4. S. S. Duffey, Sequestration of plant natural products by insects. *Ann Rev. Entomol.* **25**, 447-77 (1980).

5. G. Petschenka, A. A. Agrawal, Milkweed butterfly resistance to plant toxins is linked to sequestration, not coping with a toxic diet. *Proc. R. Soc. B Biol. Sci.* **282** (2015).

6. M. Karageorgi, *et al.*, Genome editing retraces the evolution of toxin resistance in the monarch butterfly. *Nature* **574**, 409–412 (2019).

7. A. J. Crossthwaite, S. Rendine, M. Stenta, R. Slater, Target-site resistance to neonicotinoids. *J. Chem. Biol. 2014 74* **7**, 125–128 (2014).

8. S. N. Caty, *et al.*, Molecular physiology of chemical defenses in a poison frog. *J. Exp. Biol.* **222** (2019).

9. F. Abderemane-Ali, *et al.*, Evidence that toxin resistance in poison birds and frogs is not rooted in sodium channel mutations and may rely on "toxin sponge" proteins. *J. Gen. Physiol.* **153** (2021).

10. R. D. Tarvin, *et al.*, Interacting amino acid replacements allow poison frogs to evolve epibatidine resistance. **357**, 1261-1266 (2017).

265   11.   B. L. Whitelaw, I. R. Cooke, J. Finn, K. Zenger, J. M. Strugnell, The evolution and origin
266   of tetrodotoxin acquisition in the blue-ringed octopus (genus Hapalochlaena). *Aquat.*
267   *Toxicol.* **206**, 114–122 (2019).

268   12.   S. Dobler, S. Dalla, V. Wagschal, A. A. Agrawal, Community-wide convergent evolution
269   in insect adaptation to toxic cardenolides by substitutions in the Na,K-ATPase. *Proc. Natl.*
270   *Acad. Sci. U. S. A.* **109**, 13040–13045 (2012).

271   13.   E. G. King, *et al.*, Genetic dissection of a model complex trait using the Drosophila
272   Synthetic Population Resource. *Genome Res.* **22**, 1558–1566 (2012).

273   14.   M. Chakraborty, J. J. Emerson, S. J. Macdonald, A. D. Long, Structural variants exhibit
274   widespread allelic heterogeneity and shape variation in complex traits. *Nat. Commun.* **10**,
275   1–11 (2019).

276   15.   T. N. Marriage, E. G. King, A. D. Long, S. J. Macdonald, Fine-mapping nicotine
277   resistance loci in *Drosophila* using a multiparent advanced generation inter-cross
278   population. *Genetics* **198**, 45–57 (2014).

279   16.   S. J. Macdonald, C. A. Highfill, A naturally-occurring 22-bp coding deletion in Ugt86Dd
280   reduces nicotine resistance in *Drosophila melanogaster*. *BMC Res. Notes* **13**, 1–6 (2020).

281   17.   R. M. Rizki, T. M. Rizki, Selective destruction of a host blood cell type by a parasitoid
282   wasp. *Proc. Nati. Acad. Sci. USA* **81**, 6154-6158 (1984).

283   18.   C. A. Highfill, *et al.*, Naturally segregating variation at *Ugt86Dd* contributes to nicotine
284   resistance in *Drosophila melanogaster*. *Genetics* **207**, 311–325 (2017).

285   19.   M. J. Snyder, J. I. Glendinning, Causal connection between detoxification enzyme activity
286   and consumption of a toxic plant compound. *J. Comp. Physiol. A* **179**, 255-261 (1996).

287   20.   L. Salazar-Jaramillo, B. Wertheim, Does Drosophila sechellia escape parasitoid attack by
288   feeding on a toxic resource? *PeerJ* **9** (2021).

289   21.   N. F. Milan, B. Z. Kacsoh, T. A. Schlenke, Alcohol consumption as self-medication
290   against blood-borne parasites in the fruit fly. *Curr. Biol.* **22**, 488–493 (2012).

291   22.   J. C. Fogleman, Response of Drosophila melanogaster to selection for P450-mediated
292   resistance to isoquinoline alkaloids. *Chem. Biol. Interact.* **125**, 93–105 (2000).

293   23.   P. T. Humphrey, *et al.*, Aversion and attraction to harmful plant secondary compounds
294   jointly shape the foraging ecology of a specialist herbivore. *Ecol. Evol.* **6**, 3256–3268
295   (2016).

296   24.   S. C. Groen, *et al.*, Multidrug transporters and organic anion transporting polypeptides
297   protect insects against the toxic effects of cardenolides. *Insect Biochem. Mol. Biol.* **81**, 51–
298   61 (2017).

299   25.   C. T. Hanifin, The chemical and evolutionary ccology of tetrodotoxin (TTX) toxicity in
300   terrestrial vertebrates. *Mar. Drugs* **8**, 577–593 (2010).

301   26.   G. Smagghe, L. Tirry, Insect midgut as a site for insecticide detoxification and resistance.
302   *Biochem. Sites Insectic. Action Resist.*, 293–321 (2001).

303   27.   M. A. Najarro, *et al.*, Identifying loci contributing to natural variation in xenobiotic
304   resistance in *Drosophila*. *PLOS Genet.* **11**, e1005663 (2015).

305   28.   R. W. Fitch, T. F. Spande, H. M. Garraffo, H. J. C. Yeh, J. W. Daly, Phantasmidine: an
306   Epibatidine congener from the Ecuadorian poison frog *Epipedobates anthonyi*. *J. Nat.*
307   *Prod.* **73**, 331 (2010).

308   29.   R Core Team. 2021. R: A language and environment for statistical computing. R
309   Foundation for Statistical Computing, Vienna, Austria

310   30.   W. N. Venables, B. D. Ripley, Introduction. 1–12 (2002).

311   31.   D. Bates, M. Mächler, B. Bolker, S. Walker, Fitting linear mixed-effects models using
312   lme4. *J. Stat. Softw.* **67**, 1–48 (2015).