Automated identification of piglet brain tissue from MRI images using Region-Based Convolutional Neural Networks

Kayla L. Stanke[1], Ryan J. Larsen[1,*], Laurie Rund[1], Brian J. Leyshon[2], Allison Y. Louie[3], Andrew J. Steelman[1,3,4,5]

[1]Department of Animal Sciences, University of Illinois Urbana-Champaign

[2]Abbott Nutrition, Discovery Research, Columbus, Ohio

[3]Division of Nutritional Sciences, University of Illinois Urbana-Champaign

[4]Neuroscience Program, University of Illinois Urbana-Champaign

[5]Carl R. Woese Institute for Genomic Biology, University of Illinois Urbana-Champaign

*Corresponding author

Email: larsen@illinois.edu

Abstract

Magnetic Resonance Imaging is an important tool for characterizing volumetric changes of the piglet brain during development. Typically, an early step of an imaging analysis pipeline is brain extraction, or skull stripping. Brain extractions are usually performed manually; however, this approach is time-intensive and can lead to variation between brain extractions when multiple raters are used. Automated brain extractions are important for reducing the time required for analyses and improving the uniformity of the extractions. Here we demonstrate the use of Mask R-CNN, a Region-based Convolutional Neural Network (R-CNN), for automated brain extractions of piglet brains. We validate our approach using Nested Cross-Validation on six sets of training/validation data drawn from 32 pigs. Visual inspection of the extractions shows acceptable accuracy, and

25 Dice coefficients are in the range of 0.95-0.97. These results demonstrate that R-CNNs provide a

26 viable tool for skull stripping of piglet brains.

Introduction

Piglets are an important translational model for measuring the effect of nutrition on brain

development. Not only do piglet brain stages of development correlate with human infant

development, but their nutritional requirements are also comparable (1). Magnetic resonance

imaging (MRI) is an important technique for obtaining non-invasive measurements of brain

volumes. An early step in volumetric analysis is the identification or "extraction" of the brain from

the surrounding tissue. Manual tracing has been the gold standard for brain extraction and is

performed by creating an outline that separates the surrounding skull, muscles, tissues, and fat

from the brain (1-6). However, the method is not ideal when working with large data sets because

it is time intensive and is subject to inconsistencies between raters/evaluators. Automated brain

extraction techniques are needed to overcome these limitations. However, reports of automated

extractions of pig brains are limited. A graph theory approach that makes use of prior information

of anatomical structures has been used to perform automated extraction of piglet brains (7).

However, deep learning technologies offer the possibility of automating the training of anatomical

structures to enable brain extractions (8-10). A U-Net model trained on humans, non-human

primates, and 3 pig scans has been shown to successfully perform brain extractions on 2 pig scans

(11). An alternative segmentation tool is Mask R-CNN, a Region-based Convolutional Neural

Network (R-CNN). This tool has been used to create a mouse brain atlas that is generalizable

across developmental ages and imaging modalities (12). These results suggest that Mask R-CNN

may be effective for piglet brain extraction. Here we demonstrate the use of Mask R-CNN for

automated brain extractions of piglet brains.

Methods

Animals and Care Practices

All animal care and handling procedures were approved by the University of Illinois Institutional

Animal Care and Use Committee (Protocol #18256) and were in accordance with federal

guidelines. Male and female average-for-gestational age, Yorkshire crossbred, full-term, naturally-

delivered piglets were obtained from University of Illinois Swine Farm at postnatal day 2 to allow

for colostrum consumption. All piglets remained intact but did undergo routine processing on the

farm including iron dextran (Henry Schein Animal Health, Dublin, OH, USA) and antibiotic

injection (EXCEDE®, Zoetis, Parsippany, NJ 07054, USA) per routine farm practice and

according to label. Four groups of piglets were placed individually into a caging system under

standard conditions as described in a previous publication (13) and randomly assigned to five diet

treatment groups. The control group of piglets remained with the sow until day 28 of age with free

access to suckle and were weighed daily. The current study does not distinguish between diet

groups.

Sow-raised piglets were handled daily, as were the artificially raised animals to diminish

differences between the two types of rearing. After arrival at the Edward R. Madigan Laboratory

(ERML) Animal Facility, experimental piglets received one dose of antibiotic: the first of two

cohorts received Spectraguard^{TM} (Bimeda, Inc, Oakbrook Terrace, IL 60181) on postnatal day 2

and the second cohort received Baytril® (Bayer healthcare LLC Shawnee Mission, KS 66201) on

postnatal day 4. Additional doses of antibiotic were administered during the experiment only under

the direction of the staff veterinarian. Animals were individually housed in racks of metabolism

cages specifically designed to artificially rear neonatal piglets under constant 12-h light/dark

cycles. Piglet housing at ERML was as follows: space allowance for individual piglets was 30"

deep, 23" wide, and 18.5" high, providing 4.8 square feet of floor space per piglet. Each piglet was

supplied with a heating pad, toy, and blanket. Room temperatures were kept at 85-95 ºF using

space heaters. Animals were also offered water or BlueLite® electrolyte solution ad libitum.

Cages and heating pads were disinfected, and toys and blankets replaced daily. Animal care

protocols were in accordance with National Institutes of Health Guidelines for Care and Use of

Laboratory Animals and were approved by the University of Illinois Laboratory Animal Care and

Use Committee.

MRI Acquisition

MRI data was acquired using 3 T Prisma scanner (Siemens, Erlangen) housed at the Biomedical

Imaging Center at the University of Illinois. Pigs were anesthetized using TKX (combination of

2.5 ml of xylazine (100 mg/ml) and 2.5 ml of ketamine (100mg/ml) added to a Telazol vial and

administered at a dosage of 0.02-0.03 ml/kg IM) then maintained on isofluorane (1-3%) throughout

the imaging. Animals were scanned in the supine position using a specialized piglet head coil

(Rapid Biomed, Rimpar). During scanning, the respiration rate, heart rate and blood oxygen levels

were monitored using a LifeWindow LW9x monitor (Digicare, Boynton Beach, FL).

MRI Structural Imaging: Our structural MRI scan consisted of a 3D MPRAGE acquisition (voxel

size = 0.6 x 0.6 x 0.6 mm^3, FOV = 173 x 173 mm^2, 256 slices, GRAPPA - GeneRalized

Autocalibrating Partial Parallel Acquisition - acceleration factor R = 2; TR = 2060 ms, TI = 1060

ms, flip angle = 9º, for an overall scan time of 5:21 min).

Manual brain extraction

Manual brain extraction was facilitated by first performing a rigid alignment of T_1-weighted images from all piglets to the brain atlas (14). This was done using SPM12 imaging analysis software. First, we performed a manual rotation to approximately align the T_1-weighted images with 28-day piglet template (14), without resampling. We then used the "coregistration" function of SPM12 to further align the T_1-weighted images to the template, again without resampling. The resulting alignment was accurate for all piglets, even though it was performed without first performing brain extraction, as shown by a representative image in Figure 1.

Figure 1. Results of coregistration of piglet brain to the average brain template. The top row shows images from a representative piglet brain and bottom row shows the same slices from the average brain template. Blue lines within each image indicate the locations of the perpendicular

slices.   The consistency of two sets of images confirms a good approximate alignment of the T₁-

weighted image with the average brain template.

An atlas-based brain mask was then resampled into the native space of each piglet, providing initial

estimates of the brain masks for each piglet. These initial estimates were then modified to create

individualized brain masks using Slicer3D, as shown in Figure 2. Most revisions were done in the

sagittal plane, but all three orthogonal views were reviewed and modified for improved precision.

All extractions were performed by one rater to minimize variability.

Figure 2.   Demonstration of the creation of manually-defined brain mask via modification of

the template brain mask. A representative T₁-weighted image is overlayed with the template

brain mask, creating a brighter region. The green outline shows the manual-designated brain

mask. The manual brain mask was created by editing the template brain mask for each

individual piglet.

Automated brain extraction

This study uses a deep learning instance segmentation neural network using object detection model

Mask R-CNN with Inception Resnet v2 architecture (15), pretrained on the COCO 2017 data set

(16). Our model uses a Tensor Flow 2.4.0 implementation of Mask R-CNN (15), with feature

extractor Faster R-CNN (17). Faster R-CNN utilizes a Region Proposal Network (RPN) to select

object proposals from a backbone, which for our study was generated using a combined ResNet101

and Feature Pyramid Network (FPN) (18).

Training was performed using a single NVIDIA GeForce GTX 1070 Ti GPU, with NVIDIA

developer driver 465.21, CUDA 11.0, and CUDNN library 8.0.4. The network was trained with a

cosine decay learning rate of 0.008 and a momentum optimizer value of 0.9, and a batch size of

two images, or slices, per iteration. We performed 200,000 iterations, or 48.8 epochs. The training

and segmentation were performed only in 2D sagittal planes. During evaluation, predicted masks

were binarized at a confidence parameter threshold of 0.5. The masks created from the 2D slices

were then combined into 3D datasets for final cleaning. This involved removing several small and

spurious masks, typically occurring in slices that did not include brain. Cleaning was done by first

using the Matlab function "bwconncomp" to identify all isolated masks, consisting of one or more

connected voxels, where connected voxels are defined as those with touching faces. Then we

discarded all by the largest mask, or brain mask.

Validation

Nested Cross-Validation was used to evaluate the performance of the training models. This method

has been shown to produce unbiased performance estimates, even in small datasets (19). We

randomly assigned each of the 32 pigs to one of six test groups. Four of the test groups consisted

of five pigs, and two of the test groups consisted of six pigs. For each test group, a training model

was generated using the remaining pigs, beginning with the same architecture which was naïve to

the test images. Validation of the test groups was performed by comparing the machine-generated

masks with the manually generated masks by visual inspection, by computing Dice coefficients

for each pig, and by computing Pearson correlations between the manual and machine-generated

masks. Dice coefficients were calculated using the formula $2|X \cap Y|/(|X| + |Y|)$, where $| \dots |$

indicates the number of voxels within the mask, $\cap$ indicates the union, and X and Y indicate the

manual and machine-generated masks.

Results

Visual inspection of the brain extractions reveals good accuracy of automatic brain extractions

(Figure 3). The six models were labelled with letters from A to F. We found that Model D failed

to identify the brain within several sagittal slices of one of the test pigs, as shown in Figure 4(a)

and (b). These slices included an unusually bright region in the subcutaneous fat layer near the

superior area of the head (see Figure 4(b)). This bright region was removed by manually outlining

it in one of the slices, and then removing the traced voxels from all the slices (see Figure 4(d)).

The modified structural images were then re-evaluated using the same model, producing a more

accurate brain mask (see Figure 4(c) and (d)). This brain mask was used for subsequent validation.

Dice coefficients from the 32 test scans were in the range of 0.95-0.97 (mean: 0.961, standard

deviation: 0.0036, see histogram in Figure 5). The Pearson correlation coefficient, R, of the

volumes, V_{manual} and $V_{machine}$, of the manual and machine-generated masks was R=0.90, with

p<0.001 (see Figure 6).

Figure 3. Sample extractions from two piglets. The overlay that creates a brighter region indicates machine extractions, and the green outline shows the manual brain masks. The top row shows a piglet tested using Model A, and the bottom row shows a piglet tested using Model B.

Figure 4. A segmentation error generated by Model D that was subsequently corrected. Panels

(a) and (b) show the inaccurate mask, indicated by green lines, and panels (c) and (d) show the

mask generated by the same model after image modification, with the same views and the same

slices. Axial slices are shown in (a) and (c); sagittal slices are shown in (b) and (d). The yellow

lines in the axial views, (a) and (c), show the location of the sagittal view and the red lines in

the sagittal views, (b) and (d), show the location of the axial views. For the inaccurate mask

there are several sagittal slices in which no brain regions were identified, as shown in (a). These

slices included a bright region in the fat layer of the superior region of the head, as shown in

(b). The bright region was manually traced and removed from the same voxels of all sagittal

173   *slices as shown in (d). A re-evaluation of the edited images using the same model, Model D,*

174   *produced an accurate brain extraction, as shown in (c) and (d).*

176   *Figure 5. Histogram of Dice coefficients calculated from each of the 32 pigs.*

Figure 6. The consistency of brain volumes, V_{manual} and $V_{machine}$, from manual and machine brain extractions, respectively. The solid line indicates unity and the colors denote the models used. Brain volume units are cubic centimeters.

Discussion

We have shown that Mask R-CNN trained on manually generated masks can be used to perform accurate piglet brain extractions. However, for one of the 32 test cases, a model failed to identify brain tissue within several slices. This problem was eliminated when the evaluation was repeated after the removal of an unusually bright region of subcutaneous fat was removed from the images. It is possible that that problem could have been avoided by using more training data, or by

performing a bias field correction before training. This example underscores the importance of

visual checks of machine performance.

Dice coefficients were >0.95; these are similar to values that have been achieved by neural

networks for the skull-stripping of non-human primates (11) and rodents (20).

The use of automated brain extraction has the potential to reduce analysis time because it requires

minimal supervision. This process is scalable to a high number of piglets, avoiding complications

and inconsistencies that might arise from having multiple raters perform manual brain extractions.

Improvements to our approach could be implemented in a variety of ways. Training with a larger

sample size is expected to increase performance and accuracy of machine learning algorithms (21).

Performance may also be improved by hyperparameter tuning and optimization (22, 23). Improved

performance might have been obtained by using 3D Mask R-CNN; however we employed 2D

Mask R-CNN to obtain shorter training times (24). Performance may also be improved by using

quantitative imaging techniques, such as MT saturation (25, 26), to improve the contrast between

brain and non-brain tissues. Further research is required to access whether network architectures

such as U-Net (27), may improve upon results obtained with Mask R-CNN.

The effectiveness of Mask R-CNN for performing piglet brain extractions implies that it may be a

useful tool for segmenting sub-regions of the brain. Further research is needed to assess whether

such an approach may compliment or improve upon existing methods for volumetric analysis of

piglet brain MRI data (4, 28, 29).

Acknowledgments

209   The study was funded by Abbott Nutrition, www.abbott.com, grant number 00490326 to A.S. The

210   funder aided with study design, data analysis, decision to publish, and preparation of the

manuscript. We thank the staff members of the Biomedical Imaging Center at the Beckman Institute for Advanced Science and Technology for imaging support.

Author contributions

KLS: Conceptualization, Data Curation, Formal Analysis, Investigation, Methodology, Software, Validation, Visualization, Writing – Original Draft Preparation, Writing-Review & Editing **RJL:** Conceptualization, Data Curation, Formal Analysis, Funding Acquisition, Investigation, Methodology, Project Administration, Resources, Software, Supervision, Validation, Visualization, Writing – Original Draft Preparation, Writing-Review & Editing **AL:** Investigation, Resources **BJL:** Conceptualization, Funding Acquisition, Investigation, Project Administration, Supervision, Writing-Review & Editing **LR:** Funding Acquisition, Investigation, Project Administration, Resources, Supervision **AS:** Funding Acquisition, Supervision, Writing-Review & Editing

Data Sharing

Processing scripts are available at https://github.com/kaylals2/Piglet_RCNN.

References

1. Mudd AT, Dilger RN. Early-Life Nutrition and Neurodevelopment: Use of the Piglet as a Translational Model. Advances in Nutrition. 2017;8(1):92-104.
2. Bokde AL, Teipel SJ, Schwarz R, Leinsinger G, Buerger K, Moeller T, et al. Reliable manual segmentation of the frontal, parietal, temporal, and occipital lobes on magnetic resonance images of healthy subjects. Brain Res Brain Res Protoc. 2005;14(3):135-45.
3. Conrad MS, Dilger RN, Nickolls A, Johnson RW. Magnetic resonance imaging of the neonatal piglet brain. Pediatr Res. 2012;71(2):179-84.
4. Gan H, Zhang Q, Zhang H, Chen Y, Lin J, Kang T, et al. Development of new population-averaged standard templates for spatial normalization and segmentation of MR images for postnatal piglet brains. Magn Reson Imaging. 2014;32(10):1396-402.
5. Mudd AT, Alexander LS, Berding K, Waworuntu RV, Berg BM, Donovan SM, et al. Dietary Prebiotics, Milk Fat Globule Membrane, and Lactoferrin Affects Structural Neurodevelopment in the Young Piglet. Front Pediatr. 2016;4:4.
6. Leyshon BJ, Radlowski EC, Mudd AT, Steelman AJ, Johnson RW. Postnatal Iron Deficiency Alters Brain Development in Piglets. J Nutr. 2016;146(7):1420-7.

7.   Durandeau AF, J.-B.; Bloch, I.; Mazerand, E.; Menei, P.; Montero-Menei, C.; Dinomais, M. , editor
Structural information and (hyper)graph matching for MRI piglet brain extraction. International
Conference on Pattern Recognition Systems; 2019 8-10 July 2019; Tours, France.
8.   Akkus Z, Galimzianova A, Hoogi A, Rubin DL, Erickson BJ. Deep Learning for Brain MRI
Segmentation: State of the Art and Future Directions. J Digit Imaging. 2017;30(4):449-59.
9.   Hurtz S, Chow N, Watson AE, Somme JH, Goukasian N, Hwang KS, et al. Automated and manual
hippocampal segmentation techniques: Comparison of results, reproducibility and clinical applicability.
Neuroimage Clin. 2019;21:101574.
10.   Iqbal A, Khan R, Karayannis T. Developing a brain atlas through deep learning. Nature Machine
Intelligence 2019;1:277-87.
11.   Wang X, Li XH, Cho JW, Russ BE, Rajamani N, Omelchenko A, et al. U-net model for brain
extraction: Trained on humans for transfer to non-human primates. Neuroimage. 2021;235:118001.
12.   Iqbal A, Khan R, Karayannis T. Developing a brain atlas through deep learning. Nat Mach Intell.
2019;1(6):277-87.
13.   Rytych JL, Elmore MR, Burton MD, Conrad MS, Donovan SM, Dilger RN, et al. Early life iron
deficiency impairs spatial cognition in neonatal piglets. J Nutr. 2012;142(11):2050-6.
14.   Conrad MS, Sutton BP, Dilger RN, Johnson RW. An in vivo three-dimensional magnetic
resonance imaging-based averaged brain collection of the neonatal piglet (Sus scrofa). PLoS One.
2014;9(9):e107650.
15.   He KM, Gkioxari G, Dollar P, Girshick R. Mask R-CNN. Ieee I Conf Comp Vis. 2017:2980-8.
16.   Lin T-Y, Maire M, Belongie S, Hays J, Perona P, Ramanan D, et al., editors. Microsoft COCO:
Common Objects in Context2014; Cham: Springer International Publishing.
17.   Ren SQ, He KM, Girshick R, Sun J. Faster R-CNN: Towards Real-Time Object Detection with
Region Proposal Networks. Ieee T Pattern Anal. 2017;39(6):1137-49.
18.   Lin TY, Dollar P, Girshick R, He KM, Hariharan B, Belongie S. Feature Pyramid Networks for
Object Detection. Proc Cvpr Ieee. 2017:936-44.
19.   Vabalas A, Gowen E, Poliakoff E, Casson AJ. Machine learning algorithm validation with a limited
sample size. PLoS One. 2019;14(11):e0224365.
20.   Hsu LM, Wang S, Ranadive P, Ban W, Chao THH, Song S, et al. Automatic Skull Stripping of Rat
and Mouse Brain MRI Data Using U-Net. Front Neurosci-Switz. 2020;14.
21.   Jollans L, Boyle R, Artiges E, Banaschewski T, Desrivieres S, Grigis A, et al. Quantifying
performance of machine learning methods for neuroimaging data. Neuroimage. 2019;199:351-65.
22.   Yang L, Shami A. On hyperparameter optimization of machine learning algorithms: Theory and
practice. Neurocomputing. 2020;415:295-316.
23.   Wu J, Chen X-Y, Zhang H, Xiong L-D, Lei H, Deng S-H. Hyperparameter Optimization for Machine
Learning Models Based on Bayesian Optimizationb. Journal of Electronic Science and Technology.
2019;17(1):26-40.
24.   Jeong J, Lei Y, Kahn S, Liu T, Curran WJ, Shu HK, et al. Brain tumor segmentation using 3D Mask
R-CNN for dynamic susceptibility contrast enhanced perfusion imaging. Phys Med Biol. 2020;65(18).
25.   Helms G, Draganski B, Frackowiak R, Ashburner J, Weiskopf N. Improved segmentation of deep
brain grey matter structures using magnetization transfer (MT) parameter maps. Neuroimage.
2009;47(1):194-8.
26.   Helms G, Dathe H, Kallenberg K, Dechent P. High-resolution maps of magnetization transfer with
inherent correction for RF inhomogeneity and T1 relaxation obtained from 3D FLASH MRI. Magn Reson
Med. 2008;60(6):1396-407.
27.   Ronneberger O, Fischer P, Brox T, editors. U-Net: Convolutional Networks for Biomedical Image
Segmentation2015; Cham: Springer International Publishing.
28.   Ashburner J, Friston KJ. Unified segmentation. Neuroimage. 2005;26(3):839-51.

29.   Radlowski EC, Conrad MS, Lezmi S, Dilger RN, Sutton B, Larsen R, et al. A neonatal piglet model
for investigating brain and cognitive development in small for gestational age human infants. PLoS One.
2014;9(3):e91951.

293