# Genomic events shaping epithelial-to-mesenchymal trajectories in cancer

Guidantonio Malagoli Tagliazucchi[1], Maria Secrier[1,*]

[1] UCL Genetics Institute, Department of Genetics, Evolution and Environment, University College London, UK

*To whom correspondence should be addressed (m.secrier@ucl.ac.uk)

## ABSTRACT

The epithelial to mesenchymal transition (EMT) is a key cellular process underlying cancer progression, with multiple intermediate states whose molecular hallmarks remain poorly characterized. To fill this gap, we explored EMT trajectories in 8,778 tumours of epithelial origin and identified three macro-states with prognostic and therapeutic value, attributable to epithelial, hybrid E/M (hEMT) and mesenchymal phenotypes. We show that the hEMT state is remarkably stable and linked with increased aneuploidy, APOBEC mutagenesis and hypoxia. Additionally, we provide an extensive catalogue of genomic events underlying distinct evolutionary constraints on EMT transformation, including novel pan-cancer dependencies of hEMT on driver genes *PRRX1, BCOR* and *CNOT3*, as well as links between full mesenchymal transformation and *REG3A* and *SHISA4* mutations in lung and breast cancers, respectively. This study sheds light on the aetiology of the lesser characterised hybrid E/M state in cancer progression and the broader genomic hallmarks shaping the mesenchymal transformation of primary tumours.

**INTRODUCTION**

The epithelial to mesenchymal transition (EMT) is a cellular process in which polarized

epithelial cells undergo multiple molecular and biochemical changes and lose their identity in

order to acquire a mesenchymal phenotype[1]. EMT occurs during normal embryonic

development, tissue regeneration, wound healing, but also in the context of disease[1,2]. In

cancer, it promotes tumour progression with metastatic expansion[3]. Recent studies have

uncovered that EMT is not a binary switch but rather a continuum of phenotypes, whereby

multiple hybrid EMT states underly and drive the transition from fully epithelial to fully

mesenchymal transformation[4,5].

Elucidating the evolutionary trajectories that cells take to progress through these states is key

to understanding metastatic spread and predicting cancer evolution.

The transcriptional changes accompanying EMT in cancer have been widely characterised

and are governed by several transcription factors, including Snail, Slug, Twist and zinc fingers

*ZEB1* and *ZEB2*[6,7,8]. EMT appears driven by waves of gene regulation underpinned by

checkpoints, such as *KRAS* signalling driving the exit from an epithelial state, dependent upon

*EGFR* and *MET* activation[9].

However, EMT progression is not only characterized by transcriptional alterations of

regulatory circuits; the genetic background of the cell can also impact its capacity to undergo

this transformation. Gain or loss of function mutations in a variety of genes, including *KRAS*[10],

*BRCA1*[11], *STAG2*[12], *TP53*[13,14], as well as amplifications of chromosomes 5, 7 and 13 have

been shown to promote EMT[15]. Several pan-cancer studies have also linked copy number

alterations, miRNAs and immune checkpoints with EMT on a broader level[16,17]. Mathematical

models have been developed to describe the switches between epithelial and mesenchymal

states[4] but without considering any genomic dependencies.

Despite extensive efforts to study the dynamics of EMT, some aspects of this process remain

poorly characterized. In particular, most of the studies mentioned considered EMT as a binary

switch and failed to capture evolutionary constraints that may change along the continuum of

EMT transformation. Single cell matched DNA- and RNA-seq datasets would ideally be

needed for this purpose, but they are scarce. We hypothesised that a pan-cancer survey of

EMT phenotypes across bulk sequenced samples should capture a broad spectrum of the

phenotypic variation one may expect to observe at single cell level, and this could be linked

with genomic changes accompanying EMT transformation. To explore this, we have

integrated data from the Cancer Genome Atlas (TCGA), MET500[18], MetMap[19], GENIE[20],

MSK-IMPACT[21], GDSC[22] and POG570[23] datasets to characterise and validate EMT and

linked genetic changes in cancer. By mapping 8,778 tumours of epithelial origin onto a

"timeline" of epithelial-to-mesenchymal transformation, we identified discrete EMT macro-

states and derived a catalogue of genomic hallmarks underlying evolutionary constraints of

these states. These genomic events shed light into the aetiology of the lesser characterised

hybrid E/M phenotype and could potentially act as early biomarkers of invasive cancer.

**RESULTS**

**Pan-cancer reconstruction of EMT trajectories and states in bulk tumours**

To explore the EMT process within bulk tumour samples, we employed a cohort of primary

tumours of epithelial origin (n = 8,778) spanning 25 cancer types from TCGA. The bulk RNA-

seq data from these tumours underlie multiple transcriptional programmes reflecting different

biological processes, including EMT. Inspired by McFaline-Figueroa et al[9], we quantified the

levels of EMT in these bulk tumours against a reference single cell RNA sequencing (scRNA-

seq) dataset derived from MCF10 breast cancer cells that have been profiled at different times

during the epithelial to mesenchymal transition *in vitro*. These data allowed us to reconstruct a

generic "pseudo-timeline" of spontaneous EMT transformation onto which we projected the

bulk sequenced samples from TCGA, positioning them within the continuum of EMT states

(Figure 1a). Since the bulk signal is arising from a mixture of cells, some of which would be

found in different stages of mesenchymal transformation, the overall signal would reflect the

average EMT state across the entire tumour cell population.

Using this approach, we reconstructed the EMT pseudotime trajectory across multiple cancer

tissues (Figure 1b, Supplementary Table S1). The expression of canonical epithelial and

mesenchymal markers was consistent with that observed in the scRNA-seq data and

expectations from the literature (Supplementary Figure S1a). Along the pseudotime, we

observed frequent co-expression of such markers, which could reflect a hybrid E/M state[24].

To characterise the dominant EMT states governing the continuum of transcriptional activity

described above, we discretised the pseudotime trajectory based on expression values of

canonical EMT markers using a Hidden Markov Model approach and uncovered three macro-

states: epithelial (EPI), hybrid EMT (hEMT) and mesenchymal (MES) (Figure 1b-c,

Supplementary Figure 1b). These states were robust to varying levels of gene expression

noise (Supplementary Figure S1c-d). As expected, the probability for the cancer cells to

switch from the epithelial to the hEMT (0.35) state was higher than the probability to passage

directly into the mesenchymal state (0.14). The hEMT tumours tended to remain in the same

state 44% of the times, suggesting this state could be more stable than anticipated – as

previously stipulated[25] and consistent with observations that a fully mesenchymal state is not

always reached[26].

The EMT scores progressively increased between the EPI, hEMT and MES states, as

expected (Figure 1d). Reassuringly, in an independent cohort of metastatic samples

(MET500), EMT levels were relatively elevated along the transformation timeline compared to

TCGA samples and were most abundantly falling within the hEMT state (Figure 1d,

Supplementary Figure S1e). Interestingly, we also observed possible cases of a reversion to

an epithelial state in metastatic samples, which is to be expected when colonizing a new

environmental niche.

We also applied our EMT scoring methodology to the MetMap resource, which has

catalogued the metastatic potential of 500 cancer cell lines across 21 cancer types. The

invasion potential of these cell lines increased along the pseudotime axis as expected (Figure

1e, Supplementary Figure S1f), Cell lines classified as MES by our HMM model were

predominantly metastatic, while hEMT cases had more varied invasion potential.

At tissue level, the proportion of samples in each EMT state was variable (Figure 1f,

Supplementary Figure S3g), with hEMT dominating in head and neck, oesophageal and

pancreatic carcinomas, while adenoid cystic, kidney carcinomas and melanomas were highly

mesenchymal.

As expected, the EMT classification was significantly correlated with the clinical cancer stage

(Chi-square test $p \ll 0.0001$), with transformed samples (hEMT/MES) found to be 1.6-fold and

3.3-fold enriched in late-stage tumours, respectively, while the epithelial state was 3-fold

overrepresented in early-stage cancers (Figure S1h). Overall, 18% of the profiled samples

were classified as fully transformed (MES) and only 8% of these were annotated as late-stage

tumours (Supplementary Table S2). Therefore, our method is able to detect an additional 10%

of cases presenting early evidence for the phenotypic transformation required for metastasis.

Indeed, multiple studies have demonstrated the activation of the EMT transcriptional

programme in the early stages of cancer[10,27]. Even the hEMT phenotype is hypothesised to be

sufficient for promoting metastatic dissemination[28], although our analysis of the MetMap

dataset suggests this may be tissue/context-specific.

**Tumour cell extrinsic hallmarks of EMT**

Multiple microenvironmental factors, including tumour associated macrophages, secreted

molecules (IL-1, TNF-α) or hypoxia, have been extensively described to promote EMT[29,30,31,32].

However, their macro-state specificity is less well characterised. We confirmed that

transformed samples, particularly hEMT ones, exhibited higher infiltration by non-cancer cells

(Supplementary Figure S2a-c). Endothelial, cytotoxic and γδ T cells were progressively

enriched with increased stages of EMT transformation (Figure 2a-b, Supplementary Figure

S2d-e), suggesting that the fully mesenchymal state is most often linked with "immune hot"

tumours. In line with this hypothesis, these tumours also showed the highest exhaustion levels

(Figure 2c). In contrast, the hEMT samples displayed the highest enrichment of cancer-

associated fibroblasts (Figures 2a-b, d-e,  Supplementary Figure S2f and Supplementary

Table S3), confirming previous reports[33]. To avoid confounding effects between fibroblast and

hEMT markers as highlighted by Tyler and Tirosh[34], we excluded EMT markers from the

employed fibroblast gene sets, but we acknowledge that part of the signal recovered may still

not be unambiguously attributed to either the cancer or microenvironmental component.

Additionally, samples with a transformed phenotype (MES, hEMT) presented significantly

elevated hypoxia levels (Figure 2f, Supplementary Figure S2g). Hypoxia has previously been

shown to promote EMT by modulating stemness properties[35]. We found that CD44, an

established cancer stem cell marker known to promote EMT[36,37], was most highly expressed

in the hEMT state across cancers (Figure 2g), and elevated levels of several other stemness

signatures most often accompanied the hEMT and MES macro-states (Figure S2h-i).  Unlike

mesenchymal samples, the majority of hEMT tumours (35%) were characterized by both

hypoxia and CD44 expression (Supplementary Table S4). Thus, the interplay between

hypoxia and stemness may play a greater role in attaining the hEMT state compared to the

fixation of a fully mesenchymal phenotype.

**Tumour cell intrinsic hallmarks of EMT**

In addition to environmental factors, intrinsic cell properties such as increased proliferation,

mutational burden and aneuploidy would be expected along the EMT transformation

trajectory. Across distinct tissues, these changes were most pronounced in the hEMT state

(Figure 2h). Interestingly, this group also presented higher levels of centrosome amplification,

which have been linked with increased genomic instability[38,39] and poor prognosis[40].

Such alterations to the genomic integrity of the cells result from multiple mutational processes.

These processes leave recognizable patterns in the genome termed "mutational signatures",

which in their simplest form constitute of trinucleotide substitutions and have been broadly

characterised across cancers[41]. However, their involvement in EMT transformation is poorly

understood. To investigate whether any neoplastic process introducing mutations in the

genomes was conditioned by EMT, we modelled the associations between mutational

signatures and EMT using linear mixed effects models while accounting for tissue effects

(Figure 2i-j, Supplementary Figure S3a, Supplementary Table S5). The ageing signature

SBS1 was significantly increased in non-mesenchymal tumours, while the APOBEC

mutagenesis signature SBS13 appeared specifically elevated in hEMT. It is worth noting that if

we did not account for the tissue effect, the signatures most strongly associated with EMT

progression and specifically increased in mesenchymal samples were SBS7a/b, linked with

UV light-induced damage (Supplementary Figure S3b-c). Samples exposed to these

carcinogens may progress to a mesenchymal phenotype earlier within the primary tumour, or

such mutations may be more easily fixed in transformed cells.

**Genomic driver events underlying the EMT transformation pan-cancer**

Beyond the broader hallmarks discussed above, we sought to identify specific genomic

changes creating a favourable environment for EMT transformation. We prioritised cancer

driver mutations, focal and arm-level copy number changes that may be linked with EMT, and

implemented a lasso-based machine learning framework to identify those drivers able to

discriminate between EPI, hEMT and MES states across cancers, while accounting for tissue-

specific effects (Methods, Figure 3a, Supplementary Table S6a). This model was validated

using several other machine learning approaches and demonstrated remarkably high

accuracies of 80-99% (Supplementary Figure S4a-n, Supplementary Table S6b).

Among the genomic biomarkers able to discriminate transformed tumours (hEMT, MES) from

the epithelial state (EPI), we identified well characterized driver genes such as *TP53*, *PIK3CA*,

*KRAS*, along with chromosomal arm alterations at 1q, 8q, 17q, 20q, some of which have been

previously linked with cell migration and invasion (Figure 3b, Supplementary Figure S4o,

Supplementary Table S6c). Interestingly, the fraction of cancer cells harbouring *KRAS*, *RB1*

and *FGFR2* mutations was markedly increased at the mesenchymal level, suggesting that a

clonal fixation of these events may be key for the establishment of a fully mesenchymal state

(Figure 3c-d). This is in line with findings on *KRAS* reported by McFaline-Figueroa et al[9], but

also pinpoints additional checkpoints.

The hEMT state-specific markers were mostly enriched in cell cycle, fate commitment and cell

adhesion pathways, hallmarks that are classically associated with cancer progression (Figure

3e, Supplementary Figure S4p). These alterations most often presented in a reduced cancer

cell fraction, concordant with subclonal diversification in line with phenotypic transformation

(Figure 3c). We identified several events that have not been previously linked with this

intermediate phenotype in the literature, including alterations of *PRRX1*, *BCOR*, *FAM135B*,

*CNOT3* and *ERCC3* (Figure 3e, Supplementary Figure S5a-b, Supplementary Table S6a).

*TCEA1* and *SPEN* alterations were specifically linked with high levels of aneuploidy,

stemness, centrosome amplification and hypoxia. While none of these events have been

specifically tied to an hEMT state before, *CNOT3* does not appear to have even been linked

more generally with invasion or metastasis. *CNOT3* is a translational repressor[42] upregulated

in non-small cell lung cancer[43].

Among the chromosomal arm-level events, amplifications of the 16q arm appeared to confer

the highest protective effect from transformation, with a >2-fold depletion in hEMT

(Supplementary Figure S5c). Another notable event was the deletion of the 5q arm,

associated with the EPI state. The trisomy of chromosome 5 has been previously linked with a

hybrid E/M state in a colon cancer cell line[15], hence structural changes of chromosome 5

might impose a selective pressure on EMT transformation.

The genomic markers distinguishing mesenchymal from epithelial samples were enriched in

oxidative stress and hypoxia response pathways (Figure 3f, Supplementary Figure S4q),

suggesting an evolutionary adaptation to mesenchymal-promoting changes in the

environment. All of the identified drivers have previously been linked to metastasis or EMT

transformation[44,45] (Supplementary Table S6b-c), e.g. *VHL* mutations via regulation of

hypoxia[46]. Increases in copy number amplification were generally associated with a more

mesenchymal state (Supplementary Figure S5d), as were deletions of the 6p chromosomal

arm which harbours 16 cancer drivers, several of which have been previously linked with EMT

(Supplementary Figure S5e, Supplementary Table S6a).

A good fraction of the reported alterations was independently validated in the MET500, GENIE

and MSK-IMPACT studies: 63% of point mutations and 43% of copy number events

distinguishing MES from EPI, as well as 58% of somatic mutations and 70% of the copy

number events discriminating hEMT from EPI (Figure 4a-d). Acknowledging the limitations of

these resources to metastatic samples (MET500) and targeted sequencing (GENIE, MSK-

IMPACT), our analysis was nevertheless able to recapitulate genomic events necessary for

EMT transformation that are preserved during metastatic progression, but also uncovered

some events that may not be further maintained in the new metastatic niche. Furthermore,

31% and 57% of these alterations were confirmed to be linked with the metastatic potential of

cancer cell lines, at pan-cancer and tissue specific level, respectively (Figure 4e-f). 48% of the

copy number events were validated in at least one cell-line, e.g. *COX6AC* amplifications in the

pancreas (Figure 4h). Interestingly, the impact on metastatic potential varied between

increases and decreases, which can partly be explained by the heterogeneity and complex

dependencies of cell lines, and partly by our incomplete understanding of the metastatic

potential of hEMT cells. Suppression of most of these genes strongly impacted cell viability

(Figure 4i-j) and were targets of transcription factors regulating EMT (Figure 4k-l).

Knockdowns of the hEMT-linked genes *PRRX1* and *CREBBP* were also linked with a weak

EMT phenotype in murine epithelial cells[47].

Finally, we confirmed downstream variations of the proteome accompanying EMT

transformation for the key EMT markers E-cadherin, N-cadherin, serpin E1, fibronectin

(Supplementary Figure S6a, Supplementary Table S6d). Proteins upregulated in hEMT

included components of the Hippo pathway (YAP,TAZ), regulating cell adhesion and

mechanical signals, with previously demonstrated roles in EMT transformation[48]

(Supplementary Figures S6b,d-e). In contrast, proteins differentially expressed in a fully

mesenchymal state (Supplementary Figure S6c) were linked with mTOR signaling, response

to oxygen levels, and UV damage response (Supplementary Figure S6f-g), consistent with our

previous findings linking the MES state with hypoxia and the UV mutational signature SBS7.

**Tissue-specific EMT trajectories and genomic dependencies**

EMT is a ubiquitous program, however, it is strongly influenced by tissue type and external

stimuli. Our pan-cancer reconstruction of the EMT pseudotime relied on data from single cells

captured during spontaneous EMT transformation. To further refine EMT trajectories in a

tissue-specific manner and to consider other sources of influence on the EMT transcriptional

programme, we used scRNA-seq data from two cancer cell lines (the human adenocarcinoma

alveolar basal epithelial cell line A549, and the breast cancer cell line MCF7) that were

stimulated with different exogenous molecules (TGF-β1, EGF, TNF) in a time-course

experiment[49]. As before, we mapped bulk lung and breast primary tumours from TCGA onto

the EMT pseudo-timeline derived from the scRNA-seq data and observed a finer granularity of

different activation states along the EMT trajectory in these cancers (Figure 5a-b,

Supplementary Figure S7a,d). In both cancers we identified 5 states, with multiple epithelial,

hEMT and mesenchymal-like phenotypes (Figure 5c-d).

We found evidence for positive selection in 17 genes across the 5 EMT stages in lung

adenocarcinoma, 9 of which had also been identified in the pan-cancer analysis (Figure 5e,

Supplementary Table S7a-e). Seven drivers (*RB1*, *MGA*, *ZIC1*, *NF1*, *REG3A*, *ARID2*,

*ZFP36L1*) were only enriched in the mesenchymal-like cluster M2. Among these, *REG3A*,

encoding for a secretory protein linked with inflammation-driven carcinogenesis and cell

migration in gastrointestinal cancers[50], has not been previously linked with EMT in lung cancer

and could potentially constitute a novel biomarker of mesenchymal transformation in this

cancer type, where it tends to be more highly expressed when mutated. Indeed, cell lines

carrying *REG3A* mutations show a strong increase in metastatic potential, particularly in lung

(Figure 5g-h). REG3A depletion does not impact cell viability in upper aerodigestive tissue

(Supplementary Figure S7g), making it a putative pharmacological target.

In breast cancer, the most mesenchymal state M was again linked with *NF1* mutations, but

also with alterations of *AKT1* and *SHISA4* (Figure 5f, Supplementary Table S7f-j). *AKT* has

been shown to affect epithelial cell morphology and motility[51], while the role of *SHISA4* in EMT

has not been characterised. Both genes are targets of transcription factors known to modulate

EMT (Supplementary Figure S7i).

Most of the focal copy number alteration events in both lung and breast cancers were distinct

from the pan-cancer analysis (Supplementary Figure S7b-f, Supplementary Table S7e,j), and

several of them have been previously associated with EMT, albeit not in a micro-state specific

manner. Amplifications of the non-essential *PRDM2* gene in breast cancer and deletions of

the non-essential gene *CUX1* in lung cancer, among others, were linked with metastatic

potential in cell lines and transcription factor regulation of EMT (Figure 5i, Supplementary

Figure S7h,j).

**Clinical relevance of EMT**

Finally, we show that the defined EMT states have potential clinical utility. Patients with an

hEMT phenotype had the worst overall survival outcomes (Figure 6a, Supplementary Table

S8a), while those with mesenchymal tumours showed significantly decreased progression-

free intervals (Figure 6b, Supplementary Table S8b) and disease-specific survival

(Supplementary Figure S8a, Supplementary Table S8c). As expected, patients who presented

transformed tumours and were lymph node positive had the worst prognosis (Supplementary

Figure S8b).

Among the driver events that have been linked with EMT in this study, mutations in genes

*CTNNB1* and *TSC2*, associated with a mesenchymal phenotype, and six other driver genes

associated with hEMT (*BCOR*, *RHOA*, *CDH10*, *SMAD2*, *SETD2*, *PTHCH1*) conferred worse

prognosis (Figure 6c-d, Supplementary Table S9).

Interestingly, the progression-free interval after oxaliplatin treatment was shorter in patients

with hEMT tumours (Supplementary Figure S8c), suggesting hEMT might be linked with poor

responses to this chemotherapy drug. To further explore potential links between EMT and

therapy responses, we investigated whether EMT progression might confer different levels of

sensitivity to individual cancer drugs using cell line data from GDSC[22]. We found 22

compounds whose IC50 values were significantly correlated with the EMT score (Figure 6e).

The strongest associations were observed with Acetalax, a drug used in the treatment of triple

negative breast cancers, and Sapitinib, an inhibitor of ErbB1/2/3[52]. Several hEMT and MES

biomarkers (e.g. *EP300*, *FAT1*, *NFE2L2*, *PTEN*) conferred increased sensitivity to multiple

drugs in a tissue specific manner (Supplementary Figure S8d-f), and suggested opportunities

for repurposing of non-oncology drugs (Supplementary Figure S8g-h).

Finally, we linked post-treatment EMT phenotypes with therapy responses using the POG570

dataset (Supplementary Figure S8i). The duration of treatment, used as a proxy for worsening

outcome, varied based on EMT state for the aromatase inhibitors exemestane and letrozole,

but also for the chemotherapeutics etoposide and temozolomide (Supplementary Figures S8j-

n).  Moreover, EMT potential was decreased upon chemotherapy treatment compared to

unmatched treatment-naïve cases in EPI and hEMT samples, while MES potential increased

(Figure 6f). There was a consistent drop in EMT potential in capecitabine and letrozole-treated

tumours (Figure 6g-h, Supplementary Figure S8o-p). Hence, the level of EMT transformation

may play a role in determining responses to a variety of cancer therapies.

**DISCUSSION**

Previous studies of the EMT process have suggested the existence of a phenotypic

continuum characterised by multiple intermediate states[53]. We have shown that distinct EMT

trajectories in cancer are underpinned by three macro-states, reflecting both tumour cell

intrinsic as well as tumour microenvironment associated changes. The hybrid E/M state,

characterised by the co-expression of epithelial and mesenchymal markers, was surprisingly

frequent (38%). It presented traits linked with increased neoplastic aggressiveness, such as a

tumour-promoting microenvironment and worse clinical outcomes, also in certain

chemotherapy contexts. Indeed, it has been reported that cells with hEMT features give rise to

daughter cells that are either mesenchymal or epithelial[54,36], are more prone to migrate and

promote the formation of circulating tumour cells[37]. Moreover, stable late hEMT states have

been linked with maximal metastatic potential and worse overall survival[55]. Undoubtedly, the

hEMT state can be further subdivided into sub-states, as shown by Goetz et al[4], Brown et al[56]

and also by us in the tissue-specific analyses. The true number of EMT intermediate states is

just beginning to be explored. However, the bulk sequencing data are limiting our ability to

capture them, and this will be best studied in single cell datasets.

Our study confirmed previously established molecular hallmarks of EMT, including increased

genomic instability and hypoxia in hEMT, and cytotoxicity/exhaustion in mesenchymal

tumours[17], which could inform immunotherapy strategies. We also highlighted mutational

processes that have increased activity in specific EMT states. Beyond an expected

association with ageing-induced damage along the EMT progression axis, we also found that

the signature SBS13, linked with the activity of the APOBEC3A/B cytidine deaminases, was

prevalent in hEMT tumours. In mammary epithelial cells, inflammatory signals promote the

upregulation of the activation-induced cytidine deaminase (AID) enzyme, a component of the

APOBEC family, and this has been shown to trigger EMT[57]. Furthermore, cyclic hypoxia has

recently been proven to induce APOBEC activity[58], which could further explain the

convergence of these phenotypes in the hEMT state.

While the exploration of EMT biomarkers is not new, most of the studies in this area have

been reliant on gene expression activity rather than mutational dependencies and they are

generally tissue-specific[17,30]. Pan-cancer studies generally consider EMT as a binary

switch[16,17,30]. In contrast, our study identified genomic hallmarks of three EMT macro-states

derived from the integration of bulk and single cell datasets, which provided the opportunity to

understand the evolutionary constraints on the lesser characterised hybrid E/M state. We

uncovered novel putative hEMT drivers like *PRRX1* and *BCOR,* and events in genes *CNOT3,*

*REG3A* and *SHISA4*, which have not been linked with hybrid/mesenchymal phenotypes or

metastatic expansion in the studied cancers. A causal relationship between the acquisition of

any of these genomic changes and EMT should be experimentally tested in the future. The

EMT state by itself, as well as several putative EMT biomarkers, were linked to drug

responses and could thus potentially be exploited for therapeutic benefit.

Overall, the results of this study demonstrate the complex intrinsic and microenvironmental

mechanisms that shape the landscape of EMT transformation during cancer. We have not

considered the role of chromosomal rearrangements or epigenetic changes in EMT, which

could provide further explanations to the maintenance of an hEMT phenotype. Additional

research is required to understand the biological role of the identified biomarkers, their

importance in a clinical setting, and to identify additional mechanisms that may promote EMT.

**METHODS**

**Data sources**

Bulk RNA-sequencing, copy number (segment file and focal alterations), somatic

variants(MuTect[59]) and clinical data were retrieved for 8,7778 primary tumours of epithelial

origin from the harmonized version of TCGA using the *TCGAbiolinks* R package[60]. All other

data sources employed for validation are described below.

**Reconstruction of EMT trajectories in bulk data**

The reconstruction of the EMT trajectory of the TCGA samples was performed using a

procedure that allows to map bulk samples to single cell-derived expression programmes

inspired from McFaline-Figueroa et al[9]. The workflow of the analysis consists of several steps.

The analysis requires two gene expression matrices as input, corresponding to one bulk

sequenced dataset and a single cell dataset. In our case, we used as input the bulk RNA-seq

data from TCGA samples and scRNA-seq of a spontaneous EMT model derived in MCF10

cell lines[9] with the associated trajectory (P). In the first step of the analysis the matrices were

merged; then, in order to remove the batch effects originated by the two different platforms, a

correction was applied using ComBat[61]. In the second step, principal component analysis

(PCA) was performed on the merged matrix. The MCF10 derived EMT trajectory was then

mapped onto the TCGA data using an iterative process and a mapping strategy based on k

nearest neighbours (kNN). The number of iterations (i) is equal to the number of TCGA

samples. During each i-th step of iteration, a single TCGA sample and the MCF10 scRNA-seq

data were used as input for the kNN algorithm. The procedure computed the mean of the

pseudotime values of the single cell samples that have been detected by the kNN algorithm to

be associated with the i-th TCGA sample. The implementation of the kNN algorithm is based

on get.knnx() function from the *FNN* R package.

**Segmentation of the EMT trajectory and robustness evaluation**

We used a Hidden Markov Model approach to identify of a discrete number of EMT states.

The input of this analysis was a matrix (M) where the rows were the TCGA samples (N) and

the columns the gene markers (G) of EMT (see the section "Computation of the EMT scores"

below for the list of genes). The original N columns were sorted for the t values of the

pseudotime (P). This matrix and P were provided as input for a lasso penalized regression. P

was used as response variable, the genes as the independent variables. The non-zero

coefficients obtained from this analysis were selected to create a sub-matrix of M that was

used as input for a Hidden Markov Model.

Different HMM models were tested while changing the number of states. After this tuning, and

through manual inspection, we determined that 3 states were most in line with biological

expectations. Each HMM state was assigned to a "biological group" (i.e. epithelial, hybrid

EMT, mesenchymal) by exploring the expression levels of known epithelial and mesenchymal

markers in each HMM state. The selection of the coefficients was performed with the R

package *glmnet*. The identification of the EMT states was done using the *deepmixS4* R

package.

To evaluate the "robustness" of the EMT states we applied the same procedure described

above while increasing levels of expression noise in the original dataset. We used the *jitter*

function in R to introduce a random amount of noise to the expression values of the genes

(from the default parameter of the *jitter* function to noise levels of 5500). For each noise level,

we repeated the analysis 100 times. We considered several metrics to measure the stability of

the HMM-derived EMT states. We reasoned that increasing noise could result in classification

mismatches of the samples compared to their originally assigned EMT state. Therefore, we

evaluated two metrics to assess the correct assignment of the samples to the original EMT

states. Firstly, for each level of noise added and at each iteration, we computed the change in

number of samples categorised in the new states compared to the original EMT states.

Second, we measured the assignment accuracy for the samples to the original EMT states.

**Computation of the EMT scores**

A list of epithelial and mesenchymal markers was compiled through manual curation of the

literature[30],[6],[9], as follows:

- epithelial genes: *CDH1*, *DSP*, *OCLN*, *CRB3*

- mesenchymal genes: *VIM*, *CDH2*, *FOXC2*, *SNAI1*, *SNAI2*, *TWIST1*, *FN1*, *ITGB6*,

*MMP2*, *MMP3*, *MMP9*, *SOX10*, *GSC*, *ZEB1*, *ZEB2*, *TWIST2*

EMT scores for each TCGA sample were computed in a similar manner as described by Chae

et al[62]. Briefly, the average z-score transformed expression levels of the mesenchymal

markers were subtracted from the average z-score transformed expression levels of the

epithelial markers. To segment the EMT trajectory, along with the epithelial and mesenchymal

markers we have also considered markers of hybrid EMT[63],[6]: *PDPN*, *ITGA5*, *ITGA6*, *TGFBI*,

*LAMC2*, *MMP10*, *LAMA3*, *CDH13*, *SERPINE1*, *P4HA2*, *TNC*, *MMP1*.

**EMT trajectory reconstruction of CCLE data and inference of the metastatic potential**

The RSEM gene-expression values of the Cancer Cell Line Encyclopedia[64] project were retrieved from the CCLE Data Portal. We used the same procedure described above to map the CCLE data onto the spontaneous MCF10 EMT trajectory. This allowed for the pseudotime to be quantified for each CCLE sample. A segmentation using a HMM model was performed to identify a discrete number of EMT states (n=3). The EMT scores were also computed for each cell line. These results were referenced against the metastatic potential scores from MetMap500[19]. The association between HMM states and experimentally measured metastatic potential groups in cell lines (non-metastatic, weakly metastatic and metastatic) was assessed using the *vcd* R package.

**Tumour microenvironment quantification**

The tumour purity values of TCGA samples were retrieved from Hoadley et al[65]. Immune deconvolution was performed using the ConsensusTME R package[66] and the ssGSEA method for cell enrichment analysis.

The results of ConsesusTME were used as input for a multinomial logistic regression model. The function multinom() (from the *nnet* R package) was used to determine the probability of each sample belonging to a macro-EMT state based on the cellular content of the sample.

Fibroblasts related gene-sets were manually curated as described in Supplementary Table S3. Fibroblasts enrichments scores were calculated via ssGSEA using the *GSVA*[67] R package.

**Genomic hallmark quantification**

To characterize the aneuploidy and the centromeric amplification levels of the samples in each EMT state we used the pre-computed values for TCGA from previous works [68],[40]. The hypoxia levels were quantified as described by Bandhari et al[69]. Several hypoxia gene signatures were considered, yielding similar results. Only the results obtained using the genes from Buffa et al[70] were reported. Finally, to estimate the levels of stemness in each EMT state,

we considered a catalogue of stemness gene sets[71]  and used them as input for gene set

enrichment analysis via the *GSVA* R package.

**Mutational signature analysis**

The identification of the mutational spectrum of the samples in each EMT state was performed

using a custom approach based on SigProfilerExtractor[41] and deconstructSigs[72].

SigProfilerExtractor was used for a de-novo identification of the mutational signatures.  We

selected the solutions in which the minimal stability was greater than 0.4 and the sum of the

minimal stabilities across signatures was greater than 1. The cosine similarity with mutational

signatures catalogued in the COSMIC database was computed, and only the solutions with

non-redundant signatures were selected. Next, we independently ran deconstructSigs. To

ensure consistency with Alexandrov et al[41], we evaluated the presence of the ageing-linked

SBS1 and SBS5, which have been identified in all cancers. We employed the following steps

to obtain a final list of signatures and their exposures for each tissue individually:

(1)  Considering the results obtained from deconstructSigs, the signatures with average

464             contribution (across all samples) greater than 5% were taken forward in the analysis.

465      (2) We combined the signatures obtained in (1) and by SigProfiler to obtain a final list of

466             signatures for the given tissue.  If SBS1 and SB5 were not present, we added these

467             signatures manually.

468   To identify EMT-associated mutational processes we used a similar approach to the one

469   described in Bhandari et al.[69], based on linear mixed-effect models. Cancer type was

470   incorporated as a random effect in each model.  An FDR adjustment was applied to the p-

471   values obtained from the analysis. The full model for a specific signature (SBS) is as follows:

472                            *EMT_score ~ SBS + (1|cancer)*

**Identification of cancer driver genes**

Single nucleotide variants were obtained from TCGA using the *TCGAbiolinks* R package and

the Mutect pipeline. To identify putative driver events that are positively selected in

association with an EMT state, we employed dNdScv[73], which quantified the ratio of non-

synonymous and synonymous mutations (dN/dS) in each gene and state, by tissue. We only

considered protein coding genes in the analysis, and filtered out consecutive mutations in the

genome, as recommended by the authors. All the somatic driver events with a q-value less

than of 0.10 were considered for downstream analysis.

**Definition of somatic copy number events**

Focal and arm-level copy number events across TCGA samples were identified using

GISTIC2[74], with the following parameters: armpeel = 1, brlen 0.5, cap 1.5, conf 0.99 -ta -td =

0.3, -genegistic = 1, gcm = extreme, -maxseg = -2500, qvt = 0.1, rx = 0, save gene = 1, broad

= 1.

**Identification of genomic events linked with EMT**

To search for genomic events linked with the described EMT macro-states, we considered all

somatic mutations, focal and arm-level copy number events in driver genes from the COSMIC

database that were obtained in the previous steps. Two parallel methodological approaches

based on lasso and random forest were used to identify events that could be predictive of

EMT transitions in a two-step process. First, feature selection was performed using a stability

selection approach. We used the function createDataPartition() from the *caret* R package to

generate an ensemble of vectors representing 1,000 randomly sampled training models. This

is an iterative approach, in which at each iteration a lasso analysis is performed, and the non-

negative coefficients computed by lasso are saved. This step was performed using the

cv.glmnet() function from *glmnet*. The tissue source was included as potential confounder in

the lasso model. The models were trained on 80% of the data. At the end of this stage, the

variables that were selected in at least the 80% of the iterations were taken forward and

employed as predictors. A similar approach was employed for feature selection and model

building with random forest.

In the second step, ROC curves were generated on the test dataset (20% of the data). In

addition, the predictors obtained from the two pipelines were also used as input for random

forest (*ranger* implementation), gradient boosting (*gbm*) and Naïve Bayes models. In these

cases, the trainControl() function (from the *caret* R package) was used in a 5-fold cross-

validation repeated 10 times. The function evalm() (from *MLeval* R package) was used to

compare the different machine learning methods. Only the features selected via the lasso

procedure were carried forward for downstream analysis.

Finally, to explore the relation between the outcomes (the EMT states) and the biomarkers we

used partial dependence plots generated with the function variable_effect() from the R-

package *DALEX*.

**Cancer cell fraction estimates**

The cancer cell fraction (CCF) of selected mutations was calculated using the following

formula:

$$514 \qquad CCF_i = \left(2 + \frac{[purity * (CN_i - 2)]}{purity}\right) \cdot VAF_i ,$$

where $CN_i$ stands for the absolute copy number of the segment spanning mutation *i* and $VAF_i$

is the variant allele frequency of the respective mutation. The purities of the TCGA samples

were obtained from Hoadley et al[65].

**Validation of the putative EMT biomarkers**

The following resources were used to validate the genomic associations with the EMT

programme: MET500[18], MSK-IMPACT[21], GENIE (Version 8)[20] and MetMap[19]. Different

strategies of pre-processing were employed as described below.

- **MET500**

523 Copy number regions encompassing 50% or more of the chromosomal arm were considered

524 as broad copy number events. Copy number regions below this threshold were classified as

525 focal events (based on hg19 cytoband positions and genomic coordinates obtained from

526 UCSC).

- **MSK-IMPACT**

The MSK-IMPACT resource consists of metastatic (n= 4,048) and primary tumors (n= 6,262).

First, we removed from the analysis all the samples where the metastatic site was 1) Blood

vessel, 2) Brain, 3) Spinal Cord or 4) Not available. Copy number events were classified as

broad or focal as described above for the MET500 data. Only regions where the absolute

copy number was greater than 1 and which were overlapping driver genes as annotated in

COSMIC were considered for downstream analysis. The frequency of each marker (e.g.,

TGFBR2_focal) was compared between the primary and metastatic tumors using a Chi-

square test. The resulting p-values were adjusted using Benjamini-Hochberg multiple testing

correction. Finally, only the markers with p-value<0.05 and odds ratio >1 were selected. The

same approach was applied for point mutations.

- **GENIE**

The GENIE dataset consists of metastatic (n=22,243) and primary tumors (n= 55,742). First,

we removed all the samples from 1) Glioma 2) Blood 3) Brain 4) Leukemia 5) Lymphoma 6)

Sellar 7) Nervous 8) CNS 9) Neuro 10) Neoplasms 11) Lymphoproliferative, or 12) Nerve. The

same approach described in the MSK-IMPACT section was used to analyze the GENIE

dataset. Only the copy number markers with q-value<0.10 and odds ratio >1 were selected.

Silent mutations were excluded.

- **MetMap**

To understand relevance of the hypothesised biomarkers to the metastatic dissemination of

various cancer cell lines, we downloaded the experimentally measured metastatic potential

levels for cancer cell lines from MetMap[19]. We compared metastatic potential between

samples with and without a specific EMT marker event (mutations or copy number

alterations), pan-cancer and by tissue. Only the markers that were linked with the hEMT or

MES states and that showed a statistically significant difference ($p < 0.05$) in metastatic

potential between the two groups (with and without alteration) have been considered.

**Gene essentiality evaluation using Project Achilles**

The viability of the cancer cell lines harbouring putative EMT biomarkers was evaluated based

on CRISPR screening data[75] conducted on 990 cell lines. CERES scores denoting gene

essentiality were downloaded from Project Achilles. Negative values of these scores indicate

that the depletion of a gene influences negatively the viability of a cell line. We only

considered genomic markers linked with the hEMT and MES states from our analysis and

assessed CERES scores for individual genes both pan-cancer and at tissue level.

**EMT-linked transcription factor network analysis**

Data from knockdown experiments of transcription factors regulating EMT was downloaded

from Meyer-Schaller et al[47]. Genes differentially expressed with an absolute log2 fold change

>1 and a q-value < 0.05 upon transcription factor knockdown were selected and intersected

with our list of putative EMT biomarkers.

**Proteomic biomarkers linked to EMT**

Level 4 RPPA proteomics data were retrieved from the Cancer Proteome Atlas (TCPAv3.0[76],

https://tcpaportal.org/tcpa/download.html). We identified proteins whose levels change

significantly between EMT states using logistic regression. Only the proteins with a q-value <

0.10 and odds ratio <-0.5 or >0.5 have been considered for downstream analysis.

**Tissue-specific EMT trajectory derivation**

To analyze the EMT in the context of several external stimuli we used the data generated by

Cook et al[49]. Specifically, we considered scRNA-seq data of MCF7 (breast cancer cell line)

and A549 (alveolar basal epithelial cells) under the stimulus of TNF, EGF and TGF-β1. For

each cell line, we integrated the expression data derived from all the conditions (TNF, EGF,

TGF-β1) using the FindIntegrationAnchors() and IntegrateData() functions from *Seurat*[77].

Next, we performed dimensionality reduction on the resulting data using UMAP (from the *uwot*

R package) and identified single cell expression clusters using the *mclust* R package. To

identify putative trajectories of EMT in each cell line, we used the R package *slingshot*[78] and

the function getLineages(). We manually defined the starting cluster for the trajectory as the

one with the lowest expression levels of mesenchymal markers. Using a similar bulk-to-single

cell mapping approach as described before, we mapped the RNA-seq data of LUAD and

BRCA tumours onto the trajectories derived from the single cell data (including batch effect

removal using ComBat, PCA on 25 dimensions and kNN clustering).  The EMT trajectories

derived from the single cell experiments presented different ramifications, therefore a simple

computation of the mean pseudotime values of the scRNA-seq samples associated with one

TCGA sample was not feasible. To overcome this limitation, we computed the mean of the X

and Y component values in the low dimensional matrix of the scRNA-seq data for those

scRNA-seq samples that were associated with one TCGA sample. The averaged coordinates

represent the position of the single cell samples in the UMAP space. The kNN analysis is

based on the transcriptome level of the bulk and scRNA-seq experiments. Therefore, even if

we are not able to compute pseudotime values of the TCGA samples with this analysis, we

can calculate the distance from any TCGA sample to the closest single cell RNA-seq sample,

and therefore obtain a "reference" for the single cell EMT trajectories. Using the "derived" X, Y

values of the TCGA samples, we used *mclust* to determine the number of clusters in the

TCGA cohort, then we performed a *de novo* reconstruction of the trajectory. The initial and

final clusters that determine the path of the EMT trajectory were defined manually on the basis

of increasing values of EMT scores (towards a mesenchymal phenotype) in each cluster. The

attribution of phenotype (epithelial-like, hEMT-like, mesenchymal-like) was performed based

on the lower and upper quartiles of the median EMT scores in each LUAD and BRCA cluster.

**Identification of the genomic events in the EMT tissue-specific analysis**

To identify the genes positively selected in each EMT tissue-specific cluster, we ran dNdScv[73]

in each cluster separately. To detect recurrent copy number alterations in lung

adenocarcinoma and breast cancer, we considered all genes with alterations in at least 10

patients and applied a Fisher's exact test to identify those that appeared enriched in a

particular EMT state. Only the genes with a q-value < 0.05 have been reported.

**Relation between EMT biomarkers and drug response**

We downloaded the drug sensitivity data from the Genomics of Drug Sensitivity in Cancer

database (GDSC)[22]. We considered only the mutated genes linked with EMT states and p-

value a cut-off <0.01. We also considered a second database, composed mainly of non-

oncology compounds[79]. For each compound, an ANOVA analysis was used to identify

significant changes in drug sensitivity between cell lines with or without a specific EMT

biomarker. Only the compounds with a p.value<0.01 were considered for downstream

analysis.

Finally, the POG570[23] dataset was used to study the relation between the EMT states and the

duration and effects of given cancer treatments. The EMT states in this dataset were inferred

similarly as described above using the kNN approach. To compare pre- and post-treatment

samples, we merged the TCGA and POG570 datasets and removed batch effects from the

two resources using ComBat. The resulting matrices were used to compute the EMT scores

all the samples and compare the levels of EMT between treatment-naïve and post-treatment

samples.

**Gene ontology analysis**

The characterization of the biological processes associated with the reported lists of genes

was performed using the R package *clusterProfiler*[80].

**Survival analysis**

Standardized clinical information for the TCGA cohort was obtained from Liu et al[81]. The

following end points were considered: overall survival, disease-specific survival and

progression-free interval. Patients were considered "lymph node positive" if they presented

infiltration in one or more lymph nodes. Cox proportional hazard models were used to model

survival based on variables of interest and to adjust for the following potential confounders:

tumour stage, age at diagnosis, gender and body mass index (BMI). Patients in clinical stages I-II were denoted as having "early stage tumours", while stages III-IV corresponded to "late stage tumours". The R packages *survival*, *survminer* and *ggforest* were used for data analysis and visualization.

**Data visualization and basic statistics**

Graphs were generated using the *ggplot2*, *ggpubr* and *diagram* R packages. Groups were compared using the Student's t test, Wilcoxon rank-sum test or ANOVA, as appropriate.

**Code**

All code developed for the purpose of this analysis can be found at the following repository: https://github.com/secrierlab/EMT/tree/EMTquant.v1.1 .

**ACKNOWLEDGEMENTS**

GMT was supported by a Wellcome Trust Seed Award in Science (215296/Z/19/Z). MS was supported by a UKRI Future Leaders Fellowship (MR/T042184/1) and an Academy of Medical Science Springboard award (SBF004\1042).

The results published here are in part based upon data generated by the TCGA Research Network: https://www.cancer.gov/tcga. The authors would like to acknowledge the American Association for Cancer Research and its financial and material support in the development of the AACR Project GENIE registry, as well as members of the consortium for their commitment to data sharing. Interpretations are the responsibility of study authors.

**AUTHOR CONTRIBUTIONS**

MS designed the study and supervised the analyses. GMT conducted all the analyses.

**COMPETING INTEREST STATEMENT**

None declared.

**REFERENCES**

1. Thiery, J. P., Acloque, H., Huang, R. Y. J. & Nieto, M. A. Epithelial-mesenchymal transitions in development and disease. *Cell* **139**, 871–890 (2009).

2. Kalluri, R. & Weinberg, R. A. The basics of epithelial-mesenchymal transition. *J Clin Invest* **119**, 1420–1428 (2009).

3. Pastushenko, I. & Blanpain, C. EMT Transition States during Tumor Progression and Metastasis. *Trends in Cell Biology* **29**, 212–226 (2019).

4. Goetz, H., Melendez-Alvarez, J. R., Chen, L. & Tian, X.-J. A plausible accelerating function of intermediate states in cancer metastasis. *PLOS Computational Biology* **16**, e1007682 (2020).

5. Pastushenko, I. *et al.* Identification of the tumour transition states occurring during EMT. *Nature* **556**, 463–468 (2018).

6. Puram, S. V. *et al.* Single-Cell Transcriptomic Analysis of Primary and Metastatic Tumor Ecosystems in Head and Neck Cancer. *Cell* **171**, 1611-1624.e24 (2017).

7. Karacosta, L. G. *et al.* Mapping lung cancer epithelial-mesenchymal transition states and trajectories with single-cell resolution. *Nature Communications* **10**, 5587 (2019).

8. Stemmler, M. P., Eccles, R. L., Brabletz, S. & Brabletz, T. Non-redundant functions of EMT transcription factors. *Nature Cell Biology* **21**, 102–112 (2019).

9. McFaline-Figueroa, J. L. *et al.* A pooled single-cell genetic screen identifies regulatory checkpoints in the continuum of the epithelial-to-mesenchymal transition. *Nat Genet* **51**, 1389–1398 (2019).

10. Rhim, A. D. *et al.* EMT and Dissemination Precede Pancreatic Tumor Formation. *Cell* **148**, 349–361 (2012).

678 11. Xu, J. *et al.* A novel Ubc9 -dependent pathway regulates SIRT1- ER-α Axis and BRCA1-

679 associated TNBC lung metastasis. *Integr Mol Med* **4**, (2017).

680 12. Nie, Z. *et al.* STAG2 loss-of-function mutation induces PD-L1 expression in U2OS cells.

681 *Ann Transl Med* **7**, (2019).

682 13. Chang, C.-J. *et al.* p53 regulates epithelial-mesenchymal transition and stem cell

683 properties through modulating miRNAs. *Nat Cell Biol* **13**, 317–323 (2011).

684 14. Stein, Y., Rotter, V. & Aloni-Grinstein, R. Gain-of-Function Mutant p53: All the Roads

685 Lead to Tumorigenesis. *Int J Mol Sci* **20**, (2019).

686 15. Vasudevan, A. *et al.* Single-Chromosomal Gains Can Function as Metastasis

687 Suppressors and Promoters in Colon Cancer. *Dev Cell* **52**, 413-428.e6 (2020).

688 16. Zhao, M., Liu, Y. & Qu, H. Expression of epithelial-mesenchymal transition-related

689 genes increases with copy number in multiple cancer types. *Oncotarget* **7**, 24688–24699

690 (2016).

691 17. Mak, M. P. *et al.* A patient-derived, pan-cancer EMT signature identifies global

692 molecular alterations and immune target enrichment following epithelial to mesenchymal

693 transition. *Clin Cancer Res* **22**, 609–620 (2016).

694 18. Robinson, D. R. *et al.* Integrative clinical genomics of metastatic cancer. *Nature* **548**,

695 297–303 (2017).

696 19. Jin, X. *et al.* A metastasis map of human cancer cell lines. *Nature* **588**, 331–336 (2020).

697 20. AACR Project GENIE Consortium. AACR Project GENIE: Powering Precision Medicine

698 through an International Consortium. *Cancer Discov* **7**, 818–831 (2017).

699 21. Zehir, A. *et al.* Mutational landscape of metastatic cancer revealed from prospective

700 clinical sequencing of 10,000 patients. *Nat Med* **23**, 703–713 (2017).

701    22.    Iorio, F. *et al.* A Landscape of Pharmacogenomic Interactions in Cancer. *Cell* **166**, 740–

702           754 (2016).

703    23.    Rheinbay, E. The genomic landscape of advanced cancer. *Nat Cancer* **1**, 372–373

704           (2020).

705    24.    Aiello, N. M. *et al.* EMT subtype influences epithelial plasticity and mode of cell

706           migration. *Dev Cell* **45**, 681-695.e4 (2018).

707    25.    Jolly, M. K. *et al.* Implications of the Hybrid Epithelial/Mesenchymal Phenotype in

708           Metastasis. *Front Oncol* **5**, 155 (2015).

709    26.    Plygawko, A. T., Kan, S. & Campbell, K. Epithelial–mesenchymal plasticity: emerging

710           parallels between tissue morphogenesis and cancer metastasis. *Philosophical Transactions*

711           *of the Royal Society B: Biological Sciences* **375**, 20200087 (2020).

712    27.    Sabe, H. Cancer early dissemination: cancerous epithelial-mesenchymal

713           transdifferentiation and transforming growth factor β signalling. *J Biochem* **149**, 633–639

714           (2011).

715    28.    Jolly, M. K., Ware, K. E., Gilja, S., Somarelli, J. A. & Levine, H. EMT and MET: necessary

716           or permissive for metastasis? *Mol Oncol* **11**, 755–769 (2017).

717    29.    Jing, Y., Han, Z., Zhang, S., Liu, Y. & Wei, L. Epithelial-Mesenchymal Transition in tumor

718           microenvironment. *Cell Biosci* **1**, 29 (2011).

719    30.    Gibbons, D. L. & Creighton, C. J. Pan-cancer survey of epithelial-mesenchymal

720           transition markers across the Cancer Genome Atlas. *Dev Dyn* **247**, 555–564 (2018).

721    31.    Choi, B.-J., Park, S.-A., Lee, S.-Y., Cha, Y. N. & Surh, Y.-J. Hypoxia induces epithelial-

722           mesenchymal transition in colorectal cancer cells through ubiquitin-specific protease 47-

723           mediated stabilization of Snail: A potential role of Sox9. *Sci Rep* **7**, 15918 (2017).

32.     Zhang, T., Suo, C., Zheng, C. & Zhang, H. Hypoxia and Metabolism in Metastasis. *Adv*

*Exp Med Biol* **1136**, 87–95 (2019).

33.     Aggarwal, V., Montoya, C. A., Donnenberg, V. S. & Sant, S. Interplay between tumor

microenvironment and partial EMT as the driver of tumor progression. *iScience* **24**, 102113

(2021).

34.     Tyler, M. & Tirosh, I. Decoupling epithelial-mesenchymal transitions from stromal

profiles by integrative expression analysis. *Nature Communications* **12**, 2592 (2021).

35.     Emami Nejad, A. *et al.* The role of hypoxia in the tumor microenvironment and

development of cancer stem cell: a novel approach to developing treatment. *Cancer Cell*

*Int* **21**, 62 (2021).

36.     San Juan, B. P., Garcia-Leon, M. J., Rangel, L., Goetz, J. G. & Chaffer, C. L. The

Complexities of Metastasis. *Cancers (Basel)* **11**, (2019).

37.     Mani, S. A. *et al.* The epithelial-mesenchymal transition generates cells with properties

of stem cells. *Cell* **133**, 704–715 (2008).

38.     Ganem, N. J., Godinho, S. A. & Pellman, D. A mechanism linking extra centrosomes to

chromosomal instability. *Nature* **460**, 278–282 (2009).

39.     Lingle, W. L. *et al.* Centrosome amplification drives chromosomal instability in breast

tumor development. *Proc Natl Acad Sci U S A* **99**, 1978–1983 (2002).

40.     de Almeida, B. P., Vieira, A. F., Paredes, J., Bettencourt-Dias, M. & Barbosa-Morais, N.

L. Pan-cancer association of a centrosome amplification gene expression signature with

genomic alterations and clinical outcome. *PLoS Comput Biol* **15**, e1006832 (2019).

41.     Alexandrov, L. B. *et al.* The repertoire of mutational signatures in human cancer.

*Nature* **578**, 94–101 (2020).

42.     Jing, L. *et al.* CNOT3 contributes to cisplatin resistance in lung cancer through

inhibiting RIPK3 expression. *Apoptosis* **24**, 673–685 (2019).

43.     Shirai, Y.-T. *et al.* CNOT3 targets negative cell cycle regulators in non-small cell lung

cancer development. *Oncogene* **38**, 2580–2594 (2019).

44.     Zhang, S. *et al.* Loss of VHL expression contributes to epithelial-mesenchymal

transition in oral squamous cell carcinoma. *Oral Oncol* **50**, 809–817 (2014).

45.     Duah, E. *et al.* Cysteinyl leukotriene 2 receptor promotes endothelial permeability,

tumor angiogenesis, and metastasis. *PNAS* **116**, 199–204 (2019).

46.     Liu, W., Xin, H., Eckert, D. T., Brown, J. A. & Gnarra, J. R. Hypoxia and cell cycle

regulation of the von Hippel–Lindau tumor suppressor. *Oncogene* **30**, 21–31 (2011).

47.     Meyer-Schaller, N. *et al.* A Hierarchical Regulatory Landscape during the Multiple

Stages of EMT. *Dev Cell* **48**, 539-553.e6 (2019).

48.     Yamaguchi, H. & Taouk, G. M. A Potential Role of YAP/TAZ in the Interplay Between

Metastasis and Metabolic Alterations. *Front. Oncol.* **10**, (2020).

49.     Cook, D. P. & Vanderhyden, B. C. Context specificity of the EMT transcriptional

response. *Nature Communications* **11**, 2142 (2020).

50.     Zhang, M., Wang, J. & Guo, J. Role of Regenerating Islet-Derived Protein 3A in

Gastrointestinal Cancer. *Front. Oncol.* **0**, (2019).

51.     Larue, L. & Bellacosa, A. Epithelial–mesenchymal transition in development and

cancer: role of phosphatidylinositol 3' kinase/AKT pathways. *Oncogene* **24**, 7443–7454

(2005).

52.     Mu, Z. *et al.* AZD8931, an equipotent, reversible inhibitor of signaling by epidermal

growth factor receptor (EGFR), HER2, and HER3: preclinical activity in HER2 non-amplified

inflammatory breast cancer models. *J Exp Clin Cancer Res* **33**, 47 (2014).

53.   Roche, J. The Epithelial-to-Mesenchymal Transition in Cancer. *Cancers (Basel)* **10**,

(2018).

54.   Tripathi, S., Chakraborty, P., Levine, H. & Jolly, M. K. A mechanism for epithelial-

mesenchymal heterogeneity in a population of cancer cells. *PLoS Comput Biol* **16**,

e1007619 (2020).

55.   Simeonov, K. P. *et al.* Single-cell lineage tracing of metastatic cancer reveals selection

of hybrid EMT states. *Cancer Cell* (2021) doi:10.1016/j.ccell.2021.05.005.

56.   Brown, M. S. *et al. Dynamic plasticity within the EMT spectrum, rather than static*

*mesenchymal traits, drives tumor heterogeneity and metastatic progression of breast*

*cancers*. http://biorxiv.org/lookup/doi/10.1101/2021.03.17.434993 (2021)

doi:10.1101/2021.03.17.434993.

57.   Muñoz, D. P. *et al.* Activation-induced cytidine deaminase (AID) is necessary for the

epithelial–mesenchymal transition in mammary epithelial cells. *PNAS* **110**, E2977–E2986

(2013).

58.   Bader, S. B. *et al.* Replication catastrophe induced by cyclic hypoxia leads to increased

APOBEC3B activity. *Nucleic Acids Research* **49**, 7492–7506 (2021).

59.   Cibulskis, K. *et al.* Sensitive detection of somatic point mutations in impure and

heterogeneous cancer samples. *Nature Biotechnology* **31**, 213–219 (2013).

60.   Colaprico, A. *et al.* TCGAbiolinks: an R/Bioconductor package for integrative analysis of

TCGA data. *Nucleic Acids Research* **44**, e71–e71 (2016).

61.   Johnson, W. E., Li, C. & Rabinovic, A. Adjusting batch effects in microarray expression

data using empirical Bayes methods. *Biostatistics* **8**, 118–127 (2007).

62.    Chae, Y. K. *et al.* Epithelial-mesenchymal transition (EMT) signature is inversely

associated with T-cell infiltration in non-small cell lung cancer (NSCLC). *Scientific Reports* **8**,

2918 (2018).

63.    Puram, S. V., Parikh, A. S. & Tirosh, I. Single cell RNA-seq highlights a role for a partial

EMT in head and neck cancer. *Mol Cell Oncol* **5**, e1448244 (2018).

64.    Barretina, J. *et al.* The Cancer Cell Line Encyclopedia enables predictive modelling of

anticancer drug sensitivity. *Nature* **483**, 603–607 (2012).

65.    Hoadley, K. A. *et al.* Cell-of-Origin Patterns Dominate the Molecular Classification of

10,000 Tumors from 33 Types of Cancer. *Cell* **173**, 291-304.e6 (2018).

66.    Jiménez-Sánchez, A., Cast, O. & Miller, M. L. Comprehensive Benchmarking and

Integration of Tumor Microenvironment Cell Estimation Methods. *Cancer Res* **79**, 6238–

6246 (2019).

67.    Hänzelmann, S., Castelo, R. & Guinney, J. GSVA: gene set variation analysis for

microarray and RNA-seq data. *BMC Bioinformatics* **14**, 7 (2013).

68.    Taylor, A. M. *et al.* Genomic and Functional Approaches to Understanding Cancer

Aneuploidy. *Cancer Cell* **33**, 676-689.e3 (2018).

69.    Bhandari, V. *et al.* Molecular landmarks of tumor hypoxia across cancer types. *Nature

Genetics* **51**, 308–318 (2019).

70.    Buffa, F. M., Harris, A. L., West, C. M. & Miller, C. J. Large meta-analysis of multiple

cancers reveals a common, compact and highly prognostic hypoxia metagene. *Br J Cancer*

**102**, 428–435 (2010).

71.    Miranda, A. *et al.* Cancer stemness, intratumoral heterogeneity, and immune response

across cancers. *Proc Natl Acad Sci U S A* **116**, 9020–9029 (2019).

72.  Rosenthal, R., McGranahan, N., Herrero, J., Taylor, B. S. & Swanton, C. deconstructSigs:

delineating mutational processes in single tumors distinguishes DNA repair deficiencies and

patterns of carcinoma evolution. *Genome Biology* **17**, 31 (2016).

73.  Martincorena, I. *et al.* Universal Patterns of Selection in Cancer and Somatic Tissues.

*Cell* **171**, 1029-1041.e21 (2017).

74.  Mermel, C. H. *et al.* GISTIC2.0 facilitates sensitive and confident localization of the

targets of focal somatic copy-number alteration in human cancers. *Genome Biol* **12**, R41

(2011).

75.  Dempster, J. M. *et al.* Extracting Biological Insights from the Project Achilles Genome-

Scale CRISPR Screens in Cancer Cell Lines. *bioRxiv* 720243 (2019) doi:10.1101/720243.

76.  Chen, M.-J. M. *et al.* TCPA v3.0: An Integrative Platform to Explore the Pan-Cancer

Analysis of Functional Proteomic Data. *Mol Cell Proteomics* **18**, S15–S25 (2019).

77.  Stuart, T. *et al.* Comprehensive Integration of Single-Cell Data. *Cell* **177**, 1888-1902.e21

(2019).

78.  Street, K. *et al.* Slingshot: cell lineage and pseudotime inference for single-cell

transcriptomics. *BMC Genomics* **19**, 477 (2018).

79.  Corsello, S. M. *et al.* Discovering the anti-cancer potential of non-oncology drugs by

systematic viability profiling. *Nat Cancer* **1**, 235–248 (2020).

80.  clusterProfiler: an R Package for Comparing Biological Themes Among Gene Clusters |

OMICS: A Journal of Integrative Biology.

https://www.liebertpub.com/doi/10.1089/omi.2011.0118.

81.  Liu, J. *et al.* An Integrated TCGA Pan-Cancer Clinical Data Resource to Drive High-

Quality Survival Outcome Analytics. *Cell* **173**, 400-416.e11 (2018).

**Figure 1. Pan-cancer EMT trajectories and underlying macro-states.** (a)   Workflow for

reconstructing the EMT trajectories of TCGA samples. 1: Bulk and single cell datasets are

combined and processed together to remove batch effects. 2: Dimensionality reduction using

PCA is performed. 3: A k-nearest neighbours (kNN) algorithm is used to map bulk RNA-

sequencing onto a reference EMT trajectory derived from scRNA-seq data. 4: Tumours are

sorted on the basis of their mesenchymal potential along an EMT "pseudotime" axis. (b)

Scatter plot of EMT scores along the pseudotime. Each dot corresponds to one bulk tumour

sample from TCGA. Samples are coloured according to the designated state by the HMM

model. (c) Diagram of the transition probabilities for switching from one EMT state to another,

as estimated by the HMM model. MES: fully mesenchymal state, hEMT: hybrid E/M, EPI:

epithelial state. (d) EMT scores compared across epithelial, hEMT, mesenchymal TCGA

samples, and the MET500 cohort. (e) Left: EMT pseudotime values compared between cell

lines from CCLE classified as "non metastatic" (aqua green), "weakly metastatic" (orange),"

metastatic" (red) according to the MetMap500 study. **$p<0.01$; ****$p<0.0001$. Right:

Association plot between the HMM-derived cell line states (rows) and their experimentally

measured metastatic potential (columns) ($p=2e-13$). (f) Distribution of the EMT states across

different cancer tissues. Each quarter of the pie corresponds to the 25% of the data. The

number of samples analysed is indicated for each tissue.

**Figure 2. Tumour extrinsic and intrinsic hallmarks of EMT.** (a) Heat map showcasing the

results of a multinomial logistic regression model trained to predict EMT states based on cell

infiltration in the microenvironment. Each row corresponds to a cell type and the

corresponding per-sample infiltration is highlighted via ssGSEA scores reported on the x axis.

The values reported in the heat map are the probabilities that a sample should fall into the

epithelial, hEMT or mesenchymal categories in relation to the ssGSEA score of a certain cell

type. (b) Cell abundance compared across the EMT states for significantly predicted cell types

in the multinomial analysis. (c) Levels of exhaustion quantified across the three EMT states.

(d) Heat map displaying the results a multinomial logistic regression model predicting EMT

state using fibroblast signatures from multiple publications. (e) Heat map displaying the pan-

cancer expression of the fibroblast COL11A1_FS signature genes. Highest expression is

observed in the hEMT state. (f) Median hypoxia values in the three different EMT states

across tissues. (g) Gene expression levels of the stemness marker CD44 compared across

the three EMT states. (h) Expression of the proliferation marker Ki67, mutational burden,

aneuploidy, and centromeric amplification levels compared across the three EMT states. (i)

Mutational signature exposures across TCGA samples sorted by EMT score. Only mutational

signatures that were significantly linked with EMT from the linear mixed models are displayed.

The corresponding EMT scores are displayed above. (j) Signature contributions from SBS1

and SBS13 compared between the three EMT states.

**Figure 3. Genomic driver events linked with EMT.** (a) The analytical workflow used to

detect putative EMT biomarkers. For each state and cancer type, we used dNdScv and

GISTIC to prioritise mutated genes and copy number events, respectively. These genomic

events were then employed as input for lasso modelling to classify EMT states. (b) Common

genomic features distinguishing the transformed states (hEMT, mesenchymal) from epithelial.

The individual bars depict the difference in the fraction of samples harbouring a specific

genomic alteration between every pair of EMT states: MES vs hEMT, MES vs EPI, hEMT vs

EPI. (c) Cancer cell fraction of common genomic markers between EMT states (left) and

hEMT-specific markers (right) with significantly distinct distribution between the states

(highlighted by the white-black colour gradient). (d) Scatter plot of EMT scores against the

cancer cell fraction across TCGA for KRAS, RB1 and FGFR2. (e) Top-ranked genomic

markers uniquely distinguishing hEMT from the epithelial state (notable ones highlighted in

bold). The fraction of alterations in EPI and hEMT samples, as well as the difference, are

displayed in the adjacent panels, The balloon chart on the right illustrates the association

between each marker and EMT, genome doubling (WGD), stemness index (mRNAsi),

centromeric amplification (CA20), and hypoxia. The size of the diamonds is proportional to the

significance of association, the colours report the odds ratios. (f) List of the top-ranked

genomic markers uniquely distinguishing the mesenchymal from the epithelial state and their

associated hallmarks.

**Figure 4. Validation of the pan-cancer genomic associations with EMT in external**

**datasets.** (a) UpSet chart displaying the overlap with external datasets for copy number

events derived from the MES vs EPI comparison. (b) Overlap with external datasets for

mutated gene events derived from the MES vs EPI comparison. (c-d) Same as (a-b) but for

events derived from the hEMT vs EPI comparison. (e) Fold changes in metastatic potential

across all the cell lines from CCLE harbouring point mutations in markers genes of EMT,

compared to that of cell lines without the respective mutation. The size and the colours of

the dots highlight the significance of the association ($p<0.05$). Events which increase

metastatic potential are highlighted in green. (*f*) Similar to (*e*), but with the analysis performed

at tissue level, ($p<0.05$). (*g-h*) Similar to (*e-f*), but for biomarkers harbouring copy number

alterations. (*i*) CERES essentiality scores from DepMap in individual cell lineages for genes

harbouring mutations linked with EMT. Negative values indicate increased essentiality. The

boxplots on the right indicate the CERES score distribution across all lineages. (*j*) Similar to (*i*)

but considering the genes linked with EMT via copy number alteration. (*k*) Genes showing

pan-cancer associations with EMT (rows) that are dysregulated as a result of knocking down

transcription factors relevant for EMT (columns). The colour gradient reflects the log2 fold

change in expression of the gene upon transcription factor knockdown (adjusted $p<0.05$). (*l*)

Similar to (*k*) but considering putative EMT biomarkers with alterations in copy number.

**Figure 5. EMT trajectories and associated genomic hallmarks in lung and breast**

**cancer.** (a) Lung adenocarcinoma samples from TCGA mapped onto the single cell EMT

trajectories derived in the A579 cell line. The first two principal components of the scRNA-seq

data are depicted. The hexagons represent the density of scRNA-seq samples, the TCGA

samples are coloured according to their membership to an EMT cluster. The red EMT

trajectory was computed on the scRNA-seq data. (b) Similar to (a), projecting breast cancer

samples from TCGA onto scRNA-seq from the MCF7 cell line. (c-d) EMT scores compared

across the clusters defined from (a-b) for lung and breast cancer, respectively. E1-3, H, M1-2

represent epithelial, hEMT and mesenchymal states in increasing order of transformation

(sorted by median). (e) Radial plot highlighting positively selected genes in each EMT state in

lung adenocarcinoma. Tile colours highlight the proportion of samples with a given mutated

gene. Each circle depicts the EMT score (colour) and the level of significance of association

between each driver gene and each cluster (size). (e) Similar to (f) but for the breast cancer

samples. (g) Fold change in metastatic potential across all the cell lines harbouring mutations

in marker genes of EMT in lung, compared with that of cell lines without the respective

mutation. The size and the colours of the dots highlight the significance of the association

($p < 0.05$). (h) Similar to (g), but with the analysis performed at tissue level ($p < 0.05$). (i) Similar

to (h), but for biomarkers harbouring copy number alterations.

**Figure 6. Clinical relevance of the EMT states.** (a) Overall survival compared between
MES, hEMT and EPI samples. (b) Progression free interval compared between the three
groups. (c) Mutated markers of EMT pan-cancer with a significantly worse or improved
outcome between the mesenchymal and epithelial states (q<0.001). (d) Mutated markers of
EMT pan-cancer with a significantly worse or improved outcome between the hEMT and

epithelial states. (e) Correlation between the EMT scores and IC50 values in cell lines from

GDSC treated with various drugs. The balloon chart on the left illustrates the association

between the IC50 for each compound and EMT. The size of the diamonds is proportional to

the significance of association. The IC50 ranges for all cell lines are depicted by the density

charts. (f) EMT scores compared between treatment naïve samples and those collected after

chemotherapy, by EMT state. (g-h) EMT scores compared between (unmatched) samples

before and after the treatment with capecitabine and letrozole, respectively. The tables

indicate the number of samples in each category.

## SUPPLEMENTARY MATERIAL

**Supplementary Figure S1. EMT trajectories and states in the TCGA and MET500**

**datasets**. (a) Expression of epithelial/mesenchymal markers along the EMT pseudotime

derived from single cell data. Each dot represents a TCGA sample and is coloured according

to the assigned EMT state. (b) Heat map highlighting the pan-cancer expression of known

EMT markers (coloured by their state-specificity). (c) Fraction of samples correctly assigned to

the original HMM state with increasing levels of gene expression noise in the original data. (d)

Accuracy of predicting the original HMM states with increasing levels of gene expression

noise in the original data. (e) EMT scores of the MET500 (red points) and TCGA (blue points)

samples plotted along the EMT pseudotime derived from single cell data. (f) EMT scores of

the CCLE samples plotted along the EMT pseudotime. (g) EMT scores compared across

tissues and between EMT macro-states. (h) Relation between the EMT states and the clinical

cancer stage, computed using Chi-square statistics. The circles represent the associations

between EMT states and cancer stage (early/late). The colour and size define the strength of

the association.

**Supplementary Figure S2. Tumour extrinsic properties in relation to EMT.** (a) Scatter plot

of tumour purity values (y axis) against EMT scores (x axis). Every dot corresponds to a

sample and is coloured according to the EMT macro-state. The Pearson correlation coefficient

and p-value of the association are also displayed. (b-c) Tumour purity compared between

discrete states of low and high stage of EMT transformation pan-cancer (EMT score<0 – low

stage; EMT score>0 – high stage). (c) Tumour purity compared between the three discrete

EMT macro-states pan-cancer. (d) The results of the multinomial logistic regression models

trained to predict EMT state based on non-cancer cell infiltration. The x-axis displays the

ssGSEA scores of a specific microenvironmental component. The y-axis visualizes the

probability of a sample being predicted to belong to a specific EMT state (EPI/hEMT/MES).

Every dot is a TCGA sample. (e) Tumour microenvironment composition compared between

the three EMT macro-states. (f) Active fibroblasts infiltration levels (quantified based on the

COL11A1 signature) compared between the three EMT macro-states.  (g) Hypoxia scores

compared between the three EMT macro-states pan-cancer.  (h) Correlation heat maps of the

stemness scores assessed in this study. The strength of the Pearson correlation is highlighted

by the colour gradient. (i) Stemness scores compared between each EMT state.

987
**Supplementary Figure S3. Mutational signatures across EMT states.** (a) Mutational

contribution of SBS16 compared between EMT states. (b) Mutational contributions across

TCGA samples for signatures derived as important in determining EMT by linear modelling

without accounting for the tissue effect. EMT scores are reported in the top panel and samples

are sorted according to these scores. Each value in the heatmap represents the contribution

of a specific signature to the genome in one sample. (c) Mutational contributions compared

between samples in the three EMT macro-states for selected signatures.

995
**Supplementary Figure S4. Discovery of genomic alterations linked with the EMT macro-**

**states.** (a) ROC curves of the MES vs EPI lasso model predictions based on genomic

markers obtained via the lasso procedure. (b) ROC curves of the MES vs EPI predictions

based on genomic markers obtained with the lasso procedure and tested using different

machine learning approaches. (c) ROC curves of the MES vs EPI lasso model predictions

based on genomic markers obtained with the random forest procedure. (d) ROC curves of the

MES vs EPI predictions based on genomic markers obtained with the random forest

procedure and tested using different machine learning approaches. (e-h) Same as (a-d) but

for the hEMT vs EPI model. (i-l) same as (a-d) but for the MES vs hEMT model. (m) Venn

diagram highlighting common and specific genomic markers in the MES vs EPI comparison

obtained using the lasso and random forest models, respectively. (n) Venn diagram

highlighting common and specific genomic markers in the hEMT vs EPI comparison obtained

using the lasso and random forest models, respectively. (o) Venn diagram highlighting

common and specific genomic markers between the EMT state comparison models built using

lasso. (p) Gene ontology terms enriched in the list of genes that distinguish the hEMT from the

EPI group. (q) Gene ontology terms enriched in the list of genes that distinguish the MES from

the EPI group.

1013
**Supplementary Figure S5. Characterization of the genomic alterations linked with the**

**EMT macro-states.** (a-b) Partial dependence charts of selected gene markers from the hEMT

vs EPI comparison model. The probability of a sample belonging to the hEMT group (y-axis) is

plotted against (a) different levels of copy number (numbers <0 indicate losses; 0=copy

number neutral; numbers >0 indicate gains) or (b) the presence of mutations (0/1:

absent/present). (c) Chromosomal arm events distinguishing the hEMT and EPI states.

Boxplots highlight the coefficients derived from the lasso analysis (hEMT vs EPI) of the

chromosomal arm events. The horizontal bar plots in the middle panels indicate the pan-

cancer percentage of samples with an amplification (red) or deletion (blue) of a specific

chromosomal arm. The balloon chart on the right highlights the associations between cancer

hallmarks and the chromosomal arm events. Only significant associations are displayed. (d)

Partial dependence charts of the MES vs EPI gene markers. The probability of a sample

belonging to the MES   group (y-axis) is plotted against different levels of copy number

(numbers <0 indicate losses; 0=copy number neutral; numbers >0 indicate gains) or the

presence of mutations (0/1: absent/present). (e) Chromosomal arm events distinguishing the

MES and EPI states. Boxplots depict the coefficients derived from the lasso analysis (MES vs

EPI) of the chromosomal arm events. The horizontal bar plots in the middle panels indicate

the pan-cancer percentage of samples with an amplification (red) or deletion (blue) of a

specific chromosomal arm. The balloon chart on the right highlights the associations between

cancer hallmarks and the chromosomal arm events. Only significant associations are

displayed.

1035
**Supplementary Figure S6. Proteomic profiling across the EMT states in TCGA.** (a)

Protein expression levels of canonical EMT markers compared between EPI, hEMT and MES

states. (b) Proteins with significantly different expression between hEMT and EPI states. The

x-axis and y-axis contain the OR and p-values obtained from the logistic regression analysis,

respectively. (c) Proteins with significantly different expression between MES and EPI states.

The x-axis and y-axis contain the OR and p-values obtained from the logistic regression

analysis, respectively. Panels (d-g) depict the results of gene ontology enrichment on: (d) up-

regulated proteins between hEMT and EPI, (e) down-regulated proteins between hEMT and

EPI, (f) up-regulated proteins between MES and EPI, (g) down-regulated proteins between

MES and EPI.

**Supplementary Figure S7. EMT states and genomic features of lung and breast**

**cancers.** (a) Top: results of slingshot analysis of EMT trajectories using the A549 cell line

data. The samples in each cluster are highlighted using different colours. Bottom: results of

slingshot analysis of EMT trajectories using the TCGA LUAD samples. Panels (b-c) display

circular heat maps with copy number amplifications (b) and deletions (c) significantly

associated EMT clusters in TCGA LUAD. The colour of each tile corresponds to the number of

samples with alterations in the respective cluster. The circles inside the modules highlight the

EMT scores of the TCGA samples in each cluster in the cases where the association between

the genomic event and EMT cluster is significant. (d) Top: results of slingshot analysis of EMT

trajectories using the MCF7 cell line data. Bottom: results of slingshot analysis of EMT

trajectories using the TCGA BRCA samples. Panels (e-f) display circular heat maps with copy

number amplifications (e) and deletions (f) significantly associated EMT clusters in TCGA

BRCA. (g) Essentiality across cell lines for genes whose mutations are linked with EMT in

lung or breast cancer. The heat map gradient indicates the CERES scores, with negative

values indicating increased essentiality. The boxplots in the right hand-side panel display the

CERES score distribution across all lineages. (h) Similar to (g) but considering the genes

linked with EMT via copy number alteration. (i) Genes mutated in relation to EMT in

breast/lung cancer that are dysregulated as a result of the knockdown of transcription factors

relevant for EMT. The rows display the EMT genomic markers obtained from previous

analyses, the columns indicate the transcription factors. The colour gradient reflects the log2

fold change in expression of the genes upon transcription factor knockdown. Only significant

changes are displayed (adjusted $p<0.05$). (j) Similar to (i) but considering the EMT markers

with alterations in copy number. Only lung EMT biomarkers showed changes in expression as

a result of transcription factor silencing.

**Supplementary Figure S8. Prognosis and therapeutic relevance of the EMT states.** (a)

Disease-specific survival of patients in TCGA by EMT macro-state of their corresponding

tumours. (b) Overall survival of patients in TCGA, stratified by the EMT macro-state of the

tumour and lymph node positivity. (c) Progression-free interval of the patients treated with

Oxaliplatin, by EMT macro-state of the tumour. (d) Significance of association with various

compounds plotted against the IC50 scores of cell lines from DepMap harbouring mutations in

genes identified between hEMT and EPI. E.g. cell lines harbouring mutations in the hEMT

marker ARID1A are more resistant to entinostat, vorinostat and AZD1332. (e) Significance of

association with various compounds plotted against the IC50 scores of cell lines from DepMap

harbouring mutations in genes identified between MES and EPI. (f) Significance of association

with various compounds plotted against the IC50 scores of cell lines from DepMap harbouring

mutations in genes identified commonly between transformed and EPI tumours. (g)

Differential cell viability after chemical perturbation with various compounds from the PRISM

drug repurposing resource for cell lines harbouring mutations identified by the hEMT vs EPI

comparison model. E.g. cell lines with mutations in the hEMT marker MAPK1 show an

increased cell viability after treatment with emetine. (h) Similar to (g), but only the genes

mutated in the MES vs EPI comparison have been considered. (i) EMT scores are compared

across distinct macro-states following EMT reconstruction of the post-treated samples from

POG570. The differences observed confirm the classification. (j-n) Duration of treatment (as

proxy for response) is compared between EMT macro-states for different drugs. (o-p) EMT

scores compared between unmatched but batch effect-corrected TCGA samples (treatment-

naïve) and POG570 samples after capecitabine and letrozole treatment, respectively.

Significant decreases in EMT score can be observed in both instances. Only transformed

samples (hEMT, MES) are considered for this analysis (numbers indicated in adjacent tables).

1096

**Supplementary Table S1. Pseudotime reconstruction of EMT trajectories in TCGA samples.** The results of the pan-cancer reconstruction of epithelial-to-mesenchymal trajectories is reported, along with the HMM state of the sample, and the corresponding EMT macro-state and score.

**Supplementary Table S2.** Distribution of the TCGA samples in each EMT state across clinical cancer stages.

**Supplementary Table S3. Catalogue of fibroblast gene-sets.** This table contains the gene sets used to quantify the fibroblast infiltration from bulk RNA-seq data of TCGA samples.

**Supplementary Table S4.** Stratification of samples according to hypoxia score and CD44 expression.

**Supplementary Table S5**. **Mutational signatures linked with EMT states pan-cancer.** The results of the linear mixed model trained to predict EMT state pan-cancer based on mutational signature contributions while accounting for the tissue effect are reported. Only significant mutational signatures are displayed (q < 0.05). The columns report the false discovery rate (FDR), the marginal ($R^2M$) and conditional ($R^2C$) residuals.

**Supplementary Table S6**. **Pan-cancer genomic markers linked with EMT states and protein analysis results.** (a) The results of the lasso models built to distinguish EMT states based on genomic markers. The first column indicates the comparison in which a biomarker (second column) has been identified. Additional details on the genes are provided. The coefficients from the lasso model (Median, Mean) along with the confidence intervals (Upper, Lower) are reported. The final two columns annotate if a gene is known to be relevant for EMT. (b) The results of the random forest models built to distinguish EMT states based on genomic markers. (c) Overlap between the genomic markers of EMT discovered via lasso modelling and a known list of genes with roles in cell migration from Novikov et al (Br J Cancer 2021). The cells marked with 1 highlight genes that were recapitulated by our

analysis and are known markers of migration. (d) Differentially expressed proteins between

EMT states.

**Supplementary Table S7. Genomic dependencies identified in the tissue specific EMT**

**trajectory analysis.** (a) Cancer drivers positively selected in each lung adenocarcinoma EMT

cluster. The rows depict the genes detected by dNdScv, while columns highlight the derived

EMT clusters obtained by projecting the bulk TCGA LUAD RNA-seq data onto the scRNA-seq

of the A549 cell line. Genes expressed in the cohort are indicated in bold. The last column

contains the overlap across the clusters for a given gene. (b) Number of samples containing

mutations in the positively selected genes in EMT clusters from lung adenocarcinoma. The

rows depict genes detected by dNdScv, while columns highlight the EMT clusters obtained by

projecting of the TCGA LUAD RNA-seq data onto the scRNA-seq of A549 cell line. The cells

contain the number of samples with a driver gene mutation in each cluster. (c) Overlap

between mutation events linked with EMT derived from the pan-cancer and lung-specific

analyses. Markers uniquely identified in LUAD are highlighted in bold, along corresponding

clusters in brackets. (d) Gene ontology enrichment analysis for gene markers positively

selected in different EMT states in lung adenocarcinoma. The table contains the top 10

significant gene ontology terms obtained using EnrichR. (e) Overlap between focal copy

number events linked with EMT derived from the pan-cancer and lung-specific analyses.

Markers uniquely identified in LUAD are highlighted in bold, along corresponding clusters in

brackets. The markers uniquely identified in the pan-cancer analysis have been removed. (f)

Cancer drivers positively selected in each breast cancer EMT cluster from TCGA. (g) Number

of samples containing mutations in the positively selected genes in EMT clusters from breast

cancer. (h) Overlap between mutation events linked with EMT derived from the pan-cancer

and breast-specific analyses. (i) Gene ontology enrichment analysis for gene markers

positively selected in different EMT states in breast cancer. (j) Overlap between focal copy

number events linked with EMT derived from the pan-cancer and breast-specific analyses.

**Supplementary Table S8. Cox regression model results on clinical end points.** (a)

Results of the multivariate Cox regression model for the overall survival. The mean and

standard deviation are reported for each covariate along with the hazard ratios (HR) from the

univariable and multivariable models. The p-value for the model is highly significant (p<1e-30).

(b) Results of the multivariate Cox regression model for the progression free interval (PFI).

The mean and standard deviation are reported for each covariate along with the hazard ratios

(HR) from the univariable and multivariable models. The p-value for the model is highly

significant (p<1e-10). (c) Results of the multivariate Cox regression model for the disease

specific survival. The mean and standard deviation are reported for each covariate along with

the hazard ratios (HR) from the univariable and multivariable models. The p-value for the

model is highly significant (p<1e-33).

**Supplementary Table S9. Mutation events impacting overall survival in TCGA.** The first

column indicates the EMT comparison model the marker was derived from, the second

column reports the name of the gene. Hazard ratios (log10) are reported along with

confidence intervals, p-values and outcome (Out: 'positive' indicates better prognosis for

patients harbouring the mutation in the respective gene, 'negative' indicates worse prognosis).