The cold-drought tolerance trade-off in temperate woody plants constrains range

size, but not range filling

Giacomo Puglielli[1*], Enrico Tordoni[2], Aelys M. Humphreys[3,4], Jesse M. Kalwij[5,6], Michael J.

Hutchings[7], Lauri Laanisto[1]

[1]*Institute of Agricultural and Environmental Sciences, Estonian University of Life Sciences,*
Tartu, 51006, Estonia;

[2] *Institute of Ecology and Earth Sciences, University of Tartu, Lai 40, 51005 Tartu, Estonia;*

[3] *Department of Ecology, Environment and Plant Sciences, Stockholm University, SE-106 91*
Stockholm, Sweden;

[4]*Bolin Centre for Climate Research, Stockholm University, SE-106 91 Stockholm, Sweden;*

[5] *Institute of Geography and Geoecology, Karlsruhe Institute of Technology, Karlsruhe,*
Germany;

[6] *Centre for Ecological Genomics and Wildlife Conservation, Department of Zoology, University*
of Johannesburg, Auckland Park 2006, South Africa;

[7] *School of Life Sciences, University of Sussex, Falmer, Brighton, Sussex, BN1 9QG, United*
Kingdom.

* *Author for correspondence – Email: Giacomo.Puglielli@emu.ee*

Acknowledgements: GP was supported by the Estonian Research Council grant (PSG708), and

by the Estonian University of Life Sciences (P200187PKEL). LL was supported by the Estonian

University of Life Sciences (P200190PKEL). GP also thanks Babak Naimi for helpful discussion

on the implementation of SDMs.

Abstract

Interspecific differences in plant species´ ranges are shaped by complex mechanistic interactions, which have so far remained largely beyond the reach of comprehensive models and explanations. Previous attempts to find underlying mechanisms by examining physiological tolerances to cold and heat separately have yielded contradictory results. Here we test the hypothesis that, instead of examining single stressors, abiotic stress tolerance syndromes that involve trade-offs between multiple abiotic stressors (namely drought, cold, waterlogging and shade), will provide reliable explanations.

We compiled a dataset of actual range size and range filling (the ratio between actual and potential species range) as range metrics for 331 temperate woody plants species from Europe and North America. Tolerance syndromes were expressed as two PCA axes. One axis reflects a drought-cold/waterlogging tolerance trade-off (cold/wet-drought trade-off), the second axis represents a shade tolerance spectrum. Phylogenetic generalized linear mixed models were used to model the range metric-tolerance axes relationships using latitude as an additional main effect, and phylogeny and plant functional type as random effects.

Actual range scaled negatively with the cold/wet-drought tolerance trade-off axis, mostly independently of latitude and continent. Thus, cold/wet-tolerant species had the largest ranges and drought tolerant species the smallest. The sign (-) of the relationship was independent of phylogeny and plant functional type. In contrast, range filling depended on latitude. However, deciduous and evergreen species displayed different distributions of range metrics and tolerance syndromes. No significant relationships with the shade tolerance spectrum were found.

Our findings demonstrate that the cold/wet-drought trade-off partly explains interspecific range size differences. However, this trade-off did not explain range filling. We also showed that fundamental adaptations of species also significantly influence range sizes – stress avoidance through the deciduous habit also explained interspecific differences in range size.

Keywords: abiotic stress, plant species distributions, cold tolerance, drought tolerance, Hutchinsonian niche, range filling, range size, woody plant species

Introduction

The identification of the factors shaping species range size and filling is a major focus of biogeography (Brown, 1984; Gaston *et al.*, 2009). In recent years, a number of theories aimed at explaining species distribution patterns at the global scale have been developed. For example, the mid-domain effect predicts that species close to the equator have larger distribution ranges (Colwell & Hurtt, 1994), whereas Rapoport´s rule assumes the opposite (Stevens, 1989). Several mechanistic hypotheses either addressing extrinsic factors (e.g. the climate variability hypothesis (Stevens, 1989) to explain Rapoport's rule) or intrinsic factors (e.g. the dispersal (Hanski *et al.* 1993) or niche breadth (Brown, 1984) hypotheses) have been proposed (reviewed in Sheth *et al.*, 2020). Most of the hypotheses formulated so far to explain interspecific differences in range size have emerged from zoology (Fine, 2015). In comparison, this topic has been little investigated in plant species (Sheth *et al.*, 2020), for which the determinants of range size, and its global variation, remain elusive.

Different metrics of range size are used in the literature to describe plant biogeographic patterns (Sheth *et al.*, 2020). These metrics include latitudinal or longitudinal range, extent of occurrence or area of occupancy, along with the ratio of realized range size (based on known species distributions) over potential range size (usually estimated using species distribution models), also known as range filling (Svenning & Skov, 2004; Paul *et al.*, 2009; Sheth *et al.*, 2020). Despite the use of different metrics, some consistent biogeographic patterns have emerged. For example, North American regions with strong climate instability have lower species richness and large-ranged species, while small-ranged species inhabit more species-rich regions with more stable climates (Morueta-Holme *et al.*, 2013; McFadden *et al.*, 2019). For European tree species, both potential range size and range filling increase with latitude (Svenning & Skov, 2004; Nogués-Bravo *et al.*, 2014), and both European and North American woody species tend to have larger ranges at higher latitudes (Morin & Chuine, 2006).

All these results are consistent with Rapoport´s rule, which predicts larger ranges and lower species richness at higher latitudes, possibly due to high latitude species being more tolerant of more variable environmental conditions (e.g., Morueta-Holme *et al.*, 2013), or to an increased frequency of pioneer species with broader niches at higher latitudes (Morin & Chuine, 2006). Despite these general tendencies, exceptions to Rapoport's rule have been documented.

For example, in the Americas, woody species' range size has a bimodal distribution in relation to
latitude, being largest in both north temperate and tropical areas (Weiser *et al.*, 2007). A similar
bimodal distribution has been observed for range filling in European (Svenning & Skov, 2004)
and North American tree species spanning sub-tropical to boreal climates (Seliger *et al.*, 2021).
Thus, although a positive range size-latitude relationship is well documented for plants, it is not
without exceptions and a mechanistic explanation of this pattern is still missing.

Morin & Chuine (2006) proposed that the proximate driver behind Rapoport's rule is
abiotic stress tolerance. In particular, they argued that intrinsic differences in abiotic stress
tolerance between species, and thus in their ability to persist under given resource regimes, might
explain interspecific variation in range size. Following this proposal, analysis of species'
inherent abilities to withstand extreme heat and/or cold has provided the main way to seek a link
between species' physiology and distribution ranges, chiefly expressed as latitudinal limits
(mostly for animals, e.g. Addo-Bediako *et al.*, 2000; Gaston *et al.*, 2009; Sunday *et al.*, 2011;
Araújo *et al.*, 2013). Comprehensive large-scale datasets of thermal tolerances have only
appeared recently for plant species (e.g., Lancaster & Humphreys, 2020). Nevertheless, relating
species' abiotic stress tolerances to their distribution ranges is complex. Cold and heat tolerance,
for instance, only have clear relationships with latitudinal and climatic gradients under certain
conditions. Cold tolerance seems to be more closely related to climatic conditions than heat
tolerance (Araújo *et al.*, 2013; Lancaster & Humphreys, 2020), and patterns are stronger for
northern hemisphere than southern hemisphere species. Similarly, cold and drought tolerances do
not display straightforward relationships with range filling, despite showing a clear latitudinal
pattern across Europe (Nogués-Bravo *et al.*, 2014). Thus, our comprehension of the relationship
between abiotic stress tolerance and species distributions remains poor.

Relating species' physiological tolerances to their geographical distribution patterns is
complicated by occupied ranges representing realized niches (e.g. Hutchinson, 1957), whereas
physiological tolerances should reflect species' fundamental niches. Furthermore, physiological
tolerances of different stressors might trade-off against each other due to correlation with
independent niche axes of the Hutchinsonian hypervolume (Sexton *et al.*, 2017), shrinking the
number of feasible tolerance combinations. Accounting for trade-offs between multiple
tolerances may more closely reflect species' realized physiological requirements (Sack, 2004;

Niinemets & Valladares, 2006; Laanisto & Niinemets, 2015; Puglielli *et al.*, 2021a) and possibly

reveal consistent relationships with realized range sizes. Thus, multivariate trade-off axes

between different tolerances might be needed to detect correlates with range size. Recently,

Puglielli *et al.* (2021a) examined multivariate trade-offs in woody species' ecophysiological

tolerances of four major abiotic stresses (cold, shade, drought and waterlogging). The trade-offs

were visualized in a triangular stress tolerance space (henceforth *Stress Space*). Ecophysiological

tolerance is defined as a species' ability to survive long-term extreme shortage of a given

resource in its natural environment (Niinemets & Valladares, 2006). The *Stress Space* was built

using published species-specific tolerance data (Niinemets & Valladares, 2006; Laanisto &

Niinemets, 2015) for ~800 Northern Hemisphere woody plant species. Each pair of coordinates

in the *Stress Space* reflects species-specific multi-stress tolerance syndromes shaped by trade-

offs among the different tolerances. In particular, the first *Stress Space* axis reflects a trade-off

between drought and cold/waterlogging tolerance. The second axis is a shade tolerance spectrum,

from low- to high shade tolerance. Thus, the *Stress Space* framework permits to link abiotic

stress tolerance syndromes to other aspects of species' biology, including range size. A well-

developed framework describing the relationship, and potential trade-offs, between multiple

abiotic stress tolerances is crucial for making realistic inferences about the role of abiotic stress

tolerance in shaping species distribution patterns, including global variation in range sizes

(Gaston *et al.*, 2009).

Identification and interpretation of range size correlates has the potential to increase

understanding of the factors and processes that influence species' ranges (Svenning & Skov,

2004; Estrada *et al.*, 2016). Although a complete spectrum of range size correlates is not yet

available (Estrada *et al.*, 2016, 2018), especially for plants, we employ a dataset on 300

temperate woody species from Europe and North America, to analyze the relationships between

abiotic stress tolerance syndromes, as summarized by the *Stress Space* axes, and species actual

range sizes and range filling. We decided to consider both of these range metrics because they

reflect different aspects of species' ranges. Actual range size includes historical legacies (e.g. for

temperate species, the degree to which they have been able to expand since the Last Glacial

Maximum; Svenning *et al.*, 2008; Normand *et al.*, 2011; Nogués-Bravo *et al.*, 2014; Estrada *et*

al., 2016) and ecological constraints (e.g. areas with suitable habitat, Linder *et al.*, 2013). Range

filling, on the other hand, is a measure of the extent to which species' ranges are at a climatic

equilibrium (Svenning & Skov, 2004). Thus, the traits that are positively correlated with actual

range size are expected to be those that are associated with range expansion (Estrada *et al.*,

2016, 2018), whereas traits associated with range filling reflect limits on species' distributions

imposed by historical non-climatic factors (Estrada *et al.*, 2016). Previous studies have mostly

attempted to link range filling to plant dispersal syndromes (Svenning *et al.*, 2008; Normand *et*

al., 2011; Nogués-Bravo *et al.*, 2014). However, even assuming successful migration and

dispersal, the *regeneration niche* theory (Grubb, 1977), also requires that a species is able to

survive the prevailing abiotic (and biotic) conditions to establish viable populations (Estrada *et*

al., 2018). The *Stress Space* framework can therefore provide further insights into the

determinants of species' range size and range filling.

In this study, we examined the relationships between temperate woody species range

sizes and range filling, and the *Stress Space* axes. Assuming these metrics are positively

correlated (e.g. Seliger *et al.*, 2021), we propose that species' abiotic stress tolerance syndromes

(reflected in their positioning along the *Stress Space* axes) can largely explain latitudinal

differences in range size and filling. Specifically, we hypothesized that:

1) As the first *Stress Space* axis represents a spectrum from cold/waterlogging- to drought

tolerant species, and assuming this reflects a latitudinal gradient (Nogués-Bravo *et al.*, 2014), we

expected a negative relationship between both range metrics and this axis.

2) Both range metrics will be independent of shade tolerance, which in turn is

independent of latitude. Shade tolerance data used in the *Stress Space* are measures of a species'

capacity for growth in low light conditions compared to the capacity for growth of coexisting

species (Niinemets & Valladares, 2006). Shade tolerance is also largely independent of the other

tolerances in the *Stress Space* (Puglielli *et al.*, 2021a). Therefore, shade tolerance can be either

high or low irrespective of latitude.

These hypotheses were tested by taking into account the effects of both phylogeny and

plant functional type, as defined by Puglielli *et al.* (2021b).

Methods

Actual range and occurrence data

We carried out an extensive literature search for polygons defining species' actual ranges for the

799 woody species in Puglielli *et al.* (2021a). We were able to retrieve polygons defining

species' actual range for 331 species. Specifically, spatial distributions of North American

species (*n* = 201) were obtained from the "*Digital representations of tree species range maps*

from Atlas of United States Trees" (the digitalized expert-drawn range maps by E.L. Little

Volumes 1–5, available at https://github.com/wpetry/USTreeAtlas). According to Seliger *et al.*

(2021), Little's maps are likely the best available approximation of historical realized ranges of

North America's trees and shrubs. This is because E.L. Little intentionally excluded range

alterations following Euro-American settlement Seliger *et al.,* 2021). Besides, Seliger *et al.*

(2021) also point to the fact that more recent plant occurrence databases are often limited to the

continental United States, while many of the tree species we analysed, as for Seliger *et al.*

(2021), extend into Canada and Northern Mexico.

Distributions of European species (*n* = 130) were gathered from the International Union for

Conservation of Nature (IUCN, www.iucnredlist.org), the European forest genetic resources

program (EUFORGEN, http://www.euforgen.org/species), and published papers (Kalwij *et al.*,

2014; Caudullo *et al.*, 2017; Wazen *et al.*, 2020). We consider using expert-based polygons as a

proxy of realized range size for European species set a reasonable assumption that is strongly

needed to make the results between continental species sets comparable.

Species occurrence records for the 331 species were obtained from the Global

Biodiversity Information Facility (GBIF, www.gbif.org/, accessed 21/12/2018; full list of data

sources in **Appendix S1**, **Supporting Information Table S1.1**). GBIF data were carefully

cleaned using both standardized and customized procedures (see **Appendix S1, Fig. S1.1**).

Potential range size and range filling calculation

To estimate species' potential range sizes, we used two presence-only models - i.e., Bioclim

(Busby, 1986), and Maxent (Phillips *et al.*, 2006) - to account for differences in potential range

size that may arise from algorithmic differences (Nogués-Bravo *et al.*, 2014). Nogués-Bravo et

al. (2014) also argued that, giving a possible mismatch between species' realized and potential

(i.e. modeled) species distributions, carefully implemented presence-only methods are expected

to return more reliable and parsimonious potential range estimates compared with more complex

model classes. Each model was fitted with three environmental parameters: growing degree days

at 5°C (GDD, unitless); climatic moisture index (the ratio of annual precipitation to annual

potential evapotranspiration, CMI, unitless) and mean minimum temperature of the coldest

month (T_{min}, °C). GDD and CMI data were obtained from the ENVIREM dataset (Title &

Bemmels, 2018) and T_{min} (i.e. Bio6) from WorldClim (Hijmans *et al.*, 2005); each at 10 arcmin

resolution. We decided to use only these three environmental predictors for two main reasons: (i)

these variables are historically considered as key environmental factors summarizing the

processes that limit the spatial distribution of plant species, and they were consistently used in

previous studies (e.g. Svenning & Skow, 2004; Nogués-Bravo *et al.*, 2014; Seliger *et al.*, 2021

and references herein); (ii) the use of few environmental predictors carefully selected for their

direct effects on plant species distributions avoid artificial and uninformative selection of

predictors on the sole basis of statistical techniques (e.g. variance inflation factor).

Before computing Species Distribution Models (SDMs), gathered GBIF occurrence data

were subjected to environmental filtering following Varela *et al.* (2014; see **Appendix S1, Fig.**

S1.1). All the 331 species had > 20 occurrences, which is considered a reasonable threshold for

fitting SDMs (Guisan *et al.*, 2017). The SDMs were fitted using the *sdm* R package (Naimi &

Araújo, 2016). For each run, 80% of species data was used for training, and the remaining 20%

for evaluating the model. 30 replicates per species were generated through bootstrapping and

20,000 background points were generated at each run. The Area under the ROC Curve (AUC;

Fielding & Bell, 1997) and True Skill Statistic (TSS; Allouche *et al.*, 2006) were used to

evaluate model performance. The SDMs predictions were converted into presence/absence maps

by using the threshold that maximized both sensitivity and specificity of the model. This is

considered the best option for presence-only methods (Liu *et al.*, 2013). The number of suitable

10 arcmin cells in the binary maps corresponded to potential range size while actual range was

determined by counting the 10 arcmin cells occupied by the polygons defining species' actual

range. Range filling (%) was then calculated as: (Actual range/Potential range)×100 (see

Appendix S2, Fig. S2.2). Due to a lower percentage of species with range filling estimates

greater than 100% (see **Appendix S3, Figs. S3.3-3.6, Table S3.2** for considerations on models'

performance and range filling estimates) only Bioclim derived estimates of potential range were

used in subsequent analyses, as in Nogués-Bravo *et al.* (2014). Furthermore, as pointed out by

the same authors, similar studies of range filling (e.g. Svenning & Skov, 2004) also used the

Bioclim algorithm to estimate potential range size for range filling calculation, thus ensuring that

our results are broadly comparable across a broader literature set.

In order to account for broad differences in species' adaptive syndromes, species were

classified according to three major plant functional types: deciduous angiosperms, evergreen

angiosperms and evergreen gymnosperms. For the complete list of species' actual range size

($\log_{10}$-transformed number of 10 arcmin cells), range filling, centroid latitude and species

classification according to their continental origin (N. America, Europe) and plant functional

type see **Appendix S4**.

Abiotic stress tolerance data

The species-specific estimates of tolerance of shade, drought, cold and waterlogging used to

define the *Stress Space* were obtained from the datasets of Niinemets & Valladares (2006) and

Laanisto & Niinemets (2015), which include stress tolerance scores for ~800 Northern

Hemisphere woody species. In the original data compilation (Niinemets & Valladares, 2006),

shade, drought and waterlogging tolerance were independently estimated by cross-calibrating

multiple tolerance scales reported in the literature where multiple measurements for one species

were available across tolerance scales. Cold tolerance data were extracted from USDA plant

hardiness data and represent species-specific averages gathered from multiple sources (see

Laanisto & Niinemets, 2015 for further details). In this respect, cold tolerance data do not show

any circularity with T_{min} obtained from WorldClim as it is not based on any spatially-explicit

information, and being an average species-specific estimate, it is not affected by a species range

size (i.e. species with larger ranges are more likely to encounter extreme mean temperatures).

All the stress tolerance scores vary in a continuous fashion between 1 - very intolerant species -

to 5 - very tolerant species (Niinemets & Valladares 2006 and Laanisto & Niinemets 2015).

The formalization of the *Stress Space* (Puglielli *et al.*, 2021a) revealed that two-

dimensions (principal components) capture ~80% of the variance in species-specific

combinations of shade, drought, cold and waterlogging. Each pair of coordinates in the *Stress*

Space corresponds to a species-specific stress tolerance syndrome. Stress Axis 1 is positively

correlated with drought tolerance and negatively correlated with both waterlogging and cold

tolerance. It is interpreted here as a cold-drought tolerance trade-off, where the term cold refers

to a short growing season. This interpretation stems from the positive covariance between cold

and waterlogging tolerance in our dataset: the highest cold tolerance is expected where

snowpacks are greater, resulting in later snowmelt, followed by waterlogging and consequently a

shorter growing season (Chuine, 2010). Stress Axis 2 is positively correlated with shade

tolerance, and represents a shade tolerance spectrum. Stress Axes are available in Puglielli *et al.*

(2021a).

Data analysis

Actual range size (hereafter range size) was log-transformed before analysis. Range filling was

strongly related with range size ($R^2 = 0.46$, p < 0.0001, $n = 331$), but not with $\log_{10}$-transformed

potential range size ($R^2 = 0.02$, p < 0.001, $n = 331$). Thus, we used the residuals deriving from

the relationship range size vs. range filling as a metric of range filling (Seliger *et al.*, 2021).

The relationships between stress axes and both range size and range filling residuals were

tested at different levels. First, we used Ordinary Least Square (OLS) regression analysis.

Second, we used quantile regression to provide a more comprehensive characterization of the

studied relationship (Ricotta *et al.*, 2010). In addition, quantile regression is mostly insensitive to

outliers (Ricotta *et al.*, 2010). Quantile regressions were run using the *quantreg* R package

(Koenker, 2017) using different quantiles of range metrics distribution ($\tau = 0.1$, $\tau = 0.25$, $\tau =$

0.50, $\tau = 0.75$ and $\tau = 0.90$). However, fitting models with many species while ignoring

phylogenetic relationships might lead to inflated type I errors (Freckleton *et al.*, 2002).

Consequently, as a third step, we computed Phylogenetic Generalized Linear Mixed Models

(PGLMMs, Ives & Helmus, 2011) using range metrics as response variable as a function of

stress axis 1 or 2 (considered separately), centroid latitude obtained from cleaned GBIF

occurrence data using the *geosphere* R package (Hijmans *et al.*, 2019), plus the first-order

interaction stress axis : latitude; species' phylogenetic relatedness and plant functional types

were considered as random effects. When the interaction was not found to be significant, only

the main effects were considered. PGLMMs were run using the *phyr* R package (Li *et al.*, 2020).

Phylogenetic data were retrieved for 325 species in our dataset using the mega-tree available via

the *V.PhyloMaker* R package (Jin & Qian, 2019). The mega-tree combines the phylogenies

developed by Zanne *et al.* (2014) and Smith & Brown (2018). Species nomenclature followed

The Plant List v.1.1 (2013). We also tested for potential signals of spatial autocorrelation in the

model residuals using spline correlograms from the *ncf* R package (Bjørnstad, 2020);

specifically, 95% pointwise bootstrap confidence intervals were computed from 5,000 bootstrap

samples of Pearson residuals.

Finally, we tested for differences between plant functional types in terms of the

distributions of actual range and range filling residuals, and positioning along stress axes using

the Kruskal-Wallis test. Pairwise multiple comparisons between group levels were carried using

Dunn's test by adjusting p-values with Holm correction.

All the data analysis procedures were repeated for the European and North American

species separately to account for possible geographic differences in patterns observed. All

statistical analyses were performed in R 4.0.5 (R Core Team, 2021). As we did not find any

relationship between either range size or range filling residuals with the shade tolerance

spectrum at any level of analysis, only the results relative to the cold-drought tolerance trade-off

axis are shown.

Results

Actual range scaled negatively with species positioning along the cold-drought tolerance trade-

off axis for species from both continents (Europe: slope = -0.10, R^2 = 0.06, p = 0.01, n = 130;

North America: slope = -0.18, R^2 = 0.14, p < 0.01, n = 201) (**Fig. 1 a,c**). A negative relationship

between actual range and cold-drought tolerance trade-off axis was also observed across the

considered quantiles of the response variable, but with differences between the species from the

two continents. For European species, the quantile regressions were mostly significant at average

to high values of actual range (**Fig. 1 a**). For North American species quantile regressions were

all significant, except the one fitted at the lowest quantile (**Fig. 1 c**).

Range filling residuals scaled negatively with species positioning along the cold-drought

tolerance trade-off axis only for European species (slope = -5.66, R^2 = 0.07, p < 0.01, n = 130)

and, as for actual range, the quantile regressions were mostly significant at average to high

values of range filling (**Fig. 1 b**). No significant relationship was found between range filling

residuals and the cold-drought tolerance trade-off axis for North American species, and the

relationship was not significant at any considered quantile (**Fig. 1 d**).

Fig. 1. Relationship between the cold-drought trade-off (Standard Deviation units) and actual range (log$_{10}$-transformed number of 10 arcmin cells) and range filling (residuals, see Methods) for: **(a,b)** European, and **(c, d)** North American woody plant species. Pale blue line represents the Ordinary Least Square fit. Black lines show the quantile regressions fitted at the 0.1, 0.25, 0.50, 0.75 and 0.90 quantile of the response variable distribution. Solid lines depict significant relationships at $p < 0.05$ while dashed lines represented not significant relationships. Sample size (n) is shown in panels **(a,c)** and applies also to the relationships involving range filling as the response variable. The color gradient reflects the progression from cold/wet-tolerant to dry/warm-tolerant species along the cold-drought trade-off.

The negative relationship between actual range and the cold-drought tolerance trade-off axis was not affected by including latitude as an additional main effect (**Table 1**), and the cold-drought tolerance trade-off axis effects were always greater than that of latitude for all data pooled (**see Appendix S5, Table S5.3**), and for each continent considered separately (**Table 1**). However, some differences between continents were detected. The cold-drought tolerance trade-off axis effect was only marginally significant ($p = 0.06$) for European species and the effect of latitude was not significant. The model explained 7% of the variance of actual range for European species. In contrast, for North American species, the effects of both the cold-drought tolerance trade-off axis and latitude were significant, but with opposite sign: the tolerance axis

maintained its negative relationship with actual size whereas latitude showed a positive

relationship (p < 0.01 and p < 0.05, respectively). This model explained 23% of the total

variation in actual range of North American species, and indicates that range sizes are greatest at

higher levels of cold/wet tolerance and at higher latitudes. Despite European and North

American species sets having no species in common, we explored whether differences between

continental species sets could be driven by differences in terms of genera composition.

Out of 106 genera in our entire dataset, there were 29 genera in common between the two

continents. These genera contributed 70% of the species included in the entire dataset. When the

analyses were repeated after removing genera that were unique to one or other continent, the

result remained significant for North American species, but the relationship between range size

and the cold-drought tolerance trade-off axis changed from being marginally significant to

become highly significant for European species as well ($R^2 = 0.12$, see **Appendix S5, Table**

S5.4). This indicates that differences in results between continents are partly driven by

differences in genera composition of their constituent species.

There was no significant interaction between the cold-drought tolerance trade-off axis

and latitude in any model involving actual range as the response variable for European or North

American species. Overall, species positioning along Stress Axis 1 was the main driver of

interspecific differences in actual range for both continents. Actual range data in relation to the

cold-drought tolerance trade-off axis and latitude are shown in **Appendix S5, Fig. S5.7a,c.**

The differences in results between continents were more pronounced for range filling

(**Table 1**). A significant positive interaction between the cold-drought tolerance trade-off axis

and Latitude was found in the model including range filling for European species, and it

explained approximately 19% of the variance. Despite a significant main effect of the cold-

drought tolerance trade-off axis on range filling (**Table 1**), we did not interpret this effect given

the presence of a significant interaction. We regard the interaction between the cold-drought

tolerance trade-off axis and latitude as the main driver of interspecific differences in range filling

across European species. For North American species, range filling showed a negative

relationship with latitude and no significant effect of the cold-drought tolerance trade-off axis.

This model suggests greater range filling at lower latitudes, but it explained only 4% of variance,

leaving range filling largely unexplained for North American species. Range filling data in

relation to the cold-drought tolerance trade-off axis and latitude are shown in **Appendix S5, Fig.**
S5.7b,d.

Regardless of the model, the effect of phylogenetic relatedness between species was
negligible (**Table 1**). In addition, the spline correlograms (see **Appendix S6, Fig. S6.8**) did not
reveal any evidence of spatial autocorrelation in the PGLMMs residuals. We can therefore safely
disregard spatial autocorrelation as a factor influencing model parameter estimates. Plant
functional types did not affect the strength or the sign of the relationships between the cold-
drought tolerance trade-off axis and either actual range or range filling. However, as a random
effect in PGLMMs, plant functional type had a greater effect overall than species phylogenetic
relatedness in terms of random effect variance (**Table 1**).

388   **Table 1. Results of the Phylogenetic Generalized Linear Mixed models.**

Response	Continent	Main effects	Estimate	SE	p	Model R^2	n	Random effects	Variance
Actual range	**Europe**	Intercept	3.39	0.34	< 0.0001	0.07	128	PFT	0.007
		Cold-drought trade-off	-0.07	0.04	0.06			Phylogeny	≈ 0
		Latitude	0.01	0.01	ns				
	North America	Intercept	2.9	0.22	< 0.0001	0.23	197	PFT	0.05
		Cold-drought trade-off	-0.11	0.03	< 0.001			Phylogeny	≈ 0
		Latitude	0.01	0.01	< 0.01				
Range filling	**Europe**	Intercept	27.08	17.37	ns	0.19	128	PFT	33.12
		Cold-drought trade-off	-43.18	11.77	< 0.0001			Phylogeny	0.003
		Latitude	-0.56	0.34	ns				
		Cold-drought trade-off * Latitude	0.69	0.22	< 0.001				
	North America	Intercept	22.96	9.25	0.05	0.04	197	PFT	3.97
		Cold-drought trade-off	0.01	1.56	ns			Phylogeny	0.001
		Latitude	-0.6	0.22	< 0.001				

Models were run by continent (Europe and North America) using actual range (log10-transformed number of 10 arcmin cells) and range filling (Residuals, see Methods) as

response variables, Cold-drought trade-off (Standard Deviation units) and latitude (centroid latitude, °) as main effects, and Plant Functional Type (PFT) and species

phylogenetic relatedness (Phylogeny) as random effects. Estimates of the main effects and their standard errors (SE), together with the variance explained by the model

(Model R^2) and sample size (n), are shown. The variance captured by the random effects is also shown.

The three plant functional types differed in distributions of actual range and range filling

values and in positioning along the cold-drought tolerance trade-off axis (Kruskal-Wallis test, p

≤0.05; **Fig. 2 a-f**). As a general trend, deciduous angiosperms have larger actual range and range

filling values (p ≤ 0.05; **Fig. 2 a,b,d,e**), and they occupy the cold/wet side of the cold-drought

tolerance trade-off (i.e. more negative values along the cold-drought tolerance trade-off axis)

(**Fig. 2 c,f**), compared to the other plant functional types. However, multiple comparisons

sometimes differed between continents (**Fig. 2 a-f**).

Fig. 2. Boxplots for the distribution of actual range size ($\log_{10}$-transformed number of 10 arcmin cells),
range filling (Residuals, see Methods) and positioning of species along the cold-drought trade-off
(Standard Deviation units) within each of the plant functional types (D-An = deciduous angiosperms; E-
An = evergreen angiosperms; E-Gy = evergreen gymnosperms) for: **(a-c)** European, and **(d-f)** North
American woody species. Sample size and data points for each plant functional type are shown. ***
indicates significant differences between plant functional types (Kruskal-Wallis, $p < 0.05$). '*' indicates
marginal significance ($p = 0.05$). Only significant and marginally significant differences are shown.
Multiple comparisons between groups were carried out using the Dunn's test and Holm correction for
multiple testing.

Discussion

Our results show that the cold-drought tolerance trade-off axis (hereafter cold-drought trade-off) partly explains interspecific differences in actual range size of temperate woody plants (**Fig. 1**). In particular, we found large-ranged species at the cold/wet tolerance end of the trade-off axis, and small-ranged species at the drought tolerance end. Despite some nuanced variation discussed below, this result was independent of continent, latitude, plant functional type and phylogeny, indicating its generality. In contrast, range filling showed different patterns in Europe and North America: the interaction between the cold-drought trade-off and latitude had the largest influence on range filling for European species, whereas latitude was the only driver of range filling in our North American species pool.

Determinants of range size variation in temperate woody plants

For North American species, we found that cold/wet tolerant species occurring at high latitudes (e.g. *Salix* spp., *Larix laricina* (Du Roi) K.Koch) generally have the largest ranges (**Table 1**). Consistent with this, large ranged, cold-tolerant North American tree species are known to be generally absent from regions that are consistently warm and moist, such as the southeastern regions of the continent (Pither, 2003). Palaeoecological records also provide evidence for rapid northward range shifts in North American large-ranged trees after the latest ice age (Seliger *et al.*, 2021, and references herein). Conversely, some species might have maintained relatively small realized ranges following deglaciation, perhaps due to trait syndromes guaranteeing competitive advantages only in specific ice age refugia (Seliger *et al.*, 2021). Possible examples are species with drought tolerance strategies that prevented northward range expansions after glacial retreat, such as *Juniperus deppeana* Steud., *Pinus monophylla* Torr. & Frém., and *Quercus douglasii* Hook. & Arn., among other species with relatively small ranges that are confined to SW-North America. Long-term drought, and likely adaptations to tolerate such conditions, are an important constraint on plant species distributions (e.g. Normand *et al.*, 2009). Similarly, Pither (2003) hypothesized that latitudinal patterns in the range sizes of North America woody species reflect a potential trade-off between species' cold tolerance strategies and their competitiveness in warmer environments. In support of this, our results show that the cold-drought trade-off, which largely reflect species biogeographical history, is a mechanism that actively influences interspecific differences in range size for North American plant species.

Cold-tolerant European species also have large ranges (**Fig. 1 a**), but the signal was

weaker than for North American species (**Table 1**). This difference is caused by differences in

genera composition between the continental species sets (see Results section), and by a cluster of

European species with intermediate positions along the cold-drought trade-off having large

ranges (**Fig. 1 a**). These large-ranged species include *Picea abies* (L.) H.Karst, *Pinus sylvestris*

L. and *Betula pendula*. Roth. Such species survived the last glacial maximum in central and/or

eastern Europe with easy access to Northern Europe, with possibility for rapid northward

expansion, after ice retreat (Normand *et al.*, 2011). While these species are indeed cold tolerant,

their tolerance syndromes include some degree of drought tolerance as well, shifting them

towards the center of the cold-drought trade-off axis. This effect has the potential to decouple

these species from the negative relationship between the cold-drought trade-off and range size,

indicating the importance of considering stress tolerance syndromes based on several factors

rather than single tolerances when investigating interspecific differences in ranges. Other species

that are potentially decoupled from this relationship belong to European genera that do not occur

in the North American species set. More studies are needed for these European species. In the

case of small range, drought-tolerant species, the explanation provided for North American

species is likely to apply to European species as well (e.g. Normand *et al.*, 2009).

The negative scaling of range size along the cold-drought tolerance trade-off is also

consistent with macrophysiological evidence suggesting that pre-adaptation to low temperature

or species-specific abilities to adapt to freezing temperatures may have favoured species'

northward migration after glacial retreat (e.g. Araújo *et al.*, 2013; Lancaster & Humphreys,

2020). In contrast, it has been suggested that heat tolerances might be less adaptable (e.g. Araújo

et al., 2013) and, considering that drought-tolerant species usually thrive in hot environments,

adaptations to tolerate drought might have concomitantly prevented the northward post-glacial

race for drought tolerant plants. This suggests that the cold-drought trade-off might limit niche

expansion for temperate woody species, as has been suggested for heat stress globally (Araújo *et*

al., 2013). Importantly, however, despite the fact that drought and heat stress can covary in

warm-temperate ecosystems, we argue that the effect of drought tolerance on range size should

be considered separately from that of heat tolerance. The most heat tolerant plants are from both

dry and moist environments (Lancaster & Humphreys 2020), and although transpirational

cooling generally costs moisture, plants have also evolved adaptations to reduce the detrimental

effects of heat stress under drought (Flexas *et al.*, 2014). For example, among a plethora of other

adaptations, changes in water-use efficiency (Flexas *et al.*, 2014) or leaf movements (Puglielli *et*

al., 2017) can decouple leaf physiological responses to high temperatures from those to drought

in Mediterranean woody plants.

We have shown that species positioning along the cold-drought trade-off axis imposes

fundamental constraints upon interspecific differences in the range size of woody species, and

contributes to shaping their realized niches (i.e. actual range) across continents. Species

positioning in the *Stress Space*, which is determined by trade-offs between multiple tolerances

(Puglielli *et al.*, 2021a), can be interpreted as a measure of a species' realized niche based on

abiotic resources (*sensu* Hutchinson, 1957) more than as a measure of the fundamental niche.

After all the trade-off between tolerances shrinks fundamental niche size towards that of the

realized niche (e.g. Sack, 2004). Therefore, by representing the typical multiple abiotic stress

levels to which a plant species has adapted, its position in the *Stress Space* inherently

corresponds to a set of energy constraints that define plant life history strategies. Such strategies

are thought to directly control species' geographical distributions (Morin & Chuine, 2006).

Niche position along resource-defined axes is a strong predictor of range size and occupancy in

many animal groups (Seliger *et al.*, 2021), and our results show that this is also true for

temperate woody plant species.

Range filling is not affected by the cold-drought tolerance trade-off

All species are expected to have constraints on the extent to which they fill their range because

of their specific physiological and ecological requirements (Paul *et al.*, 2009), and their inherent

trade-offs. Our results suggest that other factors that covary with latitude, not considered in this

study (e.g. dispersal syndromes, Estrada *et al.*, 2016), might be important for explaining range

filling pattern. Such historical non-climatic limitations can ultimately also covary with species'

abiotic stress tolerances. This might explain why a positive interaction between the cold-drought

tolerance trade-off and latitude drives their range filling differences. Similarly, Nogués-Bravo *et*

al. (2014) proposed that the negative correlation between seed mass and range filling found

across 38 European tree species could be due to covariation between seed mass and other factors

(e.g. drought tolerance), but further analysis including a larger number of species would be

needed in order to test this claim.

More importantly, our results highlight different range filling determinants between the

two continents, that contrarily to actual range, did not depend on differences in genera between

continents (see **Appendix S5**, **Table S5.4**). European species' range filling was driven by a

positive interaction between latitude and the cold-drought tolerance trade-off, while North

American species' range filling was influenced only by a negative effect of latitude. This

negative relationship between range filling and latitude for North American species was in

agreement with a previous study that also reported a positive relationship range filling-longitude

(Seliger *et al.*, 2021), suggesting that longitude can indeed alter the expected range filling-

latitude relationship. Differences between continents might therefore depend on their

geographical extent. Europe has a much smaller latitudinal and longitudinal range and less

gradual geographical clines than North America (Morin & Chuine, 2006). In addition, some

European species ranges also occur outside Europe (e.g., species with a Eurasian distributions)

and this might have influenced model's projections (see **Appendix S3**). According to the above,

we therefore suggest that the interaction between these contrasting continental features and

intrinsic drivers, such as dispersal syndromes, might determine continent-specific levels of range

filling.

Differences among plant functional types

Plant functional type (PFTs) did not affect the negative scaling of range size with the cold-

drought trade-off (**Table 1**). However, the distribution of range size, range filling and positioning

along the cold-drought trade-off axis differed between PFTs (**Fig. 2 a-f**). This probably explains

why PFTs generally accounted for more of the random effects variance (even though the amount

was generally low) for range metrics compared to phylogeny (**Table 1**). In general, despite some

overlap between PFTs, deciduous angiosperms showed the highest values for both actual range

and range filling (consistent with large-ranged species also having greater range filling, Seliger *et*

al., 2021), and they were located further towards the cold tolerance end of the trade-off axis.

Using a dataset of European and North American woody species, Morin & Chuine (2006) also

found that deciduous species were overrepresented among the large-ranged species.

Many deciduous angiosperms differ in trait syndromes from evergreen angiosperms or

evergreen gymnosperms (e.g. Puglielli *et al.*, 2021b), and different trait syndromes match

differences in PFTs geographical distributions (Zanne *et al.*, 2018). For example, in the northern

hemisphere, deciduous species tend to be more frequent in cold climates compared to evergreen

broad-leaved angiosperms, despite some overlap between PFTs at almost all latitudes (Zanne *et*

al., 2018). Conversely, while adaptations to tolerate drought closely match the distribution of

evergreen species, this is not always true for deciduous species (e.g. see Kunert *et al.*, 2021),

suggesting that drought does not always limit deciduous species spatial distribution.

Deciduousness *per se* has been the common explanation for the ability of deciduous

species to colonize either cold or dry environments, as it is an adaptation that permits avoidance

of unfavorable environmental conditions. This explains why deciduous angiosperms in our

dataset have larger ranges than other PFTs, and are mostly cold-tolerant. However, if

deciduousness is also a successful drought avoidance strategy that could drive to large ranges in

drought tolerant species as well, the relationship between range size and the trade-off axis should

be not significant rather than negative. We can reconcile this by considering that woody species

of the warm-temperate/temperate regions of the Northern Hemisphere are not generally drought-

deciduous. Drought tolerant deciduous species have sufficiently though leaves to widen their

growing season beyond the usual duration of periods of drought (Hallik et al. 2009). Such

adaptation is of course part of a wider trait syndrome (see Hallik *et al.*, 2009), including, for

example, greater biomass allocation to roots in species with greater drought-tolerance (Puglielli

et al., 2021b), or smaller vascular conduits (Olson *et al.*, 2018), that support the investment in

tough leaves. This is consistent with the previous discussion of a link between species' tolerance

strategies, trait syndromes and range size.

To summarize, while we recognize that adaptations to very low temperatures are

complex, and involve a combination of avoidance and tolerance strategies together with

acclimation (Schubert *et al.*, 2020), we argue that large range sizes at the cold tolerant end of the

trade-off axis is made viable because of the deciduous habit in temperate woody species. In this

instance, stress avoidance is more important as an asset than stress tolerance. However, we also

observed a greater overlap, in both range size and positioning along the trade-off axis, between

freezing-tolerant North American gymnosperms and deciduous angiosperms than with evergreen

angiosperms (**Fig. 2 d,f**). Thus, we do not exclude the additional role for freezing tolerance in

guaranteeing large ranges.

Conclusions

Our results demonstrate that the cold-drought tolerance trade-off partly explains interspecific differences in range size across temperate woody plant species and that this relationship is largely independent of latitude and consistent with woody species biogeographical histories in the considered continents. Notably, our findings also suggest that accounting for species' abiotic stress tolerance towards multiple stresses can reconcile macroecological and macrophysiological theories aimed at explaining range size differences among woody plant species, supporting a previous hypothesis by Morin & Chuine (2006). However, our results concerning the impact of abiotic stress tolerance on range filling were inconclusive, suggesting that other factors not studied here and that covary with latitude and/or abiotic stress tolerance syndromes are the main drivers of range filling in temperate woody species. Finally, our results also demonstrate the importance of considering trait syndromes, reflected here as differences between plant functional types, in clarifying the ways in which species' adaptations influence broad interspecific differences in range size in woody plant species in relation to abiotic stress tolerance.

References

Addo-Bediako, A., Chown, S.L. & Gaston, K.J. (2000) Thermal tolerance, climatic variability and latitude. *Proceedings of the Royal Society of London. Series B: Biological Sciences*, **267**, 739–745.

Allouche, O., Tsoar, A. & Kadmon, R. (2006) Assessing the accuracy of species distribution models: prevalence, kappa and the true skill statistic (TSS). *Journal of Applied Ecology*, **43**, 1223–1232.

Araújo, M.B., Ferri-Yáñez, F., Bozinovic, F., Marquet, P.A., Valladares, F. & Chown, S.L. (2013) Heat freezes niche evolution. *Ecology Letters*, **16**, 1206–1219.

Brown, J.H. (1984) On the Relationship between Abundance and Distribution of Species. *The American Naturalist*, **124**, 255–279.

Busby, J.R. (1986) A biogeoclimatic analysis of *Nothofagus cunninghamii* (Hook.) Oerst. in southeastern Australia. *Australian Journal of Ecology*, **11**, 1–7.

Caudullo, G., Welk, E. & San-Miguel-Ayanz, J. (2017) Chorological maps for the main European woody species. *Data in Brief*, **12**, 662–666.

Chuine, I. (2010) Why does phenology drive species distribution? *Philosophical Transactions of*
the Royal Society B: Biological Sciences, **365**, 3149–3160.

Colwell, R.K. & Hurtt, G.C. (1994) Nonbiological Gradients in Species Richness and a Spurious
Rapoport Effect. *The American Naturalist*, **144**, 570–595.

Estrada, A., Morales-Castilla, I., Caplat, P. & Early, R. (2016) Usefulness of Species Traits in
Predicting Range Shifts. *Trends in Ecology & Evolution*, **31**, 190–203.

Estrada, A., Morales-Castilla, I., Meireles, C., Caplat, P. & Early, R. (2018) Equipped to cope
with climate change: traits associated with range filling across European taxa.
Ecography, **41**, 770–781.

Fielding, A.H. & Bell, J.F. (1997) A review of methods for the assessment of prediction errors in
conservation presence/absence models. *Environmental Conservation*, **24**, 38–49.

Fine, P.V.A. (2015) Ecological and Evolutionary Drivers of Geographic Variation in Species
Diversity. *Annual Review of Ecology, Evolution, and Systematics*, **46**, 369–392.

Flexas, J., Diaz-Espejo, A., Gago, J., Gallé, A., Galmés, J., Gulías, J. & Medrano, H. (2014)
Photosynthetic limitations in Mediterranean plants: A review. *Environmental and*
Experimental Botany, **103**, 12–23.

Freckleton, R.P., Harvey, P.H., Pagel, M. & Losos, A.E.J.B. (2002) Phylogenetic Analysis and
Comparative Data: A Test and Review of Evidence. *The American Naturalist*, **160**, 712–
726.

Gaston, K.J., Chown, S.L., Calosi, P., Bernardo, J., Bilton, D.T., Clarke, A., Clusella-Trullas, S.,
Ghalambor, C.K., Konarzewski, M., Peck, L.S., Porter, W.P., Pörtner, H.O., Rezende,
E.L., Schulte, P.M., Spicer, J.I., Stillman, J.H., Terblanche, J.S. & Van Kleunen, M.
(2009). Macrophysiology: A Conceptual Reunification. *The American Naturalist*, **174**,
595–612.

Hijmans, R.J., Williams, E., Vennes, C. (2019). Spherical Trigonometry. R package available at
https://CRAN.R-project.org/package=geosphere.

Grubb, P.J. (1977) The Maintenance of Species-Richness in Plant Communities: The Importance
of the Regeneration Niche. *Biological Reviews*, **52**, 107–145.

Guisan, A., Thuiller, W. & Zimmermann, N.E. (2017) Habitat Suitability and Distribution
Models: With Applications in R. Cambridge: Cambridge University Press.

Hallik, L., Niinemets, Ü. & Wright, I.J. (2009) Are species shade and drought tolerance reflected
in leaf-level structural and functional differentiation in Northern Hemisphere temperate
woody flora? *New Phytologist*, **184**, 257–274.

Hanski I, Kouki J, Halkka A. 1993. Three explanations of the positive relationship between
distribution and abundance of species. In: Ricklefs R, Schluter D, eds. *Species diversity*
in ecological communities: historical and geographical perspectives. Chicago, IL, USA:
The University of Chicago Press, 108–116.

Hijmans, R.J., Cameron, S.E., Parra, J.L., Jones, P.G. & Jarvis, A. (2005) Very high resolution
interpolated climate surfaces for global land areas. *International Journal of Climatology*,
25, 1965–1978.

Hutchinson GE. 1957. Concluding Remarks. Cold Spring Harbor Symposia on Quantitative
Biology 22: 415-427.

Ives, A.R. & Helmus, M.R. (2011) Generalized linear mixed models for phylogenetic analyses of
community structure. *Ecological Monographs*, **81**, 511–525.

Jin, Y. & Qian, H. (2019) V.PhyloMaker: an R package that can generate very large phylogenies
for vascular plants. *Ecography*, **42**, 1353–1359.

Kalwij, J.M., Robertson, M.P., Ronk, A., Zobel, M. & Pärtel, M. (2014) Spatially-Explicit
Estimation of Geographical Representation in Large-Scale Species Distribution Datasets.
PLoS One, **9**, e85306.

Koenker, R. 2017. Package 'quantreg.' R Found. Stat. Comp., Vienna, Austria.

Kunert, N., Zailaa, J., Herrmann, V., Muller-Landau, H.C., Wright, S.J., Pérez, R., McMahon,
S.M., Condit, R.C., Hubbell, S.P., Sack, L., Davies, S.J. & Anderson-Teixeira, K.J.
(2021) Leaf turgor loss point shapes local and regional distributions of evergreen but not
deciduous tropical trees. *New Phytologist*, **230**, 485–496.

Laanisto, L. & Niinemets, Ü. (2015) Polytolerance to abiotic stresses: how universal is the
shade–drought tolerance trade-off in woody species? *Global Ecology and Biogeography*,
24, 571–580.

Lancaster, L.T. & Humphreys, A.M. (2020) Global variation in the thermal tolerances of plants.
Proceedings of the National Academy of Sciences USA, **117**, 13580–13587.

Li, D., Dinnage, R., Nell, L.A., Helmus, M.R. & Ives, A.R. (2020) phyr: An r package for
phylogenetic species-distribution modelling in ecological communities. *Methods in*
Ecology and Evolution, **11**, 1455–1463.

Linder, H.P., Antonelli, A., Humphreys, A.M., Pirie, M.D. & Wüest, R.O. (2013) What
determines biogeographical ranges? Historical wanderings and ecological constraints in
the danthonioid grasses. *Journal of Biogeography*, **40**, 821–834.

Liu, C., White, M. & Newell, G. (2013) Selecting thresholds for the prediction of species
occurrence with presence-only data. *Journal of Biogeography*, **40**, 778–789.

McFadden, I.R., Sandel, B., Tsirogiannis, C., Morueta-Holme, N., Svenning, J.-C., Enquist, B.J.
& Kraft, N.J.B. (2019) Temperature shapes opposing latitudinal gradients of plant
taxonomic and phylogenetic β diversity. *Ecology Letters*, **22**, 1126–1135.

Morin, X. & Chuine, I. (2006) Niche breadth, competitive strength and range size of tree species:
a trade-off based framework to understand species distribution. *Ecology Letters*, **9**, 185–
195.

Morueta-Holme, N., Enquist, B.J., McGill, B.J., Boyle, B., Jørgensen, P.M., Ott, J.E., Peet, R.K.,
Šímová, I., Sloat, L.L., Thiers, B., Violle, C., Wiser, S.K., Dolins, S., Donoghue, J.C.,
Kraft, N.J.B., Regetz, J., Schildhauer, M., Spencer, N. & Svenning, J.-C. (2013) Habitat
area and climate stability determine geographical variation in plant species range sizes.
Ecology Letters, **16**, 1446–1454.

Naimi, B. & Araújo, M.B. (2016) sdm: a reproducible and extensible R platform for species
distribution modelling. *Ecography*, **39**, 368–375.

Niinemets, Ü. & Valladares, F. (2006) Tolerance to Shade, Drought, and Waterlogging of
Temperate Northern Hemisphere Trees and Shrubs. *Ecological Monographs*, **76**, 521–
547.

Nogués-Bravo, D., Pulido, F., Araújo, M.B., Diniz-Filho, J.A.F., García-Valdés, R., Kollmann,
J., Svenning, J.-C., Valladares, F. & Zavala, M.A. (2014) Phenotypic correlates of
potential range size and range filling in European trees. *Perspectives in Plant Ecology,*
Evolution and Systematics, **16**, 219–227.

Normand, S., Ricklefs, R.E., Skov, F., Bladt, J., Tackenberg, O. & Svenning, J.-C. (2011)
Postglacial migration supplements climate in determining plant species ranges in Europe.
Proceedings of the Royal Society B: Biological Sciences, **278**, 3644–3653.

Normand, S., Treier, U.A., Randin, C., Vittoz, P., Guisan, A. & Svenning, J.-C. (2009)

Importance of abiotic stress as a range-limit determinant for European plants: insights

from species responses to climatic gradients. *Global Ecology and Biogeography*, **18**,

437–449.

Olson, M.E., Soriano, D., Rosell, J.A., Anfodillo, T., Donoghue, M.J., Edwards, E.J., León-

Gómez, C., Dawson, T., Camarero Martínez, J.J., Castorena, M., Echeverría, A.,

Espinosa, C.I., Fajardo, A., Gazol, A., Isnard, S., Lima, R.S., Marcati, C.R. & Méndez-

Alonzo, R. (2018) Plant height and hydraulic vulnerability to drought and cold.

Proceedings of the National Academy of Sciences, **115**, 7551–7556.

Paul, J.R., Morton, C., Taylor, C.M. & Tonsor, S.J. (2009) Evolutionary Time for Dispersal

Limits the Extent but Not the Occupancy of Species' Potential Ranges in the Tropical

Plant Genus Psychotria (Rubiaceae). *The American Naturalist*, **173**, 188–199.

Phillips, S.J., Anderson, R.P. & Schapire, R.E. (2006) Maximum entropy modeling of species

geographic distributions. *Ecological Modelling*, **190**, 231–259.

Pither, J. (2003) Climate tolerance and interspecific variation in geographic range size.

Proceedings of the Royal Society of London. Series B: Biological Sciences, **270**, 475–

481.

Puglielli, G., Hutchings, M.J. & Laanisto, L. (2021a) The triangular space of abiotic stress

tolerance in woody species: a unified trade-off model. *New Phytologist*, **229**, 1354–1362.

Puglielli, G., Laanisto, L., Poorter, H. & Niinemets, Ü. (2021b) Global patterns of biomass

allocation in woody species with different tolerances of shade and drought: evidence for

multiple strategies. *New Phytologist*, **229**, 308–322.

Puglielli, G., Redondo-Gómez, S., Gratani, L. & Mateos-Naranjo, E. (2017) Highlighting the

differential role of leaf paraheliotropism in two Mediterranean Cistus species under

drought stress and well-watered conditions. *Journal of Plant Physiology*, **213**, 199–208.

Ricotta, C., Godefroid, S. & Rocchini, D. (2010) Patterns of native and exotic species richness in

the urban flora of Brussels: rejecting the 'rich get richer' model. *Biological Invasions*, **12**,

233–240.

Sack, L. (2004) Responses of temperate woody seedlings to shade and drought: do trade-offs

limit potential niche differentiation? *Oikos*, **107**, 110–127.

Schubert, M., Humphreys, A.M., Lindberg, C.L., Preston, J.C. & Fjellheim, S. (2020) *To Coldly Go Where No Grass has Gone Before: A Multidisciplinary Review of Cold Adaptation in Poaceae*. In Annual Plant Reviews online, J.A. Roberts (Ed.). https://doi.org/10.1002/9781119312994.apr0739.

Seliger, B.J., McGill, B.J., Svenning, J.-C. & Gill, J.L. (2021) Widespread underfilling of the potential ranges of North American trees. *Journal of Biogeography*, **48**, 359–371.

Sexton, J.P., Montiel, J., Shay, J.E., Stephens, M.R. & Slatyer, R.A. (2017) Evolution of Ecological Niche Breadth. *Annual Review of Ecology, Evolution, and Systematics*, **48**, 183–206.

Sheth, S.N., Morueta-Holme, N. & Angert, A.L. (2020) Determinants of geographic range size in plants. *New Phytologist*, **226**, 650–665.

Smith, S.A. & Brown, J.W. (2018) Constructing a broadly inclusive seed plant phylogeny. *American Journal of Botany*, **105**, 302–314.

Stevens, G.C. (1989) The Latitudinal Gradient in Geographical Range: How so Many Species Coexist in the Tropics. *The American Naturalist*, **133**, 240–256.

Sunday, J.M., Bates, A.E. & Dulvy, N.K. (2011) Global analysis of thermal tolerance and latitude in ectotherms. *Proceedings of the Royal Society B: Biological Sciences*, **278**, 1823–1830.

Svenning, J.-C., Normand, S. & Skov, F. (2008) Postglacial Dispersal Limitation of Widespread Forest Plant Species in Nemoral Europe. *Ecography*, **31**, 316–326.

Svenning, J.-C. & Skov, F. (2004) Limited filling of the potential range in European tree species. *Ecology Letters*, **7**, 565–573.

Title, P.O. & Bemmels, J.B. (2018) ENVIREM: an expanded set of bioclimatic and topographic variables increases flexibility and improves performance of ecological niche modeling. *Ecography*, **41**, 291–307.

Varela, S., Anderson, R.P., García-Valdés, R. & Fernández-González, F. (2014) Environmental filters reduce the effects of sampling bias and improve predictions of ecological niche models. *Ecography*, **37**, 1084–1091.

Wazen, N., Garavaglia, V., Picard, N., Besacier, C. & Fady, B. (2020) Distribution maps of twenty-four Mediterranean and European ecologically and economically important forest

tree species compiled from historical data collections. *Annals of Silvicultural Research*,

44, 95–101.

Weiser, M.D., Enquist, B.J., Boyle, B., Killeen, T.J., Jørgensen, P.M., Fonseca, G., Jennings,

M.D., Kerkhoff, A.J., Jr, T.E.L., Monteagudo, A., Vargas, M.P.N., Phillips, O.L.,

Swenson, N.G. & Martínez, R.V. (2007) Latitudinal patterns of range size and species

richness of New World woody plants. *Global Ecology and Biogeography*, **16**, 679–688.

Zanne, A.E., Pearse, W.D., Cornwell, W.K., McGlinn, D.J., Wright, I.J. & Uyeda, J.C. (2018)

Functional biogeography of angiosperms: life at the extremes. *New Phytologist*, **218**,

1697–1709.