The transcription factor Zeb2 drives differentiation of age-associated B cells

Shuangshuang Gu[1, 8], Xiaxia Han[1, 8], Chao Yao[6, 8], Huihua Ding[1], Jinsong

Zhang[6], Guojun Hou[1], Bo Qu[1], Haibo Zhou[1], Zhihua Ying[3], Zhizhong Ye[3], Jie

Qian[1], Qiang Guo[1], Sheng Chen[1], Carola G. Vinuesa[2, 7, *], Dai Dai[1, *], Nan Shen[1,

2, 3, 4, 5, 9,*]

[1] Shanghai Institute of Rheumatology, Shanghai Renji Hospital, Shanghai

Jiaotong University School of Medicine, Shanghai, 200001, China.

[2] Centre for Personalised Immunology (CACPI), Renji Hospital, School of

Medicine, Shanghai Jiao Tong University (SJTUSM), Shanghai, 200001,

China.

[3] Shenzhen Futian Hospital for Rheumatic Diseases, Shenzhen, 518000,

China.

[4] State Key Laboratory of Oncogenes and Related Genes, Shanghai Cancer

Institute, Renji Hospital, Shanghai Jiaotong University School of Medicine,

Shanghai, 200001, China.

[5] Center for Autoimmune Genomics and Etiology (CAGE) and Divisions of

Rheumatology, Cincinnati Children's Hospital Medical Center, Cincinnati OH

45229, USA.

[6] Shanghai Institute of Nutrition and Health, University of Chinese Academy of

Sciences, Chinese Academy of Sciences, Shanghai 200031, China.

[7] Centre for Personalised Immunology, John Curtin School of Medical

Research, Australian National University, 131 Garran Rd, Acton, 2601, ACT,

Australia.

[8] These authors contributed equally to the work

*Correspondence to:

Nan Shen: nanshensibs@gmail.com

Dai Dai: daidaisibs@163.com

Carola G. Vinuesa: carola.vinuesa@anu.edu.au

Abstract

Age-associated B cells (ABCs) accumulate after infection, during aging, and in systemic autoimmunity, contributing to pathogenesis in the latter[1]. To date, their precise ontogeny and whether they represent a B cell subset distinct from the other major effector subsets - plasma cells and memory B cells - remain unknown. We asked whether ABC formation requires the expression of a unique transcription factor. Here, we used single-cell RNA sequencing to identify differentially-expressed transcription factors shared by ABCs from both lupus patient and lupus-prone mice. Using an arrayed CRISPR screen, we show Zeb2 is essential for ABC differentiation in vitro. Selective deficiency of Zeb2 in B cells also leads to impaired ABC formation in vivo. Zeb2 drove ABC specification by directly binding the promoters of ABC signature genes including *Itgax* and *Itgam* encoding for CD11c and CD11b respectively. Zeb2 also targeted molecules involved in cytokine receptor signaling via the JAK/STAT pathway including *Il21r* and *Socs5*. Administration of the approved JAK/STAT inhibitor tofacitinib effectively ameliorated ABC accumulation in mice and patients with autoimmunity. This study reveals ABCs represent a distinct B cell subset dependent on Zeb2 expression, and provides a rationale for the use of JAK/STAT inhibitors to treat ABC-mediated autoimmune diseases.

Main

Age-associated B cells (ABC) characterized by high expression of CD11c and T-bet are found at increased numbers in aged female mice, infection models and multiple autoimmune diseases[1]. This novel B cell subset has been given various names including ABCs, CD11c+ B cells, CD11c+T-bet+ B cells, CD21−CD23−B cells, Be1 cells, double negative B cells (DN), DN2 or atypical memory B cells[2-10]. ABCs or ABC-like B cells identified as CD11c+ T-bet+CD11b+CD21−CD23−CD86hiMHCIIhiCD95+ in mice[2,8] and CD11c+T-bet+CD27−IgD−CD38−CD21−CD95hiFCRL5+ in humans[4,6] will be referred to as ABCs henceforth, following the suggestion of a recent review[1]. In autoimmune settings, these B cells are enriched for autoantibody specificities. It is widely accepted that these cells are antigen-experienced and it has been suggested that they can rapidly differentiate into antibody-secreting cells (ASCs) upon antigen reencounter or innate stimulation[11,12].

The transcriptional factor T-bet is highly expressed in ABCs and it is required for autoantibody production in lupus-prone mice[3,13]. T-bet has been put forward as a functional regulator of ABC differentiation by putatively binding *Il21r* promoters in humans[9] and by suppressing an IFN-γ-induced inflammatory network that supports ABC terminal differentiation to ASCs in mice[10]. However, T-bet is dispensable for ABC formation or maintenance of ABC features[14,15] arguing against it being the master regulator of ABCs. In search for the TF essential for ABC formation, we screened all TFs expressed by these cells and identified Zeb2 as the key regulator required for ABC specification and differentiation in mouse and human.

Differentially-expressed TFs in autoimmune ABCs

To gain insights into the nature of ABCs, we sorted peripheral B cells from a patient with new-onset SLE (Supplementary Table 1) and performed droplet-based scRNA-seq. Seven distinct clusters containing from 94 to 1640 cells

were revealed by unsupervised clustering with a two-dimensional uniform

manifold approximation and projection (UMAP) (Fig. 1a and Extended Data Fig.

1b). These clusters were assigned to known peripheral B cell subsets including

transitional B cells (TrB), naïve B cells (NavB), activated naïve B cells (aNavB),

ABCs, memory B cells (MemB), plasmablasts (PB) and plasma cells (PC) by

comparing DEGs with established markers and previously described gene

expression profiles[6,16,17]. Landmark genes (*IGHD*, *CD27*, *CD38*, *CD24*, *CD9*,

CD79B, *VPREB3*, *MZB1*, *PRDM1*, *XBP1*, *IRF4*, *PCNA*, *MKI67*, *IGHG1*, *IGHF3*,

IGKC, *JCHAIN*) were used to identify clusters (Extended Data Fig. 1a, c).

ABCs preferentially expressed genes encoding key surface markers: *ITGAX*,

CD19, *MS4A1*, *CD86*, *FCRLA*, *FCRL3/5* and *FCGRB2,* and they lacked *CR2*,

FCER2, *CD27* and *IGHD* (Fig. 1b). Amongst transcripts encoding chemokine

receptors, *CXCR3* was highly expressed while *CXCR4* and *CXCR5* were

suppressed. Besides these, we identified 43 differentially-expressed TFs in

ABCs: 27 upregulated (*HOPX*, *ZEB2*, *POU2F2*, *TOX*, *KLF3*, *ARID3A*, *TBX21*,

LITAF, *PLEK*, *ZBTB32*, *TFEB*, *NR4A2*, *NFATC2*, *SREBF2*, *TOX2*, *JUN*B,

ZNF331, *BATF*, *KLF6*, *TFEC*, *CREM*, *BHLHE40*, *SOX5*, *JUN*, *JAZF1*, *FOXP4*,

ARID5A) and 16 downregulated (*HIF1A*, *MBD4*, *ETS1*, *SP140L*, *FLI1*, *ZNF92*,

ZFP36L1, *PLSCR1*, *AFF3*, *IRF8*, *FOS*, *MYC*, *FOXP1*, *ZFP36L2*, *IRF1*, *BACH2*)

(Fig. 1c).

We previously reported T-bet⁺CD11c⁺ B cells accumulate in Bm12-induced

lupus and contribute to autoantibody production[3]. Whether Bm12-induced T-

bet⁺CD11c⁺ B cells represent *bona fide* ABCs has not been described. Flow

cytometric staining revealed that CD11c⁺ B cells from Bm12-induced mice

exhibited ABC-specific features including a hyperactivated state

(CD19^{hi}CD86^{hi}CD62L^{low}CD44^{hi}CD95^{hi}MHCII^{hi}), characteristic markers

(CD11c⁺T-bet^{hi} CD11b^{hi}), specific lineage association (CD23⁻CD21⁻CD93⁻

CD5^{int}CD138⁻IgD^{low}IgM^{int}) and elevated expression of chemokine receptors

CXCR3 and CXCR4 (Fig. 1d). We used the surface marker CD21 to exclude

the small fraction of naive CD11c⁺ B cells. Consistent with previous reports,

Bm12-indued CD11c⁺ CD21⁻ B cells expressed high amounts of *Cxcl9* and

Cxcl10 transcripts (Fig. 1e). We then used qPCR to validate 40 differentially-

expressed TFs identified in our human scRNA-seq data excluding *ZNF331*,

SP140L and *ZNF92*, which lack orthologs in mice (Fig.1f). Amongst these, 13

upregulated TFs (*Tbx21*, *Zeb2*, *Plek*, *Litaf*, *Tfeb*, *Nfatc2*, *Zbtb32*, *Srebf2*, *Jazf1*,

Jun, *Sox5*, *Tfec*, *Batf*) and 3 downregulated TFs (*Ets1*, *Mbd4*, *Fli1*) were

identically regulated in human and mouse ABCs and were therefore considered

potential transcriptional regulators of ABC differentiation.

122 Screen for TFs directing mouse ABC differentiation

ABCs can be generated in vitro[10,18] and TLR signaling, BCR engagement, T

cell help and the T cell-derived cytokines IFN-γ and IL-21 have been shown to

be important for ABC formation in ageing, infection and autoimmune

settings[2,5,15,18,19]. We investigated the optimal combination of stimuli for efficient

ABC induction within 3 days of culture (Fig. 2a and Extended Data Fig. 2a-d).

BCR engagement with anti-CD40 and R848 promoted B cell viability (Extended

Data Fig. 2a, b). IL-21 induced CD11c and T-bet expression while IFN-γ only

facilitated T-bet expression (Extended Data Fig. 2d, e). IL-21 and IFN-γ

signaling could enhance *Cd86*, *Cxcr3* and *Cxcl10* expression (Extended Data

Fig. 2e). We therefore used a cocktail of anti-IgM, R848, anti-CD40, IL-21 and

IFN-γ for in vitro ABC induction.

To screen for functional transcriptional regulators of ABC differentiation, B cells

from Cas9 transgenic mice and congenic CD45.1 mice were cocultured in a 4:1

ratio and activated with anti-IgM, anti-CD40 and R848 for 24 hours. B cells

expressing CD45.1 and lacking Cas9 were transduced with sgRNA plasmids

and served as controls in these co-cultures. Activated B cells were then

retrovirally infected with the sgRNA plasmids co-expressed with the blue

fluorescent protein (BFP), and cultured for a further 3 days with the ABC

differentiation cocktail (Fig. 2b). We then compared the ratio between live

BFP$^+$CD45.1$^-$ (Cas9$^+$, edited) and BFP$^+$CD45.1$^+$ (Cas9$^-$, unedited) CD11c$^+$T-

bet$^+$ ABCs (Extended Data Fig. 3a). To validate our genome editing tools we

introduced a sgRNA sequence specific for *Itgax* and showed effective blockade

of CD11c$^+$T-bet$^+$ ABC cell formation from Cas9$^+$ B cells (Extended Data Fig. 3b).

We then screened the 16 candidate TFs and identified 7 (T-bet, Batf, Ets1, Fli1,

Jun, Srebf2, Zeb2) that could significantly (P<0.05) alter ABC formation (Fig. 2c

and Extended Data Fig. 3c). Except for Zeb2, ablation of the other 6 TFs

predominantly influenced cell viability (Extended Data Fig. 3d). Two different

sgRNAs targeting *Zeb2* in separate Cas9$^+$ B cell cultures lead to reduced ABC

formation (Fig. 2d), which excluded an off-target editing effect. Taken together,

we identify 7 functional TFs required for mouse ABC differentiation, with one of

them, Zeb2, acting by means other than promoting cell survival.

Functional assessment of candidate TFs in human ABCs

We next set out to determine whether besides Zeb2, any of the other 6 TFs was

required for human ABC lineage specification, using an established in vitro ABC

induction protocol used in human studies[4,6,9] (Fig. 2e). Synergistic stimulation

with anti-Ig, R848, BAFF, IL-2, anti-CD40, IL-21 and IFN-γ increased IgD$^-$CD27$^-$

DN and CD11c$^+$ T-bet$^+$ ABC percentages with a high B cell survival rate (>70%)

(Extended Data Fig. 4a-d). Compared with naïve and switched B cells, in vitro-

derived DN B cells contained the highest frequency of CD11c$^+$T-bet$^+$ cells

(Extended Data Fig. 4e, f).

We next transduced Cas9-guide RNA ribonucleoprotein (RNP) complexes by

electroporation to edit genes in primary human B cells. Using a sgRNA

sequence targeting *CD19* as a positive control, we achieved extremely efficient

gene disruption (Extended Data Fig. 5a). Primary B cells were isolated and

purified from the blood of healthy individuals. After activation for one day, cells

were electroporated with RNP and treated with the ABC differentiation cocktail

for another 3 days (Extended Data Fig. 5b). All of the 7 TFs identified in mice

led to changes in human ABC's when individually ablated (P<0.05) (Fig. 2f,g

and Extended Data Fig. 5c). Two different guide RNAs targeting *ZEB2* achieved

reduced T-bet expression (Fig. 2h). Gene editing of *ETS1*, *FLI1* and *JUN* had

opposite effects on human ABCs and were excluded. Ablation of *TBX21*, *ZEB2*,

BATF and *SREBF2* dampened ABC differentiation in both human and mouse

B cells. However, *TBX21*, *BATF* and *SREBF2* deficiency also impaired human

cell viability (Extended Data Fig. 5d, e). As a key regulator of lipid metabolism,

SREBF2 is required to supply lipids for membrane synthesis and has been

reported to influence T cell survival and the GC response[20,21]. Although there is

no published evidence that T-bet and BATF control B cell proliferation or death,

both TFs have been implicated in B lymphoproliferative disorders[22,23]. After

exclusion of the genes predominantly affecting cell viability, Zeb2 emerged as

the single TF with consistent regulatory effects in human and mouse ABC

formation and therefore the key ABC transcriptional regulator.

Zeb2 is required for ABC formation

Zeb2 belongs to the zinc-finger E homeobox-binding protein family and is

known to drive epithelial to mesenchymal transition in cancer progression[24]. In

the immune system, Zeb2 has been reported to be expressed in dendritic cells

(DC), macrophages, B, T and NK cells and participate in transcriptional

networks involved in cell differentiation, maintenance and function[25-33]. To

further explore how Zeb2 deficiency affects ABC formation we generated mice

selectively lacking Zeb2 in B cells by crossing Zeb2 floxed mice with CD19-cre

mice (Extended Data Fig. 6a-c). We first isolated B cells from Zeb2[f/f]CD19[cre/+]

conditional knockout mice (CKO) and Zeb2[f/f] littermate control mice (Ctrl) and

induced in vitro ABC differentiation for 3 days. Zeb2 deficiency significantly

impaired ABC differentiation (Fig. 3a). Zeb2 deficient B cells also exhibited

decreased CD11c and T-bet expression (Extended Data Fig. 6d, e).

We then investigated the *in vivo* effects of Zeb2 ablation in B cells on ABC

formation, using the Bm12-induced lupus model. As early as 2 weeks after

adoptive transfer of CD4+ T cell from Bm12 mice, the proportion of ABC cells

identified as CD11c⁺CD11b⁺T-bet⁺CD21⁻ or CD21⁻CD23⁻CD11c⁺CD11b⁺ was

significantly reduced in Zeb2 CKO mice (Fig. 3b, c). The absolute number of

ABCs was reduced by over 50% in Zeb2 CKO mice. By contrast, no differences

in the numbers of GC B cells or plasma cells were seen in Zeb2 CKO mice

(Extended Data Fig. 6f, g), reflecting a specific regulatory role of Zeb2 in ABC

formation. These results confirm that Zeb2 deficiency in B cells leads to

impaired ABC formation both in vitro and in vivo.

Zeb2 directly regulates ABC signature genes

The mechanisms involved in Zeb-mediated gene regulation are broad and cell-

context specific[25,26,28,31]. To investigate the consequences of Zeb2 regulation of

gene transcription we performed RNA sequencing on sorted B cells with

reduced Zeb2 (sg-*Zeb2* (CD19⁺GFP⁺BFP⁺)) or T-bet (sg-*Tbx21)* expression 4

days after in vitro activation, retroviral infection and differentiation (Extended

Data Fig. 7a). We found sg-*Zeb2* B cells expressed less *Il7r* consistent with

reports of Zeb2-mediated regulation of IL-7R expression during pro-pre B cell

development[30] (Fig. 4a). Comparative analysis identified that ABC signature

genes *Itgax*, *Tbx21*, *Itgam*, *Cxcr4* and *Cxcl10* were downregulated while *Fcer2a*

(encoding CD23a) was upregulated in sg-*Zeb2* B cells (Fig. 4a, b). In vitro-

derived ABCs from Zeb2 CKO mice also lacked expression of ABC signature

genes including *Itgax*, *Itgam* and *Cxcr4* (Extended Data Fig. 7e). T-bet ablation

also impaired some aspects of the ABC phenotype (Fig. 4b and Extended Data

Fig. 7b). Principal component analysis (PCA) distinctly separated sg-*Zeb2* and

sg-*Tbx21* B cells suggesting different regulatory roles for these two TFs during

ABC differentiation (Extended Data Fig. 7c). Clustering by signature genes

revealed Zeb2 ablation resulted in more severe impairment of ABC identity

compared with T-bet ablation (Fig. 4b).

To determine the direct gene targets of Zeb2 regulation, we performed cleavage

under targets and tagmentation (Cut & tag) followed by sequencing the

enriched DNA fragments[34]. Cut & tag-seq signals were compared among Zeb2

specific, T-bet specific and IgG peaks. Zeb2 signals were enriched in the

transcriptional start sites (TSS) especially within promoter regions (Extended

Data Fig. 8a, b). Zeb2 specific peaks were matched to motifs of Gata3, Fosl2

and Zeb2 (Extended Data Fig. 8c), consistent with Zeb2 Chip-seq data in cell

lines from the Encode project (Extended Data Fig. 8e). We found the

transcriptional regulatory regions of *Itgax* and *Itagm* were enriched in Zeb2-

specific peaks and these loci contained a Zeb2 binding motif (Fig. 4c). These

results suggest that Zeb2 orchestrates ABC specification by directly targeting

ABC signature genes including *Itgax* and *Itgam*.

Zeb2-Jak-Stat axis controls ABC differentiation

KEGG analysis revealed that the top pathway of downregulated genes in sg-

Zeb2 cells was related to Jak-Stat signaling (Extended Data Fig. 7d). Signaling

landscape changes downstream of cytokine receptor signaling via Jak-Stat

were annotated (Fig. 4a) and genes involved in Jak-Stat signaling with

significant changes were listed in a heatmap (Fig. 4d). Gene set enrichment

analysis (GSEA) also revealed sg-Zeb2 B cells had altered Jak-Stat signaling

(Fig. 4e), which is known to be critical for ABC differentiation[1]. We validated

differentially-expressed genes involved in Jak-Stat signaling using in vitro-

derived ABCs from Zeb2 CKO mouse B cells (Extended Data Fig. 7e). There

was a positive correlation between *IL21R* and *ZEB2* expression in B cells from

SLE patients (Extended Data Fig. 7f). We then compared DEGs with Zeb2

specific signals from cut-tag to narrow down the key direct targets in Jak-Stat

pathways. The promoters of *Socs5* and *Il21r* were enriched in the Zeb2-specific

group. These loci also contained Zeb2 binding motifs (Extended Data Fig. 8f).

SOCS5 is a negative regulator of IL-4 / STAT6 signaling, which dampens ABC

formation[18]. IL-21 signaling via IL21R is critical for ABC differentiation[4,18]. These

results suggest that Zeb2 directly regulates important components of Jak-Stat

signaling including *Socs5* and *Il21r* to drive ABC differentiation.

Jak-Stat inhibitors have proven to be a safe and efficient strategy to dampen

intracellular cytokine signaling[35]. Baricitinib, tofacitinib, upadacitinib and

filgotinib have been approved for oral therapy in rheumatoid arthritis (RA) and

are also commonly prescribed off-label in other autoimmune diseases including

SLE, inflammatory bowel disease and psoriasis[36]. We therefore investigated

the effects of Jak-Stat inhibitors on ABC formation. Baricitinib or tofacitinib

impaired mouse in vitro ABC differentiation in a dose-dependent manner (Fig.

4f and Extended Data Fig. 9a-d). Oral administration of tofacitinib reduced

splenomegaly and autoantibody titers in the Bm12-induced lupus model

(Extended Data Fig. 9e-g). Notably, accumulation of ABCs was effectively

ameliorated by tofacitinib (Fig. 4g). Accordingly, human ABC differentiation in

vitro was also blocked by baricitinib and tofacitinib (Fig 4h and Extended Data

Fig. 9h). Finally, we investigated the *in vivo* effects of tofacitinib on ABC

differentiation in RA patients. The frequency of human ABCs (DN2) had

decreased 4 weeks after treatment (Fig. 4i). Tofacitinib also ameliorated

systemic inflammation lowering C-reactive protein (CRP) (Extended Data Fig.

9i). These results clearly demonstrate that targeting the Jak-Stat pathway is an

efficient strategy to block ABCs in mice and humans and provide a rationale for

the use of Jak-Stat inhibition therapy in ABC-mediated autoimmunity.

In summary, our results have identified Zeb2 as an essential TF that regulates

human and mouse ABC differentiation by targeting Jak-Stat signaling.

Surprisingly, Zeb2 rather than T-bet, determines ABC specification. Mice and

humans have ABC subsets that share a similar phenotype and exert pathogenic

roles in autoimmunity[1]. It is therefore not surprising that both require Zeb2 for

ABC development. Although our TF screen takes advantage of an autoimmune

model for rapid ABC development, our results are likely to apply to other

settings as ABCs have been shown to be comparable in the context of ageing,

infection and autoimmunity.

References

Cancro, M. P. Age-Associated B Cells. *Annu Rev Immunol* **38**, 315-340, doi:10.1146/annurev-immunol-092419-031130 (2020).

Rubtsov, A. V. *et al.* Toll-like receptor 7 (TLR7)-driven accumulation of a novel CD11c(+) B-cell population is important for the development of autoimmunity. *Blood* **118**, 1305-1315, doi:10.1182/blood-2011-01-331462 (2011).

Liu, Y. *et al.* T-bet+CD11c+ B cells are critical for antichromatin immunoglobulin G production in the development of lupus. *Arthritis Research & Therapy* **19**, doi:10.1186/s13075-017-1438-2 (2017).

Wang, S. *et al.* IL-21 drives expansion and plasma cell differentiation of autoreactive CD11c(hi)T-bet(+) B cells in SLE. *Nat Commun* **9**, 1758, doi:10.1038/s41467-018-03750-7 (2018).

Manni, M. *et al.* Regulation of age-associated B cells by IRF5 in systemic autoimmunity. *Nat Immunol* **19**, 407-419, doi:10.1038/s41590-018-0056-8 (2018).

Jenks, S. A. *et al.* Distinct Effector B Cells Induced by Unregulated Toll-like Receptor 7 Contribute to Pathogenic Responses in Systemic Lupus Erythematosus. *Immunity* **49**, 725-739 e726, doi:10.1016/j.immuni.2018.08.015 (2018).

Wu, C. *et al.* Lupus-associated atypical memory B cells are mTORC1-hyperactivated and functionally dysregulated. *Ann Rheum Dis* **78**, 1090-1100, doi:10.1136/annrheumdis-2019-215039 (2019).

Hao, Y., O'Neill, P., Naradikian, M. S., Scholz, J. L. & Cancro, M. P. A B-cell subset uniquely responsive to innate stimuli accumulates in aged mice. *Blood* **118**, 1294-1304, doi:10.1182/blood-2011-01-330530 (2011).

Zumaquero, E. *et al.* IFNgamma induces epigenetic programming of human T-bet(hi) B cells and promotes TLR7/8 and IL-21 induced differentiation. *Elife* **8**, doi:10.7554/eLife.41641 (2019).

Stone, S. L. *et al.* T-bet Transcription Factor Promotes Antibody-Secreting Cell Differentiation by Limiting the Inflammatory Effects of IFN-gamma on B Cells. *Immunity* **50**, 1172-1187 e1177, doi:10.1016/j.immuni.2019.04.004 (2019).

Myles, A., Sanz, I. & Cancro, M. P. T-bet(+) B cells: A common denominator in protective and autoreactive antibody responses? *Curr Opin Immunol* **57**, 40-45, doi:10.1016/j.coi.2019.01.002 (2019).

Rubtsov, A. V., Marrack, P. & Rubtsova, K. T-bet expressing B cells - Novel target for autoimmune therapies? *Cell Immunol* **321**, 35-39, doi:10.1016/j.cellimm.2017.04.011 (2017).

Rubtsova, K. *et al.* B cells expressing the transcription factor T-bet drive lupus-like autoimmunity. *J Clin Invest* **127**, 1392-1404, doi:10.1172/JCI91250 (2017).

Du, S. W., Arkatkar, T., Jacobs, H. M., Rawlings, D. J. & Jackson, S. W. Generation of functional murine CD11c(+) age-associated B cells in the absence of B cell T-bet expression. *Eur J Immunol* **49**, 170-178, doi:10.1002/eji.201847641 (2019).

Levack, R. C., Newell, K. L., Popescu, M., Cabrera-Martinez, B. & Winslow, G. M. CD11c(+) T-bet(+) B Cells Require IL-21 and IFN-

gamma from Type 1 T Follicular Helper Cells and Intrinsic Bcl-6
Expression but Develop Normally in the Absence of T-bet. *J Immunol*
205, 1050-1058, doi:10.4049/jimmunol.2000206 (2020).

16 Kaminski, D. A., Wei, C., Qian, Y., Rosenberg, A. F. & Sanz, I.
Advances in human B cell phenotypic profiling. *Front Immunol* **3**, 302,
doi:10.3389/fimmu.2012.00302 (2012).

17 Glass, D. R. *et al.* An Integrated Multi-omic Single-Cell Atlas of Human
B Cell Identity. *Immunity* **53**, 217-232.e215,
doi:10.1016/j.immuni.2020.06.013 (2020).

18 Naradikian, M. S. *et al.* Cutting Edge: IL-4, IL-21, and IFN-gamma
Interact To Govern T-bet and CD11c Expression in TLR-Activated B
Cells. *J Immunol* **197**, 1023-1028, doi:10.4049/jimmunol.1600522
(2016).

19 Russell Knode, L. M. *et al.* Age-Associated B Cells Express a Diverse
Repertoire of VH and Vkappa Genes with Somatic Hypermutation. *J*
Immunol **198**, 1921-1927, doi:10.4049/jimmunol.1601106 (2017).

20 Kidani, Y. *et al.* Sterol regulatory element-binding proteins are essential
for the metabolic programming of effector T cells and adaptive
immunity. *Nat Immunol* **14**, 489-499, doi:10.1038/ni.2570 (2013).

21 Ricker, E. *et al.* Serine-threonine kinase ROCK2 regulates germinal
center B cell positioning and cholesterol biosynthesis. *J Clin Invest*
130, 3654-3670, doi:10.1172/jci132414 (2020).

22 Dorfman, D. M., Hwang, E. S., Shahsafaei, A. & Glimcher, L. H. T-bet,
a T-Cell–Associated Transcription Factor, Is Expressed in a Subset of
B-Cell Lymphoproliferative Disorders. *American Journal of Clinical*
Pathology **122**, 292-297, doi:10.1309/aqq2dvm75dvy0pwp (2004).

23 Care, M. A. *et al.* SPIB and BATF provide alternate determinants of
IRF4 occupancy in diffuse large B-cell lymphoma linked to disease
heterogeneity. *Nucleic Acids Res* **42**, 7591-7610,
doi:10.1093/nar/gku451 (2014).

24 Vandewalle, C. *et al.* SIP1/ZEB2 induces EMT by repressing genes of
different epithelial cell-cell junctions. *Nucleic Acids Res* **33**, 6566-6578,
doi:10.1093/nar/gki965 (2005).

25 Bagadia, P. *et al.* An Nfil3-Zeb2-Id2 pathway imposes Irf8 enhancer
switching during cDC1 development. *Nat Immunol* **20**, 1174-1185,
doi:10.1038/s41590-019-0449-3 (2019).

26 Scott, C. L. *et al.* The transcription factor Zeb2 regulates development
of conventional and plasmacytoid DCs by repressing Id2. *J Exp Med*
213, 897-911, doi:10.1084/jem.20151715 (2016).

27 Wu, X. *et al.* Transcription factor Zeb2 regulates commitment to
plasmacytoid dendritic cell and monocyte fate. *Proc Natl Acad Sci U S*
A **113**, 14775-14780, doi:10.1073/pnas.1611408114 (2016).

28 Dominguez, C. X. *et al.* The transcription factors ZEB2 and T-bet
cooperate to program cytotoxic T cell terminal differentiation in
response to LCMV viral infection. *J Exp Med* **212**, 2041-2056,
doi:10.1084/jem.20150186 (2015).

29 Omilusik, K. D. *et al.* Transcriptional repressor ZEB2 promotes terminal
differentiation of CD8+ effector and memory T cell populations during
infection. *J Exp Med* **212**, 2027-2039, doi:10.1084/jem.20150194
(2015).

30 Li, J. *et al.* The EMT transcription factor Zeb2 controls adult murine
hematopoietic differentiation by regulating cytokine signaling. *Blood*
129, 460-472, doi:10.1182/blood-2016-05-714659 (2017).
31 Scott, C. L. *et al.* The Transcription Factor ZEB2 Is Required to
Maintain the Tissue-Specific Identities of Macrophages. *Immunity* **49**,
312-325 e315, doi:10.1016/j.immuni.2018.07.004 (2018).
32 Collins, P. L. *et al.* Gene Regulatory Programs Conferring Phenotypic
Identities to Human NK Cells. *Cell* **176**, 348-360 e312,
doi:10.1016/j.cell.2018.11.045 (2019).
33 Huang, X. *et al.* Differential usage of transcriptional repressor Zeb2
enhancers distinguishes adult and embryonic hematopoiesis.
Immunity, doi:10.1016/j.immuni.2021.04.015 (2021).
34 Kaya-Okur, H. S. *et al.* CUT&Tag for efficient epigenomic profiling of
small samples and single cells. *Nat Commun* **10**, 1930,
doi:10.1038/s41467-019-09982-5 (2019).
35 O'Shea, J. J., Kontzias, A., Yamaoka, K., Tanaka, Y. & Laurence, A.
Janus kinase inhibitors in autoimmune diseases. *Ann Rheum Dis* **72**
Suppl 2, ii111-115, doi:10.1136/annrheumdis-2012-202576 (2013).
36 Traves, P. G. *et al.* JAK selectivity and the implications for clinical
inhibition of pharmacodynamic cytokine signalling by filgotinib,
upadacitinib, tofacitinib and baricitinib. *Ann Rheum Dis*,
doi:10.1136/annrheumdis-2020-219012 (2021).
37 Klarquist, J. & Janssen, E. M. The bm12 Inducible Model of Systemic
Lupus Erythematosus (SLE) in C57BL/6 Mice. *J Vis Exp*, e53319,
doi:10.3791/53319 (2015).
38 Cossarizza, A. *et al.* Guidelines for the use of flow cytometry and cell
sorting in immunological studies (second edition). *Eur J Immunol* **49**,
1457-1973, doi:10.1002/eji.201970107 (2019).
39 Heinen, A. P. *et al.* Improved method to retain cytosolic reporter protein
fluorescence while staining for nuclear proteins. *Cytometry A* **85**, 621-
627, doi:10.1002/cyto.a.22451 (2014).
40 Hsiau, T. *et al.* Inference of CRISPR edits from Sanger trace data.
BioRxiv, doi:10.1101/251082 (2019).
41 Zheng, G. X. *et al.* Massively parallel digital transcriptional profiling of
single cells. *Nat Commun* **8**, 14049, doi:10.1038/ncomms14049 (2017).
42 Stuart, T. *et al.* Comprehensive Integration of Single-Cell Data. *Cell*
177, 1888-1902.e1821, doi:10.1016/j.cell.2019.05.031 (2019).
43 Kim, D., Langmead, B. & Salzberg, S. L. HISAT: a fast spliced aligner
with low memory requirements. *Nat Methods* **12**, 357-360,
doi:10.1038/nmeth.3317 (2015).
44 Anders, S., Pyl, P. T. & Huber, W. HTSeq--a Python framework to work
with high-throughput sequencing data. *Bioinformatics* **31**, 166-169,
doi:10.1093/bioinformatics/btu638 (2015).
45 Henikoff, S., Henikoff, J. G., Kaya-Okur, H. S. & Ahmad, K. Efficient
chromatin accessibility mapping in situ by nucleosome-tethered
tagmentation. *Elife* **9**, doi:10.7554/eLife.63274 (2020).
46 Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold
change and dispersion for RNA-seq data with DESeq2. *Genome Biol*
15, 550, doi:10.1186/s13059-014-0550-8 (2014).
47 Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with

Bowtie 2. *Nat Methods* **9**, 357-359, doi:10.1038/nmeth.1923 (2012).
48   Ramírez, F. *et al.* deepTools2: a next generation web server for deep-
sequencing data analysis. *Nucleic Acids Res* **44**, W160-165,
doi:10.1093/nar/gkw257 (2016).
49   Zhang, Y. *et al.* Model-based analysis of ChIP-Seq (MACS). *Genome*
Biol **9**, R137, doi:10.1186/gb-2008-9-9-r137 (2008).

Acknowledgments

This study was supported by grants from National Natural Science Foundation of China (31630021, 31930037, 82071843 and 81901637), National Human Genetic Resources Sharing Service Platform (2005DKA21300), Shanghai Municipal Key Medical Center Construction Project (2017ZZ01024-002), Shenzhen Science and Technology Project (JCYJ20180504170414637 and JCYJ20180302145033769), Shenzhen Futian Public Welfare Scientific Research Project (FTWS2018005), and Sanming Project of Medicine in Shenzhen (SZSM201602087).

Author contributions

Conceptualization: N.S., D.D., C.G.V.; Methodology: D.D., S.G.; Formal analysis: D.D.; Experiment conduction: S.G., D.D., X.H., J.Z., G.H., B.Q., H.Z.; Resources: H.D., Q.G., S.C., Z.Y., J.Q., Z,Y., N.S.; Bioinfomatic analysis: C.Y., D.D.; Writing - Original Draft: D.D.; Writing - Review & Editing: C.G.V., D.D., N.S.; Supervision: N.S., D.D.; Funding acquisition: N.S.

Data availability

The sc-RNAseq, RNAseq and cut-tag sequencing data have been deposited in ArrayExpress database and the accession code will be provided prior to publication. Source data are available upon request.

Competing Interests

The authors declare no competing interests.

Additional information

Methods
Extended Data Figs. 1 to 9
Figure legends for Extended Data Figs. 1 to 9
Supplementary Tables 1-6

Methods

Mice and Bm12 cGVHD lupus models.

The C57BL/6J (JAX000664), Bm12 (JAX001162), Cd19-Cre (JAX006785), B6 Cd45.1(JAX002014) and Cas9-EGFP (JAX026175) mice were obtained from Jackson Laboratory. Zeb2^{flox} mice (T015122) were obtained from Gempharmatech (Nanjing). All the mouse strains were on C57BL/6 background and maintained under specific pathogen free (SPF) conditions in Renji Hospital animal facilities. All procedures involving animals were approved by Institutional Animal Care and Use Committee at Renji Hospital of Shanghai Jiaotong University School of Medicine (SJTUSM).

Bm12-cGVHD lupus model was induced following JOVE protocol[37]. Briefly, 8-12-week-old mice were intraperitoneally injected with 10 million CD4$^+$ T cells (Miltenyi Biotec, Cat#130-104-454) from Bm12 mice purified by magnetic cell sorting and mice were sacrificed for analysis at indicated time points. For Jak-Stat inhibitor treatment experiment, mice were orally gavage with 30 mg/kg tofacitinib (Selleckchem, Cat#S2789) for 2 weeks.

Patients subject

All patients were recruited from Renji Hospital. SLE patients met American College of Rheumatology criteria for SLE. Demographic and clinical information is listed (Supplementary Table 1 for scRNAseq, Supplementary Table 2 for detection of *ZEB2* and *IL-21R*). RA patients met the 2010 American College of Rheumatology criteria for RA. Clinical and laboratory data of RA patients were measured at baseline and at week 4 after tofacitinib treatment. Demographic and clinical information is listed before and after treatment (Supplementary Table 3). All procedures were fully consent under ethical and safe protocols approved by the Review Board at Renji Hospital of SJTUSM.

Cell isolation, flow cytometry and sorting.

Single cell suspensions of splenocytes were subjected to red blood cell lysis

and filtered through 70 um nylon mesh. Cells were stained for surface markers

or intracellular proteins under instructions[38]. For in vitro differentiation

experiment, B cells transfected sgRNA plasmids (BFP) were stained with

surface markers and then fixed by 2% fresh prepared formaldehyde in PBS for

5 minutes to allow efficient TF and cytokine staining and retain FP which were

optimized from previous protocol[39]. For intracellular T-bet staining, cells were

fixed and permeabilized after surface staining using Foxp3 Transcription Factor

Staining Buffer Set (ThermoFisher, Cat#00-5523-00). Samples were acquired

using LSRFortessa (BD Biosciences) and analyzed using FlowJo software

(Tree Star) .

For sorting experiment, splenic B cells were enriched by magnetic microparticle

purification using mouse B Cell Isolation Kit (Miltenyi Biotec, Cat#130-090-862)

and then further sorted by gating CD11c$^+$ B cells (CD19$^+$CD11c$^+$CD21$^-$) and

CD11c$^-$ B cells (CD19$^+$CD11c$^-$CD21$^+$) using FACSAria II (BD Biosciences).

Detailed information of antibodies were listed in Supplementary Table 4

In vitro induction of ABC and ASC.

Mouse splenic B cells were purified by negative selection and cultured in RPMI-

1640 medium with 10% fetal bovine serum, HEPES, non-essential amino acids

(NEAA), glutamine, sodium pyruvate, β-Mercaptoethanol and

penicillin/streptomycin. ABCs were induced in the presence of R848 (500 ng/ml,

Invivogen, Cat#tlrl-r848-5), anti-CD40 (1 ug/ml, Biolegend, Cat#102812), anti-

IgM F(ab')$_2$ (1 ug/ml, Jackson Immunoresearch, Cat#115-006-020), IL-21 (50

ng/ml, Peprotech, Cat#210-21), IFN-γ (1, 5 or 10ng/ml, Biotech, Cat#485-MI-

100) for 3 days. A similar procedure was done by replacing IFN-γ with IL-4 (10

ng/ml, Peprotech, Cat#214-14). To induce ASCs, B cells derived above were

transferred into new well and cultured for additional 4 days with indicated

stimulation cocktails.

Human PBMCs were isolated by density gradient centrifugation using Ficoll-

Paque (GE Healthcare, Cat#17-5442-03).   B cells were purified by negative

selection (Miltenyi Biotec, Cat#130-091-151) from PBMCs and stimulated with

R848 (1 ug/ml), CD40L (10 ug/ml, Biolegend, Cat#591708), BAFF (20 ng/ml,

Peprotech, Cat#310-13-20), IL-2 (10 ng/ml, Peprotech, Cat#200-02), Goat

Anti-Human IgA + IgG + IgM (H+L) (10 ug/ml, Jackson Immunoresearch,

Cat#109-006-064), IL-21 (10 ng/ml, Biolegend, Cat#571204), IFN-γ (20 ng/ml,

Biolegend, Cat#570206) for 3 days and stained with anti-human CD19, CD38,

CD27, IgD, CD11c, CD21, T-bet to detect ABC formation. Addition of Tofacitinib

and Baricitinib (Selleckchem, Cat#S2851) were indicated.

Vector construction, sgRNA cloning, retrovirus packaging and infection.

For the transcription factor screening, sgRNA designed by online design tool

(http://chopchop.cbu.uib.no/) and two DNA oligo annealing by T4

Polynucleotide Kinase (NEB, Cat#M0201L) and cloned into a Bbs1-digested

U6-sgRNA-PGK-Puro-BFP retrovirus vector by T4 DNA ligation. The day before

transfection, 80 million Plate-E cells were plated in 10 cm dish in DMEM with

10% FBS. Cells were transfected with 15 ug sgRNA plasmid using 30 µL of

Lipofectamine 2000 (Invitrogen, Cat#11668019). Plat-E cells were further

cultured for 48 hours. The retrovirus supernatants were centrifuged and debris

were removed. Retrovirus were spin transduced to activated B cells in the

presence of polybrene (10 ug/ml, Sigma, Cat#H9268) with 1260 RCF for 90

minutes at temperature of 32℃. The medium was removed six hours later and

replaced with fresh medium for ABC induction. At least 4 single guide RNA

(sgRNA) plasmids (co expressing blue fluorescent protein, BFP as reporter)

were constructed per gene target and validated editing efficiency in mouse

primary B cells using the online ICE tool (Inference of CRISPR Edits;

Synthego)[40]. We select the most efficient sgRNA with an editing rate over 50%

for further functional screening. The selected sgRNA sequences were listed in

Supplementary Table 5.

Human B cell editing.

Purified B cells from human PBMC were stimulated with R848(1ug/ml), IL-

2(10ng/ml), BAFF (20ng/ml), Goat Anti-Human IgA/IgG /IgM (H+L) (10ug/ml)

for one day. The next day, synthetic sgRNA oligos (GenScripts) were co-

incubated with Cas9 protein (GenScripts, Cat#Z03389) at room temperature for

20 minutes. 0.1 million cells were washed and electroporated using the Neon

transfection system (Thermofisher, Cat#MPK5000S) following manufacturer's

instruction. For candidate TFs, we selected sgRNAs with editing efficiency over

50%. The selected sgRNA sequences were listed in Supplementary Table 5.

After electronical transfection, human B cells were induced to ABC

differentiation described above.

Antibody titers.

Serum samples were collected at indicated time points to detect anti-dsDNA by

ELISA (Alpha Diagnostic International, Cat#5110) and Hep-2 cells slides (Inova

diagnostics Cat#708750) were used to detect anti-nuclear antibodies. FITC-

anti-mouse IgG were used as detection antibody for ANA. Antibody from culture

supernatant was diluted and captured by goat anti-mouse Ig (Southern

Biotechnology, Cat#5300-05B) and detected by HRP- labeled goat anti-mouse

IgG1, IgG2c, IgG3 or IgM specific Abs and TMB substrate (Biotech, Cat#DY999)

followed by sulfuric acid stop. Absorbance values at 370/450 nm (OD) were

read and absolute titers were calculated from standard curves, which were

generated from purified mouse Ig isotype (Southern Biotechnology, Cat#5300-

01B).

RNA extraction and quantitative RT-PCR analysis.

Total RNA was extracted from B cells with TRIzol reagent (ThermoFisher,

Cat#15596026) according to the manufacturer's instructions. RNA quantity and

quality were assessed using the Nanodrop 2000. cDNA was synthesized using

PrimeScript™ RT Reagent Kit (TaKaRa, Cat#RR037A) , Real-time PCR

reactions were performed using TB Green Premix Ex Taq reagent (TaKaRa,

Cat#RR420A) on QuantStudio 7 Flex (Applied Biosystems). For quantification

of gene expression, each sample was normalized to expression of an

endogenous control gene Rpl13a. All primers used were listed in

Supplementary Table 6.

Single-cell RNA library preparation and sequencing

After FACS sorting of ZA⁻CD19⁺ B cells from PBMC of patient with new-onset

SLE, 12000 cells were counted and viability was confirmed to be >90%. Single-

cell suspensions were loaded on a 10X Genomics Chromium single-cell 5′ v2

chip to prepare the libraries according to the manufacturer's protocol. The

library quality was checked by Agilent 2100 Bioanalyzer and Qubit fluorometric

quantitation. Samples were sequenced on an Illumina NovaSeq 6000 with a

sequencing depth of at least 50k reads per cell.

Processing of single-cell RNA-seq data

The raw FASTQ data were processed with Cell Ranger (v3.1.1) [41] by using

GRCm38 annotation (v1.2.0). Output from Cell Ranger was loaded into R and

used Seurat (v3.1.1) [42] package for further analysis. Total 5721 cells with fewer

than 500 or more than 5000 expressed genes, a high percentage of

mitochondrial genes (>10%) were removed. Next, the UMI counts in each cell

were normalized and scaled. The top 1500 highly variable genes were used for

dimensionality reduction using Principal Components Analysis. Upon 'Elbow

plot', we selected the first 15 PC for clustering and UMAP visualization with

default parameters. For clustering, we first selected resolution parameter = 0.8

which produced 9 clusters. To obtain cluster-specific gene signatures, we

identified differential expression analysis of each cluster against the others

using wilcox with parameter (logfc. threshold = 0.25, min.pct=0.1). Clusters

were manually merged with the nearest cluster based on the phylogenetic tree

from Seurat's Build Cluster Tree and similar marker genes. Finally, we got 7

clusters for downstream analysis.

RNAseq.

Cas9-EGFP mouse B cells were retrovirally transduced with sg-NC, sg-Zeb2,

sg-T-bet plasmids after activation for 1 day and further cultured under in vitro

ABC induction condition for 72 hours. Cells were stained withanti-mouse CD19

and CD19$^+$EGFP$^+$BV421$^+$ cells were sorted. Sorted cells were washed twice

and resuspended in TRIzol reagent. Total RNA was extracted and used as input

for Illumina TruSeq RNAseq library kit. After quality assessment, RNAseq

libraries were pooled. 50 bp paired-end sequencing was performed using

Illumina HiSeq2000.

Clean reads were aligned to mm10 reference genome with Hisat2 (v2.1.0) [43]

using default parameters. Gene expression levels were counted with HT-

seq(v0.11.2) [44] [45]using default 'union' mode. Differential expressions of genes

were conducted by DESeq2(v1.24.0) package [46] and simple plots were

produced in R(v3.3.3). GSEA was performed using the online software (Broad

Institute) (http://software.broadinstitute.org/gsea/index.jsp).

CUT-Tag

CUT&Tag libraries were generated following instructions described previously[34]

and manufacturer's protocol (Vazyme, Cat#TD901-01). Briefly, 0.5 million in

vitro derived mouse ABCs were incubated with CoA-coated magnetic beads,

and then primary antibodies (anti-Zeb2 and anti-Tbet) were added to incubate

for 1 hours at room temperature. After incubation, secondary antibodies

(incubated at RT for 1 hours) and pG-Tn5 adapter complex (incubated at RT for

1 hours) were added. Then, tagmentation took place for 1 hours, and DNAs

were extracted using phenol–chloroform–isoamyl alcohol. Libraries were

prepared and cleaned up using Ampure XP beads (Beckman Coulter) and

pooled together for paired-end sequencing.

Sequencing data were analyzed as follows. After trimming the adapters of

reads and removing low-quality bases by TrimGalore software in paired-end

mode, clean reads were mapped to mm10 reference genome with Bowtie2

(v2.3.5) [47] followed by removing PCR duplicates using Picard (v2.19.0).The

genome browser tracks in bigwig format were produced from merged replicates

using deepTools (v2.0) [48] and peaks were called using MACS2 (v2.1.2) [49] using

parameters '-f BAMPE –SPMR -nomodel'. Enriched transcription factor binding

motifs were searched by HOMER (v4.11). A set of sequences were searched

for individual matches to each of the motif using FIMO (https://meme-

suite.org/meme/tools/fimo). All analysis data were visualized using the IGV

software(Broad Institute).

Statistical analysis

Prism software (GraphPad) was used for statistical analyses. Data were shown

as mean ± SEM values. All data points represented the measurement of distinct

samples. Differences were evaluated by analysis of variance followed by

unpaired two-tailed Student's *t* test with 95% confidence intervals to compare

between two normally distributed groups with equal variances unless specified.

Spearman's test was adopted to carry out the correlation analysis. P values of

less than 0.05 were considered significant.

Fig 1. Identification of differentially-expressed transcriptional factors in ABCs.
(**a**) CD19+ B cells were sorted from PBMCs of new-onset SLE patient and performed droplet-based scRNA-seq. Unsupervised clustering was visualized with two dimensional UMAP. Seven clusters are defined and annotated as transitional B cells (TrB), naïve B cells (NavB), activate naïve B cells (aNavB), ABC, memory B cells (MemB), plasma blasts (PB) and plasma cells (PC). Each point is a single cell colored by cluster assessment. (**b**) UMAP plots of selected genes expressions to distinguish ABCs. Color scales for each gene show log-normalized expression levels. (**c**) Dot plots of 43 differentially-expressed transcription factors in ABCs with 27 upregulated TFs and 16 downregulated TFs. Dot size represents proportion of cells expressing indicated TF in particular cluster and intensity of color represents the average expression level. (**d**) C57/B6 mice were intraperitoneally injected with 10 million CD4+ T cells from Bm12 mice for 2 weeks and sacrificed for detection. Flow cytometry detection of splenic CD11c− (blue) or CD11c+ (red) B cells (n=3) for ABC specific markers. (**e-f**) QPCR analysis of ABC signature genes (*Itgam*, *Itgax, Cxcl9, Cxcl10*) and mouse genes encoding equivalent TFs which identified in **c** from splenic CD11c− (blue) or CD11c+ (red) B cells (n=3) sorted by flow cytometry at week 2 after Bm12 induction described in **d**. n refers to the measurement of distinct samples (biological repeats). Data are representative of two in **d** and three in **e,f** independent experiments with 3 mice per group. Bars indicate mean ± SEM values. Statistical significance determined by Student's t test.*p < 0.05, **P < 0.01, ***P < 0.001, ns, not significant.

Fig 2. Cas9 based screen of transcription factors for ABC differentiation.

(**a**) Flow cytometry plots of mouse ABCs (CD11c+T-bet+) derived from in vitro induction. (**b**) Flow chart of Cas9 based screen and in vitro mouse ABC differentiation assay. On day 0 (D0), splenic B cells from Cas9-EGFP and B6 CD45.1 mouse were mixed at a ratio of 4:1 and activated by anti-IgM, anti-CD40 and R848. On day 1(D1), activated B cells were retroviral infected with sgRNA-plasmids and further cultured with the ABC differentiation cocktail for 3 days. On day 4 (D4) cells were stained with surface markers and fixed/permed for fluorescent protein and intracellular T-bet detection. (**c**) Statistical analysis of ABC (CD11c+T-bet+) ratio in groups targeting indicated genes. ABC ratio was caculated as in **Extended Data Fig. 3a**. (**d**) Statistical analysis of ABC (CD11c+T-bet+) ratio in retroviral infected B cells of two distinct sgRNAs targeting *Zeb2*. (**e**) Flow cytometry plots of human DN (CD27-IgD-) and ABCs (CD27-IgD-CD11c+T-bet+) derived from in vitro induction. (**f-g**) Human B cells were activated and electroporated with Cas9-RNP targeting indicated genes and then cultures with the human ABC differentiation cocktail (anti-Ig, R848, CD40L, Baff, IL-2, IFN-γ and IL-21) for a further 3 days. Flow cytometry plots (**f**) and statistical analysis (**g**) of human ABCs (CD27-IgD-CD11c+T-bet+) after gene editing and in vitro differentiation. ABC ratio was normalized with sg-NC. (**i**) Statistical analysis of T-bet MFI in sg-*TBX21* and two distinct sg-*ZEB2* edited human B cells, comparing with sg-NC. n refers to the measurement of distinct samples (biological repeats). Data are representative of 3-4 independent experiments. Bars indicate mean ± SEM values. Statistical significance determined by Student's t test.*p < 0.05, **P < 0.01, ***P < 0.001, ns, not significant.

Fig 3. Zeb2 drives ABC differentiation.
(**a**) Splenic B cells were isolated from Zeb2[flox/flox] CD19[+/+](Ctrl) and Zeb2[flox/flox] CD19[Cre/+](CKO) mice and cultured with cocktail (anti-IgM, R848, anti-CD40 and IL-21) and different IFN-γ concentration for 3 days. Flow cytometry plots (left) and statistical analysis (right) of ABCs (CD11c[+]T-bet[+]). (**b-c**) Zeb2[flox/flox] CD19[+/+](Ctrl) and Zeb2[flox/flox] CD19[Cre/+](CKO) mice were intraperitoneally injected with 10 million CD4[+] T cells of Bm12 mice for 2 weeks. Flow cytometry plots (left) and statistical analysis (right) by different staining panels related to ABC characters. CD19[+]CD21[-]CD23[-] and CD19[+]CD21[-]CD23[-]CD11c[+]CD11b[+] were analyzed as in **b**. CD19[+]CD11c[+] and CD19[+]CD11c[+]CD21[-]T-bet[+] were analyzed in **c**. n refers to the measurement of distinct samples (biological repeats). Data are representative of 2 independent experiments. Bars indicate mean ± SEM values. Statistical significance determined by Student's t test.*p < 0.05, **P < 0.01, ***P < 0.001, ns, not significant.

Fig. 4 Zeb2 regulates ABC specification and differentiation via Jak-Stat.
(a) Volcano graph showing the transcriptional profiles in mouse B cells edited by sg-NC and sg-*Zeb2* with the ABC induction cocktail for 3 days. ABC signature genes, genes of BCR, TLR, Jak-Stat signaling from KEGG database, GO CD40 signaling and crosstalk genes were labeled with colored dots. (**b**) Heatmap showing expression of representative ABC signature genes in mouse B cells edited by sg-NC, sg-*Zeb2*, sg-*Tbx21*. Genes and groups were clustered. (**c**) IGV trunk view of open chromatin region (OCR) signals in transcriptional regulatory sites of *Itgax* and *Itgam*. IgG (gray line), Zeb2 (red line) and T-bet (blue line) specific signals at indicated loci were shown. Consensus Zeb2 (green) and T-bet (orange) binding motifs within the differentially accessible regions were indicated. (**d**) Heatmap showing expression of selected Jak-Stat signaling genes in sg-NC[+] and sg-Zeb2[+] B cells. (**e**) GSEA showing impaired receptor signaling pathway via Stat in sg-Zeb2[+] B cells compared with sg-NC. (**f**) Flow cytometry plots of in vitro induced mouse ABCs (CD11c[+]T-bet[+]) with different concentration of baricitinib (Bari) or tofacitinib (Tofa). (**g**) C57 mice were intraperitoneally injected with 10 million CD4[+] T cells of Bm12 mice for 2 weeks and treated with tofacitinib (Tofa) or dissolved solution (Ctrl) by oral gavage once a day. Flow cytometry plots and statistical analysis of splenic ABCs (CD19[+]CD11C[+]T-bet[+]CD21[-]). (**h**) Flow cytometry plots of in vitro induced human ABCs (CD11c[+]T-bet[+]) with different concentration of baricitinib or tofacitinib. (**i**) Flow cytometry plots of human ABCs (DN2, CD19[+]CD27[-]IgD[-]CD11c[+]CXCR5[-]) or (CD21[-]T-bet[+] DN, CD19[+]CD27[-]IgD[-]CD21[-]T-bet[+]) in PBMCs from patients with rheumatoid arthritis before and after Jak-Stat inhibitor tofacitinib treatment for 4 weeks. n refers to the measurement of distinct samples (biological repeats). Data are representative of 3 independent experiments in f-h. Bars indicate mean ± SEM values. Statistical significance determined by Student's t test.*p < 0.05, **P < 0.01, ***P < 0.001, ns, not significant.

Extended Data Fig. 1 B cells landscape of a new-onset SLE patient.

(**a**) Heat map representing scaled expression values of top 10 genes defining each cluster. (**b**) Number of cells per cluster. (**c**) Violin plots of selected gene expression (*IGHD, CD27, CD38, CD24, CD9, CD79B, VPREB3, MZB1, PRDM1, XBP1, IRF4, PCNA, MKI67, IGHG1, IGHG3, IGKC, JCHAIN*) in different clusters, with log-normalized expression values labeled.

Extended Data Fig. 2 In vitro induction of mouse ABCs.
(**a-b**) Flow cytometry plots (**a**) and statistical analysis (**b**) of viability from B cells cultured with indicated cocktail for 3 days described as in **Fig. 2a**. (**c**) Statistical analysis of mouse ABCs (CD11c⁺T-bet⁺) from in vitro induction described as in **Fig. 2a**. (**d**) MFIs of T-bet and CD11c of B cells described as in **Fig. 2a**. (**e**) QPCR analysis of *Tbx21*, *Cd86*, *Cxcr3* and *Cxcl10* expression from in vitro induced B cells described as in **Fig. 2a**. n refers to the measurement of distinct samples (biological repeats). Data are representative of 3 independent experiments. Bars indicate mean ± SEM values. Statistical significance determined by Student's t test.*p < 0.05, **P < 0.01, ***P < 0.001, ns, not significant.

Extended Data Fig. 3 Gene editing in mouse primary B cells and screen TFs for ABC differentiation.
(**a**) Splenic B cell were activated, retroviral infected and induced for ABC differentiation described as in **Fig. 2b**. Gating strategy for flow cytometry analysis. Cells with certain size gated by FSC-A and SSC-A following duplicates exclusion by FSC-A/H and SSC-A/W. Then viable B cells were gated from CD19+Zombie Aqua- and further divided into BFP+CD45.1- fraction and BFP+CD45.1+ fraction. ABC frequencies were gated with CD11c+T-bet+. Then ABC frequency were compared between BFP+CD45.1- fraction and BFP+CD45.1+ fraction and caculated as ABC rates. ABC rates in sg-TF group normalized with sg-NC group were set as ABC ratio. (**b**) Flow cytometry plots and frequency of ABCs (CD11c+T-bet+) in CD45+sg-*Itgax*+ (Cas9-) and CD45.1-sg-*Itgax*+ (Cas9+) cells described as in **a**. (**c**) Flow cytometry plots of ABCs (CD11c+T-bet+) in CD45+sgRNA+ (Cas9-sgRNA+) and CD45.1-sgRNA+ (Cas9+sgRNA+) cells targeting indicated genes. (**d**) Statistical analysis of cell survival ratio by comparing CD45.1-(GFP+) with CD45.1+(G-FP−) B cells and then normalized with sg-NC. n refers to the measurement of distinct samples (biological repeats). Data are representative of 4 independent experiments. Bars indicate mean ± SEM values. Statistical significance determined by Student's t test.*p < 0.05, **P < 0.01, ***P < 0.001, ns, not significant.

Extended Data Fig. 4 In vitro induction of human ABCs.
(**a-b**) Flow cytometry plots (**a**) and statistical analysis (**b**) of viability of B cells cultured in indicated cocktail described as in **Fig. 2f**. (**c**) Statistical analysis of naïve (CD19+CD27-IgD+), DN (CD19+CD27-IgD-), switched memory (CD19+CD27+CD38-IgD-) and unswitched memory (CD19+CD27+IgD+) B cells generated from B cells cultured in indicated cocktail described as in **Fig. 2f**. (**d**) Frequencies and statistical analysis of CD11c+T-bet-, CD11c-T-bet+, CD11c-T-bet- and CD11c+T-bet+ B cells from in vitro induction described as in **Fig. 2f**. (**e-f**) Flow cytometry plots (**e**) and statistical analysis (**f**) of CD11c+T-bet+ B cells in DN, naïve, switched memory B cells generated from B cells cultured with the human ABC induction cocktail for 3 days. n refers to the measurement of distinct samples (biological repeats). Data are representative of 2 independent experiments. Bars indicate mean ± SEM values. Statistical significance determined by Student's t test.*p < 0.05, **P < 0.01, ***P < 0.001, ns, not significant.

Extended Data Fig. 5 Screen TFs for human ABC differentiation.
(**a**) Cas9 RNP delivery to primary B cells. Positive control sg-*CD19* was used to verify efficiency of gene editing in human primary B cells. (**b**) Flow chart of Cas9-RNP based screen and in vitro human ABC differentiation assay. On day 0, human B cells were activated by anti-Ig, R848, CD40L, BAFF, IL-2. On day 1, activated B cells was electronical delivered of Cas9-RNP and further cultured with the human ABC differentiation cocktail. On day 4, cells were analyzed by flow cytometry. (**c**) Flow cytometry plots of ABCs (CD27⁻IgD⁻CD11c⁺T-bet⁺). Human B cells were activated and then were electroporated with Cas9-RNP targeting indicated gene described as in **b**. (**d-e**) Flow cytometry plots (**d**) and statistical analysis (**e**) of cell viability of B cells electroporated with Cas9-RNP targeting indicated gene and cultured with the human ABC differentiation cocktail. n refers to the measurement of distinct samples (biological repeats). Data are representative of 3 independent experiments. Bars indicate mean ± SEM values. Statistical significance determined by Student's t test.*p < 0.05, **P < 0.01, ***P < 0.001, ns, not significant.

Extended Data Fig. 6 Impaired ABC formation in Zeb2 KO B cells.
(**a**) Schematic diagram for construction of Zeb2 conditional knockout allele: Exon 4 to 6 are chosen as a target for inserting Loxp sites by CRISPR/Cas9 technology. The floxed mice was crossed with mice expressing Cre recombinase. QPCR primers to detect *Zeb2* expression are labeled in red. (**b**) QPCR analysis of *Zeb2* expression in splenic B cells and non-B cell fractions (Day 0) or culture for 3 days described as in **Fig. 3a**. (**c**) Western blot analysis of Zeb2 protein level in splenic B cells from Zeb2 CKO and Ctrl mice. (**d-e**) Flow cytometry and statistical analysis of CD11c (**d**) and T-bet (**e**) of B cells from Ctrl and CKO mice and cultured for 3 days described as in **Fig. 3a**. (**f**) Flow cytometry plots and statistical analysis of splenic germinal center B cells (IgD⁻GL-7⁺CD95⁺), plasma cells (TACI⁺CD138⁺) in Zeb2 CKO and Ctrl mice at week 2 after Bm-12 induction described as in **Fig. 3b**. n refers to the measurement of distinct samples (biological repeats). Data are representative of 2 independent experiments. Bars indicate mean ± SEM values. Statistical significance determined by Student's t test.*p < 0.05, **P < 0.01, ***P < 0.001, ns, not significant.

Extended Data Fig. 7 Zeb2 regulates ABC differentiation by targeting Jak-Stat.
(**a**) Cas9⁺ B cells were activated, retroviral infected with sg-RNA plasmid targeting *Tbx21* and *Zeb2* and cultured with the mouse ABC differentiation cocktail for 3 days. Infected B cells (CD19⁺GFP⁺BFP⁺) were sorted by flow cytometry with enriched efficiency >90%. (**b**) Volcano graph showing the transcriptional profiles in mouse B cells edited by sg-NC and sg-*Tbx21* described as in **a**. ABC signature genes, Immunoglobulin genes, genes of BCR, TLR, Jak-Stat signaling from KEGG database, GO CD40 signaling and crosstalk genes were labeled with colored dots. (**c**) Principal component analysis (PCA) of protein coding genes from RNA-seq data described as in **a**. (**d**) KEGG pathway enrichment analysis of downregulated genes (blue) and upregulated genes (red) in sg-*Zeb2* B cells was performed by online Enrichr tools. P-values was log-normalized. (**e**) Splenic B cells were isolated from Zeb2ᶠˡᵒˣ/ᶠˡᵒˣ CD19⁺/⁺(Ctrl) and Zeb2ᶠˡᵒˣ/ᶠˡᵒˣ CD19Cre/⁺(CKO) mice and cultured cultured with the mouse ABC differentiation cocktail for 3 days. QPCR validation of selected ABC signature genes and Jak-Stat signaling genes. (**f**) Correlation analysis of *ZEB2* mRNA expression with *IL21R* in peripheral B cells of SLE patients. n refers to the measurement of distinct samples (biological repeats). Data are representative of 2 independent experiments in **e**. Bars indicate mean ± SEM values. Statistical significance determined by Student's t test.*p < 0.05, **P < 0.01, ***P < 0.001, ns, not significant.

Extended Data Fig. 8 Zeb2 directly binds to promoter regions of *Il21r* and *Socs5*.
(**a**) Heatmap graph showing distributions of Zeb2 and T-bet specific singaling. The horizontal axis represented the normalized gene range coordinates. (**b**) Peak annotation analysis of distribution of T-bet and Zeb2 in the functional area of gene covered. (**c-d**) Motif-enrichment analysis of Zeb2 (**c**) or T-bet (**d**) specific peaks generated from CUT&Tag sequencing data.(**e**) Motif analysis of Zeb2 specific peak of Chip-seq data in cell lines from Encode project.(**f**) IGV trunk view of OCR signals in transcriptional regulatory regions of *Socs5* and *Il21r*. IgG (gray line), Zeb2 (red line) and T-bet (blue line) specific signals at indicated loci were shown. Consensus Zeb2 (green) and T-bet (orange) binding motifs within the differentially accessible regions were indicated.

Extended Data Fig. 9 Jak-stat inhibitors impair ABC formation.
(**a, c**) Statistical analysis of mouse ABCs (CD11c⁺T-bet⁺) generated from in vitro induction with different concentration (mock, 100 nM, 500 nM and 1000 nM) of tofacitinib (**a**) or baricitinib (**c**). (**b,d**) T-bet MFI of B cells generated from in vitro induction with different concentration (mock, 100 nM, 500 nM and 1000 nM) of tofacitinib (**b**) or baricitinib (**d**). (**e-g**) C57 mice injected with CD4⁺ T cells from Bm12 mice were treated with tofacitinib 30 mg/kg once a day by oral gavage for 2 weeks and then sacrificed to analyze spleen weight (**e**), ANA (**f**) and anti-dsDNA (**g**) in serum.(**h**) Statistical analysis of in vitro induced human ABCs (CD11c⁺T-bet⁺) with different concentration (mock, 100 nM, 500 nM and 1000 nM) of baricitinib or tofacitinib. (**i**) CRP level in RA patients before and after tofacitinib treatment for 4 weeks. n refers to the measurement of distinct samples (biological repeats). Data are representative of 3 independent experiments in **e-h**. Bars indicate mean ± SEM values. Statistical significance determined by Student's t test.*p < 0.05, **P < 0.01, ***P < 0.001, ns, not significant.

Supplemental Table 1. Basic clinical characteristics of a patient with new-onset SLE

	SLE patient
Age	27
Sex	Female
WBC (×10⁹/L)	4.35
RBC (×10¹²/L)	3.14
Hb(g/L)	106
PLT(×10⁹/L)	212
ESR(mm/h)	82
IgA(g/L)	4.23
IgM(g/L)	1.5
IgG(g/L)	26.5
C3(g/L)	0.149
C4(g/L)	<0.017
Anti-dsDNA(IU/ml)	292.16
SLEDAI	24
ANA	Positive
Anti-SSA	Negative
Anti-SSB	Negative
Anti-U1RNP	Negative
Anti-Sm	Negative
Anti-rRNP	Positive
Anti-ANUA	Positive
ACL	Positive
Fever	No
Rash	Yes
Alpecia	Yes
Arthritis	No
Oral ulcers	No
Cutaneous vasculitis	No
Raynaud's	No
Leukopenia and/or Thrombocytopenia	No
Anemia	No
Serositis	No
NPSLE	No
LN	III+V
Urine protein of LN patients (mg/24h)	3379.8

Supplemental Table 2. Basic clinical characteristics of the SLE cohort

	SLE patients (n=21)
Female(%)	20(0.95)
Age, years	32(20-70)
SLEDAI	12(0-22)
Anti-dsDNA (U/ml)	32.13(7.06-100)
C3(g/L)	0.646(0.218-1.25)
C4(g/L)	0.11(0.018-0.343)
ESR(mm/h)	26(1-99)
CRP(mg/L)	3.3(0.2-62.2)
ANA⁺(%)	20(0.95)
Anti-Sm⁺(%)	5(0.24)
Anti-SSA⁺(%)	11(0.52)
Anti-SSB⁺(%)	5(0.24)
Anti-RNP⁺(%)	6(0.29)
LN(%)	12(0.57)
24-urine protein (mg/24h)	2336.3(88.6-14900.7)

Supplemental Table 3. Clinical characteristics of RA patients treatment with tofacitinib

RA patients	Before (n=6)	After (n=6)
WBC(×10⁹/L)	6.30±2.92	6.30±2.03
RBC(×10¹²/L)	4.73±0.56	4.70±0.62
Hb(g/L)	136±25.25	137±23.50
PLT(×10⁹/L)	322±127.5	274±120
ESR(mm/h)	28±25.88	19±11.00
CRP(mg/L)	10.4±32.80	6.6±4.90
Cr(μmol/L)	56.2±8.93	62.2±13.85
ALT(U/L)	21±11.25	25±7.00

Supplemental Table 4. List of antibodies

Antibodies	Source	Identifier
anti-mouse CD4	biolegend	Clone GK1.5
anti-mouse CD8	biolegend	Clone 53-6.7
anti-mouse CD19	biolegend	Clone 6D5
anti-mouse CD19	biolegend	Clone 1D3/CD19
anti-mouse CD44	BD bioscience	Clone IM7
anti-mouse CD62L	BD bioscience	Clone MEL-14
anti-mouse PD-1	eBioscience	Clone J43
anti-mouse B220	biolegend	Clone RA3-6B2
anti-mouse CD95	BD bioscience	Clone Jo2
anti-mouse GL-7	BD bioscience	Clone GL7
anti-mouse IgM	BD Bioscience	Clone R6-60.2
anti-mouse IgD	BD Bioscience	Clone 11-26c.2a
anti-mouse CD138	biolegend	Clone 281-2
anti-mouse TACI	biolegend	Clone 8F10
anti-mouse CD11c	biolegend	Clone N418
anti-mouse/human T-bet	biolegend	Clone 4B10
anti-mouse CD21	biolegend	Clone 7E9
anti-mouse CD73	biolegend	Clone TY/11.8
anti-mouse CD80	biolegend	Clone 16-10A1
anti-mouse CD93	BD Bioscience	Clone AA4.1
anti-mouse CD45.1	biolegend	Clone A20
anti-mouse CD86	BD bioscience	Clone GL1
anti-mouse I-A/I-E	BD bioscience	Clone M5/114.15.2
anti-mouse CD23	BD bioscience	Clone B3B4
anti-mouse CD11b	biolegend	Clone M1/70
anti-mouse CD5	eBioscience	Clone 53-7.3
anti-mouse CXCR4	BD Bioscience	Clone 2B11/CXCR4
anti-mouse CXCR3	biolegend	Clone CXCR3-173
anti-human CD19	biolegend	Clone HIB19
anti-human CXCR5	biolegend	Clone J252D4
anti-human IgD	biolegend	Clone IA6-2
anti-human CD27	biolegend	Clone M-T271
anti-human CD11c	biolegend	Clone S-HCL-3
anti-human CD38	biolegend	Clone HIT2
anti-human CD21	biolegend	Clone Bu32
Biotin Rat Anti-Mouse CD185 (CXCR5)	BD bioscience	Clone 2G8
Zombie Aqua™ Fixable Viability Kit	biolegend	Cat#423102
PE Streptavidin	BD bioscience	Cat#554061
T-bet (4B10) Antibody	Santa Cruz Biotechnology	Cat#sc-21749
ZEB2 antibody	Novus	Cat#NBP1-82991

Supplemental Table 5. List of sgRNA sequence

Gene name	Sg sequence(5'-3')
mouse	
Zeb2 Sg1	GTACCTTCAGCGAAGCGACA
Zeb2 Sg2	TATGAATAGTAACTTGAGTG
T-bet	CGAGGACTACGCATTGCCCG
Itgax	GGGCCGTAACTCACCCTGGA
Zbtb32	CGAGGTATCGAGAGCCACAA
Tfeb	CCTCTGTGGATTACATCCGG
Litaf	GATAACAGACATACTTGCGT
Nfatc2	TTGGAGAGTGGCCACTCGAG
Srebf2	ACTCCAGTGACAGTACACTG
Jazf1	CACAGGCAGCGAGTATGATG
Jun	TGTGCCGCGGAGGTGACACT
Plek	GTTTGCCAAAGTCTTGACAA
Batf	AGAGATCAAACAGCTCACCG
Tfec	CATCAGTGGACTACATCAAG
Sox5	CGAGGGTCCGCTGGTCAGGA
Mbd4	TCAGAGTCGCCAGAAAGCAG
Ets1	TGCCTGGGGAGAGCCAGTCG
Fli1	TCACGACTGAATGTCAAGGA
human	
ZEB2 Sg1	TTGTAGCCCCGGTCGCAGTA
ZEB2 Sg2	GGCGCAAACAAGCCAATCCC
TBX21	AAACCGCCTGTACGTCCACC
BATF	AGGACTCTACCTGTTTGCCA
FLI1	ACTCAATCGTGAGGATTGGT
SREBF2	GCTGCATTCTGGTATATCAA
ETS1	CTTACTAATGAAGTAATCCG
JUN	TGAACCTGGCCGACCCAGTG
CD19	CTGTGCTGCAGTGCCTCAA

Supplemental Table 6. List of primers

	Gene name	Forward	Reverse
Mouse			
	Hopx	5'- CAACTTCAACAAGGTCAACAAGCAC-3'	5'-ACCATTTCTGCGTCTGCTCCT-3'
	Zeb2	5'-GCAGTGAGCATCGAAGAGTACC-3'	5'-GGCAAAAGCATCTGGAGTTCCAG-3'
	Tbx21	5'-TGTGGATGTGGTCTTGGTGG-3'	5'-ATTGTTGGAAGCCCCCTTGT-3'
	Cd11c	5'-TTGGCTTGTGGTCCTACTGTG-3'	5'-GGGAAGTTCTGGCTCTGCTTG-3'
	Cd11b	5'-GTGAATATGTCCTTGGGCCTGT -3'	5'-CGGAGCCATCAATCAAGAAGACA -3'
	Zbtb32	5'-TCCAGATACGGTGCTCCCTTCT-3'	5'-CCAGAGAGCTTTGGAGTGGTTC-3'
	Il7r	5'-CACAGCCAGTTGGAAGTGGATG-3'	5'-GGCATTTCACTCGTAAAAGAGCC-3'
	Cxcl10	5'-ATCATCCCTGCGAGCCTATCCT-3'	5'-GACCTTTTTTGGCTAAACGCTTTC-3'
	Cxcr3	5'-TACCTTGAGGTTAGTGAACGTCA-3'	5'-CGCTCTCGTTTTCCCCATAATC-3'
	Il2ra	5'-GCGTTGCTTAGGAAACTCCTGG-3'	5'-GCATAGACTGTGTTGGCTTCTGC-3'
	Rp13a	5'-GGGCAGGTTCTGGTATTGGAT-3'	5'-GGCTCGGAAATGGTAGGGG-3'
	Cxcl9	5'-AATGCACGATGCTCCTGCA-3'	5'-AGGTCTTTGCGGGATTTGTAGTGG-3'
	Bach2	5'-ACAGACGAAAGATGACTTGGTG-3'	5'-CTCTGCTGAGTAACAGCTTGG-3'
	Myc	5'-ACGGCCTTCTCTCCTTCCTC-3'	5'-GCCTCTTCTCCACAGACACC-3'
	Nfatc2	5'-TCATCCAACAACAGACTGCCC-3'	5'-GGGAGGGAGGTCCTGAAAACT-3'
	Junb	5'-TCACGACGACTCTTACGCAG-3'	5'-CCTTGAGACCCCGATAGGGA-3'
	Jun	5'-CCTTCTACGACGATGCCCTC-3'	5'-GGTTCAAGGTCATGCTCTGTTT-3'
	Srebf2	5'-AGAAAGAGCGGTGGAGTCCTTG-3'	5'-GAACTGCTGGAGAATGGTGAGG-3'
	Foxp1	5'-CATGCCTCTACCAATGGACAGC-3'	5'-GAAGTCGTCACAAACCGCCTCA-3'
	Zfp36l1	5'-CTTCCACACCATCGGCTTTTGC-3'	5'-CACTGGGAAACCCAGCAAAGCT-3'
	Aff3	5'-AGATGACCTGGCTTCCTCCACT-3'	5'-GTCCAGAGGATACAAGTTGCGAG-3'
	Tfeb	5'-CCACCCCAGCCATCAACAC-3'	5'-CAGACAGATACTCCCGAACCTT-3'
	Klf6	5'-GGAAGGTTGTGAGTGGCGTTTTG-3'	5'-AGGTGGTCAGACCTGGAGAAAC-3'
	Irf1	5'-TCCAAGTCCAGCCGAGACACTA-3'	5'-ACTGCTGTGGTCATCAGGTAGG-3'
	Irf8	5'-CAATCAGGAGGTGGATGCTTCC-3'	5'-GTTCAGAGCACAGCGTAACCTC-3'
	Pou2f2	5'-TCCTGGAGAAGTGGCTCAACGA-3'	5'-ATGCTGGTCCTCTTCTTGCGTC-3'
	Litaf	5'-CAAGATGATCGTGACCCAGCTG-3'	5'-GCAGTAGTGGTCCACATCCTGT-3'
	Klf3	5'-CCTCTCATGGTTTCCTTGTCGG-3'	5'-CCTCTGTGGTTCAATTCCAGGC-3'
	Jazf1	5'-CGACATAGCAGTGGCAGCCTTA-3'	5'-TCTCTGTGGTCCAGGACTCATC-3'
	Plek	5'-GGAGCAGTTCACTTGAGAGGCT-3'	5'-TGGAAGTGGCTGCCTGCAAGTA-3'
	Plscr1	5'-GACCTCTGAGATGCAGTAGCTG-3'	5'-GGAGAGTGAACTTGGGCAGACA-3'
	Zfp36l2	5'-AAGTACGGCGAGAAGTGCCAGT-3'	5'-AGAAGCCGATGGTGTGGAAGGT-3'
	Fos	5'-AGGGGCAAAGTAGAGCAGCTA-3'	5'-CAATCTCAGTCTGCAACGCA-3'
	Prdm1	5'-AAGAGGTTATTGGCGTGGTAAG-3'	5'-TAGACTTCACCGATGAGGGGT-3'
	Irf4	5'-CTTTGAGGAATTGGTCGAGAGG-3'	5'-GAGAGCCATAAGGTGCTGTCA-3'
	Tox	5'-AAGATGGCGCACTGCTCTCCAA-3'	5'-CATGCTTGCCTGCTGTCTGATG-3'
	Bhlhe40	5'-CGGATTAACGAGTGCATTGCCC-3'	5'-GCTTCAACGTAAGCTCCAGAACC-3'
	Foxp4	5'-CACCAGGATGTTCGCCTACTTC-3'	5'-TATTCCCGCTCATCCACAGTCC-3'
	Sox5	5'-CGCCAGATGAAAGAGCAACTCAG-3'	5'-TGAGTCAGGCTCTCCAGTGTTG-3'
	Crem	5'-GCAGCACAATCAGCCGATGGTA-3'	5'-AGCTCGGATCTGGTAAGTTGGC-3'
	Arid5a	5'-CGCCTCTGGAAGAACGTGTATG-3'	5'-TGGTAGGAGGCAGTGGCTTGTC-3'
	Arid3a	5'-TCCATCACCAGTGCTGCCTTCA-3'	5'-TCCCTGCGATTGCTGTCTATGG-3'
	Fli1	5'-CCATACAGACCAGTCCTCACGA-3'	5'-CATGGTCTGTGATCCTCCAAGG-3'
	Mbd4	5'-ACAGGATGGCTCTGAAATGCCC-3'	5'-ACTTGTGTCCGTGGGATGCTGT-3'
	Ets1	5'-CCAGAATCCTGTTACACCTCGG-3'	5'-CAGCGTCTGATAGGACTCTGTG-3'
	Hif1a	5'-CCTGCACTGAATCAAGAGGTTGC-3'	5'-CCATCAGAAGGACTTGCTGGCT-3'
	Il-12rb2	5'-TTGGACGGCATCAGTGTCTGCA-3'	5'-TCCGACTTTGCAGAGACCTGGT-3'
Human			
	ZEB2	5'-AATGCACAGAGTGTGGCAAGGC-3'	5'-CTGCTGATGTGCGAACTGTAGG-3'
	IL-21R	5'-ACCAGTCTGGCAACTACTCCCA-3'	5'-AGGGTCTTCGTAATCTGAGCGC-3'
	RPL13A	5'-CGAGGTTGGCTGGAAGTACC-3'	5'-CTTCTCGGCCTGTTTCCGTAG-3'