Monitoring Spore Dispersal and Early Infections of *Diplocarpon coronariae*

Causing Apple Blotch Using Spore Traps and a New qPCR Method

Clémence Boutry[1], Anne Bohr[2], Sascha Buchleither[2], Mathias Ludwig[1], Thomas Oberhänsli[1],

Lucius Tamm[1], Hans-Jakob Schärer[1] *, Pascale Flury[1] *

[1] Plant Pathology, Research Institute of Organic Agriculture FiBL, Ackerstr. 113, 5070 Frick,

Switzerland

[2] Competence center for fruit crops at the Lake of Constance (KOB), Schuhmacherhof 6, 88213

Ravensburg, Germany

* Corresponding authors, equal contribution: H.-J. Schärer, P. Flury

E-mail: hans-jakob.schaerer@fibl.org; pascale.flury@fibl.org

Keywords: TaqMan qPCR, *Marssonina coronaria*, *Diplocarpon mali*, Marssonina leaf blotch, premature leaf

fall, *Malus,* rotating-arm spore trap, impaction spore trap

Funding: Interreg V Program Number ABH003.

Abstract

Apple blotch (AB) is a major disease of apple in Asia and recently emerged in Europe and the USA.

It is caused by the fungus *Diplocarpon coronariae* (*Dc*) *(formerly: Marssonina coronaria;*

teleomorph: *Diplocarpon mali*) and leads to severe defoliation of apple trees in late summer

resulting in reduced yield and fruit quality. To develop effective disease management strategies,

a sound knowledge of the pathogen's biology is crucial. Data on the early phase of disease

development is scarce: no data on spore dispersal in Europe is available. We developed a highly

sensitive TaqMan qPCR method to quantify *Dc* conidia in spore trap samples. We monitored

temporal and spatial dispersal of conidia of *Dc*, and progress of AB in spring and early summer in

an extensively managed apple orchard in Switzerland in 2019 and 2020. Our results show that *Dc*

overwinters in leaf litter and spore dispersal and primary infections occur in late April and early

May. We provide the first results describing early-season dispersal of conidia of *Dc*, which,

combined with the observed disease progress, helps to understand the disease dynamics and will

be a basis for improved disease forecast models. Using the new qPCR method, we detected *Dc* in

buds, on bark and fruit mummies, suggesting that several apple tissues may serve as

overwintering habitats for the fungus, in addition to fallen leaves.

INTRODUCTION

41   *Diplocarpon coronariae* (*Dc*) (Ellis & Davis) Wöhner & Rossman (Crous et al. 2020),

formerly *Marssonina coronaria* (Ellis & Davis) Davis; teleomorph *Diplocarpon mali* (Harada &

Sawamura) is an ascomycete fungus that causes apple blotch (AB) (Wöhner and Emeriewen

2019). The disease can result in severe tree defoliation and weakened trees, and ultimately

decreased yield (Sharma and Thakur 2011) and fruit quality (Park et al. 2013). AB has a significant

economic impact, especially in South and East Asia. In South Korea, the loss due to AB is estimated

at US$ 29.79 M (Kwon et al. 2015). In India, AB is emerging as the most destructive disease

affecting apple trees, becoming a major bottleneck in apple cultivation in Himachal Pradesh, an

important apple producing state in the Western Himalayan region (Rather et al. 2017a; Sharma

and Gupta 2018). Recently, AB has become an issue in Europe (Wöhner and Emeriewen 2019)

and in the USA (Aćimović and Donahue 2018; Khodadadi and Aćimović 2019), especially in low-

input orchards, for example organic orchards and untreated orchards used for juice production

(Bohr et al. 2018; Hinrichs-Berger and Müller 2012; Persen et al. 2012). With the rising demand

for reduced pesticide residues, the prevalence of AB may increase in conventionally managed

apple orchards in the future.

To date, the biology and epidemiology of *Dc* and measures to control AB have been

investigated primarily in Japan, China, Korea, and India, where the disease has been an issue since

the late 1990s, while research on *Dc* and AB in Europe and the USA is nascent (Wöhner and

Emeriewen 2019). *Dc* has a hemibiotrophic lifestyle (Horbach et al. 2011; Zhao et al. 2013) and

the optimal conditions for infection are temperatures between 20 and 25°C and prolonged leaf

wetness (Sastrahidayat and Nirwanto 2016). The minimum leaf wetness period for infection is

eight hours at 15°C, and the risk of infection increases with increasing leaf wetness period and

increasing temperature (Sharma et al. 2009). During the epidemic in summer the pathogen

reproduces asexually by two-celled conidia. Additionally, in fall single celled microconidia

(spermatia) are produced and can be found in acervuli together with conidia (Harada et al. 1974;

Lindner 2012). Initial infections in spring can be caused by conidia released from acervuli or by

ascospores developing in apothecia (Wöhner and Emeriewen 2019). However, the sexual stage

of *Dc* has been described only in India, Japan, and China (Gao et al. 2011; Harada et al. 1974;

Sharma and Gupta 2018), but not in Europe (Hinrichs-Berger 2015; Wöhner and Emeriewen

2019), the USA or Korea (Back and Jung 2014). The contribution of the different spore types to

initial infections and their mode of dispersal has not been conclusively clarified, but wind as well

as splash dispersal has been reported (Dong et al. 2015; Khodadadi et al. 2022; Kim et al. 2019).

Dc overwinters in fallen leaves (Back and Jung 2014). Alternatively, buds and twigs have been

hypothesised as overwintering refugia for *Dc* (Wöhner and Emeriewen 2019). However, data on

the sources of primary infections and early disease progress is lacking in Europe and the USA.

To predict the risk of primary infections by *Dc* and subsequent outbreaks of AB, it is crucial

to understand the spore dispersal pattern of the fungus. Fungal spores can be quantified using

spore traps combined with subsequent quantification using visual (microscopic) or molecular

methods. There are various types of spore traps, and depending on the mode of pathogen

dispersal, the research question or the available resources, certain spore traps are better suited.

Spore traps either collect spores passively by gravitational deposition or actively by sampling

specific volumes of air and capturing the spores by impaction, impingement, filtration, virtual

impaction, cyclone or electrostatic attraction (West and Kimber 2015). The spores are collected

either onto a solid surface like agar on a Petri dish, filter paper, a double-sided adhesive tape,

petroleum jelly (Vaseline)-coated tape, slides or rods, or electrostatic plastic film, or into

Eppendorf tubes (Kim et al. 2018), or more rarely into a liquid. Spores captured by spore traps

have traditionally been identified and quantified by microscopy (Andersen et al. 2009; Sterling et

al. 1999). However, microscopy requires a spore trapping surface that can be examined under a

microscope and a trained investigator to identify and count the specific fungal spores accurately.

Newer methods combine microscopy with image recognition of spores and machine learning

(Basso et al. 2020; Kilin et al. 2019). Furthermore, laser-based real-time optical particle counters

(OPC) that detect particles in the air are being tested for real-time detection of spores in the air

(Basso et al. 2020; Kilin et al. 2019). Besides optical methods, molecular methods are used to

quantify fungal spores. Quantitative real-time polymerase chain reaction (qPCR), loop-mediated

isothermal amplification (LAMP) (Notomi et al. 2015; Ren et al. 2021), serial analysis of gene

expression (SAGE), and microarray technology (Aslam et al. 2017) are based on quantification of

DNA, while the enzyme-linked immunosorbent assay (ELISA) (Kennedy et al. 2000), and the

fluorescent antibody (FA) assay (Schneider et al. 2009) detect fungal proteins via antibodies. In

recent years the qPCR method has become the most frequently used quantitative molecular

method to detect and quantify fungal pathogens. It offers a more sensitive and specific

quantification compared to microscopy, as it can detect low concentrations of a fungal pathogen

against a background of particles and DNA from diverse organisms (Parker et al. 2014) and does

not depend on visual identification of a fungus spore.

Information on spore dispersal along with observations on disease progress and combined

with weather data can provide a basis to understand disease dynamics and develop disease

forecast models. Forecast models provide a valuable tool for the efficient application of crop

protection measures (Agrios 2005;   Hardwick 1998). RIMPro (RIMpro, Amsterdam, The

Netherlands) developed a first disease forecast model for AB in Europe based on literature

information from Asia. Recently, *Dc* spore dispersal was investigated in Korea from June to

October, employing spore traps and microscopy, and a disease forecast model was developed

(Kim et al. 2019). However, the authors did not investigate spore dispersal in the spring, which is

a period critical to understanding the primary infection and disease onset and which may be

subject to efficacious disease management.

This study aimed to gain a better understanding of the early phase of AB disease in spring

and summer. Thus, the specific objectives were to i) develop a qPCR method that allowed the

quantification of *Dc* spores in spore trap samples, ii) investigate temporal and spatial aspects of

Dc spore dispersal in relation to development of AB in the field in spring and early summer, and

iii) identify potential sources of primary *Dc* inoculum.

MATERIALS AND METHODS

Spore traps used in this study. Conidia and ascospores of *Dc* have been reported to be dispersed

by wind (Dong et al. 2015; Kim et al. 2019), while conidia are also reported to be splash-dispersed

(Dong et al. 2015). Two trap types, the Mycotrap and the rotating-arm spore traps, were used to

sample airborne spores in the field (Figs. 1D, E). Additionally, apple bait plants were used as in

vivo spore traps (Fig. 1F). A detailed description of the spore traps is provided in Supplementary

Materials and Methods S1. Briefly, the Mycotrap (Siegfried et al. 1996) (Fig. 1D) is an impaction

spore trap and similar to the Burkard 7-day Recording Volumetric Spore Sampler, which was used

to capture airborne conidia of *Dc* in Korea (Kim et al. 2019). The air is sucked in horizontally

through a sampling orifice and impacts a trapping surface on a cylinder inside a chamber. The

cylinder slowly rotates, completing one turn over seven days, allowing continuous sampling for

the seven day period. The rotating-arm spore traps (Fig. 1E) were constructed based on the

description of Quesada et al. (2018) (details in Supplementary Materials and Methods S2). The

rotating-arm spore traps consisted of two rods connected to a battery-powered motor (spinart™

Lightweight Hanging Motor, 30 rpm). Each rod held a microscope slide on which was mounted a

strip of Vaseline coated plastic film (Supplementary Materials and Methods S3) attached with

25 mm foldback clips (Maul, Bad König, Germany). The rotating-arm spore traps were covered

with an aluminum shield as rain protection.

To collect splash-dispersed *Dc* conidia in rain water, funnels were placed in Schott-bottles to

collect rain splash. Approximately 100 mL of rainwater was collected per Schott-bottle and

filtered through a cellulose acetate filter (25 mm in diameter, 0.8 µm pore size, Sartorius-Stedim,

Goettingen, Germany) in polycarbonate filter housings. The filters were subsequently cut into

pieces of approximately 3 × 3 mm and stored at -20°C.

Design and evaluation of qPCR primers and probes. We aimed at developing a highly specific and

sensitive TaqMan® qPCR assay for *Dc*. The primers and hydrolysis probes designed targeted the

nuclear ribosomal internal transcribed spacer (ITS1) region between the 18S and 5.8S rDNA of *Dc*

using the program Beacon Designer (V8.16, Premier Biosoft, Palo Alto, CA, USA) with the following

parameters: amplicon length between 100 and 200 bp, melting temperature (TM) of primers at

60.0°C +/- 1.0°C, and TM of the probe at 10.0°C +/- 5.0°C above TM of the primers (Table 1). The

program's default settings were used for maximum ΔG of self-complementarity, 3'-end stability,

and % GC. The specificity of the primers and probes was confirmed in silico using the National

Center for Biotechnology Information (NCBI, Bethesda, MD, USA) Primer-Blast (Ye et al. 2012) by

alignment of different ITS gene bank accessions of *Dc.*

The specificity was further tested in vitro with genomic DNA of *Dc* isolates from Switzerland, Korea

and Japan, with apple DNA, and with DNA of different fungi, i.e. fungi within the genus

Diplocarpon, fungi causing Marssonina diseases on other hosts than apple, and fungi that are

common pathogens of apple. Information on the tested isolates is provided in Supplementary

Table S1. *Dc* isolates were cultured on potato peptone carrot dextrose agar (PPCDA) adapted from

Zhao et al. (2010) and containing 100 mL carrot juice, 39 g potato dextrose agar, 10 g peptone,

and 1 L double-distilled water. Other pathogens were cultivated on potato dextrose agar (PDA).

Fungal DNA from the cultures was extracted using a ZymoBIOMICS® Quick-DNA Fungal/Bacterial

Miniprep Kit (Zymo Research, Irvine, CA, USA) according to the manufacturer's recommendations.

The qPCR was designed for use with a TaqMan® probe, but we also tested whether the

primer pair Dc_09 could be used with a DNA intercalating fluorophore (SYBR Green).

qPCR reaction conditions. All primers and probes were synthesized and purified by high-

performance liquid chromatography (HPLC) at Microsynth AG (Balgach, Switzerland). Primers and

probes (Table 1) were dissolved in a TE-dilution buffer (TE-Buffer: 10^{-3} mol $\cdot L^{-1}$ Tris, 10^{-5} mol $\cdot L^{-1}$

Na_2EDTA, pH 8.0).

TaqMan® based qPCR reactions consisted of 5 µL KAPA PROBE FAST (Sigma-Aldrich

Chemie AG, Buchs, Switzerland), 1 µL of each primermix for *Dc* and APA9 (containing primers at

a concentration of 3 µM for Dc_09 and 2 µM Africa Cassava Mosaic Virus (ACMV), and the probe

at 1 µM each), 0 to 2 µL double-distilled water and 1 to 3 µL of template DNA in a total volume of

10 µL. SYBR based *Dc* specific qPCR reactions consisted of 5 µL KAPA SYBR FAST (Sigma-Aldrich

Chemie AG, Buchs, Switzerland), 1 µL primermix (forward and reverse primers at a concentration

of 3 µM each), 3 µL double-distilled water and 1 µL of template DNA in a total volume of 10 µL.

The qPCR assay and data analysis were performed using a CFX96 Touch Real-Time PCR Detection

System (Biorad, Hercules, CA, USA). The amplification and quantification conditions used were an

initial denaturation step of 3 min at 95°C, followed by 39 and 45 cycles of 10 s at 95°C, and 20 s

at 60°C for SYBR and TaqMan® qPCR, respectively. After each SYBR qPCR, a dissociation curve

analysis was performed by gradually increasing the temperature from 65°C to 95°C by 0.5°C per

cycle.

In specificity tests, a SYBR green qPCR assay was performed with the ITS primers ITS1F/2R

in addition to the *Dc* specific qPCR, which served as an amplification control detecting all tested

fungal species (Table 1). The reactions consisted of 5 µL KAPA SYBR FAST, 1 µL primermix (primers

at a concentration of 2 µM each), 3 µL double-distilled water and 1 µL of template DNA in a total

volume of 10 µL and were subjected to an initial denaturation step of 10 min at 95°C, followed by

39 cycles of 30 s at 95°, 30 s at 50°C, and a final extension step for 1 min at 72°C.

To visualize the PCR products, 10 µL of the qPCR reaction were loaded on a 2% Tris-

acetate-EDTA buffered agarose gel stained with ROTI® GelStainRed (Carl Roth GmbH, Karlsruhe,

Germany) and run at 50 V for 60 min. Images were captured under u.v. using an Azure Biosystems

c150 (Azure Biosystems, Dublin, CA, USA) gel imaging workstation.

DNA extraction from spore trap samples and apple leaves. To extract DNA from *Dc* conidia from

Vaseline coated plastic film (Mycotrap and rotating-arm trap samples, Supplementary Materials

and Methods S3) we used the ZymoBIOMICS® Quick-DNA Fungal/Bacterial Microprep Kit (Zymo

Research, Irvine, USA) following the manufacturer's protocol with two modifications: the Bashing

Bead® tubes from the Zymo Microprep Kit were replaced by 2 mL tubes with screw caps

containing 100 mg of 0.5 mm Zirconia bashing beads (N034.1 Roth, Karlsruhe, DE) and the Bashing

Bead™ Buffer volume was reduced from 750 µL to 550 µL.

DNA from filter pieces (rain-splash samples) was extracted using a ZymoBIOMICS® Quick-

DNA Fungal/Bacterial Microprep Kit following the manufacturer's protocol, but adding skim milk

to the bashing bead buffer at a final concentration of 2% to prevent the adhesion of free DNA to

the filter membrane (Liang and Keeley 2013).

DNA from apple leaves with ambiguous AB symptoms was extracted following a CTAB buffer

protocol (described in Supplementary Materials and Methods S4).

qPCR based quantification of *D. coronariae* spores with a standard curve. The number of conidia

corresponding to a given C_q value was calculated using a standard curve based on known

quantities of conidia. Conidia suspensions were prepared by stirring AB-symptomatic apple leaves

with sporulating *Dc* for 10 min in Volvic® natural spring water (Danone S.A. Paris, France), which

is routinely used in our lab for the preparation of spore suspensions, and subsequently filtering

the spore suspension through a sieve (mesh size 0.1 mm). The spore concentration was assessed

using a haemocytometer (0.1 mm depth, 0.0025 mm^2, Neubauer®, Paul Marienfeld GmbH, Lauda-

Königshofen, Germany). A ten-fold dilution series was prepared in double-distilled water

containing 0.05% Tween® 80 as described by Mc Devitt et al. (2004), and 10^5 to 10^1 conidia were

pipetted to the material used in spore traps (Vaseline coated plastic film or filter pieces) and let

dry. DNA extraction and qPCR were performed as described above. The standard curves were

used for the test calibration, the limit of quantification (LOQ), and the calculation of spore

numbers based on Cq values.

All spore trap samples were spiked with 10^7 copies of linearized APA9 plasmid (vector

pUC19 with an insert ACMV; NCBI GenBank accession number AJ427910) (details in the

Supplementary Materials and Methods S5) (Mosimann et al. 2017) as an internal control to

account for differences in DNA extraction efficiency between different samples. The APA9

plasmid was quantified together with *Dc* using a multiplexed TaqMan® qPCR with the primers and

probe for ACMV listed in Table 1. For multiplexing, the singleplex TaqMan® qPCR protocol for *Dc*

described above was adapted by replacing 1 µL of double-distilled water with 1 µL primermix for

APA9 (containing primers at a concentration of 2 µM and the probe at 1 µM). The results were

used to normalize DNA extraction using the method of Von Felten et al. (2010).

Field sites. *Dc* spore dispersal and AB epidemiology were investigated in an extensively managed

organic apple cider orchard in Rickenbach (Zurich, Switzerland, 47°33'30.5"N 8°47'29.9"E). The

site is characterized by a mean annual temperature of 10.0°C and mean annual rainfall of 980 mm

(years 2010 to 2020). The orchard was planted in 2002 to ten different apple cultivars arranged

in 15 tree rows with 18 apple trees in each row (Fig. 1C) and an in-row planting distance of 4 m

and between-row planting distance of 9 m. In 2019, the orchard was treated with organic

fungicides (acidified clay minerals, Myco-Sin® at 8 kg/ha, Andermatt Biocontrol Suisse AG, 6146

Grossdietwil, Switzerland; in a tank mix with Sulphur, Netzschwefel Stulln at 4 kg/ha, Biofa AG,

Münsingen, Germany) on 25 May. The cultivars 'Otava', 'Rajka', 'Rubinola', 'Topaz', and 'Boskoop'

received additional sulfur with lime (Curatio®, Biofa AG, Münsingen, Germany) on 11 June. In

2020, the orchard was treated with sulfur and clay on 27 April, 9 May, 22 May (all cultivars except

'Bohnapfel', 'Schneider', and 'Tobiässler'), and on 2 June (only 'Otava', 'Rajika', 'Rubinola', and

'Boskoop'). Trees with spore traps were excluded from receiving fungicide treatments in 2019

and 2020. In addition, in 2020, the first tree of each row was left untreated.

AB epidemiology was further investigated in an organic apple orchard at the Competence

Center for Fruit Crops at Lake Constance (Kompetenzzentrum Obstbau Bodensee (KOB),

Ravensburg, Germany, 47°46'06.8"N 9°33'18.0"E). The site is characterized by a mean annual

temperature of 9.7°C and mean annual rainfall of 908 mm (years 2010-2020, data from the

weather station in Bavendorf, www.wetter-bw.de). The orchard was planted in 2003 to 'Topaz'

on M9 rootstocks with a 3.2 m between-row and 0.8 m in-row planting distance. For monitoring

AB, 120 trees in two consecutive rows were regularly assessed. No fungicide treatments were

applied to the trees assessed during the entire period of the study.

Weather data including rainfall, leaf wetness, and temperature were obtained from

weather stations close to the field sites: the Agrometeo (Agroscope, Nyon, Switzerland;

[www.agrometeo.ch)](http://www.agrometeo.ch) weather station "Liebensberg", Zurich, Switzerland (47°53'38.1"N

8°83'73.8"E, altitude 535 m), and the Agrarmeteorologie Baden-Württemberg

(Landwirtschaftliches Technologiezentrum Augustenberg, Tübingen, Baden-Württemberg,

Germany, www.wetter-bw.de) weather station "Bavendorf", Germany (47°76'84.4"N

9°55'99.0"E, altitude 481 m).

Disease assessment in the apple orchard. From 2016 to 2020, the severity of AB was assessed

every one to two weeks from the onset of first symptoms through October by scoring entire trees

using a qualitative ordinal scale of 0 to 9 (0 = no disease symptoms; 1, only few spots, no yellow

leaves; 2, few yellow leaves; 3, several clusters with yellow leaves; 4, one part of the tree with

yellow leaves, no bare branches; 5, several parts of the tree with yellow leaves, few branches

bare; 6, intermediate; 7, bare-branched areas in the tree, mass of leaves with symptoms; 8

intermediate; 9, leaves largely dropped, tree bare).

The ordinal scores were used to calculate damage (in percent) based on the McKinney

Infection Index (I) (McKinney 1923), where I (%) = [sum (class frequency × score of rating

class)]/[(total number of ratings) × (maximum grade)] × 100. The scoring scale and the results for

2016 to 2018 have been presented previously in non-peer-reviewed journals for practitioners

(Schärer et al. 2019; Wöhner et al. 2019).

Monitoring spore dispersal in an apple orchard. To investigate dispersal of *Dc* conidia in the field,

different spore traps were installed in the Rickenbach orchard from May to July and from March

to July in 2019 and 2020, respectively. Exact sampling periods for each trap are provided in

Supplementary Tables S2 and S3. To monitor the temporal resolution of the spore dispersal, one

Mycotrap was placed on the ground in the previous year's AB hotspot in the orchard within a row

of 'Otava' trees (Figs. 1C, D). A second Mycotrap was placed in the canopy of an 'Otava' tree (2.5

m above ground) above the first Mycotrap. We hypothesized that primary spores originating from

leaf litter on the ground might be detected rather by the trap on the ground, while a trap in the

tree crown would catch first spores released from other sources. We also hypothesized that later

in the season, when the leaves are heavily infected, there would be more spores in the crown

than on the ground. Container-grown, two-year-old apple trees were used as bait plants and were

placed for periods of five to fourteen days in the orchard and subsequently incubated under rain

protection to assess disease development and correlate the presence of spores in the orchard

with actual infections (Fig. 1F). In 2019, each series of bait plants consisted of three 'Topaz', three

'Gala' and three 'Kiku' apple trees. In 2020, each series of bait plants consisted of five 'Topaz'

apple trees.

To observe the spatial gradients of airborne spore catches in relation to disease severity

in the orchard, seven rotating-arm spore traps were mounted inside and at the edge of the

orchard on trees with severe (3 'Otava', 1 'Topaz' tree), mild (1 'Schneider' tree) and no *Dc*

symptoms (2 'Rubinola' trees) at one of the lowest branches of the apple trees (1.7 m above

ground) and close to the trunk. Moreover, one rotating-arm spore trap was installed at the same

height but approximately 50 m outside the orchard perimeter. The exact location of the traps is

indicated in Fig. 1C. In 2019, the traps were installed only between 22 Mai and 5 June. However,

in 2020, we covered the entire sampling period from March to July. The spore trap samples were

analyzed by the TaqMan® qPCR method described above.

To observe the spatial gradients of splash-dispersed *Dc* conidia 12 Schott-bottles with a

funnel were placed within the row directly below trees as well as between the rows under the

open sky and outside the orchard. The spore trap positions differed regarding disease severity of

nearby trees. Rainwater was collected for 16 hours from 24 to 25 September 2020. It was the first

rainy period after seven consecutive days without precipitation. The rainfall registered for the 16-

hour period was 14 mm.

Precipitation and spore number. To assess whether spore number and precipitation are

correlated, a Kendall rank correlation test was performed in RStudio version 2021.09.1+372

(http://www.rstudio.com) using R version 4.1.2. with the spore number as dependent variable

and the precipitation as independent variable. Since the origin and amount of the inoculum

changed over time the analysis was performed for each month separately.

Investigating possible sources of primary inoculum. To understand *Dc* spore dispersal, it is

essential to know all sources of primary inoculum. We first studied spore dispersal from a deposit

of overwintered *Dc* infected leaf litter. In 2019, a Mycotrap was placed in the field on a mesh cage

depot filled with heavily *Dc* infested leaf litter of the variety 'Remo' at the Competence Center for

Fruit Crops at the Lake of Constance (KOB) (Supplementary Materials and Methods S6). It sampled

for fourteen days on the same piece of plastic film. Consecutive samples were collected (one

sample per fourteen days) from 1 February to 30 June. In 2020, a Mycotrap was placed on a similar

leaf litter deposit at the Research Institute of Organic Agriculture FiBL (Frick, Switzerland) and

sampled from 1 March to 30 June onto Vaseline coated plastic film. The leaf litter for this deposit

was collected in the Rickenbach orchard on 28 February 2020. Spore numbers in the samples

were quantified by TaqMan® qPCR as previously described.

We hypothesized that beside leaf litter other inoculum sources, i.e. bark, bud or fruits,

might be relevant for disease outbreaks. On 28 February 2020, we collected bark and bud samples

from six 'Otava' and six 'Rubinola' trees in the Rickenbach orchard and on 17 March 2020 from

six 'Topaz' trees in the KOB orchard. Bark samples were collected as described by Arrigoni et al.

(2018). Briefly, bark curls of 20 mm length, 5 mm width and 1 mm thickness were taken using a

flame-sterilized scalpel. Each bark sample consisted of a pool of 30 bark curls from a single tree's

upper trunk and lower branches. Ten terminal buds per tree were collected. We further searched

for fruit mummies hanging in the trees in the Rickenbach orchard on 28 February 2020. Five pieces

of the skin of each mummy (with a diameter of about 2 to 4 mm and a thickness of 2 to 5 mm)

were pooled. Samples were lyophilized and ground in a mixer mill MM 200 (Retsch GmbH, Haan,

Germany) at 30 Hz for 30 s. A sample of 100 mg of the ground powder (or as much as was

available) was subject to DNA extraction using a NucleoSpin™ DNA Stool Kit (Macherey-Nagel,

Oensingen, Switzerland). We found only one sample of overwintered leaves hanging in a tree.

This sample was processed the same way but without prior lyophilization. Samples were assessed

for the presence of *Dc* DNA by the TaqMan® qPCR assay described above.

Finally, to assess whether the identified DNA on bark, buds and fruit mummies originates

from viable and infectious fungal tissues, we tried to infect apple leaves in the laboratory from *Dc*

PCR positive bud, bark, fruit samples. 400 to 500 mg of fruit, bark, and bud tissue were cut into

small pieces and wetted with 3.5 mL of Volvic® natural spring water (n=3). Each of the samples

was distributed onto three apple leaves. The experiment was performed twice, once with

container grown 'Topaz' trees and once with 'Topaz' apple seedlings. As controls, three apple

leaves were incubated with laboratory infected apple leaves with *Dc* acervuli (positive control)

and healthy apple leaves (negative control) processed the same way as described for bark, bud,

and fruit samples. In addition, three leaves were inoculated with pieces of a *Dc* positive leaf that

was found in the tree crown after winter. The plants were incubated at 20°C with a 16/ 8 h

day/night cycle. The first four days the relative humidity was 100 %, followed by 50 % for the rest

of the experiment. Symptoms were assessed after three weeks. Leaves showing any sort of spots

or necrosis were tested with the *Dc* qPCR.

Investigation of fruit infections. Apples with symptoms of AB were collected in September of

2020 and 2021 and stored at 4°C until March the following year, when acervuli had developed.

The acervuli were examined under a stereo microscope M205C (Leica Microsystems Switzerland,

Heerbrugg, Switzerland). The apple tissue containing acervuli were stained with cotton blue in

lactic acid and examined with a Leica DM2000 LED microscope (Leica Microsystems Switzerland,

Heerbrugg, Switzerland). Images of the apple tissue containing acervuli were captured using a

Jenoptic Gryphax Subra camera (Jenoptic AG, Jena, Germany). Conidia formed on cool-stored

fruits were tested for their infectivity to apple leaves. To this end, necrotic lesions with acervuli

were cut and added to 1 mL Volvic ® natural spring water in 2 mL Eppendorf tubes, vortexed for

30 s and the resulting spore suspension was used to inoculate leaves of six 'Topaz' seedlings.

Twelve drops of 5 µL suspension were pipetted onto two leaves per seedling. The plants were

incubated as described above for infection experiments with bark, bud and fruit mummies.

RESULTS

New qPCR allows quantification of *D. coronariae* conidia. To quantify *Dc* conidia in spore trap

samples, we developed a new TaqMan® qPCR method with the primers and probes Dc_09 (Table

1). Different primers and hydrolysis probes targeting the ITS gene of *Dc* were tested. Among them,

primers and probe Dc_09 performed best (E= 94.2 %, R^2= 0.995) also when compared to the

published primer Mc_ITS (Oberhänsli et al. 2014) (E= 83 %, R^2 = 0.88). Specificity tests for primer

Dc_09 revealed sound amplification of DNA from different isolates of *Dc* (Fig. 2A), but neither

amplification with DNA of other fungal species nor with apple DNA (Fig. 2A, Supplementary Table

S4).

The amplification of the qPCR for conidial counts on Vaseline coated plastic film, which

was the spore trapping surface used for field samples, exhibited a linear response with an

efficiency of 89.1 % (R^2 = 0.997) (average of seven qPCR runs) (Fig. 2B). The qPCR assay allowed

the consistent detection of as few as ten conidia; however, the amplification was unreliable with

three conidia per sample. Therefore, the limit of quantification is considered to be 3 to 10 conidia.

The SYBR Green based hydrolysis probe assay revealed similar sensitivity and specificity as

the TaqMan assay (Supplementary Fig. S1). However, we used the TaqMan qPCR for our field

experiments since it allowed multiplexing of the *Dc* specific qPCR with the APA9 specific qPCR

(plasmid standard), which we used to normalize the *Dc* quantification data with the DNA

extraction efficiency.

Apple blotch disease progress in two Central European apple orchards.

At the Rickenbach field site and at the KOB field site disease progression on 'Topaz' started

with the appearance of first signs of the pathogen (i.e. visible formation of acervuli) and

symptoms of AB (yellowing of the leaf), which varied between years, but occurred at the latest in

August in both orchards and all years. Progress of AB was generally characterized by a steady

increase in severity. However, differences in disease severity were observed between years. At

KOB, for example, AB was more severe in 2017 compared to other years (Fig. 1A). Rainfall was

more frequent from June through October in 2017, i.e. 78 days of rainfall compared to between

36 and 57 days in the other years (Supplementary Fig. S2). At both sites, 2018 was characterized

by an exceptionally hot and dry summer (only 36 and 33 days with rain from June to the end of

September at KOB and Rickenbach, respectively) (Supplementary Figs. S2, S3) resulting in lower

severity of AB (Fig. 1A). At Rickenbach, AB was most severe in 2019 and 2020; which was not the

case at KOB (Fig. 1A).

At the Rickenbach site we identified 'Otava', 'Rajka', 'Florina', and 'Topaz' as apple cultivars

highly susceptible to AB (Figs. 1B, C). In contrast, 'Bohnapfel', 'Blauacher', 'Tobiässler', and

'Schneider' were relatively tolerant with low AB severity, even at the end of the season (Fig. 1C).

'Rubinola', had no or very low severity, despite a high inoculum pressure in the orchard (Figs. 1B,

C).

Monitoring dispersal of *Dc* conidia in the field. In 2017 and 2018, the first clearly visible

symptoms on 'Otava' trees developed by mid-June and early July, respectively. Therefore,

considering an incubation period of about three weeks (Lee et al. 2011), the first infections were

expected to occur towards late May.

In 2019, very large numbers of spores were detected by the Mycotrap on the ground

during the first half of May (Fig. 3A). Subsequently, spore numbers generally decreased, except

for a peak in the first week of June. In May, leaves with spotting indicative of AB were collected

and tested for *Dc* by qPCR, revealing the first *Dc* positive leaf on 17 May. The first unambiguous

symptoms were observed three weeks later, on 7 June (Fig. 3A). The next spore peak was

detected on 24 June after a rainy period. At that time, 'Otava' trees already showed severe leaf

yellowing, indicating the emergence of secondary inoculum. In July, large spore numbers were

recorded in the air on most days. The second Mycotrap in the crown of an 'Otava' tree detected

more spores than the Mycotrap on the ground, except on 24 and 25 June (Supplementary Fig.

S4).

The first apparent symptoms on bait plants developed between 17 and 22 May (Fig. 3A).

From 22 May on, at least one bait plant showed typical AB symptoms of leaf yellowing, except for

the period from 3 to 7 June. The most severe leaf yellowing and leaf drop were observed on the

bait plants exposed between 7 and 13 June, followed by those exposed between 13 and 19 June.

In 2020, very few spores were captured by the trap on the ground before bud break in

March (Fig. 4). The first peak in spore production was captured on 5 May by the trap on the ground

at the end of a rainy period that had followed an extremely dry April (Figs. 3B, 4, Supplementary

Fig. S2). The bait plants standing in the orchard from 24 April to 8 May were the first to develop

AB symptoms. The first *Dc* positive leaf based on a qPCR test was on 8 May. Leaves with clear

symptoms and production of conidia were observed on 22 May 2020, which was approximately

two weeks earlier than observed in 2019.

A second peak in spore production was detected on 5 June by the spore trap in the tree

crown. As in 2019, spores were detected in the air almost every day by both traps from mid-June

to the end of July (Figs. 3, 4). Numbers of spores varied, but no significant correlation was found

between spore numbers and precipitation (Supplementary Fig. 6). In Mycotrap samples in early

June (lower trap: 5 to 19 June, upper trap: 31 May to 9 June), we experienced technical problems

resulting in poor DNA extraction efficiency. In cases with signal by the *Dc* qPCR, the low efficiency

could be taken into account by normalization with the internal plasmid standard. However, in

cases where there was no signal *Dc*, we were unable to conclude whether the zero value was due

to no spores or the low efficiency precluding detection. Thus, for some days during the first half

of June, *Dc* spores might have been present in the air, although no catch is indicated (Figs. 3B, 4).

In 2020, all bait plants placed in the orchard after 22 May developed AB symptoms. From

22 May to 5 June, bait plants exhibited approximately 50 to 80% of leaves symptomatic, and after

5 June incidence of leaves exhibiting AB exceeded 90% (Supplementary Fig. S5).

Dispersal of conidia of *Dc* with distance from an inoculum source. Rotating-arm spore traps

detected the first spores after 22 May in 2019 and 2020. Spore numbers detected in rotating-arm

spore traps were generally lower than those captured in the Mycotrap samples (Table 2, Figs. 3,

4). The highest spore numbers were generally detected in traps on 'Otava' trees, which exhibited

severe symptoms of AB, while almost no spores were detected in traps on 'Rubinola' trees.

Despite being only two rows (18 m) apart from highly diseased trees, the 'Rubinola' trees were

virtually free of AB, although some leaf yellowing and premature leaf fall was observed on

'Rubinola' trees. The symptoms on 'Rubinola' were distinctly different from typical AB, and when

such leaves were tested by qPCR no *Dc* DNA was detectable (n = 7). Furthermore, no spores were

detected in the trap 50 m beyond the orchard edge with one exception: during the period from

22 May to 3 June 2019 spores of *Dc* were detected under 'Rubinola' trees and outside the orchard.

A strong wind event with wind speed up to 37 km/h combined with rain on 21 May may have

transported some infected leaves towards the 'Rubinola' trees and outside the orchard explaining

these results.

Besides the dispersal of *Dc* spores via the air, rainwater splashes might be an important

factor for spreading the disease within a tree. Therefore, we collected rainwater below highly

infected trees ('Otava', 'Topaz'), below symptomless trees ('Rubinola'), between the rows under

the open sky and outside the orchard during a rainy period in September 2020. We detected

several hundred spores per mL rainwater below 'Otava' trees, fewer spores between rows, and

almost no spores below non-infected 'Rubinola' trees or outside the orchard (Supplementary Fig.

S6).

Origin of primary infections. At KOB in 2019, the first spores were released from a deposit of

overwintered leaf litter in the second half of April (Table 3). At the Research Institute of Organic

Agriculture FiBL in 2020, however, a few spores were detected above a leaf litter deposit in the

second half of March. In 2020, no spores were detected in April at FiBL, which was characterized

by extraordinarily dry weather (Supplementary Fig. S3), but the spores were again released from

the leaf litter deposit during May (Table 3). Spores were detected in the second half of June at

KOB and FiBL. Generally, spore numbers detected in the experiments were very low and thus not

reliably quantifiable (LOQ = 10 spores).

While all bark and bud samples of 'Rubinola' were negative for Dc, the majority of bark

and bud samples from 'Topaz' and 'Otava' trees were tested positive for Dc (Table 4). Bark

samples exhibited high Cq values close to the limit of detection corresponding to low amounts of

DNA. Bud samples contained up to 10^2 to 10^4 times higher levels of Dc DNA than bark samples.

The highest amount of Dc DNA was detected in a bud sample of an Otava tree with a Cq value of

25.6 (Table 4), which corresponds to the DNA of 10^4 conidia.

In spring 2020, four out of five fruit mummies, which were collected from trees with strong

AB infestation in 2019, were tested slightly positive (Cq values between 33.65 and 40.38) (Fig. 5C,

Supplementary Table S5), while fruit mummies from trees without AB symptoms the previous

year were negative (Supplementary Table S5). For comparison, a leaf sample still hanging in a

susceptible 'Florina' tree after the winter of 2019 was *Dc* positive (Cq, 25.83).

Attempts to infect apple leaves in the laboratory by applying wetted pieces of bark, buds,

fruit mummies and leaves from previously AB symptomatic trees did not result in successful

infections (data not shown). But *Dc* infected fruits can serve as an inoculum source for infection

of leaves after six months of storage. In fall 2020 and 2021, *Dc* infections were observed on 'Otava'

fruits in the Rickenbach orchard (Fig. 5A). In the following spring, after storing fruits at 4°C for six

months, conidia formed (Figs. 5B, D-I), and the conidia infected the leaves of apple seedlings in

the laboratory (Supplementary Fig. S7).

DISCUSSION

Development of a *D. coronariae* specific qPCR. To date, spore trap catches of *Dc* conidia have

been quantified using microscopy (Kim et al. 2019). However, microscopy is labor intensive and

identification of the *Dc* spores in field samples may be challenging. We developed a qPCR method

to provide a very sensitive means for quantification of as few as ten conidia per sample, which is

in a similar range to qPCRs developed for other fungi (Calderon et al. 2002; Dvořák et al. 2015;

Luchi   et   al.   2013).   The   qPCR   method   further   allows   diagnosis   of

AB prior to developing unambiguous visual symptoms. Early diagnosis is especially useful when

small necrotic lesions appear that are indistinguishable from symptoms from other causes. In

summary, we established a new qPCR method, which allowed the quantification of *Dc* spores in

spore trap samples, and that can be used to detect initial infections in the field, and that enabled

the identification of *Dc* DNA in various apple organs.

qPCR, though it requires several processing steps, is a rather fast method and allows the

processing of many samples concurrently. However, it has the disadvantage to only allow

conclusions regarding the presence of DNA, but not regarding the fungal organ (e.g. mycelium,

ascospore, conidia, spermatia) or its viability and infectious potential. Therefore, visual

assessments and infection experiments are needed to understand how *Dc* exists on bark, bud and

fruit samples and the role of the tree organs in the fungal life cycle.

Timing of primary infections. To determine the timing of primary infections by *Dc*, we combined

data on airborne spore catches with symptom development both in the field and on a consecutive

series of exposed bait plants. In the first three study years (2016-2018), clear AB symptoms were

observed, the earliest showing in June. However, testing leaves with ambiguous necrotic spots by

qPCR in 2019 and 2020 revealed that the first leaves of highly susceptible varieties were already

infected with *Dc* in May (17 May 2019 and 8 May 2020). The infections were most likely caused

by spores released in late April or early May. The first spore peaks were detected in the orchard

in early May in both 2019 and 2020. In 2019, spore traps were installed on 9 May 2019. Thus, it is

possible that spores were already present one or two weeks earlier. In 2020, the experiment was

initiated at the end of February, and the first clear spore peak was detected by the Mycotrap on

the ground on 5 May. We conclude that the spores collected in early May likely caused the first

infections in the field since the first bait plants in the field that developed symptoms were

exposed from 24 April to 8 May 2020.

In line with the results of the spore traps in the orchard, the spore traps located above leaf

litter caught conidia of *Dc* at the end of April and early May in both study years. The few spores

detected above the leaf litter and in the field in March 2020 were released before bud break, and

we, therefore, conclude that they were irrelevant to primary leaf infections.

Our qPCR method does not distinguish between conidia and ascospores, but apothecia of

Dc have not yet been observed in Europe (Wöhner and Emeriewen 2019), indicating it is unlikely

they were the spore type captured. Although not yet observed, the importance of ascospores as

primary inoculum in Europe should be subject to further research. A recent population genetic

study suggested a mixed sexual and asexual reproduction of *Dc* in Europe, although *Dc*

populations in Europe are genetically homogenous, clonal and dominated by a few multi-locus

genotypes (Oberhänsli et al. 2021).

The first spore peaks of the season were observed at the end of, or shortly after rainy

periods, whereas no spores were detected during dry periods before rain events, e.g. April 2020.

The result indicates that rain might be required for the release of conidia of *Dc* from leaf litter,

and acervuli might need a wetting period prior to release of conidia. This hypothesis may explain

the slightly negative correlation found for May 2019 between daily spore catches and

precipitation (Supplementary Table S6).

Based on our results, primary infections of apple with *Dc* occur in Central Europe in late

April or early May after periods of rain, which is earlier than was previously assumed (Sutton et

al. 2014). Moreover, the spore catches in March indicate that *Dc* spore discharge may occur

before April and, depending on weather conditions and phenological stage of the host, infections

could be possible. Knowing the timing of primary infections can be valuable as a basis for

management actions to preventing epidemics of AB. While protection strategies against AB in

Europe are currently focused on the summer months (Hinrichs-Berger 2015), prevention of

primary infections in the early spring might be an important additional step for more efficacious

management strategies in the future.

Leaf litter and other sources of primary inoculum. In addition to the timing of primary infections,

we were also interested in the origin of the primary inoculum. Several studies report that conidia

inside acervuli overwinter in leaf litter (Back et al. 2015; Dong et al. 2015; Gao et al. 2011; Goyal

et al. 2018; Lee et al. 2011; Sastrahidayat and Nirwanto 2016; Sharma et al. 2009). This is in

accordance with our finding of *Dc* spores above leave litter deposits in spring and also with the

observation that in May, most spores were captured in the Mycotrap on the ground, and not in

the Mycotrap in the tree canopy.

Small quantities of *Dc* DNA were detectable on fruit mummies hanging in the trees at the end

of February, and acervuli with infectious conidia were produced on *Dc* infected fruits after

overwintering at 4°C. Moreover, considerable quantities of *Dc* DNA were detected on buds, and

small amounts of *Dc* DNA were found on the bark of trees that had AB the previous season. All

these organs are reported as overwintering sites for other apple pathogens, including *Monilia*

spp. on fruit mummies, *Podosphaera leucotricha* in buds and *Neofabraea* spp. on bark (Sutton et

al. 2014). Moreover, *D. rosae*, a close relative of *Dc* causing black spot disease on roses,

overwinters, in addition to leaf litter, on bud scales and stems (Cook 1981). To date, it is unclear

what the survival structures of *Dc* on bark and buds are (e.g. mycelium, spores), and whether they

are infectious. The fact that we could not cause leaf infections from bud, bark or fruit mummy

tissues may be due to technical issues, and the role of the various apple tree organs for

overwintering of *Dc* requires further investigation. *Dc* present on buds and wood in propagation

material could be exchanged between nurseries across Europe and may explain the rapid spread

of this disease to many European apple production areas since its first detection in Northern Italy

(Oberhänsli et al. 2021). Our observation that conidia produced on fruits after six months in

storage at 4°C are able to infect apple leaves provides an indication that fruit-derived conidia

might represent an additional source of primary inoculum at least on a local scale. Moreover, our

description of fruit infection adds to our very limited knowledge regarding the role of fruit

infections in the epidemiology of AB (Wöhner and Emeriewen 2019).

Timing of secondary infections and epidemic development. By the end of May (2020) or the

beginning of June (2019), the first leaves that were producing acervuli and conidia of *Dc* were

observed, initiating the secondary phase of infection. In contrast to the first major peak of conidia

captured in May 2020, the second major peak (5 June) was not detected by the Mycotrap on the

ground but by the Mycotrap situated in the tree canopy. We hypothesize that the conidia

produced in early May originated from overwintered leaf litter, resulting in higher spore counts

in the Mycotrap on the ground. But in June, the conidia originated from the recently infected

leaves in the tree canopy, resulting in a higher spore count in the Mycotrap situated in the tree

canopy. In support of the hypothesis, the rotating-arm spore traps in the tree crown did not

capture conidia before the end of May.

From June onwards, damage caused by AB steadily increased in the orchard (Fig. 1A, B).

From the middle of June to the end of the experiment, high numbers of spores were detected

irrespective of rain events, and no correlation was found between daily precipitation and caught

spores in June and July for both years (Supplementary Table S6). This contrasts with our

observation of the early spore peaks, which occurred at the end of the rain periods and the

observations by Kim et al. (2019). They investigated dispersal of conidia of *Dc* from June to

October 2013 and 2015 in Korea and reported spore dispersal peaks during and two days after

rain events compared to dry periods. In that study, disease incidence and total spore counts were

much lower than we observed at our field site, which might explain some of the differences. Based

on our results, we hypothesize that primary inoculum may require water to ripen and discharge,

while secondary inoculum are also released on dry days. Lab experiments indicate that *Dc* conidia

can survive for several days under dry conditions after landing on a susceptible leaf, and cause

infection once the leaf becomes wet (Clémence Boutry et al., unpublished data). If so, all

secondary inoculum result in risk of infection whenever leaf wetness occurs.

The development of AB is affected by environmental factors (Li et al. 2011). Rainfall and

the subsequent relative humidity are positively correlated with the disease severity in the

summer months (Rather et al. 2017b; Sastrahidayat and Nirwanto 2016; Sharma and Sharma

2005). Under controlled conditions, the minimum leaf wetness period for infection is 8 h at 15°C

and decreases to 4 h with temperatures between 20 and 25°C (Sharma et al. 2009). Moreover,

the incubation period can vary depending on the weather conditions. At a temperature range of

15 to 25°C and seven days of leaf wetness, the incubation period is 10 to 20 days (Harada et al.

1974). In our study, the bait plants showed symptoms within one to four weeks after return from

the orchard, with a shorter incubation period for bait plants placed in the orchard later in the

season (June, July) compared to the beginning of the season (May). In 2020, the first necrotic

spots indicative of AB and the more developed symptoms were observed earlier and more rapidly

than in 2019. The faster disease progression in late spring 2020 was likely due to higher May

temperatures compared to the temperatures in 2019.

Spore dispersal in relation to distance from an inoculum source. Rotating-arm spore traps placed

in the orchard revealed high spore concentrations in AB-diseased trees, but no or few spores in

trees without symptoms. The observation could indicate that conidia of *Dc* are not dispersed over

large distances. However, as noted there were generally low spore counts captured by the

rotating-arm traps compared to the Mycotrap, indicating that the rotating-arm traps were much

less efficient in capturing spores in the field (so spores might have been there but the lower

sensitivity of the trap precluded detection). Thus, wind might transport some spores over larger

distances than we observed. Besides wind dispersal, dispersal in rain splash may contribute to the

spread of *Dc* within the tree canopy, as indicated by the very high spore counts in rainwater below

highly symptomatic trees (Supplementary Fig. S6).

Dc **life cycle.** Based on our results we contend that in Europe, the fungus overwinters primarily in

infected leaf litter (Fig. 6). However, bark, buds and fruit mummies might be alternative sources

of primary inoculum since we detected *Dc* DNA on these organs after winter in previously AB-

symptomatic trees. A few spores appear to be released in March, but primary inoculum infections

start in late April or early May, depending on the weather conditions. Secondary inoculum spores

developed in late May or early June in our field sites, which was within three to four weeks of the

primary infections. In May, spore peaks were mainly found at the end of, or after periods of rain;

whereas spore load in the air was generally high from the middle of June to the end of July. Thus,

every wet period in summer represents a risk for infection. The exact mode of pathogen dispersal

has not yet been conclusively determined. Conidia are probably dispersed by wind and rain splash

over larger distances; for the spread of *Dc* within a tree canopy or between neighboring tree

canopies, rain splash could be a major dispersal agent.

In summary, we developed a sensitive and specific qPCR method to detect and quantify

conidia of *Dc* captured in spore traps, which can be combined with the quantification of other

apple pathogens, e.g. *Venturia inaequalis* (Meitz-Hopkins et al. 2014), in spore trap samples from

apple orchards in the future. Our data contributes to our understanding of AB, especially the

timing of early spore dispersal and infections. The information can help to improve disease

forecast models for AB, such as the one by RIMpro (RIMpro, Amsterdam, The Netherlands), and

provide a basis for early disease management. However, more research is needed to test whether

preventing primary infections will reduce disease progress over the season, thereby reducing the

need for fungicide applications in the summer. Finally, the presence of *Dc* DNA on bark, buds and

fruit mummies indicates that the role of overwintering inoculum on these organs and the

anthropogenic dispersal of the pathogen on propagation material should be further investigated.

ACKNOWLEDGEMENTS

We are very grateful to Jürg and Pascale Strauss, who allowed us to perform major parts of the

study in their apple orchard and actively supported us with technical assistance in the field. We

thank David Metzger, Mona Blattner, Sumin Bae, Camilla Kappeler, Patrick Widmann, and Adrian

Rutzer for their excellent help with fieldwork, disease scoring and laboratory assistance. We are

very grateful to Lauren Dietemann for taking the time to review the manuscript for the English.

We further thank Monika Maurhofer for valuable scientific input as an examiner on a Master

thesis included in this publication. Finally, we are grateful to Marc Trapman for developing the *Dc*

infection forecast model 'RIMPro Marssonina', which gave important stimuli for our work and has

shown to be helpful in disease forecast.

LITERATURE CITED

Aćimović, S. G., and Donahue, D. 2018. Marssonina leaf blotch of apple. Pages 1-4 in: Tree Fruit News.

Agrios, G. N. 2005. Plant Pathology. Elsevier Academic Press.

Andersen, G. L., Frisch, A. S., Kellogg, C. A., Levetin, E., Lighthart, B., and Paterno, D. 2009. Aeromicrobiology/Air Quality. Pages 11-26 in: Encyclopedia of Microbiology (Third Edition). M. Schaechter, ed.

Arrigoni, E., Antonielli, L., Pindo, M., Pertot, I., and Perazzolli, M. 2018. Tissue age and plant genotype affect the microbiota of apple and pear bark. Microbiological research 211:57-68.

Aslam, S., Tahir, A., Aslam, M. F., Alam, M. W., Shedayi, A. A., and Sadia, S. 2017. Recent advances in molecular techniques for the identification of phytopathogenic fungi – a mini review. J Plant Interact 12:493-504.

Back, C.-G., and Jung, H.-Y. 2014. Biological Characterization of Marssonina coronaria Infecting Apple Trees in Korea. The Korean Journal of Mycology 42:183-190.

Back, C.-G., Lee, S.-Y., Kang, I.-K., Yoon, T.-M., and Jung, H.-Y. 2015. Occurrence and analysis of apple blotch-like symptoms on apple leaves. Kor. J. Hort. Sci. Technol. 33:429-434.

Basso, T., Tran, M., Pini, V., Schnée, S., Dubuis, P.-H., Kasparian, J., and Wolf, J.-P. 2020. Détection des spores de mildiou et d'oïdium en temps réel à haute résolution spatiale. Revue suisse Aboriculture, Viticulture, Horticulture 52:344-349.

Bohr, A., Buchleither, S., Hechinger, M., and Mayr, U. 2018. Symptom occurence and disease management of Marssonina blotch. in: 18th International Conference on Organic Fruit-Growing 19-21 February 2018, Hohenheim, Germany.

Buchleither, S. 2019. Neueste Erkenntnisse zur Blattfallkrankheit „Marssonina coronaria". Öko-Obstbau 8-11.

Calderon, C., Ward, E., Freeman, J., and McCartney, A. 2002. Detection of airborne fungal spores sampled by rotating-arm and Hirst-type spore traps using polymerase chain reaction assays. J Aerosol Sci 33:283-296.

Cook, R. T. A. 1981. Overwintering of Diplocarpon-Rosae at Wisley. T Brit Mycol Soc 77:549-556.

Crous, P. W., Wingfield, M. J., Schumacher, R. K., Akulov, A., Bulgakov, T. S., Carnegie, A. J., Jurjevic, Z., Decock, C., Denman, S., Lombard, L., Lawrence, D. P., Stack, A. J., Gordon, T. R., Bostock, R. M., Burgess, T., Summerell, B. A., Taylor, P. W. J., Edwards, J., Hou, L. W., Cai, L., Rossman, A. Y., Wohner, T., Allen, W. C., Castlebury, L. A., Visagie, C. M., and Groenewald, J. Z. 2020. New and Interesting Fungi. 3. Fungal Syst Evol 6:157-231.

Dong, X.-l., Gao, Y.-e., Li, B.-h., Yong, D.-j., Wang, C.-x., Li, G.-f., and Li, B.-d. 2015. Epidemic dynamics of apple Marssonina leaf blotch over whole growth season in the central area of Shandong Peninsula. Scientia Agricultura Sinica 48:479-487.

Dvořák, M., Rotková, G., and Botella, L. 2015. Detection of airborne inoculum of *Hymenoscyphus fraxineus* and *H. albidus* during seasonal fluctuations associated with absence of apothecia. Forests 7:1.

Gao, Y.-e., Li, B.-h., Dong, X.-l., Wang, C.-x., Li, G.-f., and Li, B.-d. 2011. Effects of temperature and moisture on sporulation of *Diplocarpon mali* on overwintered apple leaves. Scientia Agricultura Sinica 44:1367-1374.

Gardes, M., and Bruns, T. D. 1993. ITS primers with enhanced specificity for basidiomycetes - application to the identification of mycorrhizae and rusts. Molecular Ecology 2:113-118.

Goyal, A., Sharma, J. N., and Phurailatpam, S. 2018. Studies on the mode of survival of *Diplocarpon mali* causing premature leaf fall of apple. International Journal of Current Microbiology and Applied Sciences 7:2465-2474.

Harada, Y., Sawamura, K., and Konno, K. 1974. *Diplocarpon mali*, sp. nov., the perfect state of apple blotch
fungus *Marssonina coronaria*. Ann Phytopath Soc Japan 40:412-418.

Hardwick, N. V. 1998. Disease forecasting. in: The epidemiology of plant diseases. D. G. Jones, ed. Kluwer
Academic Publishers.

Hinrichs-Berger, J. 2015. Die Marssonina-Blattfallkrankheit des Apfels. Pages 72-74 in: Obstbau.

Hinrichs-Berger, J., and Müller, G. 2012. Vorzeitiger Blattfall an Apfelbäumen in Baden-Württemberg durch
Befall mit *Marssonina coronaria*. Pages 71-71 in: 58. Deutsche Pflanzenschutztagung Julius-Kühn-
Archiv, Braunschweig, DE.

Horbach, R., Navarro-Quesada, A. R., Knogge, W., and Deising, H. B. 2011. When and how to kill a plant
cell: infection strategies of plant pathogenic fungi. Journal of plant physiology 168:51-62.

Kennedy, R., Wakeham, A. J., Byrne, K. G., Meyer, U. M., and Dewey, F. M. 2000. A new method to monitor
airborne inoculum of the fungal plant pathogens *Mycosphaerella brassicicola* and *Botrytis cinerea*.
Applied and Environmental Microbiology 66:2996-3000.

Khodadadi, F., and Aćimović, S. G. 2019. Marssonina blotch of apple caused by *Marssonina coronaria* in
New York. in: Plant Health 2019 - American Phytopathological Society Annual Meeting, Cleveland,
OH, USA.

Khodadadi, F., Martin, P. L., Donahue, D. J., Peter, K. A., and Acimovic, S. G. 2022. Characterization of an
emerging disease: Apple blotch caused by Diplocarpon coronariae (syn. Marssonina coronaria) in
the Mid-Atlantic United States. Plant Disease.

Kilin, V., Pini, V., Kasparian, J., Gros, S., and Wolf, J. P. 2019. Real-time and spatially resolved assessment
of pathogens in crops for site-specific pesticide reduction strategies. BIO Web of Conferences
15:01019.

Kim, H. S., Jo, J. H., Kang, W. S., Do, Y. S., Lee, D. H., Ahn, M. I., Park, J. H., and Park, E. W. 2019. MBCAST:
A forecast model for Marssonina blotch of apple in Korea. Plant Pathology Journal 35:585-597.

Kim, K. H., Kabir, E., and Jahan, S. A. 2018. Airborne bioaerosols and their impact on human health. J
Environ Sci (China) 67:23-35.

Kwon, D., Kim, S., Kim, Y., Son, M., Kim, K., An, D., and Kim, B. 2015. An empirical assessment of the
economic damage caused by apple Marssonina blotch and pear scab outbreaks in Korea.
Sustainability 7:16588-16598.

Lacey, M. E., and West, J. S. 2006. The air spora. Springer.

Lee, D.-H., Back, C.-G., Win, N. K. K., Choi, K.-H., Kim, K.-M., Kang, I.-K., Choi, C., Yoon, T.-M., Uhm, J. Y.,
and Jung, H.-Y. 2011. Biological characterization of *Marssonina coronaria* associated with apple
blotch disease. Mycobiology 39:200-205.

Li, J., Gou, L.-x., H, X.-m., Ren, F.-p., Wei, J.-f., and An, D.-r. 2011. Effects of climate factors on the
Marssonina coronaria epidemic in Shaanxi Province and related prediction Models. Chin J Appl
Ecol 22:268-272.

Liang, Z., and Keeley, A. 2013. Filtration recovery of extracellular DNA from environmental water samples.
Environmental science & technology 47:9324-9331.

Lindner, L. 2012. Die Marssonina-Blattfleckenkrankheit jetzt auch in Südtirol. Pages 66-68 in: Obstbau
Weinbau.

Luchi, N., Ghelardini, L., Belbahri, L., Quartier, M., and Santini, A. 2013. Rapid detection of *Ceratocystis
platani* inoculum by quantitative real-time PCR assay. Applied and Environmental Microbiology
79:5394-5404.

Mc Devitt, J. J., Lees, P. S. J., Merz, W. G., and Schwab, K. J. 2004. Development of a method to detect and
quantify *Aspergillus fumigatus* conidia by quantitative PCR for environmental air samples.
Mycopathologia 158:325-335.

McCartney, H. A., Fitt, B. D., and Schmechel, D. 1997. Sampling bioaerosols in plant pathology. J Aerosol
Sci 28:349-364.

McKinney, H. H. 1923. Influence of soil temperature and moisture on infection of wheat seedlings with
Helminthosporium sativum. Journ. Agric. REs 26:195-217.
Meitz-Hopkins, J. C., G., v. D. S., Koopman, T. A., Bahramisharif, A., and Lennox, C. L. 2014. A method to
monitor airborne *Venturia inaequalis* ascospores using volumetric spore traps and quantitative
PCR. European Journal of Plant Pathology 140:527-541.
Mosimann, C., Oberhänsli, T., Ziegler, D., Nassal, D., Kandeler, E., Boller, T., Mäder, P., and Thonar, C. 2017.
Tracing of two *Pseudomonas* strains in the root and rhizoplane of maize, as related to their plant
growth-promoting effect in contrasting soils. Frontiers in Microbiology 7.
Notomi, T., Mori, Y., Tomita, N., and Kanda, H. 2015. Loop-mediated isothermal amplification (LAMP):
principle, features, and future prospects. Journal of microbiology 53:1-5.
Oberhänsli, T., Vorley, T., Tamm, L., and Schärer, H. 2014. Development of a quantitative PCR for improved
detection of *Marssonina coronaria* in field samples. Pages 187-190 in: Ecofruit. 16th International
Conference on Organic Fruit-Growing Conference Fördergemeinschaft Ökologischer Obstbau e.V.,
Hohenheim, DE.
Oberhänsli, T., Dalbosco, A., Leschenne, V., Bohr, A., Buchleither, S., Tamm, L., Wille, L., Aćimović, S. G.,
Khodadadi, F., Ryu, Y.-H., Studer, B., Schärer, H.-J., and Patocchi, A. 2021. Multiplexed SSR marker
analysis of *Diplocarpon coronariae* reveals clonality within samples from Middle Europe and
genetic distance from Asian and North American isolates. CABI Agriculture and Bioscience 2.
Park, M. Y., Sagong, D. H., Kweon, H. J., Do, Y. S., Song, Y. Y., and Lee, D. H. 2013. Influence of seasonal
incidence and defoliation degree of Marssonina blotch on fruit quality and shoot growth of
'Fuji'/M.9 Apple Tree. Korean J Hortic Sci 31:523-530.
Parker, M. L., McDonald, M. R., and Boland, G. J. 2014. Evaluation of air sampling and detection methods
to quantify airborne ascospores of *Sclerotinia sclerotiorum*. Plant Disease 98:32-42.
Persen, U., Steffek, R., Freiding, C., and Bedlan, G. 2012. Erstnachweis von *Diplocarpon mali* an *Malus*
domestica in Osterreich. Journal für Kulturpflanzen 64:168-170.
Quesada, T., Hughes, J., Smith, K., Shin, K., James, P., and Smith, J. 2018. A low-cost spore trap allows
collection and real-time PCR quantification of airborne *Fusarium circinatum* spores. Forests 9:586.
R Core Team. 2021. R: A Language and Environment for Statistical Computing. R Foundation for Statistical
Computing, Vienna, Austria, https://www.R-project.org/.
Rather, S. A., Bhat, M. A., and Mohiddin, F. A. 2017a. Influence of weather factors on development of
Marssonina leaf blotch of apple caused heterosis in cowpea for yield and quality. Indian Journal of
Ecology 44:000-000.
Rather, S. A., Bhat, M. A., and Khan, N. A. 2017b. Screening of *Malus* genotypes against Marssonina leaf
blotch. Journal of Pharmacognosy and Phytochemistry 6:134-136.
Ren, W., Liu, N., and Li, B. 2021. Development and application of a LAMP method for rapid detection of
apple blotch caused by *Marssonina coronaria*. Crop Protection 141:105452.
Sastrahidayat, I. R., and Nirwanto, H. 2016. Marssonina leaf blotch on the apple orchard in Batu, Indonesia.
AGRIVITA Journal of Agricultural Science 38.
Schärer, H.-J., Bohr, A., and Wöhner, T. 2019. Anfälligkeit von Apfelsorten gegenüber der Blattfallkrankheit
Marssonina coronaria. SZOW 19:8-12.
Schneider, R. W., Haudenshiled, J. S., Hartman, G. L., and Mahaffee, W. F. 2009. Development and use of
fluorescent antibody and qPCR protocols for the electrostatic spore trap. Phytopathology 99.
Sharma, J. N., and Gautam, D. R. 1997. Studies on premature leaf fall in apple - a new problem. Indian
Journal of Plant Protection 25:8-12.
Sharma, J. N., and Thakur, V. S. 2011. Epidemic outbreak of apple blotch disease: epidemiology and
management in Himachal Pradesh. Indian Phytopath 63:141-144.
Sharma, J. N., Thakur, V. S., Mohan, J., Paul Khurana, S. M., and Sharma, S. 2009. Epidemiology of
Marssonina blotch (*Marssonina coronaria*) of apple in India. Indian Phytopath 62:348-359.

Sharma, P. K., and Sharma, J. N. 2005. Studies on morphology and physiology of *Marssonina coronaria*
causing premature leaf fall in apple. Integrated Plant Disease Management:9-17.
Sharma, U., and Gupta, B. 2018. First report of *Diplocarpon mali*, the perfect state of apple blotch fungus
Marssonina coronaria from India. Journal of Krishi Vigyan 7:239.
Siegfried, W., Holliger, E., and Meier, H. 1996. Prognose des Rotbrenners und des Falschen
Rebenmehltaus. SZOW 132:373-374.
Sterling, M., Rogers, C., and Levetin, E. 1999. An evaluation of two methods used for microscopic analysis
of airborne fungal spore concentrations from the Burkard Spore Trap. Aerobiologia 15:9-18.
Sutton, T. B., Aldwinckle, H. S., Agnello, A. M., and Walgenbach, J. F. 2014. Compendium of apple and pear
diseases and pests.
Von Felten, A., Defago, G., and Maurhofer, M. 2010. Quantification of *Pseudomonas fluorescens* strains
F113, CHA0 and Pf153 in the rhizosphere of maize by strain-specific real-time PCR unaffected by
the variability of DNA extraction efficiency. J Microbiol Methods 81:108-115.
West, J., and Kimber, R. 2015. Innovations in air sampling to detect plant pathogens. The Annals of applied
biology 166:4-17.
White, T. J., Bruns, T. D., Lee, S. B., and Taylor, J. W. 1990. Amplification and direct sequencing of ribosomal
RNA genes for phylogenetics. PCR Protocols: A guide to methods and applications.
Wöhner, T., and Emeriewen, O. F. 2019. Apple blotch disease (*Marssonina coronaria* (Ellis & Davis) Davis)
- Review and research prospects. European Journal of Plant Pathology 153:657-669.
Wöhner, T., Bohr, A., and Schärer, H.-J. 2019. Marssonina-Anfälligkeit von Apfelsorten. Obstau 3:172-178.
Ye, J., Coulouris, G., Zaretskaya, I., Cutcutache, I., Rozen, S., and Madden, T. L. 2012. Primer-BLAST: a tool
to design target-specific primers for polymerase chain reaction. BMC bioinformatics 13:134.
Zhao, H., Huang, L., Xiao, C. L., Liu, J., Wei, J., and Gao, X. 2010. Influence of culture media and
environmental factors on mycelial growth and conidial production of *Diplocarpon mali*. Lett Appl
Microbiol 50:639-644.
Zhao, H., Han, Q. M., Wang, J., Gao, X. N., Xiao, C.-L., Liu, J., and Huang, L. L. 2013. Cytology of infection of
apple leaves by *Diplocarpon mali*. European Journal of Plant Pathology 136:41-49.
816

818   TABLE 1. Primers used in the qPCR to detect DNA of *Diplocarpon coronariae*.

Primer name[a]	Sequence (5' to 3')[b]	Mt (°C)[c]	Amplicon size (bp)	Reference
Dc_09_F	GCG TAT ACC ACC CGT GCC TA	60.4	129	This study
Dc_09_R	CTC AGA CAT CAC GTT ATT CAC ACA A	59.4		
Dc_09_P	FAM-CCT ACC TCT GTT GCT TTG GCG A-BHQ1	70.0		
ACMV_F	CCA CAG ACA AGA TCC ACT CTC C	59.7	86	Mosimann et al. (2017)
ACMV_R	CAC TCT ACT CAG GTT CCA ATC AAA G	57.9		
ACMV_P	ROX-ACA GAC AAT TCA AGA AGC GAG CCA TCC G-BHQ2	62.6		
ITS1F	CTTGGTCATTTAGAGGAAGTAA	53.2	Dependent on target species	(Gardes and Bruns 1993; White et al. 1990)
ITS2R	GCTGCGTTCTTCATCGATG	57.2		

819   [a] F, forward primer; R, reverse primer; P, fluorogenic hydrolysis probe.
820   [b] FAM, 6-Carboxyfluorescein; ROX, Carboxy-X-rhodamine; BHQ; black whole quencher
821   [c] Mt, melt (annealing) temperature.

829   TABLE 2. Numbers of conidia of *Diplocarpon coronariae* captured using rotating-arm spore traps in an apple orchard in Switzerland.[a]
830

Cultivar	Trap number	Dates (2019)[b]				Dates (2020)[c]						
		22/5 - 3/6	3/6 - 13/6	13/6 - 25/6	25/6 - 5/7	24/4 - 8/5	8/5 - 22/5	22/5 - 5/6	5/6 - 19/6	19/6 - 3/7	3/7 - 17/7	17/7– 31/7
Otava	1	7	0	335	558	0	0	70	0	0	611	41
Schneider	2	0	0	681	17	0	0	0	13	0	1	2
Otava	3	0	33	70	9	0	0	0	0	13	260	106
Otava	4	364	46	786	2412	0	0	49	16	207	171	114
Topaz	5	27	0	0	17	0	0	0	22	0	0	251
Rubinola	6	29	0	0	2	0	0	0	0	0	1	10
Rubinola	7	57	0	0	0	0	0	0	0	0	0	7
outside	8	22	0	0	0	0	0	0	0	0	0	0

831
[a] Rotating-arm spore traps were distributed within an apple orchard (Rickenbach, Switzerland) comprising 15 rows planted with different cultivars. Traps were placed
in a low branch of the apple trees (about 1.7 m). Trap Nr. 2 was installed below a 'Schneider' tree, which was at the edge of an 'Otava' tree row. The location of the
traps within the orchard is depicted in Fig. 1C. DNA of *D. coronariae* in the spore trap samples was quantified by TaqMan qPCR using primers and probe Dc_09 and
spore numbers were calculated based on a standard curve with known amounts of *D. coronariae* conidia.
[b] For 2019 the full sampling period is shown.
[c] In 2020, Rotorod traps were used for sampling from 28 February to 31 July. Since first spores were detected after 22 May, data for early series are not shown.
838
839

840
841   TABLE 3. The number of conidia of *Diplocarpon coronariae* collected in a Mycotrap spore trap above leaf litter on an
842   orchard floor in 2019 and 2020.[a]

Sample period	KOB (2019)	FiBL (2020)
15/02 – 01/03	0	-
01/03 – 15/03	0	0
15/03 – 01/04	0	14
01/04 –15/04	0	0
15/04 – 29/04	<10	0
29/04 – 16/15	<10	<10
16/05 – 04/06	0	<10
04/06 – 17/06	0	0
17/06 – 25/06	<10	<10

[a] Spore trap samples were from a Mycotrap installed above *D. coronaria*e infected leaf litter at the Competence
center for fruit crops at Lake Constance (KOB) in Ravensburg-Bavendorf (Germany) and at the Research Institute of
Organic Agriculture (FiBL) in Frick (Switzerland) in 2019 and 2020, respectively. *D. coronariae* DNA was quantified by
a TaqMan qPCR using the primers and probe Dc_09 and spore numbers were calculated based on a standard curve
with known amounts of conidia. 0, no spores detected; < 10, *D. coronariae* DNA detected, but below the limit of
quantification.

TABLE 4. Detection of *Diplocarpon coronariae* on the bark and in the buds of apple trees during winter. The apple
trees had symptoms of apple blotch the previous summer.

Apple organ	Field site[a]	Apple cultivar[b]	Tree number[c]	Normalized C_q value[d]
Bark	KOB	Topaz	18-17	38.7
Bark	KOB	Topaz	18-25	41.1
Bark	KOB	Topaz	18-37	ND
Bark	KOB	Topaz	18-38	42.3
Bark	KOB	Topaz	18-39	42.0
Bark	KOB	Topaz	18-40	ND
Bark	Rickenbach	Otava	4-10	41.1
Bark	Rickenbach	Otava	4-11	ND
Bark	Rickenbach	Otava	4-13	39.9
Bark	Rickenbach	Otava	4-14	40.71
Bark	Rickenbach	Otava	4-15	38.10
Bark	Rickenbach	Otava	4-16	38.30
Bark	Rickenbach	Rubinola	10-4	ND
Bark	Rickenbach	Rubinola	10-5	ND
Bark	Rickenbach	Rubinola	10-6	ND
Bark	Rickenbach	Rubinola	10-7	ND
Bark	Rickenbach	Rubinola	10-8	ND
Bark	Rickenbach	Rubinola	10-9	ND
Bud	KOB	Topaz	18-17	ND
Bud	KOB	Topaz	18-25	36.3
Bud	KOB	Topaz	18-37	28.0
Bud	KOB	Topaz	18-38	31.4
Bud	KOB	Topaz	18-39	36.1
Bud	KOB	Topaz	18-40	37.5
Bud	Rickenbach	Otava	4-10	33.8
Bud	Rickenbach	Otava	4-11	ND
Bud	Rickenbach	Otava	4-13	41.6
Bud	Rickenbach	Otava	4-14	33.8
Bud	Rickenbach	Otava	4-15	25.6
Bud	Rickenbach	Otava	4-16	30.7
Bud	Rickenbach	Rubinola	10-4	ND
Bud	Rickenbach	Rubinola	10-5	ND
Bud	Rickenbach	Rubinola	10-6	ND
Bud	Rickenbach	Rubinola	10-7	ND
Bud	Rickenbach	Rubinola	10-8	ND
Bud	Rickenbach	Rubinola	10-9	ND

[a] Bud and bark samples were collected from apple trees in the Rickenbach orchard (Switzerland) on 28 February
and on 13 March 2020 and from trees in the KOB orchard (Germany) on 17 March 2020.
[b] 'Topaz' and 'Otava' trees were heavily diseased with apple blotch in 2019, while 'Rubinola' trees showed no
symptoms.
[c] The tree number indicates the position of the tree within the orchard and consists of row number and tree
number within row.
[c] Cq (quantification cycle) values result from a TaqMan qPCR with primers and probe Dc_09. The APA9 plasmid was
used to normalize for DNA extraction efficiency. ND, not detected.

Figures

Fig. 1. Apple blotch (AB) development in two apple orchards and the location of the spore traps.

A, B) AB disease severity as percent damage according to the McKinney Index (I) (McKinney 1923).

A) 'Topaz' trees in two orchards (Rickenbach, Switzerland and KOB, Germany) were regularly

scored for severity of AB from June to October for five consecutive years (Rickenbach, 36 trees;

KOB, 120 trees). **B)** AB development on 'Topaz' (36 trees), 'Rubinola' (35 trees), and 'Otava' (33

trees) in the Rickenbach orchard in 2019 and 2020. **C)** AB disease scoring in the Rickenbach

orchard by middle of September 2019 and 2020. The orchard is planted with tree rows of ten

different cultivars. Each dot represents one tree. The color of the dot indicates the disease score

of the tree. The placement of the Mycotrap and rotating-arm spore traps is indicated by the

letters M and R, respectively. One rotating-arm trap was installed 50 m outside the orchard (R8).

D, E, F) Pictures of the spore traps used in this study. **D)** the Mycotrap, **E)** the rotating-arm spore

trap, **F)** potted bait plants. **G)** leaf with typical AB symptoms.

Fig. 2. *Diplocarpon coronariae* specific qPCR. **A)** Specificity test of Primer Dc_09. A PCR was

performed with DNA from different *D. coronariae* isolates, related *Diplocarpon* species (*D.*

mespili, *D. rosae*, *D. earlianum*), and with DNA of other pathogens causing Marssonina diseases

(*Drepanopeziza tremulae* and *Gnomonia leptostyla*). A 2 % TAE agaraose gel stained with ROTI®

GelStainRed (the ladder marker is a peqGOLD O'range 100bp DNA –Ladder (peqlab) **B)** *In vivo*

standard curve for calculation of spore numbers. A volume corresponding to 10^5, 10^4, 10^3, 10^2

and 10^1 conidia was pipetted onto a Vaseline coated plastic film and DNA was extracted the same

way as was done from the Mycotrap and rotating arm spore traps, i.e. using a ZymoBIOMICS®

Quick-DNA Fungal/Bacterial Microprep Kit. Samples were spiked with an APA9 plasmid for

normalization of DNA extraction and normalized Cq values are depicted. Error bars represent the

standard deviation of the mean of seven qPCR runs. The mean efficiency was 89.1 %.

Fig. 3. *Diplocarpon coronariae* (*Dc*) spore dispersal in 2019 **(A)** and 2020 **(B)**.Daily spore numbers of *Dc* captured by a Mycotrap placed within an 'Otava' tree row of apple trees. The number of spores was calculated based on normalized Cq values from the TaqMan qPCR assay with primer Dc_09 using a standard curve with known numbers of conidia. The green and the yellow lines indicate the periods when a bait plant was exposed in the orchard next to the Mycotrap and whether the bait plant developed AB symptoms (yellow) or not (green). The two blue circles indicate days where first leaves with ambiguous AB symptoms tested positive for *Dc*. The red circles indicate first unambiguous AB symptoms in the orchard. **Weather** data were obtained fromAgrometeo (location "Liebensberg") **(C, D)** including temperature (°C), relative humidity (%), precipitation (mm), and leaf wetness (light blue, <8h; dark blue, >8h).

Fig. 4. *Diplocarpon coronariae* (*Dc*) spore dispersal on the ground and in the tree crown in 2020. Daily spore numbers of *Dc* captured

by the Mycotraps placed on the ground within an 'Otava' tree row of apple trees (black bars) and within the tree canopy (grey bars).

The number of spores was calculated based on normalized Cq values from a TaqMan qPCR assay with primer Dc_09 using a standard

curve with known amounts of conidia. The green and the yellow lines indicate the periods where a bait plant was exposed in the

orchard next to the Mycotrap and whether the bait trap plants developed apple blotch symptoms (yellow) or not (green). Estimated

duration of leaf wetness is indicated as light blue (<8h) and dark blue (>8h) bands.

Fig. 5. Fruit infections with *Diplocarpon coronariae (Dc)*. A) 'Otava' apple with apple blotch symptoms hanging in a tree in September 2020 (Rickenbach, Zurich, Switzerland). B) *Dc* infected 'Otava' apple after 6 months of storage at 4°C. C) 'Florina' fruit mummy found hanging in a tree in February 2020, and tested positive for *Dc* by TaqMan qPCR. D-I) Acervuli and conidia developed on *Dc* infected 'Otava' apple collected in the field in September 2019 and stored at 4°C for six months. F-G) Longitudinal section through an acervulus. H) Conidia released from an acervulus. I) *Dc* conidia. D-F) Pictures were taken with a stereo microscope M205C (Leica Microsystems Switzerland, Heerbrugg, Switzerland). G-I) Samples were stained with cotton blue in lactic acid and pictures were taken with a Leica DM2000 LED microscope (Leica Microsystems Switzerland, Heerbrugg, Switzerland).

Fig. 6. The proposed infection cycle of *Diplocarpon coronariae* (*Dc*) in Central Europe. Based on previous knowledge and the data we presented in the current study, we hypothesize primary spores (conidia since no ascospores have been reported in Europe to date) are released in early spring and cause the first infections between the end of April and the middle of May depending on the weather conditions. Without crop protection, secondary spore inoculum is produced from the end of May onwards. The epidemic of AB spreads and leads to tree defoliation and infection of fruit in severely infested orchards. The fungus overwinters on leaf litter, but is also found on fruit mummies, in buds and on bark after winter. The fruit mummies, buds and bark might represent alternative organs for overwintering. Green, primary infections; yellow, secondary infections; blue, overwintering.