Comprehensive multi-omics single-cell data integration reveals greater

heterogeneity in the human immune system

Congmin Xu[1,†], Junkai Yang[2,†], Astrid Kosters[2], Benjamin Babcock[2], Peng Qiu[1,*], Eliver

Ghosn[2,3,4,*]

[1]Department of Biomedical Engineering, Georgia Institute of Technology and Emory

University, Atlanta, GA 30332, USA

[2]Department of Medicine, Division of Immunology, Lowance Center for Human Immunology,

Emory University School of Medicine, Atlanta, GA 30322, USA

[3]Bioinformatics Graduate Program, Georgia Institute of Technology, Atlanta, GA 30332, USA

[4]Emory Vaccine Center, Yerkes National Primate Research Center, Emory University School

of Medicine, Atlanta, GA 30322, USA

[†] Contributed equally

* Corresponding authors:

Eliver E.B. Ghosn | e-mail: eliver.ghosn@emory.edu

Peng Qiu | e-mail: peng.qiu@bme.gatech.edu

Abstract

Single-cell transcriptomics enables the definition of diverse human immune cell types across multiple tissue and disease contexts. Still, deeper biological understanding requires comprehensive integration of multiple single-cell omics (transcriptomic, proteomic, and cell receptor repertoire). To improve the identification of diverse cell types and the accuracy of cell-type classification in our multi-omics single-cell datasets, we developed SuPERR-seq, a novel analysis workflow to increase the resolution and accuracy of clustering and allow for the discovery and characterization of previously hidden cell subsets. We show that by incorporating information from cell-surface proteins and immunoglobulin transcript counts, we accurately remove cell doublets and prevent widespread cell-type misclassification. This approach uniquely improves the identification of heterogeneous cell types in the human immune system, including a novel subset of antibody-secreting cells in the bone marrow.

Introduction

Single-cell RNA sequencing (scRNA-seq) technologies have rapidly advanced in the last decade, including advances to cell-capture approaches[1-3], library preparation[4,5], and sequencing methods[1,4,6,7]. These increasingly more widely adopted technologies have significantly improved the understanding of cell heterogeneity in health and disease[4-8]. However, reliance on cellular transcriptomics alone limits the comprehensive identification of heterogenous cell populations[9]. This limitation has propelled the development of multi-omics single-cell sequencing technologies to increase the resolution and accuracy for cell subset classification.

Multi-omics single-cell sequencing technologies, such as CITE-seq[7], REAP-seq[10], and others[11], simultaneously measure gene expression (mRNA) and cell surface proteins. Additional heterogeneity of immune cell subsets can be revealed by combining single-cell gene expression with simultaneous TCR/BCR repertoire sequencing using techniques such as RAGE-seq and DART-seq[12-15]. Moreover, we show here that simultaneous measurement and comprehensive integration of transcriptomics, cell-surface protein, and cell-receptor repertoire will provide improved resolution to reveal heterogeneous cell types relevant to disease mechanisms and healthy homeostasis.

However, multi-omics technologies also present computational challenges for data integration and analysis[16-18]. Challenges include high dimensionality of the data[19], sparsity of the data[20], diversity across various omics data types[21], and technical effects between different sample batches[22,23]. Several algorithms have been developed to integrate and analyze multi-omics measurements, including linear regression models[24] and multi-omics factor analysis (MOFA)[25]. Correlated features across different -omics have been exploited by both LIGER[26] and canonical correlation analysis (CCA), as employed in the Seurat pipeline[21]. The commonality of these methods is to utilize the shared signals among different omics data types to align their distributions and achieve integration, which is an unsupervised data-driven approach. While data-driven methods have been successful for

clustering and identification of cell types, significant improvements can be made by

incorporating robust prior knowledge, such as well-established marker genes and protein

surface markers that define cell types.

This study combined our extensive expertise on high-dimensional flow cytometry

data analysis[27,28] with our own multi-omics single-cell data sets to develop SuPERR-seq

(**Su**rface **P**rotein **E**xpression, m**R**NA- and **R**epertoire-seq). SuPERR-seq is a novel,

biologically motivated approach towards the integration and analysis of multi-omics single-

cell data matrices. By combining a flow cytometry-derived "gating strategy" with the high-

dimensional analysis of scRNA-seq, SuPERR-seq increases the resolution and accuracy in

clustering algorithms and allows for the discovery of new biologically relevant cell subsets.

We first applied the "gating strategy" based on a combination of cell-surface markers and

immunoglobulin-specific transcript counts to identify major immune cell lineages. Next, we

explore the gene expression matrix following this gating strategy to resolve cell subsets

within each major immune lineage. The inclusion of this atypical "gating strategy" step also

allows for cell-doublet discrimination and dramatically enhances lineage-specific variation,

which helps better capture biological signals *within* each cell lineage. Finally, we apply

SuPERR-seq to multi-omics data sets (cell-surface protein, gene expression, VDJ repertoire)

generated in-house from human blood and bone marrow cells and directly compare its

performance to standard analysis approaches. We show that SuPERR-seq can better

leverage the power of each omics data type to identify major immune lineages more

accurately and a greater number of cell subsets within each lineage that can be confirmed

by flow cytometry, facilitating the discovery of novel immune cell types.

Results

Cell-surface proteins and immunoglobulin transcript counts identify major immune lineages in human blood and bone marrow

We generated multi-omics single-cell datasets from 12,759 human peripheral blood mononuclear cells (PBMC) and 7,426 human bone marrow (BM) cells from five individual healthy adult donors. As shown in Fig. 1A, we simultaneously captured the following three omics: total gene expression (GEX), 31 cell-surface proteins/antibody-derived tags (ADT) (Table S1), and B-cell receptor (BCR) heavy and light chain V(D)J repertoire (VDJ). The SuPERR-seq analysis pipeline can be described in two major steps shown in Fig. 1B: 1) manual biaxial gating based on the expression levels of cell-surface proteins (ADT) and total immunoglobulin (Ig) transcript counts, and 2) sub-clustering using the GEX matrix for each manually-gated population.

For the first step of the SuPERR-seq pipeline, we normalized the cell-surface (ADT) data using the DSB normalization method[29], followed by concatenating the normalized ADT matrix with the total Ig-specific UMI counts from the VDJ matrix. The integrated ADT/Ig matrix was used to identify major immune cell lineages before assessing their gene expression profile. Major immune cell lineages were identified and classified using a sequential gating strategy on biaxial plots that is similar to conventional flow cytometry data analysis (Figs. 2A and 3A). As antibody-secreting cells (ASC), also known as plasma cells, produce and secrete large quantities of immunoglobulin, they could be accurately identified based on their Ig-specific transcript counts (Figs. 2A and 3A). Of note, the SuPERR-seq pipeline was able to readily identify a rare cell cluster containing as little as eight plasma cells in the human PBMC sample. As we show below, such a rare population of plasma cells cannot be identified using current standard workflows.

In the analysis of PBMC samples, we defined gates for six major immune cell lineages: plasma cells, B cells, NK/NKT cells, Monocytes, CD4+ T cells, and CD8+ T cells (Fig. 2A, black borders). In the analysis of the BM samples, we defined gates for five major

lineages: CD138- Plasma cells, CD138+ Plasma cells, B cells, Myeloid cells, and

Hematopoietic Stem and Progenitor Cells (HSPCs) (Fig. 3A, black borders). In addition to

these main cell lineages, our gating strategy also revealed additional cell sub-clusters (Figs.

2A and 3A, gray borders), which we gated to validate downstream sub-clustering based on

the GEX data. These results were confirmed by well-established high-dimensional flow

cytometry analysis on a separate aliquot of the same samples taken prior to single-cell

encapsulation (Fig. S4)

Identifying the *major cell lineages* as the first step of the SuPERR-seq pipeline also

allows for accurate cell-doublet discrimination. We applied flow cytometry-style biaxial

feature plots on the ADT data to identify and remove cell doublets (Figs. S1 and S2). Cell

barcodes containing surface marker signals known to belong to more than one immune

lineage (e.g., co-expression of B-cell/CD19 and T-cell/CD3 markers) were considered

doublets and removed from the downstream analysis. Figs. S1 and S2 illustrate the gating

strategy used for removing doublets from the PBMC and BM data sets, respectively.

Notably, by assigning major cell lineages via a manual-gating approach before

performing the principal component analysis (PCA) of the GEX matrix, we were able to

determine that much of the data variation observed in scRNA-seq is cell-lineage specific

(Fig. 4). Key variables, including ribosomal transcript content, gene dropout rates, mRNA

abundance, and total unique gene counts, vary significantly among the cell lineages (Fig. 4,

ANOVA p<2.2e-16). We show that by gating major cell lineages before assessing variation

in the GEX matrix, we can capture biologically meaningful gene variation that occurs only

within each pre-defined major lineage. For example, the top 30 highly-variable genes

(HVGs) of total B cells are not even represented in the top 300 HVGs of the ungated total

PBMC sample. (Fig. 4D). In contrast, conventional workflows capture a range of genes

dominated by variations occurring mostly *between* cell lineages (Fig. 4D, right panel), giving

less weight to genes that would otherwise reveal further heterogeneity *within* each major

lineage. Hence, pre-selection of major cell lineages before GEX clustering enables targeted

selection of HVGs that are unique to the biology of each major cell lineage, resulting in the

identification of more and novel cell subsets.

SuPERR-seq reveals greater heterogeneity within major immune lineages

Human Peripheral Blood. Gating for major cell lineages revealed six distinct populations

within the PBMC sample (Fig. 2A, black outlines). We then further explored each major

lineage by generating subclusters using information from the GEX matrix. Briefly, we

selected a set of HVGs from within the pre-gated population. The counts of selected genes

for each cell were normalized by library size and then natural-log transformed, followed by

per-gene Z-score scaling. We then applied a singular value decomposition (SVD)

implementation of principal component analysis (PCA) to reduce the dimensionality. Left

singular values were taken as gene scores and right singular values as cell scores. Next, we

generated a K-nearest neighbors' graph (KNN, K = 30), followed by Louvain community

detection (see Methods section for detailed description). Following these steps, we obtained

the subclusters for each major lineage, herein called SuPERR-seq clusters. At a Louvain

resolution of 0.8, SuPERR-seq identified 38 clusters in human PBMC, with each major

lineage broken out into two or more subclusters, representing sub-lineage heterogeneity

(Fig. 2B).

We further investigated the 38 SuPERR-seq clusters by exploring the expression

levels of cell-surface proteins (ADT matrix) (Figs. 2C and 2D). Some ADT markers were

lineage-specific (e.g., CD19 used to classify B cells, and Ig-specific transcript counts to

classify plasma cells) and were previously used to gate major cell lineages and confirmed

the accuracy of SuPERR-seq cell classification. Other markers displayed heterogeneous

expression within lineages and were primarily used for defining and confirming subcluster

identity. We integrated the ADT and GEX data matrices by simple concatenation to generate

a joint matrix for SuPERR-seq cluster identification (Figs. 2D and S4-S9). We confirmed the

heterogenous clusters as biologically meaningful using established cell-surface lineage

markers. For example, four of the nine NK cell subclusters were identified as NKT cells

based on the co-expression of NK markers and CD3. Similarly, the monocytes were divided

into three classical and two non-classical subtypes (Fig. S4). SuPERR-seq was also able to

accurately identify T-regulatory cells (Treg) as a subtype of CD4+ T cells based on their high

expression of CD25 (ADT matrix) and FOXP3 (GEX matrix), and low expression of CD127

(ADT matrix) (Figs. 2A and S5).

Remarkably, SuPERR-seq readily identified a cluster of rare plasma cells (antibody-

secreting cells) in the blood even though we captured only eight plasma cells within this

cluster (Fig. 2). This level of resolution and accuracy in identifying rare plasma cells was

possible by analyzing the total Ig-specific transcript counts from the VDJ matrix. As plasma

cells produce large quantities of immunoglobulin transcripts, they were readily identified

based on ~2.5 $\log_{10}$-fold increase in total Ig-specific transcripts compared to B cells (Fig.

2A).

Five sub-clusters were identified within the B-cell major lineage, all of which could be

mapped back to previously described B-cell subsets[30,31] (Fig. 2D). Specifically, clusters 1

and 2 lack surface CD27, a canonical B-cell memory marker[32], and the presence of IGHD

and IGHM (IgD and IgM transcripts, respectively) identify these clusters as naïve B cells[33]

(Fig. 2D). The clusters 3 and 4 (CD27+/TACI+) are memory B cells. The presence of CD95

and the absence of IGHM further characterize the cluster 3 as class-switched memory B

cells[33]. Finally, cluster 5 is unswitched (IgM+) memory B cells (Fig. 2D). All other major

lineages were classified using the same approach, and the results are described in Table 1.

Human Bone Marrow. SuPERR-seq identified 34 unique clusters in human BM cells (Figs.

3A and 3B). Of note, SuPERR-seq was able to match the B-cell subclusters to the different

stages of their developmental pathway[34] (Fig. 5). The cell-surface expression of CD10, along

with DNTT and CD34 gene expression (GEX) transcripts, classified cluster 11 and cluster 13

as Early Pro-B (a.k.a., pre-pro-B) and Late Pro-B (a.k.a., pre-BI), respectively. The lack of

DNTT mRNA transcripts in cluster 10 indicates a Large Pre-B stage (a.k.a., pre-BII), which is

followed by the Small Pre-B stage (a.k.a., pre-BII) represented by clusters 6 (VPREB1[hi]) and

7 (VPREB1[lo]). The Transitional (T)1/T2/T3 and Naive B cells were defined by their surface

expression of CD21[35]. Finally, the mature class-switched memory B cells were identified by

the expression of IGHA1/IGHG1 immunoglobulin and surface CD27[36] (Fig. 5). The SuPERR-

seq workflow also was able to identify the developmental pathway of neutrophils,

monocytes, and erythrocytes starting from the hematopoietic stem cells (HSC) (Figs. S8 and

S9). The classification results for the other subclusters are summarized in Table 2.

SuPERR-seq workflow reveals new subsets of antibody-secreting cells in the
human bone marrow

The SuPERR-seq pipeline readily and unambiguously identified four biologically

distinct subsets of human antibody-secreting cells (a.k.a., plasma cells) in the adult bone

marrow (BM) of healthy donors (Figs. 3 and 6).

The determinant feature of a plasma cell is its ability to produce and secrete large

quantities of immunoglobulins (Ig) (i.e., antibodies). The SuPERR-seq workflow leverages

this unique plasma cell feature by quantifying the absolute UMI counts of Ig-specific genes

(IGH + IGL) from the V(D)J repertoire matrix (Fig. 3A). Unlike the GEX matrix, the V(D)J

matrix provides a more accurate count of total Ig-specific transcripts produced by plasma

cells because the V(D)J data matrix is generated from a separate library using Ig-specific

primers[8].

Remarkably, the number of total Ig-specific transcripts (Ig-specific UMI counts)

detected in plasma cells is ~2.5 $\log_{10}$-fold higher than in B cells (Figs. 3A and 3C), allowing

for unambiguous and straightforward identification of total plasma cells. Next, we separated

the total plasma cells into two subsets, based on the cell-surface protein expression (ADT

matrix) of CD138, I canonical plasma-cell marker expressed by some, but not all, BM plasma

cells[30,37]. Finally, we used the GEX data matrix to further subcluster the CD138+ and

CD138- plasma cell subsets based on their transcriptomic profile, revealing a total of 4

distinct subsets of human plasma cells (Figs. 3 and 6).

To facilitate the identification of the differential gene expression (DGE) that

distinguish each of the four plasma cell subsets, we first removed from the GEX matrix all

the mRNA transcripts (UMI) derived from immunoglobulin genes (immunoglobulin genes

represent more than 50% of the total transcripts/UMI recovered from I plasma cell). We then

log-normalized the immunoglobulin-depleted GEX matrix and performed DGE analysis using

the Wilcoxon Rank-Sum Test. The resulting differentially-expressed genes readily defined

unique biological processes for each plasma cell cluster (Fig. 6). Cluster 1: CD138-

/Ig^{hi}/PRDM1-; cluster 2: CD138-/HLA-DR^{hi}/SDC1^{lo}/G2-M phase; cluster 3: CD138+/IRF4^{hi};

cluster 4: CD138+/XBP1^{hi}/SDC1^{hi} (Figs. 6A and 6C). Clusters 2 and 4 show characteristics

that are similar to previously identified human BM plasma cell subsets, in which they were

described as Fraction I and Fraction B, respectively[37]. Notably, cluster 1 represents a unique

plasma cell subset, in that >75% of its total mRNA transcripts represent immunoglobulin

genes (Fig. 6B). This large proportion of immunoglobulin gene transcripts suggests these

cells are allocating most of their metabolic activity to produce and secrete antibodies.

Indeed, pathway analysis (Reactome Pathway Database)[38] using the differentially expressed

genes (DGE) for cluster 1 revealed signals mainly for protein metabolism (Fig. 6F).

Plasma cells develop from activated B cells through a dynamic cell-differentiation

process, leading to the down-regulation of B-cell identity genes, such as PAX5, and up-

regulation of well-described plasma cell genes, including PRDM1, SDC1, XBP1, and IRF4[37].

These genes are considered canonical plasma cell genes, and, hence, they are used in

scRNA-seq experiments to identify and classify plasma cells based on their transcriptomics

(GEX matrix). Surprisingly, we found that not all plasma cell clusters express the canonical

plasma cell genes. Of note, cluster 1 does not express IRF4 and PRDM1. The absence of

these canonical genes in cluster 1 was not due to the overrepresentation of immunoglobulin

(Ig) genes because IRF4 and PRDM1 still were not detected after normalizing the GEX

matrix without the immunoglobulin genes (Fig. 6C).

By comparing the immunoglobulin isotypes and subclasses (IgM, IgD, IgG1-4, IgA1-

2, IgE) expressed by each plasma cell subset, we found that cluster 2 contains plasma cells

of multiple isotypes/subclasses. In contrast, cluster 1 is more homogeneous, containing

mainly IgG1 and IgA1 (Fig. 6D). Furthermore, we observed that cluster 1 is composed of

plasma cell repertoire clones (defined by their BCR CDR3 amino acid sequences) shared

among cluster 3 and cluster 4 (Fig. 6E). Finally, pathway analysis (Reactome Pathway

Database) revealed unique biological processes and genetic programs that define each

plasma cell subset (Fig. 6F). Notably, cluster 4 expresses genes involved in the cell cycle

and programmed cell death, while cluster 3 appears to be actively responding to

environmental stimulation.

In sum, the SuPERR-seq pipeline readily and unambiguously identified four

biologically distinct human plasma cell subsets in the adult BM. These findings further

support the need for a comprehensive multi-omics single-cell data integration and reveal the

potential shortcomings of relying solely on one omics data type (i.e., transcriptomics) to

identify and classify cell types as done by existing approaches in the literature. In the

following section, we provide specific examples of how the SuPERR-seq workflow

outperforms existing approaches.

267 Benchmarking of SuPERR-seq against conventional methods reveals reduced
268 cell-type misclassification, and superior ability to resolve cell doublets and
269 classify new cell types

As we described above, the first step of the SuPERR-seq workflow, in which we use

cell-surface proteins in sequential biaxial plots to identify major cell lineages, also allows for

accurate cell-doublet discrimination (Figs. S1, S2, 7A, and 7B). Our SuPERR-seq approach

successfully identified and removed 370 cell doublets, while the conventional approach[39] of

trimming cells with very high mRNA transcripts (i.e., cell barcodes with greater than the

mean UMI value + 4 standard deviation) as potential cell doublets identified only 42 doublets

in PBMC. For BM, the SuPERR-seq approach identified 108 cell doublets, and the

conventional approach identified only one cell doublet. (Fig. 7A). Strikingly, every plasma cell

we identified in PBMCs would have been trimmed/removed by the conventional approach of

removing cells with mean + 4 SD UMI counts (Fig. 7A). Unlike the standard/conventional

approach, the SuPERR-seq workflow readily recognized the PBMC plasma cells as single

cells (confirmed by their unique VDJ repertoire) containing high UMI counts (Fig. 7A). The

true cell doublets removed by SuPERR-seq would otherwise have been missed by

conventional approaches and erroneously included as single cells in downstream GEX

analysis, as visualized on the UMAP plot (Fig. 7C).

We further compared our approach with current computational methods explicitly

designed to remove cell doublets in scRNA-seq datasets. We tested the standard workflow

of DoubletFinder[40] and compared it with the cell doublets defined by the SuPERR-seq

method (Fig. 7D). Although DoubletFinder outputs a similar number of cells compared to the

SuPERR-seq method, the results of the two methods did not overlap, mainly due to false

positives and false negatives from DoubletFinder. Notably, SuPERR-seq identified hundreds

of true cells doublets (i.e., cell barcodes that co-express markers of more than one cell

lineage) that were missed by the DoubletFinder workflow (Fig. 7D). Interestingly,

DoubletFinder also identified PBMC plasma cells as cell doublets due to their high UMI

counts. However, we confirmed that PBMC plasma cells represent singlets by both their

surface markers (ADT matrix) and their unique V(D)J repertoire (VDJ matrix).

Next, we compared our SuPERR-seq pipeline with the commonly used Seurat

package V3[22] using default parameters. For the PBMC sample, Seurat identified 24 clusters

compared to 38 clusters identified by the SuPERR-seq pipeline (Fig. S10). Seurat workflow

identified 22 clusters for the BM sample compared to 34 clusters identified by the SuPERR-

seq pipeline (Fig. S11). When we directly compared the cluster assignment of the same cell

barcodes from each pipeline (Figs. S10 and S11), multiple Seurat clusters were further

divided by the SuPERR-seq workflow into several biologically meaningful clusters (Figs. 5, 6,

S4-S9). The ability of the SuPERR-seq pipeline to identify greater and novel subsets of

immune cells can be explained by the advantages of pre-gating major immune cell lineages

based on cell-surface markers and Ig-specific transcripts before exploring the GEX matrix.

As we show in Fig. 4, pre-gating major lineages reveal the gene variation that occurs only

within the pre-defined lineage instead of gene variation observed *across all* major lineages.

The select list of highly variable genes (HVGs) that occur only within the pre-defined

lineages generates clusters of biologically meaningful cell types dominated by lineage-

specific genes (Fig. 4D).

Importantly, our benchmarking analysis reveals a significant number of cell

misclassifications (i.e., a cluster containing cells from different major lineages) generated by

the standard Seurat workflow. For example, the Seurat workflow generated three clusters (5,

9, and 15) (Fig. S10) containing a mixture of cell lineages, including CD8+ T cells and NK

cells. In contrast, SuPERR-seq correctly clustered and classified these cells separately.

Similarly, Seurat cluster 3 mixed CD4+ and CD8+ T cells, while SuPERR-seq correctly

identified and separated these different cell types (Fig. S10). Notably, such cell

misclassification artifacts are not rare and commonly occur when clustering cells in high-

dimensional space using the GEX data matrix as done by most, if not all, scRNA-seq

analysis workflow[41,42].

To quantify the extent of cell misclassification in conventional data analysis

approaches, we developed a new scoring system named Cell Misclassification Statistics

(CMS) score[43]. Briefly, we consider the biaxial feature plots on the cell surface (ADT) protein

data (Figs. S1 and S2) to represent a "gold standard," as this iterative nature of SuPERR-

seq clustering and cell-type assignment prohibits cells from inappropriately co-clustering with

a separate lineage. This approach is borrowed from the well-established biaxial gating

strategy of flow cytometry analyses. We then compare these cell-type identities to those

from a standard Seurat workflow and consider the proportion of cells that change

classification. This is reported as a "CMS score," which provides a statistical measure of

uncertainty in the cell-type assignment steps of the Seurat workflow. The PBMC and BM

clusters generated by Seurat showed a CMS score of 0.0984 and 0.0594, respectively,

indicating that 9.84% of PBMCs and 5.94% of BM cells were misclassified by the

conventional approach (Fig. 8A).

To determine whether cell misclassification in Seurat clusters could confound

interpretation of downstream differential gene expression (DGE) analysis, we explored the

PBMC cluster 5 generated by Seurat (Fig. 8C). DGE analysis is a commonly used method to

interpret biological differences between clusters or cell types[44]. We hypothesized that

running DGE analysis on a cluster with contaminating cell types could generate misleading

results even at low contamination numbers. The Seurat PBMC cluster 5 contains mainly NK

cells, but it is also contaminated/mixed with both NKT and T cells. Indeed, by analyzing the

DGE list of cluster 5 before removing the contaminating cells (NKT and T), the *TRDV1* gene

appeared as highly expressed for this cell population (Fig. 8C). However, *TRDV1* is

expressed on gamma-delta T cells[45], not on NK cells. These results suggest that widespread

cell misclassification often observed in conventional data analysis workflow can potentially

confound data interpretation.

Finally, when comparing the new SuPERR-seq-identified plasma cells with

conventional Seurat workflow, we observed major differences. While SuPERR-seq readily

and accurately identified a cluster of eight plasma cells in PBMCs (Fig. 2A), the conventional

workflow fails to identify these rare cells. The inability of conventional approaches to

identifying these rare plasma cells can be explained by: a) the conventional approach of

removing cell doublets based on total UMI counts incorrectly removes plasma cells due to

their naturally high mRNA content (Fig. 7A), and b) even when plasma cells are not removed

as cell doublets and instead included in downstream clustering analysis, the plasma cells are

assigned to a cluster of mixed cell types (Fig. S10B).

Unlike the PBMC, BM cells contain significantly more plasma cells. In this case,

SuPERR-seq identified four biologically distinct plasma cell subsets, while Seurat workflow

identified one cluster (Figs. 6 and S11). Of note, the plasma cell cluster generated by the

Seurat workflow contained a mixture of cell types, including B cells and HSPCs (Fig. S11B).

In addition, a fraction of the plasma cells identified by the SuPERR-seq based on Ig-specific

transcripts (VDJ matrix) were misclassified by the Seurat workflow and appeared spread

across multiple clusters on the UMAP (Fig. S11B). To determine whether the conventional

workflow could still identify the four SuPERR-seq plasma cell subsets, we performed a

second iteration of clustering (re-cluster) on the original Seurat plasma cell cluster.

Surprisingly, even after the second iteration of clustering, the conventional workflow was

able to identify only three BM plasma cell subsets (Fig. S11C). More importantly, the three

plasma cells clusters identified by the conventional approach remained a mixture of B cells

and HSPCs (Fig. S11C). For the fairness of comparison, the parameters used for running

the Seurat and SuPERR-seq workflows were set as the default parameters described in

Methods. The re-clustering approach on the original Seurat clusters was limited to two

iterations.

Taken together, the results from the benchmarking analysis show that the SuPERR-

seq workflow not only identifies more and novel clusters but also improves the clustering

purity and the accuracy of biological interpretation.

Discussion

A core component of any single-cell RNA-seq analysis is the identification and classification of cell types. The most popular and widely-used approach relies on principal component analysis (PCA) derived from a set of highly variable genes (HVGs), followed by nearest-neighbors graph-based clustering. The resulting clusters are then assigned to a cell type and biological function via manual annotation, using the mRNA transcript counts as a guide. In existing approaches, the selection of HVGs for downstream clustering analysis is often dominated by lineage-specific markers. These markers often capture variance that exists *between* cell lineages (e.g., the variance that distinguishes T cells from B cells) but often misses variance that exists *within* each cell lineage (e.g., the variance that distinguishes the various T-cell subsets, including effector memory from central memory, stem memory, naïve, etc.) (Fig. 4). Thus, an improved method to select HVGs *within* each cell lineage is needed to allow the discovery of new biologically meaningful cell types.

Moreover, selecting genes with high variance across all lineages (instead of *within* lineages) as the very first step in data analysis (as in many existing approaches) carries the risk of inappropriately grouping together cells with similar gene expression but arising from separate cell lineages, resulting in cell misclassification (Fig. 8). Cell misclassification is a common occurrence in conventional scRNA-seq analyses that rely on a single omics data type (i.e., transcriptomics) and can be caused by a mathematical phenomenon known as the curse of dimensionality[41,42], inadvertently misguiding biological interpretations (Fig. 8, see CMS scores). Another contributing factor to cell misclassification is the presence of cell doublets in the final dataset. Despite recent advances in microfluidics that precisely generate droplets containing single cells, a significant fraction of the droplets can still contain more than one cell[2,71]. Current approaches to remove cell doublets bioinformatically simply remove droplets that contain very high mRNA transcript counts (i.e., outliers)[21]. However, as we show here, many cell doublets contain average mRNA counts and can mistakenly be carried over to downstream analysis (Fig. 7).

To address these pitfalls of existing methods, we presented here the SuPERR-seq

pipeline. In SuPERR-seq, we simultaneously apply information gained from triple-omics

sequencing, namely gene expression (GEX), surface-protein markers (ADT), and antibody

production/repertoire (VDJ). As a quality control measure and to prevent downstream

misclassification, the first step of SuPERR-seq is to perform a "manual gating" similar to

standard flow cytometry analysis. ADT and V(D)J matrices provide measurements of both

protein surface markers and antibody production that can reliably classify cells into clearly

distinct major lineages by canonical markers. This gating step ensures that any subsequent

cell clusters generated are composed of a single lineage, allowing for a more accurate

investigation of clusters as a representation of cell type. The second step of the SuPERR-

seq is to examine each lineage independently, both from one another and from the first

gating steps. Rather than selecting genes with high variance across all cell lineages, we

select HVGs only from within each manually-gated lineage. These HVGs inform a PCA and

subsequent nearest-neighbors graph-based clustering. In our analysis of PBMC and BM

samples, we show that the SuPERR-seq approach reveals additional cell types (and likely

cell states) as compared to a standard approach. We reason that this is mainly due to the

ability of SuPERR-seq to select only the relevant HVGs for each well-defined lineage and

ignore extra sources of variance that may not be informative within each cell lineage.

We recognize the difficulty and expense of generating triple-omics data for every

sample, as well as the limitations of applying the SuPERR-seq approach to older data sets

for which ADT and V(D)J matrices are unavailable. Also, some samples might not include B

and/or T cells, which means there will not be V(D)J matrix. Thus, to broaden the application

of SuPERR-seq principles to datasets without triple-omics available, we propose a gene-

based (GEX) recursive analysis approach, which carries some of the same benefits of

SuPERR-seq. We applied this approach to the same BM dataset explored by SuPERR-seq

and identified improvements to standard approaches but also limitations. Focusing on the

plasma cells, the recursive application of Seurat, which only identified one contaminated

(misclassified) plasma cell cluster in the first iteration, now generated three sub-clusters of

plasma cells. However, this recursive application of Seurat was not able to accurately isolate

the plasma cells from the cell contaminants, in that the three resulting sub-clusters of plasma

cells represented a mixture of plasma cells, B cells, and HSPCs (Fig. S11C).

In contrast, SuPERR-seq readily and accurately identified four subsets of plasma

cells that were confirmed based on the three omics (GEX, ADT, VDJ), including the high

antibody transcript counts (~2.5 $\log_{10}$-fold higher antibody transcripts than B cells). The

additional plasma cell subset identified by SuPERR-seq is of great biological importance, as

revealed by the Reactome pathway analysis generated on the set of DGE that distinguishes

each plasma cell subset (Fig. 6). Therefore, the integration of three omics in the SuPERR-

seq provides unique information that cannot be achieved by recursive strategies for defining

HVGs. This conclusion is in agreement with the latest implementation of Seurat Version 4,

which also integrates ADT measurements in the analysis pipeline[21], further supporting the

SuPERR-seq approach of multi-omics integration for defining cell clusters.

Taken together, we developed a comprehensive multi-omics single-cell data

integration and analysis pipeline that mitigates or resolves pitfalls in existing approaches and

allows for the discovery of novel biologically meaningful cell subtypes in the human immune

system.

Methods

Subjects and specimen collection

Peripheral blood was collected from healthy adult donors through Emory University's
Children's Clinical and Translational Discovery Core. Bone marrow samples were collected
from healthy adult donors through Emory University under IRB00066294 or obtained from
AllCells (Alameda, CA).

Cell preparation

Peripheral blood mononuclear cells (PBMCs) were isolated from peripheral venous blood
using EasySep™ Direct Human PBMC Isolation Kit (# 19654). B cells were enriched from
hip bone marrow (BM) using EasySep™ Direct Human PBMC Isolation Kit (# 19654)
followed by EasySep Human B-Cell Enrichment Kit II without CD43 Depletion (# 17963)
following manufacture's protocol. Up to $1x10^6$ cells per donor were incubated with Fc block
(Myltenyi) on ice for 10 min. followed by staining with a mix of 31 oligo-conjugated
(barcoded) antibodies (see Table S1 for full list) for 30 min on ice, in custom-made RPMI-
1640 media deficient in biotin, L-glutamine, phenol red, riboflavin, and sodium bicarbonate,
and containing 3% newborn calf serum (def-RPMI-1640), followed by two washes in def-
RPMI-1640/0.04% BSA. Cells were resuspended at a concentration of 1200-1500 cells/uL in
custom RPMI-1640/0.04% BSA and passed through a 20-um nylon filter prior to loading onto
a Chromium Controller (10X Genomics, Pleasanton, CA).

Single-cell encapsulation, library generation, and sequencing

Cells were loaded to target encapsulation of 6000 cells. Gene expression (GEX), Antibody
Derived Tag (ADT), and V(D)J libraries were generated using the Chromium Single Cell 5′
Library & Gel Bead Kit v1 with feature barcoding (10X Genomics, Pleasanton, CA) following
the manufacturer's instructions. Gene expression libraries were sequenced together on a
NovaSeq 6000 platform (Illumina, San Diego, CA). Pooled ADT and pooled V(D)J libraries
were sequenced separately on a Next-seq platform (Illumina, San Diego, CA). The

479 sequencing depth is shown in Table S2.

Multi-omics single-cell data preprocessing

481 10X Genomics CellRanger V3.1.0 was used to perform barcode processing and single-cell

482 5′ unique molecular identifier (UMI) counting. Reads from GEX and ADT libraries were

483 processed simultaneously using "cellranger count," while reads from V(D)J libraries were

484 aligned by running "cellranger vdj."

Gene expression (GEX)

The scRNA-seq expression datasets were integrated into two final matrices, one for PBMC

and another for BM. To remove potential batch effects, we used the canonical correlation

analysis (CCA) to integrate the three individual PBMC samples into one final PBMC dataset

and combine the two individual BM samples into one final BM dataset. Before integration, we

performed library-size scaling and log transformation for each individual sample. Next, we

identified the top 2,000 highly variable genes for each individual sample using regularized

negative binomial regression[72], followed by CCA to find anchors and integrate cells from

individual samples into one integrated dataset. Finally, we performed per-gene z-score

normalization for each integrated dataset. Genes that were expressed in less than three

cells, cells with less than 200 genes, and cells with more than 20% of mitochondrial gene

counts were removed from downstream analysis. GEX raw count matrices are available

through NCBI GEO, accession numbers XXX (PBMC), and XXX (BM).

Antibody-derived tags (ADT)

The surface protein expression (ADT) data were normalized using the DSB normalization[29].

Cell barcodes in the CellRanger unfiltered matrix that expressed less than 40 genes and less

than 100 total UMIs were considered background. The barcodes from CellRanger

"filtered_feature_bc_matrix" were considered true cells. By subtracting the mean value of the

background population and then regressing out the cell-cell technical variation[29], the ADT

values representing background expression levels were centered around zero. ADT raw

count matrices are available through NCBI GEO, accession numbers XXX (PBMC), and

XXX (BM).

Antibody repertoire (VDJ)

The CellRanger-generated "all_contig_annotations.csv" (unfiltered) and

"filtered_contig_annotations.csv" files were used to generate new features that can be

merged with the corresponding ADT matrix. Briefly, we summed the total UMIs of each cell

barcode in the "all_contig_annotations.csv" file to generate a feature called "Total Ig

Transcripts" that represent the total UMIs of immunoglobulin heavy and light chains. In

addition, we generated a binary feature called "Productive VDJ" that identifies whether a cell

barcode with V(D)J information is contained in the CellRanger "filtered" annotation file and

hence considered true B cell/plasma cell. B-cell receptor repertoire V(D)J raw count matrices

are available through NCBI GEO, accession numbers XXX (PBMC), and XXX (BM).

Manual gating

Manual gating was performed based on the two VDJ features (i.e., total Ig-specific

transcripts and productive VDJ) and the 31 surface protein features in the normalized ADT

data. The major cell lineages (six for PBMCs and five for BM cells) were identified and

manually gated using a customized strategy of biaxial plots implemented by MATLAB script

(https://github.com/LindaXucongmin/SuPERR-seq.git). The markers and the gating hierarchy

were determined by prior knowledge of variations of surface protein expression across

various major cell lineages (Figs. 2A and 3A). Cell doublets were manually "gated out"

based on the expression of two or more major cell-lineage markers (Figs. S1 and S2).

Clustering on gene expression matrix

Each major cell lineage identified by the manual gating was next clustered based on the

integrated gene expression matrix. Highly variable genes were computed for cells in each

major cell lineage using the variance-stabilizing transformation (VST) method[72], and

expression data of each gene was transformed to zero mean and unit variance. We then

performed PCA on the highly variable genes of each main lineage, and the top 30 PCs were

used for downstream analysis. In the low-dimensional PCA space, a K-nearest neighbor

(KNN) graph was constructed based on the Euclidean distance between cells, with K=30.

The KNN graph was further converted by Jaccard to derive a refined version of the cell-cell

similarity matrix, followed by Louvain community detection algorithm[73] to define cell clusters

in each major cell lineage.

Differential gene expression analysis

The differential gene expression (DGE) analysis was done using the "FindAllMarkers"

function of Seurat V3. Statistical significance was tested using the Wilcoxon Rank-Sum test,

with p_val_adj<0.05 (p-value after Bonferroni correction). For the DGE analysis of the BM

plasma cell clusters, we first removed the immunoglobulin-specific UMIs from the GEX

matrix and log-normalized the resulting matrix to allow for the discovery and exploration of

non-immunoglobulin plasma cell genes.

Pathway analysis

545   The DGE list for each BM plasma cell cluster was uploaded to the Reactome pathway

546   database[38] (https://reactome.org/PathwayBrowser/) to visualize active biological pathways

547   and regulatory processes in each plasma cell cluster. The firework diagrams were cropped

548   so that only the hit pathways were preserved.

Comparison of cell doublets identified by DoubletFinder and SuPERR-seq

The DoubletFinder workflow was run independently on the gene expression matrices from

each PBMC and BM sample, following the standard workflow described in the original

publication[40]. Briefly, we used the bimodality coefficient (BC_{MVN}) maximization to select the

optimal proportion of K-nearest neighbors (pK) for each sample. We then set the proportion

of artificial doublets to pN=25% and selected the top 10 PCs to identify cell doublets. The

estimated doublet formation rate from the microfluidics was determined for each sample

based on the 10X Genomics user guide[74]. Next, the clustering results generated by the

Seurat integration workflow were used to reduce the false-positive rate. Since we did not

explicitly classify every cell barcode during manual gating, we removed the cell barcodes

that were not present in the final singlet and doublet population defined by the SuPERR-seq

workflow.

Cell Misclassification Statistic

To generate the Cell Misclassification Statistic (CMS) score[43], we first assign cell type labels

to each cluster in a merged, multi-sample dataset. This can be accomplished either by

following a gating strategy and assigning cell types to entire clusters which fall within each

gate or by transferring cluster labels from a previously gated dataset containing shared cells.

The transferred labels can be used to assist cell type identification but should always be

validated by marker expression following the gating strategy. After cell types have been

recorded for the merged samples (target dataset), we match a vector of cell identities from

the same sample, processed by a different workflow (reference dataset). We then compare

the target cell IDs to a vector of the reference cell type IDs. The CMS score generated is 1 -

the sum of matching values (total number of cell type ID matches) between the two cell-

typing replicates, divided by the total number of cells. The result is a measure of the fraction

of cells that have changed cell-type ID between single-cell analysis workflows, such that a

CMS=0 indicates that no cells have changed their classification/cell-type ID.

Acknowledgments

This study was supported in part by Georgia Clinical and Translational Science Alliance (CTSA) through the National Center for Advancing Translational Sciences of the National Institutes of Health under Award number NIH UL1TR002378 (E.G.); National Science Foundation CCF1552784 and CCF2007029 (P.Q.); Pediatric Research Alliance, Center for Transplantation, and Immune-Mediated Disorders, Children's Healthcare of Atlanta (E.G.); and Lowance Center for Human Immunology (E.G.). P.Q. is an ISAC Marylou Ingram Scholar and a Carol Ann and David D. Flanagan Faculty Fellow. We thank Sachin Kumar (Emory University) for providing the Python script used to generate the Circos plots and Matthew C. Woodruff (Emory University) for the R script used to generate the V(D)J input files for the Circos plots. Flow cytometry data were collected at the Emory's Pediatrics/Winship Flow Cytometry Core (access supported in part by Children's Healthcare of Atlanta). Single-cell libraries were sequenced at the Emory Integrated Genomics Core (EIGC), which is subsidized by the Emory University School of Medicine and is one of the Emory Integrated Core Facilities; Parker H. Petit Institute for Bioengineering and Bioscience at the Georgia Institute of Technology; PerkinElmer Genomics Inc. We thank Dr. F. Eun-Hyung Lee (Emory University), Sang N. Le, and Mindy R. Hernández for kindly providing the bone marrows aspirates from healthy adult donors through Dr. Lee's Emory IRB protocol, and Emory University's Children's Clinical and Translational Discovery Core (CCTDC) for providing peripheral blood samples from healthy donors.

Author Contributions

600 **Author Contributions**
601
E.G. and P.Q. conceived the idea and experimental design, interpreted the data, and wrote

the final draft. C.M. and J.Y. generated, analyzed, and interpreted the multi-omics single-cell

datasets and generated the figures. J.Y. performed the preprocessing workflows to obtain

final count matrices for each single-cell dataset. A.K. received and processed the human

samples and generated the single-cell libraries and flow cytometry datasets, and analyzed

the data. B.R.B. provided input into data analysis and generated the CMS scores. All authors

contributed to the writing of the manuscript and approved the final version.

Competing Interests

610 **Competing Interests**
611
612 The authors declare no competing interests.

References

613 **References**
614
1 Evan *et al.* Highly Parallel Genome-wide Expression Profiling of Individual
Cells Using Nanoliter Droplets. *Cell* **161**, 1202-1214,
doi:10.1016/j.cell.2015.05.002 (2015).
2 Allon *et al.* Droplet Barcoding for Single-Cell Transcriptomics Applied to
Embryonic Stem Cells. *Cell* **161**, 1187-1201, doi:10.1016/j.cell.2015.04.044
(2015).
3 Utada, A. S., Fernandez-Nieves, A., Stone, H. A. & Weitz, D. A. Dripping to
Jetting Transitions in Coflowing Liquid Streams. *Physical Review Letters* **99**,
doi:10.1103/physrevlett.99.094502 (2007).
4 Picelli, S. *et al.* Smart-seq2 for sensitive full-length transcriptome profiling in
single cells. *Nature Methods* **10**, 1096-1098, doi:10.1038/nmeth.2639 (2013).
5 Hashimshony, T., Wagner, F., Sher, N. & Yanai, I. CEL-Seq: Single-Cell RNA-
Seq by Multiplexed Linear Amplification. *Cell Reports* **2**, 666-673,
doi:10.1016/j.celrep.2012.08.003 (2012).
6 Habib, N. *et al.* Massively parallel single-nucleus RNA-seq with DroNc-seq.
Nature Methods **14**, 955-958, doi:10.1038/nmeth.4407 (2017).
7 Stoeckius, M. *et al.* Simultaneous epitope and transcriptome measurement in
single cells. *Nature Methods* **14**, 865-868, doi:10.1038/nmeth.4380 (2017).
8 Zheng, G. X. Y. *et al.* Massively parallel digital transcriptional profiling of
single cells. *Nature Communications* **8**, 14049, doi:10.1038/ncomms14049
(2017).
9 Tran, H. T. N. *et al.* A benchmark of batch-effect correction methods for
single-cell RNA sequencing data. *Genome Biology* **21**, doi:10.1186/s13059-
019-1850-9 (2020).
10 Peterson, V. M. *et al.* Multiplexed quantification of proteins and transcripts in
single cells. *Nature Biotechnology* **35**, 936-939, doi:10.1038/nbt.3973 (2017).
11 Lee, J., Hyeon, D. Y. & Hwang, D. Single-cell multiomics: technologies and
data analysis methods. *Experimental & Molecular Medicine* **52**, 1428-1442,
doi:10.1038/s12276-020-0420-2 (2020).
12 Singh, M. *et al.* High-throughput targeted long-read single cell sequencing
reveals the clonal and transcriptional landscape of lymphocytes. *Nature*
Communications **10**, doi:10.1038/s41467-019-11049-4 (2019).

13 Horns, F., Dekker, C. L. & Quake, S. R. Memory B Cell Activation, Broad Anti-
influenza Antibodies, and Bystander Activation Revealed by Single-Cell
Transcriptomics. *Cell Reports* **30**, 905-913.e906,
doi:10.1016/j.celrep.2019.12.063 (2020).

14 Zemmour, D. *et al.* Single-cell gene expression reveals a landscape of
regulatory T cell phenotypes shaped by the TCR. *Nature Immunology* **19**,
291-301, doi:10.1038/s41590-018-0051-0 (2018).

15 Yermanos, A. *et al.* Single-cell immune repertoire and transcriptome
sequencing reveals that clonally expanded and transcriptionally distinct
lymphocytes populate the aged central nervous system in mice. *Proc Biol Sci*
288, 20202793, doi:10.1098/rspb.2020.2793 (2021).

16 Luecken, M. D. & Theis, F. J. Current best practices in single‐cell RNA‐seq
analysis: a tutorial. *Molecular Systems Biology* **15**, e8746,
doi:10.15252/msb.20188746 (2019).

17 Stuart, T. & Satija, R. Integrative single-cell analysis. *Nature Reviews*
Genetics **20**, 257-272, doi:10.1038/s41576-019-0093-7 (2019).

18 Colomé-Tatché, M. & Theis, F. J. Statistical single cell multi-omics integration.
Current Opinion in Systems Biology **7**, 54-59, doi:10.1016/j.coisb.2018.01.003
(2018).

19 Yu, P. & Lin, W. Single-cell Transcriptome Study as Big Data. *Genomics,*
Proteomics & Bioinformatics **14**, 21-30, doi:10.1016/j.gpb.2016.01.005 (2016).

20 Qiu, P. Embracing the dropouts in single-cell RNA-seq analysis. *Nature*
Communications **11**, doi:10.1038/s41467-020-14976-9 (2020).

21 Hao, Y. *et al.* Integrated analysis of multimodal single-cell data. *Cell* **184**,
3573-3587 e3529, doi:10.1016/j.cell.2021.04.048 (2021).

22 Stuart, T. *et al.* Comprehensive Integration of Single-Cell Data. *Cell* **177**,
1888-1902 e1821, doi:10.1016/j.cell.2019.05.031 (2019).

23 Ma, T. *et al. Protracted yet coordinated differentiation of long-lived SARS-*
CoV-2-specific CD8+ T cells during COVID-19 convalescence (Cold Spring
Harbor Laboratory, 2021).

24 Cao, J. Joint profiling of chromatin accessibility and gene expression in
thousands of single cells. *Science (New York, N.Y.)* (2019).

25 Argelaguet, R. *et al.* Multi‐Omics Factor Analysis—a framework for

680		unsupervised integration of multi‑omics data sets. *Molecular Systems Biology*
681		**14**, e8124, doi:10.15252/msb.20178124 (2018).
682	26	Welch, J. D. *et al.* Single-Cell Multi-omic Integration Compares and Contrasts
683		Features of Brain Cell Identity. *Cell* **177**, 1873-1887.e1817,
684		doi:10.1016/j.cell.2019.05.006 (2019).
685	27	Ghosn, E., Yoshimoto, M., Nakauchi, H., Weissman, I. L. & Herzenberg, L. A.
686		Hematopoietic stem cell-independent hematopoiesis and the origins of innate-
687		like B lymphocytes. *Development* **146**, dev170571, doi:10.1242/dev.170571
688		(2019).
689	28	Orlova, D. Y. *et al.* QFMatch: multidimensional flow and mass cytometry
690		samples alignment. *Scientific Reports* **8**, doi:10.1038/s41598-018-21444-4
691		(2018).
692	29	Mulè, M. P., Martins, A. J. & Tsang, J. S. Normalizing and denoising protein
693		expression data from droplet-based single cell profiling. *bioRxiv,*
694		doi:10.1101/2020.02.24.963603 (2020).
695	30	Garimalla, S. *et al.* Differential transcriptome and development of human
696		peripheral plasma cell subsets. *JCI Insight* **4**, doi:10.1172/jci.insight.126732
697		(2019).
698	31	Glass, D. R. *et al.* An Integrated Multi-omic Single-Cell Atlas of Human B Cell
699		Identity. *Immunity* **53**, 217-232 e215, doi:10.1016/j.immuni.2020.06.013
700		(2020).
701	32	Kaminski, D. A., Wei, C., Qian, Y., Rosenberg, A. F. & Sanz, I. Advances in
702		human B cell phenotypic profiling. *Front Immunol* **3**, 302,
703		doi:10.3389/fimmu.2012.00302 (2012).
704	33	Tiller, T. *et al.* Autoreactivity in Human IgG+ Memory B Cells. *Immunity* **26**,
705		205-213, doi:10.1016/j.immuni.2007.01.009 (2007).
706	34	Mehtonen, J. *et al.* Single cell characterization of B-lymphoid differentiation
707		and leukemic cell states during chemotherapy in ETV6-RUNX1-positive
708		pediatric leukemia identifies drug-targetable transcription factor activities.
709		*Genome Medicine* **12**, doi:10.1186/s13073-020-00799-2 (2020).
710	35	Zhou, Y. *et al.* Transitional B cells involved in autoimmunity and their impact
711		on neuroimmunological diseases. *Journal of translational medicine* **18**,
712		doi:10.1186/s12967-020-02289-w (2020).

36   Becker, S. C. *et al.* A comparative analysis of human bone marrow–resident
and peripheral memory B cells. *Journal of Allergy and Clinical Immunology*
141, 1911-1913.e1917, doi:10.1016/j.jaci.2017.12.983 (2018).

37   Halliley, J. L. *et al.* Long-Lived Plasma Cells Are Contained within the
CD19(-)CD38(hi)CD138(+) Subset in Human Bone Marrow. *Immunity* **43**,
132-145, doi:10.1016/j.immuni.2015.06.016 (2015).

38   Jassal, B. *et al.* The reactome pathway knowledgebase. *Nucleic Acids Res*
48, D498-d503, doi:10.1093/nar/gkz1031 (2020).

39   Ocasio, J. *et al.* scRNA-seq in medulloblastoma shows cellular heterogeneity
and lineage expansion support resistance to SHH inhibitor therapy. *Nature*
Communications **10**, doi:10.1038/s41467-019-13657-6 (2019).

40   McGinnis, C. S., Murrow, L. M. & Gartner, Z. J. DoubletFinder: Doublet
Detection in Single-Cell RNA Sequencing Data Using Artificial Nearest
Neighbors. *Cell Syst* **8**, 329-337 e324, doi:10.1016/j.cels.2019.03.003 (2019).

41   Altman, N & Krzywinski, M. The curse(s) of dimensionality. *Nat Methods* **15**,
397, doi:10.1038/s41592-018-0013-3 (2018).

42   Orlova, D. Y., Herzenberg, L. A. & Walther, G. Science not art: statistically
sound methods for identifying subsets in multi-dimensional flow and mass
cytometry data sets. *Nature Reviews Immunology* **18**, 77-77,
doi:10.1038/nri.2017.150 (2018).

43   Babcock, B. *et al.* Data Matrix Normalization and Merging Strategies Minimize
Batch-specific Systemic Variation in scRNA-Seq Data. *bioRxiv* (2021).

44   Anjum, A. *et al.* Identification of Differentially Expressed Genes in RNA-seq
Data of Arabidopsis thaliana: A Compound Distribution Approach. *Journal of*
Computational Biology **23**, 239-247, doi:10.1089/cmb.2015.0205 (2016).

45   Uldrich, A. P. *et al.* CD1d-lipid antigen recognition by the gammadelta TCR.
Nat Immunol **14**, 1137-1145, doi:10.1038/ni.2713 (2013).

46   Shlomchik, M. J. & Weisel, F. Germinal center selection and the development
of memory B and plasma cells. *Immunological Reviews* **247**, 52-63,
doi:10.1111/j.1600-065x.2012.01124.x (2012).

47   Poli, A. *et al.* CD56bright natural killer (NK) cells: an important NK cell subset.
Immunology **126**, 458-465, doi:10.1111/j.1365-2567.2008.03027.x (2009).

48   Evans, J. H. *et al.* A distinct subset of human NK cells expressing HLA-DR
expand in response to IL-2 and can aid immune responses to BCG. *Eur J*

Immunol **41**, 1924-1933, doi:10.1002/eji.201041180 (2011).

49   He, Y. *et al.* EBV promotes human CD8 NKT cell development. *PLoS*

pathogens **6**, e1000915, doi:10.1371/journal.ppat.1000915 (2010).

50   Lawand, M., Déchanet-Merville, J. & Dieu-Nosjean, M. C. Key Features of

Gamma-Delta T-Cell Subsets in Human Diseases and Their

Immunotherapeutic Implications. *Front Immunol* **8**, 761,

doi:10.3389/fimmu.2017.00761 (2017).

51   Cai, Y. *et al.* Single-cell transcriptomics of blood reveals a natural killer cell

subset depletion in tuberculosis. *EBioMedicine* **53**, 102686,

doi:10.1016/j.ebiom.2020.102686 (2020).

52   Wong, E. B. *et al.* TRAV1-2(+) CD8(+) T-cells including oligoconal expansions

of MAIT cells are enriched in the airways in human tuberculosis. *Commun Biol*

2, 203, doi:10.1038/s42003-019-0442-2 (2019).

53   Villani, A. C. *et al.* Single-cell RNA-seq reveals new types of human blood

dendritic cells, monocytes, and progenitors. *Science (New York, N.Y.)* **356**,

doi:10.1126/science.aah4573 (2017).

54   Blaser, C. *et al.* Beta-galactoside-binding protein secreted by activated T cells

inhibits antigen-induced proliferation of T cells. *Eur J Immunol* **28**, 2311-2319,

doi:10.1002/(sici)1521-4141(199808)28:08<2311::Aid-immu2311>3.0.Co;2-g

(1998).

55   Hashimoto, K. *et al.* Single-cell transcriptomics reveals expansion of cytotoxic

CD4 T cells in supercentenarians. *Proceedings of the National Academy of*

Sciences of the United States of America **116**, 24242-24251,

doi:10.1073/pnas.1907883116 (2019).

56   Juno, J. A. *et al.* Cytotoxic CD4 T Cells-Friend or Foe during Viral Infection?

Front Immunol **8**, 19, doi:10.3389/fimmu.2017.00019 (2017).

57   Kumar, B. V., Connors, T. J. & Farber, D. L. Human T Cell Development,

Localization, and Function throughout Life. *Immunity* **48**, 202-213,

doi:10.1016/j.immuni.2018.01.007 (2018).

58   Zhu, J., Yamane, H. & Paul, W. E. Differentiation of effector CD4 T cell

populations (*). *Annu Rev Immunol* **28**, 445-489, doi:10.1146/annurev-

immunol-030409-101212 (2010).

59   Braun, J., Frentsch, M. & Thiel, A. Hobit and human effector T-cell

differentiation: The beginning of a long journey. *Eur J Immunol* **45**, 2762-2765,

doi:10.1002/eji.201545959 (2015).

60   Martin, M. D. & Badovinac, V. P. Defining Memory CD8 T Cell. *Front Immunol*
9, 2692, doi:10.3389/fimmu.2018.02692 (2018).

61   Jin, X. *et al.* Characterization of dendritic cell subtypes in human cord blood
by single-cell sequencing. *Biophysics Reports* **5**, 199-208,
doi:10.1007/s41048-019-00096-5 (2019).

62   Kuramasu, A., Saito, H., Suzuki, S., Watanabe, T. & Ohtsu, H. Mast cell-
/basophil-specific transcriptional regulation of human L-histidine
decarboxylase gene by CpG methylation in the promoter region. *J Biol Chem*
273, 31607-31614, doi:10.1074/jbc.273.47.31607 (1998).

63   Källberg, E. & Leanderson, T. A subset of dendritic cells express joining chain
(J-chain) protein. *Immunology* **123**, 590-599, doi:10.1111/j.1365-
2567.2007.02733.x (2008).

64   Dzierzak, E. & Philipsen, S. Erythropoiesis: development and differentiation.
Cold Spring Harb Perspect Med **3**, a011601,
doi:10.1101/cshperspect.a011601 (2013).

65   Pellin, D. *et al.* A comprehensive single cell transcriptional landscape of
human hematopoietic progenitors. *Nat Commun* **10**, 2395,
doi:10.1038/s41467-019-10291-0 (2019).

66   Xie, X. *et al.* Single-cell transcriptomic landscape of human blood cells.
National Science Review **8**, doi:10.1093/nsr/nwaa180 (2020).

67   Lai, S. *et al.* Mapping Human Hematopoietic Hierarchy at Single Cell
Resolution by Microwell-seq. *bioRxiv*, 127217, doi:10.1101/127217 (2017).

68   Yang, J., Zhang, L., Yu, C., Yang, X. F. & Wang, H. Monocyte and
macrophage differentiation: circulation inflammatory monocyte as biomarker
for inflammatory diseases. *Biomark Res* **2**, 1, doi:10.1186/2050-7771-2-1
(2014).

69   Kawamura, S. *et al.* Identification of a Human Clonogenic Progenitor with
Strict Monocyte Differentiation Potential: A Counterpart of Mouse cMoPs.
Immunity **46**, 835-848.e834, doi:10.1016/j.immuni.2017.04.019 (2017).

70   Evrard, M. *et al.* Developmental Analysis of Bone Marrow Neutrophils Reveals
Populations Specialized in Expansion, Trafficking, and Effector Functions.
Immunity **48**, 364-379.e368, doi:10.1016/j.immuni.2018.02.002 (2018).

71   Cao, J. *et al.* Comprehensive single-cell transcriptional profiling of a

multicellular organism. *Science (New York, N.Y.)* **357**, 661-667,

doi:10.1126/science.aam8940 (2017).

72   Hafemeister, C. & Satija, R. Normalization and variance stabilization of single-

cell RNA-seq data using regularized negative binomial regression. *Genome*

Biology **20**, 296, doi:10.1186/s13059-019-1874-1 (2019).

73   Traag, V. A., Waltman, L. & Van Eck, N. J. From Louvain to Leiden:

guaranteeing well-connected communities. *Scientific Reports* **9**,

doi:10.1038/s41598-019-41695-z (2019).

74   Chromium Single Cell 5' Reagent Kits User Guide (v2 Chemistry Dual Index).

https://support.10xgenomics.com/permalink/user-guide-chromium-single-cell-

5-reagent-kits-user-guide-v2-chemistry-dual-index.

Main lineage	SuPERR-seq cluster	Cell classification	Phenotype	Refs
Plasma cells	1	Plasma cells	CD19+/CD20-/highest levels of Ig-specific transcripts	46
B cells (CD19+, CD20+)	2	Naïve-I	IgD+/IgM+/CD27-/CD95-	30-33
	3	Naïve-II	IgD+/IgM+/CD27-/CD95-/HLA-DQA2+	
	4	Switched Memory	IgD-/IgM-/CD27+/CD95+	
	5	Unswitched Memory	IgD+/IgM+/CD27+	
	6	IgM^{hi}	IgM^{hi}/CD27-/CD95+/CD24-	
NK/NKT/MAIT/gdT (CD56+)	7	NK-I CD16+	KLRC3+/CD11b^{hi}	47-52
	8	NK-II CD16^{hi}	CX3CR1+	
	9	NK-III CD16+	CX3CR1+	
	10	NK CD56^{hi}	CD16-	
	11	MAIT cells	TRAV1-2+	
	12	NKT	CD4+/CCR7+	
	13	iNKT	TRAV24+/CD8+	
	14	NK-IV CD16+	CD11b^{hi}	
	15	γδ T cells	TRGV9+/TRDV2+	
	16	NK-V CD16+ (dividing)	MKI67+	
Monocytes (CD14$^{lo/+}$)	17	Classical-I	CD16-/CD68+/HLA-Drhi	53
	18	Classical-II	CD16-/CD68+/HLA-DR^{lo}	
	19	Classical-III	CD16-/CD68+/HLA-DR^{lo}/CD11b^{hi}	
	20	Non-classical	CD16+/CD14$^{-/lo}$/CD68+/HLA-DR^{hi}	
	21	Intermediate/DCs	CD16^{int}/CD14^{int}/CD68+/HLA-DR^{hi}/CD11c+	
CD4 T cells (CD3+, CD4+)	22	Naïve-I	CCR7+/SELL+/CD27+/CD95-/CD3^{hi}	54-58
	23	Naïve-II	CCR7+/SELL+/CD27+/CD95-	
	24	Tcm→Tem	CCR7^{lo}/SELL+/CD27+/CD95+	
	25	CTL-I	KLRB1+	
	26	CTL-II	KLRB1+/GZMA+/GZMK+	
	27	Tem	CCR7^{lo}/SELL-/CD27-/CD95+/LGALS1+/S100A4^{hi}	
	28	Treg	FOXP3+/CTLA4+/CD95+/HLA-DRB1+	
	29	Treg Naïve	FOXP3+/CD45RA+/CD95-	
	30	Temra	CD45RA+/NKG7+/GNLY+/GZMB+	
CD8 T cells (CD3+, CD8a+)	31	Naïve-I	CCR7+/CD45RA+/CD27+/CD127+	59,60
	32	Tcm-I	CCR7+/SELL+/CD27+/CX3CR1-	
	33	Tem	CCR7-/SELL+/CX3CR1+	
	34	TLE	CCR7-/SELL+/CX3CR1+/ZNF683+	
	35	MAIT cells	TRAV1-2+	
	36	Naïve-II	CCR7+/CD45RA+/CD27^{hi}/CD127+	
	37	Temra	CD45RA+/CCR7-/CD27-/CD127-	
	38	Tcm-II	CCR7+/SELL+/CD27+/CX3CR1-/TNFSF10+	

Table 1. Cell classification and phenotype of PBMC clusters generated by the SuPERR-seq pipeline. Gene expression (GEX) and cell-surface (ADT) markers used to classify the lymphoid and myeloid cell populations in the human peripheral blood. NK: Nature Killer cells; NKT: NK T cells; iNKT: invariant NKT; γδ T: gamma-delta T cells. Tcm: central memory; Tem: effector memory; CTL: cytotoxic T lymphocytes; Treg: regulatory T cells; Temra: effector memory expressing CD45RA; TLE: long-lived effector memory; MAIT: mucosal associated invariant T cells.

Main lineage	SuPERR-seq cluster	Cell classification	Phenotype	Refs
Plasma cells (CD138-)	1	Ig^{hi}/PRDM1-	Ig^{hi}/PRDM1-	30, 37
	2	HLA-DR^{hi}/SDC1^{lo}/G2-M phase	HLA-DR^{hi}/SDC1^{lo}/G2-M phase	
Plasma cells (CD138+)	3	IRF4^{hi}	IRF4^{hi}	
	4	XBP1^{hi}/SDC1^{hi}	XBP1^{hi}/SDC1^{hi}	
B cells (CD19+, CD20+)	5	T3/Naïve	CD21+/IL4R+	34-36
	6	Small pre-B/pre-BII-I	RAG+/IGKC+	
	7	Small pre-B/pre-BII-II	RAG+/IGLC3+	
	8	Switched Memory	CD21+/CD27+/IGHA1+/IGHG1+	
	9	T1/T2	RAG-/CD10+/CD20$^{-/lo}$	
	10	Large pre-B/pre-BII	CD34-/DNTT-/MKI67+	
	11	Early pro-B/pre-pro-B	CD34+/DNTT^{hi}	
	12	Activated naïve	CD21+/NR4A1+/DUSP2+	
	13	Late pro-B/pre-BI	CD34+/DNTT^{int}	
HSPCs (Lineage-, CD34+/-)	14	Pre-reticulocytes	GYPA^{int}/HBB+	61-67
	15	Pre-pDCs	CD123+/CD304+/CD303+/CSF2RA+	
	16	GMP	CD34+/FLT3+/CD164+/CD45RA^{hi}/Cell cycle	
	17	MEP	CD34+/GYPA-/ITGA2B+	
	18	Reticulocytes	GYPA-/HBB+/HBM-	
	19	pro-Neutrophil	CD34+/MPO+/ELANE+	
	20	Reticulocytes GYPA^{hi}	GYPA^{hi}/HBB+	
	21	HSC/MPP	CD34+/AVP+/CD38-	
	22	Erythrocytes	GYPA-/HBB+/HBM-	
	23	Pro-erythoblast	GYPA^{lo}/HBB+/HBM+	
	24	CMP	CD34+/FLT3+/CD164+	
	25	Basophil/Mast cell Progenitors	CD34+/CLC+/HDC+	
Myeloid/Granuloid (CD11c+)	26	Neutropoiesis	MPO^{hi}/ELANE^{hi}	68-70
	27	Monopoiesis	CD68^{hi}/CD14^{hi}	
	28	Monopoiesis	CD68^{lo}/CD14^{int}	
	29	Neutropoiesis	MPO^{int}/ELANE^{int}	
	30	Monopoiesis	CD68^{lo}/CD14^{lo}	
	31	Monopoiesis	CD68^{int}/CD14^{int}	
	32	Neutropoiesis	MPO^{lo}/ELANE^{lo}/MKI67^{hi}	
	33	Monopoiesis	CD68^{hi}/CD14^{hi}	
	34	Progenitors	FLT3+/CD74+	

Table 2. Cell classification and phenotype of BM clusters generated by the SuPERR-seq pipeline. Gene expression (GEX) and cell-surface (ADT) markers used to classify the plasma cells, B cells, myeloid and granuloid cells, and the hematopoietic stem and progenitor cells (HSPCs) in the human bone marrow. Ig: immunoglobulin production; G2-M phase: genes involved in cell cycle; T1/2/3: Transitional B cells; pDCs: plasmacytoid dendritic cells; GMP: granulocyte-monocyte progenitor; MEP: megakaryocyte-erythroid progenitor; HSC: hematopoietic stem cell; MPP: multi-potent progenitor; CMP: common myeloid progenitor.

Figure 1. SuPERR-seq workflow. (A) Schematic overview of the experimental design. Blood and bone marrow aspirates were encapsulated with barcoded microspheres. We generated three libraries for each sample corresponding to gene expression (GEX), cell-surface protein/antibody-derived tags (ADT), and cell-receptor repertoire (VDJ). Libraries were sequenced to a target depth, and count matrices were assembled for each -omic data separately. (B) SuPERR-seq workflow is composed of two main steps. At the first step, major cell lineages are manually gated by integrating information from both the ADT and V(D)J data matrices. Then, the manually-gated cell lineages are further sub-clustered based on information from the GEX data. The V(D)J matrix can be used to further identify the diversity of heavy (VH) and light (VL) variable genes among the plasma cell clusters. PCs: plasma cells.

Figure 2. SuPERR-seq workflow applied to PBMCs. (A) "Gating strategy" approach to identify major cell lineages on biaxial plots based on surface markers (ADT) and V(D)J data. Gates for major lineages are indicated as black outlines and black text. Gates for downstream cell identity validation are indicated as golden outlines and golden text. (B) Cross comparison between the manually-gated major lineages and the final SuPERR-seq clusters. (C) The average expression levels of surface markers (ADT) and Ig-specific transcripts (V(D)J) for the final SuPERR-seq clusters. (D) UMAP representation of the B-cell clusters identified by the SuPERR-seq workflow. PCs: plasma cells. usM B: unswitched memory B cells. sM B: switched memory B cells.

Figure 3. SuPERR-seq workflow applied to BM cells. (A) "Gating strategy" approach to identify major cell lineages on biaxial plots based on surface markers (ADT) and V(D)J data. Gates for major lineages are indicated as black outlines and black text. Gates for downstream cell identity validation are indicated as golden outlines and golden text. (B) Cross comparison between the manually-gated major lineages and the final SuPERR-seq clusters. (C) The average expression levels of surface markers (ADT) and Ig-specific transcripts (V(D)J) for the final SuPERR-seq clusters. PCs: plasma cells.

Figure 4. Cell-type specific variations in gene expression count matrices. (A) "Gene count" represents the number of unique genes expressed by each cell type. (B) "UMI count" represents the total mRNA abundance expressed by each cell type. (C) "Percent of Ribosomal" represents the percentage of ribosomal gene-derived UMI counts expressed by each cell type. Grey line shows the mean expression level for each feature in the total PBMC and BM samples. (D) Left panel: top 30 highly-variable genes from PBMC-derived B cells (green dots) displayed together with the top 300 highly-variable genes from total PBMC (grey dots). Right panel: top 30 highly-variable genes from total PBMC (red dots) displayed together with the top 300 highly-variable genes from total PBMC (grey dots). The dots under the red dashed line fall below the top 300 HVGs of the total PBMC. Student's t-test was used to compare the mean of each cell type with the mean of the total PBMC/BM. *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001, unpaired, two-tailed. Multiple-group ANOVA test for (A), (B), and (C): p<2.2e-16. PCs: plasma cells. HVGs: highly-variable genes.

Figure 5. B-cell developmental pathway in human adult BM identified by the SuPERR-seq workflow.
(A) Cell-type classification of BM B cells based on three -omics, surface protein (ADT), V(D)J matrix, and gene expression profile (GEX). (B) The expression level of surface proteins and genes used to define the B-cell developmental stages. (C) Violin plots illustrate the differences in expression levels (ADT and GEX) between clusters. Grey line shows the mean expression level across all clusters.

Figure 6. SuPERR-seq workflow identifies four subsets of human plasma cells in the BM. (A) UMAP representation of the four BM plasma cells. (B) Top panel: percentage of Ig-specific transcripts (UMI) expressed in each plasma cell subset. Bottom panel: expression levels (sum) of plasma cell genes (IRF4, PRDM1, XBP1, SDC1) after removing Ig-specific UMIs and re-normalizing the data matrix. (C) Expression levels of individual plasma cell genes after removing Ig-specific UMIs and re-normalizing the data matrix. Grey line shows the mean expression level across all clusters. (D) The antibody isotypes and subclasses expressed by each plasma cell subset. (E) Circus plot showing shared clones between clusters (clonal lineage was identified by the identical V and J gene usage, identical CDR3 nucleotide length, and ≥85% homology within the CDR3 nucleotide sequence). (F) The Reactome Pathway Database analysis show the unique biological processes that define each plasma cell subset.

Figure 7. Cell-doublet identification by SuPERR-seq and other conventional approaches. (A) Distribution (unique gene count x total UMI) of cell doublets (left) and singlets (right) detected by the SuPERR-seq approach. The red dashed lines show the threshold used by the conventional approaches to exclude cells that express higher than mean+4SD of gene count and total UMI. Only the cells above the dashed lines would have been excluded from the downstream analysis in conventional approaches (i.e., plasma cells in PBMCs, highlighted in red circle, would have been excluded from downstream analysis). (B) Number of unique genes (left panels) and number of total UMIs (right panels) expressed by singlets and doublets in PBMC (top panels) and BM (bottom panels). Grey line shows the mean expression level across all clusters. (C) Cell doublets identified by SuPERR-seq and projected on a UMAP show these cells are spread across multiple clusters. (D) Venn diagram comparing the cell doublets identified by the SuPERR-seq and DoubletFinder. Many cell doublets confirmed by SuPERR-seq (blue) were missed by DoubletFinder.

Figure 8. Cell-type misclassification by conventional workflow. (A) and (B) *Left panels*: PBMC (A) and BM (B) clustering result by conventional workflow (i.e., based solely on gene expression matrix). *Middle panels*: PBMC (A) and BM (B) major cell lineages identified by SuPERR-seq and projected onto the UMAP plot of the conventional workflow. *Right panels*: red dots show the PBMC (A) and BM (B) cells that were misassigned/misclassified by the conventional Seurat workflow. The Cell Misclassification Statistic (CMS) score calculate the percentage of cells that were misclassified (9.84% in PBMC and 5.94% in BM) by the conventional workflow. (C) The Seurat cluster 5 contains misassigned/misclassified cells (i.e., mixture of NK, NKT, and T cells) and was further explored using the cell-surface markers CD56 and CD3 (left panel). The differentially-expressed genes (DEGs) for cluster 5 (pink circle) show *TRDV1* as a top hit (Venn diagram). TRDV1 is expressed on CD3+ gamma-delta T cells only (right panel) and absent in NK cells, showing that the contaminated cells in cluster 5 have contributed to the top DEGs.

Antigen	Clone	Vendor	Cat. no.
CCR7 (CD197)	G043H7	Biolegend	353251
CD11b	ICRF44	Biolegend	301359
CD127 (IL-7Ra)	A019D5	Biolegend	351356
CD14 (LPS receptor)	M5E2	Biolegend	301859
CD16 (FcγRIII)	3G8	Biolegend	302065
CD19 (B4)	HIB19	Biolegend	302265
CD20 (B1)	2H7	Biolegend	302363
CD21 (C3dR)	Bu32	Biolegend	354923
CD25	BC96	Biolegend	302649
CD27 (S152)	O323	Biolegend	302853
CD3 (T3, CD3ε)	UCHT1	Biolegend	300479
CD38 (T10)	HIT2	Biolegend	303543
CD4	RPA-T4	Biolegend	300567
CD45	2D1	Biolegend	368545
CD45RA	HI100	Biolegend	304163
CD56	QA17A16	Biolegend	392425
CD69	FN50	Biolegend	310951
CD80	2D10	Biolegend	305243
CD86	IT2.2	Biolegend	305447
CD8a	RPA-T8	Biolegend	301071
CD95 (APO-1)	DX2	Biolegend	305651
HLA-DR	L243	Biolegend	307663
PDL1 (CD274)	29E.2A3	Biolegend	329751
Custom oligo-conjgated in house			
Antigen	**Clone**	**Vendor**	**Cat. no.**
CD11c (αX integrin)	S-HCL-3	Biolegend	371502
CD133 (Prominin-1)	clone 7	Biolegend	372802
CD138 (Syndecan-1)	MI15	Biolegend	356502
CD34 (gp105-120)	581	Biolegend	343602
CD41 (gpIIb)	HIP8	Biolegend	303702
CD43 (Ly-48)	CD43-10G7	Biolegend	343202
CD45R (B220)	RA3-6B2	Biolegend	103202
CD5 (Ly-1)	UCHT2	Biolegend	300602
CD20 (B1)	2H7	Biolegend	302302

Table S1. List of antibody-derived tags (ADT) used in the multi-omics single-cell assays. Cell-surface protein antigens measured in both human peripheral blood and bone marrow samples.

	Sample ID	Mean Reads per Cell	Cell count
Gene expression	PBMC1	138,274	4,034
	PBMC2	150,868	2,302
	PBMC3	45,805	6,423
	BM1	149,030	3,480
	BM2	149,622	3,946
Surface protein expression	PBMC1	1,399	4,034
	PBMC2	2,273	2,302
	PBMC3	928	6,423
	BM1	2,762	3,480
	BM2	1,703	3,946
V(D)J expression	PBMC1	126,518	273
	PBMC2	601,049	64
	PBMC3	56,584	582
	BM1	22,991	1,669
	BM2	17,418	2,553

Table S2. Sequencing depth. Mean reads per cell and total number of cells analyzed in each sample. PBMC: peripheral blood mononuclear cells; BM: bone marrow cells.

Figure S1. Biaxial feature plots to identify and remove cell doublets from PBMCs. Supplement for Figure 2. Cell doublets were gated using red boxes. The visualization was generated by SeqGeq™ (FlowJo, LLC).

Figure S2. Biaxial feature plots to identify and remove cell doublets from BM cells. Supplement for Figure 3. Cell doublets were gated using red boxes. The visualization was generated by SeqGeq™ (FlowJo, LLC).

Figure S3. Gating strategy to identify major lineages by high-dimensional flow cytometry. An aliquot of each PBMC (top row) and BM (bottom row) samples was stained with a panel of fluorescently-labeled antibodies in parallel to the SuPERR-seq workflow. The gating strategy (red gates) shows the percentage of live, single B cells and plasma cells.

Figure S4. Blood monocytes identified by the SuPERR-seq workflow. UMAP and violin plots show surface markers and gene expression levels used to identify and classify five subsets of blood (PBMC) monocytes. Grey line shows the mean expression level across all clusters.

Figure S5. Blood CD4 T-cell subsets identified by the SuPERR-seq workflow. UMAP and violin plots show surface markers and gene expression levels used to identify and classify nine subsets of blood (PBMC) CD4 T cells. Grey line shows the mean expression level across all clusters. Tcm: central memory; Tem: effector memory; CTL: cytotoxic T lymphocyte; Treg: regulatory T cells; Temra: effector memory expressing CD45RA.

Figure S6. Blood CD8 T-cell subsets identified by the SuPERR-seq workflow. UMAP and violin plots show surface markers and gene expression levels used to identify and classify eight subsets of blood (PBMC) CD8 T cells. Grey line shows the mean expression level across all clusters. Tcm: central memory; Tem: effector memory; TLE: long-lived effector memory; MAIT: mucosal associated invariant T cells; Temra: effector memory expressing CD45RA.

Figure S7. Blood NK, NKT, and gd T cell subsets identified by the SuPERR-seq workflow. UMAP and violin plots show surface markers and gene expression levels used to identify and classify ten subsets of blood (PBMC) NK, NKT, and gd T cells. Grey line shows the mean expression level across all clusters. MAIT: mucosal associated invariant T cells; NKT: natural killer T cells; iNKT: invariant NKT; gd T: gamma-delta T cells.

Figure S8. Monocyte and neutrophil development in the BM identified by the SuPERR-seq workflow. UMAP and violin plots show surface markers and gene expression levels used to identify and classify nine subsets of BM monopoiesis and granulopoiesis. Grey line shows the mean expression level across all clusters. The red arrow shows potential direction of developmental pathway for neutrophils. The cyan arrow shows potential direction of developmental pathway for monocytes.

Figure S9. BM Hematopoietic Stem and Progenitor Cells (HSPCs) identified by the SuPERR-seq workflow. UMAP and violin plots show surface markers and gene expression levels used to identify and classify twelve subsets of BM HSPCs. Grey line shows the mean expression level across all clusters. pDCs: plasmacytoid dendritic cells; GMP: granulocyte-monocyte progenitor; MEP: megakaryocyte-erythroid progenitor; HSC: hematopoietic stem cell; MPP: multi-potent progenitor; CMP: common myeloid progenitor.

Figure S10. PBMC cell-type misassignment by conventional workflow. (A) Cross comparison of PBMC clustering results between conventional (Seurat) and SuPERR-seq workflows. For the sake of comparison, each workflow analyzed the same set of cell barcodes. The plasma cells identified by the SuPERR-seq "manual gating" using the Ig-specific transcripts were mostly removed from the conventional workflow (see Fig. 7A), but were included here as SuPERR-seq cluster 1. (B) UMAP and clustering results from the conventional workflow (Seurat). SuPERR-seq-defined major lineages were projected onto UMAP to reveal the misassignment/misclassification of cells by the conventional workflow. For example, the cluster 9 from the conventional workflow (Seurat) represent a mixture of NK cells (blue), monocytes (red), CD4+ T cells (aqua) and CD8+ T cells (pink). Similarly, the conventional workflow misclassified NK cells, illustrated here by the spreading of the well-defined SuPERR-seq NK cells (blue) across multiple clusters in the conventional workflow.

Figure S11. BM cell-type misassignment by conventional workflow. (A) Cross comparison of BM clustering results between conventional (Seurat) and SuPERR-seq workflows. For the sake of comparison, each workflow analyzed the same set of cell barcodes. (B) UMAP and clustering results from the conventional workflow (Seurat). SuPERR-seq-defined major lineages were projected onto UMAP to reveal the misassignment/misclassification of cells by the conventional workflow. For example, the cluster 3 from the conventional workflow (Seurat) represent a mixture of plasma cell (green), B cells (red), and various HSPCs (blue), and myeloid cells (pink). Similarly, the conventional workflow misclassified B cells, illustrated here by the spreading of the well-defined SuPERR-seq B cells (red dots) across multiple clusters in the conventional workflow. (C) Cell-type composition in each "plasma cell" sub-cluster generated by iterative/recursive Seurat clustering.