N-VEGF, the autoregulatory arm of VEGF-A

Marina Katsman[1], Aviva Azriel[1], Guy Horev[2], Yitzhak Reizel[1†], and Ben-Zion Levi[1*]

[1]Faculty of Biotechnology and Food Engineering, Technion-Israel, Institute of Technology, Haifa, Israel

[2]Bioinformatics Knowledge Unit, The Lorry I. Lokey Interdisciplinary Center for Life Sciences and Engineering, Technion-Israel, Institute of Technology, Haifa, Israel

*Corresponding author, e-mail: blevi@technion.ac.il

†Co-corresponding author, e-mail: y.reizel@technion.ac.il

Abstract

Vascular endothelial growth factor A (VEGF-A) is a secreted protein that stimulates angiogenesis in response to hypoxia. Under hypoxic conditions, a non-canonical long isoform called L-VEGF is concomitantly expressed with VEGF-A. Once translated, L-VEGF it is proteolytically cleaved to generate N-VEGF and VEGF-A. Interestingly, while VEGF-A is secreted and affects the surrounding cells, N-VEGF is mobilized to the nucleus. This suggests that N-VEGF participates in transcriptional response to hypoxia. In this study, we performed a series of complementary experiments to examine the functional role of N-VEGF. Strikingly, we found that the mere expression of N-VEGF followed by its hypoxia-independent mobilization to the nucleus was sufficient to induce key genes associated with angiogenesis, such as Hif1α, VEGF-A isoforms, as well as genes associated with cell survival under hypoxia. Complementarily, when N-VEGF was genetically depleted, key hypoxia-induced genes were downregulated and cells were significantly susceptible to hypoxia-mediated apoptosis. This is the first reports of N-VEGF serving as an autoregulatory arm of VEGF-A. Further experiments will be needed to determine the role of N-VEGF in cancer and embryogenesis.

Introduction

VEGF-A (or VEGF) is a key regulator of angiogenesis in normal physiology as well as in pathological events such as cancer (for review see [1, 2]). When translated under hypoxic conditions, a longer isoform of VEGF (L-VEGF) is generated from a non-canonical start site, CUG. L-VEGF adds 180 aa to the VEGF coding sequence and has been detected in VEGF-producing cells, including various cancer cells [3-5]. Once translated, L-VEGF is subjected to a proteolytic cleavage upstream to the classical translation start site of VEGF, yielding N-VEGF and VEGF (see schematic illustration in Fig. 1A). While VEGF-A is known to dramatically affect angiogenesis in surrounding cells the functional role of N-VEGF is not known. Interestingly, ectopically expressed N-VEGF shuttles to cell nuclei only under hypoxic conditions [6]. Furthermore, immunohistochemical staining of normal angiogenic tissue samples, such as kidney and breast, demonstrated sporadic staining of N-VEGF in the nuclei of renal tubular epithelium and of epithelium cells coating the mammary ducts. Finally, renal cell carcinoma and breast cancer cell nuclei exhibited strong immunostaining of N-VEGF in all cells [6]. These observations imply the involvement of N-VEGF in transcriptional regulation of VEGF-secreting cells. However, its actual impact on gene expression and on the response to hypoxia remains a mystery.

We hypothesize that N-VEGF and VEGF play complementary regulatory roles. While VEGF is secreted and mainly affects neighboring cells, N-VEGF is retained in the cell and shuttles to the nucleus where it regulates the hypoxic transcriptional response. To characterize the role of N-VEGF in regulating transcription and in the adaptation to hypoxic stress, we performed a series of experiments. N-VEGF was mobilized to the nucleus, independent of hypoxia. Strikingly, its expression under normoxia was sufficient to upregulate key genes involved in the response to hypoxia and in the induction of angiogenesis. Complementarily, genetic depletion of N-VEGF

inhibited the upregulation of hypoxia-induced genes and increased the apoptosis rate upon hypoxia. All these are sound proof for the key role of N-VEGF in the response to hypoxia.

Results

Expression and hypoxia-independent-mobilization of N-VEGF

To examine if the mere presence N-VEGF in the nucleus is sufficient to induce a hypoxia-associated transcriptional response, we fused N-VEGF at its N-terminus to the nuclear localization sequence (NLS) of the Cas9 nuclease (NLS-N-VEGF). We chose to introduce N-VEGF to the nucleus under normoxic conditions to avoid the notable change that occurs in the transcriptome upon hypoxia [7]. For this purpose, N-VEGF and NLS-N-VEGF were cloned under the inducible Tet-on promoter in a lentiviral vector [8]. Subsequently, NIH3T3 cells expressing N-VEGF or NLS-N-VEGF, were grown in the presence or absence of the Tet-on promoter inducer, doxycycline (dox). NIH3T3 cells are mouse fibroblasts cells shown in many studies to activate hypoxia-associated genes, including VEGF-A and Hif1α, upon exposure to hypoxic stress [9, 10]. As seen in Fig. 1B, in the presence of dox, 25% of NLS-N-VEGF was identified in the nuclear fraction. The reason for the partial nuclear localization is probably because Cas9-NLS fusion to N-VEGF resulted in reduced translocation efficiency. It is important to note that under hypoxia, only 50% of the induced N-VEGF shuttled to the nucleus (Fig. 3A). Thus, only a 50% reduction in its nuclear mobilization was noted on addition of the NLS tag. Further, immunofluorescence analysis confirmed that N-VEGF exhibited nuclear localization only following hypoxia (Fig. 1C, second row), while NLS-N-VEGF showed nuclear expression even without hypoxia (Fig. 1C, third and fourth rows).

Hypoxia-independent nuclear mobilization of N-VEGF induces part of the hypoxic transcriptional program

To study the effect of nuclear NLS-N-VEGF on cellular transcription, RNA was extracted from cells expressing either N-VEGF or NLS-N-VEGF and subjected to RNA-Seq. In efforts to identify altered gene expression resulting from shuttling of N-VEGF to the nucleus, while avoiding the transcriptional dynamics affected by the mere expression of N-VEGF in the cytoplasm, we only monitored genes exhibiting differential expression in dox-induced NLS-N-VEGF as compared to N-VEGF expressing cells and their dox untreated counterparts. Cells were collected after a 16-h treatment with dox to allow the translation and mobilization of N-VEGF to the nucleus. The differentially expressed genes were divided to up- and downregulated groups (fold change ≥ 2, adjusted p-value<0.05) and each group was functionally annotated using Metascape [11]. 123 upregulated and 32 downregulated differentially expressed genes resulting from the shuttling of N-VEGF to the nuclei were identified (see genes list and annotations in Table S1).

The annotated groups of the upregulated genes (Fig. 2A) clustered into three main categories. The first and most relevant category (Fig. 2A, purple bars), represent genes promoting angiogenesis, including Hif1α, a major hypoxia-induced transcriptional regulator [12], and VEGF isoforms 120 and 164 (Fig. 2B). Additional signature genes mediating angiogenesis, such as Adgrg1, Itgav, Nrp, Flna and Pik3c2a, were upregulated by mobilization of NLS-N-VEGF to the nucleus (Fig. 2B, and Table S1) [13]. Another dominant annotation involved genes associated with anti-apoptotic pathways rescuing cells from hypoxic stress (Fig. 2A, brown bars). Indeed, signature genes of anti-apoptotic pathways such as Sh3rf1 [14], Lrrk2 [15] and Src [16], were induced by N-VEGF shuttling (Fig. 2B, and Table S1). The last and major category of annotated gene groups was associated with integrity, structure, and maintenance of cellular cytoskeleton,

which is highly relevant during hypoxic stress (Fig. 2A, red bars), including Cav2 [17], Lamc2 [18], Cfl2 [17], Rock2 [19], and more (Fig. 2B). Strikingly, many of the 123 upregulated genes are known targets of Hif1α, pointing to a key pathway activated by N-VEGF. Genes downregulated following NLS-N-VEGF mobilization to the nucleus included genes associated with regulation of translation, (Fig. S1). This aligns with the role of N-VEGF as an inducer of adaptive mechanism to hypoxic crisis by inhibiting the translational machinery [20].

In order to examine whether the expression pattern of the 155 differentially expressed genes overlapped with genes induced by hypoxia, we compared N-VEGF-induced genes to hypoxia-mediated gene expression (elaborated hereafter, Fig. 4A). Overall, 42% of the genes exhibited similar trends, with 51 genes induced and 14 genes downregulated in both hypoxia-induced cells and in cells showing hypoxia-independent mobilization of N-VEGF (Fig. 2B). The annotated groups of upregulated genes that exhibited similar expression, grouped into the three main categories described above (Fig. S2A). These observations were verified by qPCR on a set of key genes in the three groups (Fig 2C). Exposure of NLS-N-VEGF-expressing cells to both dox and hypoxia resulted in a synergistic effect, with increased expression levels of these key genes (Fig. 2C, dark blue columns). Last, the critical role of nuclear mobilization of N-VEGF in regulating the expression of these genes was demonstrated by the lack of an effect of dox on N-VEGF-infected cells. Taken together, this focused gene profile expression analysis highlighted the role of nuclear N-VEGF in initiating part of the main hypoxic signaling cascade.

Genetic deletion of N-VEGF using CRISPR-Cas9

To show the significance of endogenous N-VEGF expression on cellular physiology, we took advantage of the CRISPR-Cas9 genome editing system to delete only the coding region of N-VEGF (178 aa in mice) without affecting the translation start site of the "classical" VEGF. To

measure VEGF and N-VEGF protein expression levels, we isolated soluble proteins from cytoplasmic and nuclear extracts of ΔN-VEGF cells and control NIH3T3 cells and subjected them to Western blot and immunofluorescence analyses. As clearly demonstrated in Fig. 3A and C, N-VEGF was entirely absent following CRISPR-Cas9 deletion. Hence, hypoxia-induced expression and nuclear mobilization of N-VEGF occurred only in wild type cells. Importantly, N-VEGF deletion did not affect VEGF levels (Fig. 3A).

Impact of N-VEGF deletion on hypoxia-induced genes

To show the effect of N-VEGF deletion on gene expression levels, RNA-seq analysis was performed on deleted cells and wild type cells before and following hypoxia (see Materials and Methods). A gene expression heatmap focusing on the 3,000 genes modified following hypoxia in wild type versus ΔN-VEGF cells before and after hypoxia is presented in Fig. 4A. The genes are grouped according to expression pattern, similarity or dissimilarity, between the two cell types before or following hypoxia (G1-G6, indicated Fig. 4A, and annotations in Table S2). Most genes (66%) in wild type and ΔN-VEGF cells exhibited similar expression patterns following hypoxia, as indicated in groups 1 and 6. Thus, deletion of N-VEGF led to a partial yet significant cellular response to hypoxia. Interestingly, 34% of the genes induced or repressed by hypoxia showed differential gene expression in ΔN-VEGF compared with control NIH3T3 cells (groups G2-G5). Of note, most of these genes showed differential expression in ΔN-VEGF cells even before hypoxia was induced. Following hypoxia, most of these genes showed only partial elevation or downregulation and did not reach the levels of expression observed in normal cells (groups G2 and G5). A small proportion of the genes displayed opposite expression trends under normoxia as compared to hypoxia in the two cell lines (groups G3 and G4). For example, group G3 showed higher levels of expression in the knockout compared to the wild type under normal conditions,

while following hypoxia, expression levels were downregulated in the knockout and elevated in the wild type. This observation shows that N-VEGF deletion had an immense impact on gene expression even before and more massively after hypoxic stress (see annotations in Table S2). Since the main focus of this study was to characterize the nuclear role of N-VEGF, we analyzed the impact of its knockout on the 155 genes modified following N-VEGF translocation to the nucleus (Fig. 4B, NLS-N-VEGF vs. ΔNVEGF). Interestingly, out of the 155 genes modified in NLS-N-VEGF cells, 53% exhibited opposite expression patterns in hypoxia-challenged ΔN-VEGF cells, which is much higher to what is expected by random. Annotations of the genes that were upregulated by nuclear mobilization of N-VEGF and downregulated in N-VEGF-depleted cells included angiogenesis and cell survival pathways (Fig. 4B), including genes associated with negative regulation of apoptotic signaling pathway, such as Aqp1, Msn, Sgk1, Src, Nrp1, and Pik3c2a [16, 21-25]. This suggests that N-VEGF expression protects cells against death elicited by stress signals such as hypoxia.

N-VEGF deletion leads to increased hypoxia-mediated cell death

To examine if N-VEGF plays a role in defending cells against apoptosis-induced hypoxic stress, the effects of three different yet similar stress signals on ΔN-VEGF vs. wild type cells were compared. More specifically, cells were exposed to hypoxia, $CoCl_2$ [26], and serum deprivation [27], three stressors that mimic lack of blood supply. As a control, we induced Endoplasmic Reticulum stress (ER) [28], which does not mimic lack of blood supply and does not lead to VEGF expression. All stressors resembling lack of blood supply led to statistically significant more apoptosis in ΔN-VEGF cells in comparison to NIH3T3 cells (Fig. 4C, compare second and fourth columns in Hypoxia, $CoCl_2$ and Starvation panels). In contrast, ER stress did not elicit a significant change in the apoptotic index between N-VEGF-deleted cells as compared to their WT counterpart

cells (Fig. 4C, Tunicamycin panel). Taken together, N-VEGF is essential for mediation of the anti-apoptotic state in stressed cells to overcome the harsh conditions until oxygen or blood supplies are improved. Overall, N-VEGF has a significant nuclear role on hypoxia-induced genes and physiology, as per the following proposed model (Fig.5). Upon hypoxia, the non-canonical isoform L-VEGF is translated and cleaved to VEGF and N-VEGF. Then, N-VEGF is mobilized to the nucleus and participates in the transcriptional regulation of the hypoxic response and in induction of anti-apoptotic genes. While VEGF is secreted and regulates the hypoxic response in neighboring cells, N-VEGF serves as the autoregulatory arm of VEGF.

Discussion

Hypoxia is a severe cellular stress that leads to cellular injury and cell death. It leads to the upregulation of expression of gene sets that are involved in pathways aimed at minimizing cellular damage and enabling survival. Among the many rescue pathways activated in response to hypoxia are genes involved in the initiation of angiogenesis, in inhibiting apoptotic pathways, and in the maintenance of cellular structure [29]. This was demonstrated in meta-analysis of hypoxic transcriptomes of many cell types [10]. Our data demonstrated typical changes in the transcriptome profile in the NIH3T3 cell response to hypoxia (Table S2). Likewise, nuclear localization of N-VEGF in various normal as well as cancer cell lines and tissues was also reported [3-6, 30]. Therefore, our model of an autoregulatory role for N-VEGF (Fig. 5) is physiologically relevant and represents its genuine potential.

The lack of effect of N-VEGF depletion on VEGF expression levels is explained by the existence of two separate Internal Ribosome Entry Site (IRES) in the 5' end of VEGF. These two IRES) sequences are functional for non-canonical translation under hypoxic conditions when

normal ribosome scanning mechanism is halted (see illustration in Fig. 1A) [9]. N-VEGF deletion results in impaired IRES A and likely IRES B is sufficient for effective translation of VEGF under these conditions [31].

This study provided complementary proof for the essential role of N-VEGF in activating the hypoxic response. We identified an exclusive gene set whose expression was modified by the mere mobilization of N-VEGF to the nucleus. Strikingly, although the hypoxic response is regulated by multiple factors, a significant number of the genes whose expression was modified when N-VEGF was shuttled to the nucleus had the inverse impact when hypoxia was induced in ΔN-VEGF cells. This set of genes included Hif1α and VEGF and a set of genes relevant to hypoxia. Functionally, N-VEGF deletion impaired cell death defense against hypoxic stress.

Although VEGF has been extensively studied in numerous physiological systems over more than three decades and N-VEGF was identified about 20 years ago, our study provides the first evidence of the functional role of N-VEGF. The regulatory architecture of the gene is comprised of two separate arms activated by the same transcript. While VEGF is secreted and activates angiogenesis in surrounding cells, N-VEGF shuttles to the nucleus and regulates intracellular gene responses, thereby serving as the autoregulatory arm of VEGF. This regulatory architecture is beneficial because it provides simultaneous regulation of both intra- and inter-cellular responses by a single promoter (see schematic illustration in Fig. 5). While other biological systems, such as the pro-opiomelanocortin transcript that generates different secreted hormones by proteolytic cleavage (for review see [32]) or the delta-notch signaling in which a segment of the notch receptor is sent to the nucleus and regulates transcription (for review see [33]), to our knowledge, this is the first report of a mechanism in which the same transcript regulates intra- and

inter- cellular response. It would be interesting to identify additional transcripts having similar regulatory architecture.

The extent to which N-VEGF plays a regulatory role in hypoxic responses and in angiogenesis in physiological and pathophysiological contexts remains to be determined. Circumstantial evidence suggests involvement of N-VEGF in various pathologies. Genomic polymorphism resulting in IRES B dysfunction was shown to reduce L-VEGF levels in ALS, in macular retinal thickness, and found to be associated with an increased risk of breast cancer aggressiveness, gastric as well as prostate cancers (reviewed in [1]). The role of N-VEGF in models of these diseases as well as in normal development remains to be examined.

Since this study uncovered a previously unknown regulatory mechanism of VEGF, many elementary questions regarding the manner in which N-VEGF exerts its effect remain to be addressed. Future experiments will determine if N-VEGF binds to the chromatin, elucidate the genomic loci it interacts with and identify its partners. In addition, deciphering its three-dimensional structure will help in understanding its specific role.

Acknowledgements

We are grateful to the Gellman-Lasser Fund for Medical & Biomedical Research and innovation (grant No. 907679) to B.Z.L, and to the Technion's Genome Center and the Russell Berrie Nanotechnology Institute for their support. We are also deeply grateful to Prof. Gera Neufeld, Faculty of Medicine, Prof. Yuval Shoham, Faculty of Biotechnology and Food engineering and Dr. Nitsan Fourier, Genomic Centre, Technion, for their input and critical reading of the manuscript. We would like to thank Dr. Yehuda Shabtai from UPenn for fruitful discussions. The initial construction of the Tet-On lenti-vectors by Mrs Tania Vorobyov, research assistant, is

acknowledged. B.Z.L. is an incumbent of the Lily and Silvian Marcus Chair in Life Sciences, Technion.

References

1.   Arcondeguy T, Lacazette E, Millevoi S, Prats H, Touriol C. VEGF-A mRNA processing, stability and translation: a paradigm for intricate regulation of gene expression at the post-transcriptional level. Nucleic Acids Res. 2013;41(17):7997-8010. doi: 10.1093/nar/gkt539. PubMed PMID: 23851566; PubMed Central PMCID: PMCPMC3783158.

2.   Apte RS, Chen DS, Ferrara N. VEGF in Signaling and Disease: Beyond Discovery and Development. Cell. 2019;176(6):1248-64. Epub 2019/03/09. doi: 10.1016/j.cell.2019.01.021. PubMed PMID: 30849371; PubMed Central PMCID: PMCPMC6410740.

3.   Meiron M, Anunu R, Scheinman EJ, Hashmueli S, Levi BZ. New isoforms of VEGF are translated from alternative initiation CUG codons located in its 5'UTR. Biochem Biophys Res Commun. 2001;282(4):1053-60.

4.   Tee MK, Jaffe RB. A precursor form of vascular endothelial growth factor arises by initiation from an upstream in-frame CUG codon. Biochem J. 2001;359(Pt 1):219-26.

5.   Huez I, Bornes S, Bresson D, Creancier L, Prats H. New vascular endothelial growth factor isoform generated by internal ribosome entry site-driven CUG translation initiation. Mol Endocrinol. 2001;15(12):2197-210.

6.   Rosenbaum-Dekel Y, Fuchs A, Yakirevich E, Azriel A, Mazareb S, Resnick MB, et al. Nuclear localization of long-VEGF is associated with hypoxia and tumor angiogenesis. Biochem Biophys Res Commun. 2005;332(1):271-8. doi: 10.1016/j.bbrc.2005.04.123. PubMed PMID: 15896327.

7.   Shih SC, Claffey KP. Hypoxia-mediated regulation of gene expression in mammalian cells. Int J Exp Pathol. 1998;79(6):347-57. Epub 1999/05/13. doi: 10.1046/j.1365-2613.1998.00088.x. PubMed PMID: 10319016; PubMed Central PMCID: PMCPMC3220374.

8.   Gossen M, Freundlieb S, Bender G, Muller G, Hillen W, Bujard H. Transcriptional activation by tetracyclines in mammalian cells. Science. 1995;268(5218):1766-9. PubMed PMID: 7792603.

9.   Lum JJ, Bui T, Gruber M, Gordan JD, DeBerardinis RJ, Covello KL, et al. The transcription factor HIF-1alpha plays a critical role in the growth factor-dependent regulation of both aerobic and anaerobic glycolysis. Genes Dev. 2007;21(9):1037-49. Epub 2007/04/18. doi: 10.1101/gad.1529107. PubMed PMID: 17437992; PubMed Central PMCID: PMCPMC1855230.

10.   Bono H, Hirota K. Meta-Analysis of Hypoxic Transcriptomes from Public Databases. Biomedicines. 2020;8(1). Epub 2020/01/16. doi: 10.3390/biomedicines8010010. PubMed PMID: 31936636; PubMed Central PMCID: PMCPMC7168238.

11.   Zhou Y, Zhou B, Pache L, Chang M, Khodabakhshi AH, Tanaseichuk O, et al. Metascape provides a biologist-oriented resource for the analysis of systems-level datasets. Nature communications. 2019;10(1):1523. Epub 2019/04/05. doi: 10.1038/s41467-019-09234-6. PubMed PMID: 30944313; PubMed Central PMCID: PMCPMC6447622.

12.   Ferrara N, Gerber HP, LeCouter J. The biology of VEGF and its receptors. Nat Med. 2003;9(6):669-76. doi: 10.1038/nm0603-669. PubMed PMID: 12778165.

13.   Stelzer G, Rosen N, Plaschkes I, Zimmerman S, Twik M, Fishilevich S, et al. The GeneCards Suite: From Gene Data Mining to Disease Genome Sequence Analyses. Curr Protoc Bioinformatics. 2016;54:1 30 1-1 3. Epub 2016/06/21. doi: 10.1002/cpbi.5. PubMed PMID: 27322403.

14.   Wilhelm M, Kukekov NV, Schmit TL, Biagas KV, Sproul AA, Gire S, et al. Sh3rf2/POSHER protein promotes cell survival by ring-mediated proteasomal degradation of the c-Jun N-terminal kinase scaffold POSH (Plenty of SH3s) protein. J Biol Chem. 2012;287(3):2247-56. Epub 2011/12/01. doi: 10.1074/jbc.M111.269431. PubMed PMID: 22128169; PubMed Central PMCID: PMCPMC3265902.

15.   Bae YH, Joo H, Bae J, Hyeon SJ, Her S, Ko E, et al. Brain injury induces HIF-1alpha-dependent transcriptional activation of LRRK2 that exacerbates brain damage. Cell Death Dis. 2018;9(11):1125. Epub 2018/11/14. doi: 10.1038/s41419-018-1180-y. PubMed PMID: 30420654; PubMed Central PMCID: PMCPMC6232134.

16.   Abu-Ghazaleh R, Kabir J, Jia H, Lobo M, Zachary I. Src mediates stimulation by vascular endothelial growth factor of the phosphorylation of focal adhesion kinase at tyrosine 861, and migration and anti-apoptosis in endothelial cells. Biochem J. 2001;360(Pt 1):255-64. Epub 2001/11/07. doi: 10.1042/0264-6021:3600255. PubMed PMID: 11696015; PubMed Central PMCID: PMCPMC1222225.

17.   Varela-Guruceaga M, Belaidi E, Lebeau L, Aka E, Andriantsitohaina R, Giorgetti-Peraldi S, et al. Intermittent Hypoxia Mediates Caveolae Disassembly That Parallels Insulin Resistance Development. Front Physiol. 2020;11:565486. PubMed PMID: 33324235.

18.   Huang B, Xiao E, Huang M. MEK/ERK pathway is positively involved in hypoxia-induced vasculogenic mimicry formation in hepatocellular carcinoma which is regulated negatively by protein kinase A. Med Oncol. 2015;32(1):408. Epub 2014/12/10. doi: 10.1007/s12032-014-0408-7. PubMed PMID: 25487444.

19. Pandey P, Kumari S, Ali Z, Pasha MQ. Acute hypobaric hypoxia augments ROCK2 protein level and activity. Exp Lung Res. 2017;43(4-5):181-6. Epub 2017/06/29. doi: 10.1080/01902148.2017.1333174. PubMed PMID: 28657365.

20. Koritzinsky M, Magagnin MG, van den Beucken T, Seigneuric R, Savelkouls K, Dostie J, et al. Gene expression during acute and prolonged hypoxia is regulated by distinct mechanisms of translational control. Embo j. 2006;25(5):1114-25. Epub 2006/02/10. doi: 10.1038/sj.emboj.7600998. PubMed PMID: 16467844; PubMed Central PMCID: PMCPMC1409715.

21. Thompson MA, Edmonds MD, Liang S, McClintock-Treep S, Wang X, Li S, et al. miR-31 and miR-17-5p levels change during transformation of follicular lymphoma. Hum Pathol. 2016;50:118-26. Epub 2016/03/22. doi: 10.1016/j.humpath.2015.11.011. PubMed PMID: 26997445; PubMed Central PMCID: PMCPMC4804154.

22. Chao LM, Sun W, Chen H, Liu BY, Li PF, Zhao DW. MicroRNA-31 inhibits osteosarcoma cell proliferation, migration and invasion by targeting PIK3C2A. Eur Rev Med Pharmacol Sci. 2018;22(21):7205-13. Epub 2018/11/24. doi: 10.26355/eurrev_201811_16254. PubMed PMID: 30468462.

23. Yamazato Y, Shiozaki A, Ichikawa D, Kosuga T, Shoda K, Arita T, et al. Aquaporin 1 suppresses apoptosis and affects prognosis in esophageal squamous cell carcinoma. Oncotarget. 2018;9(52):29957-74. Epub 2018/07/26. doi: 10.18632/oncotarget.25722. PubMed PMID: 30042826; PubMed Central PMCID: PMCPMC6057448.

24. Bai JA, Xu GF, Yan LJ, Zeng WW, Ji QQ, Wu JD, et al. SGK1 inhibits cellular apoptosis and promotes proliferation via the MEK/ERK/p53 pathway in colitis. World J Gastroenterol.

2015;21(20):6180-93. Epub 2015/06/03. doi: 10.3748/wjg.v21.i20.6180. PubMed PMID: 26034353; PubMed Central PMCID: PMCPMC4445095.

25.   Han Z, Jiang G, Zhang Y, Xu J, Chen C, Zhang L, et al. Effects of RNA interference-mediated NRP-1 silencing on the proliferation and apoptosis of breast cancer cells. Mol Med Rep. 2015;12(1):513-9. Epub 2015/03/05. doi: 10.3892/mmr.2015.3405. PubMed PMID: 25738638.

26.   Sandner P, Wolf K, Bergmair U, Gess B, Kurtz A. Hypoxia and cobalt stimulate vascular endothelial growth factor receptor gene expression in rats. Pflugers Arch. 1997;433:803-8.

27.   Li Y, Zhang H, Xie M, Hu M, Ge S, Yang D, et al. Abundant expression of Dec1/stra13/sharp2 in colon carcinoma: its antagonizing role in serum deprivation-induced apoptosis and selective inhibition of procaspase activation. Biochem J. 2002;367(Pt 2):413-22. Epub 2002/07/18. doi: 10.1042/BJ20020514. PubMed PMID: 12119049; PubMed Central PMCID: PMCPMC1222902.

28.   Bull VH, Thiede B. Proteome analysis of tunicamycin-induced ER stress. Electrophoresis. 2012;33(12):1814-23. Epub 2012/06/29. doi: 10.1002/elps.201100565. PubMed PMID: 22740470.

29.   Bristow RG, Hill RP, Kaplan AR, Glazer PM. Hypoxia and metabolism. Hypoxia, DNA repair and genetic instability

Impact of hypoxia on DNA repair and genome integrity. Nat Rev Cancer. 2008;8(3):180-92. Epub 2008/02/15. doi: 10.1038/nrc2344. PubMed PMID: 18273037.

30.   Bastide A, Karaa Z, Bornes S, Hieblot C, Lacazette E, Prats H, et al. An upstream open reading frame within an IRES controls expression of a specific VEGF-A isoform. Nucleic Acids Res. 2008;36(7):2434-45. doi: 10.1093/nar/gkn093. PubMed PMID: 18304943; PubMed Central PMCID: PMCPMC2367723.

31.   Bornes S, Prado-Lourenco L, Bastide A, Zanibellato C, Iacovoni JS, Lacazette E, et al. Translational induction of VEGF internal ribosome entry site elements during the early response to ischemic stress. Circ Res. 2007;100(3):305-8. Epub 2007/01/27. doi: 10.1161/01.RES.0000258873.08041.c9. PubMed PMID: 17255526.

32.   Cawley NX, Li Z, Loh YP. 60 YEARS OF POMC: Biosynthesis, trafficking, and secretion of pro-opiomelanocortin-derived peptides. J Mol Endocrinol. 2016;56(4):T77-97. Epub 2016/02/18. doi: 10.1530/jme-15-0323. PubMed PMID: 26880796; PubMed Central PMCID: PMCPMC4899099.

33.   Henrique D, Schweisguth F. Mechanisms of Notch signaling: a simple logic deployed in time and space. Development. 2019;146(3). Epub 2019/02/03. doi: 10.1242/dev.172148. PubMed PMID: 30709911.

34.   Schneider CA, Rasband WS, Eliceiri KW. NIH Image to ImageJ: 25 years of image analysis. Nat Methods. 2012;9(7):671-5. Epub 2012/08/30. doi: 10.1038/nmeth.2089. PubMed PMID: 22930834; PubMed Central PMCID: PMCPMC5554542.

Figures

Fig. 1 Hypoxia-independent nuclear mobilization of N-VEGF using the Cas9-NLS system. **(A)** Schematic illustration of the VEGF mRNA transcript, IRES domains and the translated products. Two translational products are generated. The non-canonical L-VEGF translation start site is CUG, and the canonical start site of VEGF is ATG. The locations of IRES A and B promoting translation under hypoxic stress are indicated. VEGF and L-VEGF proteins are illustrated as well as the proteolytic cleavage site of L-VEGF resulting in N-VEGF. **(B)** NIH3T3 cells were transduced with either N-VEGF or NLS-N-VEGF expression constructs without or with the inducer of the Tet-on promoter (Dox). Nuclear as well as cytoplasmic fractions were obtained from these cells. Western blot with anti N-VEGF antibodies was performed on these extracts. Subsequent Western blots on the same membrane were done using antibodies against LDH and histone H3 to show the purity of cytoplasmic and nuclei fractions, respectively. Ponceau S staining was carried out to demonstrate equal protein loading for all cytoplasmic and nuclear fractions. **(C)** Immunofluorescence of N-VEGF and NLS-N-VEGF cells before and following hypoxia (green) and nuclear staining with DAPI (red) is demonstrated.

Fig. 2 Transcriptional impact of hypoxia-independent mobilization of N-VEGF to the nucleus. (A) The top gene ontology terms of the 123 genes upregulated when N-VEGF is shuttled to the nucleus. **(B)** Qualitative comparison between 155 differentially expressed genes following mobilization of N-VEGF to the nucleus and their expression trend in wild type cells exposed to hypoxia. The columns represent a qualitative heatmap showing the direction of change in gene expression level (up-red, down-blue, no change-white) in NLS-N-VEGF cells following dox treatment (NLS-NVEGF, left column) in comparison to NIH3T3 cell under hypoxia (3T3 H, right column). The numbers within the right column represent the number of genes in each group that were annotated (detailed annotations in Fig. S2 A-D). **(C)** Validation of RNA seq results using qPCR. Real-time RT-qPCR analyses were performed on select

upregulated genes from each functional annotation group. The data is presented for L-VEGF-expressing cells before (yellow) and following treatments with dox (dark yellow), NLS-L-VEGF cells before (light blue) and following treatments with dox (blue) and NLS-L-VEGF cells treated with dox plus hypoxia (dark blue). The data were compared to control NIH3T3 cells before (light green) and following hypoxia (green). Genes associated with angiogenesis (Angio), Apoptosis (Apo) and cell structure (Cyto) are indicated. Fold changes in mRNA levels in triplicates from three independent experiments were determined. Statistical significance was determined by two-way ANOVA, * p≤0.05, ** p≤0.01). Values are mean± AVEDEV.

Fig. 3 Genomic deletion of N-VEGF using CRISPR-Cas9. (A) Cytoplasmic and nuclear fractions were prepared from NIH3T3 and N-VEGF deleted cells (ΔNVEGF). Western blots were performed with antibodies directed against N-VEGF and VEGF showing complete deletion of endogenous N-VEGF using CRISPR-Cas9. Numbers below lanes indicate the average relative density of the band from three different gels as determined using ImageJ software [34]. Statistical significance was determined by Student's t-test, p≤0.05. **(B)** To demonstrate the purity of soluble cytoplasmic and nuclear protein extracts, Western blots and Ponceau S staining were performed as elaborated in Fig.1 **(C)** Immunostaining showing complete absence of N-VEGF in deleted cells was performed as elaborated in Fig 1.

Fig. 4 Impact of N-VEGF deletion on hypoxia-induced genes and on cellular physiology. (A) Shown is a heatmap demonstrating gene expression levels in NIH3T3 cells (3T3) and ΔN-VEGF counterpart cells (ΔNVEGF) under normoxia (N) and following hypoxia (H). The heatmap presents only the 3,000 genes modified following hypoxia in the wild type. The genes were divided into six groups (G1-G6) of upregulated (red), downregulated (blue) expression in hypoxia. The numbers of differential genes in each group is indicated next to the

name of each group (red-up and blue-down). **(B)** A qualitative comparison between the 155 genes that exhibited differential expression when N-VEGF was shuttled to the nucleus and the expression trend of these genes in ΔN-VEGF cells following hypoxia. The numbers of genes in each group is indicated in the right column. The annotations of genes (68) that were upregulated in NLS-N-VEGF cells following dox and downregulated in ΔN-VEGF cells are presented. **(C)** To induce stress signals, NIH3T3 cells and their N-VEGF-deleted counterparts were exposed to hypoxia, $CoCl_2$, starvation, and tunicamycin for 18 h. Apoptosis was determined using the ApoTox-Glo^{TM} Triplex Assay. The results represent three different experiments that were performed in triplicates. Statistical significance was determined by Student's t-test, * $p<0.05$, ** $p<0.01$).

Fig. 5 N-VEGF as an autoregulator

Schematic illustration of L-VEGF processing and autoregulation by N-VEGF.

Supporting Information

Fig. S1 Annotation and pathway analysis of the 32 differentially expressed genes that were down regulated following hypoxia independent nuclear mobilization of NLS-N-VEGF. The annotations for the downregulated genes was performed as elaborated under Fig.1.

Fig. S2 Annotation of 155 genes that exhibited differential expression in NLS-N-VEGF cells and NIH3T3 cells following hypoxia. (A) Annotation of genes (51) that were upregulated in both cell types. (B) annotation of genes (72) that were upregulated in NLS-N-VEGF cells following dox and not changed or downregulated in NIH3T3 cells. (C) annotation of genes (18) that were downregulated in NLS-N-VEGF cells following dox and upregulated or not changed in NIH3T3 cells. (D) Annotation of genes (14) that were downregulated in both cell types. Bars below dotted line represent less significant annotation groups.

Fig. S3. Schematic illustrations. Schematic illustration of the two-steps protocol for CRISP-Cas9 mediated biallelic deletion of N-VEGF.

Fig. S4. PCR analysis of NIH3T3 with a replacement of the N-VEGF gene with a reporter cassette. (A) Schematic illustration of the reporter cassette harboring a reporter gene followed by Puromycin or Hygromycin resistance genes. Primer pairs for each junction were used for clones analysis. **(B)** Agarose gel showing the expected 435bp PCR product from 5' junction of isolated clone 1 (panel 1) obtained using the 5'-del NVEGF forward primer and reverse primer EF1-R from the insert. **(C)** Agarose gel showing levels of the expected 482bp PCR product from the 3' junction (panel 1), obtained using the SV40 forward primer and 3' del NVEGF reverse primer.

Fig. S5. PCR analysis to verify genomic N-VEGF deletion from NIH3T3. (A) Schematic illustration of the reporter cassette harboring a reporter gene followed by Puromycin or Hygromycin resistance genes. Primer pairs for each junction were used for clones analysis. **(B)** Agarose gel showing levels of the expected band of 435 bp PCR product from the 5' junction of clone 1 before (panel 1) and following VCre expression (panel 3). Removal of the reporter cassette was determined using 5'-del NVEGF forward primer and reverse primer EF1-R from the inset. **(C)** Agarose gel showing levels of the expected band of 482 bp PCR product from the 3' junction, obtained using the SV40 forward primer and 3'-del NVEGF reverse primer before (panel 1) versus after VCre expression (panel 3).

Table S1. Gene list and annotation of 155 genes the exhibited differential expression in NLS-N-VEGF cells treated with dox. Annotation of 123 upregulated genes (up tab) and 32 downregulated genes (down tab) in NLS-N-VEGF cells treated with dox.

Table S1.xlsx

Table S2. Annotation of genes that exhibited differential expression between Δ-N-VEGF and NIH3T3 cell following hypoxia. The genes are arranged in six groups (G1-G6). G1 and G6, similar expression pattern in both cell types of genes either induced or repressed following hypoxia, respectively. G3 and G4, inverse expression pattern in Δ-N-VEGF cells, either upregulated (G3) or downregulated (G4) following hypoxia in comparison to NIH3T3 under the same treatment. Last, genes that exhibited no change in expression pattern in ΔN-VEGF cells that were either upregulated or downregulated in NIH3T3 cells following hypoxia, groups G2 and G5, respectively. Statistically significant annotation groups, Log q < 0.05.

Table S2.xlsx

Table S3. List of primers used in this study.

Table S3.xlsx