Understanding associative false memories in aging using multivariate analyses

Nancy A. Dennis[1], Amy A. Overman[2], Catherine M. Carpenter[1], &

Courtney R. Gerver[1]

[1]Department of Psychology, The Pennsylvania State University, University Park, PA

[2] Department of Psychology, Elon University, Elon, NC

Corresponding author:

Nancy A. Dennis

Professor

Department of Psychology

The Pennsylvania State University

450 Moore Building

University Park, Pennsylvania 16802

nad12@psu.edu

Keywords: associative memory, false memory, aging, ERS, RSA

ABSTRACT

Age-related declines in associative memory are ubiquitous, having been observed across a wide array of stimuli and experimental paradigms. Research further shows that such decreases in behavioral discriminability arise from increases in false memories for recombined lures. The current study examined the underlying neural basis of associative false memories by using representational similarity analyses to examine both age differences in the reactivation of encoded representations during retrieval and the neural overlap in the similarity of neural patterns underlying targets and related lures during retrieval. Behaviorally, we observed an age-related reduction in d', which was shown to be driven by increased false alarms in the older adults. While we found no age difference in the relationship between patterns of neural activity underlying hits across memory phases (as measured by ERS), the similarity of neural patterns underlying targets and lures was affected by age. Specifically, while younger and older adults exhibited overall higher pattern similarity between hits and CRs compared to hits and FAs (as measured by RSA), this difference was reduced in older adults within occipital cortices. Additionally, greater Hit-FA representational similarity correlated with increases in associative FAs across occipital, frontal, and parietal cortices. Results suggest that while neural representations underlying targets may not differ across age, greater pattern similarity between the neural representation of targets and lures may reflect reduced distinctiveness of the information encoded in memory, such that old and new items are more difficult to discriminate, leading to more false alarms.

Introduction

Associative memory is essential to daily life. It requires the ability to not only remember items and events of the past, but to remember the specific relationship between details of a given event. For example, it is important to remember both the faces and the names of new people you meet, as well as remembering which name is associated to each face. Misremembering one or more of these associative details, such as calling someone by the wrong name, can feel like a relatively minor memory error, but even these minor false memory errors can cause embarrassment and frustration. Associative memory success is most critical in cases where the consequences of errors are more serious, such as remembering medications and their accompanying dosages, a task that many older adults regularly encounter in their daily routines. Unfortunately, while associative false memories occur throughout the lifespan, they have been shown to be more frequent in older individuals (Chen & Naveh-Benjamin, 2012; Kilb & Naveh-Benjamin, 2007; Naveh-Benjamin, 2000; Naveh-Benjamin et al., 2004). Given that successfully navigating the world depends on making and retrieving accurate associations, it is important to understand how such associative false memories arise and what leads to increased error rates in older adults.

Age deficits in associative memory have been observed using a wide array of stimuli and experimental paradigms (e.g., Chalfonte & Johnson, 1996; Chen & Naveh-Benjamin, 2012; Kausler & Puckett, 1981a, 1981b; Kilb & Naveh-Benjamin, 2007; Naveh-Benjamin, 2000; Naveh-Benjamin et al., 2004; Naveh-Benjamin & Craik, 1995; Park et al., 1982; Park & Puglisi, 1985; Spaniol et al., 2006). The focus of this prior work has largely been on associative encoding and has examined how age-related deficits in binding contribute to memory errors (Earles et al., 2016; Nashiro & Mather, 2010; Naveh-Benjamin, 2000; Old & Naveh-Benjamin, 2008).

Specifically, behavioral and hippocampal evidence suggests that older adults exhibit a reduced ability to encode the relationship amongst discrete pieces of information, leading to associative information not being as well remembered as individual items (e.g., Bastin & Van der Linden, 2006; Chalfonte & Johnson, 1996; Dennis, Hayes, et al., 2008; Giovanello, Kensinger, Wong, & Schacter, 2009; Mitchell, Raye, Johnson, & Greene, 2006; Naveh-Benjamin, 2000; Sperling et al., 2003). While this deficit is largely conceptualized as an age-related reduction in associative hits (i.e., correctly recognizing, at retrieval, an associative pair previously seen at encoding), a closer look at the results of prior studies, our own included (Dennis, Hayes, et al., 2008; Overman, McCormick-Huhn, Dennis, Salerno, & Giglio, 2018), suggests that older adults' associative memory deficit also arises from the endorsement of recombined lures (i.e., test pairs in which each individual stimulus had originally appeared as part of two different encoding pairs) during retrieval. This results in higher false alarm (FA) rates in older compared to younger adults (e.g., Castel & Craik, 2003; Healy, Light, & Chung, 2005; Naveh-Benjamin, Hussain, Guez, & Bar-On, 2003; Old & Naveh-Benjamin, 2008b; Overman & Stephens, 2013; Rhodes, Castel, & Jacoby, 2008).

False memories become increasingly pronounced in older adults when there is a high degree of perceptual overlap between studied targets and lures presented at retrieval (Koutstaal & Schacter, 1997; Norman & Schacter, 1997). This overlap leads to a reclassification of related lures as "remembered" at a similar rate as targets (Glanzer & Adams, 1985; Hockley, 2008). Univariate analyses assessing the neural basis of perceptual false memories in both young and older adults have shown there to be a high amount of overlap in the physical location of activated voxels across true and false memories. For example, both true and false memories activate regions within medial prefrontal cortex, angular gyrus, occipital cortices and the medial temporal

lobe (e.g., Dennis, Bowman, & Peterson, 2014; McDermott, Gilmore, Nelson, Watson, & Ojemann, 2017; Schacter, Buckner, Koutstaal, Dale, & Rosen, 1997; Webb & Dennis, 2019). Such overlap in neural activation has been taken as an indication of the similarity in neural processes across the two trial types. But given the nature of univariate analyses, they have not been able to speak to an overlap in the nature of the neural processing across trial types.

One reason associative memory tasks are especially challenging to older adults is that the *items* in recombined lures are highly familiar to the participant, having been presented previously. Yet an individual must decide not simply that they have seen the items previously, but whether that specific *pair of items* was previously presented together. It has been suggested that higher rate of false associative memories in older adults is due to their failure to make use of successfully encoded perceptual details when confronted with lures at retrieval, particularly when those lures are novel pairs comprised of previously seen items (Bulevich & Thomas, 2012; Koutstaal, 2003; Mitchell, Ankudowich, Durbin, Greene, & Johnson, 2013).

Reflecting this difficulty, and consistent with recent work showing that reductions in neural distinctiveness underlie perceptual false memories (Bowman, Chamberlain, & Dennis, 2019; Lee, Samide, Richter, & Kuhl, 2019), we posit that older adults will show greater overlap in the similarity of neural patterns underlying hits and false alarms, compared to patterns underlying hits and correct rejections. That is, we predict that representational similarity will be greater for retrieval pairs that are behaviorally labeled as "old" compared to the overlap of targets and lures that are correctly differentiated. We also posit that this representational overlap between hits and false alarms will be related to false alarm rates in older adults. To this point, neural pattern similarity analysis in younger adults has shown that false memories are more prevalent when there is higher overlap in the representational similarity between targets and

related lures in with cortical regions critical to memory processing (Chadwick et al., 2016; Wing et al., 2020). The above-mentioned multivariate studies examining false memories, as well as our own univariate meta-analysis of false memories (Kurkela & Dennis, 2016), suggests that perceptual processing regions, including inferior and middle occipital cortex, lateral temporal cortices, and frontoparietal cortices may be critical regions for investigating representational similarity differences between age groups, based on the importance of these regions in mediating false memories across a wide range of memory paradigms.

Given this proposed overlap between targets and lures, the transfer of information from encoding to retrieval is a key component to associative memory success, including the ability to discount new information containing some familiar aspects (i.e., lure pairs comprised of familiar items) by retrieving strong encoding traces of target information. Specifically, with regard to associative memory, not only do the individual items need to be encoded well and recapitulated at retrieval, but critically the relationship between them needs to be recapitulated in order to endorse the *target* pair (a re-presentation of target items in the exact same pairing) and reject a *lure* pair (a recombination of target items in a novel pairing). When the encoded association is not sufficiently reinstated at retrieval, a lure may be mistakenly endorsed, resulting in a false memory. The reactivation of this encoded representation can be examined by assessing the similarity of neural patterns between individual target trials across memory phases (i.e., encoding/retrieval) using encoding–retrieval similarity (ERS) analysis (e.g., Jonker, Dimsdale-Zucker, Ritchey, Clarke, & Ranganath, 2018; Kuhl, Rissman, Chun, & Wagner, 2011; Ritchey, Wing, LaBar, & Cabeza, 2013; Wing, Ritchey, & Cabeza, 2015). Previous work using ERS analysis within the domain of episodic memory has suggested that overlapping patterns of activation between encoding and retrieval correlates with both item and source memory success.

For example, research shows that ERS is capable of detecting the reinstatement of individual episodic memories in the hippocampus (Tompary, Duncan, & Davachi, 2016), as well as in other temporal cortex regions, the occipital lobe, and the frontoparietal cortex (Wing et al., 2015; Xiao et al., 2017). Additionally, within the domain of associative memory, work from our group suggests that one's ability to generate similar neural patterns across memory phases differs with respect to the similarity of stimulus configuration across those memory phases (Gerver, Overman, Babu, Hultman, & Dennis, 2020). Examining age differences in ERS of targets helps us understand whether there are age differences in how the encoded representations of associative information are recapitulated at retrieval, and whether reductions in pattern similarity across memory phases contributes to age-related increases in false associative memories. We posit that, if older adults experience difficulty binding associative information during encoding, this would lead to deficits in recapitulating associative information during the retrieval process. To this end, we would expect to see age-related reductions in ERS within regions critical to perception and binding processes, including regions in the ventral visual cortex and hippocampus, with these reductions related to increases in false memory errors. Further, based on our previous work showing that ERS is modulated by the configural congruency between encoding and retrieval (Gerver et al., 2020), we posit that age-related changes in ERS may be greater for targets in which configural changes between encoding and retrieval reduce the familiarity of the associative pair across memory phases. Thus, the present study will investigate the neural basis of age-related increases in associative false memory by using representational similarity analysis to not only examine the overlap between hits and false alarms during retrieval, but also to examine whether reductions in target recapitulation is related to false memory errors.

Methods

Participants. Thirty-one right-handed young adults (the same sample as reported in Gerver et al.

2020) and thirty right-handed older adults were recruited from the Pennsylvania State University

community and greater State College area, respectively. All participants provided written

consent and received compensation for their participation. Participants were screened for history

of psychiatric and neurological illness, head injury, stroke, learning disability, medication that

affects cognitive and physiological function, and substance abuse. On the day of the study, all

participants provided written informed consent for a protocol approved by the Pennsylvania State

University Institutional Review Board. All participants were native English speakers with normal

or corrected-to-normal vision. One older adult was excluded due to an inability of the analysis

program to process their data and one older adult because they received the wrong version of the

memory task. Three younger adults were removed from the study due to incomplete data (e.g.,

they stopped the task early), one because of excess movement in the scanner, and one because of

misunderstanding the paradigm. Thus, the reported results are based on data from 26 young adult

participants ranging from 18 to 25 years old (19 female; age: M = 20.5 years, SD = 1.98 years)

and 28 older adult participants ranging from 63 to 83 years old (21 female; age: M= 71.04 years,

SD = 6.02 years)[1].

Cognitive Assessments. All participants provided demographic information. In addition, older

participants completed a short battery of cognitive assessments on the day of the study prior to

being scanned to assess general cognitive function. The total time taken to complete all tests was

---

[1] The current sample size in in line with several previous studies examining age-related differences in neural similarity during episodic memory (Folville et al., 2020; Hill, King, & Rugg, 2021; Srokova, Hill, Elward, & Rugg, 2021; Trelle, Henson, & Simons, 2019)

less than 30 minutes and a break was given after tests and before scanning. The cognitive assessment battery included MMSE (*M*=29.52, *SD*=0.98), Digit Symbol Coding (*M*=12.43, *SD*=2.29), Symbol Copy (*M*=102.5, *SD*=23.76), Digit Span (*M*=12.48, *SD*=2.64), Letter-Number Sequencing (*M*=12.09, *SD*=3.11), WAIS-III Vocabulary (*M*=11.39, *SD*=1.68), and the Geriatric Depression Scale (*M*=0.86, *SD*=1.14) (Kurlowicz & Wallace, 1999; Wechsler, 1981; Yesavage et al., 1982). The stimuli and procedure were the same as that reported in Gerver et al. 2020, and are replicated here for completeness.

Stimuli. Stimuli consisted of 170 color photographs of faces and 170 color photographs of scenes paired together. Face stimuli consisted of both male and females faces, each exhibiting a neutral expression, taken from the following online databases: the Color FERET Database (Phillips, Moon, Rizvi, & Rauss, 2000), adult face database from Dr. Denise Park's laboratory (Minear & Park, 2004), the AR Face Database (Martinez & Benavente, 1998), and the FRI CVL face data-base (Solina, Peer, Batageli, Juvan, & Kovac, 2003). Scene stimuli consisted of outdoor and indoor scenes collected from an Internet image search. Using Adobe Photoshop CS2 Version 9.0.2 and Irfanview 4.0 (www.irfanview.com/), we edited face stimuli to a uniform size (320 × 240 pixels) and background (black), and scene stimuli were standardized to 576 × 432 pixels.

Figure 1. Task design. Examples of encoding and retrieval conditions for both congruent and incongruent targets and lures.

During encoding, half of the face–scene pairs were presented side by side; and the other half, with the face superimposed on top of the scene (see Figure 1). At retrieval, half of the pairs were presented in a manner congruent with their original contextual configuration; and the other half, in a contextual configuration that was incongruent with how it had been originally presented (e.g., at encoding was superimposed, at retrieval was side-by-side). Each encoding and retrieval block consisted of 34 total pairs. Within the 34 pairs per retrieval block, 10 were lures (five of which were side-by-side configuration and five of which were superimposed configuration) in which the face was rearranged with a different scene than was originally presented at encoding. During encoding, pairs were presented for 4 sec and for 4 sec at retrieval. A jittered ISI (2–8 sec) separated the presentation of each image. Each encoding and retrieval block lasted 4 min and 18 sec. A second version of this task was created to counterbalance the design such that the same stimuli were presented in the alternate configuration across versions (e.g., a face–scene pair

presented side-by-side at encoding in Version A was presented as superimposed at encoding in Version B).

Procedure. Before scanning, all participants were provided the full set of instructions for encoding and retrieval and completed a practice session of the full task. Participants were also asked to explain the instructions verbally before proceeding with the experiment to verify an accurate understanding of the task. They were encouraged to ask questions during this time. The scanning session began with a structural scan (magnetization prepared rapid gradient echo) that took approximately 7 min. During this time, the participants were asked to remain as still as possible. After the structural scan, participants completed five encoding and five retrieval blocks presented in an alternating order. Instruction screens, mirroring those used in the practice session, were presented before each block reiterating the verbal instructions participants received before entering the scanner. Presentation of all instruction screens was self-paced, meaning that the participants pressed "1" on the handheld button box to advance to the next screen when they read the instructions and were ready to begin the task.

After advancing past the instruction slides in encoding, the participant was presented with a series of face and scene pairings displayed on the screen in either a side-by-side or superimposed configuration. Each pair was presented for 4 sec, during which time the participant responded to the question "How welcoming are the scene and face together?" Text of the question was presented below each pair by using a rating scale from 1 to 4 (1= not at all, 4=very) and making a key-press on their handheld button box (See Figure 1a). The instructions emphasized that the participants should choose the rating based on how welcoming the face and scene pairing was collectively in order to facilitate encoding of both the face and the scene, rather than only one or the other, or them separately. This question also helped ensure that

participants paid attention to the scene, even when it was configured behind the face in the superimposed condition because we did not want the scene to be incidentally encoded while the face was intentionally encoded. Across versions, faces and scenes were counterbalanced for their inclusions in either pair configuration. No differences across versions were noted, and all analyses were collapsed across versions.

Each encoding block was followed by a retrieval block. Similar to encoding, each face–scene pair at retrieval was presented for 4 sec. During this time, participants were asked to respond to the question: "Please identify the pairings that have been presented previously." Displayed below the question were the following choices: 1 = remember, 2 = know, and 3 = new. A remember–know–new design was chosen to isolate recollection-related activity, associated with "remember" responses, from that of familiarity, associated with "know" responses. This distinction has shown to be critical when assessing memory-related activity particularly within the MTL (Yonelinas, 2002; Yonelinas et al., 2005, 2007). Participants were instructed to make their memory judgments based on the co-occurrence of the face and scene and not to base their judgments on the configuration of the display. Specifically, participants were instructed to select "Remember" if they were able to retrieve details of the face and scene appearing together at study, select "Know" if they thought the face and scene appeared together previously but they could not remember specific details about the original appearance, and select "New" if they believe that the face–scene pair did not appear together previously. Thus, whether a retrieval configuration was congruent or incongruent with how it had been presented at encoding, the Remember/Know/New labels applied equally to both types of configurations, with a participant's response simply dependent on the vividness of their memory. Similar to the encoding phase, all responses were made using the button. With respect to retrieval

configurations, half of the trials were congruent with their encoding configuration and half were incongruent (such that a side-by-side trial at encoding was presented as a superimposed trial at retrieval) (See Figure 1).

Image Acquisition. Structural and functional images were acquired using a Siemens 3-T scanner equipped with a 12-channel head coil, parallel to the AC–PC plane. Structural images were acquired with a 1650-msec repetition time, a 2.03-msec echo time, a 256-mm field of view, a 2562 matrix, 160 axial slices, and a 1.0-mm slice thickness for each participant. Echo-planar functional images were acquired using a descending acquisition, a 2500-msec repetition time, a 25-msec echo time, a 240-mm field of view, a 802 matrix, a 90° flip angle, and 42 axial slices with a 3.0-mm slice thickness resulting in 3.0-mm isotropic voxels.

Image Processing. Raw anatomical and functional brain images were first skull stripped using the Brain Extraction Tool (Smith, 2002) in the FMRIB Software Library (FSL) Version 5.0.10 (www.fmrib.ox.ac.uk/fsl). FSL's MCFLIRT function (Jenkinson et al., 2002, 2012) was then applied for realignment and motion correction within each functional run. All volumes were aligned to the middle volume of the middle run of encoding. The realigned functional images were then processed by FSL's fMRI Expert Analysis Tool (Woolrich et al., 2001), where they were high-passed filtered and spatially smoothed at 6-mm FWHM. These data were then pre-whitened to account for temporal autocorrelations within voxels. Finally, the structural data underwent nonlinear transformation into the standardized Montreal Neurological Institute (MNI) space by using the warping function in FSL's FNIRT (Andersson et al., 2007). For multivariate analyses, the raw data underwent the exact same steps as above, absent smoothing.

Behavioral Analyses. All behavioral analyses were conducted in Rstudio using base R functions

and rstatix (www.rstudio.com; Kassabara, 2020). Because this study aimed to examine false

associative memory, behavioral and neuroimaging analyses were focused on accurately and

inaccurately recollected memory decisions, in the form of d' (calculated in Excel as

norm.s.inv(hits) – norm.s.inv(FAs)). To test for age differences in memory, three 2 (Between

Age Group: Old, Young) x 2 (Within Encoding Configuration: side-by-side or superimposed) x 2

(Within Retrieval Congruency: congruent or incongruent) mixed model ANOVAs were

performed. To examine significant main effects and interactions, paired-sample t-tests were run,

with Bonferroni correction.

Multivariate Pattern Similarity Analyses. To estimate neural activity associated with individual

trials, separate GLMs were estimated in SPM12 defining one regressor for each trial at encoding

and retrieval (170 in total for each respective memory phase) (Mumford, Davis, & Poldrack,

2014). An additional six nuisance regressors were included in each run corresponding to motion.

Whole-brain beta parameter maps were generated for each trial at retrieval for each participant.

In any given parameter map, the value in each voxel represents the regression coefficient for that

trial's regressor in a multiple regression containing all other trials in the run and the motion

parameters. These beta parameter maps were next concatenated across runs and submitted to the

CoSMoMVPA toolbox (Oosterhof et al., 2016) for similarity analyses (representational

similarity analysis: RSA and encoding-retrieval similarity analysis: ERS).

ROIs. Based on our focus on using ERS and RSA to better understand age-related increases in

false memories, all analyses were computed in ROIs previously identified as critical to false

memories. Specifically, we drew on regions identified both in past multivariate studies of false

memories (Bowman et al., 2019; Lee et al., 2019; Wing et al., 2020) and our meta-analysis of

neural activation underlying false memories (Kurkela & Dennis, 2016). ROIs used in the current study included the medial superior PFC (Medial SFG; N_{voxels} = 9,644), angular gyrus (AG; N_{voxels} = 801), inferior (IOC; N_{voxels} = 1,930) and middle occipital cortex (MOC; N_{voxels} = 5,368), and medial temporal gyrus (MTG; N_{voxels} = 10,013). We also included a subregion of the MTL, the bilateral hippocampus (HC; N_{voxels} = 1,878), that has been shown to be critical to associative memory and binding. While the general ROIs were drawn from this past work, specific ROIs in the current investigation were defined using AAL PickAtlas in SPM12 using anatomically defined boundaries identified by the anatomical labeling. The medial superior PFC region was created in the AAL PickAtlas based upon coordinates and activation identified in the Kurkela & Dennis meta-analysis.

ERS analysis. Encoding-retrieval similarity analysis (Kriegeskorte, Mur, & Bandettini, 2008; Ritchey et al., 2013) was conducted to examine whether there was an age deficit in how encoded information was re-represented at retrieval and whether this influenced false memory performance, we directly compared neural patterns of activation between encoding and retrieval. On the basis of our previous work indicating that encoding configuration (i.e., side-by-side, superimposed) significantly influences patterns of neural activity during encoding (Dennis et al., 2019) and configural congruency between encoding and retrieval was critical to ERS (Gerver et al., 2020), we separated all targets into four conditions of interest: side-by-side congruent targets, superimposed congruent targets, side-by-side incongruent targets, and superimposed incongruent targets. To compute pattern similarity, we used the beta estimates extracted from the single trial models for encoding and retrieval described above. Activation for each individual trial for a given condition at encoding was Fisher's z transformed and correlated with every trial of the same type at retrieval (e.g., targets at encoding that were presented as a side-by-side

configuration that were congruently re-presented as side-by-side at retrieval). This resulted in similarity scores, as operationalized by Pearson's *r* correlation values, for each trial. The correlations were then averaged within condition for each participant.

From these values, group averages were computed and age differences were assessed using a mixed factorial ANOVA, using the rstatix package in R (Kassambara, 2021), with age (young, old) as the between subjects variable and ROI (IOC, MOC, AG, MTG, medial SFG, hippocampus) and condition (side-by-side congruent, superimposed congruent, side-by-side incongruent and superimposed incongruent) as within subjects variables. To examine significant main effects and interactions, paired-sample t-tests were run and Bonferroni corrected using the emmeans R package (Lenth, 2021). Noted below in results, the ANOVA revealed no age differences in ERS. Using the BayesFactor package in R (Morey, Rouder, Jamil, & Morey, 2015) we provide Bayes factors that indicate how strong the evidence is for the null hypothesis (no age difference) relative to the alternative hypothesis (age difference) for ERS, collapsed across condition and ROI using an approach similar to Bowman, Ashby, and Zeithamova (2021). We did the same for our follow up ERS ANOVAs with ROIs when examining the absence of age effects in the main analyses.

RSA. A representational similarity analysis (Kriegeskorte et al., 2008) was conducted to examine overlap in the representation of stimuli associated with different behavioral outcomes. Specifically, given our interest in understanding whether older adults' increase in false memories was related to heightened pattern similarity across hits and FAs, we examined the similarity of patterns of neural activity between targets and related lures as a function of behavior.

Associative false memories

Specifically, we correlated neural pattern similarity between Hits and FAs (representing pattern similarity between different types of "old" responses; i.e., veridical and false memories) and between Hits and CRs (representing pattern similarity between different types of veridical memories) within each ROI described above. Given trial counts across each condition and the fact that false memories were not affected by contextual congruency, we collapsed across this factor in the current analysis. To compute pattern similarity, we used the beta estimates extracted from the retrieval single trial model described above to run two separate Pearson correlations between Hits and FAs and Hits and CRs. The resulting correlation matrices were Fisher's z transformed then averaged to get a single correlation value per trial pair (Hits and FAs and Hits and CRs; see Trelle et al., 2019 for a similar methodological approach) for each participant. From these values group averages were computed and age differences assessed using a mixed factorial ANOVA, using the rstatix package in R (Kassambara, 2021), with age (young, old) as the between subjects variable and ROI (IOC, MOC, AG, MTG, medial SFG, hippocampus) and trial pair (Hit and CR RSA, Hit and FA RSA) as within subjects variables. To examine significant main effects and interactions, paired-sample t-tests were run and Bonferroni corrected.

Results

Behavioral results

For the *d'* measures, the ANOVA revealed a significant main effect of age group, $F(1,52)=16.95$, $p<.001$, $\eta^2 = 0.25$, such that young adults ($M=1.40$, 95% C.I. [1.32, 1.49]) had better discrimination than older adults ($M= 0.68$, 95% C.I. [0.60, 0.75]).The main effect of encoding configuration was also significant, $F(1,52)=5.80$, $p=.020$, $\eta^2 = 0.10$, such that superimposed trials ($M=1.12$, 95% C.I. [1.03, 1.20]) had better discrimination between targets

and lures than side-by-side trials (M=0.93, 95% C.I. [0.84, 1.02]). The main effect of retrieval congruency and the two and three-way interactions were not significant (all F's < 3.14, all p's > .08).

Breaking the d' measure down by hit and FA rates we found that the hit ANOVA revealed a significant main effect of encoding configuration, $F(1,52)$=6.10, p=.017, η^2 =0.11, such that superimposed configurations (M=0.84, 95% C.I. [0.83, 0.85]) had higher hits than side-by-side (M=0.82, 95% C.I. [0.81, 0.83]). Additionally, there was a significant main effect of retrieval configuration congruency, $F(1,52)$=53.83, p<.001, η^2 =0.51, such that congruent trials (M=0.87, 95% C.I. [0.86, 0.88]) showed higher hit rates than incongruent trials (M=0.79, 95% C.I. [0.77, 0.80]). The main effect of age group and the two and three-way interactions were not significant (all F's<1.26, all p's>.27). With regard to FA rates, the ANOVA revealed a significant main effect of age group, $F(1,52)$=19.27, p<.001, η^2 = 0.27, such that older adults (M=0.63, 95% C.I. [0.60, 0.65]) had higher false alarms than younger adults (M= 0.39, 95% C.I. [0.37, 0.41]). The main effect of retrieval configural congruency was also significant, $F(1,52)$=22.18, p<.001, η^2 =0.30, such that congruent trials (M=0.55, 95% C.I. [0.52, 0.57]) showed higher false alarms than incongruent trials (M=0.48, 95% C.I. [0.45, 0.50]). The main effect of encoding congruency and the two and three-way interactions were not significant (all F's < 3.67, all p's >.06). [See Table 1 for means].

	Congruent		Incongruent	
	Item-Item	*Item-Context*	*Item-Item*	*Item-Context*
Young Adults	Mean(SD)	Mean(SD)	Mean(SD)	Mean(SD)
Hits	.86 (.08)	.89 (.07)	.77 (.10)	.81 (.11)
False Alarms	.45 (.24)	.39 (.21)	.35 (.22)	.36 (.23)
d'	1.31 (.85)	1.64 (.79)	1.22 (1.00)	1.43 (.81)
Older Adults				
Hits	.86 (.14)	.87 (.11)	.77 (.15)	.79 (.14)
False Alarms	.70 (.24)	.64 (.21)	.57 (.25)	.60 (.23)
d'	.65 (1.05)	.84 (.70)	.59 (.59)	.62 (.75)

Table 1. Behavioral performance broken down by condition and age group. Mean and standard deviation reported.

<u>*ERS*</u>. The main objective in analyses of ERS was to examine age differences across conditions. Thus, the post-hoc analyses based on the ERS ANOVA were focused on meaningful comparisons with respect to those factors. Critically, the ERS ANOVA revealed both a significant main effects of condition and ROI and a significant 3-way interaction between condition, age, and ROI. With respect to the main effect of condition, $F(3,153)=28.42$, $p<.001$, both congruent side-by-side *(M=0.010, SD=0.019)* and congruent superimposed *(M=0.011, SD=0.020)* ERS metrics were greater than either of the incongruent ERS metrics *(*side-by-side $M=0.0077$, $SD=0.021$; superimposed $M=0.0081$, $SD=0.021$). Specifically, superimposed congruent had greater ERS than superimposed incongruent, $t(52)=6.24$, $p<.001$, and side-by-side incongruent, $t(52)=6.99$, $p<.001$. Additionally, side-by-side congruent showed greater ERS than

superimposed incongruent, $t(52)=5.05$, $p<.001$, and side-by-side incongruent, $t(52)=5.83$,

$p<.001$. There was also a significant main effect of ROI, $F(2.13, 108.73)=21.20$, $p<.001$. There

was also a significant two-way interaction between Condition and ROI, $F(4.89, 249.32)$, $p<.001$

and a three-way interaction between age, condition and ROI was significant $F(4.89,$

$249.32)=2.46$, $p=.035$. The main effect of age $F(1,51)=0.75$, $p=.39$, the two-way interaction

between age group and condition $F(3, 153)=1.10$, $p=0.35$, interaction between age group and

ROI $F(2.13, 108.73)=0.85$, $p=.44$, was not significant. Consistent with the findings from the

ANOVA, a Bayesian analysis of age differences across ERS collapsed across condition and ROI

yields modest evidence in favor of the null hypothesis (JZS $BF_{01} = 1.11$).

To examine whether the three-way interaction between condition and ROI and age was

meaningful, we conducted a repeated measures ANOVA within each ROI. There was a

significant main effect of condition within the IOC, $F(3,156)=17.75$, $p<.001$ and MOC,

$F(3,156)=43.06$, $p<.001$. Follow-up paired-sample t-tests revealed that, within the IOC, ERS

significantly differed between the side-by-side congruent ($M=0.002$, 95% C.I. [0.002, 0.003])

and side-by-side incongruent targets ($M=-0.002$, 95% C.I. [-0.003, -0.001]), $t(52)= 4.59$, $p<.001$,

as well as superimposed incongruent targets ($M=-0.002$, 95% C.I. [-0.002, -0.001]), $t(52)= 4.07$,

$p<.001$. Additionally, ERS significantly differed between the superimposed congruent targets (M

$=0.004$, 95% C.I. [0.003, 0.005]) and the superimposed incongruent targets ($M=-0.002$, 95% C.I.

[-0.002, -0.001]), $t(52)= 5.08$, $p<.001$ as well as side-by-side incongruent targets ($M=-0.002$,

95% C.I. [-0.003, -0.001]), $t(52)= 5.33$, $p<.001$. See Figure 2a. A similar pattern was found in

the MOC where ERS significantly differed between side-by-side congruent targets ($M=0.005$,

95% C.I. [0.005, 0.006]) and side-by-side incongruent targets ($M=-0.004$,) 95% C.I. [-0.005, -

0.003], $t(52)= 7.43$, $p<.001$, superimposed congruent targets ($M=0.008$, 95% C.I. [0.007,

0.009]), $t(52)= -3.01$, $p<.001$, and superimposed incongruent targets (M=-0.004, 95% C.I. [-0.005, -0.002]), $t(52)= 6.58$, $p<.001$. Additionally, ERS significantly differed between superimposed congruent targets (M=0.008, 95% C.I. [0.007, 0.009]) and superimposed incongruent targets (M=-0.004, 95% C.I. [-0.005, -0.002]), $t(52)= 6.73$, $p<.001$ and side-by-side incongruent targets (M=-0.004, 95% C.I. [-0.005, -0.003]), $t(52)= 7.78$, $p<.001$. See Figure 2a. While neither ANOVA exhibited a main effect of age, a Bayesian analysis of age differences across ERS within the IOC collapsed across condition yields moderate evidence in favor of the alternative hypothesis (age differences) (JZS $BF_{10} = 8.96$), while a Bayesian analysis of age differences across ERS within the MOC collapsed across condition yields modest evidence in favor of the null hypothesis (no age differences) (JZS $BF_{01} = 1.57$).

There was no main effect of age in any ROI, suggesting that neural pattern similarity of all four contextual and configural conditions across encoding and retrieval yielded comparable ERS between young and older adults across all *a priori* regions of interest (all p's > .05). This also suggests that the main effect of age within the 3-way interaction resulted from differences across condition and ROI which would not be interpretable within the current analysis/design. See Supplemental Table A for ERS values reported for each age group for each ROI.] Finally, we found no correlation between ERS metrics and memory performance in either age group. [See Figure 2].

Figure 2. ERS in the Inferior Occipital Cortex (IOC) and Middle Occipital Cortex (MOC), broken down by the four target conditions in the younger and older adults. Error bars represent standard error of the mean.

<u>RSA</u>. Similar to the ERS analysis, Critically, the RSA ANOVA revealed both a significant main effects of trial type and ROI and a significant 3-way interaction between trial type, age, and ROI. With respect to the main effect of trial type ($F(1, 52) = 130.34$, $p < .001$), the average pattern similarity was higher for Hits and CRs ($M = 0.22$, $SD = 0.19$) than for Hits and FAs ($M = 0.13$, $SD = 0.10$). There was also a main effect of ROI ($F(1.66, 86.12) = 176.25$, $p < .001$), with no main effect of age group $F(1, 52) = 0.12$, $p = 0.73$ ($M_{old} = 0.17$, $SD_{old} = 0.15$; $M_{young} = .18$, $SD_{young} = 0.16$). Given the collapsed RSA means (i.e., trial type) are not interpretable, we chose not to evaluate these findings further, but to focus on interactions with trial type and age.

To that end, there was a significant trial type by ROI interaction ($F(1.44, 74.85) = 95.47$, $p < .001$), such that the pattern similarity between Hits and CRs was greater than that of Hits and FAs in the AG (Hit and CR: $M = 0.18$, 95% C.I. [0.17, 0.20]; Hit and FA: $M = 0.11$, 95% C.I. [0.10, 0.12]; $t(53) = 8.55$, $p < .001$), IOC, (Hit and CR: $M = 0.37$, 95% C.I. [0.35, 0.40]; Hit and

FA: M=.19, 95% C.I. [0.18, 0.21]; $t(53)$ =9.00 , $p < .001$), MOC (Hit and CR: M=0.47, 95% C.I. [0.44, 0.49]; Hit and FA: M= 0.24, 95% C.I. [0.22, 0.26]; $t(53)$ =10.63, $p < .001$), medial SFG (Hit and CR: M =0.12, 95% C.I. [0.12, 0.13]; Hit and FA: M= 0.08, 95% C.I. [0.08, 0.09]; $t(53)$ =7.06, $p < .001$), and MTG (Hit and CR: M= 0.13, 95% C.I. [0.12, 0.0.13]; Hit and FA: M= 0.09, 95% C.I. [0.08, 0.09]; $t(53)$ = 9.07, p=.03).

There was also an age by trial type interaction ($F(1, 52)$ = 9.15, p=.004), as well as a significant 3-way interaction ($F(1.44, 74.85)$ = 12.03, $p < .001$). The age by RSA type interaction showed that younger adults exhibited higher neural similarity across Hits and CRs (M= 0.24, SD = 0.20, 95% C.I. [0.22, 0.25]) than Hits and FAs (M= 0.12, SD= 0.07, 95% C.I. [0.11, 0.12]; $t(25)$ = 9.32, $p < .001$). Older adults similarly exhibited higher similarity across Hits and CRs (M= 0.21, SD= 0.17, 95% C.I. [0.19, 0.22]) than Hits and FAs (M= 0.14, SD= 0.12, 95% C.I. [0.13, 0.15]; $t(27)$ = 6.54, $p < .001$). Breaking down the foregoing 3-way interaction, results showed that the difference between the two RSA metrics was reduced in older adults compared to younger adults in the IOC (M_{OA}=0.12, M_{YA}=0.25), $t(52)$ = 3.41, $p < .005$, MOC (M_{OA}=0.16, M_{YA}=0.30), $t(52)$ = 3.62, $p < .001$, and medial SFG (M_{OA}=0.027, M_{YA}=0.052), $t(52)$ = 2.36, p = .022 (the difference in the medial SFG did not survive Bonferroni correction)[2]. [See figure 3].

Next, we ran a series of linear regressions, including Age, Hit-FA similarity, and the interaction term, to examine how age and Hit and FA similarity may predict false alarm rates. In line with the foregoing behavioral results, age was a significant predictor of false alarm rates in all ROIs such that older adults (M=0.29) had higher false alarms than younger adults (M=0.16), all p's<.001. In the MOC (b=0.36, p=.017), MTG (b=1.49, p=.003), and medial SFG (b=2.52,

---

[2] The RSA values broken down by age group are reported in Supplemental Table A for descriptive purposes.

$p<.001$), similarity was a significant predictor of FAs such that as similarity increases, FA rates

increase. The age by similarity interaction was not significant in any ROI[3].

A similar set of linear regressions using Hit-CR similarity in place of Hit-FA similarity,

were run within the ROIs to predict corrected recognition rates. Neither Hit-CR similarity nor the

interaction term was found to predict corrected recognition rates (all *p's*>.15). Mirroring the

effect of age in the FA regression above, age was a significant predictor of correct rejection rates

in all ROIs such that younger adults had higher correct rejections (*M*=0.45) compared to older

adults (*M*=0.26), all *p*'s<.001.

Discussion

The current study aimed to shed light on the neural processes underlying the role of false

memories within age-related associative memory deficits. Behaviorally, our results revealed the

observed associative memory deficit in *d'* was driven by greater false alarm rates in older adults

than younger adults, rather than by an age-related reduction in hit rates. Our ERS analysis reveal

a significant main effect of ROI and configural congruency condition, such that we observed an

advantage for representing and processing targets as a function of intact configural congruency

across memory phases within occipital regions. Yet, absent of age effects in these regions, the

results suggest that older adults, like young, exhibited greater correspondence of neural activity

between encoding and retrieval for targets that maintained the same configural context across

memory phases than targets that did not. With regard to the retrieval-related RSA, we saw an

age-related difference in the similarity of neural patterns underlying accurate and inaccurate

---

[3] Despite the lack of age by similarity interactions, we report correlations with behavior for each age group in Supplemental Table A for descriptive purposes. We also plot age group regression lines alongside group regression lines displayed in Figure 4 for illustrative purposes.

memory retrieval. Specifically, results revealed that pattern similarity between Hits and FAs and Hits and CRs differed as a function of age in occipital ROIs, such that older adults exhibited a smaller difference between the two similarity metrics than did younger adults. Additionally, pattern similarity between Hits and FAs correlated with FA rates across a number of regions in older adults. Behavioral and multivariate findings will be discussed in turn.

Behavioral Results. Consistent with a wealth of past research (Chalfonte & Johnson, 1996; Naveh-Benjamin, 2000; Naveh-Benjamin, Guez, Kilb, & Reedy, 2004; Old & Naveh-Benjamin, 2008a; Overman & Becker, 2009; Overman et al., 2018; Rhodes et al., 2008), older adults exhibited deficits in associative memory compared to younger adults. Specifically, older adults showed overall lower behavioral discriminability, measured by *d'* (see Figure 3). When breaking down *d'* and examining hit and false alarm rates, results showed that the foregoing age reduction in *d'* was driven not by age differences in hit rates, but by age-related increases in associative false alarms. Absent of any interactions with age, results suggest that the foregoing age differences were not affected by either encoding or retrieval conditions but reflect a general decline in the ability to identify novel information during retrieval. While age-related associative memory deficits are often conceptualized with respect to errors in target detection, the current finding is consistent with past research that also finds that errors in lure rejection (i.e., false memories) underlie age-related associative memory deficits (e.g., Castel & Craik, 2003; Healy et al., 2005; Naveh-Benjamin et al., 2003; Old & Naveh-Benjamin, 2008b; Overman & Stephens, 2013; Rhodes et al., 2008).

Moreover, this pattern of results is consistent with the fact that older adults tend to make memory responses based on a feeling of familiarity opposed to recalling specific details during

memory retrieval (e.g., Bastin & Van der Linden, 2006; Davidson & Glisky, 2002). In the present memory study, lure familiarity arises due to the fact that the lures were composed of recombined pairs from encoding, such that all *items* within a lure pair are 'old', yet not the specific association amongst the items. In order to identify a recombined lure as new, one must overcome the item familiarity and respond based on the novel combination of items (Cohn & Moscovitch, 2007; Lepage, Brodeur, & Bourgouin, 2003; Rotello & Heit, 2000). The current behavioral results suggest that there is an age-related decline in the ability to undertake this relatively difficult task of identifying the novel association as new. This is also consistent with an age-related reduction in strategic processing at retrieval (Cohn, Emrich, & Moscovitch, 2008), as well as with previous work showing that older adults have difficulty employing strategic processing at retrieval to reject similar but novel information (i.e., lack of recall-to-reject processing) (Bowman & Dennis, 2015; Pierce, Waring, Schacter, & Budson, 2008). Future research may look at different response profiles across age based on whether the option of responding 'rearranged' is offered (i.e., 'intact' vs. 'rearranged'), as is done in some associative memory tasks.

ERS analyses. To understand why older adults endorse associative lures as 'old' pairs at higher rates than younger adults, we first examined whether there were age differences in how encoded information related to targets was recapitulated during retrieval. Specifically, following earlier work with younger adults (Gerver et al., 2020) we examined the similarity of neural patterns of target pairs as a function of both the encoding configuration and configural congruency across memory phases. We hypothesized that, consistent with prior research in item and source memory (Chamberlain, Bowman, & Dennis, under review; Folville et al., 2020; Hill et al., 2021; Trelle et

al., 2019), older adults would exhibit reduced ERS of target pairs compared to younger adults.

We theorized that such a reduction in the transfer of associative information from encoding to

retrieval would contribute to the reduced distinctiveness of targets and lures within retrieval

itself. Our results showed that in older, like younger adults (Gerver et al., 2020), both configural

context and configural congruency were critical factors in ERS within the IOC and MOC (see

figure 2). Specifically, these findings suggest that when an associative target is presented in the

same configural context across both memory phases, it evokes greater recapitulation of visual

processes associated with the encoding episode than if the configural context is altered for the

item-item pair. This pattern of ERS across target conditions in older adults suggests that, like

younger adults (Gerver et al., 2020), older adults take advantage of stability in the configural

context across memory phases when reinstating memory representations associated with

associative pairs across memory phases. This finding aligns with the behavioral results showing

higher hit rates in the congruent compared to incongruent configural condition across age groups.

While we didn't find that that ERS correlated with behavioral metrics of associative memory

success or false memory rates, it is likely that this consistency for configurally-congruent pairs

supports identification of targets using both item-item details and general familiarity of the

associative configurations.

Noted above, our Bayes Factor analysis examining age differences within the two

occipital ROIs found that while there is moderate evidence supporting the null hypothesis (no

age differences) within the MOC, there is moderate evidence supporting the alternative

hypothesis (age differences) within the IOC. A closer look at the ERS values within IOC

suggests that this age difference may lie in the fact that, despite exhibiting the same pattern of

ERS across the four conditions, older adults exhibit somewhat lower overall ERS for the two

incongruent conditions compared to younger adults. Thus, the current results offer mixed evidence for an age-reduction in target reinstatement within the occipital cortex. Prior work in the field of memory and aging has observed age-related decline in the reinstatement of target processing, (Chamberlain et al., under review; Folville et al., 2020; Hill et al., 2021; Trelle et al., 2019) within occipital cortices. This prior memory research concludes that aging is associated with reduced reinstatement of neural activity associated with the encoding of detailed stimuli. Irrespective of this deficit, prior findings also suggest that this decrease in recapitulation does not affect the subjective vividness of these memories in older adults (Folville et al., 2020) and may be related to neural dedifferentiation during encoding (Hill et al., 2021). Given the complicated nature of age differences within the ERS results, future work should continue to examine how congruency across memory phases affects the reinstatement of associative information in aging.

One important point to be considered with respect to any group comparisons in ERS metrics is that ERS metrics do not account for the specificity of the information that is carried over from encoding to retrieval or whether the same type of processing is occurring within the same ROIs across age groups. They merely account for similarity in neural patterns within an individual across memory phases. Thus, it cannot be concluded that both age groups encode and maintain the same information from encoding to retrieval, simply that both age groups exhibit relatively similar patterns of encoding and retrieval activity across the two memory phases. Thus, it may be that younger adults encode and maintain more nuanced or detailed associative representations compared to older adults, whereas older adults process information in a more generic or gist-based manner (e.g., Pierce, Sullivan, Schacter, & Budson, 2005; Schacter, Koutstaal, & Norman, 1997; Stephens & Overman, 2018; Tun, Wingfield, Rosen, & Blanchard, 1998) and that we cannot capture these subtle differences in with the current ERS analysis

approach. With more advanced analyses, we will hopefully be able to assess the specificity of the memory representations across age groups.

RSA analyses. While the investigation of target-related patterns of activity indicates stability of the target representation across age groups, the investigation of age differences in pattern similarity related to false alarms during retrieval told a different story. Specifically, to investigate the neural processes underlying older adults' reduced ability to distinguish between targets and lures, we used pattern similarity analyses to determine how the relationship between hits and false memory processing differs from that of hits and correctly rejected lures. We also directly investigated whether this relationship was affected by age and whether neural similarity across targets and lures related to false memory rates within our participants. Results showed that while the neural pattern similarity underlying hits and correct rejections was greater than neural pattern similarity underlying hits and false alarms, this difference between the two similarity scores was reduced for older adults. Specifically, the difference between pattern similarity underlying Hits and CRs and Hits and FAs was reduced in older compared to younger adults (see Figure 3), with overall age differences driven by neural processing within both the IOC and MOC (see Figure 3).

Figure 3. RSA for Hits & False Alarms (FA) and Hits & Correct Rejections (CRs) in Inferior Occipital Cortex (IOC) and Middle Occipital Cortex (MOC) across younger adults (YAs) and older adults (OAs).

The finding that neural patterns underlying veridical memories, in any form, are more similar than those corresponding to a behavioral response of 'old', may be reflective of overlap that exists between target identification and recall-to-reject strategies in recognition memory (Brainerd, Reyna, Wright, & Mojardin, 2003; Gallo, Bell, Beier, & Schacter, 2006; Rotello & Heit, 2000). That is, in order to overcome the familiarity inherent in targets and recombined lures in associative memory retrieval, one needs to recall the specific correspondence between the items within the association and not merely the past occurrence of the individual items. The current results suggest that when this process of lure rejection is successful, individuals, irrespective of age, exhibit higher similarity between the identification of intact target pairs and the identification of the recombined novel pairs as lures than they do with novel lure pairs that are erroneously identified as old.

Critically, with respect to aging, the results also identified a reduction in the difference between the RSA metrics in the IOC and MOC in older compared to younger adults. Driven by

an increase in the neural overlap across Hits and FAs, within older adults, this suggests that the

two trial types may be less distinct in how they are processed in older adults, specifically within

occipital regions. This finding is consistent with a wealth of literature showing that older adults

exhibit dedifferentiation with respect to the similarity of neural processing across distinct trial

types (e.g., Deng et al., 2021; Koen, Hauck, & Rugg, 2019; Koen & Rugg, 2019; St-Laurent,

Abdi, Bondad, & Buchsbaum, 2014). While age-related dedifferentiation is usually reported in

the context of distinctiveness of object processing (e.g., face/scenes), our results extend this past

work, given the current need to differentiate between two fairly similar sets of objects,

specifically, intact and recombined pairs. This finding is also consistent with prior work in our

lab indicating that patterns of neural activation between targets and lures are less distinguishable

in visual cortices in aging (Bowman et al., 2019). In this past work, age deficits in memory were

related to reduced discriminability of cognitive processes (i.e., old/new recognition). To that

point, and in line with our predictions, the similarity in neural patterns of activity underlying Hits

and FAs predicted FA rates in our sample within several ROIs, including MOC, MTG, and

medal PFC (see Figure 4). While activity in the foregoing regions has been linked to false

memory processing across a large number of studies in both young and older adults (e.g.,

Dennis, Bowman, et al., 2014; Dennis, Kim, & Cabeza, 2008; Dennis & Turney, 2018; Duarte,

Graham, & Henson, 2010), the current finding supports the idea that similarity in neural

processing across Hits and FAs is likely to underlie the fact that lures are erroneously labeled as

"old".

Figure 4. Correlation between RSA for Hits & False Alarms (FA) with FA rates in each ROI collapsed across age (YAs: younger adults; OAs: older adults). Group regression lines are also included for illustrative purposes. Middle Occipital Cortex (MOC); Middle Temporal Gyrus (MTG); Medial Superior Frontal Gyrus (SFG).

This conclusion is consistent with a wealth of past research examining false memories and aging that points to perceptual relatedness, shared semantic gist, and overlapping mnemonic representations across targets and lures as underlying mechanisms accounts for age-related increases in false memory rates (e.g., Dennis, Bowman, et al., 2014; Devitt & Schacter, 2016; Tun et al., 1998). In accord with the factors described above, associative lures, much like related lures within semantic and perceptual false memory paradigms (e.g., DRM test, perceptual

relatedness studies; see Dennis, Bowman, & Turney, 2015), share a high degree of similarity with target items, as both items within the rearranged pair have been previously encountered. Thus, the lure incurs a high perceptual overlap with encoded information and is likely to evoke a similar perceptual and conceptual gist as to that of target items. In the absence of clear and distinct representations across trial types, such shared perceptual and gist-related representations are likely to lead to a similar lure being incorrectly labeled as "old". Regarding episodic representations, the MTG has been shown to be involved in processing the gist associated with episodes (Simons et al., 2005; Wise & Price, 2006) and MOC in processing more general properties of stimuli (Slotnick & Schacter, 2004, 2006; Vaidya, Zhao, Desmond, & Gabrieli, 2002). Given the correspondence in the relationship between Hit and FA representations and false memory rates in these regions, results suggest that the extent to which related lures invoke the general feeling of oldness associated with past veridical item-item associations, the more likely they will evoke an erroneous response of "old" at retrieval.

Yet the similarity of representations within these lower-level processing regions does not solely account for the creation of this memory error. The current results also suggest that similarities in the manner in which a Hit and FA are processed within higher level monitoring and evaluation regions, including both the medial SFG and parietal cortex, may contribute to increased false alarm rates in older adults. The correlation between Hit and FA RSA and false memories in the medial SFG is particularly interesting given that this region has consistently been identified in false memory studies across a wide variety of false memory paradigms (for a metanalysis see Kurkela & Dennis, 2016), including studies involving older adults (e.g., Dennis, Bowman, et al., 2014; Dennis & Turney, 2018; Webb & Dennis, 2019). Noted above, the role of the medial SFG during false memory retrieval has been linked to part of a frontal-parietal

cognitive control network responsible for the evaluation and monitoring of critical lures (Atkins & Reuter-Lorenz, 2011; Dennis, Bowman, & Vandekar, 2012; Dennis, Johnson, & Peterson, 2014; Garoff-Eaton, Kensinger, & Schacter, 2007; Hofer et al., 2007; Iidaka, Harada, Kawaguchi, & Sadato, 2012; Kensinger & Schacter, 2006; Slotnick & Schacter, 2004; von Zerssen, Mecklinger, Opitz, & von Cramon, 2001). Prior findings have established a link between processing within the medial prefrontal cortex and top-down retrieval monitoring in the absence of a strong sensory signal (Dennis, Bowman, et al., 2014) in the face of competing representations (Iidaka et al., 2012), and when there is a strong conceptual similarity between targets and lures (Garoff-Eaton et al., 2007). The current results extend this univariate work from younger adults to show not only that activity within this region support false associative memories in aging, but also that the greater the extent to which lure processing within the medial SFG corresponds to that underlying a veridical memory response, the greater the likelihood of a false memory.

General Conclusions

The current results shed light on the neural processes underlying age-related associative memory deficit. Behaviorally, our results highlight the role of false memories of recombined pairs at retrieval as a significant, contributing factor to age deficits in associative memory performance. That is, the associative memory deficit was driven by greater false alarm rates in older adults, rather than by a reduction in the hit rate. Paralleling the absence of age deficits in hit rates, we also found no age-related differences in the relationship between patterns of neural activity underlying targets across memory phase (as measured by ERS), though we did see an advantage for representing and processing targets as a function of intact configural congruency across

memory phases. Specifically, older adults, like young (Gerver et al., 2020), exhibited greater correspondence of neural activity between encoding and retrieval for targets that maintained the same configural context across memory phases than targets that did not. The results highlight the strength of configural arrangements as part of the associative memory trace and the importance of maintaining configural congruency across memory phases as a means to bolster the transfer of associative information across phases in older adults. This finding is consistent with earlier behavioral work from our collaborative group (Overman, Dennis, McCormick-Huhn, Steinsiek, & Cesar, 2019; Overman et al., 2018) where we found that both the manner of encoding presentation as well as congruency of configural display was critical to memory success. Importantly, the current results suggest that the strength of this representation is preserved across the adult lifespan.

At the same time, both age groups exhibited overall lower pattern similarity for hits and FAs when compared to hit and CRs. This representational difference between a consistent behavioral response (i.e., "old") and a correct memory decision suggests that there is more neural uniformity associated with veridical memory decisions across the associative memory network than for trials that share a similar behavioral response but differ with regard to memory veracity. Despite this overall difference, results also showed this difference to be reduced in older compared to younger adults, specifically withing occipital regions. Together with prior examining dedifferentiation and neural distinctiveness within false memory paradigms, this finding is indicative of an age-related increase in the similarity in processing across targets and lures that underlies lures being erroneously identified as targets in older adults. To this point, it was the overlap in neural pattern similarity across hits and FAs that correlated with false memory rates across several ROIs. This latter finding speaks directly to the idea that, when both young

and older adults experience a high correspondence in neural processing between targets and related lures, they are more apt to erroneously identify the lure as previously studied. Thus, the current results continue to underscore the importance of understanding errors of commission in aging in order to account for age-related memory decline.

Acknowledgements: We thank Kayla McGraw, M. Andrew Rowley, Joanna Salerno, Harini Babu, and Chloe Hultman for help with data collection and analyses. We also thank Jordan Chamberlain, Dan Elbich, and Ashley Steinkrauss for analysis support and help with earlier versions of this work. This work was supported by the National Institutes of Health under grant R15AG052903 awarded to A.A.O. and N.A.D. In addition, N.A.D. was also supported in part by National Science Foundation grants BCS1025709 and BCS2000047. Portions of the research in this article used the Color FERET (Facial Recognition Technology) database of facial images collected under the FERET program, sponsored by the Department of Defense Counterdrug Technology Development Program Office.

References

Andersson, J. L., Jenkinson, M., & Smith, S. (2007). *Non-linear registration, aka Spatial normalisation FMRIB technical report TR07JA2. 2*(1).

Atkins, A. S., & Reuter-Lorenz, P. A. (2011). Neural mechanisms of semantic interference and false recognition in short-term memory. *Neuroimage, 56*(3), 1726-1734.

Bastin, C., & Van der Linden, M. (2006). The effects of aging on the recognition of different types of associations. *Experimental Aging Research, 32*(1), 61-77.

Bowman, C. R., Ashby, S. R., & Zeithamova, D. (2021). Age effects on category learning, categorical perception, and generalization. *Memory*, 1-18. doi:10.1080/09658211.2021.2003818

Bowman, C. R., Chamberlain, J. D., & Dennis, N. A. (2019). Sensory Representations Supporting Memory Specificity: Age Effects on Behavioral and Neural Discriminability. *Journal of Neuroscience, 39*(12), 2265-2275. doi:10.1523/JNEUROSCI.2022-18.2019

Bowman, C. R., & Dennis, N. A. (2015). Age differences in the neural correlates of novelty processing: The effects of item-relatedness. *Brain Research, 1612*, 2-15. Retrieved from <Go to ISI>://WOS:000356551700002

Brainerd, C. J., Reyna, V. F., Wright, R., & Mojardin, A. H. (2003). Recollection rejection: False-memory editing in children and adults. *Psychological Review, 110*(4), 762-784. doi:10.1037/0033-295x.110.4.762

Bulevich, J. B., & Thomas, A. K. (2012). Retrieval effort improves memory and metamemory in the face of misinformation. *Journal of Memory and Language, 67*(1), 45-58. Retrieved from <Go to ISI>://WOS:000306673500004

Castel, A. D., & Craik, F. I. M. (2003). The Effects of Aging and Divided Attention on Memory for Item and Associative Information. *Psychology & Aging, 18*(4), 873-885.

Chalfonte, B. L., & Johnson, M. K. (1996). Feature memory and binding in young and older adults. *Memory & Cognition, 24*(4), 403-416.

Chamberlain, J. D., Bowman, C. R., & Dennis, N. A. (under review). Age-related differences in encoding-retrieval similarity and their relationship to false memory. *https://www.biorxiv.org/content/10.1101/2021.07.12.451838v1*.

Cohn, M., Emrich, S. M., & Moscovitch, M. (2008). Age-related deficits in associative memory: the influence of impaired strategic retrieval. *Psychol Aging, 23*(1), 93-103.

Cohn, M., & Moscovitch, M. (2007). Dissociating measures of associative memory: Evidence and theoretical implications. *Journal of Memory & Language, 57*(3), 437-454.

Davidson, P. S., & Glisky, E. L. (2002). Neuropsychological correlates of recollection and familiarity in normal aging. *Cognitive, Affective & Behavioral Neuroscience 2*(2), 174-186.

Deng, L., Davis, S. W., Monge, Z. A., Wing, E. A., Geib, B. R., Raghunandan, A., & Cabeza, R. (2021). Age-related dedifferentiation and hyperdifferentiation of perceptual and mnemonic representations. *Neurobiol Aging, 106*, 55-67. doi:10.1016/j.neurobiolaging.2021.05.021

Dennis, N. A., Bowman, C. R., & Peterson, K. P. (2014). Age-related differences in the neural correlates mediating false recollection. *Neurobiology of Aging, 35*, 395-407.

Dennis, N. A., Bowman, C. R., & Turney, I. C. (2015). Functional Neuroimaging of False Memories. In D. R. Addis, M. Barense, & A. Duarte (Eds.), *The Wiley Handbook on the Cognitive Neuroscience of Memory*. Chichester, UK: John Wiley & Sons, Ltd.

Dennis, N. A., Bowman, C. R., & Vandekar, S. N. (2012). True and phantom recollection: an

fMRI investigation of similar and distinct neural correlates and connectivity.

Neuroimage, 59(3), 2982-2993.

Dennis, N. A., Hayes, S. M., Prince, S. E., Huettel, S. A., Madden, D. J., & Cabeza, R. (2008).

Effects of aging on the neural correlates of successful item and source memory encoding.

Journal of Experimental Psychology: Learning, Memory and Cognition, 34(4), 791-808.

Dennis, N. A., Johnson, C. E., & Peterson, K. M. (2014). Neural correlates underlying true and

false associative memories. *Brain and Cognition, 88*, 65-72. Retrieved from <Go to

ISI>://000337777700009

Dennis, N. A., Kim, H., & Cabeza, R. (2008). Age-related differences in brain activity during

true and false memory retrieval. *Journal of Cognitive Neuroscience, 20*(8), 1390-1402.

Dennis, N. A., Overman, A. A., Gerver, C. R., McGraw, K. E., Rowley, M. A., & Salerno, J. M.

(2019). Different types of associative encoding evoke differential processing in both

younger and older adults: Evidence from univariate and multivariate analyses.

Neuropsychologia, 135, 107240. doi:10.1016/j.neuropsychologia.2019.107240

Dennis, N. A., & Turney, I. C. (2018). The influence of perceptual similarity and individual

differences on false memories in aging. *Neurobiol Aging, 62*, 221-230.

doi:10.1016/j.neurobiolaging.2017.10.020

Devitt, A. L., & Schacter, D. L. (2016). False memories with age: Neural and cognitive

underpinnings. *Neuropsychologia, 91*, 346-359. Retrieved from <Go to

ISI>://WOS:000387194200031

Duarte, A., Graham, K. S., & Henson, R. N. (2010). Age-related changes in neural activity

associated with familiarity, recollection and false recognition. *Neurobiology of Aging,*

31(10), 1814-1830. Retrieved from

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citati
on&list_uids=19004526

Folville, A., Bahri, M. A., Delhaye, E., Salmon, E., D'Argembeau, A., & Bastin, C. (2020). Age-related differences in the neural correlates of vivid remembering. *Neuroimage, 206*, 116336. doi:10.1016/j.neuroimage.2019.116336

Gallo, D. A., Bell, D. M., Beier, J. S., & Schacter, D. L. (2006). Two types of recollection-based monitoring in younger and older adults: Recall-to-reject and the distinctiveness heuristic. *Memory, 14*(6), 730-741. doi:10.1080/09658210600648506

Garoff-Eaton, R. J., Kensinger, E. A., & Schacter, D. L. (2007). The neural correlates of conceptual and perceptual false recognition. *Learning & Memory, 14*(10), 684-692.

Gerver, C. R., Overman, A. A., Babu, H. J., Hultman, C. E., & Dennis, N. A. (2020). Examining the Neural Basis of Congruent and Incongruent Configural Contexts during Associative Retrieval. *J Cogn Neurosci*, 1-17. doi:10.1162/jocn_a_01593

Giovanello, K. S., Kensinger, E. A., Wong, A. T., & Schacter, D. L. (2009). Age-related neural changes during memory conjunction errors. *Journal of Cognitive Neuroscience, 22*(7), 1348-1361.

Glanzer, M., & Adams, J. K. (1985). The Mirror Effect in Recognition Memory. *Memory & Cognition, 13*(1), 8-20. doi:Doi 10.3758/Bf03198438

Healy, M. R., Light, L. L., & Chung, C. (2005). Dual-process models of associative recognition in young and older adults: evidence from receiver operating characteristics. *J Exp Psychol Learn Mem Cogn, 31*(4), 768-788. doi:10.1037/0278-7393.31.4.768

Hill, P. F., King, D. R., & Rugg, M. D. (2021). Age Differences In Retrieval-Related Reinstatement Reflect Age-Related Dedifferentiation At Encoding. *Cereb Cortex, 31*(1), 106-122. doi:10.1093/cercor/bhaa210

Hockley, W. E. (2008). The effects of environmental context on recognition memory and claims of remembering. *J Exp Psychol Learn Mem Cogn, 34*(6), 1412-1429. doi:10.1037/a0013016

Hofer, A., Siedentopf, C. M., Ischebeck, A., Rettenbacher, M. A., Verius, M., Golaszewski, S. M., . . . Fleischhacker, W. W. (2007). Neural substrates for episodic encoding and recognition of unfamiliar faces. *Brain Cogn, 63*(2), 174-181. doi:10.1016/j.bandc.2006.11.005

Iidaka, T., Harada, T., Kawaguchi, J., & Sadato, N. (2012). Neuroanatomical substrates involved in true and false memories for face. *Neuroimage, 62*(1), 167-176. Retrieved from http://www.ncbi.nlm.nih.gov/pubmed/22575420

Jonker, T. R., Dimsdale-Zucker, H., Ritchey, M., Clarke, A., & Ranganath, C. (2018). Neural reactivation in parietal cortex enhances memory for episodically linked information. *Proc Natl Acad Sci U S A, 115*(43), 11084-11089. doi:10.1073/pnas.1800006115

Kassabara, A. (2020). rstatix: pipe-friendly framework for basic statistical tests. R package version 0.4. 0. Retrieved from https://cran. r-project. org/web/packages/rstatix/index.

Kassambara, A. (2021). rstatix: Pipe-Friendly Framework for Basic Statistical Tests (Version R package version 0.7.0). Retrieved from https://CRAN.R-project.org/package=rstatix

Kensinger, E. A., & Schacter, D. L. (2006). Amygdala activity is associated with the successful encoding of item, but not source, information for positive and negative stimuli. *Journal of Neuroscience, 26*(9), 2564-2570.

Koen, J. D., Hauck, N., & Rugg, M. D. (2019). The Relationship between Age, Neural

Differentiation, and Memory Performance. *J Neurosci, 39*(1), 149-162.

doi:10.1523/JNEUROSCI.1498-18.2018

Koen, J. D., & Rugg, M. D. (2019). Neural Dedifferentiation in the Aging Brain. *Trends Cogn

Sci, 23*(7), 547-559. doi:10.1016/j.tics.2019.04.012

Koutstaal, W. (2003). Older adults encode--but do not always use--perceptual details: intentional

versus unintentional effects of detail on memory judgments. *Psychol Sci, 14*(2), 189-193.

Koutstaal, W., & Schacter, D. L. (1997). Gist-based false recognition of pictures in older and

younger adults. *Journal of Memory & Language, 37*, 555-583.

Kriegeskorte, N., Mur, M., & Bandettini, P. (2008). Representational similarity analysis -

connecting the branches of systems neuroscience. *Front Syst Neurosci, 2*, 4.

doi:10.3389/neuro.06.004.2008

Kuhl, B. A., Rissman, J., Chun, M. M., & Wagner, A. D. (2011). Fidelity of neural reactivation

reveals competition between memories. *Proc Natl Acad Sci U S A, 108*(14), 5903-5908.

doi:10.1073/pnas.1016939108

Kurkela, K. A., & Dennis, N. A. (2016). Event-related fMRI studies of false memory: An

Activation Likelihood Estimation meta-analysis. *Neuropsychologia, 81*, 149-167.

doi:10.1016/j.neuropsychologia.2015.12.006

Lee, H., Samide, R., Richter, F. R., & Kuhl, B. A. (2019). Decomposing Parietal Memory

Reactivation to Predict Consequences of Remembering. *Cereb Cortex, 29*(8), 3305-3318.

doi:10.1093/cercor/bhy200

Lenth, R. V. (2021). emmeans: Estimated Marginal Means, aka Least-Squares Means.

Lepage, M., Brodeur, M., & Bourgouin, P. (2003). Prefrontal cortex contribution to associative

    recognition memory in humans: an event-related functional magnetic resonance imaging

    study. *Neuroscience Letters, 346*(1-2), 73-76.

Martinez, A. M., & Benavente, R. (1998). The AR face database. *CVC Technical Report #24*.

McDermott, K. B., Gilmore, A. W., Nelson, S. M., Watson, J. M., & Ojemann, J. G. (2017). The

    parietal memory network activates similarly for true and associative false recognition

    elicited via the DRM procedure. *Cortex, 87*, 96-107. doi:10.1016/j.cortex.2016.09.008

Minear, M., & Park, D. C. (2004). A lifespan database of adult facial stimuli. *Behavior Research

    Methods, Instruments, & Computers, 36*(4), 630-633.

Mitchell, K. J., Ankudowich, E., Durbin, K. A., Greene, E. J., & Johnson, M. K. (2013). Age-

    related differences in agenda-driven monitoring of format and task information.

    Neuropsychologia, 51(12), 2427-2441.

Mitchell, K. J., Raye, C. L., Johnson, M. K., & Greene, E. J. (2006). An fMRI investigation of

    short-term source memory in young and older adults. *Neuroimage, 30*(2), 627-633.

Morey, R. D., Rouder, J. N., Jamil, T., & Morey, M. R. D. P. b. (2015). Package 'bayesfactor'.

Mumford, J. A., Davis, T., & Poldrack, R. A. (2014). The impact of study design on pattern

    estimation for single-trial multivariate pattern analysis. *Neuroimage, 103*, 130-138.

    doi:10.1016/j.neuroimage.2014.09.026

Naveh-Benjamin, M. (2000). Adult age differences in memory performance: tests of an

    associative deficit hypothesis. *Journal of Experimental Psychology: Learning, Memory

    and Cognition, 26*(5), 1170-1187.

Naveh-Benjamin, M., Guez, J., Kilb, A., & Reedy, S. (2004). The associative memory deficit of older adults: further support using face-name associations. *Psychol Aging, 19*(3), 541-546.

Naveh-Benjamin, M., Hussain, Z., Guez, J., & Bar-On, M. (2003). Adult age differences in episodic memory: Further support for an associative-deficit hypothesis. *Journal of Experimental Psychology: Learning, Memory, & Cognition, 29*(5), 826-837.

Norman, K. A., & Schacter, D. L. (1997). False recognition in younger and older adults: exploring the characteristics of illusory memories. *Memory & Cognition, 25*(6), 838-848.

Old, S. R., & Naveh-Benjamin, M. (2008a). Differential effects of age on item and associative measures of memory: a meta-analysis. *Psychol Aging, 23*(1), 104-118.

Old, S. R., & Naveh-Benjamin, M. (2008b). Memory for people and their actions: further evidence for an age-related associative deficit. *Psychol Aging, 23*(2), 467-472. doi:10.1037/0882-7974.23.2.467

Overman, A. A., & Becker, J. T. (2009). The Associative Deficit in Older Adult Memory: Recognition of Pairs Is Not Improved by Repetition. *Psychology and aging, 24*(2), 501-506. doi:Doi 10.1037/A0015086

Overman, A. A., Dennis, N. A., McCormick-Huhn, J. M., Steinsiek, A. B., & Cesar, L. B. (2019). Same face, same place, different memory: manner of presentation modulates the associative deficit in older adults. *Aging, Neuropscyhology, and Cognition, 26*(1), 44-57. doi:10.1080/13825585.2017.1397097

Overman, A. A., McCormick-Huhn, J. M., Dennis, N. A., Salerno, J. M., & Giglio, A. P. (2018). Older adults' associative memory is modified by manner of presentation at encoding and retrieval. *Psychol Aging, 33*(1), 82-92. doi:10.1037/pag0000215

Overman, A. A., & Stephens, J. D. W. (2013). Synergistic Effects of Encoding Strategy and Context Salience on Associative Memory in Older Adults. *Psychology and Aging, 28*(3), 654-665. doi:Doi 10.1037/A0031441

Phillips, P. J., Moon, H., Rizvi, S. A., & Rauss, P. J. (2000). The FERET evaluation methodology for face-recognition algorithms. *Ieee Transactions on Pattern Analysis and Machine Intelligence, 22*(10), 1090-1104. doi:Doi 10.1109/34.879790

Pierce, B. H., Sullivan, A. L., Schacter, D. L., & Budson, A. E. (2005). Comparing source-based and gist-based false recognition in aging and Alzheimer's disease. *Neuropsychology, 19*(4), 411-419. doi:10.1037/0894-4105.19.4.411

Pierce, B. H., Waring, J. D., Schacter, D. L., & Budson, A. E. (2008). Effects of distinctive encoding on source-based false recognition: further examination of recall-to-reject processes in aging and Alzheimer disease. *Cognitive and behavioral neurology : official journal of the Society for Behavioral and Cognitive Neurology, 21*(3), 179-186. doi:10.1097/WNN.0b013e31817d74e7

Rhodes, M. G., Castel, A. D., & Jacoby, L. L. (2008). Associative recognition of face pairs by younger and older adults: the role of familiarity-based processing. *Psychol Aging, 23*(2), 239-249. doi:10.1037/0882-7974.23.2.239

Ritchey, M., Wing, E. A., LaBar, K. S., & Cabeza, R. (2013). Neural similarity between encoding and retrieval is related to memory via hippocampal interactions. *Cereb Cortex, 23*(12), 2818-2828. doi:10.1093/cercor/bhs258

Rotello, C. M., & Heit, E. (2000). Associative recognition: a case of recall-to-reject processing. *Mem Cognit, 28*(6), 907-922. doi:10.3758/bf03209339

Schacter, D. L., Buckner, R. L., Koutstaal, W., Dale, A. M., & Rosen, B. R. (1997). Late onset of anterior prefrontal activity during true and false recognition: an event-related fMRI study. *Neuroimage, 6*(4), 259-269.

Schacter, D. L., Koutstaal, W., & Norman, K. A. (1997). False memories and aging. *Trends in cognitive sciences, 1*(6), 229-236. doi:10.1016/S1364-6613(97)01068-1

Simons, J. S., Verfaellie, M., Hodges, J. R., Lee, A. C., Graham, K. S., Koutstaal, W., . . . Budson, A. E. (2005). Failing to get the gist: reduced false recognition of semantic associates in semantic dementia. *Neuropsychology, 19*(3), 353-361.

Slotnick, S. D., & Schacter, D. L. (2004). A sensory signature that distinguishes true from false memories. *Nature Neuroscience, 7*(6), 664-672.

Slotnick, S. D., & Schacter, D. L. (2006). The nature of memory related activity in early visual areas. *Neuropsychologia, 44*(14), 2874-2886.

Smith, A. (2002). Finding maps in the cortex. *Trends Cogn Sci, 6*(3), 114.

Solina, F., Peer, P., Batageli, B., Juvan, S., & Kovac, J. (2003, March 10-11). *Color-based face detection in the "15 seconds of fame" art installation.* Paper presented at the Conference on Computer Vision / Computer Graphics Collaboration for Model-based Imaging, Rendering, Image Analysis and Graphical Special Effects, INRIA Rocquencourt, France.

Sperling, R. A., Bates, J. F., Chua, E. F., Cocchiarella, A. J., Rentz, D. M., Rosen, B. R., . . . Albert, M. S. (2003). fMRI studies of associative encoding in young and elderly controls and mild Alzheimer's disease. *Journal of Neurol Neurosurg Psychiatry, 74*(1), 44-50.

Srokova, S., Hill, P. F., Elward, R. L., & Rugg, M. D. (2021). Effects of age on goal-dependent modulation of episodic memory retrieval. *Neurobiol Aging, 102*, 73-88. doi:10.1016/j.neurobiolaging.2021.02.004

St-Laurent, M., Abdi, H., Bondad, A., & Buchsbaum, B. R. (2014). Memory reactivation in healthy aging: evidence of stimulus-specific dedifferentiation. *J Neurosci, 34*(12), 4175-4186. doi:10.1523/JNEUROSCI.3054-13.2014

Stephens, J. D. W., & Overman, A. A. (2018). Modeling age differences in effects of pair repetition and proactive interference using a single parameter. *Psychol Aging, 33*(1), 182-194. doi:10.1037/pag0000195

Tompary, A., Duncan, K., & Davachi, L. (2016). High-resolution investigation of memory-specific reinstatement in the hippocampus and perirhinal cortex. *Hippocampus, 26*(8), 995-1007. doi:10.1002/hipo.22582

Trelle, A. N., Henson, R. N., & Simons, J. S. (2019). Neural evidence for age-related differences in representational quality and strategic retrieval processes. *Neurobiol Aging, 84*, 50-60. doi:10.1016/j.neurobiolaging.2019.07.012

Tun, P. A., Wingfield, A., Rosen, M. J., & Blanchard, L. (1998). Response latencies for false memories: gist-based processes in normal aging. *Psychology and Aging, 13*(2), 230-241.

Vaidya, C. J., Zhao, M., Desmond, J. E., & Gabrieli, J. D. (2002). Evidence for cortical encoding specificity in episodic memory: memory-induced re-activation of picture processing areas. *Neuropsychologia, 40*(12), 2136-2143.

von Zerssen, G. C., Mecklinger, A., Opitz, B., & von Cramon, D. Y. (2001). Conscious recollection and illusory recognition: an event-related fMRI study. *The European Journal of Neuroscience, 13*(11), 2148-2156.

Webb, C. E., & Dennis, N. A. (2019). Differentiating True and False Schematic Memories in Older Adults. *J Gerontol B Psychol Sci Soc Sci, 74*(7), 1111-1120. doi:10.1093/geronb/gby011

Wing, E. A., Geib, B. R., Wang, W. C., Monge, Z., Davis, S. W., & Cabeza, R. (2020). Cortical
Overlap and Cortical-Hippocampal Interactions Predict Subsequent True and False
Memory. *J Neurosci, 40*(9), 1920-1930. doi:10.1523/JNEUROSCI.1766-19.2020

Wing, E. A., Ritchey, M., & Cabeza, R. (2015). Reinstatement of individual past events revealed
by the similarity of distributed activation patterns during encoding and retrieval. *J Cogn
Neurosci, 27*(4), 679-691. doi:10.1162/jocn_a_00740

Wise, R. J. S., & Price, C. J. (2006). Functional imaging of language. In R. Cabeza & A.
Kingstone (Eds.), *Handbook of functional neuroimaging of cognition (2nd ed)* (pp. 191-
228). Cambridge, MA: MIT Press.