1

Svep1 orchestrates distal airway patterning and alveolar differentiation in murine lung
development

Foxworth N[1*], Wells J[2*], Ocaña-Lopez S[1], Muller S[1], Bhayani P[1], Denegre J[2], Palmer K[2],
Memishian W[2], McGee T[2], Murray SA[2], Donahoe PK[1,3,+], Bult CJ[2,+], and Loscertales M[1,3,+]

[1]Pediatric Surgical Research Laboratories, Massachusetts General Hospital, Boston, MA 02114

[2]The Jackson Laboratory, Bar Harbor, ME 04609

[3]Department of Surgery, Harvard Medical School, Boston, MA 02114

[*]Contributed equally

[+]Co-senior authors

Correspondence: Maria Loscertales, mloscertales@mgh.harvard.edu (ORCID 0000-0002-7585-
1235); Carol J. Bult, carol.bult@jax.org (ORCID 0000-0001-9433-210X)

Abstract

Disruptions in airway branching or alveolar differentiation during lung development can lead to
severe respiratory deficiencies and neonatal death. The molecular mechanisms governing
branching patterning and early alveolar formation remain elusive. Loss of *Svep1* function in
mice results in various developmental defects, including lung hypoplasia and perinatal lethality.
Our examination of the lungs of *Svep1* knockout (*Svep1$^{-/-}$*) mouse embryos, both *in vivo* and *in
vitro*, revealed that *Svep1* mutants exhibit an increase in the number of disorganized distal
airway tips and progressively greater disruption of lung lobe morphology over time and saccular
development. *Svep1* interacts with FGF signaling to regulate smooth muscle differentiation and,
together with *Fgf9,* guides airway branching patterning. Transcriptomic data from the lungs of
Svep1$^{-/-}$ embryos revealed dysregulated gene expression affecting saccular maturation. Our
findings demonstrate that *Svep1* is a key extracellular matrix player shaping airway morphology
and influencing alveolar fate. These insights offer potential avenues for therapeutic interventions
in congenital lung disorders.

Introduction

The architectural blueprint of the lungs emerges from two fundamental developmental

processes: branching morphogenesis, responsible for constructing the intricate airway tree

structure, and differentiation, which generates specialized cells in distinct lung regions,

predominantly after the formation of the tracheobronchial tree [1]. The cessation of molecular

genetic signaling promoting branching is a prerequisite for distal alveolar cell differentiation [2, 3].

In mice, lung development begins at embryonic day 9.5 (E9.5) [4, 5] when two primary buds

emerge from the ventral foregut. During the pseudoglandular stage (E12.5 – E16.5), epithelial

progenitors at the bud tips undergo an iterative process of budding and bifurcation into the

surrounding mesenchyme and this is followed by expansion of the diameter and length of the

respiratory tree during the canalicular stage (E16.5-E17.5). During the saccular stage (E17.5-

P5), the vasculature develops along the airway and terminal airway tips widen. The alveolar

epithelium subsequently differentiates into alveolar type 2 (AT2) cells, responsible for surfactant

protein C (SFPC) production, and the thin, gas-exchanging alveolar type 1 (AT1) cells. Primitive

septa arise at this stage. Secondary septation and alveolar maturation occur during the

postnatal alveolar stage (P5-P30) [5].

The geometry of the airway tree is generated by three local branching modes: domain, planar,

and orthogonal [6]. During domain branching, new branches grow perpendicularly around the

circumference of the parent branch and generate the initial scaffold of each lobe and the overall

shape of the lung. Planar bifurcation occurs at the tips of the airway epithelium and generates

branches in the same plane, resulting in the thin edges of the lung lobes. Orthogonal bifurcation

generates branches perpendicular to prior branches, thus filling in the interior of the lung.

Trifurcations are rare and appear principally at the lung surface [7, 8]. As branching proceeds,

proximal-distal patterning is established, and cellular specification follows [3, 9, 10, 11]. Lung bud tip

progenitors persist and proliferate and subsequently undergo differentiation into alveolar type 1

(AT1) and alveolar type 2 (AT2) cells [3]. The cells of the primal airway stalk differentiate into

multiple proximal cell types [12, 13, 14, 15].

Studies in laboratory mice have shown that branching morphogenesis requires mesenchymal-

epithelial interactions, and various signaling pathways, particularly the FGF pathway, play

important roles in branching [16]. Fibroblast growth factor 10 (FGF10) is secreted by

mesenchymal cells and acts via the *FGFR2* receptor to maintain a permissive environment for

branching morphogenesis. FGF10 maintains distal epithelial SOX9$^+$ progenitor cells in an

undifferentiated state. These cells play a pivotal role in subsequent alveolar and mesenchymal

differentiation [17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28]. FGF10$^+$ cells give rise to bronchial airway smooth

muscle cells during early branching [29, 30, 31, 32]. FGF9 is also required for lung development and

plays different roles depending on its spatial and temporal expression. In the mesothelium,

FGF9 plays a primary role in mesenchymal growth; in the epithelium, FGF9 controls epithelial

branching [33, 34, 35, 36]. The extracellular matrix (ECM) also plays important roles in epithelial

branching and alveolarization [37, 38, 39, 40, 41, 42] and contributes to organ shape [43].

Svep1 (Sushi, Von Willebrand Factor Type A, EGF and pentratix domain-containing 1), also

known as Polydom, encodes an ECM protein that is ubiquitously expressed in the tissues of

adult mice and plays roles in cell adhesion [44], epidermal differentiation [45], and lymphatic vessel

formation and remodeling [46, 47]. *Svep1* homozygous mutant mice die at birth with edema and

body abnormalities [45, 47, 48, 49]. We investigated the role of *Svep1* in the development of lungs of

mice homozygous for the *Svep1*$^{tm1b(EUCOMM)Hmg}$ allele (hereafter, *Svep1*$^{-/-}$) generated by the

Knockout Mouse Phenotyping Program (KOMP2) program [49]. We observed that the loss of

Svep1 disrupts the basic lung branching program resulting in irregular lung lobe shape due to

an aberrant increase in distal airway tips. Smooth muscle differentiation is also disrupted in the

lungs of *Svep1*$^{-/-}$ embryos. Through the treatment of lung explant cultures with exogenous

SVEP1 protein, we demonstrated that *Svep1* functions as an inhibitor of branching

morphogenesis. Furthermore, we demonstrated that inactivation of FGF signaling recovers

smooth muscle differentiation in *Svep1*$^{-/-}$ lungs and that exogeneous FGF9 treatment restores

SVEP1 branching inhibition, implying a potential role for SVEP1-FGF9 signaling specific to

branching.

Results

Svep1 is expressed in lung mesenchyme with high expression at the lung periphery

Svep1 is expressed in the lung mesenchyme, especially at the tip and edges of the lung (Fig.

1a-*d* and Suppl. Fig. 1a-b). *Svep*1 mRNA expression was strong in the mesenchyme adjacent

to the distal epithelial airways and around bifurcating tips (Fig.1a and Suppl. Fig.1a). We

detected SVEP1 protein expression on the epithelium membrane (ECAD+) of very distal lung

buds (Fig.1b and Suppl. Fig. 1b) and microvasculature (Suppl. Fig.1c). At the saccular stage

(E18.5), transcript and protein expression of *Svep1* was detected in the primary septa of lung

parenchyma and in the mesenchyme next to the proximal epithelium (Fig.1d and Suppl. Fig.1d).

Consistent with previous findings, *Svep1* expression during lung development was high

between E14.5 and E18.5 [50] (Suppl. Fig.1e). Single cell RNA-seq data demonstrated that

expression of *Svep1* in the lungs of E18.5 wild type embryos was prominent in matrix fibroblasts

(Suppl. Fig. 3a).

Hypoplastic and aberrant lung shape in Svep1$^{-/-}$ embryos

Consistent with previous reports [49], *Svep1*$^{-/-}$ embryos displayed perinatal lethality, whole-body

edema, and lung hypoplasia that was prominent at E18.5 by microCT (Fig. 1e and Suppl. Fig.

1f). *In situ* hybridization confirmed the absence of *Svep1* expression in the lungs of homozygous

knockout animals (Suppl. Fig. 1g). Examination of gross examination of embryos at E18.5

revealed small lungs with lobes that failed to enclose the heart and had a highly lobulated,

cauliflower-like appearance (Fig. 1f). Histological analysis at E16.5 showed small lungs in

mutants with irregular shape; at E18.5, small saccules were observed along a dorsal-ventral

gradient, accompanied by a reduced lumen primarily on the ventral side (Fig. 1g,h). Noticeable

differences, including a thickening along the lobulated ventral edges of the lung composed of

mesothelium, were observed in *Svep1*$^{-/-}$ embryos compared to wild type embryos reflecting

developmental abnormalities in the lung mesothelium of mutant mice (Fig. 1h).

Saccular development defects in Svep1$^{-/-}$ embryos

Sacculation begins in the murine lung shortly after E16.5, with the flattening of AT1 cells at the

proximal-to-distal opening of the distal airway lumen. Cells in the developing distal airways at

the lung edge are the last to differentiate [11, 30, 51]. Examination of distal airway morphology of

Svep1$^{-/-}$ embryos at E18.5, through labeling the lung epithelium with ECAD and the distal

progenitor marker SRY-box transcription factor 9 (SOX9), revealed that the densely packed

distal airways at the ventral edge exhibited a morphology typical of the earlier pseudoglandular

stage instead of the saccular stage (Fig. 1h). Quantitative real-time PCR (qPCR) confirmed

elevated expression of *Sox9* in the lungs of *Svep1* knockout embryos at E18.5 but not at earlier

time points (Fig. 1i). Co-staining of SOX9 with the cell proliferation market KI67 showed

proliferating epithelial progenitor cells in E18.5 lungs, mainly at the lung periphery in *Svep1*$^{-/-}$

mutants (Suppl. Fig. 1h). Our results indicate that the distal epithelium in *Svep1* lungs is in an

earlier developmental stage, characterized by a higher proportion of epithelial cells in a

progenitor stage, rather than in the saccular period when cells are differentiating into specialized

lung epithelial type I and II cells. Our finding highlights the essential role of Svep1 in the

maturation of the distal lung epithelium.

Lungs of Svep1^{-/-} embryos show an aberrant increase of distal airways altering lung lobe shape

To determine whether *Svep1^{-/-}* embryos have branching defects, we stained E16.5 lungs with

SOX9 and ECAD which revealed densely packed distal airways in *Svep1^{-/-}* embryos. Mutant

lobe tips had multiple distal branches and rosettes on the ventral lobe side (Fig. 2a). To further

examine defects in the branching program modes[6] (Fig.2b) in *Svep1* mutant lungs, we stained

whole lung and individual lobes with SOX9 at E13.5, E14.5, and E15.5 and showed that *Svep1*

branching defects accumulate over time (Fig. 2c-e and Suppl. Fig. 2a,b). We observed

branching and lobe shape anomalies in *Svep1^{-/-}* lungs as early as E13.5, primarily at the lung

tips and edges (Fig. 2c,d). Moreover, the branching abnormalities observed in the mutants

were not uniform across the lung lobes, leading to different outcomes in the shape of the lobes.

At E13.5 for example, the shape of the Right Caudal (RCa) lobe was narrower in *Svep1^{-/-}*

embryos compared to wild-type embryos. The Right Accessory (RAc) and Right Medial (RMe)

lobes of *Svep1* mutant embryos terminated in two or more buds instead of a single tip typically

observed in lungs of wild type embryos (Fig. 2c). Further examination of the left lobes showed

that *Svep1^{-/-}* embryos at E13.5 exhibited peripheral branch mis-localization and additional

budding, disrupting the normal planar bifurcation of the lung (Fig. 2d and Suppl. Fig. 2a). When

planar bifurcation is disrupted at the lobe tip, ectopic branching angles occur at the lung tips in

mutants (Fig. 2d). This observation suggests that *Svep1* deficiency impairs lung branching

angles early in development.

The orthogonal bifurcation mode of the branching program fills the lung's exterior and interior

surfaces in a specific pattern where each new bud maintains its boundaries with others and

creates the surface of the lung lobes [6]. However, this orthogonal pattern was disrupted in

mutant embryos at E14.5, with the airways rearranged into a "row of beads" pattern, resulting in

abnormal rounded tips (Fig. 2e). Moreover, the left lung lobes of E14.5 mutant embryos were

8% more elongated than those of wild type *Svep1* embryos (length/width ratio (mm) *Svep1^{+/+}*

=1.60; *Svep1^{-/-}* =1.90 *p <0.001 t test;* Suppl. Fig. 2c). Tissue cleared left lobes stained with

SOX9 at E14.5 confirmed abnormal mutant left lobe morphology as well as multi budding of the

distal airways and peripheral trifurcations (Fig. 2f).

At E15.5, aberrations persist in lung branching patterns along the margins and ventral side,

leading to substantial alterations in lobe elongation in *Svep1^{-/-}* embryos (Suppl. Fig. 2b,c,d).

Staining left lung lobes of *Svep1^{-/-}* embryos at E15.5 with DAPI (4',6-diamidino-2-phenylindole)

revealed prominent abnormalities on the lung surface (Suppl. Fig. 2e).

Quantification of SOX9+ in a whole lung Z-stack imaging revealed a significant increase in both
the total number of SOX9+ distal tips and the ratio of tips to area by E14.5 (Fig. 2g; $p < 0.05$ t-
test). Overall, these findings demonstrate that *Svep1* is crucial for normal branching
morphogenesis, and its absence leads to disorganization of airway patterning, primarily
affecting the tips, edges, and ventral areas of the lung, which in turn alters the overall shape of
the lung lobes.

Lungs of Svep1$^{-/-}$ embryos have ectopic branching, decreased planar bifurcations, and
increased trifurcations

To investigate which of the main branching programs are affected in *Svep1* mutants, we
modeled the branching patterns of normal and mutant lungs. We labeled tissue-cleared whole
lungs from E14.5 embryos with ECAD and SOX9 and generated a 3D reconstruction composed
of Z-stack images. We observed significant structural differences in the airways of wild type and
Svep1$^{-/-}$ lungs, confirming the disorder of the developing airway tree as well as the aberrant
duplication of the distal airway tips in *Svep1$^{-/-}$* embryos (Fig. 3a).

To quantify the differences in branching morphology in the lungs of mutant vs wild type
embryos, we generated a branching lineage diagram for the right accessory (RAc) lobe at
E14.5. Each branch was characterized as one of three branching modes: domain, planar, or
orthogonal [6]. Branches and direction of growth were mapped to create a lineage diagram in
which trifurcations were also indicated (Fig. 3b). Orthogonal branching was significantly
increased in mutants, contributing 63% of all branches compared to 49% ($p < 0.01$, t-test) in the
lungs of *Svep1$^{+/+}$* embryos (Fig. 3c). In contrast, the planar branching in the RAc lobes was
significantly decreased in mutant (5%) compared to wild type (13%) embryos ($p < 0.05$, t-test;
Fig. 3c). Trifurcations were more prevalent in *Svep1$^{-/-}$* lungs (10% of all branches) compared to
the lungs of *Svep1$^{+/+}$* embryos (4% of all branches) ($p < 0.05$ t-test; Fig. 3c and Fig.3e). An
average of 17% of first-level domain branches originating from the primary bronchus were
ectopic in *Svep1$^{-/-}$* lungs, while no ectopic branching was observed in the lungs of wild type
embryos (Fig. 3d and Fig. 3f). In RAc lobes of wild-type lungs, daughter branches of the primary
bronchus sprouted from one of only three directions: anterior, ventral-posterior, and dorsal-
posterior. In all the *Svep1$^{-/-}$* lungs analyzed, daughter branches sprouted from the primary
bronchus in random directions (Fig. 3f). In summary, loss of *Svep1* disrupts airway patterning
through decreased planar bifurcations, increased trifurcations, and ectopic domain branching
that arise in random directions.

To determine if the ectopic and trifurcated branches in $Svep1^{-/-}$ lungs was the result of loss of

Svep1, we analyzed time-lapse images of E12.5 lung explants for 48 hrs. (Fig. 3g, Video1,

Video2, Video 3). We observed ectopic branching arising from the main and secondary

branches and trifurcations at the lung tips. Taken together, these results suggest that *Svep1*

controls the position and orientation of new airway epithelial buds, defines distal bifurcations

regulating the form and structure of the growing epithelium, and is, therefore, critical for

determining the bronchial tree architecture and, ultimately, the final shape of the lobes of the

lung.

SVEP1 recombinant protein inhibits airway branching in normal E12.5 lung explants and

restores normal branching in Svep1$^{-/-}$ lungs.

The increased branching observed in the distal airways of $Svep1^{-/-}$ embryos suggests that

SVEP1 normally acts as an inhibitor of lung branching morphogenesis and is essential for

regulating lung airway budding. To further investigate the function of *Svep1* in branching

morphogenesis, we treated lung explants of wild type E11.5 -E12.5 embryos for 48hrs with a

recombinant SVEP1 (rSVEP1) protein. Time-lapse images revealed significantly decreased

branching in the treated lung explants (Fig. 4a; Video 4 and Video 5), quantified at the

peripheral airway tips (Fig. 4b). SVEP1 branching inhibition was more profound at earlier stages

(11.5) (data not shown). When lung explants from E12.5 $Svep1^{-/-}$ embryos were treated with

rSVEP, distal ectopic and trifurcated branches were reduced, and observed in only one of six

explants (Fig.4c). The *other Svep1$^{-/-}$* explants exhibited distal bifurcation that resembled wild

type lungs (Fig.4c).,

The loss of Svep1 leads to defects in smooth muscle and bronchial airway development

To characterize further the lungs of $Svep1^{-/-}$ embryos at the saccular stage, we conducted RNA

sequencing (RNA-seq) of whole lungs of $Svep1^{+/+}$ and $Svep1^{-/-}$ embryos. Term enrichment

analysis of the functional annotations for the differentially expressed genes revealed that genes

associated with cell mobility and adhesion, epithelial proliferation, extracellular matrix

organization, and collagen binding were significantly downregulated in $Svep1^{-/-}$ lungs (Fig. 5a

and Fig. 5b).

The RNA-seq data from the whole RCa lobe at E18.5 revealed that genes associated with

muscle development were significantly downregulated in $Svep1^{-/-}$ lungs, including *Acta2* (actin

alpha 2) which is necessary for smooth muscle differentiation, and *Cnn1* (calponin1), a marker

for mature smooth muscle (Fig. 5b). Lungs labeled with ACTA2 and particularly CNN1

confirmed the reduction in distal smooth muscle differentiation in the mutant at E18.5 (Fig. 5c)

evident around the proximal airways and adjacent blood vessels, particularly CNN1 expression

(Fig. 5c). The expression of ACTA2 was noticeably reduced also in lobular bronchioles in *ex*

vivo organ culture of E12.5 *Svep1*$^{-/-}$ lung explants that were allowed to grow in vitro for six days

(Fig. 5e) confirming that *Svep1* is necessary for proper peribronchial smooth muscle

differentiation which is defective in the lungs of *Svep1* mutant embryos before saccular

development.

To investigate lung airway development in *Svep1*$^{-/-}$ mutants further, we examined cellular

differentiation of the proximal epithelium by staining with the proximal epithelium progenitor

SRY-box 2 (SOX2) marker at the end of branching morphogenesis (E16.5)(Fig. 5f) and with

both club cell secretoglobin 1a1 (SCGB1A1) and ciliated cell forkhead box protein J1 (FOXJ1)

markers at E18.5 in the proximal airway epithelium (Fig. 5g). In mutants, SOX2 protein was

predominantly localized in the lung core (Fig. 5f), suggesting that the airway epithelium follows a

normal developmental pattern from proximal to distal at E16.5, with most proximal epithelium

undergoing typical development. SCGB1A1 staining revealed that *Svep1*$^{-/-}$ deficient embryos

displayed smaller distal bronchioles, accompanied by disorganized and densely packed

SCGBA1+ epithelial cells, leading to an increase in SCGBA1+ cells (Fig. 5g). In contrast, the

number of FOXJ1+ cells in mutants did not appear to differ from those in wild type lungs (Fig.

5g). Although *Scgba1* mRNA expression did not significantly differ between mutant and wild-

type animals, likely due to normal epithelial development in the most proximal airways, the

single cell RNA sequencing data indicated that the club cell cluster was enriched in *Svep1*$^{-/-}$

lungs (Suppl Fig. 3b). These finding highlights that normal airway structure relies on functional

Svep1 as the absence of *Svep1* leads to anomalies in bronchial epithelium morphology and an

increased number of club cells.

Altered FGF signaling is a contributing factor to the abnormalities in smooth muscle

development observed in Svep1$^{-/-}$ *lungs.*

RNA-seq data from whole lungs revealed that *Fgfr2* expression was significantly increased in

Svep1$^{-/-}$ embryos at E18.5. *Fgfr2* is expressed in the epithelium of the normal lung with

decreased expression after E17.5 [50]. In *Svep1*$^{-/-}$ embryos, however, *in situ* hybridization showed

that levels of expression of *Fgfr2* transcripts persisted in the lungs of mutants, even at E18.5,

compared to levels in lung of *Svep1*$^{+/+}$ embryos (Fig. 6a). Results from RT-qPCR showed a

significant increase in RNA expression levels of *Fgfr2* and other members of the FGF signaling

pathway, including *Fgf10*, *Fgf9*, and *Kras,* in the lungs of *Svep1* mutants at E18.5 (Fig. 6b) but

not at E14.5 or E16.5 (data not shown). *Fgfr2* has two isoforms, *Fgfr2b* and *Fgfr2c*. Only *Fgfr2b*

was significant at E18.5 (Fig. 6b). Furthermore, *Svep1$^{-/-}$* mutants showed no change in the

expression levels of important genes involved in lung development, such as *Tgfb1*, *Bmp*4, and

Shh at E14.5, E16.5, and E18.5 (Data not shown).

FGF10/ FGFR2b signaling is necessary for airway outgrowth and promotes epithelial progenitor

proliferation and migration [52, 53]. FGF10+ cells are progenitors to parabronchial smooth muscle

cells, and this progenitor status is maintained by mesothelial-derived FGF9, which inhibits

smooth muscle differentiation. Smooth muscle progenitors become gradually displaced from

distal to a proximal position alongside the bronchial tree in the mouse lung [35, 54] during

development. To investigate if FGF signaling is involved in *Svep1$^{-/-}$* branching defects and

airway smooth muscle differentiation, we treated lung explants with the FGF pathway inhibitor

SU5402 at a low enough dose to allow for some branching activity. SU5402 is a fibroblast

growth factor receptor 1 (FGFR1) tyrosine kinase inhibitor that blocks the FGF signaling

necessary for smooth muscle differentiation, leading to more smooth muscle around the

airway[55]. As expected, we observed branching defects in the control and *Svep1$^{-/-}$* lung explants

after 48hr of treatment (Fig. 6c). Immunofluorescence SOX9 and ACTA2 revealed normal distal

airway bifurcation in explants of *Svep1* wild type lungs and trifurcations in lung explants from

mutant embryos along with ectopic branching and budding anomalies (Fig. 6c). After 4 days of

FGF signaling inhibition, an increase in smooth muscle coverage around the distal airways,

wrapping around the distal tips, was observed in both normal lungs and, to a lesser degree, in

Svep1$^{-/-}$ lungs (Fig. 6d). Overall, these results suggest that the branching pattern defects in

Svep1 mutants are independent of the canonical FGF pathway, but that smooth muscle

differentiation defects are partially mediated by increased FGF signaling.

SVEP1 and FGF9 interact during branching to direct normal lung development

Fgf9 is essential for mouse lung development, and both loss and gain of function alleles of *Fgf9*

result in branching defects [33, 35, 36, 56]. *Fgf9* is expressed in both the lung mesothelium and distal

epithelium during early-to-mid branching morphogenesis. *Fgf9* levels in the mesothelium are

high at E12.5, but at E14.5 *Fgf9* is limited to the distal epithelium [33, 35]. We found that *Fgf9* is

normally expressed in mesothelium and epithelium at E14.5 and exclusively in the distal airway

tip epithelium of E16.5 lungs (mid-to-late branching) (Fig. 6e).

Overexpression of *Fgf9* causes branching defects that are characterized by dilatation of the

distal airways [35, 36, 53, 57], a phenotype that is opposite to the multi-budding phenotype of *Svep1*[-/-]

lungs at the pseudoglandular stage (Fig. 2c). Furthermore, overexpression of *Svep1* (Fig. 4a,b)

and loss of *Fgf9* function [33, 35] inhibit branching. To investigate whether SVEP1 and FGF9

interact during branching morphogenesis, we treated normal lung explants with exogenous

SVEP1 and FGF9 alone and in combination for two days. SVEP1 cultured explants showed a

significant decrease in branching (Fig. 6f) FGF9 cultured explants exhibited dilated distal

airways (Fig. 6f). Lung explants treated with both SVEP1 and FGF9 resulted in normal

branching and distal airways (Fig. 6f). That the treatment with combined SVEP1 and FGF9

restored branching and distal airway morphologies suggests that an interaction between SVEP1

and FGF9 is necessary for normal airway development.

Svep1 is necessary for cellular differentiation and saccule maturation

In *Svep1* mutants, alveolar saccules are either absent or underdeveloped, indicating that there

is a disruption in the normal alveolar differentiation process. Our transcriptomics data indicate

an increase in expression of AT2-related genes and a decrease in expression of AT1-related

genes in *Svep1*[-/-] lungs at E18.5 (Fig. 7a). Immunolabeling with AT1 markers (PDPL and HOPX)

and AT2 markers (SFTPC and ABCA3) reveals reduced PDPL and HOPX1 positive domains,

expanded SFTPC expression, and intensified expression of ABCA3 in the central saccules at

E18.5 (Fig. 7b). Additionally, SFTPC expression correlates with SOX9, indicating an immature

AT2 cell fate phenotype in mutants (Suppl. Fig. 4a). These findings underscore that *Svep1*

deficiency disrupts the balance of alveolar epithelial differentiation, favoring an AT2 cell fate.

This effect is particularly pronounced in the lung periphery which retains an AT2 progenitor

phenotype reminiscent of the pseudoglandular stage.

Alveolar ACTA2 positive myofibroblasts transiently differentiate from their PDGFRA positive

progenitors during septation which, during normal murine lung development, are located at the

base of saccules in the regions of future alveoli, where they invaginate toward the saccule

lumen [58, 59]. To assess if *Svep1*[-/-] lungs exhibit defects in alveolar myofibroblast progenitor

formation, we stained E18.5 tissues with PDGFRA, revealing a significant increase in the

number of PDGFRA[+] cells per saccular tissue in mutants (Fig. 7c). While RNA-seq analysis

showed no significant difference in *Pdgfra* gene expression between mutant and wild-type

lungs, the scRNA-seq data revealed a deficiency in *Pdgfra*[+] cells in the lungs of *Svep1*[-/-]

embryos (Suppl. Fig. 3b). The expression of *Pdgfa*, the PDGRA ligand crucial for septation,

myofibroblast differentiation, proliferation, and migration [60, 61], was reduced in *Svep1*$^{-/-}$ embryos

at E18.5 in the bulk RNA-seq data (Log2(FC)= - 0.19318, adjusted p value < 0.05 t test).

PDGFRA+ cells, which are known to give rise to alveolar lipofibroblasts characterized by lipid-

filled vesicles, exhibited no differences in *Plin2* (*perilipin 2*, also known as adipophilin; *Adfp*)

expression between mutant and wild-type lungs at E18.5 (Suppl. Fig. 4b). These findings show

that *Svep1* expression is essential for regulating alveolar myofibroblast progenitor formation.

Notably, despite the absence of primary septation, *Svep1*$^{-/-}$ mutants exhibit a broad

myofibroblast progenitor fate (PDGFRA+ cells). This increase may arise from an augmented

number of distal airways and/or defects in the myofibroblast differentiation process itself.

The lungs are covered by a monolayer of flat mesothelial cells that differentiate from round

Wilms' tumor transcription factor 1 (WT1) positive cells that surround the lung buds during early

branching [62]. *Wt1* expression normally decreases during development [62] whereas the

hypercellular mesothelium of the *Svep1*$^{-/-}$ lungs were enriched in WT1+ mesothelial progenitors

and had increased *Wt1* gene expression (Fig. 7d and Suppl. Fig. 5). To determine if the

expansion of mesothelial cells was due to an increase in cellular proliferation, we co-labeled

with WT1 and KI67 (Fig. 7d). Although there was a significant increase in the number of WT1+

cells in *Svep1*$^{-/-}$ lungs (Fig. 7d, left), the percentage of WT1+ that were also KI67+ was not

significantly different between normal and mutants at E16.5 (Fig. 7d, right), indicating that the

disruption of mesothelial differentiation in mutants is not due to an increase in proliferation.

RNA-seq data from the whole lung revealed that vascular development genes are

downregulated in the lungs of *Svep1*$^{-/-}$ embryos compared to wild-type controls (Fig. 6a). To

determine, therefore, if lung microvasculature formation is affected in *Svep1*$^{-/-}$ embryos, we

stained E16.5 (Suppl. Fig. 5) and E18.5 (Fig. 7e) lungs with PECAM1 (CD31). At E16.5 lung

microvasculature looks normal (Suppl. Fig. 5); however, at E18.5, the microvasculature was

disorganized and failed to form the double layer pattern characteristic of the saccular stage (Fig.

7e). These results indicate that *Svep1* is necessary for lung microvasculature plexus formation,

which is consistent with previous findings that *Svep1* plays a role in lymphatic vascular

development and remodeling in mice and explains why disruption of *Svep1* results in severe

edema in embryos [46, 47].

Discussion

While previous studies have led to important insights into the genetic and molecular processes that underlie lung development, our understanding of the factors that regulate the bronchial tree's stereotypic structure and the subsequent transition from branching to saccular formation remains incomplete. The study reported here revealed, for the first time, that *Svep1* function is necessary to preserve the stereotypic pattern of the bronchial tree, contributes to the transition required to end branching morphogenesis, and promotes alveolar differentiation and maturation (Fig. 8). The results also suggest that *Svep1* regulates FGF signaling pathway for smooth muscle differentiation and that the interaction between SVEP1 and FGF9 is essential for normal airway development. Although further investigation is needed to uncover molecular mechanisms, the findings reported here provide new insights into the role the extracellular matrix plays in governing lung morphogenesis.

Our work shows that *Svep1* plays a significant role in regulating the branching program in the developing lung. The loss of *Svep1* expression leads to disordered airway tips, particularly in the distal regions where *Svep1* expression is normally high. Our results demonstrate that SVEP1 acts as a negative regulator of lung branching, inhibiting the growth of new branches in lung explants. The effects of SVEP1 on the lung bud epithelium suggests a role in shaping the lobes by restricting the potential for aberrant branching. We hypothesize that early in lung development, the local genetic program driven by FGF from its signaling centers is sufficient to direct local branching; however, as lung airway complexity increases, this local signaling becomes diffuse and may activate the entire susceptible SOX9$^+$ directed epithelial branching. Therefore, negative regulators, like SVEP1 act to restrict excessive branching while ensuring proper airway patterning. Notably, at the pseudoglandular stage, lungs of *Svep1$^{-/-}$* embryos show no significant differences in the expression of essential branching genes. This suggests that *Svep1* may not be the primary regulator of the overall genetic branching program but rather a modulator of the timing and extent of branching events. *Svep1* most likely directs branching through localized signaling without significantly impacting genes fundamental for branching morphogenesis *(e.g., Fgf10*, *Shh*, and *Bmp4*), allowing early lung developmental processes while maintaining later appropriate branching morphology.

The recovery of SVEP1 branching patterns in lung explant cultures following exogenous treatment of FGF9 suggests a potential physical interaction between SVEP1 and FGF9 in the regulation of branching similar to the SVEP1-ANG2 during the remodeling of lymphatic vessels [47, 63]. It is plausible that SVEP1 interacts with FGF9 signaling locally in a particular distal epithelial progenitor to modify the overall branching program..

The disordered airways in the lungs of $Svep1^{-/-}$ embryos disrupts the shape of the lung lobes.

The disrupted branching program manifests as ectopic budding and trifurcations of distal

airways. The ordinarily thin and flat margins of the lung are rounded, and the tips of the lungs

are foreshortened in $Svep1^{-/-}$ embryos. Our 3D model of the right auxiliary (RAc) lobe showed

that branching in $Svep1^{-/-}$ embryos results in a pattern of dense rosettes generated at multiple

branching sites, resulting in a cauliflower-like appearance of the lungs at E18.5. In the lungs of

$Svep1^{-/-}$ embryos, we observed a dorsal-to-ventral maturation gradient with small immature

saccules on the dorsal side and narrow airways on the lobulated ventral side. The dorsal-ventral

spatial variation in aberrant lung lobe morphology is consistent with previous work showing that

tissue geometry and biomechanical forces influence the genetic branching program during lung

development [7, 64]. Our data indicate that the lobe-specific phenotypes in $Svep1^{-/-}$ lungs are due

to a complex interplay of genetic and physical factors and that the ventral lung needs tighter

control than the dorsal lung to develop properly to fill the chest cavity together with the heart and

diaphragm (ventral) and the body wall (dorsal). Hence, $Svep1$ is especially crucial for preserving

the basic structure of lung tips and edges during development.

The lungs of $Svep1^{-/-}$ embryos exhibit impaired cellular differentiation and saccular maturation.

$Svep1^{-/-}$ pups die shortly after birth, with a lung morphology more typical of the pseudoglandular

stage with enrichment of SOX9+ progenitor cells. The alveolar epithelium in the lungs of $Svep1$

mutant embryos is biased toward AT2 cells, which correlates with the proliferative SOX9+ cell

population which, in turn, is maintained by the persistent expression of genes involved in FGF

signaling that impedes differentiation of AT1 cells [27, 60]. Furthermore, $Svep1^{-/-}$ lungs are deficient

in HOPX+ and PDPL+ AT1 cells with reduced lung bud stalks potentially leading to proximal

airway hypoplasia. $Svep1^{-/-}$ lungs have reduced smooth muscle differentiation that can be

rescued by the inhibition of FGF signaling. The observed increase in SCGBA1+ club cells and

the concomitant reduction in bronchiole lumen diameter point to a profound influence of the loss

of $Svep1$ on airway morphology, in addition to the smooth muscle development defects around

the proximal airways. Furthermore, $Svep1^{-/-}$ lungs show reduced expression of genes related to

smooth muscle, including $Acta2$. This deficiency may contribute to the impact of the loss of

$Svep1$ on septa and airway hypoplasia, influencing the distalization of the airway.

Other studies have found that mechanical pressures, in conjunction with growth factors, are

essential for proper alveolar differentiation during development [65]. $Svep1^{-/-}$ embryos are

characterized by chest cavity edema secondary to lymphatic defects at E18.5 which may cause

atypical internal lung pressures that could, in turn, contribute to the observed aberrant cell

differentiation. Mesothelial cells and alveolar myofibroblasts at E18.5 in $Svep1^{-/-}$ are also

maintained in a progenitor state of immature saccular development marked by increased WT1

and PDGFRA expression in mutants.

Maintenance of the extracellular matrix has been shown previously to be essential for normal

organ and epithelial tube morphogenesis [38, 66] and cellular differentiation processes [66]. Our

transcriptome analysis indicated that the absence of *Svep1* disrupts the expression of genes

associated with cell adhesion, cell motility, and ECM organization. Our transcriptomics data

showed that *Col1a1* and the basement membrane collagen *Col4a4*, are significantly reduced in

E18.5 $Svep1^{-/-}$ lungs, disruption of which may contribute to the proliferation and differentiation

defects that we observed in the lungs of $Svep1^{-/-}$ *embryos in this study.*

The transition from branching to differentiation in lung development is essential for establishing

the foundation for alveologenesis and has significant clinical ramifications for neonates born

with such respiratory deficiencies. Our findings demonstrate the importance of the ECM in

directing airway architecture and the transition from branching programs to alveolar

differentiation. Additional research into the specific role that *Svep1* plays in preparing the lung

for alveologenesis is required to understand more fully the molecular, cellular, and mechanical

mechanisms critical to this transition and its potential for targeted therapeutic interventions to

improve the outcomes of patients born with impaired or delayed lung development.

Methods

Svep1 mutant mice

Svep1 mutant mice (B6N(Cg)-*Svep1*$^{tm1b(EUCOMM)Hmgu}$/J; MGI:5509058) were generated by the

Knockout Mouse Phenotyping Program (KOMP2) by inserting the L1L2_Bact_P cassette

upstream of exon 8 of *Svep1* [49]. This cassette contains a flippase recombinase target (FRT) site

followed by a lacZ sequence and a loxP site. The first loxP site is followed by a neomycin

resistance gene, a second FRT site, and a second loxP site. A third loxP site is inserted

downstream of exon 8. The construct was introduced into embryonic cells and embryonic stem

cell clone HEPD0747_6_B06 was injected into B6(Cg)-Tyr^{c-2J}/J blastocysts. Resulting chimeric

males were bred to C57BL/6NJ female mice to generate heterozygous tm1a (i.e., knockout first)

animals[67]. No homozygous mice were recovered at weaning from heterozygous intercrosses.

Heterozygous tm1a mice were then bred to B6N.Cg-Tg (*Sox2*-cre)1Amc/J mice to remove the

floxed neomycin sequence and exon 8 of *Svep1*. Resulting offspring were bred to C57BL/6NJ

mice to remove the *cre*-expressing transgene. Genotyping was performed as described in

Supplementary Methods.

Histology

Mouse embryos were euthanized by decapitation at E14.5, E16.5, or E18.5. Tails were

collected for genotyping. Whole lungs were excised surgically and fixed in 4%

paraformaldehyde (PFA) and then embedded in paraffin. Paraffin tissue blocks were sectioned

(5 μm) and stained for Hematoxylin and Eosin (H&E) using standard procedures.

In situ hybridization (ISH)

For whole mount *in situ* hybridization of *Svep1*, a 725bp segment of the *Svep1* transcript

(NM_153366.4; 746-11179) was PCR amplified (PCR Master Mix, Promega). *In situ*

hybridization was performed using methods published previously[68] and developed using BM

purple AP substrate (Roche) per the manufacturer's instructions. 5 μm thick sections of paraffin

embedded lung tissues were used for ISH of *Fgf10* (NM_446371, 862-1978 nt), *Fgf9*

(NM_013518.4, 27-1198 nt), *Fgfr2* (NM_010207, 2-1677nt), and *Svep1* (NM_022814.2, 2879 -

3726 nt) using RNAscope 2.0 Red Detection Kit (Advanced Cell Diagnostic) according to the

manufacturer's instructions.

A *Svep1* riboprobe was synthesized with one pair of exon-exon boundary overlapping primers

(5' TGTGGTCCTCCAAGTCACGTA 3'; 5' CCAGCAGACAGCAGAGTATGT 3') designed using

NCBI's Primer-BLAST[69]. The purified PCR fragment was cloned into the pCR™II-TOPO® TA

vector (TOPO® TA Cloning® Kit, Dual Promoter, ThermoFisher), and transformed into One

Shot® TOP10 Chemically Competent Cells, ThermoFisher). Ampicillin resistance transformed

colonies were selected. The orientation of the insert after linearization of the vector with SpeI

restriction digestion (New England Biolabs, Inc.) was determined by Sanger sequencing. Sense

and anti-sense Digoxigenin-11-UTP labeled probes (DIG RNA Labeling Mix, Sigma-Aldrich)

were synthesized with SP6 and T7 RNA polymerases, respectively.

Immunohistochemistry (IHC) on sectioned lung tissue was performed using standard

techniques. Details on the antibodies used for IHC are provided in Supplementary Methods.

Antigen retrieval was achieved by heat treatment in a microwave oven for 20 mins at low power

in 0.01 M sodium citrate buffer at pH 6. For tissues imaged using fluorescence, Alexa Fluor

secondary antibodies were used (Life Technology). Images were obtained with a Nikon80i or

Keyence (BZX800) microscope.

Quantification of lung distal epithelial tips

481 Fixed whole lungs were washed with PBS and permeabilized in 0.5% Triton-X, 5% BSA in PBS

482 for 1 hour at room temp. SOX9 primary antibody diluted in 1X PBS was incubated overnight at

483 room temperature. Samples were washed and incubated with an Alexa fluor (Life Technology,

484 USA) secondary for 2 hrs at room temperature, mounted with DAPI mounting medium. and fixed

485 with 4% PFA in PBS for 2–4 hrs.

Tissue-clearing for evaluation of branching patterns

487 Fixed whole lungs from E13.5 embryos were dehydrated in a methanol series, washed with

488 20% DMSO in methanol, and then re-hydrated with a methanol series. Lungs were then

489 permeabilized with 0.2% Triton-X in PBS. Lung tissue clearing was performed using the

490 Visikol® Histo™ kit (Visikol, Inc; NJ) according to the manufacturer's instructions. Lungs

491 mounted in silicone-imaging chambers were imaged with a Keyence (BZX800) microscope.

Whole mount branching immunofluorescence

Lung explants were fixed with 4%PFA, dehydrated in a PBS/methanol series, bleached with 6%

hydrogen peroxide (H1009, Sigma) in methanol overnight at 4° C, then re-hydrated in a

methanol series. Samples were blocked in PBS + 0.3% Triton X-100 + 5% normal goat serum

for 2 hrs and incubated with primary antibodies into blocking solution overnight at 4° C. Samples

were washed with PBS + 0.1% Triton X-100 + 0.1% Tween-20, followed by addition of Alexa

Fluor antibodies (Life Technology) diluted in PBS + 0.3% Triton X-100 at 4 °C overnight.

Samples were mounted with DAPI mounting medium. washed, then fixed with 4% PFA in PBS

for 2–4 hrs. Whole lungs were imaged using a Keyence microscope. A Z-stack image series

(BZ-H4XD) was generated for quantification and analyses of branching patterns of embryonic

whole lungs. Time lapse videos were generated to trace the lineages of airway branches for

orthogonal and planar bifurcation, and for quantification of trifurcation and ectopic branching.

Statistical significance was evaluated using a two-tailed t test with significance at p <0.05.

Embryonic lung explant imaging

Lungs from E12.5 or E13.5 embryos were dissected and transferred to the inserts of Corning

transwells (Millipore Sigma) containing 2 ml of DMEM/F12 medium supplemented with 1 U/ml

penicillin-streptomycin. Lungs were incubated in the chamber and time-lapse images were

captured every 10 min over 48 hrs using the Keyence BZX800 software (BZ-H4CXT).

Morphometric analysis of treated lung explants

Lung explants from E12.5 or E13.5 *Svep1*[+/+] and *Svep1*[-/-] embryos were treated for 2 days with

a 146 amino acid recombinant protein antigen of *Svep1* (10 µg/ml, NP_0735725.2, region 874-

971, Novus Biologicals), for 2 to 4 days with and without FGFR2 kinase inhibitor (2.5 µM

SU5401, Pfizer) signaling and for two days with FGF9 (200 µM; R&D system) or FGF9+SVEP1

recombinant protein

The fold change of epithelial branches was determined by manual counting of individual

branches compared to epithelial buds emerging from the lung epithelium at various intervals

throughout a time lapse experiment using ImageJ [70]. Lung buds were categorized as peripheral

branches only if they appeared at the most distal end of the secondary bronchi. Quantification

was performed for the left lobe only. Statistical significance of the counts was evaluated using a

two-tailed *t* test with significance at p <0.05.

RNA sequencing and analyses

Total RNA was isolated from whole embryonic lungs at E18.5 using the RNeasy Mini Kit

(Qiagen, Valencia, CA, USA) according to the manufacturer's instructions. Double stranded

cDNA was synthesized from total mouse lung RNA using the iScriptTM cDNA Synthesis Kit

from BIORAD following the manufacturer's instructions. RNA samples were sent to Novogene

Co. Ltd for sequencing on the Illumina platform (see Supplementary Methods). Annotation term

enrichment for pathways and function for differentially expressed genes was performed using

Metascape (http://metascape.org) [71], VisuaL Annotation Display (VLAD) [72]

(http://proto.informatics.jax.org/prototypes/vlad/), and the Gene Ontology term enrichment tool

(http://geneontology.org/docs/go-enrichment-analysis/). Heat maps of gene expression were

generated using the Morpheus matrix visualization and analysis software

(https://software.broadinstitute.org/morpheus).

Single cell RNA-seq (scRNA-seq) sequencing and analysis

Single cell suspensions for scRNA-seq were prepared from E18.5 embryos, sequenced, and

processed as described in Supplementary Methods. The Seurat software package was used to

analyze the processed data [73]. To determine the cell populations most impacted by the loss of

Svep1, the distribution of cell populations in each experimental group was determined by

calculating the ratio of each Seurat cluster to total number of cells from an experimental group.

A t-test of the ratios was conducted to determine populations significantly enriched or depleted

in the mutant tissue. Clusters demonstrating a segregation between mutant and wild type origin

cells were compared to the nearest-related clusters using the Wilcoxon Rank Sum test.

Quantitative PCR

Real-time PCR was performed on cDNA with IQ SYBR Green Supermix (Biorad,) and gene-

specific primers (Supplementary Methods) and beta-actin (*Actb*) as the reference. The data

were analyzed using the ΔΔCt method on the BioRad CFX manager v1.5. RT-qPCR mRNA

expression was analyzed by the Wilcoxon rank-sum test.

Data availability

RNA-seq and scRNA-seq data from this study will be submitted to a public archive upon

publication. The *Svep1* mouse is available from the Mutant Mouse Resource and Research

Center (MMRRC:049943-UCD).

Acknowledgements

The authors thank Drs. Martin Ringwald and Vidhya Munnamalai for their comments on an early

draft of this manuscript. Sandy Daigle (The Jackson Laboratory) provided essential technical

support and advice for the scRNA-seq experiment. Pete Finger (The Jackson Laboratory)

performed the toluidine blue staining.

Funding

Funding was provided by the National Institute of Child Health and Human Development

(NICHD) P01HD068250 to PKD, CB, and ML.

References

1.   Rawlins EL. The building blocks of mammalian lung development. *Dev Dyn* **240**,
463-476 (2011).

2.   Chang DR*, et al.* Lung epithelial branching program antagonizes alveolar
differentiation. *Proc Natl Acad Sci U S A* **110**, 18042-18051 (2013).

3.   Frank DB*, et al.* Early lineage specification defines alveolar epithelial ontogeny in
the murine lung. *Proc Natl Acad Sci U S A* **116**, 4362-4371 (2019).

4.   Ten Have-Opbroek AAW. Lung development in the mouse embryo. *Experimental*
Lung Research **17**, 111-130 (1991).

5.   Warburton D*, et al.* Lung organogenesis. *Curr Top Dev Biol* **90**, 73-158 (2010).

6.   Metzger RJ, Klein OD, Martin GR, Krasnow MA. The branching programme of
mouse lung development. *Nature* **453**, 745-750 (2008).

7.   Blanc P*, et al.* A role for mesenchyme dynamics in mouse lung branching
morphogenesis. *PLoS One* **7**, e41643 (2012).

8.   Zhao X*, et al.* Bronchial anatomy of left lung: a study of multi-detector row CT.
Surg Radiol Anat **31**, 85-91 (2009).

9.   Rawlins EL, Clark CP, Xue Y, Hogan BL. The Id2+ distal tip lung epithelium
contains individual multipotent embryonic progenitor cells. *Development* **136**,
3741-3745 (2009).

10.   Yang Y*, et al.* Comparative transcriptome analysis reveals conserved branching
morphogenesis related genes involved in chamber formation of catfish
swimbladder. *Physiol Genomics* **50**, 67-76 (2018).

11.   Zhang K, Aung T, Yao E, Chuang PT. Lung patterning: Is a distal-to-proximal
gradient of cell allocation and fate decision a general paradigm?: A gradient of
distal-to-proximal distribution and differentiation of tip progenitors produces
distinct compartments in the lung. *Bioessays* **46**, e2300083 (2024).

12.   Conway FM, Garner JL, Orton CM, Srikanthan K, Kemp SV, Shah PL.
Contemporary Concise Review 2018: Lung cancer and pleural disease.
Respirology **24**, 475-483 (2019).

13.   Rock JR, Hogan BL. Epithelial progenitor cells in lung development,
maintenance, repair, and disease. *Annu Rev Cell Dev Biol* **27**, 493-512 (2011).

14.   Herriges M, Morrisey EE. Lung development: orchestrating the generation and
regeneration of a complex organ. *Development* **141**, 502-513 (2014).

15. Whitsett JA, Kalin TV, Xu Y, Kalinichenko VV. Building and Regenerating the
Lung Cell by Cell. *Physiol Rev* **99**, 513-554 (2019).
613

614 16. Morrisey EE, Hogan BL. Preparing for the first breath: genetic and cellular
615   mechanisms in lung development. *Dev Cell* **18**, 8-23 (2010).
616

617 17. Bellusci S, Grindley J, Emoto H, Itoh N, Hogan BL. Fibroblast growth factor 10
618   (FGF10) and branching morphogenesis in the embryonic mouse lung.
619   *Development* **124**, 4867-4878 (1997).
620

621 18. Min H*, et al.* Fgf-10 is required for both limb and lung development and exhibits
622   striking functional similarity to Drosophila branchless. *Genes Dev* **12**, 3156-3161
623   (1998).
624

625 19. Sekine K*, et al.* Fgf10 is essential for limb and lung formation. *Nat Genet* **21**, 138-
626   141 (1999).
627

628 20. De Moerlooze L, Spencer-Dene B, Revest JM, Hajihosseini M, Rosewell I,
629   Dickson C. An important role for the IIIb isoform of fibroblast growth factor
630   receptor 2 (FGFR2) in mesenchymal-epithelial signalling during mouse
631   organogenesis. *Development* **127**, 483-492 (2000).
632

633 21. Weaver M, Dunn NR, Hogan BL. Bmp4 and Fgf10 play opposing roles during
634   lung bud morphogenesis. *Development* **127**, 2695-2704 (2000).
635

636 22. Abler LL, Mansour SL, Sun X. Conditional gene inactivation reveals roles for
637   Fgf10 and Fgfr2 in establishing a normal pattern of epithelial branching in the
638   mouse lung. *Dev Dyn* **238**, 1999-2013 (2009).
639

640 23. Ramasamy SK*, et al.* Fgf10 dosage is critical for the amplification of epithelial cell
641   progenitors and for the formation of multiple mesenchymal lineages during lung
642   development. *Dev Biol* **307**, 237-247 (2007).
643

644 24. Warburton D. Developmental biology: order in the lung. *Nature* **453**, 733-735
645   (2008).
646

647 25. Hyatt BA, Shangguan X, Shannon JM. FGF-10 induces SP-C and Bmp4 and
648   regulates proximal-distal patterning in embryonic tracheal epithelium. *Am J
649   Physiol Lung Cell Mol Physiol* **287**, L1116-1126 (2004).
650

651 26. Ornitz DM, Yin Y. Signaling networks regulating development of the lower
652   respiratory tract. *Cold Spring Harb Perspect Biol* **4**, (2012).
653

27. Volckaert T, Campbell A, Dill E, Li C, Minoo P, De Langhe S. Localized Fgf10
expression is not required for lung branching morphogenesis but prevents
differentiation of epithelial progenitors. *Development* **140**, 3731-3742 (2013).

28. Park WY, Miranda B, Lebeche D, Hashimoto G, Cardoso WV. FGF-10 is a
chemotactic factor for distal epithelial buds during lung development. *Dev Biol*
201, 125-134 (1998).

29. Warburton D*, et al.* Molecular mechanisms of early lung specification and
branching morphogenesis. *Pediatr Res* **57**, 26R-37R (2005).

30. Li J, Wang Z, Chu Q, Jiang K, Li J, Tang N. The Strength of Mechanical Forces
Determines the Differentiation of Alveolar Epithelial Cells. *Dev Cell* **44**, 297-312
e295 (2018).

31. Mailleux AA*, et al.* Fgf10 expression identifies parabronchial smooth muscle cell
progenitors and is required for their entry into the smooth muscle cell lineage.
Development **132**, 2157-2166 (2005).

32. Yuan T, Volckaert T, Chanda D, Thannickal VJ, De Langhe SP. Fgf10 Signaling
in Lung Development, Homeostasis, Disease, and Repair After Injury. *Front*
Genet **9**, 418 (2018).

33. Colvin JS, Feldman B, Nadeau JH, Goldfarb M, Ornitz DM. Genomic
organization and embryonic expression of the mouse fibroblast growth factor 9
gene. *Dev Dyn* **216**, 72-88 (1999).

34. Colvin JS, White AC, Pratt SJ, Ornitz DM. Lung hypoplasia and neonatal death in
Fgf9-null mice identify this gene as an essential regulator of lung mesenchyme.
Development **128**, 2095-2106 (2001).

35. del Moral PM*, et al.* Differential role of FGF9 on epithelium and mesenchyme in
mouse embryonic lung. *Dev Biol* **293**, 77-89 (2006).

36. Yin Y, Wang F, Ornitz DM. Mesothelial- and epithelial-derived FGF9 have distinct
functions in the regulation of lung development. *Development* **138**, 3169-3177
(2011).

37. Zhou Y*, et al.* Extracellular matrix in lung development, homeostasis and
disease. *Matrix Biol* **73**, 77-104 (2018).

38. Mecham RP. Elastin in lung development and disease pathogenesis. *Matrix Biol*
73, 6-20 (2018).

39. Nguyen NM, Miner JH, Pierce RA, Senior RM. Laminin alpha 5 is required for
lobar septation and visceral pleural basement membrane formation in the
developing mouse lung. *Dev Biol* **246**, 231-244 (2002).

40. Sakai T, Larsen M, Yamada KM. Fibronectin requirement in branching
morphogenesis. *Nature* **423**, 876-881 (2003).
704

705 41. Chen J, Krasnow MA. Integrin Beta 1 suppresses multilayering of a simple
706   epithelium. *PLoS One* **7**, e52886 (2012).
707

708 42. Loscertales M*, et al.* Type IV collagen drives alveolar epithelial-endothelial
709   association and the morphogenetic movements of septation. *BMC Biol* **14**, 59
710   (2016).
711

712 43. Walma DAC, Yamada KM. The extracellular matrix in development.
713   *Development* **147**, (2020).
714

715 44. Sato-Nishiuchi R*, et al.* Polydom/SVEP1 is a ligand for integrin alpha9beta1. *J
716   Biol Chem* **287**, 25615-25630 (2012).
717

718 45. Samuelov L*, et al.* SVEP1 plays a crucial role in epidermal differentiation. *Exp
719   Dermatol* **26**, 423-430 (2017).
720

721 46. Karpanen T*, et al.* An Evolutionarily Conserved Role for Polydom/Svep1 During
722   Lymphatic Vessel Formation. *Circ Res* **120**, 1263-1275 (2017).
723

724 47. Morooka N*, et al.* Polydom Is an Extracellular Matrix Protein Involved in
725   Lymphatic Vessel Remodeling. *Circ Res* **120**, 1276-1288 (2017).
726

727 48. Kishore R*, et al.* Automated generation of gene summaries at the Alliance of
728   Genome Resources. *Database (Oxford)* **2020**, (2020).
729

730 49. Dickinson ME*, et al.* High-throughput discovery of novel developmental
731   phenotypes. *Nature* **537**, 508-514 (2016).
732

733 50. Beauchemin KJ*, et al.* Temporal dynamics of the developing lung transcriptome
734   in three common inbred strains of laboratory mice reveals multiple stages of
735   postnatal alveolar development. *PeerJ* **4**, e2318 (2016).
736

737 51. Desai TJ, Brownfield DG, Krasnow MA. Alveolar progenitor and stem cells in
738   lung development, renewal and cancer. *Nature* **507**, 190-194 (2014).
739

740 52. El Agha E, Bellusci S. Walking along the Fibroblast Growth Factor 10 Route: A
741   Key Pathway to Understand the Control and Regulation of Epithelial and
742   Mesenchymal Cell-Lineage Formation during Lung Development and Repair
743   after Injury. *Scientifica (Cairo)* **2014**, 538379 (2014).
744

745 53. Yin Y, Ornitz DM. FGF9 and FGF10 activate distinct signaling pathways to direct
746   lung epithelial specification and branching. *Sci Signal* **13**, (2020).
747

54. Yi L, Domyan ET, Lewandoski M, Sun X. Fibroblast growth factor 9 signaling
inhibits airway smooth muscle differentiation in mouse lung. *Dev Dyn* **238**, 123-
137 (2009).

55. Goodwin K*, et al.* Smooth muscle differentiation shapes domain branches during
mouse lung development. *Development* **146**, (2019).

56. Lin Y, Liu G, Wang F. Generation of an Fgf9 conditional null allele. *Genesis* **44**,
150-154 (2006).

57. White AC, Xu J, Yin Y, Smith C, Schmid G, Ornitz DM. FGF9 and SHH signaling
coordinate lung growth and development through regulation of distinct
mesenchymal domains. *Development* **133**, 1507-1517 (2006).

58. McGowan SE, Grossmann RE, Kimani PW, Holmes AJ. Platelet-derived growth
factor receptor-alpha-expressing cells localize to the alveolar entry ring and have
characteristics of myofibroblasts during pulmonary alveolar septal formation.
Anat Rec (Hoboken) **291**, 1649-1661 (2008).

59. Branchfield K, Li R, Lungova V, Verheyden JM, McCulley D, Sun X. A three-
dimensional study of alveologenesis in mouse lung. *Dev Biol* **409**, 429-441
(2016).

60. Gouveia L, Betsholtz C, Andrae J. PDGF-A signaling is required for secondary
alveolar septation and controls epithelial proliferation in the developing lung.
Development **145**, (2018).

61. Betsholtz C. Insight into the physiological functions of PDGF through genetic
studies in mice. *Cytokine Growth Factor Rev* **15**, 215-228 (2004).

62. Dixit R, Ai X, Fine A. Derivation of lung mesenchymal lineages from the fetal
mesothelium requires hedgehog signaling for mesothelial cell entry.
Development **140**, 4398-4406 (2013).

63. Sato-Nishiuchi R, Doiguchi M, Morooka N, Sekiguchi K. Polydom/SVEP1 binds
to Tie1 and promotes migration of lymphatic endothelial cells. *J Cell Biol* **222**,
(2023).

64. Lang C, Conrad L, Iber D. Organ-Specific Branching Morphogenesis. *Front Cell
Dev Biol* **9**, 671402 (2021).

65. Rozario T, DeSimone DW. The extracellular matrix in development and
morphogenesis: a dynamic view. *Dev Biol* **341**, 126-140 (2010).

66. Streuli C. Extracellular matrix remodelling and cellular differentiation. *Curr Opin
Cell Biol* **11**, 634-640 (1999).

67. Skarnes WC*, et al.* A conditional knockout resource for the genome-wide study of
mouse gene function. *Nature* **474**, 337-342 (2011).

68. Riddle RD, Johnson RL, Laufer E, Tabin C. Sonic hedgehog mediates the
polarizing activity of the ZPA. *Cell* **75**, 1401-1416 (1993).

69. Ye J, Coulouris G, Zaretskaya I, Cutcutache I, Rozen S, Madden TL. Primer-
BLAST: a tool to design target-specific primers for polymerase chain reaction.
BMC Bioinformatics **13**, 134 (2012).

70. Schneider CA, Rasband WS, Eliceiri KW. NIH Image to ImageJ: 25 years of
image analysis. *Nat Methods* **9**, 671-675 (2012).

71. Zhou Y*, et al.* Metascape provides a biologist-oriented resource for the analysis
of systems-level datasets. *Nat Commun* **10**, 1523 (2019).

72. Richardson JE, Bult CJ. Visual annotation display (VLAD): a tool for finding
functional themes in lists of genes. *Mamm Genome* **26**, 567-573 (2015).

73. Hao Y*, et al.* Integrated analysis of multimodal single-cell data. *Cell* **184**, 3573-
3587 e3529 (2021).

Figure 1. Svep1 expression pattern and Svep1-/- hypoplastic lung lobes anomalies. **(a)** *In*

situ hybridization of lungs from E14.5 wild type embryos showed strong *Svep1* mRNA

expression in the mesenchyme at the lung periphery and around bifurcating distal airway tips

(right inset outlined in red, arrowhead). **(b)** Arrowhead shows Mesenchymal expression of

SVEP1 protein in normal E12.5 lung. White arrows indicate the location of the SVEP1 protein on

the epithelial membrane stained with ECAD (purple) in a distal airway bud. **(c)** SVEP1

expression is strong in the mesenchyme towards the lobe tip of the E14.5 lungs and branching

cleft sites (arrowheads). **(d)** SVEP1 protein localizes in the lung parenchyma including the

primary septa (upper box) and adjacent to the proximal airway epithelium (lower box). **(e)**

microCT scan of E18.5 embryos showing multiple defects including thinning of the diaphragm

and lung hypoplasia (red arrows). **(f)** Ventral view of E18.5 of $Svep1^{+/+}$ and $Svep1^{-/-}$ lungs. The

arrowheads indicate the tips of the lung lobes, which are short and rounded (highlight in black)

in $Svep1^{-/-}$ mice and have like a cauliflower morphology. **(g)** H&E-stained lung sections embryos

demonstrate that E16.5 lungs from $Svep1^{-/-}$ embryos are smaller and have irregular lobe edges.

Black arrowheads indicate left lobes with highlighted surface. **(h)** The red arrowheads show the

dorsal side of the E18 lungs, whereas the black arrows highlight the ventral side. $Svep1^{-/-}$

embryos show defective saccular development mainly at the edge on the ventral side of the

lung (inserts arrowheads) . Co-localization of ECAD and SOX9 confirms high expression of

SOX9 in distal airways (white arrowheads) in the lung of $Svep1^{-/-}$ embryos at E18.5. **(i)** Plots

showing *Sox9* relative mRNA expression at E14.5, E16.5, and E18.5. mRNA expression is

significantly high at E18.5 in $Svep1^{-/-}$ embryos (average number of tips per area ± SEM; p <

0.05; n ≥ 5). Right cranial (RCr), Right Caudal (RCa), Right Medial (RMe), and Right Accessory

(RAc) lobes. *Scale bars: 50µm (a, b, c, d), 100 µm.*

Figure 1

a   E14.5 *Svep1^{+/+}*

b   E12.5 *Svep1^{+/+}*   SVEP1 ECAD

c   E14.5 *Svep1^{+/+}*   SVEP1 DAPI

d   E18.5 *Svep1^{+/+}*   SVEP1 DAPI

e   E18.5 *Svep1^{+/+}*   E18.5 *Svep1^{-/-}*

f   ventral view   E18.5 *Svep1^{+/+}*
RCr   RAc   Heart
RMe
RCa   Left

ventral view   E18.5 *Svep1^{-/-}*
RCr   RAc   Heart
RMe
RCa   Left

g   H&E   E16.5 *Svep1^{+/+}*
E16.5 *Svep1^{-/-}*

h   H&E   E18.5 *Svep1^{+/+}*   ECAD SOX9 DAPI
E18.5 *Svep1^{-/-}*

i   *Svep1^{+/+}*   *Svep1^{-/-}*
Relative *Sox9* mRNA abundance (2^(-ΔΔCt))
E14.5   E16.5   E18.5

Figure 2. Airway organization defects and abnormal branching patterns in *Svep1^{-/-}*

embryonic lungs resulting in lung lobe shape anomalies. (a) Co-localization of epithelial

(ECAD) and epithelial progenitor (SOX9) markers showing distal airways in lungs of E16.5 mice.

(a') The blue highlights in inserts represent the distal tip of the normal and mutant RAc lung

lobes, with the mutant left lobe showing multiple branching. **(a")** Inserts illustrate airway density

at the lung periphery in WT and mutants, with mutants having higher density. **(a''')** In *Svep1^{-/-}*

lungs, distal airway tips are organized in packed airway rosettes on the ventral side of the lung

lobe. **(b)** Illustration of the three branching modes that pattern the airways of the lung (domain

branching, planar bifurcation, orthogonal bifurcation) and distal trifurcation. **(c-f)** Representative

Z-stack images show SOX9 immunofluorescence **(c)** E13.5 Whole lungs of normal and *Svep1^{-/-}*

embryos. The images show *Svep1^{-/-}* lungs with abnormal RCa lobe morphology (arrowheads),

RAc lobe distal tip branching (arrows), and secondary branching pattering (orange arrowheads).

(d) Pictures of E13.5 *Svep1^{+/+}* and *Svep1^{-/-}* embryos showing ectopic branching/budding

anomalies in mutant embryos (areas outlined in pink in lower inserts). **(e)** E14.5 left lung lobes

of *Svep1^{+/+}* and *Svep1^{-/-}* embryos. SOX9⁺ buds highlighted in pink and cartoon show airway tip

arrangement in normal and mutants. Arrow shows defective tip arrangement resulting in

abnormal lung tip morphology in *Svep1^{-/-}* embryos. **(f)** Cleared tissue immunofluorescence of

SOX9 in the left lung lobe at E14.5 reveals airway disorganization in *Svep1^{-/-}* embryos vs

Svep1^{+/+} embryos. Distal lung tips highlighted show branching trifurcations in the *Svep1^{-/-}*

embryo (white arrow) and aberrant distal tip airway. **(g)** Representation of the left lung lobe Z-

stack for distal airway quantification. Whole lungs of *Svep1^{-/-}* embryos at E14.5 have a

statistically significant increase in the total number of SOX9 positive airway tips and ratio of tip

and area compared to wild type embryos (average number of tips for area $\pm$ SEM; $p < 0.05$; n $\geq$

5). *Right cranial (RCr), Right Caudal (RCa), Right Medial (RMe), and Right Accessory (RAc)*

lobes. Scale bars: 100µm (a-e).

Figure 2

Figure 3. Disrupted branching programs and altered airway architecture in _Svep1^{-/-}_ RAc lung lobes at E14.5. (a) Compilation of a Z-stack image series for tissue-cleared whole RAc lung lobes labeled with ECAD (green) and SOX9 (purple) highlight disorganized branching in _Svep1^{-/-}_ embryos at E14.5. Arrows (lower right) point to branching airways with rosette-like arrangements and ectopic airway tips growing in random directions. Arrowheads (left panels) at the RCa lobe tips indicate a single tip in the lobe of _Svep1^{+/+}_ embryos but multi branched tips in _Svep1_ knockout embryos. **(b)** Schematic illustrating representative domain (blue), planar (green), and orthogonal (red) branching program types of _Svep1^{+/+}_ and _Svep1^{-/-}_ RAc lobes. Lungs of _Svep1^{-/-}_ embryos lack planar bifurcations and have many trifurcating branch tips (red highlighted in black). **(c-d)** Graphs displaying the percentages of branching modes, orthogonal bifurcations and trifurcations **(c)** and ectopic budding **(d)** in E14.5 Rac lobes of wild type and _Svep1^{-/-}_ embryos. **(e)** Representative single Z-stack images of RAc lobes at E14.5, stained with ECAD (green) and SOX9 (purple), show bifurcations (white dotted lines and white arrowhead) in _Svep1^{+/+}_ embryos, but trifurcations (white dotted lines and white arrows) in _Svep1^{-/-}_ mice. **(f)** Single Z-stack image of RAc lobes stained with ECAD (green) and SOX9 (red) display ectopic domain branching in _Svep1^{-/-}_ mice (arrow). **(g)** Lung explants from E12.5 embryos after 48 hours in organ culture. Explants from _Svep1^{+/+}_ embryos show normal bifurcation (black arrowheads) whereas explants from _Svep1^{-/-}_ embryos show trifurcations (black arrows) and ectopic budding (red arrowheads). (Average number of tips per area ± SEM; $p < 0.05$; $n \geq 5$) (c,d). _Scale bars 100µm (a, g); 50µm (e)._

Figure 3

Figure 4. SVEP1 treatment modulates airway branching in E12.5 lung explants

(a) Images of lung explants from E12.5 *Svep1$^{+/+}$* embryos cultured for 48 hrs. Images of lung

explants from E12.5 *Svep1$^{+/+}$* embryos cultured for 48 hrs. show a significant reduction of

branching (control, top row) when treated with SVEP1 peptide (bottom row). **(b)** Plot showing

peripheral airway counts of a left lung lobe after 48 hrs cultured with and without SVEP1

treatment. **(c)** *Svep1$^{-/-}$* lung explants treated with SVEP1 for 48 hours show a reduction in

trifurcations compared to untreated lung explants . Black arrowheads indicate bifurcation;

arrows indicate trifurcations. (Average counts ± SEM; *t* test $p < 0.05$ n$\leq$ 6). *Scale bars 100 µm*

(a,c).

Figure 4

a

0h   24h   48h

E12.5 Control

0h   24h   48h

E12.5 rSVEP1

b

Left lobe

Peripheral airway fold change

*

● Control   ● rSVEP1

c

E12.5 *Svep1⁺/⁺*   E12.5 *Svep1⁻/⁻*, rSVEP1

48h   48h

Figure 5. Impaired smooth muscle and airway epithelium differentiation in

Svep1$^{-/-}$ embryonic lungs. (a) Interactome of downregulated genes in lungs of E18.5 *Svep1$^{-/-}$*

embryos with clusters labeled by enriched annotation terms. **(b)** Gene expression heatmap from

RNA sequencing shows significantly downregulated genes in lungs from *Svep1$^{-/-}$* embryos are

associated with collagen and smooth muscle genes. **(c)** ACTA2 and Calponin staining at E16.5.

(d) ACTA2 and calponin localization in lung at E18.5 (white arrowheads indicate distal

respiratory bronchioles; orange arrows indicate proximal bronchioles; white arrows indicate

blood vessels). **(e)** ACTA2 staining (red) in wild type and *Svep1$^{-/-}$* E12.5 lung explants each

cultured for six days, showed a reduction of smooth muscle cell formation in secondary and

tertiary bronchi of the mutants (white arrows). **(f)** SOX2 Immunofluorescence staining marking

the progenitor cells of the epithelium in the proximal airways. Arrow marks the proximal airway

at the core of the *Svep1$^{-/-}$* lung. **(g)** Immunofluorescence of the club cell marker SCGB1A1

(green) highlights the small proximal airways in mutant lungs. SCGB1A1 (green) colocalization

with the ciliated cell marker FOXJ1 (pink) reveals a multicellular airway epithelium composed

primarily of club cells. In *Svep1$^{-/-}$* embryos, the bronchiolar lumina are narrowed than *Svep1$^{+/+}$*

(highlighted in white)*.* Quantification shows an increased average of number of club cells

(SCGB1A1) but not of ciliated cells (FOXJ1). (Average counts ± SEM; *t* test p < 0.05 n≤ 3).

Scale bars 200 µm or 100 µm

Figure 5

Figure 6. Enhanced FGF signaling in *Svep1*^{-/-} embryos contributes to smooth muscle

differentiation and branching defects and FGF9 rescues SVEP1 inhibition of branching.

(a) *In situ* hybridization demonstrates increased expression of *Fgfr2* in *Svep1*^{-/-} mutants. **(b)**

Relative transcript abundance of *Fgfr2* (and its isoforms *Fgf2b* and *Fgfr2c*), *Fgf10*, *Fgf9,* and

Kras showing increased expression in *Svep1*^{-/-} mutant embryos at E18.5. **(c)** *Svep1*^{+/+} and

Svep1^{-/-} lung explants from E12.5 embryos treated *ex vivo* with the FGF signaling inhibitor

SU5402 (2.5 µg/ml). Black arrows point to normal tip morphology in *Svep1*^{+/+} lungs. The black

arrowhead points to distal tip trifurcations that persist in lung explants from *Svep1*^{-/-} embryos.

SOX9 (green) and ACTA2 (purple) staining confirm the branching anomalies (highlighted in the

white dotted areas) in *Svep1*^{-/-} lung explants. White arrows mark normal bifurcations, white

arrowheads indicate trifurcations and distal budding; the red arrowhead highlights ectopic

budding not affected by the induction of FGF. **(d)** E12.5 lung explants treated with SU5402 (2.5

µg/ml) for 4 days demonstrate reduced ACTA2 localization in distal airways of *Svep1*^{-/-} lungs

compared to wild type lungs . **(e)** *Fgf9* mRNA expression in wild type (E14.5) and mutant lung

(E16.5). **(f)** Wild type E12.5 lung explants after 48 hours: untreated control (top left); treated with

SVEP1 alone (top right); FGF9 alone (bottom left); combination of SVEP1 and FGF9 (bottom

right). The combination restores a normal bud tip phenotype. (Average counts ± SEM; *t*-test p <

0.01 and <0.05, n >3). *Scale bars 100 µm (a, c, d, e).*

Figure 6

Figure 7. Saccular cellular and structural defects in *Svep1*$^{-/-}$ embryos at E18.5. (a)

Expression heatmaps of selected genes associated with alveolar AT1 and AT2 cells show that

Svep1$^{-/-}$ lungs have lower expression of AT1 genes compared to *Svep1*$^{+/+}$ lungs and higher

expression of AT2 associated genes. **(b)** Immunofluorescence of PDPN, HOPX, Pro-SFPC and

ABCA3 in the lungs of E18.5 *Svep 1*$^{+/+}$ and *Svep1*$^{-/-}$ embryos. Arrows indicate airway bud stalks.

(c) PDGRFA+ (green) staining at E18.5. Plots of the average number of PDGFRA+ cells

(average counts ± SEM; *t* test p < 0.05 n=3). **(d)** Immunofluorescence of the mesothelial

progenitor marker WT1 (red) with KI67 (green) staining indicating actively dividing WT1+

progenitors (white arrows). Plot of the number of WT1+ and WT+/KI67+ cells at E16.5 and

E18.5. *Svep1*$^{-/-}$ lungs have more WT1+ cells than *Svep1*$^{+/+}$ lungs at E18.5; however, the number

of WT1+ KI67 lungs is unchanged between *Svep1*$^{+/+}$ and *Svep1*$^{-/-}$ lungs. (**e**) labeling of

endothelial cells with CD31 illustrating the double layer of the saccular microvasculature in

normal lungs (arrowhead; and disorganized microvasculature in *Svep1*$^{-/-}$ lungs at E18.5. *Scale*

bars 250 µm (B); 100 µm (c,d) 5 µm (e,h). Scale bars 25 µm (c,e); 50 µm (b) 100 µm (c).

Figure 7

969 **Figure 8. Graphical summary o normal and *Svep1*$^{-/-}$ embryonic lung phenotypes during**
970 **branching morphogenesis and saccular formation.** (Created with BioRender©).
971

Figure 8