Uncoupling of the Diurnal Growth Program by Artificial Genome Relaxation in *Synechocystis* sp. PCC 6803

Anna Behle[1,3,*], Maximilian Dietsch[1,*], Louis Goldschmidt[2], Wandana Murugathas[1], David Brandt[4], Tobias Busche[4], Jörn Kalinowski[4], Oliver Ebenhöh[2], Ilka M. Axmann[1], Rainer Machné[1,2,#]

1: Institut f. Synthetische Mikrobiologie, and
2: Institut f. Quantitative u. Theoretische Biologie, Heinrich-Heine Universität Düsseldorf, Universitätsstrasse 1, 40225 Düsseldorf, Germany
3: Photanol B.V, Science Park 406, 1098 XH Amsterdam, The Netherlands
4: Centrum für Biotechnologie (CeBiTec), Universität Bielefeld, Universitätsstrasse 27, 33615 Bielefeld, Germany

* AB and MD contributed equally, # correspondance to RM, machne@hhu.de

Abstract

In cyanobacteria DNA supercoiling varies over the diurnal light/dark cycle and is integrated with temporal programs of transcription and replication. We manipulated DNA supercoiling in *Synechocystis* sp. PCC 6803 by CRISPRi-based knock-down of gyrase subunits and overexpression of topoisomerase I, and characterized the phenotypes. Cell division was blocked, most likely due to inhibition of genomic but not plasmid DNA replication. Cell growth continued to 4-5x of the wildtype cell volume, and metabolic flux was redirected towards glycogen in the topoI overexpression strain. Topoisomerase I induction initially lead to down-regulation of GC-rich and up-regulation of AT-rich genes. The response quickly bifurcated and four diurnal co-expression cohorts (dawn, noon, dusk and night) all responded differently, in part with a circadian ($\approx 24\,\mathrm{h}$) pattern. We suggest a model where energy- and gyrase-gated transcription of growth genes at the dark/light transition (dawn) generates DNA supercoiling which then directly facilitates DNA replication and initiates the diurnal transcriptome program.

Figure 1. DNA Supercoiling & Transcription: Homeostasis and *Twin-Domain* Models. **A**: Global homeostasis of supercoiling by direct feedback on expression of topoisomerases (GYR: Gyrase holoenzyme; TopoI: topoisomerase I) and GC-rich anabolic/growth genes and AT-rich catabolic and stress-response genes. **B**. Transcription-dependent supercoiling downstream (positive) and upstream (negative) of an RNA or DNA polymerase, widely known as the *twin-domain* model. Gyrase-activity downstream can prevent polymerase stalling and induce transcriptional bursts while Topoisomerase I activity upstream prevents R-loop formation. **C**: The torsional stress exerted by transcription can lead to long-distance cooperative and antagonistic effects, and gene order on the chromosome can underlie temporal programs of gene expression.

Introduction

In vivo, the DNA double helix exists in a torsionally strained and underwound state, often denoted as "negative DNA supercoiling". A homeostatic feedback system of DNA supercoiling is coupled to differential expression of large gene groups in many different bacterial species [1, 2, 3, 4, 5, 6]. The general picture is that DNA supercoiling is high during times of high metabolic flux, such as during exponential growth, and required to express rRNA and GC-rich growth genes and allow for DNA replication [7]. The ATP/ADP ratio has direct effects on DNA supercoiling [8, 9, 10]. Supercoiling would allow an analog modulation of transcription factor and polymerase binding [11] and thereby qualifies as a potential mechanism for the long known and recently re-discovered monotonous relation of rRNA and growth gene expression with growth rate[12, 13, 14]. However, the relation of RNA transcription and DNA replication to DNA supercoiling is mutual and complex [7]. According to the twin-domain model of transcription-dependent supercoiling, negative supercoiling accumulates upstream and positive supercoiling downstream of polymerases [15], leading to cooperative and antagonistic long-range effects between transcription loci [16]. Strong transcriptional activity requires downstream activity of gyrase to set the elongation rate and avoid polymerase stalling [17, 18, 19] and upstream activity of topoisomerase I (TopoI) to avoid R-loop formation and genome instability [20, 21]. Such cooperative long-range effects can underpin temporal expression programs; locally in the *leu* operon [22, 23] and globally as a spatio-temporal gradient along origin-terminus axis of the *Escherichia coli* (*E.coli*) genome [24].

In cyanobacteria, this system forms an integral part of the diurnal (light/dark cycles) changes in metabolism and transcription, and is integrated with the output of the cyanobacterial circadian clock [25, 3]. Supercoiling of chloroplast genomes, the endosymbiotic descendants of cyanobacteria, was observed to fluctuate with the diurnal light/dark (LD) cycle [26], and plants encode for gyrase enzymes [27]. Cyanobacteria themselves traverse through a well defined transcriptional program during diurnal LD cycles in several species [28, 29, 30, 31]. Mori and Johnson first suggested [32] that diurnal DNA supercoiling may be involved in the genome-wide nature of diurnal transcription in cyanobacteria. Genome compaction was found to fluctuate with LD cycles in *Synechococcus elongatus* PCC 7942 [33], and an endogenous plasmid showed diurnal fluctuations of DNA supercoiling [25]. The temporal transcriptome program during entrained circadian cycles correlated with plasmid supercoiling states and depended on gyrase activity [3]. In *Synechocystis sp.* PCC 6803 (hereafter abbreviated as *Synechocystis*), cold, heat and salt stress all lead to similar changes in the transcriptome and all were enhanced by treatment with the gyrase inhibitor novobiocin (NB) [4]. Gene groups with coherent response to stress and NB overlapped significantly with diurnal co-expression cohorts [34]. Supercoiling of the endogenous plasmid pCA2.4_M increased within $30\,\mathrm{min}$ after the transition to light phase, and continuously decreased during $12\,\mathrm{h}$ and further during a prolonged $24\,\mathrm{h}$ dark phase [35][1].

In this work, we manipulate DNA supercoiling in *Synechocystis* by inducible overexpression [36] and CRISPRi-based knock-down [37] of the key genes involved in modulation of supercoiling: topoisomerase I (gene: *topA*), and gyrase (subunit genes: *gyrA* and *gyrB*. We can confirm both, the homeostasis and transcription-dependent supercoiling models in *Synechocystis*. All manipulations that should decrease supercoiling lead to a strong pleiotropic phenotype, where cell division is blocked but cell volume growth continues. Especially *topA* over-expression induces overflow metabolism and glycogen production, and uncouples the diurnal transcription program. Metabolism and transcriptome appear to be locked in a state between late night and early day.

Results and Discussion

Artificial Genome Relaxation Blocks Division but not Growth

Manipulation of Gyrase and Topoisomerase Expression. To study DNA supercoiling in *Synechocystis*, we used the dCas9-mediated CRISPR-interference system [37] to repress (knock-down) transcription of topoisomerase I (gene *topA*, slr2058), and gyrase subunits *gyrA* (slr0417) and *gyrB* (sll2005), or *gyrA* and *gyrB* simultaneously, yielding strains named <gene>[KD]. Additionally, we constructed a tunable expression plasmid

---

[1]*** ref. [35] is a preprint by our group at bioRxiv: https://doi.org/10.1101/2021.07.26.453679 ***

Figure 2. Batch Culture Endpoint Measurements. Overexpression and knock-down strains of this study where grown for 5 days in BG11 medium supplemented with all required antibiotics, and all inducers for the plasmid constructs in each experiment (100 ng/mL aTc, 1 mM L-rhamnose). **A:** The optical density at 750 nm (OD_{750}) was measured daily and cell dry weight (CDW) determined directly after the last measurement on day 5. **B:** The cell volume distribution was measured daily in the CASY cell counter and plotted as a gray-scale gradient (black: more cells at this volume). **C:** Absorption spectra after the harvest on day 5. See Figure S1B for spectra at inoculation time. The absorption at 750 nm was subtracted from each spectrum. **D:** ATP and ATP+ADP contents at harvest time were determined by a luciferase-based assay, and boxplots of six technical replicates (3 samples and 2 measurements) are shown. **E:** Glycogen content at harvest time was determined by a colorimetric assay after harvest, and boxplots of 18 technical replicates (3 samples, each measured 3x in 2 assays) are shown.

pSNDY [36] where a copy of the native *Synechocystis topA* is under the control of a rhamnose-inducible promoter, strain topA^{OX} (Tab. 1). All six strains were induced with anhydro-tetracycline (aTc) and rhamnose and cultured for five days, then harvested for quantification of cell dry weight, ATP, glycogen, and plasmid supercoiling. Reverse transcription quantitative PCR (RT-qPCR) verified the functionality of our inducible genetic constructs (Fig. S1A). All knockdown strains showed an abundance reduction to 8 %–15 % of the wild-type level, and *topA* induction was ≈30 fold. We also measured all three transcripts in all strains and observed compensatory upregulation and downregulation of the non-manipulated supercoiling enzymes, a first verification of the homeostatic control model in *Synechocystis*.

Cell Volume Growth, Adenosine and Glycogen Content. Initially, all cultures showed comparable growth (Fig. 2A). After three days all strains except topA^{KD} grew slower than the EVC; and topA^{OX} showed the

strongest growth defect. The cell dry weight (CDW) at harvest time correlated with the final OD_{750} of the cultures (Fig. 2A), but was relatively higher for the EVC and topA[KD] strains. Cell volume distributions of the EVC and topA[KD] strains showed a transient small increase ($\approx 10\%$) on the first day of cultivation and were stable thereafter (Fig. 2B). In contrast, cell volumes of the gyr[KD] and topA[OX] strains increased over time, from $4\,fL$–$5\,fL$ to $12\,fL$–$15\,fL$ after four days of cultivation. Total cell numbers increased only slightly. Thus, strains where gyrase subunits were knocked down or topoisomerase I overexpressed showed inhibition of cell division but not of cell growth. Pigmentation was strongly affected. Cultures of the topA[OX] strain appeared pale and gyr[KD] strains blue, compared to their uninduced state and to EVC. Absorption spectra (Fig. 2C, S1B) confirmed an overall decrease of all pigments in topA[OX]. The gyr[KD] strains showed a stronger decrease at chlorophyll-specific wavelengths than at phycocyanin-specific wavelengths, explaining their blue appearance.

All knock-down strains showed glycogen levels similar to the EVC, with 25 % of the total CDW (Fig. 2D), which is consistent with literature values for the wild type [38]. In contrast, topA[OX] showed more than twice as much glycogen with 55 % of CDW. Values of up to $60\,\%$ are reported for nitrogen-starved cells [39, 40]. The topA[OX] strain accumulated more than four times as much ATP+ADP as the EVC (Fig. 2E). gyrB[KD] and gyrAB[KD] accumulated about twice as much ATP+ADP as the EVC; topA[KD] and gyrA[KD] showed no difference to the EVC control. Thus, the strains repressing the ATPase subunit of gyrase, gyrB[KD] and gyrAB[KD], had slightly elevated levels of ATP+ADP.

Figure 3. Plasmid Supercoiling. **A:** Baseline-corrected electropherograms from CQ agarose gels (Fig. S1D) of samples from the cultures shown in Figure 2, taken at harvest time (5 d). At $20\,\mu g\,mL^{-1}$ CQ, originally (*in vivo*) more relaxed (rel.) plasmids migrate further (higher migration distance) than more negatively supercoiled (sc.) plasmids [35][*** 2 ***]. Two distinct plasmid topoisomer distributions can be distinguished in the EVC, and we assume the less far migrated bands to correspond to the longer plasmid pCA2.4_M, and the further migrated bands to the shorter plasmid pCB2.4_M. **B**: same as (A) but samples taken as a time series from the topA[OX] strain, induced with rhamnose at $0\,d$ (Fig. S2). **C:** as (B) but for a population of topoisomers from a larger plasmid which could be separated by running the gel longer than the gel analyzed in (A). We tentatively assign these bands to the endogenous plasmid pCC5.2_M, based on its relative migration distance.

Plasmid Hyper-Supercoiling in the topA[OX] Strain. Agarose gel electrophoresis in the presence of chloroquine (CQ) was used to analyse plasmid supercoiling at harvest time (5 d) of the batch cultures, and interpreted as outlined previously [35][2]. The gels show three sets of topoisomer bands (Fig. S1D). These likely stem from the three annotated small plasmids of *Synechocystis*, pCA2.4_M, pCB2.4_M and pCC5.2_M. Electropherograms of the two smaller plasmids indicate that only strains gyrA[KD] and gyrAB[KD] showed plasmid relaxation (Fig. 3A). In the gyrB[KD] and topA[OX] strains plasmids appeared to have a higher level of DNA supercoiling. We could not extract plasmids from the topA[KD] strain. Increased plasmid supercoiling in topA[OX] could results from a long-term adaptation and compensatory up-regulation of gyrase subunits. We thus tested plasmid supercoiling as a time series after inoculation in fresh medium with and without the inducer (Fig. S2). The gel

---

[2]*** ref. [35] is a preprint by our group at bioRxiv: `https://doi.org/10.1101/2021.07.26.453679` ***

run time was increased to also separate topoisomers of pCC5.2_M. All three plasmids were more relaxed 83
(less negative supercoiling) after 3 h of growth (Fig. 3B,C). Already after 8 h the trend had reversed, and at 84
20 h plasmids were more supercoiled than at time 0 h and in the uninduced control time series (Fig. S2E). 85
Then plasmids became further supercoiled to an extent where topoisomers were not separable anymore. 86

Reduction of rRNA Abundances in gyr^{KD} Strains.   To test effects on transcription, we selected three 87
strains, gyrA^{KD}, gyrB^{KD} and topA^{OX}, and the EVC for RNA-seq analysis. Strains were grown in triplicate and 88
harvested for flow cytometry, RNA extraction and RNA-seq analysis 5 d after induction and 3 d after a culture 89
dilution step (Fig. S3A). Flow cytometry confirms the growth phenotype (Fig. 4A, S4): forward scatter (FSC) 90
was increased in all strains, and most in topA^{OX}. Side scatter (SSC) additionally revealed two cell populations 91
in topA^{OX} which could reflect 8-shaped cells in division. Total nucleic acid content increased with cell size. 92
Total RNA composition and the relative abundances of rRNA and mRNA were analyzed by cappillary gel 93
electrophoresis (Fig. S3B,C, S5). Ribosomal RNA species were strongly reduced in the gyrA^{KD} and gyrB^{KD} 94
strains and less reduced in topA^{OX} (Fig. 4B), except for the 600 bp fragment of the 23S rRNA (Fig. S6C). 95

Figure 4. rRNA and mRNA Abundances in Induced Strains. **A:** Flow Cytometry confirms the cell growth
phenotype. The natural logarithms of forward scatter, side scatter and nucleic acid stain Syto9 were calculated
and 2D distributions plotted as contour plots; see Figure S4 for all data and details. **B:** Relative abundances
of the unfragmented 23S (B) and the 16S (C) rRNA species were calculated from the electropherograms
of capillary gel electrophoresis of extracted RNA (Fig. S5). Figure S6 shows all rRNA species. Mean and
standard deviation of the three replicates are indicated as error bars.**C & D:** Relative expression changes of
coding genes were derived as the $\log_2$ ratio of RPKM normalized read-counts in the induced over-expression
or knock-down strains to values in the control strain (EVC) and compared between the three different strains
by 2D histograms (yellow: highest and purple: lowest local density of genes). The Pearson correlations (r)
are indicated in the bottom right corner. **C**: gyrA^{KD} (y-axis) *vs.* gyrB^{KD} (x-axis) strains. **D**: gyrA^{KD} (y-axis)
vs. topA^{OX} (x-axis). The induction/repression and the homeostatic responses of *gyrA*, *gyrB* and *topA* are
highlighted by arrows from the origin to indicate the direction of change. The gene *sll1941* is a homolog or
paralog of *gyrA* (blue diamond) and its transcript showed no response in either experiment.

Consistent Changes in mRNA Abundances & Homeostatic Regulation of Supercoiling Enzymes.   The 96
same RNA samples were further processed (rRNA species depleted) and sequenced on the Illumina platform, 97
and transcript abundances relative to the EVC evaluated with DESeq2 [41]. All strains showed overall similar 98
expression changes, but the extent was lower in topA^{OX} (Fig. 4C,D). However, this difference could also 99
just reflect normalization effects by the decreased rRNA content in the gyrase knock-downs. In all strains 100
the targeted manipulation was still observable at harvest times (arrows in Fig. 4C,D), *i.e.*, *gyrA* transcripts 101
were reduced in gyrA^{KD}, *gyrB* transcripts in gyrB^{KD} and *topA* transcripts were increased in topA^{OX}. The 102
non-manipulated genes showed the compensatory response expected from homeostatic regulation, *i.e.*, *topA* 103
was repressed in both gyr^{KD} strains, and all non-manipulated gyrase subunits were induced in all experiments. 104

In contrast, the *sll1941* gene, annotated either as a second gyrase A subunit or as the topoisomerase IV ParC subunit, showed no response in either experiment.

Overexpression of *topA* Uncouples Diurnal Co-expression Cohorts

Figure 5. Pulsed Induction in Continuous Culture. **A:** Photobioreactor growth of the topA^{OX} strain (1 L BG11 medium, 0.5% CO_2, illumination $\approx$90 µmol m^{-2} s^{-1} per OD$_{750}$). Optical density was recorded online (OD$_\lambda$) and post-calibrated to offline OD$_{750}$. The arrows indicate inoc.: inoculation; cnt.: onset of continuous culture (rate $\phi = 0.01$ h^{-1}); IND.: induction of *topA* by pulse-addition of rhamnose to 2 mM (0.33 g L^{-1}) at time 0 d; and batch: switch-off of dilution. The dashed black line is the theoretical wash-out curve of rhamnose (g L^{-1}). Cell dry weight (CDW, g L^{-1}, red) and glycogen content (g L^{-1}, blue) of the culture were measured at the indicated times (points), and LOESS regressions are shown (solid lines) with 95% confidence intervals (dashed lines). **B:** Cell numbers (blue points) and volume distributions (gray scale) were recorded daily, and at higher resolution after induction, with the CASY cell counter. The peaks of the cell volume distributions are shown as yellow points. LOESS regressions with 95% confidence intervals are shown as lines. One outlier of the CASY measurement (x) was due to cell lysis during a washing step and was not included for regression. **C:** growth rates μ were calculated by local (piecewise) linear regressions of the OD$_\lambda$ (A), and cell count (B) and total cell volume (D) measurements and subtraction of the culture dilution rate (Fig. S9). **D:** The total cell volume (V_{total}) was calculated by integrating the single cell volume distributions in (B), and the CDW density was calculated by dividing the OD$_\lambda$ signal, calibrated to the CDW measurements (A, Fig. S8C), by V_{total}.

Transient Increase in Cell Volume and Density. To study the dynamic response to transient *topA* induction, the topA^{OX} strain was grown in a Lambda Minifor bioreactor (Fig. S7) with continuous (online) monitoring of turbidity (OD$_\lambda$, Fig . S8A,B). Continuous culture dilution was initiated at OD$_\lambda \approx 2.9$ and with dilution rate $\phi \approx 0.24$ d^{-1}. The culture stabilized around OD$_\lambda \approx 2.7$. Notably a subtle ≈ 24 h pattern of OD$_\lambda$ was observed in both, batch and pre-induction continuous growth phases. Then rhamnose was injected to 2 mM to induce overexpression of *topA*. The *topA* transcript was upregulated to $\approx$45-fold over the pre-induction level within 4 h, as measured by RT-qPCR and confirmed by RNA-seq (Fig. S11) and decreased slowly over the course of the experiment. The OD$_\lambda$ initially increased for 1 d post-induction, then slowly decreased. Cell dry weight (CDW) measurements were noisy but matched the OD$_\lambda$ signal over the sampled period

(Fig. 5A, S8C). In contrast, cell numbers started to decrease immediately, and cell volumes increased (Fig. 117
5B). We calculated growth rates of OD_λ, cell numbers and the total cell volume (Fig. 5C, S9). Cell division 118
was not completely blocked but severely reduced to a division time of $\approx 10\,d$ ($\mu_{count} \approx 0.07\,d^{-1}$). Total cell 119
volume growth was much less affected and remained stable ($\mu_{volume} \approx 0.18\,d^{-1}$) throughout continuous 120
culture operation until $12\,d$ post-induction. Thus, artificial *topA* overexpression blocked cell division but not cell 121
volume growth. OD_λ growth remained highest ($\mu_{OD} \approx 0.23\,d^{-1}$) and stable over the first $5\,d–6\,d$. In parallel, 122
glycogen content increased to about $35\,\%–40\,\%$ of the CDW (Fig. 5A). We further noticed that sampled cells 123
started to sediment much faster, indicating increased intracellular density. By calibrating the OD_λ signal to 124
the CDW measurements (Fig. S8C) and dividing by the total cell volume we can estimate a CDW density 125
and this value also increased over time from 0.3 to $0.5\,g_{DCW}/mL_{cell}$ (Fig. 5D). This range is consistent with 126
data from *E. coli* [42, 43]. However, the CDW per OD_{750} was relatively lower for the enlarged strains in the 127
endpoint measurement (Fig. 2A), and thus, the calibration to OD_λ may overestimate true CDW density. The 128
enlarged cells also became increasingly fragile: in the CASY cell counter data a small population of varying 129
intensity appeared at $<2\,fL$. This peak was highest at $7\,d$ (outlier x in Fig. 5B), where cells were lysed during 130
centrifugation in a washing step. The washing step was skipped thereafter, and the peak of small cells (dead 131
or fragmented) remained small but increased towards the end of the continuous culture. 132

Maximal cell volumes $>20\,fL$ were reached $10\,d–15\,d$ post-induction. From day 14 a population of smaller 133
cells, $\approx 7.5\,fL$, appeared. On $16\,d$ this population was the majority, and cell volume further decreased to $5\,fL$. 134
Cell pigmentation recovered and the culture appeared greener again. We then switched off dilution, and the 135
culture resumed growth, although at lower growth rates than pre-induction. 136

Upregulation of Plasmid & Growth Genes, Downregulation of Photosynthesis Genes. Samples for 137
total RNA analysis and RNA-seq were taken $1\,d$ and $0.5\,h$ before induction, and then over the next $25\,d$ in 138
decreasing temporal resolution to roughly capture three time-scales of the response. Coding gene transcript 139
abundances were calculated, the resulting time series clustered (Fig. 6A, S12), clusters sorted along significant 140
overlaps with diurnal cohorts (Fig. 6C), and functional annotation enrichments (`CyanoBase` "categories": Fig. 141
6B, S14A; gene ontology: Fig. S15) calculated. Cluster 2 (red) comprises the majority of the ribosomal protein 142
category, enzymes of RNA and DNA synthesis, amino acid biosynthesis and the ATP synthase. Transcript 143
abundances increased slightly over the three days after induction, with a notable circadian ($\approx 24\,h$) pattern. 144
Cluster 3 (yellow) comprises the majority of photosynthesis-related genes, including the two photosystems. Its 145
transcript abundances decreased continuously. The remaining clusters showed less clear functional profiles. 146
Clusters 1 (green) and 4 (blue) were enriched in genes with unknown function, and cluster 4 strongly enriched 147
with transposase genes. Gene Ontology analysis additionally shows enrichments with 7 of 8 genes annotated 148
with "DNA polymerase activity" and 60 "DNA binding"proteins (Fig. S15). Both clusters 1 and 4 showed 149
decreased transcript abundances on the first day post-induction, then only cluster 4 increased steeply. Cluster 150
1 transcripts showed a subtle circadian pattern, peaking anti-phase to cluster 2 transcripts. Cluster 6 (cyan) 151
showed a similar profile to cluster 4 and comprises transcripts that showed the strongest abundance increase. 152
Both clusters 4 and 6, were strongly enriched for plasmid-encoded genes (Fig. S16), and both also contained 153
significant fractions of the ribosomal protein category. Cluster 5 (gray) showed decreased abundances but had 154
the least change over time, was slightly enriched with "hypothetical", "Hydrogenase" and "plasma membrane" 155
gene annotations. 156

Clusters Reflect Co-expression Cohorts. Next, we compared the time series clustering from topA[OX] 157
and the endpoint transcript abundances from the topA[OX] and gyr[KD] strains with previously characterized 158
co-expression cohorts (Fig. 6D). Prakash *et al.* clustered genes into three groups by their response to salt, 159
heat and cold stress, each with or without the gyrase inhibitor novobiocin (NB) [4]. Zavrel *et al.* measured 160
protein abundances at different growth rates in constant light conditions and presented 7 clusters [14] which 161
we summarized into cohorts upregulated or downregulated with growth rate, with a complex response and 162
stable proteins without growth rate-dependance. Saha et al. and Lehmann *et al.* analyzed transcriptome 163
time series over light-dark cycles [44, 34]. Saha *et al.* did not cluster their data, thus we clustered the data 164
into 5 clusters and assigned them names that correspond to their order of expression during the diurnal 165
light-dark cycle (Fig. S17). Clusters 2 (red) and 3 (yellow) show the clearest functional profiles, and are 166

Figure 6. Co-Expressed Gene Cohorts. **A:** Cluster medians (colored solid lines) and 25%/75% quantiles (transparent ranges) of relative transcript abundances (rel. ab.: $\log_2$ ratio (TPM) to the mean TPM of two pre-induction samples); points indicate sampling times (ticks on upper axis). Cluster labels and sizes are indicated in the legend. **B:** Sorted enrichment profile of co-expression cohorts with the `CyanoBase` "categories" gene annotation; black field: $p \leq 10^{-10}$; white text: $p \leq 10^{-5}$, row sorting and filter: $p_{sort} \leq 0.01$. Figure S14 shows unfiltered results for all categories. Numbers are counts of genes in both categories. **C, top:** Cluster medians as in (A) but zoomed in on the first five hours after induction and without - the quantile ranges. **C, bottom:** Cluster enrichment profile (black: $p \leq 10^{-10}$; white text: $p \leq 10^{-5}$) with genes upregulated (up), downregulated (down) or without change (nc) immediately (5 min–20 min) after induction. **D:**. Enrichment profiles (gray scale) with other published gene classifications (see text) and t-value profiles (red-blue scale) of clusters in the end-point transcriptome experiments. Blue: $t > 0$, red: $t < 0$; $p_{min} = 10^{-10}$, white text: $p \leq 10^{-5}$. The text indicates total counts as in (B), or t-values in t-test profiles. **E:** GC-content profiles around start codons (ATG, top panel) and transcription sites (TSS, bottom panel) from S23. GC-content was calculated in 66 bp windows at each position. Point sizes scale with $-\log_2(p)$ from local motif enrichment tests (filled points) and deprivation (open circles) tests, and the minimal p-values are indicated in the legends. These significance points are only shown every 10^{th} (top) or 3^{rd} position (bottom).

found co-expressed with opposite patterns in all tested experiments, and at dawn and noon, respectively, in the diurnal datasets. Cluster 2 was most enriched with transcripts that showed a mixed response to the applied stress conditions (`Stress+NB`) and with genes whose protein abundances correlated positively with growth rate (`GROWTH`). Cluster 3 was most enriched with transcripts downregulated in stress and whose protein abudances correlated negatively with growth rate. In the diurnal data set cluster 2 transcript abundances peaked in early day (dawn) and were followed by a peak of cluster 3 transcripts at noon. These clusters likely reflect biological co-expression cohorts and from hereon we denote cluster 2 as cohort `RB/dawn` (ribosomal proteins and biosynthesis) and cluster 3 as `PS/noon` (photosynthesis), according to their annotation and experiment enrichment profiles. Cluster 1 is enriched with proteins of unknown function. In both, stress and growth experiments, we found only enrichments in the NA classes, ie. the original papers did not report values. However, in the diurnal data set we found enrichment with genes clusters that peak at noon or after noon. We

denote this cohort as UK/dusk (unknown function, expressed at dusk). The remaining clusters 5, 4 and 6 all 178
show weak functional enrichments. Only cluster 4 is enriched with transposase genes, DNA polymerase and 179
other DNA binding proteins. All of them, and strongest in cluster 4, show weak enrichment with transcript 180
cohorts that peak during the dark phase. Cluster 6 is enriched in the late night cohort in the diurnal data set, 181
and further shows some overlaps in enrichment with the RB/dawn cohort, thus closing the (diurnal) cycle of 182
expression. We tentatively label cluster 4 as the DNA/night cohort and cluster 6 as plasmids. 183

To compare the time series with the endpoint transcriptome data, we calculated t-test profiles of all clusters 184
(red/blue squares in Fig. 6D and S13). RB/dawn, PS/noon and UK/dusk showed consistent behaviour in all 185
strains, although at lower t-values for the gyrase knock-downs. DNA/night and plasmids, both enriched with 186
plasmid genes, were upregulated only in topA^{OX} but not in the gyr^{KD} strains. Transcripts of the large cluster 5, 187
which showed the weakest response in the time series, were upregulated only in the gyr^{KD} strains. Likewise, 188
the diurnal co-expression cohorts from ref. [44] (Fig. S17)) all show subtle but notable differences between 189
the strains, only PS/noon was consistently downregulated. 190

Uncoupling the Diurnal Program at Dawn. Genes with supercoiling-activated transcription were found to 191
be GC-rich in both upstream non-coding and coding regions of several species, and *vice versa*, supercoiling- 192
repressed genes are AT-rich [1, 3, 45, 6]. Thus, we aligned nucleotide sequences of all genes from the main 193
chromosome at their start codons (ATG), calculated average nucleotide content in moving windows for each 194
cluster, and performed a statistical enrichment test (cumulative hypergeometric distribution) at each position 195
(Fig. 6E). The up-regulated RB/dawn cohort and the down-regulated PS/noon cohort are GC-rich downstream 196
and upstream of the start codon. The UK/dusk (downregulated) and DNA/night cohorts (upregulated) are 197
AT-rich. Zooming in to the immediate response after induction shows that the response of the co-expression 198
cohorts switches within the first h post-induction (Fig. 6C). The immediate response is consistent with data 199
from other species. Transcript abundances of both GC-rich cohorts, RB/dawn and PS/noon, decreased, while 200
those of the AT-rich UK/dusk and DNA/night cohorts increased. Already 1 h post-induction, the two GC-rich 201
and two AT-rich cohorts have bifurcated. Transcript abundances of RB/dawn increased while those of PS/noon 202
continue to decrease. Similary, the AT-rich UK/dusk transcripts now decreased while DNA/night transcripts 203
continued to increase in abundance. 204

Transcription Unit Promoter Structure: A GC-rich Discriminator. To analyze actual promoter structures 205
we mapped the coding genes on previously described transcription units (TU) [46], calculated average 206
temporal abundance profiles of TUs, and clustered TU profiles by k-means using the average profiles of the 207
gene-based clustering as cluster centers (Fig. S23). TU were then aligned at their transcription start sites 208
(TSS) and again average nucleotide content profiles of clustered TU calculated (Fig. 6E). Downstream of 209
the TSS the general differences in GC/AT content were similar to those of start codon-aligned coding genes, 210
albeit with lower stastistical power. TSS alignment reveals a distinct GC-rich peak at ca. -50 bp. This peak 211
corresponded better to the mid-term trend of transcript abundances. Cohorts that were upregulated within the 212
first 3 d post-induction (DNA/night and RB/dawn) had a higher, and downregulated cohorts (PS and UK/dusk) 213
had the lowest GC-content at this putative "discriminator" position. 214

Genes of Interest. In Figures S18–S22, we analyze transcript abundances of a few specific gene sets: 215
circadian clock genes (*kai*), response regulators, sigma factors, and photosynthesis and metabolic genes. 216
However, these patterns are hard to interpret, due to the unknown function of upregulated genes and differential 217
regulation of paralogs and enzyme complex subunits. In short: most *kai* genes were down-regulated, while 218
kaiC3 is upregulated with RB/dawn, followed by *kaiB2* and *kaiC2* from 3 d (Fig. 7A, S18A). KaiC3 is required 219
for chemoheterotrophic growth in constant darkness [47]. The response regulator *rpaB* is downregulated, and 220
rpaA slightly upregulated (Fig. S18B). Of the tested regulators, only *pmgA* is strongly upregulated over the first 221
three days, with the diurnal pattern of the RB/dawn cohort. Sigma factors of unknown function [48], *sigH→segI*, 222
were upregulated until 3 d; the stress factor *sigB* is up-regulated only at 10 d (Fig. S18C), where cells are 223
enlarged. Photosystem, phycobilisome, and carboxysome genes are predominantly downregulated (Fig. S19– 224
S20). In contrast, RuBisCo and genes of the carbon concentrating mechanisms are predominantly upregulated 225

Figure 7. Explaining the Phenotype? A: Relative transcript abundances of the Kai circadian clock genes in the topA^{OX} time series, only *kaiC3* is upregulated over the first three days post-induction. The gray background shows the 92.5 % and 7.5 % quantiles of all data as a reference. **B:** Flow cytometry of batch culture endpoint samples, as Fig. 4A but for nucleic acid content (Syto9 marker) *vs.* the side scatter signal, an indicator of cell morphlgical features. **C:** Relative transcript abundances (as in (A)) of genes from the plasmid pCC5.2_M. **D:** RNA and DNA extraction yields in the topA^{OX} time series experiment, divided by the OD$_\lambda$ signal calibrated to CDW. OD$_\lambda$ and cell volume distributions (Fig. 5A,B) are shown (gray) as reference. Lines show a LOESS regression with 95% confidence interval. **E:** Plasmid extraction yields in the plasmid supercoiling time series (Fig. 3) from the induced and uninduced cultures. **F:** Cell volume and count development after re-induction of of cells isolated after harvest from the topA^{OX} time series experiment in fresh BG11 medium.

over the first three days (Fig. S21). The glycogen degrading enzyme *glgP1* was strongly downregulated; glucose-1-phosphate adenylyltransferase *glgC* was upregulated, and glycogen debranching enzymes *glgX*/*X2* peaked at 3 d (Fig. S22A). NAD synthesis genes are generally upregulated, and we find strong differential response of certain subunits of the NADH dehydrogenase (Fig. S22B,C). Together with the upregulation of the redox and photomixotrophic growth regulator *pgmA* [49], these responses point to changes in redox metabolism. The thioredoxin *trxM1* is strongly down-regulated, and *trxA* has only a short and low peak at 2 d–3 d (Fig. S22D).

Reversible Inhibition of DNA Replication Explains the Growth Phenotype. Manipulation of DNA super-coiling has highly pleiotropic effects, and can also interfere with DNA replication and lead to DNA damage [50]. Here, we observed a block of division while cell growth continued. Could an inhibition of genome replication be involved? In endpoint RNA-seq experiments relative rRNA abundances were decreased but the total nucleic acid content (Syto-9 stain) increased with cell size and morphology (Fig. 4A, 7B, S4). To differentiate nucleic acids, we extracted RNA and DNA from the topA^{OX} time series experiment (Fig. 7C). DNA extraction yields

per CDW were lower while RNA yield was constant over the three days post-induction. From 4 d post-induction ₂₃₉
DNA yield increased steeply ans surpassed RNA yields. This could be due to increased genome or plasmid ₂₄₀
copy numbers. The transcript abundances from all plasmid-derived genes became high at this time (Fig. 7D, ₂₄₁
S16). Plasmid DNA extraction yields from the topA^{OX} plasmid supercoiling time series increased strongly ₂₄₂
in the induced but not the control culture, but this bias was lost when treating with the T5 exonuclease to ₂₄₃
remove all linear or nicked circular DNA (Fig. 7E, S2B,C). Large plasmids and the genome are easily nicked ₂₄₄
during the extraction procedure. Thus, the most parsimonious explanation for the division block phenotype is ₂₄₅
impaired DNA replication of the genome, while plasmid copy numbers increased during the late stages of ₂₄₆
cell volume growth. A block of genome replication could be due to permanent DNA damage. The volume ₂₄₇
growth and division block phenotype was reversible. All strains recovered readily when re-inoculated in fresh ₂₄₈
medium without the inducer (Fig. S1C). In the continuous culture experiment, cells recovered as the inducer ₂₄₉
was washed out (Fig. 5A). Re-inoculation with and without inducer showed that in some cells our constructs ₂₅₀
remained intact and induced cells again grew in volume, while other cells were refractive to induction, and ₂₅₁
overgrew the division-blocked cells (Fig. 7E, S10). Thus, we conclude that the putative block in genome DNA ₂₅₂
replication was not due to irreversible and lethal genome damage. ₂₅₃

Conclusion ₂₅₄

A

B

Figure 8. Summary and Model. A: Summary tables of physiological characteristics of the tested strains
(top) and the co-expression cohorts observed after topA^{OX} induction (bottom). The top table indicates
changes in cell volume, ATP+ADP, glycogen, rRNA content and plasmid supercoiling. The bottom table
summarizes enriched properties of the co-expression cohorts (clusters): the initial and late response of
transcript abundances, functional enrichments, overall GC content and the presence of a putative GC-rich
discriminator at -50 bp upstream of the TSS. **B:** Development of an Integrative Model: at dawn energy from
photosynthesis becomes available, allowing for transcription of GC-rich growth genes (RB/dawn), gyrase
activity downstream of these genes can gate their transcription, *e.g.* tune bursting frequency [51] or set the
elongation rate [52]. This is inhibited in the gyrase knock-downs and not inhibited or even enhanced through
gyrase upregulation in the *topA* over-expression experiments. The combination of transcription and gyrase
activity leads to genome-wide increase in negative supercoiling. The accumulated torsional strain can then
be channeled into transcription of the second GC-rich cohort, PS/noon. This latter process is inhibited in all
strains. In normal conditions topoisomerase I could then be involved in resolving R-loops [20] that may have
been generated by the dawn/noon program. During night-metabolism ppGpp signaling [53] activates the night
transcription program, at low levels of transcription [31], and supercoiling globally decreases. Clock-mediated
glycogen mobilization at the end of the night [54] could kick-start the RB/dawn program already before onset
of light [44].

We presented a comprehensive characterization of DNA supercoiling effects in *Synechocystis*. Several ₂₅₅

models of the role of this regulatory system in bacterial cell biology can be confirmed in this species. The enzymes directly responsible for supercoiling are under homeostatic control [55]. Increased gyrase levels and transcription conspire to hyper-supercoil plasmids in the topA^{OX} strain, in agreement with the twin-domain model of transcription-dependent supercoiling [15, 56]. Differential transcription of GC-rich and AT-rich genes [1, 3, 45, 6] was observed immediately (5 min–20 min) after induction of *topA* over-expression. Already after 60 min the pattern had changed. The compensatory upregulation of gyrase and other regulatory mechanisms, such as the ppGpp mediated repression of day time transcription [53], may underlie this quick bifurcation of the response. A GC-rich discriminator -50 bp upstream of TSS could be involved. Large gene cohorts developed similarly over the first three days post-induction, and these revealed a tight link to the diurnal transcription program, consistent with results from *Synechococcus elongatus* PCC 7942 [3].

Cooperative and antagonistic long-range effects of transcription *via* local accumulation and diffusion of DNA torsional strain [16] underlies temporal expression sequences in operons [22, 23], and genome-wide along the origin-terminus axis during *E. coli* growth phases [24]. We suggest a simple model of supercoiling-mediated cooperativity (Fig. 8), where energy-driven transcription of ribosomal RNA and growth genes around dawn (RB/dawn) requires downstream gyrase activity. This process would generate the supercoiling required for both, transcription of the GC-rich PS/noon genes and DNA replication between dawn and noon [57]. This coupling is affected differently by our interventions, and reflected in the subtly different response of diurnal cohorts in the three strains (Fig. 6B). In the gyrase knock-downs it is directly inhibited, resulting in drastic reduction of rRNA abundances (Fig. 4B). In the topo I overexpression experiments the gyrase genes were upregulated and downstream gyrase activity could locally be enhanced. Globally, the high abundance of topo I would constantly remove accumulated negative supercoiling. In all strains, the supercoiling required for the onset of the program beyond dawn gene expression could not be generated: DNA replication is blocked, and transcripts of the PS/noon and UK/dusk cohorts are downregulated. The overall decreased genomic supercoiling level results in up-regulation of night-specific transcripts, including *kaiC3*, a homolog of the circadian clock gene and required for heterotrophic growth in constant darkness [47]. The cells overall appear "stuck" between night and dawn states. Next, our strains and interventions should be studied in this natural diurnal context, that is, in cultures grown in regular day/light cycles, and interference initiated at different times of the cycle. Our tunable induction/repression constructs could further be combined in one strain, topA^{OX}+gyr^{KD}, to avoid adaptive effects of gyrase overexpression. This would allow better and tunable external control of DNA supercoiling and the transcriptional response. Finally, we note that topoisomerase I overexpression caused an increase of cellular glycogen content of up to 60 % of the cell dry weight. This phenotype could be exploited in biotechnological applications, *e.g.* for synthesis of high value products or directly as a fermentation substrate for yeast [40].

Data Availability

The clustering and time series data from the topA^{OX} strain (both as raw abundances in TPM and as the log2 ratios to the mean of two pre-induction values, as plotted in this manuscript), and endpoint measurements (log2 ratio of abundances in the gyrA^{KD}, gyrB^{KD} and topA^{OX} strains to the EVC strain) are available as Datatable_S1.tsv.

*** The raw sequencing reads will be made available at an appropriate facility (SRA) upon publication. ***

Funding

RM was funded by the *Deutsche Forschungsgemeinschaft* (DFG), grants AX 84/4-1 & STA 850/30-1 (COILseq) and EXC-2048/1–project ID 390686111 (CEPLAS). MD, DB and TB were financially supported by the CLIB Competence Centre Biotechnology (CKB) funded by the European Regional Development Fund ERDF (EFRE-0300096 and EFRE-0300095).

Acknowledgments

The parental strain *Synechocystis* sp. PCC 6803, encoding aTc-inducible dCas9, and the gyrB^{KD} strain were a gift from P. Hudson and L. Yao. We are grateful to Nic Schmelling for critical discussion of the manuscript.

Materials and Methods

Strains and Plasmids

The *Synechocystis* parent strain used for all genetic alterations was kindly provided by Lun Yao and Paul Hudson, and based on a strain they had obtained from Dr. Martin Fulda (Göttingen, Germany). It contains a TetR cassette, as well as dCas9 under the promoter PL$_{22}$, inducible with anhydrotetracycline (aTc), at the genomic insertion site psbA1. Construction of this strain was described in ref. [37]. For overexpression of genes in *Synechocystis*, relevant genes were directly PCR-amplified from the *Synechocystis* genome, fused to the rhamnose-inducible promoter [36] *via* overlap extension PCR, and integrated into a modified variant of the conjugative vector pSHDY containing a nourseothricin resistance instead of spectinomycin, termed pSNDY. The vector backbone also contained the activator *rhaS* from [36]. Lun Yao and Paul Hudson further provided a strain with *gyrB*-targeting sgRNA. For the construction of the additional sgRNA constructs, sgRNA sequences were designed using CHOPCHOP [58], constructed *via* overlap extension PCR and integrated into the vector designed by [37] (Addgene #73224), which inserts into the slr0230 site of the *Synechocystis* genome. Integrative sgRNA plasmids were integrated *via* transformation. Briefly, 10 mL of exponentially grown culture was concentrated to 250 µL, 1-2 µg of pure plasmid was added and the mixture was incubated up to 5 hours before plating the entire mixture on BG-11 plates. After drying the plates, agar was underlaid with 300 µL of 1 mg/mL kanamycin stock using a sterile spatula, thereby forming a diffusion gradient. After 1-2 weeks of incubation at 30 °C with the lid facing upward, isolated green colonies were carefully transferred to a fresh plate. Over time, positive clones were gradually shifted to higher concentrations of kanamycin (*i.e.*, 4, 8, 12, 20, 40 µg/mL final concentration in the plate). Complete segregation of mutants was ensured *via* colony PCR. Replicative vectors were introduced into the dCas9 background strain *via* conjugation as described (dx.doi.org/10.17504/protocols.io.ftpbnmn). Clones were selected using nourseothricin (Jena Bioscience, #AB-102L) at a final concentration of 50 µg/mL and verified *via* colony PCR.

Culture Conditions

Batch Culture Conditions. Plates were freshly streaked before liquid cultivation. In the case of noursethricin, liquid cultures were maintained with 25 µg/mL while solid media was supplemented with 50 µg/mL. Spectinomycin and Kanamycin were supplemented at concentrations of 20 µg/mL and 25 µg/mL, respectively. For pre-culturing and growth experiments, *Synechocystis* strains were cultivated in BG11 medium [59]. Cultivation was performed at $30\,°C$ with 150 rpm shaking and continuous illumination of $\approx 80\,\mathrm{µmol\,m^{-2}\,s^{-1}}$ (16% setting in Infors HT). Aeration was ensured by continuous shaking and CO_2 enriched air (0.5 %) in an Infors HT multitron chamber. Pre-culturing was performed in 100 mL baffle-free Erlenmeyer shaking flasks with 20 mL cell suspension for three days. After adjusting all different strains to the same $OD_{750}750 \approx 0.4$, growth experiments were performed after one additional day of pre-culturing. For this, 30 mL cultures were incubated in Erlenmeyer shaking flasks for 5 days with a start $OD_{750} \approx 0.25$ in biological triplicates.

Continuous Culture, Online Measurements and Calculations

Continuous culture was performed in a Lambda Photobioreactor and with additional measurement and control devices (Fig. S7) in BG11 medium, supplemented with the required antibiotics, at culture volume $V_\ell = 1\,L$, aeration with $1\,L\,min^{-1}$ of CO_2-enriched (0.5 %) air, agitation by the Lambda fish-tail mixing system at $5\,Hz$, temperature control at $30\,°C$, and pH 8, with $0.5\,M$ NaOH and $0.5\,M$ H_2SO_4 as correction solutions. After equilibration to these conditions the reactor was inoculated to a start $OD_{750} \approx 0.5$, from $100\,mL$ pre-culture.

White light from the Lambda LUMO module was initially increased as a ramp from 42 to 250 photons, then kept constant, and manually decreased to maintain light intensity approximately at $\sim$90 μmol m^{-2} s^{-1} per OD$_{750}$ (Fig. S8F). After the switch to batch culture light was again increased in ramp from 70 to 250 μmol m^{-2} s^{-1}. For evaporation control and continuous culture, the total weight of the reactor setup was kept constant using the built-in Lambda reactor mass control module and automatic addition of fresh culture medium through the feed pump. In batch mode this controls for evaporation of medium. Continuous culture was performed by setting the waste pump to a fixed speed.

Calibrations of Measured & Controlled Data. The Lambda Photobioreactor (Fig. S7) was equipped with online monitoring of dissolved O$_2$ and pH, and additional monitoring of optical density by a DasGip OD4 module and monitoring of offgas O$_2$ and CO$_2$ concentrations and the weights of feed and pH control bottles by Arduino-based custom-built data loggers (Fig. S7). The signal from the OD4 probe was calibrated to offline OD$_{750}$ measurements (Fig. S8A-B). For normalizations of glycogen measurements by biomass and for estimation of the biomass density of cells (g$_{DCW}$/mL$_{cell}$, Fig. 5D) a LOESS regression of the OD$_\lambda$ signal was calibrated to CDW (Fig. S8C-D). The Lambda LUMO light module consistst of a strip of white LEDs and was calibrated to light intensity in μmol m^{-2} s^{-1} with a Licor light meter (LI-250A) with a spherical sensor bulb (LI-193) (Fig. S8E-F).

Calculation of Dilution and Growth Rates. All rates were calculated as slopes of measured data using piecewise linear segmentation [60], implemented in the CRAN R package dpseg (version 0.1.2) [61] (Fig. S9A-D). Growth rates were then calculated as the difference between slopes of measured biomass rate changes (OD$_\lambda$, CASY cell counts) and the culture dilution rate corrected for evaporation loss, see Figure S9E-F for details. Cell volume growth rate was calculated as the rate of change of the peaks of the CASY cell volume distributions.

Biomass & Metabolite Measurements

Cell Dry Weight Measurement. To determine the cell dry weight (CDW) 5 mL cell culture was filtered through a pre-dried and pre-weighed cellulose acetate membrane (pore size 0.45 μm) using a filtering flask. After that the membrane was dried at 50 °C for 24 h and weighed after cooling. 5 mL of filtered and dried growth medium served as a blank.

Optical Density OD$_{750}$ and Absorption Spectra. The optical density (OD$_{750}$) and absorbance spectra were measured on a Specord200 Plus (Jena Bioscience) dual path spectrometer, using BG11 as blank and reference. Samples were appropriately diluted with BG11 before measuring. All topA^{OX} time series samples were diluted 1:4 before recording OD$_{750}$. For absorbance spectra the OD$_{750}$ was adjusted to 0.5 and the absorbances at 750 nm were subtracted from each spectrum.

Glycogen Measurement. To determine the glycogen content, 0.5 mL of cell culture was harvested into reaction vessels that had been pre-cooled on ice. After centrifugation at maximum speed for 5 min at 4 °C the pellets were flash-frozen in liquid nitrogen and stored at -80 °C until further processing. To start the glycogen extraction, the pellet was resuspended in 400 μL KOH (30% w/v) and incubated at 95 °C for 2 h. For precipitation, 1200 μL ice cold ethanol was added and the mixture was incubated at -20 °C overnight. After centrifugation at 4 °C for 10 min at 10 000 g, the pellet was washed once with 70 % ethanol and again with pure ethanol. Afterwards the pellets were dried in a Concentrator Plus speed-vac (Eppendorf) for 20 min at 60 °C. Then the pellet was resuspended in 1 mL 100 mM sodium acetate (pH 4.5) supplemented with amyloglucosidase powder (Sigma-Aldrich, 10115) at a final concentration of 35 U/mL. For enzymatic digestion samples were incubated at 60 °C for 2 h. For the spectrometric glycogen determination the Sucrose/D-Glucose Assay Kit from Megazyme (K-SUCGL) was applied according to the manufacturer's specifications, but omitting the fructosidase reaction step and scaling down the total reaction volume to 850 μL. Absorbance at 510 nm was measured using a BMG Clariostar photospectrometer.

ATP and ADP Measurement. 2 mL tubes were preloaded with 250 µL of buffer BI (3 M HClO$_2$, 77 mM EDTA). 1 mL culture sample was added, briefly vortexed and incubated on ice for 15 min. 600 µL of BII (1 M KOH, 0.5 M KCl, 0.5 M Tris) were added, vortexed and incubated on ice for 10 min. After centrifugation at 0 °C for 10 min at 12 000 g, samples were flash-frozen in liquid nitrogen and stored at −80 °C until further processing. Extracts were thawed on ice and centrifuged again at 0 °C for 10 min at 12 000 g. 200 µL samples were added either to 320 µL of BIII/PEP (100 mM HEPES, 50 mM MgSO$_4$ x 7H$_2$O, adjusted to pH=7.4 with NaOH, and 1.6 mM phosphoenolpyruvate (Sigma-Aldrich, 860077) for ATP quantification or BIII/PEP+PK (BIII/PEP with 2 U/µL pyruvate kinase (Sigma-Aldrich, P1506) for ATP + ADP quantification, and incubated for 30 min at 37 °C. All samples were heat-inactivated at 90 °C for 10 min. For luminescence-based quantification, the Invitrogen ATP determination kit was used (ThermoFisher: A22066). 10 µL of each PEP or PEP+PK-treated sample was loaded in a white 96 well plate with solid bottom and kept on ice until the reaction was started. The luciferase master mix was scaled down in volume, and 90 µL of master mix was added to each well. Luminescence was recorded using a BMG Clariostar. ATP concentrations were calculated using a standard curve with commercially available ATP stock solution (Invitrogen).

Cell Count and Size Distribution To determine the cell count 10 µL cyanobacteria culture, pre-diluted for OD$_{750}$ measurement, were dispensed in 10 mL CASYton and measured with a Schaerfe CASY Cell Counter (Modell TTC) using a diameter 45 µm capillary. Cell size was recorded in the diameter range 0 µm–10 µm. Each sample was measured with 400 µL in triplicate runs.

Analysis of the raw data was performed in R. Counted events in the CASY are a mix of live cells, dead cells, cell debris and background signals. Only counts with diameter $d > 1.5$ µm and $d < 5$ µm were considered for the time series experiment (Fig. 5B) while a lower cutoff $d > 1.25$ µm was used for the endpoint measurements (Fig. 2C) to avoid cutting the distribution of the slightly smaller topA^{KD} cells. Since *Synechocystis* cells are spherical, the cell volumes were calculated from the reported cell diameters d as $V_{cell} = (\frac{d}{2})^3 \pi \frac{4}{3}$.

Flow Cytometry and Analysis Samples were fixed in 4 % para-formaldehyde in 1xPBS (phosphate buffered saline), washed three times in 1xPBS, and stained with the SYTO9 green fluorescent nucleic acid stain from the LIVE/DEAD kit (thermo) according to manufacturer's instructions. The flow cytometric measurements were taken at the FACS Facility at the Heinrich-Heine University (Dipl.-Biol. Klaus L. Meyer) using a BD FACSAria III. Forward scatter (FSC) and side-scatter (SSC) were recorded. Syto9 was measured with a 530/30 nm filter, and chlorophyll fluorescence was measured with 695/40 nm filter. For each sample 10,000 events (cells, debris and background) were recorded.

Data was exported in `.fcs` format, parsed and analyzed using the `flowCore` R packge [62], and plotted using functions from our in-house `segmenTools` R package.

DNA and Plasmid Extraction Agarose Gels

DNA Extraction. To isolate the DNA, 1 ml culture was centrifuged at maximum speed for 10 min at 4 °C, flash-frozen in liquid nitrogen and stored at −80 °C. After thawing, the samples were then resuspended in 1 mL 1x TE buffer by pipetting up and down and 100 µL lysozyme (50 mg/mL stock solution) was added, inverted, and incubated for 1h at 37 °C. Then 10 µL Proteinase K (20 mg/mL stock solution) and 100 µL 20 % SDS were added to the samples and incubated at 37 °C for 20 h. The lysed cell suspension was completely transferred to Phasemaker Tubes (ThermoFisher: A33248) and one volume of phenol/chloroform/isoamyl alcohol was added to each tube. The two resulting phases were thoroughly mixed and then separated by centrifugation for 10 min at maximal speed at 4 °C and the upper (nucleic acid-containing) phase transferred to a new tube. This was mixed with 100 ng/µL RNAse A and incubated for 15 min at 37 °C for RNA degradation. Followed by the addition of 1 volume of chloroform/isoamyl alcohol, the centrifugation step was repeated. The upper phase was again transferred and precipitated with 1 volume 2-propanol at −20 °C over night. After centrifugation at maximal speed and 4 °C for 10 min to pellet the DNA, the supernatant was discarded. The pellet was washed twice with 500 µL ice-cold 70 % EtOH and centrifuged for 10 min at maximal speed at 4 °C. After drying the pellets at room temperature, the pellets were dissolved in 30 µL water and the concentration was determined *via* Nanodrop (Thermo Scientific NanoDrop 2000c) .

Plasmid Extraction. 20 mL of cell culture were mixed with 20 mL of undenatured 99.5 % ethanol, precooled to $-80\,°C$, in 50 mL centrifuge tubes and stored at $-80\,°C$ until processing. After thawing on ice, the supernatant was discarded after centrifugation for 10 min at $4\,°C$ and 4000 g. The QIAprep Spin miniprep kit was used for the following steps and adapted and expanded for individual steps. The cell pellet was resuspended in 250 µL Qiagen P1 solution and transferred to 1.5 mL reaction tubes. Then 50 µL lysozyme solution (50 mg mL^{-1}) was added, mixed, and incubated for 1 h at $37\,°C$. After the addition of 55 µL of 20 % SDS and 3 µL of proteinase K (20 mg mL^{-1}), the reaction mixture was incubated at $37\,°C$ for 16 h. Starting with the alkaline lysis with the Qiagen P2 solution, all further steps according to the QIAprep Spin Miniprep Kit were carried out with amounts that were adjusted to the initial volume. Next, the concentrations and quality (260/280, 230/280 ratio) was determined using the Nanodrop (Thermo Scientific NanoDrop 2000c). The T5 exonuclease (NEB: M0363) was used for removal of linear and open circular DNA according to the manufacturer's protocol and incubated at $37\,°C$ for 30 min. The isolated plasmid DNA was further purified with the QIAprep modules. For this, 5 volumes of the PB buffer were added to 1 volume of DNA solution. All subsequent steps were carried out using the QIAprep Spin miniprep kit and the plasmid DNA concentration was determined with the Nanodrop.

Gel Electrophoresis and Analysis

Chloroquine Agarose Gel Electrophoresis. Agarose gels with chloroquine diphosphate (CQ, Sigma: C6628-50G, CAS: 50-63-5) were used to determine the relative migration speed of supercoiled topoisomers. A 1.2% agarose gel in 0.5x TBE (Roth: 3061.2) was prepared by heating to boiling. After cooling (hand-warm) CQ was added to 20 µg mL^{-1} and the mixture poured into the gel chamber. As running buffer served 0.5x TBE buffer with the same CQ concentration as the gel. For each sample, 120 ng DNA was mixed with loading dye and filled up to 30 µL with water. Next, the voltage source was adjusted 1.8 V cm^{-1}. Gels were run for 20 h–24 h covered in foil to protect from light. Next, the gel was washed two times for 30 min in 250 mL 0.5x TBE buffer to remove the CQ. After that, the gel is stained with 25 µL SYBR Gold (ThermoFisher: S11494) in 225 mL 0.5x TBE buffer for 3 h. Then the CQ agarose gel was imaged by a BioRad Imaging System (ChemiDocTM MP). Washing and staining were also performed light-protected. *** This method and its calibration are introduced in more detail in our preprint manuscript at `bioRxiv` [35]. ***

Analysis of Gel Electropherograms. Electropherograms of agarose gels of plasmids were extracted in ImageJ for each lane. Electropherograms from capillary gel electophoresis of total RNA samples (Agilent Bioanalyzer) were parsed from exported XML files using the R package `bioanalyzeR` (v 0.7.3, obtained from `https://github.com/jwfoley/bioanalyzeR`) [63]. Electropherograms were then processed in R, using LOESS smoothing and peak detection functions from the `msProcess` R package (version 1.0.7) (`https://cran.r-project.org/web/packages/msProcess/`). A baseline was determined in two steps using the `msSmoothLoess` function . The first step used the full signal and served to determine the coarse positions of peaks. The final baseline was then calculated from the signal after removal of peak values. This baseline was subtracted from the total signal to detect peaks (bands) with the `msPeakSimple` function from `msProcess` and calculate peak areas. For the total RNA analysis, the baseline signal stems from mRNA and rRNA degradation fragments, and was used to calculate ratios of rRNA peak areas to the "baseline" area.

Transcriptome Analysis: Sampling, RNA Extraction and Sequencing

RNA Extraction and Processing. Cellular activity was stopped by adding 1 mL culture directly to 250 µL 100 % ethanol supplemented with 5 % phenol, flash-frozen in liquid nitrogen and stored at $-80\,°C$ until further processing. For RNA extraction, a protocol modified from [64] was used. Briefly, frozen samples were centrifuged at maximum speed at $4\,°C$. After discarding the supernatant, the pellet was resuspended in 1 mL PGTX and incubated at $95\,°C$ for 5 min. After cooling on ice for 2 min, 700 µL chloroform:isoamyl alcohol (24:1) were added and the mixture was incubated shaking gently at room temperature for 10 min.The mixture was centrifuged for 10 min at maximal speed at $4\,°C$. The upper phase was transferred to a fresh tube and 1 volume chloroform:isoamyl alcohol was added. After repeating the centrifugation step, the upper phase was

again transferred and precipitated with 3 volumes of 100 % ethanol and 1/2 volume 7.5 M ammonium acetate at $-20\,^{\circ}$C over night. In cases where low RNA concentrations were to be expected, 1 μL RNA-grade glycogen was added to the precipitation mixture. The RNA was pelleted for 30 min at maximum speed and 4 °C, washed twice with 70 % ethanol and resuspended in 30 μL RNase-free water. RNA was DNaseI-digested using commercial DNaseI (ThermoFisher: EN0525), according to the manufacturer's specifications, but using twice the concentration of reaction buffer. DNaseI-digested RNA was phenol/chloroform extracted again to remove the DNaseI. For precipitation after DNaseI-digest, 1/10 volume of 3 M sodium acetate (pH 5.3), was used instead of ammonium acetate.

Quantitative RT-PCR. For qRT-PCR, DNaseI-digested RNA was reverse-transcribed to cDNA using the commercial RevertAid RT (ThermoFisher: K1621) according to the manufacturer's specifications in a reaction volume of 20 μL. To the 20 μL reaction, 60 μL RNase-free water was added. Of this cDNA dilution, 2 μL per reaction well were loaded, and 8 μL Master Mix prepared from the DyNAmo ColorFlash SYBR Green qPCR-Kit (ThermoFisher: F416L) were added to each 2 μL cDNA reaction. The thermal cycling conditions were as follows: 7 min at 95 °C, followed by 40 cycles of 5 s at 95 °C and 30 s at 60 °C. Data was recorded after each cycle. RT-negative controls and no-template-controls (distilled water) were included for each run. Each sample was loaded in technical triplicates.

Gene expression changes at indicated time points, usually after induction, were then quantified by the $\Delta\Delta Ct$ method [65]:

$$\Delta Ct = Ct_{\text{goi}} - Ct_{\text{ref}} \tag{1}$$
$$\Delta\Delta Ct = \Delta Ct_t - \Delta Ct_{t_0}, \tag{2}$$

where Ct_{goi} and Ct_{ref} are the Ct values of a gene of interest (*gyrA*, *gyrB*, *topA*) and of a reference gene (*rpoA*), respectively. The $\Delta\Delta Ct$ value then compares expression at time points t after induction with expression at a time point t_0 before induction, and is equivalent to a $\log_2$ fold change.

RNAseq: Total RNA Analysis, Library Generation and Sequencing. RNA quality was evaluated spectrometrically by Trinean Xpose (Gentbrugge, Belgium) and by fragment size distribution on an Agilent 2100 Bioanalyzer with the RNA Nano 6000 kit (Agilent Technologies, Böblingen, Germany). Electropherograms for the endpoint RNAseq samples were exported as XML files for further analysis; see paragraph *Analysis of Gel Electropherograms.* for details.

The Illumina Ribo-Zero Plus rRNA Depletion Kit was used to remove the ribosomal RNA molecules from the isolated total RNA. Removal of rRNA was evaluated with the RNA Pico 6000 kit on the Agilent 2100 Bioanalyzer. RNA was free of detectable rRNA. Preparation of cDNA libraries was performed according to the manufacturer's instructions for the TruSeq stranded mRNA kit (Illumina, San Diego, CA, United States). Subsequently, each cDNA library was sequenced on an Illumina NextSeq 500 system (2 x 75 nt PE high output v2.5).

RNAseq: Read Mapping. The resulting sequence reads were quality trimmed with Trimmomatic v0.33 [66] using standard setting. The quality trimmed reads were subsequently mapped to coding genes of the Synechocystis sp. PCC 6803 reference genome (NC_000911) including the plasmids pCA2._M, pCB2.4_M, pCC5.2_M, pSYSM, pSYSA, pSYSG and pSYSX (CP003270, CP003271, CP003272, NC_005229, NC_005230, NC_005231, NC_005232), and the constructed plasmid pSNDY_Prha_topA-6_119rhaS_20210310 using Bowtie 2 [67].

For the endpoint measurements from batch cultures the log2-fold changes with respect to the controls (EVC) were calculated with the DESeq2 algorithm [41] *via* the ReadXplorer software version 2.0 [68], based on three replicate measurements for each strain ("M-value"). For the time series read-count data were normalized by library sizes to the Transcripts Per kilobase Million (TPM) unit. Missing values at indivdual time points were interpreted as 0 TPM.

Transcriptome Analysis: Batch and Time-Series Analysis

Cluster Analysis. For clustering the time series into co-expressed cohorts, a previously established pipeline was used [69, 70]. Briefly, the time-series of TPM values was `arcsinh`-transformed:

$$x' = \ln\left(|x| + \sqrt{|x|^2 + 1}\right), \tag{3}$$

then the Discrete Fourier Transform (DFT) was calculated:

$$X_k = \sum_{n=0}^{N-1} x'_n e^{-2\pi i \frac{kn}{N}} \quad, \quad k = \{0, \ldots, N-1\}, \tag{4}$$

where $x'_n = \{x'_0, \ldots, x'_{N-1}\}$ are the (transformed) expression values at time points $\{t_0, \ldots, t_{N-1}\}$, and X_k, the DFT, is a vector of complex numbers representing the decomposition of the original time series into a constant (mean) component (at $k = 0$) and a series of harmonic oscillations around this mean. The input time series x_n were RNA-seq samples 2 to 16 (from -0.5 h to 72 h around the time of induction at 0 h), *i.e.*, without the first pre-induction time-point and ignoring the two long-term response samples. Components $k > 1$ were further scaled by the mean of amplitudes at all other components $k > 1$:

$$X'_{k>0} = \frac{X_{k>0}}{|X|_{k \neq \{0,k\}}} . \tag{5}$$

Real and imaginary parts of selected components $X_{k=1,\ldots,6}$ of the DFT were then clustered with the `flowClust` algorithm [71] over cluster numbers $K = 2, \ldots, 10$. The clustering with the maximal Bayesian Information Criterion, as reported by `flowClust` (Fig. S12A), was selected for further analyses. Data transformation and clustering were performed by the `processTimeseries` and the `clusterTimeseries2` functions of `segmenTier` and `segmenTools` packages, respectively. The resulting clusters were sorted and colored based on the comparison with diurnal co-expression cohorts (Fig. 6 and S17) for informative plots of the subsequent analyses.

Immediate Response Analysis. To estimate the immediate transcriptional response to *topA* overexpression (Fig. 6C, bottom panel) the difference of read counts (TPM) between the means of the two pre-induction time points (-1 d, -35 min) and the two post-induction time points (5 min, 20 min) was calculated. Transcripts with negative values were labelled as "down", with positive values as "up", and 0 or not available values as "nc".

Clustering of Transcription Units. Average expression was calculated for transcription units (TU) reported by ref. [46] from the expression of coding genes they encompass (*via* the "Sense.tags" column of the original data set). The resulting TU time-series was clustered by k-means, using cluster centers from the CDS clustering (Fig. 5) and identical time-series processing. This way protein-coding transcription units could be assigned to the same cluster labels (Fig. S23). Clusters were then sorted by their mean expression peaks, and colored and named according to the diurnal time of their peaks.

Clustering of Diurnal Transcriptome Data. Diurnal expression data from [44] were obtained from GEO (GSE79714) and genes summarized as the mean over all associated probes. These expression values were clustered the same was as described for the RNA-seq data: all time-series were DFT-transformed and amplitude-scaled DFT components $X'_{k=1,\ldots,7}$ (eq. 4–5) were clustered with `segmenTier` functions `processTimeseries` and `flowclusterTimeseries`, using the maximal BIC clustering with $K = 5$ clusters.

Cluster-Cluster Overlap Tests. Categorical enrichments, *e.g.*, coding gene co-expression cohorts vs. gene ₅₅₉ annotations, were analyzed by cumulative hypergeometric distribution tests (R's `phyper`) using `segmenTools`'s ₅₆₀ `clusterCluster` function and the `clusterAnnotation` wrapper for GO and and protein complex analysis, ₅₆₁ which compares overlaps of each pair of two distinct classifications into multiple classes, and stores overlap ₅₆₂ counts and p-values (enrichment tables) for informative plots (see "Enrichment Profile Plots"). ₅₆₃

Enrichment Table Sorting. For intuitively informative plots the enrichment tables were sorted. Table rows ₅₆₄ were sorted along the other dimension (table columns) such that all categories enriched above a certain ₅₆₅ threshold p_{sort} in the first column cluster are moved to the top, and, within, sorted by increasing p-values. Next, ₅₆₆ the same sorting is applied to all remaining row clusters for the second column cluster, and so on until the ₅₆₇ last column cluster. Remaining row clusters are either plotted unsorted below a red line or removed. This is ₅₆₈ especially useful to visualize enrichment of functional categories along the temporal program of co-expression ₅₆₉ cohorts, *e.g.*, Figure 6B. This sorting is implemented in `segmenTools`' function `sortOverlaps`. ₅₇₀

Cluster t-Test Profiles. To compare clusters (co-expression cohorts) with numerical data, here the log_2 ₅₇₁ fold-changes of transcript abundances in induced gyr^KD and topA^OX strains, we developed the `segmenTools`' ₅₇₂ function `clusterProfile`. For each cluster a two-sided `t-test` was performed (R base function `t.test`, ₅₇₃ incl. Welch approximation for different sample sizes), comparing the distribution of values of the cluster with ₅₇₄ all other values. The reported t statistic and the p-value were stored for each test. The resulting t-test profile ₅₇₅ was stored for informative plots (see "Enrichment Profile Plots"). ₅₇₆

Enrichment Profile Plots. The results of cluster enrichment tests (cluster-cluster overlap tests and t-test ₅₇₇ profiles) were visualized as colored table plots (*e.g.* Fig. 6B, C), using `segmenTools`' function `plotOverlaps`. ₅₇₈ For the categorical overlap tests, the total counts of overlapping pairs are plotted as text, and for t-test profils ₅₇₉ the rounded t statistic. The text color is black or white based on a p-value cutoff p_{txt} (as indicated). ₅₈₀

The field background colors scale with $\log_2(p)$ of the reported p-values, where the full color corresponds ₅₈₁ to a minimal p-value p_{min} cutoff (as indicated). For categorical enrichment tests the full color is black and other ₅₈₂ colors are selected from a gradient to white. For numerical tests, the sign of the t statistic is used to determine ₅₈₃ a color to indicate the direction of change: red for negative values (downregulated) and blue for positive values ₅₈₄ (upregulated). ₅₈₅

Motif Scan and DNA Structure Analysis. The genome was scanned for short sequence motifs using ₅₈₆ custom-built R code based on the `str_locate_all` function of the `stringr` R package [72] into a vector ₅₈₇ of 0 and 1 for each genome position, where 1 indicates occurence of the motif under consideration. Motif ₅₈₈ occurence vectors upstream and downstream of start codons or transcription start sites were extracted from ₅₈₉ the genome vector and aligned into a matrix (columns: positions around the alignment anchor, rows: all ₅₉₀ genomic sites under consideration). The occurence of a motif in all sequences of a cluster were counted ₅₉₁ at each position i in 66 bp windows surrounding the position. Cumulative hypergeometric distribution tests ₅₉₂ (R's `phyper`) were performed to analyze statistical enrichment or deprivation of a given motif within the 66 bp ₅₉₃ window of all genes in a cluster vs. the same window in all genes in the total analyzed set (all clustered and ₅₉₄ aligned sequences), and the p-values p_i at position i stored. In the resulting plots of the mean position-wise ₅₉₅ motif occurence the size of the plotted data point at position i was scaled by the enrichment or deprivation ₅₉₆ p-values to emphasize regions of significant difference (Fig. 6D). The maximal size was determined by the ₅₉₇ minimum p-value in each test series (Figure panels). The point style (closed or open circles) indicates the ₅₉₈ directionality of the test (enriched or deprived). The significance points are only shown at every third or tenth ₅₉₉ position to avoid overlaps. ₆₀₀

Other Data Sources. Genome sequences and annotation were downloaded from NCBI, see "RNAseq: ₆₀₁ Read Mapping" for the RefSeq IDs. The gene "categories" annotation was downloaded on 2017-09- ₆₀₂ 23 from `CyanoBase` [73]: `http://genome.annotation.jp/cyanobase/Synechocystis/genes/category.` ₆₀₃ `txt`. Gene Ontology annotation was downloaded from the `UniProt` database (2021-03-20, `organism:1111708`) ₆₀₄

[74]. Datasets from other publications were all obtained from the supplemental materials of the indicated publications.

References

1. B.J. Peter, J. Arsuaga, A.M. Breier, A.B. Khodursky, P.O. Brown, and N.R. Cozzarelli. Genomic transcriptional response to loss of chromosomal supercoiling in *Escherichia coli*. *Genome Biol*, 5(11):R87, 2004.

2. N. Blot, R. Mavathur, M. Geertz, A. Travers, and G. Muskhelishvili. Homeostatic regulation of supercoiling sensitivity coordinates transcription of the bacterial genome. *EMBO Rep*, 7(7):710–715, Jul 2006.

3. V. Vijayan, R. Zuzow, and E.K. O'Shea. Oscillations in supercoiling drive circadian gene expression in cyanobacteria. *Proc Natl Acad Sci U S A*, 106(52):22564–22568, Dec 2009.

4. J.S. Prakash, M. Sinetova, A. Zorina, E. Kupriyanova, I. Suzuki, N. Murata, and D.A. Los. DNA supercoiling regulates the stress-inducible expression of genes in the cyanobacterium *Synechocystis*. *Mol Biosyst*, 5(12):1904–1912, Dec 2009.

5. A.G. de la Campa, M.J. Ferrandiz, A.J. Martin-Galiano, M.T. Garcia, and J.M. Tirado-Velez. The transcriptome of *Streptococcus pneumoniae* induced by local and global changes in supercoiling. *Front Microbiol*, 8:1447, 2017.

6. M.J. Szafran, M. Gongerowska, T. Malecki, M. Elliot, and D. Jakimowicz. Transcriptional response of *Streptomyces coelicolor* to rapid chromosome relaxation or long-term supercoiling imbalance. *Front Microbiol*, 10:1605, 2019.

7. C.J. Dorman. DNA supercoiling and transcription in bacteria: a two-way street. *BMC Mol Cell Biol*, 20(1):26, Jul 2019.

8. L.S. Hsieh, R.M. Burger, and K. Drlica. Bacterial DNA supercoiling and [ATP]/[ADP]. changes associated with a transition to anaerobic growth. *J Mol Biol*, 219(3):443–450, Jun 1991.

9. M. van Workum, S.J. van Dooren, N. Oldenburg, D. Molenaar, P.R. Jensen, J.L. Snoep, and H.V. Westerhoff. DNA supercoiling depends on the phosphorylation potential in *Escherichia coli*. *Mol Microbiol*, 20(2):351–360, Apr 1996.

10. J.L. Snoep, C.C. van der Weijden, H.W. Andersen, H.V. Westerhoff, and P.R. Jensen. DNA supercoiling in *Escherichia coli* is under tight and subtle homeostatic control, involving gene-expression and metabolic regulation of both topoisomerase I and DNA gyrase. *Eur J Biochem*, 269(6):1662–1669, Mar 2002.

11. A.A. Travers, G. Muskhelishvili, and J.M. Thompson. DNA information: from digital code to analogue structure. *Philos Transact A Math Phys Eng Sci*, 370(1969):2960–2986, Jun 2012.

12. M. Schaechter, O. Maaløe, and N.O. Kjeldgaard. Dependency on medium and temperature of cell size and chemical composition during balanced growth of *Salmonella typhimurium*. *J Gen Microbiol*, 19(3):592–606, Dec 1958.

13. M. Scott, C.W. Gunderson, E.M. Mateescu, Z. Zhang, and T. Hwa. Interdependence of cell growth and gene expression: origins and consequences. *Science*, 330(6007):1099–1102, Nov 2010.

14. T. Zavrel, M. Faizi, C. Loureiro, G. Poschmann, K. Stuhler, M. Sinetova, A. Zorina, R. Steuer, and J. Cerveny. Quantitative insights into the cyanobacterial cell economy. *Elife*, 8, Feb 2019.

15. L.F. Liu and J.C. Wang. Supercoiling of the DNA template during transcription. *Proc Natl Acad Sci U S A*, 84(20):7024–7027, Oct 1987.

16. S. Kim, B. Beltran, I. Irnov, and C. Jacobs-Wagner. Long-distance cooperative and antagonistic RNA polymerase dynamics via DNA supercoiling. *Cell*, 179(1):106–119.e16, Sep 2019.

17. N.S. Rovinskiy, A.A. Agbleke, O.N. Chesnokova, and N.P. Higgins. Supercoil levels in *E. coli* and *Salmonella* chromosomes are regulated by the C-terminal 35(-)38 amino acids of GyrA. *Microorganisms*, 7(3), Mar 2019.

18. J. Ma, L. Bai, and M.D. Wang. Transcription under torsion. *Science*, 340(6140):1580–1583, Jun 2013.

19. D. Sutormin, N. Rubanova, M. Logacheva, D. Ghilarov, and K. Severinov. Single-nucleotide-resolution mapping of DNA gyrase cleavage sites across the *Escherichia coli* genome. *Nucleic Acids Res*, 47(3):1373–1388, Feb 2019.

20. M. Drolet, S. Broccoli, F. Rallu, C. Hraiky, C. Fortin, E. Masse, and I. Baaklini. The problem of hypernegative supercoiling and R-loop formation in transcription. *Front Biosci*, 8:d210–21, Jan 2003.

21. W. Ahmed, C. Sala, S.R. Hegde, R.K. Jha, S.T. Cole, and V. Nagaraja. Transcription facilitated

genome-wide recruitment of topoisomerase I and DNA gyrase. *PLoS Genet*, 13(5):e1006754, May 2017.

22. H.Y. Wu and M. Fang. DNA supercoiling and transcription control: a model from the study of suppression of the leu-500 mutation in *Salmonella typhimurium* topA- strains. *Prog Nucleic Acid Res Mol Biol*, 73:43–68, 2003.

23. X. Zhi, S. Dages, K. Dages, Y. Liu, Z.C. Hua, J. Makemson, and F. Leng. Transient and dynamic DNA supercoiling potently stimulates the leu-500 promoter in *Escherichia coli*. *J Biol Chem*, 292(35):14566–14575, Sep 2017.

24. P. Sobetzko, M. Glinkowska, A. Travers, and G. Muskhelishvili. DNA thermodynamic stability and supercoil dynamics determine the gene expression program during the bacterial growth cycle. *Mol Biosyst*, Mar 2013.

25. M.A. Woelfle, Y. Xu, X. Qin, and C.H. Johnson. Circadian rhythms of superhelical status of DNA in cyanobacteria. *Proc Natl Acad Sci U S A*, 104(47):18819–18824, Nov 2007.

26. M.L. Salvador, U. Klein, and L. Bogorad. Endogenous fluctuations of DNA topology in the chloroplast of *Chlamydomonas reinhardtii*. *Mol Cell Biol*, 18(12):7235–7242, Dec 1998.

27. M.K. Wall, L.A. Mitchenall, and A. Maxwell. *Arabidopsis thaliana* DNA gyrase is targeted to chloroplasts and mitochondria. *Proc Natl Acad Sci U S A*, 101(20):7821–7826, May 2004.

28. H. Ito, M. Mutsuda, Y. Murayama, J. Tomita, N. Hosokawa, K. Terauchi, C. Sugita, M. Sugita, T. Kondo, and H. Iwasaki. Cyanobacterial daily life with Kai-based circadian and diurnal genome-wide transcriptional control in *Synechococcus elongatus*. *Proc Natl Acad Sci U S A*, 106(33):14168–14173, Aug 2009.

29. J.R. Waldbauer, S. Rodrigue, M.L. Coleman, and S.W. Chisholm. Transcriptome and proteome dynamics of a light-dark synchronized bacterial cell cycle. *PLoS One*, 7(8):e43432, 2012.

30. R. Lehmann, R. Machné, J. Georg, M. Benary, I. M. Axmann, and R. Steuer. How cyanobacteria pose new problems to old methods: challenges in microarray time series analysis. *BMC Bioinformatics*, 14(1):133, Apr 2013.

31. C. Beck, S. Hertel, A. Rediger, R. Lehmann, A. Wiegard, A. Kolsch, B. Heilmann, J. Georg, W.R. Hess, and I.M. Axmann. Daily expression pattern of protein-encoding genes and small noncoding RNAs in *Synechocystis* sp. strain PCC 6803. *Appl Environ Microbiol*, 80(17):5195–5206, Sep 2014.

32. T. Mori and C.H. Johnson. Circadian programming in cyanobacteria. *Semin Cell Dev Biol*, 12(4):271–278, Aug 2001.

33. R.M. Smith and S.B. Williams. Circadian rhythms in gene transcription imparted by chromosome compaction in the cyanobacterium *Synechococcus elongatus*. *Proc Natl Acad Sci U S A*, 103(22):8564–8569, May 2006.

34. R. Lehmann, R. Machné, and H. Herzel. The structural code of cyanobacterial genomes. *Nucleic Acids Res*, 42(14):8873–8883, Oct 2014.

35. Salima Rüdiger, Anne Rediger, Adrian Kölsch, Dennis Dienst, Ilka Maria Axmann, and Rainer Machné. Plasmid supercoiling decreases during the dark phase in cyanobacteria: a clarification of the interpretation of chloroquine-agarose gels. *bioRxiv*, 2021.

36. A. Behle, P. Saake, A.T. Germann, D. Dienst, and I.M. Axmann. Comparative dose-response analysis of inducible promoters in cyanobacteria. *ACS Synth Biol*, 9(4):843–855, Apr 2020.

37. L. Yao, I. Cengic, J. Anfelt, and E.P. Hudson. Multiple gene repression in cyanobacteria using CRISPRi. *ACS Synth Biol*, 5(3):207–212, Mar 2016.

38. R. Damrow, I. Maldener, and Y. Zilliges. The multiple functions of common microbial carbon polymers, glycogen and PHB, during stress responses in the non-diazotrophic cyanobacterium *Synechocystis* sp. PCC 6803. *Front Microbiol*, 7:966, 2016.

39. T. Hasunuma, F. Kikuyama, M. Matsuda, S. Aikawa, Y. Izumi, and A. Kondo. Dynamic metabolic profiling of cyanobacterial glycogen biosynthesis under conditions of nitrate depletion. *J Exp Bot*, 64(10):2943–2954, Jul 2013.

40. K.B. Mollers, D. Cannella, H. Jorgensen, and N.U. Frigaard. Cyanobacterial biomass as carbohydrate and nutrient feedstock for bioethanol production by yeast fermentation. *Biotechnol Biofuels*, 7:64, 2014.

41. M.I. Love, W. Huber, and S. Anders. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol*, 15(12):550, 2014.

42. S.B. Zimmerman and S.O. Trach. Estimation of macromolecule concentrations and excluded volume effects for the cytoplasm of escherichia coli. *J Mol Biol*, 222(3):599–620, Dec 1991.

43. Enno R. Oldewurtel, Yuki Kitahara, Baptiste Cordier, Gizem Özbaykal, and Sven van Teeffelen. Bacteria control cell volume by coupling cell-surface expansion to dry-mass growth. *bioRxiv*, 2019.

44. R. Saha, D. Liu, A. Hoynes-O'Connor, M. Liberton, J. Yu, M. Bhattacharyya-Pakrasi, A. Balassy,

F. Zhang, T.S. Moon, C.D. Maranas, and H.B. Pakrasi. Diurnal regulation of cellular processes in the cyanobacterium *Synechocystis* sp. strain PCC 6803: Insights from transcriptomic, fluxomic, and physiological analyses. *MBio*, 7(3), 2016.

45. M.J. Ferrandiz, A.J. Martin-Galiano, J.B. Schvartzman, and A.G. de la Campa. The genome of *Streptococcus pneumoniae* is organized in topology-reacting gene clusters. *Nucleic Acids Res*, 38(11):3570–3581, Jun 2010.

46. M. Kopf, S. Klahn, I. Scholz, J.K. Matthiessen, W.R. Hess, and B. Voss. Comparative analysis of the primary transcriptome of *Synechocystis* sp. PCC 6803. *DNA Res*, 21(5):527–539, Oct 2014.

47. A. Wiegard, C. Kobler, K. Oyama, A.K. Dorrich, C. Azai, K. Terauchi, A. Wilde, and I.M. Axmann. *Synechocystis* KaiC3 displays temperature- and KaiB-dependent ATPase activity and is important for growth in darkness. *J Bacteriol*, 202(4), Jan 2020.

48. A. Srivastava, M.L. Summers, and R. Sobotka. Cyanobacterial sigma factors: Current and future applications for biotechnological advances. *Biotechnol Adv*, 40:107517, May 2020.

49. Y. Nishijima, Y. Kanesaki, H. Yoshikawa, T. Ogawa, K. Sonoike, Y. Nishiyama, and Y. Hihara. Analysis of spontaneous suppressor mutants from the photomixotrophically grown pmgA-disrupted mutant in the cyanobacterium *Synechocystis* sp. PCC 6803. *Photosynth Res*, 126(2-3):465–475, Dec 2015.

50. K.S. Jeong, Y. Xie, H. Hiasa, and A.B. Khodursky. Analysis of pleiotropic transcriptional profiles: a case study of DNA gyrase inhibition. *PLoS Genet*, 2(9):e152, Sep 2006.

51. S. Chong, C. Chen, H. Ge, and X.S. Xie. Mechanism of transcriptional bursting in bacteria. *Cell*, 158(2):314–326, Jul 2014.

52. N. Rovinskiy, A.A. Agbleke, O. Chesnokova, Z. Pang, and N.P. Higgins. Rates of gyrase supercoiling and transcription elongation control supercoil density in a bacterial chromosome. *PLoS Genet*, 8(8):e1002845, 2012.

53. A.M. Puszynska and E.K. O'Shea. ppGpp controls global gene expression in light and in darkness in *S. elongatus*. *Cell Rep*, 21(11):3155–3165, Dec 2017.

54. S.A. Angermayr, P. van Alphen, D. Hasdemir, G. Kramer, M. Iqbal, W. van Grondelle, H.C. Hoefsloot, Y.H. Choi, and K.J. Hellingwerf. Culturing *Synechocystis* sp. strain PCC 6803 with N2 and CO2 in a diel regime reveals multiphase glycogen dynamics with low maintenance costs. *Appl Environ Microbiol*, 82(14):4180–4189, Jul 2016.

55. R. Menzel and M. Gellert. Regulation of the genes for *E. coli* DNA gyrase: homeostatic control of DNA supercoiling. *Cell*, 34(1):105–113, Aug 1983.

56. M. Drolet, X. Bi, and L.F. Liu. Hypernegative supercoiling of the DNA template during transcription elongation in vitro. *J Biol Chem*, 269(3):2068–2074, Jan 1994.

57. A. Werner, C.D. Broeckling, A. Prasad, and C.A.M. Peebles. A comprehensive time-course metabolite profiling of the model cyanobacterium *Synechocystis* sp. PCC 6803 under diurnal light:dark cycles. *Plant J*, 99(2):379–388, Jul 2019.

58. K. Labun, T.G. Montague, M. Krause, Y.N. Torres Cleuren, H. Tjeldnes, and E. Valen. CHOPCHOP v3: expanding the CRISPR web toolbox beyond genome editing. *Nucleic Acids Res*, 47(W1):W171–W174, Jul 2019.

59. Rosmarie Rippka, Josette Deruelles, John B. Waterbury, Michael Herdman, and Roger Y. Stanier. Generic assignments, strain histories and properties of pure cultures of cyanobacteria. *Journal of General Microbiology*, 111(1):1–61, 1979.

60. Richard Bellman. On the approximation of curves by line segments using dynamic programming. *Communications of the ACM*, 4(6):284, 1961.

61. Rainer Machné and Peter F. Stadler. *dpseg: Piecewise Linear Segmentation by Dynamic Programming*. R package version 0.1.2.

62. B Ellis, P Haaland, F Hahne, N Le Meur, N Gopalakrishnan, J Spidlen, M Jiang, and G Finak. *flowCore: Basic structures for flow cytometry data*, 2019. R package version 1.48.1.

63. Joseph Foley. *bioanalyzeR: Analysis of Agilent electrophoresis data*, 2021. R package version 0.7.3.

64. F.L. Pinto, A. Thapper, W. Sontheim, and P. Lindblad. Analysis of current and alternative phenol based RNA extraction methodologies for cyanobacteria. *BMC Mol Biol*, 10:79, Aug 2009.

65. K.J. Livak and T.D. Schmittgen. Analysis of relative gene expression data using real-time quantitative PCR and the $2^{-\Delta\Delta C_T}$ method. *Methods*, 25(4):402–408, Dec 2001.

66. A.M. Bolger, M. Lohse, and B. Usadel. Trimmomatic: a flexible trimmer for illumina sequence data. *Bioinformatics*, 30(15):2114–2120, Aug 2014.

67. B. Langmead and S.L. Salzberg. Fast gapped-read alignment with Bowtie 2. *Nat Methods*, 9(4):357–359, Mar 2012.

68. R. Hilker, K.B. Stadermann, O. Schwengers, E. Anisiforov, S. Jaenicke, B. Weisshaar, T. Zimmermann,

and A. Goesmann. ReadXplorer 2 – detailed read mapping analysis and visualization from one single source. *Bioinformatics*, 32(24):3702–3708, Dec 2016.

69. R. Machné and D.B. Murray. The yin and yang of yeast transcription: elements of a global feedback system between metabolism and chromatin. *PLoS One*, 7(6):e37906, 2012.

70. R. Machné, D.B. Murray, and P.F. Stadler. Similarity-based segmentation of multi-dimensional signals. *Sci Rep*, 7(1):12355, Sep 2017.

71. K. Lo, F. Hahne, R.R. Brinkman, and R. Gottardo. flowClust: a Bioconductor package for automated gating of flow cytometry data. *BMC Bioinformatics*, 10:145, 2009.

72. Hadley Wickham. *stringr: Simple, Consistent Wrappers for Common String Operations*, 2019. R package version 1.4.0.

73. T. Fujisawa, R. Narikawa, S.I. Maeda, S. Watanabe, Y. Kanesaki, K. Kobayashi, J. Nomata, M. Hanaoka, M. Watanabe, S. Ehira, E. Suzuki, K. Awai, and Y. Nakamura. CyanoBase: a large-scale update on its 20th anniversary. *Nucleic Acids Res*, 45(D1):D551–D554, Jan 2017.

74. UniProt Consortium. UniProt: the universal protein knowledgebase in 2021. *Nucleic Acids Res*, 49(D1):D480–D489, Jan 2021.

75. S. Sakata, N. Mizusawa, H. Kubota-Kawai, I. Sakurai, and H. Wada. Psb28 is involved in recovery of photosystem II at high temperature in *Synechocystis* sp. PCC 6803. *Biochim Biophys Acta*, 1827(1):50–59, Jan 2013.

76. J. Fu and X. Xu. The functional divergence of two glgP homologues in *Synechocystis* sp. PCC 6803. *FEMS Microbiol Lett*, 260(2):201–209, Jul 2006.

Supporting Information

Strain name	Chromosomal genotype	Plasmid
EVC	P_{L22} dCas9	pSNDY
gyrA^{KD}	P_{L22} dCas9; P_{L22} sgRNA_*gyrA*	pSNDY (EVC)
gyrB^{KD}	P_{L22} dCas9; P_{L22} sgRNA_*gyrB*	pSNDY (EVC)
gyrAB^{KD}	P_{L22} dCas9; P_{L22} sgRNA_*gyrA+gyrB*	pSNDY (EVC)
topA^{KD}	P_{L22} dCas9; P_{L22} sgRNA_*topA*	pSNDY (EVC)
topA^{OX}	P_{L22} dCas9	pSNDY P_{J23119} *rhaS*; P_{rha} *topA*

Table 1. Overview of strains investigated in this work. The parental strain for all strains listed here (*Synechocystis* sp. PCC 6803 encoding aTc-inducible dCas9) was a gift from P. Hudson and L. Yao [37]

Gene	Direction	Sequence
gyrA	forward	GAACTTTGGCTCCGTGGATAA
	reverse	GCCTCAATGTCCCGCAATAA
gyrB	forward	TGCCCGTAAGCGCAATAA
	reverse	ATTCTGGGTCCGGTACTTTAAC
topA	forward	AGACCGGGAAGGAGAAAGTA
	reverse	CGAATGGCTTCCTGGGTAAT
rpoA	forward	CCATGAGTTCGCCACTATTCT
	reverse	GGCTGATCGGTGTAGCTTT

Table 2. qPCR Primers.

Gene	sgRNA Sequence
gyrA	TCAGTCATGCAATTACTCCA
gyrB	CTGGCTTCAACCCATCCCGTGCAT
topA	GATAGTGCGGGCTTTAGTGG

Table 3. sgRNA Sequences.

Figure S1. Batch Culture Endpoint Measurements. See Figure 2 for details. **A:** RT-qPCR results using *rpoA* as reference gene. Boxplots of 9 technical replicates (3 samples, each measured 3x) . **B:** Absorption spectra at incolution and harvest times. **C:** Recovery without inducer: all cultures were re-inoculated in fresh BG11 with antibiotics but without the inducers (aTc and rhamnose) and the cell size distributions measured at indicated time points. **D:** Chloroquine-agarose gels (1.2% agarose, 0.5x TBE and $20\,\mu g\,mL^{-1}$ CQ) of plasmids extracted at harvest time (5 d).

Figure S2. Plasmid Supercoiling Gels. **A:** Growth curves of topA^{OX} strain, induced with 1 mM rhamnose at time 0 h and uninduced control. A starter culture was split into 8 cultures at 0 h, each harvested at the indicated time points for OD$_{750}$ measurement and plasmid extraction. **B & C**: Yields of plasmid extraction over time, each normalized to the OD$_{750}$ (A), and before (B) and after (C) treatment with the T5 exonuclease to remove all non closed circular DNA. **D & E**: Chloroquine-agarose gels (1.2% agarose, 0.5x TBE and 20 µg mL^{-1} CQ) of plasmids extracted from topA^{OX} strain cultures (A), induced (B) and uninduced (C). Note that only the gel of the induced culture (B) was run for a longer time (XYZ) to get a better separation of topoisomers of the pCC5.5_M plasmid. Samples are ordered by sampling times (see A) from left to right on both gels. The 120 h sample is missing in (E).

Figure S3. Growth Curves & RNA Extraction for RNA-seq Experiment. **A:** Growth curves of triplicate cultures (split upon induction at 0 d).**B & C:** Total RNA compositions were analysed by a formaldehyde-agarose gel (B, 500 ng RNA per well) and by capillary gel electrophoresis (Agilent Bioanalyzer); electropherograms of these samples were analyzed for Figures 4, S5 and S6. Sample lane order are identical in (B) and (C) and comprise of triplicates from EVC (lanes 1-3), gyrA^{KD} (lanes 4-6), gyrB^{KD} (lanes 7-9) and topA^{OX} (lanes 10-12).

Figure S4. Flow Cytometry. Cells from the cultures used for RNA-seq and total RNA analysis (Fig. 4) were fixed in para-formaldehyde and stained with Syto-9, a nucleic acid fluorescence marker, and analyzed by flow cytometry. The data was gated by the side scatter signal (SSC-A> 2000) and the forward scatter signal (FSC-A> 10) to filter debris and background signals. Forward scatter (FSC-A) is proportional to cell size, side scatter (SSC-A) reflects cytoplasmic granularity and morphology; the FITC fluorescence channel (530/30 nm) excites the Syto-9 stain, and the PI channel (695/40 nm) excites chlorophyll. The natural logarithm ($\ln$) of all data was plotted. Colors reflect local density (red: high, blue: low). The bottom panels show a zoom into the data, comparing the two strains with the most extreme values, EVC and topA^{OX} (white: high local density).

Figure S5. Total RNA Electropherogram Analysis. Electropherograms of the capillary gel electrophoresis (Fig. 4A), exported as XML files from the 2100 Bioanalyzer software. **A–D:** are each the triplicate samples for the indicated strain. Samples are the same as shown on the formaldehyde-agarose gel in Figure S3B and subsequently used for RNAseq analysis (Fig. 4B,C). Data was parsed into R with `bioanalyzeR` (v 0.7.3) [63]. Baseline and peak detection and quantification were performed in R, see Methods paragraph *Analysis of Gel Electropherograms* for details. The areas of the blue peaks were further analyzed as relative rRNA species abundances (see top axis annotation). r23S.S and r23S.L are short and long fragments of the 23S rRNA typically seen in *Synechocystis*, the other peaks were assigned to 5S, 16S and (full length) 23S rRNA. The red area under the baseline (green) and > 200 bp was used as relative mRNA abundance for the ratios in Figure 4B,C and S6. The 200 nt length cut-off was based on the size calibration to the RNA ladder lane as reported by the `bioanalyzeR` parser. The peak on the left is the 25 nt lower size marker.

Figure S6. rRNA vs. mRNA: Relative Abundances. Ratios of the indicated rRNA peak areas to the baseline (mRNA) area, the blue and red areas in Figure S5, respectively. The error bars indicate the mean and standard deviation of the three replicates.

Figure S7. Photobioreactor Setup. A schematic overview of the cultivation setup showing the Lambda minifor bioreactor in front and top-down views alongside its external components and custom expansions. The gas input mixture is generated by a Lambda MASSFLOW 5000 gas flow controller and a Voegtlin red-y smart controller, which regulate the flow of compressed air and CO_2 respectively. This input gas mixture is then introduced into the cultivation vessel via the sparger at the end of the agitation unit. The offgas condenser as well as the reactor's cooling finger are part of a water cooling circuit which is regulated by a Lauda Eco Silver thermostat set to $16\,°C$. An Aalborg Massflow Meter monitors the flow rate of the culture's offgas before it is lead through a custom microcontroller-based gas sensor array in order to evaluate its O_2 and CO_2 content. The reactor actively regulates the culture's pH and temperature values by controlling its heating compartment as well as the Lambda Preciflow peristaltic pumps which are attached to NaOH and H_2SO_4 stock bottles, each $0.5\,M$. Additional culture parameters are monitored by a dissolved O_2 probe attached to the reactor and an OD4 probe connected to a DASGIP OD4 device. An additional set of peristaltic pumps is attached to the culture's medium stock and waste containers in order to control the reactor's volume and medium turnover. The reactor weighting module enables the system to operate under chemostatic conditions. This is achieved by manually configuring the medium feed peristaltic pump at a constant speed in order to achieve a desired medium turnover rate while automatically regulating the waste pump speed to keep the total reactor weight constant. Additionally, a custom microcontroller-based scale setup is monitoring the weight of both the medium and NaOH stock bottles, which allows for the calculation of medium and base pump rates from the recorded data. The culture's illumination is provided by the Lambda LUMO modules, an LED strip fitted around the cultivation vessel.

Figure S8. Calibrations. **A:** LOESS regression (R `loess`) of the raw signal (resolution ca. 1 sec) from the DasGip OD4 module ($OD_{\lambda,raw}$). **B:** calibration of the LOESS fit of the $OD_{\lambda,raw}$ signal to offline $OD_{750\,nm}$ by linear regression (R *lm*). The calibrated signal is used throughout the document and denoted OD_λ. **C:** calibration cell dry weight (CDW) to the OD_λ signal. Data points marked by X were removed as outliers. **D:** the LOESS fit of the OD4 signal was then used to estimate CDW for all time points. **E:** calibration of the Lambda LUMO light module with a Licor light meter (LI-250A) with a spherical sensor bulb (LI-193). **F:** time-series of set and calibrated light intensities (black line, left y-axis) compared to the OD_λ time-series (gray line, right axis). The light intensity was manually adjusted to avoid high-light stress in the culture during biomass decrease. **G:** The Arduino-based scales where calibrated prior to the experiment (not shown). During the experiment the liquid level on the 5L feed bottle was marked regularly, and the mass of water filled to these marked was recorded on a benchtop scale (Kern XZY) after the experiment to test consistent performance. The recorded mass was reproduced sufficiently well (red line: linear regression): the intercept of the linear regression corresponds to the mass of the empty feed bottle and the slope was ≈ 1. Since the manual marks on the bottle are more error prone than the pre-calibration, we did not re-calibrate the data but relied on the recorded mass for calculation of the dilution rate.

Figure S9. Calculation of Dilution and Growth Rates. All rates were calculated from the slopes of measured data (or of their natural logarithms as indicated) using piecewise linear segmentation with the R package dpseg. The plots in **A-D** were generated by dpseg and the vertical lines indicate borders of the piecewise segments, and the used penality parameter P is shown in the plot title on the top axis. The minimal segment length paramater minl was only used in (C). **A**: the calibrated OD_λ signal (1 sec resolution) was smoothed with a moving average and window size 15 and interpolated at 300 sec intervals. **B**: sum of the recorded weights of edium feed and pH control bottle weight; outliers (faulty measurements or bottle changes) were removed and data interpolated at 300 sec intervals. **C:** the total cell count for each CASY measurement, single measurements and means of technical duplicates. **D:** the total cell volume, calculated as the integral of the single cell volume distribution, for each CASY measurement, single measurements and means of technical duplicates. **E:** Observed rates. The (negative) slopes of the summed bottle weight changes (B) reflect the amounts added to the reactor culture by the Lambda reactor mass control system, assuming 1 g/mL density. The total culture dilution rate (dashed gray line, "dilution + evaporation") is obtained by division by the culture volume ($V_\ell = 1$ L). The liquid loss by evaporation is seen at times before onset of continuous culture (time -4 d) and is subtracted to obtain the actual dilution rate ϕ (black line). The slopes of the change of the natural logarithms of the OD_λ signal (A), the total integrated cell volume (B), and the cell counts (C) are the observed change rates $\mu_{obs,OD}$ (gray line), $\mu_{obs,volume}$ (red line) and $\mu_{obs,count}$ (blue line), respectively. **F:** The culture growth rates μ_{OD} (gray line) and μ_{count} (blue line) and μ_{volume} (red line) were calculated as the difference between observed change rates and the culture dilution rate: $\mu = \mu_{obs} - \phi$.

Figure S10. Re-Induction of Bioreactor Time-Series Cultures. After harvest, cells from the bioreactor timeseries experiment (strain topA^{OX}) were spinned down, washed and re-inoculated in fresh BG11 with antibiotics and without (top panels,-rhamnose) or with (bottom panels, +rhamnose) in shake flasks (batch culture), and cell count and volume distribution measured daily with the CASY cell counter. **A:** cell volume distributions as gray-scale (left axis) and cell count and total cell volume with open and closed points (right y-axis). **B:** Distributions of the cell diameter as reported by the CASY cell counter and from which cell volumes were calculated. Cells grew normally without further topA^{OX} induction and showed the initial transient volume increase that was observed in all experiments without induced volume growth (*e.g.* EVC and topA^{KD} in Fig. 2B and Fig. S1C). After re-induction of topA^{OX}, only a fraction of cells again showed in increase in cell volume.

Figure S11. RT-qPCR vs. RNA-seq. **A:** The topA, gyrA and gyrB genes were also measured by RT-qPCR using *rpoA* as a reference "house-keeping" gene. **B-D:** The tested genes and the house-keeping gene in the RNA-seq data at different x-axis zoom levels and for raw TPM read-count data in (B) and log2-fold change over the mean of the two pre-induction samples (C, D). Notably, *rpoA* expression in RNA-seq data increases with a periodic pattern and this should have affected the RT-qPCR measurements. Considering this, the RT-qPCR and RNA-seq data are consistent: gyrA initially decreases and gyrB increases (< 1 h); later (< 3 d) gyrA increases more (than *rpoA*) and gyrB less (than *rpoA*); followed by a phase of roughly equal expression (4-5 d) and again bifurcation of expression values (> 6 d).

Figure S12. Clustering & Total Read-Count Distribution. A: Bayesian Information Content (BIC) as reported by `flowClust` for clustering of selected scaled components $X'_{k=1,\ldots,6}$ of the Discrete Fourier Transform (DFT, eq. 5)) of the raw TPM data over varying number of cluster centers (K). The maximal BIC was reached for a classification into $K = 6$ distinct clusters (co-expression cohorts). This clustering was chosen for further analysis. **B:** real and imaginary parts of the DFT that were used for clustering (R package `flowClust` [71]). Colors already indicate the final cluster assignments of each transcript at $K = 6$ (A). **C:** Cluster medians (solid lines) of the relative transcript abundances (rel. abund.). For each transcript the $\log_2$ of the ratio of read-counts at time points i to mean of the two samples before induction ($i = 1, 2$, at $-1\,\mathrm{d}$ and $-1\,\mathrm{h}$) was calculated (points indicate the sampling time points i). The transparent ranges indicate the 25% and 75% quantiles of each cluster. Only the time points within to two vertical lines were used for clustering. **D:** Cluster-wise distributions (boxplots) of minimal (left) and maximal (right) read-count values (TPM) of each transcripts.

Figure S13. Co-Expression Cohorts in the Endpoint RNA-seq and Construction of t-Test Profiles. A: Distributions of the $\log_2$ fold-change of transcript abundances in the three strain endpoint experiment for each of the co-expression clusters derived from the topA^OX time series data. The gray background shows the distribution of all other transcripts. The y-axis are the counts for the colored distributions, while the gray background distributions are densities (without axis). For each cluster a t-test was performed (base R function `t.test`) against all transcripts not in the cluster, and the cluster sizes n, and the t-values and the p-values from each test are shown in each plot. The total number of transcripts with expression values were 3676 for gyrA^KD and gyrB^KD, and 3680 for topA^OX. **B:** A t-test profile plot is constructed from the t-test results in (A). A negative t-value indicates that the tested cluster transcripts have a lower mean abundance than all other transcripts and this is indicated by a red color field, the rounded t-value is shown in the fields; blue indicates a positive t-value and higher mean abundance. The p-value is converted to a transparency value for the red and blue colors (along a color palette from red/blue to white), such that the full color is reached for $p \leq p_{\text{min}}$, and for higher p-values the transparency scales with $\log_2(p)$. Both, for visibility of the text and to indicate an additional p-value cut-off the text (t-values) is plotted in white if $p \leq p_{\text{text}}$. The bottom legend shows 5 p-values (text: $\log_{10}(p)$) and the resulting field and text colors. Here $p_{\text{min}} = 10^{-10}$ and $p_{\text{text}} = 10^{-5}$.

Figure S14. `CyanoBase` **Category Analysis of Co-Expressed Cohorts**. **A:** Sorted enrichment profile of functional category annotations as in Figure 5E (colored with $p_{min} = 10^{-10}$ and $p_{text} = 10^{-5}$) but sorted at $p_{sort} = 0.1$. All categories below the red line had only $p > p_{sort}$ and are unsorted. Some abbreviations of the original annotation terms are used for readability of the plot: synth. - synthesis, mod. - modification, repl. - replication, transcr. - transcription, recomb. - recombination, restr. - restriction, s. - saccharides, assim. - assmilation, & - and. **B:** Overlap enrichement and t-test profiles with clusterings as Figure 6C but for additional gene classifications from other publications; from top to bottom: experimental conditions with maximal expression of transcription units from ref. [46], stress and novobiocin (NB) treatment (same as in Fig. 6C) from ref. [4], the original non-collapsed clustering of protein abundance level response to growth rate from ref. [14], and a clustering of a diurnal transcriptome data set from the supplemental material of ref. [34].

Figure S15. GO Analysis of Co-Expressed Cohorts. Sorted enrichment profile as in Figures 5E and S14A (colored with $p_{\text{min}} = 10^{-10}$ and $p_{\text{text}} = 10^{-5}$) but for Gene Ontology (GO) terms, downloaded from the `UniProt` database (2021-03-20, `organism:1111708`). Rows are cut and sorted along columns at $p_{\text{sort}} = 0.01$.

Figure S16. Transcriptome Time Series - Plasmids. Top left panel: Enrichment profile of time series clusters with the locations on the chromosome, one of the seven endogenous plasmids, or our construct pSNDY [36] (Table 1). All other panels show the temporal transcript abundance profiles for the coding genes of each plasmid (see top right legends for plasmid names); each transcript is colored according to its cluster label.

Figure S17. Diurnal Co-Expression Cohorts. Clustering of diurnal transcriptome data from ref. [44] into 5 co-expression cohorts, see Methods for details. **A:** Cluster medians of the normalized (to mean 0) expression values with an additional moving average over 3 samples. Transparent ranges show the 10% and 90% quantiles, *i.e.* they encompass 80% of all values in a cluster. Cluster labels and sizes (number of genes) are indicated on the right y-axis. The gray and white bars on the top indicate dark and light phases of the experiment. **B:** Enrichment profiles of co-expressed cohorts with `CyanoBase` functional categories as for Figure 5E (colored with $p_{\text{min}} = 10^{-10}$ and $p_{\text{text}} = 10^{-5}$), but cut and sorted at $p_{\text{sort}} = 0.05$.

A: *kai* Clock Genes

B: Circadian Response Regulators

C: Sigma Factors

Figure S18. Genes of Interest: Regulators. Transcript abundance profiles of a selected groups of genes; from left to right: the full time-course, a zoom on the first three days, and expression changes with respect to the EVC in the endpoint experiments (as in Fig. 4D,E). The gray background in the time series plots shows the 92.5 % and 7.5 % quantiles of all data as a reference, the gray dots in the endpoint experiments are all other genes and the gray scale indicates local density. Some genes with strong response are specifically mentioned in the following. **A, Clock Genes:** During the first half day post-induction *kaiB1* was upregulated, then quickly downregulated with most other *kai* genes. Only *kaiC3* was upregulated, reflecting the pattern of the RB/dawn cohort, and then remained slightly over-expressed until the last sampled time point. At 3 d *kaiB2* and *kaiC2* were upregulated. **B, Signaling:** The circadian clock output regulator *rpaA* was only slightly upregulated, while its paralog *rpaB* was downregulated. **C, Sigma Factors:** Most sigma factors [48] were downregulated, except for the group 3/4 factors *sigH*, peaking at 1 d, followed by *sigI* (2 d–3 d), and a late peak (10 d, when cells were already enlarged) of the group 2 and stress-response sigma factor *sigB*.

Figure S19. Genes of Interest: Photosystem I & II. As Figure S18 but for genes encoding for the photo-systems. **B/C, PSII:** *psb28-2*: photosystem II reaction center protein Psb28 homologue, extrinsic protein of photosystem II, *psb28-1* but not *psb28-2* was required in PSII recovery after high-light [75].

Figure S20. Genes of Interest: Phycobilisome. As Figure S18 but for genes encoding for the phycobilisome. Phycobilisome genes were downregulated in all experiments, except for the topA^{OX} endpoint measurements (right panels).

A: RuBisCo

B: Carbon Concentrating Mechanism

Figure S21. Genes of Interest: RuBisCo, Carbon Concentrating Mechanism. As Figure S18 but for genes encoding for RuBisCo and carbon concentrating mechanisms. **A, RuBisCo:** large and small subunits of RuBisCo (*rbcL/S* and the assembly factor (*rbcX* were all upregulated. *ecaB*: carbonic anhydrase; *slr1347*: carbonic anhydrase; *sll1636*: gamma carbonic anhydrase. One carbonic anhydrase *slr1347* was upregulated, while the other two were downregulated.

Figure S22. Genes of Interest: Metabolism. As Figure S18 but for genes that encode for metabolic enzymes/pathways. **A, Glycogen:** The glycogen degrading enzyme *glgP1* [76] was strongly downregulated, but *glgP2* upregulated at 3 d. The glucose-1-phosphate adenylyltransferase *glgC* was upregulated with the `RB/dawn` cohort, but both glycogen synthase genes, *glgA/A2*, where downregulated. The glycogen debranching enzymes, *glgX/X2*, were both upregulated at 3 d. **B, NADH dehydrogenase:** selected genes from the NADH dehydrogenase complex that showed a strong response, notably, only in the topA^{OX} strain but not the gyr^{KD} strains; their annotations are *ndhD2 (slr1291)*: electron transfer from NADH to plastoquinone; *ndhD5 (slr2007)*: Na-proton antiporter; *ndhH/ndhF1 (slr0261/slr0844)*: NAD(P)H-quinone oxidoreductase subunit H (chain 7) and chain 5. Notably, a knock-out of *pgmA* had impaired growth in photomixotrophic conditions, with frequent revertant mutations in NADH dehydrogenase subunits [49]. **C, NAD metabolism:** *nadA (sll0622)*: quinolinate synthasegenes involved in NAD metabolism, *pntA/B (slr1239/slr1434)*: transhydrogenase, transfers H from NADPH to NAD. **D, Thioredoxins:** The thioredoxin *trxM1* is strongly down-regulated, while *trxA* has a short peak at 2 d–3 d.

Figure S23. Transcription Start Site Analysis. A: Clustering of transcription units (TU) defined by [46]. Average expression was calculated for all TU from the expression of coding genes they encompass (*via* the "Sense.tags" column of the original data set), and the resulting TU time-series was clustered by k-means, using cluster centers from the CDS clustering (Fig. 5) and identical time-series processing. **B:** enrichment profile of the original CDS clustering (y-axis) with the TU-based re-clustering; colored with $p_{min} = 10^{-10}$ and $p_{text} = 10^{-5}$ and with the original order. **C:** enrichment profile of the time-series clusters with the original TU classification by condition of maximal expression (column "Max.cond.") by [46]; colored with $p_{min} = 10^{-10}$ and $p_{text} = 10^{-5}$ but using the original orders. Note, that many CDS were re-assigned to different clusters but the overall pattern is reproduced.