Effect of stimulation time on the expression of human macrophage polarization markers

Duygu Unuvar Purcu[1,4], Asli Korkmaz[2,¶], Sinem Gunalp[2,¶], Derya Goksu Helvaci[3], Yonca Erdal[4], Yavuz Dogan[5], Gerhard Wingender[4,6], Duygu Sag[2,4,7*]

[1]Department of Molecular Medicine, Health Sciences Institute, Dokuz Eylul University, Izmir, Turkey

[2]Department of Genomic Sciences and Molecular Biotechnology, Izmir International Biomedicine and Genome Institute, Dokuz Eylul University, Izmir, Turkey

[3]Faculty of Medicine, Dokuz Eylul University, Izmir, Turkey

[4]Izmir Biomedicine and Genome Center, Izmir, Turkey

[5]Department of Microbiology, Faculty of Medicine, Dokuz Eylul University, Izmir, Turkey

[6]Department of Biomedicine and Health Technologies, Izmir International Biomedicine and Genome Institute, Dokuz Eylul University, Izmir, Turkey

[7]Department of Medical Biology, Faculty of Medicine, Dokuz Eylul University, Izmir, Turkey

* Corresponding author

E-mail: duygu.sag@ibg.edu.tr

¶These authors contributed equally to this work.

Abstract

Macrophages are highly plastic cells that can polarize into functionally distinct subsets *in vivo* and *in vitro* in response to environmental signals. The development of protocols to model macrophage polarization *in vitro* greatly contributes to our understanding of macrophage biology. Macrophages are divided into two main groups: Pro-inflammatory M1 macrophages (classically activated) and anti-inflammatory M2 macrophages (alternatively activated), based on several key surface markers and the production of inflammatory mediators. However, the expression of these common macrophage polarization markers is greatly affected by the stimulation time used. Unfortunately, there is no consensus yet regarding the optimal stimulation times for particular macrophage polarization markers in *in vitro* experiments. This situation is problematic, (i) as analysing a particular marker at a suboptimal time point can lead to false-negative results, and (ii) as it clearly impedes the comparison of different studies. Using human monocyte-derived macrophages (MDMs) *in vitro*, we analysed how the expression of the main polarization markers for M1 (CD64, CD86, CXCL9, CXCL10, HLA-DR, IDO1, IL1β, IL12p70, TNF), M2a (CD200R, CD206, CCL17, CCL22, IL-10, TGM2), and M2c (CD163, IL-10, TGFβ) macrophages changes over time at mRNA and protein levels. Our data establish the most appropriate stimulation time for the analysis of the expression of human macrophage polarization markers *in vitro*. Providing such a reference guide will likely facilitate the investigation of macrophage polarization and its reproducibility.

Introduction

Macrophages are phagocytes that play essential roles in both innate and adaptive immunity and are involved, for example, in host defence, regulation of metabolism, and tissue remodelling [1–3]. An essential feature of macrophages is their plasticity, whereby varying environmental signals can induce the differentiation of macrophages into functional distinct subsets [4–6]. Mirroring the Th1/Th2 paradigm, macrophages can polarize into classically activated M1 or alternatively activated M2 phenotypes [4,7,8]. Although the M1/M2 dichotomy appears to be a simplification of the *in vivo* situation, it is a helpful experimental distinction that greatly contributed to our understanding of macrophage biology [9].

M1 macrophages are proinflammatory cells that display anti-tumour and anti-microbial activities [10,11]. M1 activation is induced by IFNγ alone or in combination with LPS or TNF [8,12]. M1 macrophages secrete large amounts of pro-inflammatory cytokines such as TNF, IL-1β, IL-6, IL-12, and IL-23, as well as chemokines like CXCL9 and CXCL10. They also express MHC class II (e.g. HLA-DR in humans) and costimulatory molecules including CD80 and CD86 [13–15]. In addition, M1 macrophages upregulate the expression of the tryptophan catabolizing enzyme IDO1 and of SOCS3, a suppressor of M2 cytokine signaling [8,16].

M2 macrophages are anti-inflammatory cells that play a role in angiogenesis, wound healing, and tissue remodelling [17,18]. M2 macrophages can further be divided into M2a, M2b, M2c, and M2d subtypes [14,19,20]. M2a macrophages can be induced by IL-4 and/or IL-13 and they express CD206 (MRC1), CD200R, the decoy receptor IL-1RII, TGM2 (transglutaminase 2) and the chemokines CCL17, CCL22 and CCL24. M2b macrophages can be induced by immune complexes, and TLR agonists, or IL-1

receptor ligands and they can express IL-6, IL-10, IL-1β, and TNF [12,20,21]. M2c macrophages can be induced by IL-10, glucocorticoids, or TGFβ and express CD163, IL-10, and TGFβ [12,22]. Finally, M2d macrophages can be induced by TLR agonists or adenosine receptor activation and they express IL-10, IL-12, TGFβ, and TNF [23,24].

Significant differences between mice and humans for the expression of macrophage polarization markers have been noted. For instance, while murine M1 macrophages express inducible NO synthase (iNOS), which is involved in L-arginine metabolism, human M1 macrophages do not [25]. Moreover, murine M2a macrophages induced by IL-4 and/or IL-13 express Arg1, Ym1, and Fizz1, but these genes do not have human homologs [14,19,26,27]. Due to these differences, it is unclear to which extent data on murine M1/M2 polarization mimic human macrophage biology. Although monocytic cell lines such as THP-1 are commonly used for human macrophage studies, these cells display some substantial differences compared to primary human macrophages [28]. For example, PMA-treated THP-1 macrophage-like cells do not express HLA-DR and CD206 but show higher expression of CD14 and IL-1β compared to primary human macrophages [29]. Furthermore, it was reported that the response to LPS stimulation by THP-1 cells is weaker than by primary cells [30] and the expression of CD14 and cytokines by THP-1 cells was sensitive to the culture conditions such as the cell density [31]. Therefore, experiments using primary human macrophages are most suitable to understand macrophage polarization in humans.

While there is largely agreement on which markers are best for the analysis of macrophage polarization [8,24,32,33], this is not the case for the optimal time point to do so. However, the consideration of the stimulation time is important as the expression of common macrophage polarization markers greatly changes over time.

This leads to the unfortunate situation that many studies report on the inability of a particular macrophage subset to respond, that may well be false negative data as a suboptimal time point for the analysis was chosen. This lack of consensus on the optimal stimulation time of each polarization marker, greatly complicates the comparison of different studies and their findings. However, to our best knowledge, no comprehensive time course analysis of the common polarization markers has been published. Here we report a detailed time course of the expression of the main M1, M2a, and M2c polarization markers of human monocyte-derived macrophages (MDMs) in vitro at both mRNA and protein levels. Our data establish the optimal time points for the analysis of the expression of these polarization markers, which will greatly facilitate their investigation.

Materials and Methods

Generation of human monocyte-derived macrophages (MDMs)

The ethical approval for buffy coats was obtained from the "Non-Interventional Research Ethics Committee" of the Dokuz Eylül University (Approval number: 2019/02-39). The buffy coats from healthy donors were obtained from the Dokuz Eylul University Blood Bank (Izmir, Turkey) after written consent. Human monocytes were purified from buffy coats by two gradient centrifugations: PBMCs were initially isolated with Ficoll-Paque (GE Healthcare, Pittsburgh, PA) and subsequently peripheral monocytes were isolated with Percoll (GE Healthcare, Pittsburgh, PA) [34]. To generate human monocyte-derived macrophages (MDMs), monocytes were treated for 7 days with 10 ng/mL human recombinant M-CSF (PeproTech, Rocky Hill, NJ) in

R5 medium (RPMI 1640 (Gibco, Thermo Fisher Scientific, Waltham, MA, USA) supplemented with 5% heat inactivated fetal bovine serum (Gibco, Thermo Fisher Scientific, Waltham, MA, USA), 1% Penicillin-Streptomycin (Gibco, Thermo Fisher Scientific, Waltham, MA, USA)) in ultra-low attachment six-well plates (Corning Life Sciences, Tewksbury, MA) [35]. On day 7, the mature macrophages were collected from the low attachment plates and verified to be more than 90% CD68[+] by flow cytometry.

Polarization of human MDMs

Human MDMs were plated in 24 well plates at 6.5×10^5 cells per well in 1 mL R5 medium and allowed to rest overnight at 37 °C before stimulation. Then, the human MDMs (M0 macrophages) were either left unstimulated or stimulated for 4h, 8h, 12h, 24h, 48h, or 72 h to induce macrophage polarization: (i) for M1 polarization with 100 ng/mL LPS (Ultrapure; InvivoGen, San Diego, CA) and 20 ng/mL IFNγ (R&D, Minneapolis, MN), (ii) for M2a polarization with 20 ng/mL IL-4 (R&D, Minneapolis, MN), or (iii) for M2c polarization with 20 ng/mL IL-10 (R&D, Minneapolis, MN).

ELISA

After stimulations, supernatants were collected and TNF (Invitrogen, Thermo Fisher Scientific, Waltham, MA, USA), IL-12 (BioLegend, San Diego, CA), IL-10 (BioLegend, San Diego, CA), IL-1β (BioLegend, San Diego, CA), and TGFβ (R&D, Minneapolis, MN) concentrations were determined by ELISA according to the manufacturer's instructions.

Quantitative Real Time PCR

Total RNA of macrophages was isolated using the Nucleo-Spin RNA kit (Macharey Nagel, Germany). RNA purity and quantity were evaluated with Nanodrop spectrophotometer (Thermo Scientific, Rockford, IL). cDNA synthesis was performed using EvoScript Universal cDNA Master, using 1000 ng of RNA per sample (Roche, Basel, Switzerland). Quantitative PCR (qPCR) was performed with a LightCycler 480 II real time system (Roche, Basel, Switzerland) using Fast Start Essential DNA Probes Master (Roche, Basel, Switzerland) and the following RealTime ready Single Assays (Roche, Basel, Switzerland) for IDO1 (#100134768), CXCL10 (#100134759), TNF (#100134777), CXCL9 (#100137998), MRC1 (#100134731), TGM2 (#100134722), CCL17 (#100138007), CCL22 (#100134713), IL-10 (#100133437), CD163 (#100134801), and ACTB (#100074091). RNA isolation, cDNA synthesis, and the quantitative PCR were performed according to the manufacturers' recommendations. Relative mRNA expression was calculated using the $2^{-\Delta\Delta Ct}$ method [36] and β-actin was used for normalization as housekeeping gene.

Flow cytometry

Macrophages were detached from the cell plate surface using StemPro Accutase Cell Dissociation Reagent (Gibco; Thermo Fisher Scientific, Waltham, MA, USA) according to manufacturer's instructions. Single-cell suspensions were stained in flow cytometry staining buffer (PBS, 1% bovine serum albumin, 0.1% sodium azide). Zombie UV Fixable Viability Kit (BioLegend, San Diego, CA, USA) was used to determine cell viability. Fcγ receptors were blocked with Human TruStain FcX antibody

(BioLegend, San Diego, CA) for 10 min and surface antigens on cells were stained for 40 min at 4 °C. Following anti-human antibodies were used, with their fluorochrome, clone, and dilution: CD86-BV605 (IT2.2; 1:400), HLA-DR-APC-Cy7 (L243; 1:400), CD64-PerCP-Cy5.5 (10.1; 1:200), CD200R-PE-Dazzle594 (OX-108; 1:400), CD206-AF700 (15-2; 1:400), CD163-PE-Cy7 (GHI/61; 1:200). Antibodies were purchased from BioLegend (San Diego, CA, USA). Cell fluorescence was measured using an LSR Fortessa (BD Biosciences) and the data were analyzed with FlowJo software (TreeStar, Ashland, OR). Forward- and side-scatter parameters were used to exclude doublets from the analysis.

Statistical Analysis

Results are expressed as mean ± SEM. All experimental data were analyzed and graphs were generated using Graph Pad Prism 8 (GraphPad Software, CA, USA). The p-values were calculated by two-tailed paired Student's t test or Wilcoxon matched-pairs signed rank test when appropriate. P values less than 0.05 was considered statistically significant.

Results

Time course dependent changes in the expression of polarization markers at mRNA level in M1, M2a, and M2c macrophages

Primary human MDMs were polarized into an M1 (100 ng/mL LPS, 20 ng/mL IFNγ), an M2a (20 ng/mL IL-4), or an M2c (20 ng/mL IL-10) phenotype for 4, 8, 12, 24, 48, or 72 h. The expression of M1, M2a, or M2c markers was analyzed at mRNA level by qPCR. Unpolarized human MDMs were used as control. In M1 macrophages, the mRNA expression of the M1 markers CXCL9, CXCL10, and TNF reached their highest level after 4 h of stimulation with a gradually decrease thereafter (Figs 1A-1C). In contrast, the IDO1 expression in M1 macrophages continuously increased during the time course reaching its highest value at 48-72 h (Fig 1D).

In M2a macrophages, the M2a markers MRC1 (CD206) and TGM2 were induced after 4 h without major changes until the 48 h time point. A further increase for both was noted at 72 h (Figs 1E and 1F). Compared to the other M2a markers, CCL17 expression was very high at all time points (Fig 1G). The expression of CCL22 in M2a macrophages gradually increased from 4h to 48h (Fig 1H). In M2c macrophages, the M2c markers IL-10 and CD163 were analyzed. The expression of IL-10 mRNA peaked at 4h (Fig 1I), whereas the expression of CD163 mRNA was comparable between 4 h and 72 h (Fig 1J).

These data indicate that 4 h is the optimal stimulation time at the mRNA level for M1 markers except IDO1, which peaks after 48 h. While for M2c markers 4-8 h of stimulation seems ideal, for M2a markers longer stimulation is needed as these markers peaked around 48-72 h (Table 1).

Table 1. Suggested stimulation times for the detection of M1, M2a, and M2c polarization markers in human monocyte-derived macrophages.

Macrophage Phenotype	Stimulant	Detection Method	Marker	Type	Suggested Stimulation Time for Optimal Detection
M1	LPS + IFNγ	qPCR	CXCL9	Chemokine	4 h
			CXCL10	Chemokine	4 h
			TNF	Cytokine	4 h
			IDO1	Enzyme	48-72 h
		Flow Cyt.	CD86	Co-stimulatory M.	8-12 h
			CD64	Fc Receptor	48-72 h
			HLA-DR	MHC class II M.	8-12 h
		ELISA	TNF	Cytokine	8 h
			IL12p70	Cytokine	12 h
			IL-1β	Cytokine	8-12 h
M2a	IL-4	qPCR	MRC1	Mannose R.	72 h
			TGM2	Enzyme	72 h
			CCL17	Chemokine	4-72 h
			CCL22	Chemokine	48-72 h
		Flow Cyt.	CD200R	Inhibitory R.	48-72 h
			CD206	Mannose R.	48-72 h
		ELISA	IL-10	Cytokine	48-72 h
M2c	IL-10	qPCR	IL-10	Cytokine	4 h
			CD163	Scavenger R.	8-24 h
		Flow Cyt.	CD163	Scavenger R.	24-72 h
		ELISA	TGFβ	Cytokine	72h

Flow Cyt., flow cytometry; R, receptor; P, protein; M, molecule.

Time course depended changes in the expression of surface markers at protein level in M1, M2a and M2c macrophages

Next, to determine the impact of stimulation time on the expression of surface markers, the primary human MDMs were analyzed for the expression of M1, M2a or M2c surface markers by flow cytometry. In human MDMs polarized into an M1 phenotype, the CD86 expression was, surprisingly, bimodal, with peaks after 12 h and 72 h of stimulation and at background levels at the 24 h time point (Fig 2A). The CD64 expression increased from 8 h to 72 h, with the highest levels observed after 48 h and 72 h (Fig 2B). However, no significant changes for the expression of HLA-DR were noted on M1 macrophages (Fig 2C).

In M2a polarized macrophages, expression of CD200R and CD206 (MRC1) significantly increased starting from 8h of stimulation with the highest levels reached after 48 - 72 h (Figs 2D and 2E). In M2c polarized macrophages, the expression of CD163 was significantly increased after 24 h of stimulation and remained stable thereafter (Fig 2F).

These data indicate that 12 h of stimulation is best for the M1 surface markers CD86 and HLA-DR, whereas 48-72 h of stimulation is needed for the M1 marker CD64 and for the M2 markers CD163, CD200R, and CD206. (Table 1).

Time course depended changes in cytokine production by M1, M2a and M2c macrophages

Finally, the production of key cytokines by the M1, M2a, or M2c polarized human MDMs was analyzed by flow cytometry. TNF production by M1 macrophages peaked

at 4, 8, and 12 h of stimulation, with a slight decrease at later time points (Fig 3A). IL-12 production was low at 4 h but prominent and stable between 8 h to 72 h of stimulation (Fig 3B). In contrast, IL-1β production by M1 macrophages was only significantly increased at the 8 h and 12 h time points (Fig 3C). In M2a macrophages, a significant production of cytokines was only observed after 48 h (IL-10) and 72 h (IL-10, TGFβ) of stimulation (Figs 3D and 3E).

These data indicate that 8-12 h of stimulation is best for the detection of the M1 cytokines IL-1β, IL-12p70, and TNF, whereas 72 h of stimulation is needed for the detection of the M2 cytokines IL10 and TGFβ (Table 1).

Discussion

Macrophages can polarize into different functional phenotypes according to the signals they receive from the environment. Environmental signals that induce macrophage polarization in vivo can be mimicked in in vitro studies [5,6,37]. Although the stimulation time affects the expression levels of most markers used to analyze M1 and M2 polarization, there is no consensus in the literature regarding the timing to analyze particular markers during in vitro experiments. Here we report a detailed time course of M1, M2a, and M2c polarization markers of primary human macrophages that determine the optimal time points for their detection at mRNA and protein levels.

Our results demonstrate a dramatic increase in the expression of the chemokines CXCL9 and CXCL10 and the cytokine TNF at mRNA level in the first 4 h of stimulation of primary MDMs polarized into the M1 phenotype with LPS-IFNγ (Figs 1A-1C). In M1 polarization, the upward trend in gene expression decreased with prolonged stimulation, for these genes. Interestingly, in contrast to the other M1 markers tested, IDO1 expression was increased with prolonged stimulation, reaching its highest value between 48 and 72 h (Fig 1D). Since IDO1 is a regulatory enzyme, that is induced after inflammation [38], it is understandable that it is induced later than cytokines and chemokines. Our results are in accordance with previous studies, in which macrophages were stimulated for 48 h to determine IDO1 expression [39,40]. Based on our results, 4 h LPS-IFNγ stimulation, seems to be the optimum time point to observe the changes in M1 markers at mRNA level. Although IDO1 mRNA expression peaked at late stimulation points, its expression was also quite high at 4 h of stimulation (Fig 1D). In M2a macrophages, CCL17 expression was high at all time points studied, while CCL22 expression was increased from 4 h to 48 h of stimulation. (Figs 1G-1H).

The surface markers CD64, CD86, and HLA-DR are commonly used as M1 polarization markers [19]. In our study, the expression of CD86 on M1 macrophages showed the strongest increase at the 8-12 h time points (Fig 2A). Although we observed a slight increase of HLA-DR expression after 4 h to 12 h of stimulation, this did not reach statistical significance (Fig 2C). This was most probably due to the high variation between donors. On the contrary, HLA-DR expression at 24 and 48 h of stimulation was lower in M1 macrophages compared to the unpolarized MDMs, similar to previous studies [41]. We observed a similar pattern for CD86 expression. Stimulation of macrophages with LPS leads to the production IL-10 that inhibits CD86 in an autocrine manner to prevent overexpression of CD86 [42]. Therefore, the observed decrease in CD86 and HLA-DR could be due to the down-regulation of these receptors by IL-10 [42–44]. It has been reported that CD86 expression is higher in both M1 and M2 macrophages than unpolarized MDMs at 48 h [41], and 72 h [13] of stimulation. Based on our results, the optimum stimulation time for CD86 and HLA-DR to be considered as M1 polarization markers is 12 h, and their expression in M1 macrophages decreases thereafter. In contrast, CD64 expression on M1 macrophages was increased from 8 h to 72 h of stimulation (Fig 2B), unlike HLA-DR and CD86 expression. CD64 expression has been reported to be consistently high on M1 macrophages over a wide range of stimulation times [13,37,45]. CD64 has been defined as one of the most distinctive M1 polarization markers [37].

While CD206 is a recognized M2a marker for murine macrophages [46], it was suggested that it is not a specific marker for M2a polarization in humans. [45,47]. It has been reported that MDMs polarized into M2a phenotype for 24 h, upregulated CD200R but not CD206 [47]. According to our results, the expression of the surface markers CD200R and CD206 gradually increased over time in human MDMs polarized into M2a

with IL-4, reaching a maximum level at 48 and 72 h of stimulation (Figs 2D and 2E). Expression of the mannose receptor CD206 (MRC1) at mRNA level also peaked at 72 h of stimulation in M2a macrophages (Fig 2E).

The scavenger receptor CD163 is a specific M2c polarization marker (Vogel et al., 2014). In our study, the CD163 expression on IL-10 stimulated M2c macrophages was significantly increased after 24-72 h of stimulation (Fig 2F), which is consistent with the previous studies [37,41,45]. In M2c macrophages, CD163 gene expression was also increased at 8-24 h of stimulation (Fig 1J), in accordance with this, CD163 surface marker expression peaked at later hours as expected (Fig 2F)

We also examined the impact of time on the cytokine production of polarized MDMs. The production of IL-1β, IL12p70, and TNF was high after a wide range of stimulation times in M1 macrophages with 8-12 h of stimulation being optimal (Figs 3A-3C). A previous study has reported TNF and IL12p40 production in M1 macrophages after 48 h of stimulation, while no IL-1β production was detected [37]. In M2a macrophages, we detected IL-10 after 48 to 72 h of stimulation (Fig 3D). It has been reported that IL-10 production by M1 macrophages was higher than M2 macrophages at 24 h of stimulation [47]. In our hands, the production of IL-10 in M2 macrophages at 24 h of stimulation was still at background levels, only after 48 h the IL-10 production was prominent (Fig 3D). In M2c macrophages, 72 h of stimulation was necessary for prominent production of TGFβ (Fig 3E).

In summary, we report here the optimal time points for the analysis of the expression of the main M1, M2a, and M2c polarization markers. It is unfortunate that there is no consensus yet regarding the optimal stimulation times for particular macrophage polarization markers in in vitro experiments, as this can lead to false-negative results and impedes the comparison of different studies. Therefore, our study provides useful

guideline for the optimal polarization times of primary human macrophages, which will likely facilitate the investigation of macrophage polarization and its reproducibility.

Acknowledgements

We thank Dr. Safiye Nese Atabey and Dr. Mehtap Yuksel Egrilmez for valuable scientific contributions. We thank the Flow Cytometry and Cell Sorting Facility at the Izmir Biomedicine and Genome Center for excellent technical assistance. This work was funded by the Co-Funded Brain Circulation Scheme (MSCA/TUBITAK, #115C074), the Dokuz Eylul University (#2017.KB.SAG.011), the Turkish Academy of Sciences (TUBA-GEBIP award), and the Science Academy, Turkey (BAGEP award) (all to D.S).

References

1. Stout RD, Suttles J. Functional plasticity of macrophages: reversible adaptation to changing microenvironments. J Leukoc Biol. 2004;76: 509–513. doi:10.1189/jlb.0504272

2. Mosser DM, Edwards JP. Exploring the full spectrum of macrophage activation. Nat Rev Immunol. 2008;8: 958–969. doi:10.1038/nri2448

3. Wynn TA, Chawla A, Pollard JW. Macrophage biology in development, homeostasis and disease. Nature. Nature Publishing Group; 2013. pp. 445–455. doi:10.1038/nature12034

4. Biswas SK, Mantovani A. Macrophage plasticity and interaction with

lymphocyte subsets : cancer as a paradigm. Nat Immunol. 2010;11: 889–896. doi:10.1038/ni.1937

5.   Sica A, Mantovani A. Macrophage plasticity and polarization: In vivo veritas. Journal of Clinical Investigation. American Society for Clinical Investigation; 2012. pp. 787–795. doi:10.1172/JCI59643

6.   Akira S, Misawa T, Satoh T, Saitoh T. Macrophages control innate inflammation. Diabetes, Obesity and Metabolism. Blackwell Publishing Ltd; 2013. pp. 10–18. doi:10.1111/dom.12151

7.   Mills CD, Kincaid K, Alt JM, Heilman MJ, Hill AM. M-1/M-2 macrophages and the Th1/Th2 paradigm. J Immunol. 2000;164: 6166–73. Available: http://www.ncbi.nlm.nih.gov/pubmed/10843666

8.   Martinez FO, Gordon S. The M1 and M2 paradigm of macrophage activation: time for reassessment. F1000Prime Rep. 2014;6: 13. doi:10.12703/P6-13

9.   Locati M, Curtale G, Mantovani A. Diversity, Mechanisms, and Significance of Macrophage Plasticity. Annu Rev Pathol Mech Dis. 2020;15: 123–147. doi:10.1146/annurev-pathmechdis-012418-012718

10. Sica A, Larghi P, Mancino A, Rubino L, Porta C, Totaro MG, et al. Macrophage polarization in tumour progression. Seminars in Cancer Biology. 2008. pp. 349–355. doi:10.1016/j.semcancer.2008.03.004

11. Atri C, Guerfali FZ, Laouini D. Role of human macrophage polarization in inflammation during infectious diseases. International Journal of Molecular Sciences. MDPI AG; 2018. doi:10.3390/ijms19061801

12. Martinez FO, Sica A, Mantovani A, Locati M. Macrophage activation and polarization. Front Biosci. 2008;13: 453–61. doi:10.2741/2692

13. Beyer M, Mallmann MR, Xue J, Staratschek-Jox A, Vorholt D, Krebs W, et al.

High-Resolution Transcriptome of Human Macrophages. Zirlik A, editor. PLoS One. 2012;7: e45466. doi:10.1371/journal.pone.0045466

14. Wang N, Liang H, Zen K. Molecular mechanisms that influence the macrophage M1-M2 polarization balance. Frontiers in Immunology. Frontiers; 2014. p. 614. doi:10.3389/fimmu.2014.00614

15. Martinez FO, Gordon S, Locati M, Mantovani A. Transcriptional Profiling of the Human Monocyte-to-Macrophage Differentiation and Polarization: New Molecules and Patterns of Gene Expression. J Immunol. 2006;177: 7303–7311. doi:10.4049/jimmunol.177.10.7303

16. Wilson HM. SOCS proteins in macrophage polarization and function. Frontiers in Immunology. Frontiers Research Foundation; 2014. doi:10.3389/fimmu.2014.00357

17. Gordon S. Alternative activation of macrophages. Nature Reviews Immunology. Nature Publishing Group; 2003. pp. 23–35. doi:10.1038/nri978

18. Mantovani A, Biswas SK, Galdiero MR, Sica A, Locati M. Macrophage plasticity and polarization in tissue repair and remodelling. J Pathol. 2013;229: 176–185. doi:10.1002/path.4133

19. Mantovani A, Sozzani S, Locati M, Allavena P, Sica A. Macrophage polarization: Tumor-associated macrophages as a paradigm for polarized M2 mononuclear phagocytes. Trends in Immunology. 2002. pp. 549–555. doi:10.1016/S1471-4906(02)02302-5

20. Mantovani A, Sica A, Sozzani S, Allavena P, Vecchi A, Locati M. The chemokine system in diverse forms of macrophage activation and polarization. Trends in Immunology. Oxford University Press, Oxford; 2004. pp. 677–686. doi:10.1016/j.it.2004.09.015

21.   Wang L xun, Zhang S xi, Wu H juan, Rong X lu, Guo J. M2b macrophage polarization and its roles in diseases. Journal of Leukocyte Biology. John Wiley and Sons Inc.; 2019. pp. 345–358. doi:10.1002/JLB.3RU1018-378RR

22.   Huang X, Li Y, Fu M, Xin HB. Polarizing macrophages in vitro. Methods in Molecular Biology. Humana Press Inc.; 2018. pp. 119–126. doi:10.1007/978-1-4939-7837-3_12

23.   Shapouri-Moghaddam A, Mohammadian S, Vazini H, Taghadosi M, Esmaeili SA, Mardani F, et al. Macrophage plasticity, polarization, and function in health and disease. Journal of Cellular Physiology. Wiley-Liss Inc.; 2018. pp. 6425–6440. doi:10.1002/jcp.26429

24.   Rőszer T, Roszer T. Understanding the Mysterious M2 Macrophage through Activation Markers and Effector Mechanisms. Mediators Inflamm. 2015;2015: 1–16. doi:10.1155/2015/816460

25.   Murray PJ, Wynn TA. Obstacles and opportunities for understanding macrophage polarization. J Leukoc Biol. 2011;89: 557–563. doi:10.1189/jlb.0710409

26.   Gordon S, Martinez FO. Alternative activation of macrophages: Mechanism and functions. Immunity. 2010;32: 593–604. doi:10.1016/j.immuni.2010.05.007

27.   Mantovani A, Locati M. Tumor-associated macrophages as a paradigm of macrophage plasticity, diversity, and polarization lessons and open questions. Arterioscler Thromb Vasc Biol. 2013;33: 1478–1483. doi:10.1161/ATVBAHA.113.300168

28.   Tarique AA, Logan J, Thomas E, Holt PG, Sly PD, Fantino E. Phenotypic,functional,and plasticity features of classical and alternatively activated human macrophages. Am J Respir Cell Mol Biol. 2015;53: 676–688.

doi:10.1165/rcmb.2015-0012OC

29.   Kohro T, Tanaka T, Murakami T, Wada Y, Aburatani H, Hamakubo T, et al. A

        comparison of differences in the gene expression profiles of phorbol 12-

        myristate 13-acetate differentiated THP-1 cells and human monocyte-derived

        macrophage. J Atheroscler Thromb. 2004;11: 88–97. doi:10.5551/jat.11.88

30.   Schildberger A, Rossmanith E, Eichhorn T, Strassl K, Weber V. Monocytes,

        peripheral blood mononuclear cells, and THP-1 cells exhibit different cytokine

        expression patterns following stimulation with lipopolysaccharide. Mediators

        Inflamm. 2013;2013. doi:10.1155/2013/697972

31.   Aldo PB, Craveiro V, Guller S, Mor G. Effect of culture conditions on the

        phenotype of THP-1 monocyte cell line. Am J Reprod Immunol. 2013;70: 80–

        86. doi:10.1111/aji.12129

32.   Murray PJ, Allen JE, Biswas SK, Fisher EA, Gilroy DW, Goerdt S, et al.

        Macrophage Activation and Polarization: Nomenclature and Experimental

        Guidelines. Immunity. 2014;41: 14–20. doi:10.1016/j.immuni.2014.06.008

33.   Martinez FO, Gordon S, Locati M, Mantovani A. Transcriptional Profiling of the

        Human Monocyte-to-Macrophage Differentiation and Polarization: New

        Molecules and Patterns of Gene Expression. J Immunol. 2006;177: 7303–

        7311. doi:10.4049/jimmunol.177.10.7303

34.   Menck K, Behme D, Pantke M, Reiling N, Binder C, Pukrop T, et al. Isolation of

        human monocytes by double gradient centrifugation and their differentiation to

        macrophages in Teflon-coated cell culture bags. J Vis Exp. 2014 [cited 13 Aug

        2020]. doi:10.3791/51554

35.   Mukundan L, Bishop GA, Head KZ, Zhang L, Wahl LM, Suttles J. TNF

        Receptor-Associated Factor 6 Is an Essential Mediator of CD40-Activated

Proinflammatory Pathways in Monocytes and Macrophages. J Immunol. 2005;174: 1081–1090. doi:10.4049/jimmunol.174.2.1081

36. Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time quantitative PCR and the 2-ΔΔCT method. Methods. 2001. doi:10.1006/meth.2001.1262

37. Vogel DYS, Glim JE, Stavenuiter AWD, Breur M, Heijnen P, Amor S, et al. Human macrophage polarization in vitro: Maturation and activation methods compared. Immunobiology. 2014;219: 695–703. doi:10.1016/j.imbio.2014.05.002

38. Kim S, Miller BJ, Stefanek ME, Miller AH. Inflammation-induced activation of the indoleamine 2,3-dioxygenase pathway: Relevance to cancer-related fatigue. Cancer. John Wiley and Sons Inc.; 2015. pp. 2129–2136. doi:10.1002/cncr.29302

39. Wang XF, Wang HS, Wang H, Zhang F, Wang KF, Guo Q, et al. The role of indoleamine 2,3-dioxygenase (IDO) in immune tolerance: Focus on macrophage polarization of THP-1 cells. Cell Immunol. 2014;289: 42–48. doi:10.1016/j.cellimm.2014.02.005

40. Zhao Q, Kuang D-M, Wu Y, Xiao X, Li X-F, Li T-J, et al. Activated CD69 + T Cells Foster Immune Privilege by Regulating IDO Expression in Tumor-Associated Macrophages . J Immunol. 2012;188: 1117–1124. doi:10.4049/jimmunol.1100164

41. Iqbal S, Kumar A. Characterization of In vitro Generated Human Polarized Macrophages. J Clin Cell Immunol. 2015;06: 380. doi:10.4172/2155-9899.1000380

42. De Vries JE. Immunosuppressive and anti-inflammatory properties of

interleukin 10. Ann Med. 1995;27: 537–541. doi:10.3109/07853899509002465

43. Groux H, Bigler M, de Vries JE, Roncarolo MG. Inhibitory and stimulatory effects of IL-10 on human CD8+ T cells. J Immunol. 1998;160.

44. De Waal Malefyt R, Abrams J, Bennett B, Figdor CG, De Vries JE. Interleukin 10(IL-10) inhibits cytokine synthesis by human monocytes: An autoregulatory role of IL-10 produced by monocytes. J Exp Med. 1991;174: 1209–1220. doi:10.1084/jem.174.5.1209

45. Ambarus CA, Krausz S, van Eijk M, Hamann J, Radstake TRDJ, Reedquist KA, et al. Systematic validation of specific phenotypic markers for in vitro polarized human macrophages. J Immunol Methods. 2012;375: 196–206. doi:10.1016/j.jim.2011.10.013

46. Chroneos Z, Shepherd VL. Differential regulation of the mannose and SP-A receptors on macrophages. Am J Physiol - Lung Cell Mol Physiol. 1995;269. doi:10.1152/ajplung.1995.269.6.l721

47. Jaguin M, Houlbert N, Fardel O, Lecureur V. Polarization profiles of human M-CSF-generated macrophages and comparison of M1-markers in classically activated macrophages from GM-CSF and M-CSF origin. Cell Immunol. 2013;281: 51–61. doi:10.1016/j.cellimm.2013.01.010

Figure Legends

Fig 1. Time depended changes in the expression of macrophage polarization markers at mRNA level. Primary human MDMs (M0) were left unstimulated or were polarized to M1 (100 ng/mL LPS, 20 ng/mL IFNγ), M2a (20 ng/mL IL-4), or M2c (20 ng/mL IL-10) for 4, 8, 12, 24, 48, or 72 h. The expression of **(A-D)** CXCL9, CXCL10, TNF, and IDO1 mRNA for M1 polarization; of **(E-H)** MRC1, TGM2, CCL17, and CCL22 mRNA for M2a polarization; and of **(I-J)** IL-10 and CD163 mRNA for M2c polarization were analyzed by qPCR. Data shown are mean ± SEM of biological replicates of 3 donors and representative of two independent experiments with similar results. Polarized macrophages (M1, M2a, or M2c) at all time points were compared with unstimulated (US) M0 macrophages. Statistical analyses were performed with paired Student's t test, * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$ (Statistical comparison of all time points are shown in the S1 Table).

Fig 2. Time depended changes in the expression of macrophage polarization markers at protein level. Primary human MDMs (M0) were left unstimulated or were polarized to M1 (100 ng/mL LPS, 20 ng/mL IFNγ) or M2a (20 ng/mL IL-4) for 4, 8, 12, 24, 48, or 72 h. Surface marker expression **(A-C)** of CD86, CD64 and HLA-DR for M1 polarization; **(D-E)** of CD200R and CD206 (MRC1) for M2a polarization, and **(F)** of CD163 for M2c polarization was analyzed by flow cytometry. Shown are representative histograms (left) and summary data of one of two independent experiments with biological replicates of 3 donors (right). Polarized macrophages (M1, M2a, or M2c) at all time points were compared with unstimulated (US) M0 macrophages. Statistical analyses were performed with paired Student's t test, * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$ (Statistical comparison of all time points are shown in the S2 Table).

Fig 3. Time depended changes in cytokine production by M1, M2a, and M2c macrophages. Primary human MDMs (M0) were left unstimulated or were polarized to M1 (100 ng/mL LPS, 20 ng/mL IFNγ), M2a (20 ng/mL IL-4), or M2c (20 ng/mL IL-10) for 4, 8, 12, 24, 48, or 72 h . Production of **(A-C)** TNF, IL-12 and IL-1β for M1 polarization; of **(D)** IL-10 for M2a polarization, and of **(E)** TGFβ for M2c polarization was analyzed by ELISA. The data shown are mean ± SEM of biological replicates of 3 donors and representative of two independent experiments with similar results. Polarized macrophages (M1, M2a, or M2c) at all time points were compared with unstimulated (US) M0 macrophages. Statistical analyses were performed with paired Student's *t* test or Wilcoxon matched-pairs signed rank test, * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$ (Statistical comparison of all time points are shown in the S3 Table).

Fig 1

Fig 2

Fig 3