Untangling Microbiota Diversity and Assembly Patterns in the World's Largest Water Diversion Canal

Lu Zhang[1,2], Wei Yin[3], Chao Wang[3], Aijing Zhang[4], Hong Zhang[5], Tong Zhang[6] and Feng Ju[1,2*]

[1] Key Laboratory of Coastal Environment and Resources of Zhejiang Province, School of Engineering, Westlake University, 18 Shilongshan Road, Hangzhou 310024, Zhejiang Province, China

[2] Institute of Advanced Technology, Westlake Institute for Advanced Study, 18 Shilongshan Road, Hangzhou 310024, Zhejiang Province, China

[3] Changjiang Water Resources Protection Institute, 515 Qintai Street, Wuhan 430051, Hubei Province, China

[4] Construction and Administration Bureau of South-to-North Water Diversion Middle Route Project, 1 Yuyuantan South Road, Beijing 100038, China

[5] State Key Laboratory of Environmental Aquatic Chemistry, Research Center for Eco-Environmental Sciences, Chinese Academy of Sciences, 18 Shuangqing Road, Beijing 100085, China

[6] Environmental Microbiome Engineering and Biotechnology Laboratory, Department of Civil Engineering, Pokfulam Road, The University of Hong Kong, Hong Kong 999077, China

* Corresponding to:

Dr. Feng Ju (Assistant Professor)

Address:

Environmental Microbiome and Biotechnology Lab (EMBLab)

Westlake University, 18 Shilongshan Road, Hangzhou 310024, China

Tel.: 571-87963205 (lab), 571-87380995 (office)

Fax: 0571-85271986

E-mail: jufeng@westlake.edu.cn

Abstract

Large water diversion projects are important constructions for reallocation of human-essential

water resources. Deciphering microbiota dynamics and assembly mechanisms underlying

canal water ecosystem services especially during long-distance diversion is the prerequisite

for water quality monitoring, biohazard warning and sustainable management. Using a

1432-km canal of the South-to-North Water Diversion Projects as a model system, we answer

three central questions: how bacterial and micro-eukaryotic communities spatio-temporally

develop, how much ecological stochasticity contributes to microbiota assembly, and which

immigrating populations better survive and navigate across the canal. We applied quantitative

ribosomal RNA gene sequence analyses to investigate canal water microbial communities

sampled over a year, as well as null model- and neutral model-based approaches to

disentangle the microbiota assembly processes. Our results showed clear microbiota dynamics

in community composition driven by seasonality more than geographic location, and

seasonally dependent influence of environmental parameters. Overall, bacterial community

was largely shaped by deterministic processes, whereas stochasticity dominated

micro-eukaryotic community assembly. We defined a local growth factor (LGF) and

demonstrated its innovative use to quantitatively infer microbial proliferation, unraveling

taxonomically dependent population response to local environmental selection across canal

sections. Using LGF as a quantitative indicator of immigrating capacities, we also found that

most micro-eukaryotic populations (82%) from the source lake water sustained growth in the

canal and better acclimated to the hydrodynamical water environment than bacteria (67%).

Taxa inferred to largely propagate include *Limnohabitans* sp. and *Cryptophyceae*, potentially

contributing to water auto-purification. Combined, our work poses first and unique insights

into the microbiota assembly patterns and dynamics in the world's largest water diversion

canal, providing important ecological knowledge for long-term sustainable water quality

maintenance in such a giant engineered system.

Key words: Engineered system; Bacterial community; Micro-eukaryote; Deterministic

assembly; Stochastic assembly; Population growth

Introduction

Microorganisms are fundamental components of ecosystems and play vital roles in global biogeochemical cycling (Arrigo, 2005; Battin et al., 2016; Du et al., 2021). Knowledge about composition and dynamics of microbial communities across space and time is a prerequisite for predicting and manipulating the microorganisms in both natural and engineered systems (Bell et al., 2005; Bru et al., 2011; Ju and Zhang, 2015). Due to artificial system design and control, microbial communities in engineered ecosystems have received increasing attention to unravel the community dynamics underlying their eco-environmental or biotechnological services (Briones and Raskin, 2003; Ju et al., 2017; Ju et al., 2014; Vanwonterghem et al., 2014; Wu et al., 2019). Specifically, microorganisms in drinking water systems are closely linked with water auto-purification and eventually to public health (Berry et al., 2006). Thus, comprehensive microbiome monitoring is needed to differentiate baseline and disturbed microbial communities in engineered water systems to facilitate system management (Hull et al., 2019; Sagova-Mareckova et al., 2021).

The South-to-North Water Diversion (SNWD) Projects were built to aid in water supply in the drought-stricken north China, which represent the world's largest water diversion projects. The projects consist of the Eastern Route Project, the Middle Route Project (MRP) and the Western Route Project. The 1432-km MRP went into service in 2014 and has drawn 40 billion m^3 of water from the Danjiangkou Reservoir to He'nan and Hebei Provinces, and Beijing and Tianjin Municipalities, benefiting tens of millions of people. Despite their

important roles in controlling geochemical processes and indicating water quality (Chen et al.,

2018; Dang et al., 2018), little has been known about the microbial constituents throughout

the system (Luo et al., 2019), neither for other engineered long-distance water diversion

canals. Especially, the MRP is a closed engineered system where microorganisms are subject

to potentially less complex biotic and abiotic interactions, while the numerous hydraulic

structures and engineered perturbations (e.g., inverted siphon, aqueduct and tunnels) along the

main canal add to challenges to model microbial community dynamics, calling for

comprehensive investigation of microbial community patterns in the MRP canal, to assist in

establishing integrative assessment of water quality with microbial community monitoring

(Sagova-Mareckova et al., 2021).

Understanding processes and mechanisms driving the community patterns underlying

ecosystem services is one of the key issues in microbial ecology (Zhou et al., 2014).

Microbial community composition in aquatic ecosystems can be affected deterministically by

both abiotic or environmental factors, e.g., water temperature, pH, dissolved oxygen, and

nutrient availability (Hu et al., 2017; Liu et al., 2020a; Zhang et al., 2019b), and biotic factors,

such as competition and predation. In addition to deterministic mechanisms, stochastic

processes, for instance, dispersal and ecological drift, also contribute to microbial community

assembly. Efforts have been made to disentangle the relative importance of deterministic vs.

stochastic processes based on a variety of null and neutral modeling methods (Ning et al.,

2019; Sloan et al., 2006; Stegen et al., 2013). Depending on which ecosystems or organism

types these methods are applied to, the relative importance of deterministic vs. stochastic

processes may greatly vary (Chen et al., 2019; Liu et al., 2018b; Wang et al., 2020; Yan et al.,

2020; Niederdorfer et al., 2021). For instance, bacterial communities in lake water were

revealed to be predominantly structured by determinism, whereas micro-eukaryotic

communities largely by stochasticity (Logares et al., 2018). For the MRP, the closed channel,

the environmental gradients and the variety of hydraulic constructions along the canal make it

obscure to what extent ecological stochasticity determines the assembly of microbiota

community, which is essential for an improved understanding of the ecosystem processes and

microbial dynamics prediction.

Other emerging questions in such large water-diversion canal and the comparable

systems, like river continuum (Gweon et al., 2021), include which immigrating

microorganisms are adapted to local niches in the receiving water and how they functionally

contribute to the ecological service. Especially in the MRP, how many of the microbial

populations originated from a natural reservoir may survive in the contrastingly engineered

canal remains an open question. Although population growth associated with microbial

immigration has been quantified in other engineered water systems (Mei and Liu, 2019), such

as activated sludge reactors (Mei et al., 2019), efforts in estimating microbial growth in

long-distance and evolving systems are still lacking.

In the present study, we followed the spatio-temporal dynamics of prokaryotic and

micro-eukaryotic community along the main canal of MRP of the SNWD project over a year

to address: 1) How does microbial community develop with the water flow? 2) How is the

relative importance of deterministic and stochastic processes on shaping the community

assembly in this closed turbulent engineered system? 3) How do the immigrating bacteria and

micro-eukaryotes from water source survive, grow, or die off in the canal? Based on

high-throughput marker gene (i.e., 16S and 18S) amplicon and metagenomic sequencing,

bioinformatics, and statistical modeling of water microbiota, we elucidated the microbial

diversity and population dynamics along the canal over four seasons and uncovered the key

patterns and processes driving community assembly. By overcoming the limitations of widely

used relative abundance measures with estimated absolute abundance (Barlow et al., 2020;

Tkacz et al., 2018), we demonstrated the first use of local growth factor to predict growth

capacity of the immigrating microbial populations along the water canal and revealed the

divergent microbial fates and population lifestyles throughout the world's largest water

diversion canal.

Materials and Methods

A full version of the Materials and methods are available in the Supplementary Information.

Study sites, sampling and physiochemical monitoring

The Middle Route Project (MRP) of the South-to-North Water Diversion (SNWD) Projects

started from Danjiangkou reservoir and diverted water northwards until Beijing and Tianjin,

which is an artificial system without direct connection with surface waters in the area. Fifty

samples were collected from 19 sites (Fig. 1), including one at the Danjiangkou reservoir

(S01) and 18 along the main canal of the MRP (S02-S18). The 18 sampling sites along the

canal were selected to capture the spatial dynamics of microbial communities from canal head

to end across the subtropical and warm-temperate climate zones. The sites were grouped to

three sections: S02-S08 (Section1), S09p-S14 (Section2) and S15-S18 (Section3). Canal

water was sampled and filtered through 0.22-μm cellulose nitrate filters (Sartorius, Germany).

Physiochemical properties, including water temperature (T), pH, dissolved oxygen (DO), total

nitrogen (TN) and fluoride (F⁻), were measured according to the environmental quality

standard for surface water of China (GB3838-2002). Details about the sampling sites and

physiochemical measurements are in the Supplementary Method S1.

DNA extraction, PCR amplification and sequencing

Genomic DNA was extracted from filters using FastDNA Spin Kit for Soil (MP Biomedicals,

USA). Bacterial 16S rRNA gene V3-V4 region and eukaryotic 18S rRNA gene V4 region

were amplified and sequenced on the Illumina platforms producing 250 bp paired-end reads at

the Guangdong Magigene Biotechnology Co., Ltd. (Guangzhou, China). Real-time

quantitative PCR (qPCR) was additionally performed to quantify the 16S rRNA gene copies

(as a proxy for bacterial biomass) in each genomic DNA sample, as described in Ju et al.

(2019). Detailed methodologies are provided in the Supplementary Method S2.

Each genomic DNA was also used for shot-gun metagenomic sequencing on the

Illumina's Novaseq platform using a 150 bp paired-end sequencing strategy at the Novogene

(Beijing, China). The sequencing data generated in this study have been deposited into CNGB

Sequence Archive (CNSA) (Guo et al., 2020) of China National GeneBank DataBase

(CNGBdb) (Chen et al., 2020) with accession number CNP0001593.

Bioinformatics and statistical analyses of microbial diversity

The 16S and 18S rRNA gene amplicon sequence data were processed using Qiime2 pipeline

(Bolyen et al., 2019) and DADA2 algorithm (Callahan et al., 2016) as described in the

Supplementary Method S3, generating final abundance tables of amplicon sequence variants

(ASVs) consisting of 6705 bacterial and 3217 eukaryotic ASVs, respectively. The

alpha-diversity metrics of bacterial and micro-eukaryotic communities were further computed

in Qiime2. All the univariate and multivariate statistics were performed in R (R Core Team,

2020). A set of spatial factors was generated for the sampling sites based on the geographic

coordinates using the principal coordinates of neighboring matrices (PCNM) analysis

(Borcard and Legendre, 2002).

The metagenomic data were pretreated and searched against the SILVA SSU132 database

to identify rRNA gene reads. This information was further used to estimate the 16S rRNA and

18S rRNA gene copies detected in each metagenome based on the sum for the read coverage

over the full length of each matched reference sequence in the database. Equation and details

are available in the Supplementary Method S3. The copies of 18S rRNA genes in each water

sample were estimated as 16S rRNA gene copies (determined by the aforementioned qPCR

analysis) divided by ratios of 16S rRNA to 18S rRNA gene copy (Table S1) as sequenced and

estimated in the metagenomic data.

Neutral and null model analyses of microbial community patterns

To disentangle the relative contribution of stochastic and deterministic processes to microbial

community assembly, Sloan neutral model (Sloan et al., 2006) and null model-based NST

approach (Ning et al., 2019) were applied to the 16S and 18S community data. Fitting of the

model were performed in R according to Burns et al. (2016). Accordingly, ASVs were

classified into three groups depending on whether their observed occurrence falling higher

than ("above prediction"), within ("as predicted") or lower than ("below prediction") the 95%

confidence interval of the neutral model predictions. Normalized stochasticity ratio (NST)

was used to quantify ecological stochasticity in communities within canal sections (Ning et al.,

2019). NST analysis was performed in R using the NST package (Ning et al., 2019). More

details about the methodology are in the Supplementary Method S4.

Population lifestyle and local growth factor analyses

The absolute abundance of bacterial and micro-eukaryotic ASVs was calculated by

multiplying relative abundance of each ASV computed from amplicon sequence data by the

overall 16S and 18S rRNA gene copies detected in a given metagenome for all the Oct

samples (see the Supplementary Method S5 for more details).

To characterize fate of microorganisms originated from natural reservoir after transported

into the closed canal, we defined a local growth factor (LGF) to depict whether the

populations grew or restrained, by comparing their mean absolute abundance in each canal

196   section to the abundance at the canal head, as follows:

$$\text{LGF}_{\text{X}} = \log_{10} \frac{\text{AA}_{\text{x,N}} + 1}{\text{AA}_{\text{x,OctS02}} + 1} \qquad (1)$$

where $\text{AA}_{\text{x, N}}$ is the average absolute abundance of ASV x in Section N of sampling sites and

$\text{AA}_{\text{x, OctS02}}$ is the absolute abundance of ASV x at OctS02, the canal head. OctS02 was

excluded from Section1 for the LGF calculation. An average LGF > 0 across the three

sections was considered to indicate growth in the canal. The potentially pathogenic bacterial

ASVs were identified referring to a list of potentially pathogenic genera (Table S2)

summarized from Woolhouse et al. (2015), and LGFs of these ASVs were examined.

Results

Microbiota biomass and diversity shifted with seasonality

Microbial communities showed clear seasonal shifts (Fig. 2), whether for their biomass (A

and B) or alpha (C-F) and beta diversity (Fig. 2G, 2H and Fig. S1). Both the bacterial and

micro-eukaryotic biomass, as measured by 16S rRNA (A) and 18S rRNA (B) gene copies,

were the highest in August and the lowest in October (Fig. 2). PERMANOVA showed that

seasonality significantly (P < 0.001) influenced both the bacterial (F = 22.78) and

micro-eukaryotic (F = 25.57) community composition. Moreover, the seasonal impacts were

much more pronounced than the spatial effects along the canal, whether for bacteria (R^2 =

0.52 vs. 0.14) or micro-eukaryotes (R^2 = 0.55 vs. 0.14). Accordingly, Bray Curtis-based

NMDS analyses of all canal water samples (n = 52) revealed strong clustering of bacterial

(Fig. 2G) and micro-eukaryotic (Fig. 2H) communities in relation to seasons, of which May

and August had witnessed greater intra-season community variations along the canal.

Seasonal variation was also observed for specific bacterial and micro-eukaryotic lineages (Fig.

S2 and S3).

Microbiota composition and diversity changed along the canal

Both the source water from Danjiangkou reservoir (S01) and the canal water were dominated

by phyla *Actinobacteria* (39.7 - 65.0%, mainly hgcI clade and CL500-29 marine group),

Proteobacteria (17.9 - 38.3%), *Bacteroidetes* (0.9 - 29.5%), *Verrucomicrobia* (1.2 - 9.2%) and

Cyanobacteria (0.2 - 8.6%, Fig. 3A). Overall, bacterial communities at the canal head (S02)

were most structurally similar to the source water (Fig. 3A and Fig. S1A). However,

compared with the source water, relative abundance of *Proteobacteria* at the canal head (i.e.,

S02) decreased whereas *Actinobacteria* and *Bacteroidetes* increased. Within the canal, both

Cyanobacteria (mostly *Cyanobium*_PCC-6307, $F = 16.77$, $P < 0.001$) and *Bacteroidetes* ($F =$

6.06, $P < 0.05$) showed significant decrease in relative abundance with water flow (Fig. 3A).

In contrast, *Limnohabitans* ($F = 19.87$) and *Candidatus* Methylopumilus ($F = 58.69$) within

Proteobacteria became increasingly abundant along the canal ($P < 0.001$). Likewise, both the

overall bacterial biomass (Fig. S4A) and Chao1 richness (Fig. S5A) exhibited highly dynamic

variation patterns along the canal. The variation trend of Shannon diversity index largely

agreed with (though relatively weaker than) that of Chao1 richness (Fig. S5B).

Unlike bacterial communities, micro-eukaryotic communities appeared more dynamic

along the canal water flow (Fig. 3B). The relative abundance of *Metazoa*, mostly *Calanoida*,

was as high as 24.0% in Danjiangkou Reservoir. These organisms were drastically enriched

by 2.3 times upon entering the canal head site, but largely declined in the downstream sites

(S03 to S18). Two predominant lineages of *Cryptophyceae*, i.e., *Cryptomonas* ($F = 4.11$, $P =$

0.06) and *Teleaulax ($F = 9.50$, $P < 0.01$)*, were more abundant in the source water and canal

upstream (S02-S08) than downstream sites (S09p-S14), accompanied by their gradual

replacement by two uncultured *Cryptophyceae* lineages ($F = 30.53$, $P < 0.001$; $F = 24.60$, $P <$

0.001, respectively), especially close to the canal end (S15-S18). The canal head communities

were distinct from water source and the downstream communities (Fig. 3B and Fig. S1B).

This special pattern was manifested by the dominance of *Calanoida* (72.0%) at the canal head

site where micro-eukaryotic biomass was much higher (3.2×10^9 vs. average 5.9×10^8 18S

rRNA gene copies L^{-1}, Fig. S4B) whereas alpha diversity metrics much lower compared with

the other sites (Fig. S6), suggesting intriguing local micro-eukaryotic selection in the canal

head.

Environmental factors influenced canal microbiota diversity and biomass

The influence of five environmental parameters, namely water temperature, pH, dissolved

oxygen (DO), total nitrogen (TN) and fluoride (Table S3), and spatial factors (i.e., ten PCNM

variables representing the quantitative spatial patterns, Table S4) on the microbial

communities was tested by redundancy analysis. All the above-mentioned environmental

variables and PCNM1, 2 and 4 were identified to exert significant impacts on bacterial

community composition ($P < 0.05$, Fig. 4A). These variables except for PCNM4 also

represented the significant influential factors for micro-eukaryotic communities ($P < 0.05$; Fig.

4B). Seasonal variation of environmental factors appeared based on the RDA plots (Fig. 4A

and 4B), for instance, fluoride concentration was especially higher in March, which was

supported by the significant seasonal difference in the environmental variables by

Kruskal-Wallis one-way ANOVA (Table S5). The abundant actinobacterial hgcI clade ($r_s =$

0.63, $P < 0.01$) and CL500-29 ($r_s = -0.68$, $P < 0.01$) marine group were among the bacterial

lineages positively and negatively related to fluoride concentration, respectively (Table S6).

For micro-eukaryotes, for instance, an ASV affiliated to *Mediophyceae* ($r_s = 0.66$, $P < 0.01$)

was positively correlated with fluoride (Table S7).

The identified significant variables were employed in further variation partitioning

analysis (VPA). The result showed clearly stronger influence of seasonality than that of

spatial and examined environmental factors for both bacterial and eukaryotic communities

(Figure 4C and 4D). Seasonality alone could explain 19% and 17% of the bacterial and

micro-eukaryotic community variance, respectively, while only 6 - 9% of the variance could

be explained exclusively by environmental or spatial factors. Notably high proportions of the

community variance could be attributed to the co-effects of seasonality and environmental

factors (29% for bacteria and 23% for micro-eukaryotes), implying their strong interplay in

shaping microbial communities. Besides, temperature, DO and TN were found to be

significantly correlated with bacterial biomass, while only DO for micro-eukaryotic biomass

(as measured by 16S rRNA and 18S rRNA gene copies, Table S8).

Bacterial and micro-eukaryotic communities exhibited distinct and spatio-temporally dynamic stochasticity

The apparent dispersal of microorganisms in the canal water and engineered hydraulic manipulations (i.e., inverted siphon, aqueduct, and tunnels) along the main canal drove us to question on the relative contribution of stochastic vs. deterministic processes to the canal bacterial and micro-eukaryotic community assembly, which was first quantitatively assessed with a recently proposed metric, normalized stochasticity ratio (NST) (Ning et al., 2019). This new metric quantifies the relative importance of stochasticity in governing community assembly ranging from 0 to 1, with a value > 0.5 indicating more stochastic assembly and < 0.5 more deterministic assembly. NST analysis revealed contrasting trends between bacteria and micro-eukaryotes. In the upstream section, a highest NST of 0.78 was estimated for bacteria (Fig. 5A). This value decreased to 0.46 in the middle section, and to as low as 0.32 close to the canal end. On the contrary, NST was generally above 0.5 for micro-eukaryotes throughout the canal, with the highest value of 0.75 in the middle section (Fig. 5A).

While NST provides rigorous quantification of relative importance of stochasticity, the Sloan neutral model analysis (Sloan et al., 2006) allows for assessing how well the observed community patterns could be explained by neutral processes (e.g., birth, death, and immigration) and attributing the observed community-level assembly pattern to the divergent population behavior. According to seasonal comparison of the model fitness, neutral model could explain much higher proportions of community patterns in October (R^2 = 0.73 and 0.67

for bacteria and micro-eukaryotes, respectively) and March (R^2 = 0.77 and 0.72), but lowest

in May (R^2 = 0.42 and 0.13, Fig. S7).

To link community patterns to the underlying population dynamics, bacterial and

micro-eukaryotic ASVs were classified into three groups based on the neutral model analysis

of the October samples. Depending on whether they empirically occurred more frequently

than, as frequently as or less than predicted by the neutral model, ASVs were assigned to

"above prediction", "as predicted" and "below prediction" groups that were different in

abundance and composition (Fig. 5B-D). We found that 7.8% of the bacterial community

richness (i.e., number of ASVs) accounted by above-prediction group corresponded to 67.9%

of the total number of sequences, which was dominated by *Actinobacteria* and *Proteobacteria*,

whereas the neutrally distributed group contributed 86.8% of the richness but only 23.9% of

the abundance (Fig. 5B and Fig. S8). Similar trend was detected for micro-eukaryotes (Fig.

5B). The three partitioning groups of ASVs also showed distinct taxonomic composition (Fig.

5C, 5D and Fig. S8). *Cryptomonas* constituted 13.7% of the above-prediction group in

abundance, but they were almost absent in the neutral and below-prediction groups. *Metazoa*,

in contrast, accounted for only 0.6% of the above-prediction group, but as high as 32.9% and

62.3% of the abundance of the neutral and below-prediction groups (Fig. 5D).

Selective growth of immigrating bacterial and micro-eukaryotic populations along the

canal

To resolve which populations of the water communities died off, sustained, or proliferated

during dispersal over 1432-km canal, we proposed local growth factor (LGF) (see equation 1

in Methods) as a quantitative measure of variation in population abundance in each canal

section compared to the abundance at the canal head, with an average LGF > 0 across the

three sections indicating growth in the canal. LGF analysis was applied with the Oct sample

dataset composed of 1389 bacterial and 1229 eukaryotic ASVs, using absolute abundance of

ASVs estimated by multiplying relative abundance of ASVs at each canal site with the total

copies of bacterial 16S rRNA (7.89×10^8 to 2.93×10^9 copies L^{-1}) or micro-eukaryotic 18S

rRNA (3.62×10^8 to 3.19×10^9 copies L^{-1}) genes therein. Micro-eukaryotes showed higher

overall occurrence (0.313±0.008 vs. 0.216±0.007; $P < 0.001$) while lower variability

(2.879±0.033 vs. 3.055±0.032; $P < 0.001$) in absolute abundance compared to bacteria.

In total, 67.1% of the bacterial ASVs (Table S9) and 82.3% of the micro-eukaryotic

ASVs (Table S10) displayed positive average LGFs. The ASVs were further observed to form

clearly three clusters based on their LGFs in each section: LGF > 3, -3 < LGF < 3 and LGF <

-3, which could indicate substantial growth, survival, and die-off of the microbial populations

(Fig. 6). The survival group represented the largest proportion of the ASVs (45.6 - 60.2%),

while only 4.1 - 21.7% of the ASVs went extinction in the canal. We also found that both the

contrastingly high and low LGFs tended to be attributed to high-abundance microbial lineages

(Fig. 6A and B). For example, the abundant *Luteolibacter* sp. (ASV106) and *Limnohabitans*

sp. (ASV27) exhibited the highest LGF values in Section 2 and Section 3, respectively (Table

S9). In stark contrast, ASV138 assigned to actinobacterial CL500-29 marine group and two

Cyanobium PCC-6307 lineages (ASV144 and ASV250) were especially restrained in growth

close to the canal end (Table S9). A total of 53 bacterial ASVs in the canal were related to 16

genera included in a published reference pathogen list (Table S2, summarized from

Woolhouse et al., 2015). While LGF of most (39 out of 53) of these potentially pathogenic

ASVs fell between -3 and 3, three *Acinetobacter*-associated ASVs (ASV794, ASV1108 and

ASV 1261) showed LGFs ranging from -5.9 to -6.1, indicating notable depletion along the

canal. For micro-eukaryotes, two uncultured *Cryptophyceae* lineages (ASV54 and ASV80) as

well as two ASVs related to *Acineta flava* (ASV224) and *Spumella* sp. (ASV59) were

elevated throughout the canal (Table S10).

Discussion

Diversion canals widely built for reallocation of water resources represent an ideal engineered

system to model microbial communities fulfilling key eco-environmental services, as well as

reflecting changes in ecosystem structure and function. Based on quantitative metagenomic

approaches that enable absolute quantification of microbial taxon, we uncovered the

microbiota spatio-temporal dynamics, key processes driving community assembly, and

navigation of immigrating microorganisms through the world's largest water diversion

project.

Microbiota composition and biomass are dynamic along the canal

Both bacterial and micro-eukaryotic communities in the source water showed marked

variations in taxonomic composition upon entering the canal, which could be attributed to the

readily comprehensible differences in the environmental conditions between the natural

reservoir and its receiving engineered canal and potential influences exerted by the approach

channel through which source water enters the canal. Meanwhile, we observed intriguingly

decreased water microbial diversity after water transportation through an inverted siphon (S09

vs. S09p, Fig. S5 and S6). Taken together, these suggest that hydraulic structures along the

canal apply considerable influence on the microbial community diversity and composition,

which needs to be fully considered when setting baseline communities for long-term

microbial monitoring in the water diversion canal.

The prevalence of uncultured actinobacterial hgcI clade and CL500-29 marine group

echoed what was previously reported in other freshwater habitats (Keshri et al., 2018; Yang et

al., 2017). While both groups of yet-to-be-cultured lineages have been shown to favor lower

level of dissolved organic carbon and temperature (Lindh et al., 2015), the hgcI clade was

additionally implicated to be competitively advantageous in utilizing low-concentration

organic carbon (Glöckner et al., 2000) and to harbor potentially photoheterotrophic species

showing preferential utilization of nitrogen-rich organic compounds in freshwater (Ghylin et

al., 2014), implying its potential role in the nutrient uptake and water auto-purification.

Unlike bacterial communities, micro-eukaryotic communities in the canal have not yet

been explored before. By virtue of the high-throughput molecular analyses, we detected not

only *Rotifea*, copepods and protozoa previously microscopically observed as abundant in the

Danjiangkou Reservoir (Li et al., 2012), but also thousands of micro-eukaryotic populations

(as ASVs), of which 6.7% are uncultured or unclassified even at the class level, implying

detection of numerous novel populations in the canal water. Notably, metazoan *Calanoida*

was repeatedly found to show drastic increase in abundance in the micro-eukaryotic

communities of canal head, compared with in the source water in both October and August.

This big order of aquatic crustaceans belonging to *Copepoda* inhabiting marine and

freshwater environments are found as the most successful among zooplankton colonizing

pelagic habitats (Blanco-Bercial et al., 2011; Bradford-Grieve, 2002). We further noted that

the SAR clade and *Cryptophyceae* showed substantially decreased relative abundance as a

result of high enrichment of *Calanoida* compared with the source water. However, their

absolute abundance or biomass, as we estimated as their 16S rRNA gene copies, remained

comparable (2.78 vs. 3.43×10^8 16S-copies L^{-1} for the SAR clade, and 3.68 vs. 4.25×10^8

16S-copies L^{-1} for *Cryptophyceae*). This strong decoupling pattern between relative and

absolute abundance cannot have been resolved by currently popular sequence-based relative

measures alone which have been shown to provide evidently misleading information on the

variation of absolute abundance (Stämmler et al., 2016; Vieira-Silva et al., 2019). It also well

demonstrates the power of our quantitative metagenomics approaches as an alternative to a

more direct but difficult-to-achieve quantitative PCR assay approach (due to the lack of 18S

rRNA gene universal primers) for estimating total micro-eukaryotic biomass.

Microbiota assembly is deterministically driven by seasonally dependent habitat filtering effects

Elucidating mechanisms and factors shaping biogeographic patterns has become a central goal in microbial ecology (Hanson et al., 2012). Environment exerts influence on community composition by selecting for populations that better adapted to the corresponding ecological niches. We found that temperature, pH, dissolved oxygen, total nitrogen and fluoride all showed significant effects on the community dynamics in the main canal. Fluoride affects bacterial growth and metabolic activity through targeting enzyme inhibition (Liao et al., 2015). Fluoride has been reported to alter microbial community in high-fluoride groundwater (Zhang et al., 2019a) and to exhibit inhibitory effect on bacterial populations involved in various wastewater treatment processes (Ochoa-Herrera et al., 2009). Our results showed that fluoride with concentration falling within the range of 0.14-0.22 mg L^{-1} (Table S3) could also affect microbial community structure in a drinking water diversion system, where fluoride could be mainly contributed by natural processes, such as mineral weathering (Nong et al., 2019), and atmospheric pollutants (Pu et al., 2017). Interestingly, we found positive response of the hgcI clade to elevated fluoride concentration (Table S6). While the mechanisms of their fluoride resistance still await exploration, such evidence is in line with their favorable distribution in a wide range of water ecosystems.

Although bacterial and micro-eukaryotic communities are both significantly influenced by the examined environmental variables and spatial segregation, their overall variation were

explained more by seasonality. Meanwhile, the strong co-influence of temporal and

environmental factors revealed by variation partitioning analysis could be at least partially

explained by seasonal variation of the examined variables (Table S5), suggesting

season-driven habitat filtering effects on microbiota. These results together imply a strong

interplay between temporal, spatial and environmental factors on influencing microbial

communities in the canal water, as previously noticed in natural lakes (Hu et al., 2017).

Although potential contribution from unmeasured environmental factors cannot be ruled out,

the remaining unexplained variation could possibly be generalized and linked to the influence

of stochastic processes.

Contrasting proportion of stochasticity contributes to bacterial and micro-eukaryotic

community assembly

Microbial community assembly is typically governed by four processes: selection, drift,

speciation and dispersal (Vellend and Agrawal, 2010). While selection is inherently

deterministic and drift is stochastic, speciation and dispersal could involve both deterministic

and stochastic mechanisms (Chase and Myers, 2011). For the MRP main canal, an

investigation period of one year is probably short for resolving speciation. Thus, selection

(deterministic processes), drift (stochastic processes) and dispersal (involving both) are

expected to shape the observed community assembly patterns, and their relative contributions

could be disentangled to help answer when, where, why and how much each process

contribute to the observed patterns (Hanson et al., 2012).

Based on NST estimation and neutral modeling, we found that relative importance of

stochastic and deterministic processes in microbial community assembly differ between

bacteria and micro-eukaryotes, and among seasons and the canal sections. Neutral processes

explained ~67% to 76% of the community patterns in October and March, but notably less in

May (~41% and 13% for bacteria and eukaryotes, respectively). Liu et al. (2018a) reported a

prominent shortage in water storage of Danjiangkou Reservoir related to seasonal flood

control in May, which could have possibly imposed higher environmental selection effect to

the microbiota in the reservoir and across the canal, making the observed decreased

community stochasticity (or increased determinism) plausible. Although ecological

stochasticity adds to challenge to predict microbial communities, the new knowledge

produced in this study undoubtedly contribute to an improved understanding on when the

canal water microbiota is more prone to stochastic processes, or under which conditions

stochasticity is of more relevance.

The NST values suggested that the relative contribution of deterministic processes is

larger to bacterial community assembly compared with micro-eukaryotic communities, where

stochasticity prevails. Recently, Gweon et al. (2021) reported that stochastic processes played

a more important role in the planktonic bacterial communities in the River Thames,

presumably due to the influence of various natural and anthropogenic sources of the surface

water. What we found showed that in a water ecosystem almost isolated from other sources

than resource water, bacterial communities are predominantly shaped by deterministic abiotic

and biotic processes. The micro-eukaryotes, however, possess structurally and functionally

more complex cells than prokaryotes (Massana and Logares, 2013), which could facilitate

niche adaptation and diminish the environmental filtering effect at the community level. This

was supported by our observation that the micro-eukaryotes were distributed with higher

occurrence frequency and lower variability across the canal compared to bacteria. The

different controlling mechanisms between bacterial and micro-eukaryotic community

assembly found here has been observed in other natural habitats, such as Antarctic (Logares et

al., 2018) and Tibetan lakes (Liu et al., 2020b). Our comparative analyses between bacteria

and micro-eukaryotes across canal sections and over seasons thus provide first informative

insights into the domain-specific, and space- and season-dependent patterns in the canal water

community assembly.

Divergent population lifestyles underly microbiota dynamics revealed by absolute

quantification

A synthesis of population and community ecological perspectives can provide valuable

insights into how microorganisms drive ecosystem processes (Koch et al., 2018). Microbial

populations in this giant system displayed interestingly divergent lifestyles reflecting the

difference in their adaptability to the canal environment. The Sloan neutral model sorted the

microbial ASVs to above-prediction, below-prediction (jointly non-neutral) and as-predicted

(neutral) groups, which were different in community composition (Fig. 5C and 5D),

indicating taxonomically dependent competitive fitness in the investigated canal. In addition,

we observed contrasting low (~20%) contribution of neutral group to community abundance

despite their high proportion in richness (~80%) for both bacteria and micro-eukaryotes (Fig.

5B). Chen et al. (2019) reported for micro-eukaryotic communities in a subtropical river as

source of drinking water the dominance of neutral groups in both richness and abundance of

the overall communities as well as across taxonomic groups. These contrasting patterns

suggested that microorganisms are prone to more selective processes in an engineered canal

compared with natural habitats.

In particular, we demonstrate the use of local growth factor (LGF) as a meaningful

metric to infer growth of immigrating microbial taxa in the canal by employing absolute

quantification of microbial populations. This leads to the first estimation that 67.1% of

bacterial and 82.3% of micro-eukaryotic populations were capable of sustaining growth

(LGF > 0), implying again better acclimation of eukaryotes to the environmental conditions in

the canal. It is worth clarifying that the microbial growth estimated here was aimed to

characterize fate of immigrating microorganisms across the canal, calculated based on relative

fluctuation in absolute abundance at local sites compared with canal head, and thus is not

equivalent to per-capita or population growth in the Sloan neutral model (Sloan et al., 2006).

Limnohabitans sp., ubiquitous in a wide variety of freshwater habitats, showed remarkable

growth across the canal. Members of *Limnohabitans* typically display high growth rate as

well as metabolic flexibility, and they actively utilize alga-derived dissolved organic

substances and are vulnerable to protist grazing (Horňák et al., 2017; Šimek et al., 2013;

Šimek et al., 2011). *Cryptophyceae* were among the micro-eukaryotes that displayed

prominent growth throughout the canal, which are small flagellates as important food of many

protists and micro-crustaceans (Horňák et al., 2017; Posch et al., 2015), and are perceived as

characteristic of clean water (Lackey, 1941). Thus, *Limnohabitans* and *Cryptophyceae* are

potentially important constituents of water food webs in the investigated canal. Two

cyanobacterial lineages (*Cyanobum* PCC-6307) and *Acinetobacter* spp., on the contrary, were

clearly depressed in the canal downstream sections, demonstrating the elimination of these

microorganisms of water security concern along the canal. *Acinetobacter* spp. are ubiquitous

in nature of major ecological and clinical significance, featuring, for instance,

multi-antimicrobial resistance and rapid resistance development (Jung and Park, 2015). The

diminishing of the versatile *Acinetobacter* spp. along the canal could possibly be attributed to

unfavorable environmental conditions (Al Atrouni et al., 2016) or composite intra- and

inter-domain biological interactions. Further research efforts with genome-centric

metagenomic approaches are needed to elaborate the mechanisms behind such die-off patterns.

It is noteworthy that due to the restriction in taxonomic resolution of the ribosomal RNA gene

sequencing analysis employed in this study, species-level confirmation is still needed to

evaluate the relevance of these potentially pathogenic microorganisms. Together, our results

demonstrated the power of local growth factor to identify key players in the investigated

system and facilitate predicting fate of the microbes traversing such thousand-kilometer water

diversion canal.

Conclusions

The main canal of the Middle Route Project of the South-to-North Water Diversion projects is

an engineered system where the aquatic ecosystem is still at its early forming stage (since

2014). Various aspects of hydrology, geochemistry and ecology need to be intricately

elaborated to jointly contribute to sustained system's eco-environmental services, especially

water auto-purification ensuring its quality and competence for designed social-economic

aims. The present study uncovered the underlying microbial community assembly

mechanisms and population dynamics in a giant engineered water diversion water canal, such

as the sharp decline of potentially harm-producing microbes (e.g., *Cyanobacteria*) and

potential pathogens (e.g., *Acinetobacter* spp.), and the favorable enrichment of alga-derived

organic scavengers (e.g., *Limnohabitans*) as potential key microbial food-web players,

demonstrating the valuable insights provided by microbial community and population

assessments for water monitoring and their potential in indicating ecosystem disturbance.

Furthermore, the bacterial community shift driven by combined environmental effects as well

as the stochasticity in micro-eukaryotic community assembly needs to be considered in water

quality monitoring along the canal, towards an appropriate microbe-based biomonitoring

which enables deeper understanding of processes behind water quality parameters.

Considering possibly occasional pollution into the canal water caused by, for example, traffic

accidents on highway bridges across the canal, and the central roles microorganisms play in

the water quality maintenance, future efforts are needed to explore the metabolic functions of

microbial community contributing to water purification (e.g., carbon, nitrogen, and

phosphorus removal) during diversion and sustaining water quality.

Acknowledgments

We would like to thank Yuanzhen He, Guoqing Zhang, Zhipeng Zhang, Shengguang Yuan, Aoxiang Fan and Qinglin Jia for the assistance in water sampling. This work was supported by National Natural Science Foundation of China (Grant No. 51908467 and No. U2040210), and Zhejiang Provincial Natural Science Foundation of China (Grant No. LQ20C030002).

References

Al Atrouni, A., Joly-Guillou, M.-L., Hamze, M. and Kempf, M. 2016. Reservoirs of non-baumannii
Acinetobacter species. Frontiers in Microbiology 7, 49.

Arrigo, K.R. 2005. Marine microorganisms and global nutrient cycles. Nature 437(7057), 349-355.

Barlow, J.T., Bogatyrev, S.R. and Ismagilov, R.F. 2020. A quantitative sequencing framework for
absolute abundance measurements of mucosal and lumenal microbial communities. Nature
Communications 11(1), 2590.

Battin, T.J., Besemer, K., Bengtsson, M.M., Romani, A.M. and Packmann, A.I. 2016. The ecology
and biogeochemistry of stream biofilms. Nature Reviews Microbiology 14(4), 251-263.

Bell, T., Newman, J.A., Silverman, B.W., Turner, S.L. and Lilley, A.K. 2005. The contribution of
species richness and composition to bacterial services. Nature 436(7054), 1157-1160.

Berry, D., Xi, C. and Raskin, L. 2006. Microbial ecology of drinking water distribution systems.
Current Opinion in Biotechnology 17(3), 297-302.

Blanco-Bercial, L., Bradford-Grieve, J. and Bucklin, A. 2011. Molecular phylogeny of the Calanoida
(Crustacea: Copepoda). Molecular Phylogenetics and Evolution 59(1), 103-113.

Bolyen, E., Rideout, J.R., Dillon, M.R., Bokulich, N.A., Abnet, C.C., Al-Ghalith, et al. 2019.
Reproducible, interactive, scalable and extensible microbiome data science using QIIME 2.
Nature Biotechnology 37, 852-857.

Borcard, D. and Legendre, P. 2002. All-scale spatial analysis of ecological data by means of principal
coordinates of neighbour matrices. Ecological Modelling 153(1), 51-68.

Bradford-Grieve, J.M. 2002. Colonization of the pelagic realm by calanoid copepods. Hydrobiologia
485(1), 223-244.

Briones, A. and Raskin, L. 2003. Diversity and dynamics of microbial communities in engineered
environments and their implications for process stability. Current Opinion in Biotechnology
14(3), 270-276.

Bru, D., Ramette, A., Saby, N.P.A., Dequiedt, S., Ranjard, L., Jolivet, C., Arrouays, D. and Philippot, L.
2011. Determinants of the distribution of nitrogen-cycling microbial communities at the
landscape scale. The ISME Journal 5(3), 532-542.

Burns, A.R., Stephens, W.Z., Stagaman, K., Wong, S., Rawls, J.F., Guillemin, K. and Bohannan, B.J.M.
2016. Contribution of neutral processes to the assembly of gut microbial communities in
the zebrafish over host development. The ISME Journal 10, 655.

Callahan, B.J., McMurdie, P.J., Rosen, M.J., Han, A.W., Johnson, A.J.A. and Holmes, S.P. 2016.
DADA2: High-resolution sample inference from Illumina amplicon data. Nature Methods
13(7), 581-583.

Chase, J.M. and Myers, J.A. 2011. Disentangling the importance of ecological niches from
stochastic processes across scales. Philosophical Transactions of the Royal Society B:
Biological Sciences 366(1576), 2351-2363.

Chen, F.Z., You, L.J., Yang, F., Wang, L.N., Guo, X.Q., Gao, F., Hua, C., Tan, C., Fang, L., Shan, R.Q., Zeng,
W.J., Wang, B., Wang, R., Xu, X. and Wei, X.F. 2020. CNGBdb: China National GeneBank
DataBase. Yi Chuan 42(8), 799-809.

Chen, W., Ren, K., Isabwe, A., Chen, H., Liu, M. and Yang, J. 2019. Stochastic processes shape
microeukaryotic community assembly in a subtropical river across wet and dry seasons.
Microbiome 7(1), 138.

Chen, Z., Yuan, J., Sun, F., Zhang, F., Chen, Y., Ding, C., Shi, J., Li, Y. and Yao, L. 2018. Planktonic
fungal community structures and their relationship to water quality in the Danjiangkou
Reservoir, China. Scientific Reports 8(1), 10596.

Dang, C., Liu, W., Lin, Y., Zheng, M., Jiang, H., Chen, Q. and Ni, J. 2018. Dominant role of
ammonia-oxidizing bacteria in nitrification due to ammonia accumulation in sediments of
Danjiangkou reservoir, China. Applied Microbiology and Biotechnology 102(7), 3399-3410.

Du, M., Peng, X., Zhang, H., Ye, C., Dasgupta, S., Li, J., Li, J., Liu, S., Xu, H., Chen, C., Jing, H., Xu, H., Liu,
J., He, S., He, L, Cai, S., Chen, S. and Ni, J. 2021. Geology, environment, and life in the
deepest part of the world's oceans. The Innovation 2(2), 100109.

Ghylin, T.W., Garcia, S.L., Moya, F., Oyserman, B.O., Schwientek, P., Forest, K.T., et al. 2014.
Comparative single-cell genomics reveals potential ecological niches for the freshwater acI
Actinobacteria lineage. The ISME Journal 8, 2503.

Glöckner, F.O., Zaichikov, E., Belkova, N., Denissova, L., Pernthaler, J., Pernthaler, A. and Amann, R.
2000. Comparative 16S rRNA analysis of lake bacterioplankton reveals globally distributed
phylogenetic clusters including an abundant group of *Actinobacteria*. Applied and
Environmental Microbiology 66(11), 5053-5065.

Guo, X., Chen, F., Gao, F., Li, L., Liu, K., You, L., Hua, C., Yang, F., Liu, W., Peng, C., Wang, L., Yang, X.,
Zhou, F., Tong, J., Cai, J., Li, Z., Wan, B., Zhang, L., Yang, T., Zhang, M., Yang, L., Yang, Y., Zeng,
W., Wang, B., Wei, X. and Xu, X. 2020. CNSA: a data repository for archiving omics data.
Database (Oxford) 2020.

Gweon, H.S., Bowes, M.J., Moorhouse, H.L., Oliver, A.E., Bailey, M.J., Acreman, M.C. and Read, D.S.
2021. Contrasting community assembly processes structure lotic bacteria

metacommunities along the river continuum. Environmental Microbiology 23(1), 484-498.

Hanson, C.A., Fuhrman, J.A., Horner-Devine, M.C. and Martiny, J.B.H. 2012. Beyond biogeographic
patterns: processes shaping the microbial landscape. Nature Reviews Microbiology 10(7),
497-506.

Horňák, K., Kasalický, V., Šimek, K. and Grossart, H.P. 2017. Strain-specific consumption and
transformation of alga-derived dissolved organic matter by members of the Limnohabitans-C
and Polynucleobacter-B clusters of Betaproteobacteria. Environmental Microbiology 19(11),
4519-4535.

Hu, A., Ju, F., Hou, L., Li, J., Yang, X., Wang, H., Mulla, S.I., Sun, Q., Bürgmann, H. and Yu, C.-P. 2017.
Strong impact of anthropogenic contamination on the co-occurrence patterns of a riverine
microbial community. Environmental Microbiology 19(12), 4993-5009.

Hull, N.M., Ling, F., Pinto, A.J., Albertsen, M., Jang, H.G., Hong, P.-Y., Konstantinidis, K.T., LeChevallier,
M., Colwell, R.R. and Liu, W.-T. 2019. Drinking water microbiome project: Is it time?
Trends in Microbiology 27(8), 670-677.

Ju, F., Beck, K., Yin, X., Maccagnan, A., McArdell, C.S., Singer, H.P., Johnson, D.R., Zhang, T. and
Bürgmann, H. 2019. Wastewater treatment plant resistomes are shaped by bacterial
composition, genetic exchange, and upregulated expression in the effluent microbiomes. The
ISME Journal 13(2), 346-360.

Ju, F., Lau, F. and Zhang, T. 2017. Linking microbial community, environmental variables, and
methanogenesis in anaerobic biogas digesters of chemically enhanced primary treatment
sludge. Environmental Science and Technology 51(7), 3982-3992.

Ju, F., Xia, Y., Guo, F., Wang, Z. and Zhang, T. 2014. Taxonomic relatedness shapes bacterial
assembly in activated sludge of globally distributed wastewater treatment plants.
Environmental Microbiology 16(8), 2421-2432.

Ju, F. and Zhang, T. 2015. Bacterial assembly and temporal dynamics in activated sludge of a
full-scale municipal wastewater treatment plant. The ISME journal 9(3), 683.

Jung, J. and Park, W. 2015. *Acinetobacter* species as model microorganisms in environmental
microbiology: current state and perspectives. Applied Microbiology and Biotechnology 99,
2533-2548.

Keshri, J., Pradeep Ram, A.S., Nana, P.A. and Sime-Ngando, T. 2018. Taxonomical resolution and
distribution of bacterioplankton along the vertical gradient reveals pronounced
spatiotemporal patterns in contrasted temperate freshwater lakes. Microbial Ecology 76(2),
372-386.

Koch, B.J., McHugh, T.A., Hayer, M., Schwartz, E., Blazewicz, S.J., Dijkstra, P., van Gestel, N., Marks, J.C.,
Mau, R.L., Morrissey, E.M., Pett-Ridge, J. and Hungate, B.A.   2018.   Estimating
taxon-specific population dynamics in diverse microbial communities. Ecosphere 9(1),
e02090.

Lackey, J.B.   1941.   Two groups of flagellated algae serving as indicators of clean water. Journal
(American Water Works Association) 33(6), 1099-1110.

Li, Y.-Y., Zhang, Z., Yin, W., Hu, L., Liang, Z., Wang, F. and Lei, G.   2012.   Zooplankton diversity in the
water resource area of the Mid-Line Project of South to North Water Division. Advanced
Materials Research 356-360, 199-203.

Liao, Y., Chen, J., Brandt, B.W., Zhu, Y., Li, J., van Loveren, C. and Deng, D.M.   2015.   Identification
and functional analysis of genome mutations in a fluoride-resistant *Streptococcus mutans*
strain. PLOS ONE 10(4), e0122630.

Lindh, M.V., Lefébure, R., Degerman, R., Lundin, D., Andersson, A. and Pinhassi, J.   2015.
Consequences of increased terrestrial dissolved organic matter and temperature on
bacterioplankton community composition during a Baltic Sea mesocosm experiment. AMBIO
44(3), 402-412.

Liu, H., Yin, J. and Feng, L.   2018a.   The dynamic changes in the storage of the Danjiangkou
Reservoir and the influence of the South-North Water Transfer Project. Scientific Reports 8(1),
8710.

Liu, J., Zhu, S., Liu, X., Yao, P., Ge, T. and Zhang, X.-H.   2020a.   Spatiotemporal dynamics of the
archaeal community in coastal sediments: assembly process and co-occurrence relationship.
The ISME Journal 14(6):1463-1478.

Liu, K., Liu, Y., Hu, A., Wang, F., Chen, Y., Gu, Z., Anslan, S. and Hou, J.   2020b.   Different community
assembly mechanisms underlie similar biogeography of bacteria and microeukaryotes in
Tibetan lakes. FEMS Microbiology Ecology 96(6).

Liu, T., Zhang, A.N., Wang, J., Liu, S., Jiang, X., Dang, C., Ma, T., Liu, S., Chen, Q., Xie, S., Zhang, T. and Ni,
J.   2018b.   Integrated biogeography of planktonic and sedimentary bacterial communities
in the Yangtze River. Microbiome 6(1), 16.

Logares, R., Tesson, S.V.M., Canbäck, B., Pontarp, M., Hedlund, K. and Rengefors, K.   2018.
Contrasting prevalence of selection and drift in the community structuring of bacteria and
microbial eukaryotes. Environmental Microbiology 20(6), 2231-2240.

Luo, Z., Li, S., Hou, K. and Ji, G.   2019.   Spatial and seasonal bacterioplankton community dynamics
in the main channel of the Middle Route of South-to-North Water Diversion Project. Research
in Microbiology 170(1), 24-34.

Massana, R. and Logares, R.　2013.　Eukaryotic versus prokaryotic marine picoplankton ecology.
675   Environmental Microbiology 15(5), 1254-1261.

676   Mei, R., Kim, J., Wilson, F.P., Bocher, B.T.W. and Liu, W.-T.　2019.　Coupling growth kinetics modeling
677   with machine learning reveals microbial immigration impacts and identifies key
678   environmental parameters in a biological wastewater treatment process. Microbiome 7(1),
679   65.

680   Mei, R. and Liu, W.-T.　2019.　Quantifying the contribution of microbial immigration in engineered
681   water systems. Microbiome 7(1), 144.

682   Ning, D., Deng, Y., Tiedje, J.M. and Zhou, J.　2019.　A general framework for quantitatively assessing
683   ecological stochasticity. Proceedings of the National Academy of Sciences 116(34),
684   16892-16898.

685   Niderdorfer, R., Fragner, L., Yuan, L., Hausherr, D., Wei, J., Magyar, P., Joss, A., Lehmann, M.F., Ju, F. and
686   Bürgmann, H.　2021.　Distinct growth stages controlled by the interplay of deterministic
687   and stochastic processes in functional anammox biofilms. Water Research 200, 117225.

688   Nong, X., Shao, D., Xiao, Y. and Zhong, H.　2019.　Spatio-temporal characterization analysis and
689   water quality assessment of the South-to-North Water Diversion Project of China.
690   International Journal Environmental Research and Public Health 16(12), 2227.

691   Ochoa-Herrera, V., Banihani, Q., León, G., Khatri, C., Field, J.A. and Sierra-Alvarez, R.　2009.　Toxicity
692   of fluoride to microorganisms in biological wastewater treatment systems. Water Research
693   43(13), 3177-3186.

694   Posch, T., Eugster, B., Pomati, F., Pernthaler, J., Pitsch, G. and Eckert, E.　2015.　Network of
695   interactions between ciliates and phytoplankton during spring. Frontiers in Microbiology 6,
696   14.

697   Pu, W., Quan, W., Ma, Z., Shi, X., Zhao, X., Zhang, L., Wang, Z. and Wang, W.　2017.　Long-term trend of
698   chemical composition of atmospheric precipitation at a regional background station in Northern
699   China. Science of The Total Environment 580,1340-1350.

700   R Core Team 2020　R: A language and environment for statistical computing, R foundation for
701   statistical computing.

702   Sagova-Mareckova, M., Boenigk, J., Bouchez, A., Cermakova, K., Chonova, T., Cordier, et al.　2021.
703   Expanding ecological assessment by integrating microorganisms into routine freshwater
704   biomonitoring. Water Research 191, 116767.

Šimek, K., Kasalický, V., Jezbera, J., Horňák, K., Nedoma, J., Hahn, M.W., Bass, D., Jost, S. and Boenigk, J.

2013.   Differential freshwater flagellate community response to bacterial food quality with a
focus on *Limnohabitans* bacteria. The ISME Journal 7(8), 1519-1530.

Šimek, K., Kasalický, V., Zapomělová, E. and Horňák, K.   2011.   Alga-derived substrates select for
distinct betaproteobacterial lineages and contribute to niche separation in *Limnohabitans*
strains. Applied and Environmental Microbiology 77(20), 7307-7315.

Sloan, W.T., Lunn, M., Woodcock, S., Head, I.M., Nee, S. and Curtis, T.P.   2006.   Quantifying the roles
of immigration and chance in shaping prokaryote community structure. Environmental
Microbiology 8(4), 732-740.

Stämmler, F., Gläsner, J., Hiergeist, A., Holler, E., Weber, D., Oefner, P.J., Gessner, A. and Spang, R.
2016.   Adjusting microbiome profiles for differences in microbial load by spike-in bacteria.
Microbiome 4(1), 28.

Stegen, J.C., Lin, X., Fredrickson, J.K., Chen, X., Kennedy, D.W., Murray, C.J., Rockhold, M.L. and
Konopka, A.   2013.   Quantifying community assembly processes and identifying features
that impose them. The ISME Journal 7, 2069.

Tkacz, A., Hortala, M. and Poole, P.S.   2018.   Absolute quantitation of microbiota abundance in
environmental samples. Microbiome 6(1), 110.

Vanwonterghem, I., Jensen, P.D., Dennis, P.G., Hugenholtz, P., Rabaey, K. and Tyson, G.W.   2014.
Deterministic processes guide long-term synchronised population dynamics in replicate
anaerobic digesters. The ISME Journal 8(10), 2015-2028.

Vellend, M. and Agrawal, A.   2010.   Conceptual synthesis in community ecology. The Quarterly
Review of Biology 85(2), 183-206.

Vieira-Silva, S., Sabino, J., Valles-Colomer, M., Falony, G., Kathagen, G., Caenepeel, C., Cleynen, I., van
der Merwe, S., Vermeire, S. and Raes, J.   2019.   Quantitative microbiome profiling
disentangles inflammation- and bile duct obstruction-associated microbiota alterations
across PSC/IBD diagnoses. Nature Microbiology 4(11), 1826-1831.

Wang, W., Ren, K., Chen, H., Gao, X., Rønn, R. and Yang, J.   2020.   Seven-year dynamics of testate
amoeba communities driven more by stochastic than deterministic processes in two
subtropical reservoirs. Water Research 185, 116232.

Woolhouse, M. E. J., Gowtage-Sequeria, S. and Evans, B.   2015.   Quantitative analysis of the
characteristics   of   emerging   and   re-emerging   human   pathogens.
http://webarchive.nationalarchives.gov.uk/20121206154522/http://www.bis.gov.uk/assets/f
oresight/docs/infectious-diseases/t16.pdf 2015.

Wu, L., Ning, D., Zhang, B., Li, Y., Zhang, P., Shan, et al.   2019.   Global diversity and biogeography of
bacterial communities in wastewater treatment plants. Nature Microbiology 4, 1183-1195.

Yan, L., Herrmann, M., Kampe, B., Lehmann, R., Totsche, K.U. and Küsel, K.   2020.   Environmental
selection shapes the formation of near-surface groundwater microbiomes. Water Research
170, 115341.

Yang, C., Wang, Q., Simon, P.N., Liu, J., Liu, L., Dai, X., Zhang, X., Kuang, J., Igarashi, Y., Pan, X. and Luo, F.
2017.   Distinct network interactions in particle-associated and free-living bacterial
communities during a *Microcystis aeruginosa* bloom in a plateau lake. Frontiers in
microbiology 8, 1202-1202.

Zhang, X., Gao, X., Li, C., Luo, X. and Wang, Y.   2019a.   Fluoride contributes to the shaping of
microbial community in high fluoride groundwater in Qiji County, Yuncheng City, China.
Scientific Reports 9, 14488.

Zhang, X., Li, B., Xu, H., Wells, M., Tefsen, B. and Qin, B.   2019b.   Effect of micronutrients on algae in
different regions of Taihu, a large, spatially diverse, hypereutrophic lake. Water Research 151,
500-514.

Zhou, J., Deng, Y., Zhang, P., Xue, K., Liang, Y., Van Nostrand, J.D., Yang, Y., He, Z., Wu, L., Stahl, D.A.,
Hazen, T.C., Tiedje, J.M. and Arkin, A.P.   2014.   Stochasticity, succession, and environmental
perturbations in a fluidic ecosystem. Proceedings of the National Academy of Sciences 111(9),
E836-E845.

 Figure Legends

 Fig. 1 Four sampling campaigns at 19 sites along the 1432-km main canal of the MRP of the

 SNWD project selected for microbial and physicochemical analyses. Colored fans indicate

 months of sampling at each site. The map was created using QGIS (version 3.10.5,

 www.qgis.org).

 Fig. 2 Seasonal variation of bacterial and micro-eukaryotic communities in the canal. (A-F)

 Ribosomal RNA gene copies (A and B), Chao1 richness (C and D) and Shannon diversity

 index (E and F) across seasons. Different letters above boxes indicate statistically significant

 differences between months of sampling (Mann-Whitney *U* test with the Benjamini-Hochberg

 correction, $P < 0.05$). (G-H) Ordination analysis of communities using non-metric

 multidimensional scaling (NMDS) based on Bray-Curtis dissimilarities. Trajectories indicate

 locations of sampling sites along the canal.

 Fig. 3 Microbial community composition and diversity along the MRP main canal (~1432 km)

 of the SNWD project in October 2018. (A) Bacterial composition with abundant phyla (> 1%)

 and 10 most abundant genera shown. (B) Micro-eukaryotic composition with abundant phyla

 (> 1%) and 10 most abundant orders shown.

 Fig. 4 Redundancy analysis (RDA) and variation partitioning analysis (VPA) demonstrating

 the relative influence of seasonality, spatial and environmental factors on the canal microbial

 communities. The RDA biplots show factors impacting bacterial (A) and micro-eukaryotic (B)

 communities. The VPA diagrams show the percentage contribution of seasonal, spatial and

environmental factors to bacterial (C) and micro-eukaryotic (D) community variations. Only

those factors identified to significantly influence bacterial and micro-eukaryotic communities

are shown in the RDA and included in the VPA analyses. PCNMs, geographic factors

generated by principal coordinate analysis of neighborhood matrices. T, temperature; F,

fluoride; TN, total nitrogen; DO, dissolved oxygen.

Fig. 5 Normalized stochasticity ratio (NST) estimation and microbiota partitioning by

population compliance to the neutral model prediction. (A) NST estimated stochasticity in

bacterial and micro-eukaryotic community assembly. (B) Proportions in richness (ASV

number) and abundance (sequence number) of the three fitting groups (occurrence frequency

above prediction, as predicted and below prediction) within the microbial communities to the

Sloan neutral model. (C and D) Taxonomic composition of the three fitting partitions in

richness (Ri) and abundance (Ab) for bacterial (C) and micro-eukaryotic (D) communities.

Only abundant bacterial phyla (> 1%) and 10 most abundant micro-eukaryotic orders are

shown.

Fig. 6 Microbial population proliferation across canal sections evaluated by local growth

factor analysis. Distribution patterns of population local growth factors in the different canal

sections are shown with violin plots for bacteria (A) and micro-eukaryotes (B). Each dot

represents an ASV with colors indicating its average abundance across the canal.

(A)

(B)

(A)

(B) Bacteria   Micro-eukaryotes

(C) Proportion (%)

(D) Proportion (%)

(A)

Section3

Section2

Section1

Log$_{10}$(average16S copies)

Local growth factor

(B)

Section3

Section2

Section1

Log$_{10}$(average18S copies)

Local growth factor