Title: Cell-specific expression buffering promotes cell survival and cancer robustness

Hao-Kuen Lin[a,b,1], Jen-Hao Cheng[a,c,1], Chia-Chou Wu[a,1], Feng-Shu Hsieh[a], Carolyn Dunlap[a],

and Sheng-hong Chen[a,c*]

[a] Lab for Cell Dynamics, Institute of Molecular Biology, Academia Sinica, Taipei 115,

Taiwan

[b] College of Medicine, National Taiwan University, Taipei 106, Taiwan

[c] Genome and Systems Biology Degree Program, Academia Sinica and National Taiwan

University, Taipei, Taiwan

[1] equal contribution

[*] Correspondence: shengchen@gate.sinica.edu.tw

Summary blurb

This study explores a genome-wide functional buffering mechanism, Cell-specific

Expression Buffering (CEBU), where gene expression contributes to functional buffering in

specific cell types and tissues. CEBU is prevalent in humans, linking to genetic interactions,

tissue homeostasis and cancer robustness.

Abstract

Functional buffering ensures biological robustness critical for cell survival and physiological

homeostasis in response to environmental challenges. However, in multicellular organisms,

the mechanism underlying cell- and tissue-specific buffering and its implications for cancer

development remain elusive. Here, we propose a Cell-specific Expression-BUffering (CEBU)

mechanism, whereby a gene's function is buffered by cell-specific expression of a buffering

gene, to describe functional buffering in humans. The likelihood of CEBU between gene

pairs is quantified using a C-score index. By computing C-scores using genome-wide

CRISPR screens and transcriptomic RNA-seq of 684 human cell lines, we report that C-

score-identified putative buffering gene pairs are enriched for members of the same pathway,

protein complex and duplicated gene family. Furthermore, these buffering gene pairs

contribute to cell-specific genetic interactions and are indicative of tissue-specific robustness.

C-score derived buffering capacities can help predict patient survival in multiple cancers. Our

results reveal CEBU as a critical mechanism of functional buffering contributing to cell

survival and cancer robustness in humans.

Running title: Expression buffering for cancer robustness

Keywords: functional buffering, expression buffering, buffering capacity, genetic interaction,

tissue homeostasis, cancer robustness

Introduction

Robustness in biological systems is critical for organisms to carry out vital functions in the face of environmental challenges [1, 2]. A fundamental requirement for achieving biological robustness is functional buffering, whereby the biological functions performed by one gene can also be attained via other buffering genes. These buffering genes thereby enhance the robustness of the biological system. Multiple mechanisms underlie this functional buffering, either via the responsive or intrinsic activities of buffering genes. Responsive buffering mechanisms activate buffering genes only if the function of a buffered gene is compromised. Therefore, the mechanism incorporates a control system that senses a function has been compromised and then activates the buffering genes, usually by up-regulating their gene expression [3]. One classical responsive buffering mechanism is genetic compensation among duplicated genes, whereby functional compensation is achieved by activating expression of paralogs when the functions of the original duplicated genes is compromised [4]. Genetic analyses of duplicated genes in *Saccharomyces cerevisiae* have revealed up-regulated gene expression in ~10% of paralogs when cell growth is compromised due to deletions of their duplicated genes [5-7]. Apart from duplicated genes, non-orthologous/analogous genes can also be activated for functional buffering [8]. For example, multiple growth signaling pathways coordinate cellular growth in parallel, and inactivation of one pathway can lead to activation of others for better survival [3, 4]. In cases of both duplicated and analogous genes, expression of their buffering genes is not activated unless their buffered functions are compromised [3, 7], so responsiveness follows a "need-based" principal. This responsive buffering mechanism has been observed for stress responses of both unicellular and multicellular organisms [3].

Although numerous studies over recent decades have focused on investigating such

responsive buffering mechanisms, a few recent studies have revealed another less-

characterized type of buffering mechanism in multicellular organisms, i.e. intrinsic buffering

through cell-specific gene expression. In *Caenorhabditis elegans*, essentiality of a gene can

depend on the natural variation in expression of other genes in the same pathway, suggesting

that functionally analogous genes in the same pathway can buffer each other [9]. The

essentiality of duplicated genes across different human cells can be associated with the

expression levels of their paralogs [10-12]. Specifically, higher expression of the buffering

paralogs is associated with diminished essentiality of their corresponding buffered duplicated

genes, indicating that cell-specific expression of paralogs can contribute to different levels of

functional buffering. Thus, expression of buffering genes, either non-duplicated or duplicated,

is constitutively active even if the cell does not experience compromised function.

Accordingly, this scenario represents an intrinsic buffering mechanism whereby expression

of the buffering genes is constitutive in specific cells and tissue types. It is likely that this

intrinsic buffering mechanism is controlled by cell- and tissue-specific transcriptional or

epigenetic regulators, leading to variable essentiality across different cells and tissues.

In this study, we directly investigated if cell-specific gene expression can act as an intrinsic

buffering mechanism (which we have termed "Cell-specific Expression-BUffering" or CEBU,

Fig. 1A) to buffer functionally related genes in the genome, thereby strengthening cellular

plasticity for cell- and tissue-specific tasks. To estimate buffering capability, we quantified

the proposed CEBU mechanism using a quantitative index, the C-score. This index calculates

the adjusted correlation between expression of a buffering gene and the essentiality of the

buffered gene (**Fig. 1B**), utilizing transcriptomics data [13] and gene dependency data from

the DepMap project [13, 14] across 684 human cell lines. Our results suggest that intrinsic

buffering by CEBU plays a critical role in cell-specific survival, tissue homeostasis, and

cancer robustness.

Results

Development of the C-score to infer cell-specific expression buffering

In seeking an index to infer cell-specific expression buffering, we postulated a buffering

relationship where the essentiality of a buffered gene (G1) increases when expression of its

buffering gene (G2) decreases. Given that G2 expression differs among cells, the strength of

functional buffering varies across cells, thereby conferring on G1 cell-specific essentiality

(**Fig. 1A**). This cell-specific expression buffering mechanism, here named CEBU, is the basis

for our development of C-score. The C-score of a gene pair is derived from the correlation

between the essentiality of a buffered gene (G1) and expression of its buffering gene (G2)

(see C-score plot, **Fig. 1B**), and is formulated as

$$\text{C-score} = \rho_{G1,G2}\left(1 + b\,\frac{slope_{min}}{slope_{G1,G2}}\right)$$

where ρ denotes the Pearson correlation coefficient between essentiality of G1 and

expression of G2. Their regression slope ($slope_{G1,\,G2}$) is normalized to $slope_{min}$, which denotes

the minimum slope of all considered gene pairs in the human genome (see **Methods**). The

normalized slope can be weighted by cell- or tissue-type specific b. In our current analysis, b

is set as 1 for a pan-cell- and pan-cancer-type analysis. Gene essentiality is represented by

gene dependency data (dependency scores, D.S.) from the DepMap project [14], where the

effect of each gene on cell proliferation was quantified after its knockout using the

CRISPR/Cas-9 approach. Specifically, a more negative dependency reflects slower cell

proliferation when the gene is knocked out, thus reflecting stronger essentiality. Expression

data was obtained through RNA-seq [13]. We anticipated that the higher the C-score of a

gene pair, the more likely G2 would buffer G1 based on the proposed CEBU mechanism.

We conducted a genome-wide analysis to calculate C-scores for all gene pairs across 684

human cell lines. The calculated C-score was compared to a bootstrapped null distribution

generated by random shuffling of G2 expression among cell lines (**Fig. S1A**). The

bootstrapped null distribution can be modeled as a normal distribution (**Fig. S1B**). For our

analysis, we considered gene pairs to have a high C-score with a strong likelihood of intrinsic

buffering when their C-scores were > 0.25 (0.058% of gene pairs in the human genome,

significant with a q-value < 2.2e-16 after multiple testing correction, **Fig. S1A**). Based on our

C-score definition, a high C-score should be indicative of marked variability in cell-specific

essentiality and expression. Indeed, we observed higher variability in both G1 dependency

and G2 expression for high C-score gene pairs (**Fig. S2A**). Nevertheless, high variation alone

is insufficient to grant a high C-score. A high C-score requires consistent pairing between G1

and G2 across cell lines and, as anticipated, disrupting the pairing between G1 dependency

and G2 expression by shuffling G2 expression amongst cell lines (without changing

variability) abolished the C-score relationship (**Fig. S2B**). Moreover, both mean G1

dependency and mean G2 expression in high C-score gene pairs were lower than those

parameters in all gene pairs (**Fig. S2C**), implying that G1s and G2s in high C-score gene

pairs tend to be more essential and less expressed, respectively.

<u>Characterization of C-score-identified gene pairs</u>

Since several duplicated gene pairs have been implicated to have functional buffering via

gene expression [10-12], we started out by characterizing the duplicated genes among C-

score-identified gene pairs. We found that duplicated gene pairs are enriched among gene

pairs with C-scores > 0.085 (p-value = 0.05 using a hypergeometric test), suggesting that

CEBU is a prevalent buffering mechanism among duplicated genes. Interestingly, the

majority of high C-score gene pairs are non-duplicated (>90%, **Fig. 2A**). In these cases, G2s

can be G1s' functional analogs as surrogate genes. Accordingly, we examined if the high C-

score gene pairs are more likely to participate in the same function or biological pathway or

physically interact. We calculated the enrichment of curated gene sets in terms of Gene

Ontology (GO) [15] and Kyoto Encyclopedia of Genes and Genomes (KEGG) [16] from the

Molecular Signatures Database [17] (**Fig. 2B**). Gene pairs with higher C-scores consistently

exhibited greater functional enrichment. Likewise, we observed a monotonic increase in the

enrichment of protein-protein interactions (PPI, STRING [18] and CORUM [19] databases)

between G1 and G2 in accordance with increasing C-score cutoff (**Fig. 2C**). The enriched

functions mostly include house-keeping functions such as nitrogen metabolism and

biosynthetic processes (**Fig. 2D**). Moreover, the observed pathway and PPI enrichments are

mostly due to the non-duplicated gene pairs since duplicated gene pairs appear to be

independent of C-scores (**Fig. S3A** and **S3B)**. In summary, duplicated gene pairs are enriched

among high C-score gene pairs and, for the non-duplicated gene pairs having high C-scores,

they are likely members of the same biological pathways and/or are encoding proteins with

physical interactions, supporting the possibility that functional buffering between these gene

pairs via CEBU.

Experimental validation of C-score-inferred cell-specific expression buffering

To validate putative C-score-inferred buffering gene pairs, we conducted experiments on the

highest C-score gene pair, i.e., *FAM50A-FAM50B*, which is a duplicated gene pair. Based on

the C-score plot of *FAM50A-FAM50B* (**Fig. 3A**), we expected that *FAM50B* would have a

stronger buffering effect for *FAM50A* in cell lines located at the top-right of the plot (e.g.

A549 and MCF7) relative to those at the bottom-left (e.g. U2OS). Consequently, growth of

the cell lines at top-right would be more sensitive to dual suppression of *FAM50A* and

FAM50B. Using gene-specific small hairpin RNAs (shRNAs), we suppressed expression of

FAM50A and *FAM50B*, individually and in combination. We observed stronger suppression

in the A549 and MCF7 cell lines relative to the U2OS cell line, as indicated by relative cell

growth (**Fig. 3B**). Using these results, we quantified genetic interactions between *FAM50A*

and *FAM50B* using Bliss score [20], whereby lower scores indicate stronger synergistic

interactions (see **Methods**). Consistent with our C-score prediction, the *FAM50A-FAM50B*

gene pair in the A549 and MCF7 cell lines exhibited stronger synergy than in the U2OS cell

line (Bliss score: 1.07 in A549, 1.06 in MCF7 and 1.44 in U2OS). Note that a recently study

showed synthetic lethality between *FAM50A* and *FAM50B* in the A375 cell line [21], which

is also located at the top-right of the *FAM50A-FAM50B* C-score plot (**Fig. 3A**).

Although duplicated genes are better recognized for their buffering relationship, there is no

direct evidence of intrinsic buffering among non-duplicated genes. Thus, we also examined a

non-duplicated gene pair, *POP7-RPP25*, representing one of the high C-score gene pairs with

physical interaction. These two genes encode protein subunits of the ribonuclease P/MRP

complex. In the C-score plot of *POP7-RPP25* (**Fig. 3C**), the HT29 cell line lies in the top-

right region and the U2OS and LN18 cell lines are in the bottom-left region, indicating the

likelihood of a stronger buffering effect in the HT29 cell line. Indeed, when we suppressed

expression of *POP7* and *RPP25* using gene-specific shRNAs in these three cell lines, we

observed that dual suppression of *POP7* and *RPP25* resulted in stronger synergistic effects

for the HT29 cell line but not for the U2OS or LN18 cell lines (Bliss scores for *POP7-RPP25*

genetic interactions are 1.02 in HT29, 1.13 in U2OS, and 1.19 in LN18, **Fig. 3D**).

<u>**Harnessing C-score to infer cell-specific genetic interactions**</u>

C-score describes the likelihood of functional buffering between two genes across hundreds

of cell lines. Thus, for a particular cell line, it is possible to use C-scores to infer genetic

interactions. As revealed by our experimental results in **Figure 3**, cell lines located at the

upper right of a C-score plot are more sensitive to dual gene suppression. We reasoned that

these cells can have greater buffering capacity relative to other cell lines, so they are more

likely to exhibit synergistic genetic interactions. To infer cell-specific genetic interactions

using C-score, we calculated buffering capacities as the relative G2 expression (compared to

that of all other cell lines) of the cell line of interest adjusted by the C-score of the gene pair

(**Fig. S4A**). We validated these C-score-derived buffering capacities for predicting genetic

interactions using experimental results from four independent studies in human cells (**Table**

S1) [22-25]. Using the receiver operating characteristic (ROC) curve to assess the

performance of buffering capacity predictions, the area under curve (AUC) is significantly

larger than random (Mann-Whitney U test with p-value < 0.05). Furthermore, the prediction

performance increased with higher C-score cutoffs, as indicated by the increase in AUC (**Fig.**

4). Moreover, in addition to the qualitative predictions described above, C-score-derived

buffering capacity appears to be quantitatively related to the strength of genetic interaction.

We observed a negative correlation between C-score-derived buffering capacities and

experimentally validated genetic interactions (C-score cutoff = 0.25, correlation = -0.231, p-

value = 0.034, **Fig. S4B**), and this correlation is stronger for higher C-score cutoffs (**Fig.**

S4C). Note that this inference on genetic interaction is robust since changing threshold for

calculating buffering capacity did not qualitatively alter the results (**Methods** and **Fig. S4D**).

Accordingly, based on these results, the CEBU mechanism can be used to infer cell-specific

genetic interactions in human cells.

<u>Tissue-specificity of C-score-derived buffering capacity</u>

Typically, the same cell and tissue types share similar gene expression profiles. Thus, next

we explored if expression of G2s in high C-score gene pairs exhibit tissue specificity. For

each G1 and G2 gene pair, we grouped cell lines based on their tissue types and applied a

tissue specificity index, τ [26], which quantifies tissue specificity (low τ, broadly expressed

across tissues; high τ, only expressed in one or a few specific tissues). As shown in **Figure**

5A, G2s generally presented higher tissue specificity compared to G1s (significant with *t*-test,

$p < 2.2e-16$) and compared to the control of randomly shuffled cell lines in G2 (**Fig. S5**).

Since G2 expression is a critical determinant of our C-score-derived buffering capacity (**Fig.**

S4A), we wondered if buffering capacity may also be tissue-specific. To investigate this

possibility, we calculated the average of buffering capacity for each tissue type based on high

C-score gene pairs (**Fig. 5B**). Our results showed distinct C-score-derived buffering

capacities across different tissues, with the top three buffered tissues being the central

nervous system, bone and the peripheral nervous system. In contrast, blood cells—including

multiple myeloma, lymphoma, and leukemia cell lines—exhibited the lowest buffering

capacities.

<u>C-score-derived buffering capacity is indicative of cancer aggressiveness</u>

If tissue-specific buffering is indeed a natural phenomenon, we wondered if cancers derived

from different tissues utilize the buffering capacities endowed by the CEBU mechanism for

more robust proliferation. In other words, would higher buffering capacity render cancers

more robust and aggressive, thereby resulting in a poorer prognosis? To test this hypothesis,

we aimed to gain a ground truth of expression-based cancer patient prognosis by analyzing

patient gene expression and survival data for all available cancer types from The Cancer

Genome Atlas (TCGA) [27] and then used this ground truth to examine the performance of

buffering capacity predictions. As an example, **Figure 6A** presents the C-score plot of

NAMPT dependency and *CALD1* expression, indicating stronger buffering capacity in cell

lines of the central nervous system (yellow) and weaker buffering capacity in leukemia cell

lines (blue). Importantly, when we stratified patients based on *CALD1* expression, we

observed a considerable difference in survival for patients with lower grade glioma (LGG,

and see **Table S2** for cross-referencing between cell lines and TCGA cancers and for the full

names of cancer abbreviations), but not for patients with acute myeloid leukemia (LAML,

Fig. 6B left panel for LGG and right panel for LAML), suggesting that buffering capacity

derived from the expression of *CALD1* and its C-score with *NAMPT* in human cell lines

could be used to infer differences in cancer patient survival for LGG but not LAML.

To gain a ground truth of expression-based cancer patient prognosis for all cancer types, we

assessed differential patient survival against gene expression using Cox regression and

controlling for clinical characteristics including age, sex, pathological stage, clinical stage,

and tumor grade, followed by multiple testing correction (false discovery rate < 0.2). We then

systematically assessed how buffering capacity from C-score-identified gene pairs could help

predict cancer patient survival for all cancer types. In **Figure 6C**, we show examples of ROC

curves for two cancer types: pancreatic adenocarcinoma (PAAD) and mesothelioma (MESO).

Examining all 30 cancer types, we found that for 20 of them, expression of at least one gene

predicted patient survival with statistical significance and, for 10 out of those 20 cancer types,

performance of buffering capacity for at least one C-score cutoff was significantly better than

random (AUC > 0.5, false discovery rate < 0.2) (**Fig. S6A**). In addition, buffering capacity is

also predictive of pathological stage (**Fig. 6D** and **S6B**), clinical stage (**Fig. S6C**), and tumor

grade (**Fig. S6D**) in multiple cancer types. In general, the buffering capacity-based

predictions performed better for greater C-score cutoffs. Taken together, our results show that

buffering capacity derived from our C-score index can be indicative of cancer aggressiveness,

as illustrated by patient survival, cancer pathological stage, clinical stage and tumor grade.

Discussion

Unlike the responsive buffering mechanism whereby the buffering gene is only activated

when its buffered function is compromised, the intrinsic buffering or CEBU mechanism

proposed herein maintains a constitutively active state with both cell- and tissue-specificity

and, thus, displays variability in buffering capacity, as illustrated in **Figure 1**. Since the

buffering gene (G2) is continuously expressed in certain cell types, there is no need for a

control system. Thus, the response time of intrinsic buffering is much quicker than responsive

buffering. By means of the CEBU mechanism, intrinsic buffering capacity can be adjusted by

altering the expression level of buffering genes, likely via epigenetic regulation, thereby

meeting cell- or tissue-specific functional needs. Therefore, CEBU describes a simple,

efficient and versatile mechanism for functional buffering in humans and other multicellular

organisms. It is likely that as an intrinsic buffering mechanism, CEBU serves distinct kinds

of functions compared to those of responsive buffering. Due to its integral and "hard-wired"

nature of cell- and tissue-specific buffering, we suspect that CEBU could play an important

role in maintaining tissue- and physiological homeostasis.

Based on our analyses, the G1s of high C-score gene pairs exhibit stronger essentiality and

are broadly expressed across tissues, whereas the G2s of such gene pairs are less essential

and expressed in specific tissues (**Fig. 5A** and **Fig. S2A**). Previously, the essentiality of genes

has been correlated with their expression level and tissue specificity [28-30]. Housekeeping

genes that are broadly expressed in most cells exhibit stronger essentiality, whereas genes

expressed in specific cell types are considered to have weaker essentiality. However, it

remains elusive as to how these two types of genes are functionally associated with each

other. Here, CEBU represents a putative mechanistic link between these two types of genes,

enabling their cooperation to regulate cellular functions through intrinsic buffering. In

particular, house-keeping functions like metabolic and cell-cycle-related processes are highly

enriched among high C-score gene pairs (**Fig. 2E**), indicating that house-keeping functions

can be more robustly maintained via CEBU-mediated regulation. Moreover, we found that

the tissues predicted to have the strongest buffering capacities are the nervous and bone

systems (**Fig. 5B**), both of which exhibit relatively low regenerative capacities. In contrast,

blood cells, which are frequently regenerated in humans, are predicted to have the weakest

buffering capacities (**Fig. 5B**). It is possible that cell types with weaker regenerative

capacities can be sustained robustly through the stronger functional buffering afforded by

CEBU, thereby maintaining their tissue homeostasis.

CEBU describes an intrinsic buffering mechanism that is expected to function under normal

physiological conditions. Indeed, when we examined if our C-score could be biased due to

our usage of cancer cell lines, we found that only a low percentage (2.3% per gene pair, **Fig.**

S7A) of mutant cell lines contributed to our C-score measurements. Moreover, excluding

mutant cell lines did not qualitatively affect our C-score measurements, especially for high C-

score gene pairs (**Fig. S7B**). The same trend holds for cancer-related genes (**Fig. S7B**). These

results suggest mutant cell lines are not the major determinants of C-scores. Similarly, since

copy number variation (CNV) is a major mechanism for oncogenic expression, we checked if

CNV contributes to G2 expression. As shown in **Figure S7C**, the correlation between G2

expression and copy number decreases with increasing C-score, indicating that CNV is not a

primary mechanism regulating G2 expression. Thus, the C-score measurement is likely not

biased by the utilization of cancer cell lines.

The high C-score gene pairs are enriched with duplicated genes, supporting the notion that

duplicated genes contribute to the context-dependent essentiality of their paralogous genes

[10-12]. Although redundant duplicated genes are thought to be evolutionarily unstable [31],

divergence in expression provides a mechanism to retain them [5-7, 12]. It is possible that

this divergence in duplicated gene expression facilitates the cell- and tissue-specific

expression of buffering genes, thus representing a critical evolutionary mechanism for CEBU

in multicellular organisms. On the other hand, despite numerous observations of functional

buffering among duplicated genes [32-34], how widely a gene can buffer another non-

orthologous gene is less well known. Our C-score index identified a high percentage of non-

duplicated gene pairs with high buffering capacities (**Fig. 2A**), and these non-duplicated gene

pairs tend to belong to the same pathways and/or protein complexes (**Fig. 2B** and **2C**).

Therefore, it is possible that many of these G1s and G2s represent non-orthologous functional

analogs. One simple scenario could be that G1 and G2 physically interact with each other to

form a protein complex, wherein G1's function can be structurally substituted by G2. Indeed,

we identified the *POP7* and *RPP25* gene pair as an example of this scenario (**Fig. 3C** and

3D). More sophisticated and indirect functional buffering can also occur between G1s and

G2s given the complex interactions among biological functions [35]. We expect that

additional types of molecular mechanisms can account for the buffering effects of CEBU,

which remain to be tested experimentally.

C-score-derived cell-specific buffering capacities comply well with experimentally validated

genetic interactions in human cells (**Fig. 4** and **Fig. S4**), indicating that CEBU may represent

a critical mechanism for synthetic lethality in human cells. In practice, it remains a daunting

challenge to systematically characterize genetic interactions in organisms with complex

genomes due to large numbers of possible gene pairs, i.e. ~200 million gene pairs in humans.

Previous efforts have used computational approaches on conserved synthetically lethal gene

pairs in the budding yeast *Saccharomyces cerevisiae* or employ data mining on multiple large

datasets to infer human synthetic lethalities [36, 37]. Nevertheless, predictions emanating

from different studies exhibit little overlap [38], evidencing the marked complexity of

synthetic lethality in humans. The CEBU mechanism proposed here can contribute both

experimentally and computationally to a better characterization of human genetic interactions.

Using G2 expression of a single high C-score gene pair to stratify cancer patients, we

observed a significant difference in cancer patient survival, indicating that stronger buffering

capacity (as derived from C-scores and relative G2 expression in human cell lines) could be

predictive of cancer aggressiveness in patients (see **Fig. 6A** and **6B** for an example). Indeed,

buffering capacity helps predict cancer patient survival in ten cancer types, and generally the

performance of prediction is better with a higher C-score cutoff (**Fig. 6C** and **S6A**). Apart

from patient survival, buffering capacity is also indicative of pathological stage, clinical stage,

and tumor grade of cancers (**Fig. 6D** and **S6B-6D**). These results support our hypothesis that

stronger buffering capacity via higher G2 expression contributes to cancer robustness in

terms of proliferation and drug resistance. Given the complexity of cancers, it is surprising to

see such general predictivity of cancer prognosis by individual high C-score gene pairs.

Accordingly, we suspect that some cancer cells may adopt this cell- and tissue-specific

buffering mechanism to enhance their robustness in proliferation and stress responses by

maintaining or even upregulating the gene expression of buffering genes. Clinically, the

expression of such buffering genes could represent a unique feature for evaluating cancer

progression when applied alongside other currently used clinical characteristics. Finally,

experimental validation of C-score-predicted genetic interactions in cancers will help identify

potential drug targets for tailored combination therapy based on cancer types and their

expression profiles.

Materials and Methods

Dependencies and gene expression data retrieval and processing

Data on dependencies and CCLE (Cancer Cell Line Encyclopedia) gene expression were

downloaded from the DepMap database (DepMap Public 19Q4) [13, 14]. Dependencies

modeled from the CERES computational pipeline based on a genome-wide CRISPR loss-of-

function screening were selected. CCLE expression data was quantified as $\log_2$ TPM

(Transcripts Per Million) using RSEM (RNA-seq by Expectation Maximization) with a

pseudo-count of 1 in the GTEx pipeline (https://gtexportal.org/home/documentationPage).

Only uniquely mapped reads in the RNA-seq data were used in the GTEx pipeline.

Integrating and cross-referencing the dependency and expression datasets yielded 18239

genes and 684 cell lines. Genes lacking dependencies for any one of the 684 cell lines were

discarded from our analyses.

C-score calculation

Our C-score index integrates the dependencies of buffered genes (G1) and gene expression of

buffering genes (G2) to determine the buffering relationship between gene pairs. Genes with

mean dependencies > 0 or mean gene expression < 0.5 $\log_2$ TPM were discarded, yielding

9196 G1s and 13577 G2s. The C-score integrates the correlation (ρ) and slope between the

dependency of gene $G1$ and the gene expression of gene $G2$, defined as:

$$\text{C-score} = \rho_{G1,G2} \left(1 + b \frac{slope_{min}}{slope_{G1,G2}} \right)$$

where ρ denotes the Pearson correlation coefficient and $slope_{min}$ denotes the minimum slope

of all considered gene pairs that present a statistically significant positive correlation. The

normalized slope can be weighted by cell- and tissue-type specific b. In our analysis, b is set

as 1 for a pan-cell or pan-cancer analysis.

Duplicated gene assignment

392   Information on gene identity was obtained from ENSEMBL (release 98, reference genome

393   GRCh38.p13) [39]. Two genes are considered duplicated genes if they have diverged from

394   the same duplication event.

Enrichment analysis for buffering gene pairs

397   For the enrichment analysis of gene pairs, we used a previously described methodology [40].

398   Briefly, gene sets of GO and KEGG were downloaded from the Molecular Signatures

399   Database. The number of total possible gene pairs is 9196 (G1) x 13577 (G2). The condition

400   of G1 and G2 being the same gene was not considered as a buffering gene pair under all C-

401   score cutoffs. Enrichment was calculated as:

$$\log \frac{\frac{e_{ac}}{e_a}}{\frac{e_c}{e_t}}$$

where e_{ac} represents the number of gene pairs that are both annotated and with buffering

capability, e_a is the number of annotated gene pairs, e_c is the number of buffering gene pairs,

and e_t is the total number of gene pairs.

Protein-protein interaction (PPI) data was downloaded from the STRING database (version

11) [18]. Only high-confidence interactions (confidence > 0.7) in human were considered.

The STRING database determines confidence by approximating the probability that a link

exists between two enzymes in the KEGG database. Data on protein core complexes were

downloaded from CORUM (http://mips.helmholtz-muenchen.de/corum). The enrichment

calculation is the same as for GO and KEGG, except that e_{ac} represents the number of gene

pairs that have PPI or are in the same complex and have buffering capability, and e_a is the

number of gene pairs that have PPI or are in the same complex.

Experimental validation

A549, H4, HT29, LN18, MCF7, and U2OS cell lines were selected based on their locations

in the C-score plots (**Fig. 2A and 2C**), indicating different buffering capacity. All cell lines

were purchased from ATCC and they were cultured in Dulbecco's Modified Eagle Media

(H4 and LN18), Ham's F-12K Medium (A549), or RPMI 1640 media (HT29, MCF7, and

U2OS) supplemented with 5% fetal bovine, serum, 100 U/mL penicillin, 100 μg/mL

streptomycin, and 250 ng/mL fungizone (Gemini Bio-Products). Cell growth was monitored

by time-lapse imaging using Incucyte Zoom, taking images every 2 hours for 2-4 days. To

suppress *FAM50A, FAM50B*, *POP7* and *RPP25* expression, lentivirus-based shRNAs were

delivered individually or in combination. The gene-specific shRNA sequences are: *FAM50A*

- CCAACATTGACAAGAAGTTCT and GAGCTGGTACGAGAAGAACAA; *FAM50B* –

CACCTTCTACGACTTCATCAT; *POP7* – CTTCAGGGTCACACCCAAGTA and

CGGAGACCCAATGACATTTAT; and *RPP25* – CCAGCGTCCAAGAGGAGCCTA. To

ensure better knockdown of gene expression, shRNAs were delivered twice (7 days and 4

days before seeding). Equal numbers of cells were seeded for cell growth measurements by

time-lapse imaging using Incucyte Zoom. The lentivirus-based shRNAs were purchased from

the RNAi core of Academia Sinica. The growth rate of each condition was measured by

fitting cell confluence to an exponential growth curve using the Curve Fitting Toolbox in

MATLAB.

Bliss independence model

The synergy of cytotoxicity was measured using the Bliss independent model [20]. The Bliss

model is presented as a ratio of the expected additive effect to the observed combination

effect:

$$E_{bliss} = \frac{E_A + E_B - E_A \times E_B}{E_{AB}}$$

where E is the effect of drug A, B, or a combination of A and B. Effect was measured by the

relative cell growth, based on the fold-change of confluency between 0 and 72 hours upon

suppression of $FAM50A$ and $FAM50B$ or suppression of $POP7$ and $RPP25$ in all cell lines

except MCF7, and between 0 and 96 hours upon suppression of $FAM50A$ and $FAM50B$ in

MCF7.

Cell-specific buffering capacity and comparison with experimental genetic interactions

Cell-specific buffering capacity was derived from the C-score of a given gene pair and gene

expression of the buffering gene (G2) in the cell line of interest following the equation:

$$\text{buffering capacity} = \frac{\text{cell line expression} - \text{25th percentile of all expression (G2)}}{slope_{mod}}$$

$$\text{where } slope_{mod} = \text{C-score} \times \frac{sd(G2\ expression)}{sd(G1\ dependency)}$$

where sd = standard deviation. The 25th percentile cutoff for expression is determined

empirically, although different percentile cutoffs do not qualitatively affect the measurements

of buffering capacities (**Fig. S4A**).

Combinatorial CRISPR screen-derived genetic interaction scores were pooled from four

literature sources [22-25] (**Table S1**). We only considered cell lines that appear in DepMap

CERES 19Q4. There were two C-scores for each gene-pair of the experimental dataset (either

gene could be a G1), and we assigned the higher C-score for that gene-pair. Overall, we

curated 10,222 genetic interaction scores in various cell lines from the literature, and 1986

out of 10,222 genetic interaction scores had a C-score > 0.1. To evaluate the validity of

buffering capacity, we generated a ground truth table by assigning gene-pairs with a positive

genetic interaction as false for buffering and a negative genetic interaction as true for

buffering. The qualitative performance of buffering capacity to the ground truth data was

assessed by ROC curve. Additionally, we correlated the buffering capacity directly via a

ground truth genetic interaction score for quantitative evaluation.

Tissue specificity

To calculate tissue-specificity, cell lines were grouped by their respective tissues, and

expression of genes in cell lines of the same tissue were averaged. Tissue specificity was

calculated as tau (τ) [26], where τ is defined as:

$$\tau = \frac{\sum_{i-1}^{N}(1 - \frac{x_i}{x_{max}})}{N - 1}$$

with N denoting the number of tissues, x_i denoting the expression of a gene, and x_{max} denoting

the highest gene expression across all tissues. Note, expression values were log-transformed,

so $\log_2$ TPM < 1 was considered as 0 in tissue specificity calculations [41].

Cancer-specific survival prediction according to C-score gene pairs

Gene expression and survival data from The Cancer Genome Atlas (TCGA) [27] was

retrieved from Xena [42]. The DepMap cancer cell lines were mapped to TCGA cancers

based on the annotation in **Table S2** (cancers that do not have a matched cancer type in

CERES 19Q4 were not analyzed). To systematically analyze cancer prognosis, we first

performed a multiple test correction on the p-values from Cox regression controlling for age,

sex, pathological stage, clinical stage and tumor grade. We calculated the false discovery rate

(FDR) using the Benjamini–Hochberg procedure with a threshold < 0.2. The ground truth

table for each cancer was constructed using the adjusted p-value. AUC of ROC curves were

used to assess the performance of survival based on buffering capacity. AUCs and ROCs

were generated using python. The statistical significance of AUC was assessed by using the

Mann-Whitney U test [43] to evaluate whether gene expression with a positive Cox

coefficient (poorer prognosis) reflected significantly higher buffering capacities in each

cancer with different C-score cut-offs. The p-values of the Mann-Whitney U test were

adjusted using the Benjamini-Hochberg procedure with a threshold < 0.2. We conducted a

similar approach to the prognosis analysis for buffering capacities and clinical features. We

calculated the p-values of correlation of gene expression and clinical features, staging and

grade, followed by correcting the p-values using the Benjamini-Hochberg procedure with a

threshold < 0.2. We then used Mann-Whitney U tests to evaluate if gene pairs with a

significant positive correlation between gene expression and tumor aggressiveness presented

a significantly higher buffering capacity in each cancer for different C-score cut-offs.

Data Availability

All data and code used for this article are available in the FigShare repository

https://figshare.com/s/6f8929c6543687a6062f, and

https://figshare.com/s/b778489bb2f6fc3b0069, respectively.

Acknowledgement

We thank members of the Lab for Cell Dynamics for helpful discussions. We are grateful to

Jose Reyes, Hannah Katrina Co, and Jun-Yi Leu for their comments and suggestions on the

manuscript, and Ann Mikaela Lynne Ong Co, Jo-Hsi Huang, Su-Ping Lee, the Imaging Core

at the Institute of Molecular Biology, and the RNAi Core at Academia Sinica for their

technical support.

Author Contributions

H.-K.L., J.-H.C., and S.-h.C. conceived the project. H.-K.L., J.-H.C., C.-C.W., and S.-h.C.

designed and conducted the computational and statistical analyses. H.-K.L., J.-H.C., C.-C.W.,

and S.-h.C. wrote the manuscript. F.-S.H., C.D. and S.-h.C. designed the experiments, and F.-

S.H., and C.D. conducted the experiments.

Conflict of Interests

J.-H.C. is an employee of ACT Genomics

References and Citations
518
1.   Kitano H. Biological robustness. Nat Rev Genet. 2004;5(11):826-37. Epub
2004/11/03. doi: 10.1038/nrg1471. PubMed PMID: 15520792.
2.   Masel J, Siegal ML. Robustness: mechanisms and consequences. Trends Genet.
2009;25(9):395-403. doi: 10.1016/j.tig.2009.07.005. PubMed PMID:
WOS:000270185900004.
3.   El-Brolosy MA, Stainier DYR. Genetic compensation: A phenomenon in search of
mechanisms. PLoS Genet. 2017;13(7):e1006780. Epub 2017/07/14. doi:
10.1371/journal.pgen.1006780. PubMed PMID: 28704371; PubMed Central PMCID:
PMCPMC5509088.
4.   Diss G, Ascencio D, DeLuna A, Landry CR. Molecular mechanisms of paralogous
compensation and the robustness of cellular networks. J Exp Zool B Mol Dev Evol.
2014;322(7):488-99. Epub 2014/01/01. doi: 10.1002/jez.b.22555. PubMed PMID: 24376223.
5.   Kafri R, Bar-Even A, Pilpel Y. Transcription control reprogramming in genetic
backup circuits. Nat Genet. 2005;37(3):295-9. Epub 2005/02/22. doi: 10.1038/ng1523.
PubMed PMID: 15723064.
6.   Kafri R, Levy M, Pilpel Y. The regulatory utilization of genetic redundancy through
responsive backup circuits. P Natl Acad Sci USA. 2006;103(31):11653-8. doi:

10.1073/pnas.0604883103. PubMed PMID: WOS:000239616400042.
7.   Kafri R, Springer M, Pilpel Y. Genetic Redundancy: New Tricks for Old Genes. Cell.
2009;136(3):389-92. doi: 10.1016/j.cell.2009.01.027. PubMed PMID:
WOS:000263120600006.
8.   Koonin EV, Mushegian AR, Bork P. Non-orthologous gene displacement. Trends
Genet. 1996;12(9):334-6. Epub 1996/09/01. PubMed PMID: 8855656.
9.   Vu V, Verster AJ, Schertzberg M, Chuluunbaatar T, Spensley M, Pajkic D, et al.
Natural Variation in Gene Expression Modulates the Severity of Mutant Phenotypes. Cell.
2015;162(2):391-402. Epub 2015/07/18. doi: 10.1016/j.cell.2015.06.037. PubMed PMID:
26186192.
10.   Wang T, Birsoy K, Hughes NW, Krupczak KM, Post Y, Wei JJ, et al. Identification
and characterization of essential genes in the human genome. Science. 2015;350(6264):1096-
101. Epub 2015/10/17. doi: 10.1126/science.aac7041. PubMed PMID: 26472758; PubMed
Central PMCID: PMCPMC4662922.
11.   Dandage R, Landry CR. Paralog dependency indirectly affects the robustness of
human cells. Mol Syst Biol. 2019;15(9):e8871. Epub 2019/09/27. doi:
10.15252/msb.20198871. PubMed PMID: 31556487; PubMed Central PMCID:
PMCPMC6757259.
12.   De Kegel B, Ryan CJ. Paralog buffering contributes to the variable essentiality of
genes in cancer cell lines. Plos Genetics. 2019;15(10). doi: ARTN e1008466
10.1371/journal.pgen.1008466. PubMed PMID: WOS:000503176900046.
13.   Ghandi M, Huang FW, Jane-Valbuena J, Kryukov GV, Lo CC, McDonald ER, et al.
Next-generation characterization of the Cancer Cell Line Encyclopedia. Nature.
2019;569(7757):503-+. doi: 10.1038/s41586-019-1186-3. PubMed PMID:
WOS:000468844100040.
14.   Meyers RM, Bryan JG, McFarland JM, Weir BA, Sizemore AE, Xu H, et al.
Computational correction of copy number effect improves specificity of CRISPR-Cas9
essentiality screens in cancer cells. Nature Genetics. 2017;49(12):1779-+. doi:
10.1038/ng.3984. PubMed PMID: WOS:000416480600018.
15.   Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, et al. Gene
ontology: tool for the unification of biology. The Gene Ontology Consortium. Nat Genet.
2000;25(1):25-9. Epub 2000/05/10. doi: 10.1038/75556. PubMed PMID: 10802651; PubMed
Central PMCID: PMCPMC3037419.
16.   Kanehisa M, Goto S. KEGG: kyoto encyclopedia of genes and genomes. Nucleic
Acids Res. 2000;28(1):27-30. Epub 1999/12/11. doi: 10.1093/nar/28.1.27. PubMed PMID:
10592173; PubMed Central PMCID: PMCPMC102409.
17.   Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA, et al.
Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide
expression profiles. Proc Natl Acad Sci U S A. 2005;102(43):15545-50. Epub 2005/10/04.
doi: 10.1073/pnas.0506580102. PubMed PMID: 16199517; PubMed Central PMCID:
PMCPMC1239896.
18.   von Mering C, Jensen LJ, Snel B, Hooper SD, Krupp M, Foglierini M, et al.
STRING: known and predicted protein-protein associations, integrated and transferred across
organisms. Nucleic Acids Res. 2005;33(Database issue):D433-7. Epub 2004/12/21. doi:
10.1093/nar/gki005. PubMed PMID: 15608232; PubMed Central PMCID: PMCPMC539959.
19.   Ruepp A, Waegele B, Lechner M, Brauner B, Dunger-Kaltenbach I, Fobo G, et al.
CORUM: the comprehensive resource of mammalian protein complexes--2009. Nucleic
Acids Res. 2010;38(Database issue):D497-501. Epub 2009/11/04. doi: 10.1093/nar/gkp914.
PubMed PMID: 19884131; PubMed Central PMCID: PMCPMC2808912.
20.   Bliss CI. The toxicity of poisons applied jointly. Annals of Applied Biology.

1939;26(3):585–615.

21.   Thompson NA, Ranzani M, van der Weyden L, Iyer V, Offord V, Droop A, et al.
Combinatorial CRISPR screen identifies fitness effects of gene paralogues. Nat Commun.
2021;12(1):1302. Epub 2021/02/28. doi: 10.1038/s41467-021-21478-9. PubMed PMID:
33637726; PubMed Central PMCID: PMCPMC7910459.

22.   Rosenbluh J, Mercer J, Shrestha Y, Oliver R, Tamayo P, Doench JG, et al. Genetic and
Proteomic Interrogation of Lower Confidence Candidate Genes Reveals Signaling Networks
in beta-Catenin-Active Cancers. Cell Syst. 2016;3(3):302-+. doi: 10.1016/j.cels.2016.09.001.
PubMed PMID: WOS:000395775300012.

23.   Shen JP, Zhao DX, Sasik R, Luebeck J, Birmingham A, Bojorquez-Gomez A, et al.
Combinatorial CRISPR-Cas9 screens for de novo mapping of genetic interactions. Nat
Methods. 2017;14(6):573-+. doi: 10.1038/nmeth.4225. PubMed PMID:
WOS:000402291800016.

24.   Najm FJ, Strand C, Donovan KF, Hegde M, Sanson KR, Vaimberg EW, et al.
Orthologous CRISPR-Cas9 enzymes for combinatorial genetic screens. Nat Biotechnol.
2018;36(2):179-89. Epub 2017/12/19. doi: 10.1038/nbt.4048. PubMed PMID: 29251726;
PubMed Central PMCID: PMCPMC5800952.

25.   Zhao D, Badur MG, Luebeck J, Magana JH, Birmingham A, Sasik R, et al.
Combinatorial CRISPR-Cas9 Metabolic Screens Reveal Critical Redox Control Points
Dependent on the KEAP1-NRF2 Regulatory Axis. Mol Cell. 2018;69(4):699-708 e7. Epub
2018/02/18. doi: 10.1016/j.molcel.2018.01.017. PubMed PMID: 29452643; PubMed Central
PMCID: PMCPMC5819357.

26.   Yanai I, Benjamin H, Shmoish M, Chalifa-Caspi V, Shklar M, Ophir R, et al.
Genome-wide midrange transcription profiles reveal expression level relationships in human
tissue specification. Bioinformatics. 2005;21(5):650-9. Epub 2004/09/25. doi:
10.1093/bioinformatics/bti042. PubMed PMID: 15388519.

27.   Hoadley KA, Yau C, Hinoue T, Wolf DM, Lazar AJ, Drill E, et al. Cell-of-Origin
Patterns Dominate the Molecular Classification of 10,000 Tumors from 33 Types of Cancer.
Cell. 2018;173(2):291-304 e6. Epub 2018/04/07. doi: 10.1016/j.cell.2018.03.022. PubMed
PMID: 29625048; PubMed Central PMCID: PMCPMC5957518.

28.   Hastings KEM. Strong evolutionary conservation of broadly expressed protein
isoforms in the troponin I gene family and other vertebrate gene families. J Mol Evol.
1996;42(6):631-40. doi: Doi 10.1007/Bf02338796. PubMed PMID:
WOS:A1996UU51200002.

29.   Subramanian S, Kumar S. Gene expression intensity shapes evolutionary rates of the
proteins encoded by the vertebrate genome. Genetics. 2004;168(1):373-81. doi:
10.1534/genetics.104.028944. PubMed PMID: WOS:000224393700029.

30.   Zhang LQ, Li WH. Mammalian housekeeping genes evolve more slowly than tissue-
specific genes. Mol Biol Evol. 2004;21(2):236-9. doi: 10.1093/molbev/msh010. PubMed
PMID: WOS:000220083300005.

31.   Lynch M, Conery JS. The evolutionary fate and consequences of duplicate genes.
Science. 2000;290(5494):1151-5. Epub 2000/11/10. doi: 10.1126/science.290.5494.1151.
PubMed PMID: 11073452.

32.   Gu ZL, Steinmetz LM, Gu X, Scharfe C, Davis RW, Li WH. Role of duplicate genes
in genetic robustness against null mutations. Nature. 2003;421(6918):63-6. doi:
10.1038/nature01198. PubMed PMID: WOS:000180165500037.

33.   DeLuna A, Springer M, Kirschner MW, Kishony R. Need-Based Up-Regulation of
Protein Levels in Response to Deletion of Their Duplicate Genes. Plos Biol. 2010;8(3). doi:
ARTN e1000347
10.1371/journal.pbio.1000347. PubMed PMID: WOS:000278125400021.

34.   Deutscher D, Meilijson I, Kupiec M, Ruppin E. Multiple knockout analysis of genetic
robustness in the yeast metabolic network. Nat Genet. 2006;38(9):993-8. Epub 2006/08/31.
doi: 10.1038/ng1856. PubMed PMID: 16941010.
35.   Rancati G, Moffat J, Typas A, Pavelka N. Emerging and evolving concepts in gene
essentiality. Nat Rev Genet. 2018;19(1):34-49. Epub 2017/10/17. doi: 10.1038/nrg.2017.74.
PubMed PMID: 29033457.
36.   Jerby-Arnon L, Pfetzer N, Waldman YY, McGarry L, James D, Shanks E, et al.
Predicting cancer-specific vulnerability via data-driven detection of synthetic lethality. Cell.
2014;158(5):1199-209. Epub 2014/08/30. doi: 10.1016/j.cell.2014.07.027. PubMed PMID:
25171417.
37.   Srivas R, Shen JP, Yang CC, Sun SM, Li JF, Gross AM, et al. A Network of
Conserved Synthetic Lethal Interactions for Exploration of Precision Cancer Therapy.
Molecular Cell. 2016;63(3):514-25. doi: 10.1016/j.molcel.2016.06.022. PubMed PMID:
WOS:000381619900016.
38.   Liu L, Chen X, Hu C, Zhang D, Shao Z, Jin Q, et al. Synthetic Lethality-based
Identification of Targets for Anticancer Drugs in the Human Signaling Network. Sci Rep.
2018;8(1):8440. Epub 2018/06/02. doi: 10.1038/s41598-018-26783-w. PubMed PMID:
29855504; PubMed Central PMCID: PMCPMC5981615.
39.   Cunningham F, Achuthan P, Akanni W, Allen J, Amode MR, Armean IM, et al.
Ensembl 2019. Nucleic Acids Res. 2019;47(D1):D745-D51. Epub 2018/11/09. doi:
10.1093/nar/gky1113. PubMed PMID: 30407521; PubMed Central PMCID:
PMCPMC6323964.
40.   Boyle EA, Pritchard JK, Greenleaf WJ. High-resolution mapping of cancer cell
networks using co-functional interactions. Molecular Systems Biology. 2018;14(12). doi:
ARTN e8594
10.15252/msb.20188594. PubMed PMID: WOS:000453861400001.
41.   Kryuchkova-Mostacci N, Robinson-Rechavi M. A benchmark of gene expression
tissue-specificity metrics. Brief Bioinform. 2017;18(2):205-14. Epub 2016/02/20. doi:
10.1093/bib/bbw008. PubMed PMID: 26891983; PubMed Central PMCID:
PMCPMC5444245.
42.   Vivian J, Rao AA, Nothaft FA, Ketchum C, Armstrong J, Novak A, et al. Toil enables
reproducible, open source, big biomedical data analyses. Nat Biotechnol. 2017;35(4):314-6.
Epub 2017/04/12. doi: 10.1038/nbt.3772. PubMed PMID: 28398314; PubMed Central
PMCID: PMCPMC5546205.
43.   Mason SJ, Graham NE. Areas beneath the relative operating characteristics (ROC)
and relative operating levels (ROL) curves: Statistical significance and interpretation.
Quarterly Journal of the Royal Meteorological Society. 2002;128(584):2145-66. doi:
10.1256/003590002320603584.
44.   Mi H, Ebert D, Muruganujan A, Mills C, Albou LP, Mushayamaha T, et al.
PANTHER version 16: a revised family classification, tree-based classification tool, enhancer
regions and extensive API. Nucleic Acids Res. 2020. Epub 2020/12/09. doi:
10.1093/nar/gkaa1106. PubMed PMID: 33290554.

678
679
680

Figure Legends

Figure 1. Genome-wide CEBU analysis using C-score

C-score plot: the x-axis is the dependency of the buffered gene (G1) and the y-axis is the

expression of the buffering gene (G2). G1 is considered as being potentially buffered by G2,

as quantified by C-score, which is an adjusted correlation for a given gene based on the C-

score plot.

Figure 2. Properties of high C-score gene pairs

(A) Percentage of C-score-identified non-duplicated and duplicated gene pairs with different

C-score cut-offs. (B-C) Functional enrichment of C-score gene pairs increases with C-score

cutoff. (B) Enrichment of pairs of genes annotated with the same gene ontology biological

process (GO:BP) term or KEGG pathway in C-score gene pairs, and enrichment increases

with C-score cut-off. (C) Enrichment of pairs of genes with annotated protein-protein

interactions from STRING and within the same protein complex from CORUM in C-score

gene pairs, and enrichment increases with C-score cut-off. (D) Top 10 GO:BP enrichment of

high C-score gene pairs with the same functional annotation. GO was annotated using

PANTHER 15.0 [44] , and selected using GO Slim terms.

Figure 3. Experimental validation of cell-specific expression buffering between

FAM50A* and *FAM50B*, and *POP7* and *RPP25

(A) C-score plot of the highest C-score gene pair, *FAM50A* (dependency score - D.S.) and

FAM50B (G2) expression, labeled with the U2OS (predicted not synergistic), A549

(predicted synergistic), and MCF7 (predicted synergistic) cell lines. (B) Relative cell growth

based on fold-change in confluency of the U2OS, A549, and MCF7 cell lines with or without

FAM50A or *FAM50B* suppression. Bliss scores indicate strength of synergy between double

suppression of *FAM50A* and *FAM50B* compared to either gene alone. Error bars indicate

standard deviation of six technical repeats. (C) C-score plot of the non-duplicated gene pair,

POP7 D.S. and *RPP25* expression, labeled with the LN18 (predicted not synergistic), U2OS

(predicted not synergistic), and HT29 (predicted synergistic) cell lines. (D) Relative cell

growth based on fold-change in confluency of the HT29, U2OS and LN18 cell lines with or

without shRNA-based *POP7* or *RPP25* suppression. Bliss scores indicate strength of synergy

between double suppression of *POP7* and *RPP25* compared to either gene alone. Error bars

indicate standard deviation of six technical repeats.

Figure 4. C-score-derived cell-specific buffering capacity predicts genetic interaction

and cancer patient survival

Predictive performance shown as ROC curves for predicting genetic interactions using cell-

specific buffering capacity. Prediction sets consist of 84 data-points (37 unique genetic

interaction gene-pairs) across 8 cell lines with a C-score cut-off of 0.25.

Figure 5. C-score-derived tissue-specific buffering capacity

(A) Tissue specificity (τ) of G1 and G2 pairs. τ was calculated for G1 and G2 from high C-

score gene pairs. Statistical significance was assessed by paired-*t* test. (B) Mean buffering

capacity for each tissue type.

Figure 6. Harnessing cell-specific high C-score gene pairs for cancer patient prognosis

(A) C-score plot of *NAMPT* D.S. and *CALD1* expression (C-score = 0.447). Yellow circles

represent central nervous system (CNS) cell lines and blue circles denote leukemia cell lines.

(B) Kaplan-Meier overall survival plots for CNS (LGG, lower grade glioma, left panel) and

leukemia (LAML, acute myeloid leukemia, right panel) cancer patients. Patients were

stratified by high (>75%) or low (<25%) expression of *CALD1*, and *p*-values were calculated

using Cox regression controlling for age, sex, pathological staging, clinical staging, and

tumor grade, and corrected for multiple testing (false discovery rate < 0.2). (C) ROC curves

based on C-score gene pair-based buffering prediction of survival for PAAD (pancreatic

adenocarcinoma, left panel) and MESO (mesothelioma, right panel) with different C-score

cutoffs. (D) ROC curves based on C-score gene pair-based buffering prediction of

pathological stage for PAAD with different C-score cutoffs.

Supporting Information Captions

Figure S1. Distribution of C-scores

Figure S2. Correlation properties of G1 dependency and G2 expression according to

increasing C-score

Figure S3. Functional and pathway enrichments of C-score-identified duplicated genes

Figure S4. C-score-based prediction of cell-specific genetic interaction using buffering

capacity

Figure S5. Shuffling the G2-tissue relationship disrupts G2 tissue-specificity

Figure S6. C-score-based prediction of cancer patient survival, pathological stage,

clinical stage and tumor grade

Figure S7. Decreasing effects of mutational variation as C-score increases

Table S1. List of combinatorial CRISPR-screened gene pairs in published literature

Table S2. Cross-reference table for TCGA cancer type and DepMap cancer type

Figure 1

A   **C**ell-specific **E**xpression-**BU**ffering mechanism (**CEBU**)

B   C-score plot

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6