Barcoded reciprocal hemizygosity analysis via sequencing illuminates the complex genetic basis of yeast thermotolerance

Melanie B. Abrams* [1], Julie N. Chuong* [†] [1], Faisal AlZaben*[2], Claire A. Dubin*[2], Jeffrey M. Skerker^[3], and Rachel B. Brem* [‡]

*Department of Plant and Microbial Biology, UC Berkeley, Berkeley, CA; [†]Ph.D. Program in Biology, New York University, New York, NY; ^ Environmental Genomics and Systems Biology Division, Lawrence Berkeley National Laboratory, Berkeley, CA; [‡]Buck Institute for Research on Aging, Novato, CA

[1]These authors contributed equally to this work.
[2]These authors contributed equally to the work
[3]Current address: Zymergen, Inc., 5980 Horton St #105, Emeryville, CA 94608

17   Barcoded RH-seq and yeast thermotolerance
18

19   Keywords: Evolution, Genetics, Saccharomyces, Thermotolerance, Adaptation

21   Corresponding authors:
22
Rachel Brem, Ph.D.
Department of Plant and Microbial Biology
University of California, Berkeley
312F Innovative Genomics Institute Building
2151 Berkeley Way
Berkeley, CA 94730
rbrem@berkeley.edu
30
31   Jeffrey Skerker, Ph.D.
32   Zymergen, Inc.
33   5980 Horton St #105
34   Emeryville, CA 94608
35   skerker1@gmail.com

ABSTRACT

Decades of successes in statistical genetics have revealed the molecular underpinnings of traits as they vary across individuals of a given species. But standard methods in the field can't be applied to divergences between reproductively isolated taxa. Genome-wide reciprocal hemizygosity mapping (RH-seq), a mutagenesis screen in an inter-species hybrid background, holds promise as a method to accelerate the progress of interspecies genetics research. Here we describe an improvement to RH-seq in which mutants harbor barcodes for cheap and straightforward sequencing after selection in a condition of interest. As a proof of concept for the new tool, we carried out genetic dissection of the difference in thermotolerance between two reproductively isolated budding yeast species. Experimental screening identified dozens of candidate loci at which variation between the species contributed to the thermotolerance trait. Hits were enriched for mitosis genes and other housekeeping factors, and among them were multiple loci with robust sequence signatures of positive selection. Together, these results shed new light on the mechanisms by which evolution solved the problems of cell survival and division at high temperature in the yeast clade, and they illustrate the power of the barcoded RH-seq approach.

INTRODUCTION

Understanding how and why organisms from the wild exhibit different traits is a central goal of modern genetics. Linkage and association mapping have driven decades of success in dissecting trait variation across individuals of a given species (Ott et al. 2015; Tam et al. 2019). But since these methods can't be applied to reproductively isolated taxa, progress in the field of interspecies genetics has lagged behind. However, newer statistical-genetic methods appropriate to comparisons between species have been proposed in the recent literature (Weiss and Brem 2019), which hold promise for elucidating the genetics of ancient traits. For most such methods, limitations accruing from throughput and/or coverage issues remain to be refined.

The budding yeast *Saccharomyces cerevisiae* grows better at high temperature than any other species in its clade (Sweeney et al. 2004; Gonçalves et al. 2011; Salvadó et al. 2011; Hittinger 2013; Weiss et al. 2018), in keeping with its likely ecological origin in hot, East Asian locales (Peter et al. 2018). This derived and putatively adaptive trait serves as a model for the genetic study of deep evolutionary divergences. Thermosensitivity, the ancestral phenotype in the clade, is borne out in *S. paradoxus*, a close sister species to *S. cerevisiae*, making the former a useful point of comparison. Our group previously used this system as a testbed to develop RH-seq (Weiss et al. 2018), a genomic version of the reciprocal hemizygosity test (Stern 2014) that is well-suited to the mapping of natural trait variation between sister species. This technique starts with the generation of large numbers of random transposon mutant clones of a viable but sterile interspecies hybrid. In a given clone, loss of function from a transposon insertion in one species' allele of a gene reveals the function of the uncovered allele from the other species. These hemizygotes are competed en masse in a condition of interest; the abundance of each hemizygote in turn in the selected pool is quantified by bulk sequencing, and used in a test for allelic impact on the focal trait. In previous work, we identified eight genes through this approach at which species divergence contributed to thermotolerance (Weiss et al. 2018).

Against a backdrop of successful biological and evolutionary inference from our yeast RH-seq pilot (Weiss et al. 2018; Abrams et al. 2021), we noted that the combination of *S. cerevisiae* alleles of all eight genes mapped to thermotolerance recapitulated only <20% of the difference between the species (AlZaben et al. 2021). Thus, many of the determinants of yeast thermotolerance likely remain undetected. If so, boosting the replication and throughput of

genetic mapping, to enable higher statistical power, could help meet the challenge. In our initial
implementation of RH-seq, we had quantified the abundance of hemizygotes in a sample by
sequencing across the transposon junction with the genome, using one universal primer that
recognized the transposon and another recognizing a ligated adapter at DNA fragment ends
(Weiss et al. 2018). This protocol, though rigorous, is labor-intensive and expensive, limiting the
potential for throughput and coverage. A higher-throughput alternative starts with the tagging of
transposon sequences by random short DNA barcodes (Wetmore et al. 2015). After
mutagenesis of a genotype of interest by these barcoded transposons, and then selection of the
mutants in bulk in a challenging condition, mutant abundance can be quantified from
sequencing of DNA straight from the pool with a simple PCR. We set out to adapt this barcoding
strategy to enable highly replicated RH-seq, with application to yeast thermotolerance as a test
case to achieve a deeper exploration of the complex genetics of the trait.
100
101   **MATERIALS AND METHODS**
102
103 **Construction of a randomly barcoded piggyBac transposase pool**
104
105 For barcoded RH-seq, we constructed a pool of plasmids, each harboring the piggyBac
106 transposase and a randomly barcoded copy of the piggyBac transposon, via Golden Gate
107 cloning of random 20bp barcodes flanked by universal priming sites into a plasmid backbone
108 containing the piggyBac machinery, modified from pJR487 (Weiss et al. 2018) as follows
109 (Figure S1).
110
111 *Preparation of the backbone vector*
112
To allow the use of BbsI as the Type IIS restriction enzyme for Golden Gate cloning of barcodes
into pJR487 (see below), we first removed all three BbsI cut sites from pJR487 by introducing
silent mutations that disrupted the restriction enzyme's recognition pattern. The resulting
plasmid was called pCW328. We next modified pCW328 to make a Golden-Gate-ready vector,
with the final identifier pJC31, by replacing transposon nucleotides with those of a stuffer at a
location 70 nucleotides from the end of the right arm of the transposon (Table S1); see
Supplementary Note and Figure S2 for a description of this choice. The stuffer contained two
BbsI cut sites with custom Type IIS overhang sequences from (Lee et al. 2015), and a NotI cut
site in between the two BbsI cut sites. All cloning steps were carried out by GenScript, Inc.
122
123 *Preparation of barcode oligonucleotides*
124
To make barcodes, we acquired an oligonucleotide pool from IDT that contained random 20 bp
sequences (from hand-mixed random nucleotides) flanked by universal priming regions, U1 and
U2 (Wetmore et al. 2015, Coradetti et al. 2018). These custom oligos were produced and PAGE
purified by IDT. Additionally, we designed forward (FW_BbsI_JC) and reverse (REV_BbsI_JC)
primers which each contained a BbsI cut site, BbsI overhang sequences complementary to the
backbone vector, and either universal priming sequence (Table S2) (Coradetti et al. 2018). We
set up 50 µL amplification PCR reactions with 1 µL of random 20 bp barcodes as template, from
a 2.5 µM stock, and 0.25 µL of each of the forward and reverse primers from a 100 µM stock.
Amplification used Phusion High Fidelity polymerase (NEB) and the following cycling protocol:
98°C for 30 seconds, (98°C for 10 seconds, 58°C for 30 seconds, 72°C for 60 seconds) × 6,
72°C for five minutes. PCR products were purified (Zymo DNA Clean & Concentrator kit) and
then combined. This yielded the final donor barcodes: random 20bp barcodes flanked by
universal priming regions, with BbsI cut sites at the extreme edges.
138

139   *Cloning barcodes into plasmids*
140
141   To clone barcodes into pJC31, we proceeded in two barcoding reactions.
142
The first reaction contained 2:1 molar ratio of vector to barcodes (4 µg of pJC31 and 128 ng of
donor barcodes), 5 µL of 10X T4 Ligase Buffer (ThermoFisher), 2.5 µL of T4 Ligase
(ThermoFisher), 2.5 µL FastDigest BpiI (ThermoFisher), and sterile water up to 50 µL. The
cycling program was: 37°C for five minutes, (37°C for two minutes, 16°C for five minutes) x 25,
65°C for 10 minutes. Then a mixture containing 5 µL 10X FastDigest Buffer (ThermoFisher),
3.13 µL BSA 2 mg/mL (NEB), 12.5 µL FastDigest NotI (ThermoFisher), and 12.5 µL FastDigest
BpiI (ThermoFisher) was spiked into the reaction and incubated at 37°C for 16 hours to digest
unbarcoded backbone vectors. Ten of these reactions were combined, purified, and eluted in
H_2O (Zymo DNA Clean & Concentrator). To spot-check this cloning, 5 µL of this product was
transformed into 25 µL of *E. coli* 10beta electrocompetent cells (NEB). Sanger sequences
across the barcode regions of 20 individually miniprepped *E. coli* colonies showed 95%
barcoding efficiency.
155
The second reaction contained 2:1 molar ratio of vector to donor barcodes (4µg of pJC31 and
128 ng of donor barcodes), 5 µL of 10X T4 Buffer (ThermoFisher), 2.5 µL T4 Ligase
(ThermoFisher), 2.5 µL BpiI (ThermoFisher), and sterile water up to 50 µL. The cycling program
was: 37°C for five minutes, (37°C for two minutes, 16°C for five minutes) x 25, 65°C for 10
minutes. Then a mixture containing 2.5 µL 10X FastDigest Buffer (ThermoFisher), 2.5 µL G
Buffer, (ThermoFisher), 3.13 µL BSA 2 mg/mL (NEB), 12.5 µL FastDigest NotI (ThermoFisher),
and 12.5 µL BpiI (ThermoFisher) was spiked in the reaction and incubated at 37°C for 16 hours
to digest remaining unbarcoded backbone vectors. Six of these reactions were combined,
purified, and eluted in H_2O (Zymo DNA Clean & Concentrator). Then every 5 µL of cleaned
eluted product was redigested with 5µL of NotI-HF (NEB), 5 µL 10X CutSmart buffer (NEB), and
35 µL H_2O at 37°C for 16 hours then 80°C for 20 minutes. The reactions were purified again
(Zymo DNA Clean & Concentrator) and pooled. Spot checks of this cloning reaction proceeded
as above, and Sanger sequences across the barcode regions of 20 individually miniprepped *E.*
coli colonies showed 95% barcoding efficiency.
170
Purified plasmids from the two reactions were combined in a master tube of DNA before
transforming into electrocompetent *E. coli* cells (NEB) to generate the final barcoded piggyBac
pool (final identifier P58). Each electroporation cuvette (BTX) contained 25 µL of 10beta
electrocompetent cells (NEB) and 5 µL of cleaned master tube DNA from the previous golden
gate barcoding step. We performed 21 electroporation reactions in total using the Bio-Rad
GenePulser Xcell machine set to 2.0 kV, 200 Ohms, 25 µF. After electroporation, each culture
was recovered in provided outgrowth media (NEB) by shaking at 37°C at 250 rpm for 1.5 hours.
After recovery, all independent 21 electroporation reactions were combined.
179
The combined recovered transformation *E. coli* culture was used to inoculate two 1L fresh LB
cultures containing carbenicillin at 100 µg/mL to select for *E. coli* cells containing barcoded
piggyBac plasmids. Each culture was incubated for 15.5 hours at 37°C, 250 rpm (overnight) to
expand the barcoded piggyBac *E. coli* pool. Then the two cultures were combined yielding the
final barcoded transposon plasmid pool, P58. This was aliquoted into 1 mL volumes with 15%
glycerol and stored at -80°C.
186
Sequencing verification of barcoded piggyBac pool plasmid DNA for barcode diversity
To verify barcode diversity in the barcoded piggyBac plasmid pool (P58), we sequenced
barcodes as follows. One frozen aliquot of P58 was inoculated into 1.25 L of LB containing

carbenicillin 100 µg/mL and grown for 16 hours 37°C, 250 rpm or until it reached an OD_{600} of
2.1. This culture was gigaprepped on using a column kit (Invitrogen) to generate 5 mg of
plasmid. We used this as input into a PCR with primers (Table S2) annealing to the universal
priming regions flanking the barcode. These primers were dual-indexed, although in this work
we only carried out sequencing of the resulting amplicon from one end (see below), such that
only one index was used. The generic form of the forward primer was
AATGATACGGCGACCACCGAGATCTACACTCTTTCCCTACACGACGCTCTTCCGATCT(N1-
4)xxxxxxGTCGACCTGCAGCGTACG, where the N1-4 represent variable amounts of random
bases from 1-4 to help samples cluster on the Illumina lane and the (x6) represent a unique 6-
bp index sequence for multiplexing samples. The generic reverse primer was
CAAGCAGAAGACGGCATACGAGATxxxxxxGTGACTGGAGTTCAGACGTGTGCTCTTCCGAT
CTGATGTCCACGAGGTCTCT . Four PCR reactions used 50 ng of prepped P58 plasmid
template each. Amplification used Q5 High Fidelity Polymerase (NEB) and a cycling program
98°C for four minutes, (98°C for 30 seconds, 55°C for 30 seconds, 72°C for 30 seconds) x 25,
72°C for five minutes. Each PCR product was purified on a column (Zymo DNA Clean &
Concentrator-5 Kit) and eluted in 10 µL prewarmed 65°C provided elution buffer (Zymo). Six µL
of each were then combined and sequenced off the U2 region via Illumina amplicon
sequencing, on one lane of HiSeq4000 SR50 at the Genomics Sequencing Laboratory at UC
Berkeley. Reads sequenced per library are reported in Table S3. Sequencing of the *E. coli*
vector pool p58 revealed 27,538,142 barcodes with an estimated sequencing error rate of
1.38% analyzed as described (Coradetti et al. 2018).
211
212 **Yeast hemizygote pool construction via barcoded transposon mutagenesis**
213
214 We constructed our yeast hemizygote pool essentially as described (Weiss et al. 2018) but with
215 modifications as follows.
216
217 To prepare plasmid DNA for mutagenesis, one frozen aliquot of P58 was inoculated into 1.25L
218 of LB containing carbenicillin 100 µg ml^{-1} and grown for 16 hours at 37°C, 250 rpm or until it
219 reached an OD_{600}/mL of 2.1. This culture was gigaprepped on using a column kit (Invitrogen) to
220 generate 5 mg of plasmid.
221
Next, we transformed yeast in several, smaller subpools which we combined to form a final pool
as follows. We carried out mutagenesis of CW27, an F1 hybrid from the mating of *S. cerevisiae*
DBVPG1373 with *S. paradoxus* Z1 (Weiss et al. 2018) across the first two days. The first day,
we generated one subpool in a single 50 mL culture and one subpool in five 50 mL cultures at
OD_{600}/mL ~0.9 (~45 OD_{600} units of cells each). The second day, we generated two subpools in
five 50 mL cultures each at OD_{600}/mL ~0.9 (~45 OD_{600} units of cells).
228
To generate subpools consisting of a single 50 mL culture, one colony of CW27 was inoculated
into 5 mL of YPD and incubated at 28°C 200 rpm. 24 hours later, the OD_{600}/mL of the overnight
culture was 3.86. It was backdiluted to an OD_{600}/mL of 0.1 in 50 mL of YPD in a 250 mL
Erlenmeyer flask and grown with shaking at 28°C, 200 rpm for 5.5 hours. After 5.5 hours, it had
reached OD_{600}/mL ~0.9 and cells were at mid-log phase. This 50 mL culture was gently pelleted
at 1000xg for three minutes. The pellet was washed with 25 mL sterile water and then 5 mL
of 0.1 M lithium acetate (Sigma) mixed with 1X Tris-EDTA buffer (10 mM Tris-HCl and 1.0 mM
EDTA); after spin-down, to the tube was added a solution of 0.269 mg of P58 mixed 5:1 by
volume with salmon sperm DNA (Invitrogen), followed by 3 mL of 39.52% polyethylene glycol,
0.12 M lithium acetate and 1.2X Tris-EDTA buffer (12 mM Tris-HCl and 1.2 mM EDTA). The
tube was rested for 10 minutes at room temperature, then heat-shocked in a water bath at 37°C
for 26 minutes. The tube was gently spun at 1000g for three minutes after which supernatant

was removed. We transferred the cells to a flask and added YPD to attain an OD_{600}/mL of
~0.35–4 in ~70 mL. Each such culture was recovered by shaking at 28°C and 200 rpm for two
hours. G418 (Geneticin; Gibco) was added to each at a concentration of 300 µg/mL to select for
those cells that had taken up the plasmid, and cultures were incubated with 200 rpm shaking at
28°C for two days until each reached an OD_{600}/mL of ~2.5. We transferred cells from this
culture, and YPD + G418 (300 µg/mL), to new 250 mL flasks at the volumes required to attain
an OD_{600}/mL of 0.2 in 50 mL each. We cultured each flask with 200 rpm. shaking at 28°C
overnight until each reached an OD_{600}/mL of 3.43. To cure transformants of the P58 URA3+
plasmid, we spun down 10% of this master culture, and resuspended in water with the volume
required to attain a cell density of 1.85 OD_{600}/mL. Four mL of this resuspension were plated (1
mL per 24.1 cm x 24.1 cm plate) onto plates containing complete synthetic media with 5-
fluorooritic acid (0.2% dropout amino acid mix without uracil or yeast nitrogen base (US
Biological), 0.005% uracil (Sigma), 2% D-glucose (Sigma), 0.67% yeast nitrogen base without
amino acids (Difco), 0.075% 5-fluorooritic acid (Zymo Research)). After incubation at 28°C to
enable colony growth, colonies were scraped off all four plates and combined into water at the
volume required to attain 44 OD_{600}/mL, yielding the transposon mutant hemizygote subpool.
This was aliquoted into 1 mL volumes with 10% dimethylsulfoxide and frozen at −80°C.
258
To generate subpools consisting of five 50 mL cultures, one colony of CW27 was inoculated to
100 mL of YPD in a 250 mL Erlenmeyer flask and incubated shaking at 28°C, 200 rpm. Twenty-
four hours later, the OD_{600}/mL of the overnight culture was OD_{600}/mL 3.89. The overnight culture
was backdiluted to OD_{600}/mL 0.1 in 250 mL of YPD and incubated for 5.5 hours at 28°C, 200
rpm. After 5.5 hours, the OD_{600}/mL reached 0.9 and cells were split into five 50 mL conical
tubes, and subjected each to heat shock as above. We then transferred all cells from this post-
transformation culture to one 1L flask and added fresh YPD to attain OD_{600}/mL 0.4 in ~750 mL
YPD. The transformed culture was recovered by shaking at 28°C, 200rpm, for two hours. G418
(300mg/ul) was added to select for the transposed cells. The culture continued shaking for 48
hours or until the OD_{600}/mL reached 2.1. This culture was then backdiluted to create a new
culture at OD_{600}/mL 0.2 in 500 mL of YPD with 300mg/µL G418 shaking for 24 hours at 28°C,
200 rpm until it reached OD_{600}/mL ~3.4 The curing, scraping, and freezing steps were the same
as above.
272
To combine the four subpools to yield the final 160X hemizygote pool (final identifier P75), three
1 mL aliquots of each subpool were thawed on ice for one hour. They were transferred to each
of four 1L flasks with 500 mL YPD to OD_{600}/mL 0.2, cultured at 28°C, 200 rpm for 17 hours upon
which the OD_{600}/mL was 3.5-4. They were gently pelleted, combined, and resuspended in two
ways to reach OD_{600}/mL of 44: YPD with 15% glycerol and YPD with 7% DMSO, aliquoted to 1
mL volumes, and frozen at -80°C.
279
280 **Tn-seq mapping of yeast hemizygote pool**
281
282 *Tn-seq library preparation*
283
To associate barcoded transposon insertions to genomic location in the hemizygote pool, which
we refer to as Tn-seq, we first sequenced barcoded transposon insertions according to the
methods of (Weiss et al. 2018) as follows. Each 44 OD_{600}/mL aliquot of each subpool or final
pool was thawed on ice, and its genomic DNA (gDNA) was harvested with the ZR
Fungal/Bacterial DNA MiniPrep Kit (Zymo Research). gDNA was resuspended in DNA elution
buffer (Zymo Research) prewarmed to 65°C, and its concentration was quantified using a Qubit
4.0 fluorometer. Illumina transposon sequencing (Tn-seq) library construction was as described
previously. Briefly, gDNA was sonicated and ligated with common adapters, and for each

fragment deriving from a barcoded transposon insertion in the genome, a sequence containing
a barcode, a portion of the transposon, and a portion of its genomic context (the barcoded
transposon–genome junction) was amplified using one primer homologous to the U1 region
immediately upstream of barcode and another primer homologous to a region in the adapter.
See Table S2 for the transposon-specific primer ("forward primer"), where Ns represent random
nucleotides, and the indexed adapter-specific primer ("reverse primer"). Amplification used
Jumpstart polymerase (Sigma) and the following cycling protocol: 94°C for two minutes, (94°C
for 30 seconds, 65°C for 20 seconds, 72°C for 30 seconds) × 25, 72°C for 10 minutes.
Sequencing of paired-end reads of 150 bp was done over two lanes on a HiSeq4000 at
Novogene Corporation (Sacramento, CA) and one lane on a NovaSeq SP at the Genomics
Sequencing Laboratory at UC Berkeley (Berkeley, CA). Reads sequenced per library are
reported in Table S4.
304
305 *Tn-seq data analysis*
306
Tn-seq data of the hemizygote pool was analyzed, to infer transposon insertions on the basis of
barcodes detected in reads as junctions with genomic sequence, essentially as described
(Coradetti et al. 2018) (https://github.com/stcoradetti/RBseq/tree/master/Old_Versions/1.1.4),
with the following modifications. For each barcode, instead of scanning positions for the end of
the insertion from a sequence specified by a model file, we searched for the final 22 base pairs
of the right arm of the piggyBac transposon allowing for two mismatches. For annotation, we
converted the annotation file from https://github.com/weiss19/rh-seq for the *S.* cerevisiae D1373
x *S. paradoxus* Z1 hybrid to a compliant GFF3 file using Another GFF Analysis Toolkit (AGAT) -
Version: v0.4.0 (https://github.com/NBISweden/AGAT). Then, we used a custom Jupyter
notebook to annotate the file generated by the RBseq mapping software.
317
Quality control for Tn-seq, to eliminate barcodes whose junction genomic sequence mapped to
multiple insertion locations in the hybrid genome, and to minimize the proportion of sequencing
errors included in final tallies, was as described (Coradetti et al. 2018). Briefly, we eliminated
from further consideration any case where a barcode observed in Tn-seq sequencing data
differed from another, much more abundant, barcode by a single base (a total of 2,024,812 off-
by-one barcodes in 2,888,129 reads). We also filtered out off-by-two barcodes (280,949
barcodes in total). Separately, we eliminated barcodes that were detected in sequencing data
as a junction with more than one genomic context, suggesting the respective transposon had
inserted into multiple locations in one or many clones (98,669 barcodes where this inference
was based on multiple strong mapping matches, and an additional 46,583 barcodes where this
inference was ambiguous, with one strong mapping match with reads outnumbered by those
assigned to weaker mapping matches). The final filtered barcode set comprised 548,129
uniquely barcoded and mapped inferred transposon insertions in the P75 hemizygote pool, at
an average read depth of 308.6 reads, and a median read depth of 47 reads; 166,834 of these
insertions were mapped as genic. The annotation script, GFF3 file, and modified mapping script
are available at https://github.com/melanieabrams-pub/RH-seq_with_barcoding.
334
335 **Competition cultures**
336
For the thermotolerance competition at 37°C (Table S5), one aliquot of the yeast hemizygote
pool was thawed and inoculated into 150 mL of YPD in a 250 mL unbaffled Erlenmeyer flask
and grown for six hours at 28°C, 200 rpm. This pre-culture (T_0, at OD_{600}/mL of 1.22) was
backdiluted into 12 10 mL competition cultures at 200 rpm at each of 28°C and 37°C, with a
starting OD_{600}/mL of 0.02 or 0.05 in at 28°C and 37°C respectively. These competition cultures
were maintained within logarithmic growth through back-dilutions into fresh tubes of 10 mL of

YPD at the same optical density as the starting culture, for a total of 10-15 generations.
Dilutions for the 28°C competition cultures were performed after 8.5, 18.5, and 25.5 hours after
the T_0 timepoint, and dilutions for the 37°C competition cultures were performed after 8.5, 18.5,
25.5 hours and 32.5 hours after the T_0 timepoint. The entire cell culture was harvested from
each of these biological replicate tubes for sequencing as biological replicates.
348
For thermotolerance competition at 36°C (Table S6), competition cultures were grown as above
with the following differences. The high temperature was 36°C, instead of 37°C. The pre-culture
(T_0, at OD_{600}/mL of 0.693 after 5.5 hours at 28°C, 200 rpm) was backdiluted to a starting
OD_{600}/mL of 0.02 for competition cultures at 36°C. Dilutions for both the 28°C and 36°C
competition cultures were performed after 8.5, 18.5 and 25.25 hours after the T_0 timepoint.
Eleven instead of 12 replicates were carried out at 28°C.
355
356   **Barcode quantification from competition cultures**
357
358   *Bar-seq library preparation*
359
To determine the abundance of barcoded transposon mutant hemizygote clones after selection,
we sequenced barcodes insertions as follows. Each cell pellet from a selection sample was
thawed on ice, and its genomic DNA (gDNA) was harvested with the Zymo QuickDNA Kit
(Zymo#D6005). gDNA was resuspending in DNA elution buffer (Zymo Research) prewarmed to
65°C, and its concentration was quantified using a Qubit 4.0 fluorometer. The barcode insertion
was amplified as above (see *Sequencing verification of barcoded piggyBac pool plasmid DNA*
for barcode diversity). Each PCR product was purified on a column (Zymo DNA Clean &
Concentrator) and eluted in 10 µL prewarmed 65°C provided elution buffer (Zymo). Six µL of
each were then combined and sequenced off the U2 region by Illumina sequencing on one lane
of HiSeq4000 SR50 at the QB3 Genomics Sequencing Laboratory at UC Berkeley.
370
371   *Bar-seq data analysis*
372
Bar-seq mapping and quantification were as described (Coradetti et al. 2018)
(https://github.com/stcoradetti/RBseq/tree/master/Old_Versions/1.1.4), wherein only barcodes
that passed quality control in Tn-seq (see *Tn-seq data analysis* above) were analyzed for
quantitative measures of abundance via Bar-seq. Thus we did not use in our screen any
barcode that was detected in Bar-seq sequence data but not Tn-seq data (the product of *e.g.*
sequencing errors in Bar-seq, or a failure to observe in Tn-seq a barcode associated with a
bona fide transposon insertion that could be detected in Bar-seq). A total of 301,349 barcodes
conformed to these criteria from across all replicates of Bar-seq in competitions for the
dissection of determinants of growth at 37°C relative to 28°C, with an average read depth of
305.3 reads and a median of 12 reads; 89,772 of these Bar-seq detected barcodes
corresponded to inferred transposon insertions in genes and were analyzed as input to the
reciprocal hemizygosity testing pipeline described below. In a given replicate competition culture
we detected a median 1×10^5 barcodes. The latter represented a fifth of the size of the total
pool of hemizygotes detectable after quality control by Tn-seq (5.5×10^5; see *Tn-seq data*
analysis above). Thus, the extent of bottlenecking in any given competition experiment was
modest, with diversity retained at the order of magnitude of the mutant pool size.
389
390   Competitions for the dissection of growth at 36°C relative to 28°C (Table S6) used the same
391   procedures as above, mapping a total of 230,469 barcodes, 68,523 of which corresponded to
392   inserts in genes and were analyzed as input to the reciprocal hemizygosity testing pipeline

393	described below. In a given replicate competition culture, we detected a median 5×10^4
394	barcodes.
395	

Reciprocal hemizygosity testing

398	The tabulated counts of abundance from Bar-seq for each barcode in each replicate were used
399	as input into reciprocal hemizygosity tests essentially as in (Abrams et al. 2021), with slight
400	changes as follows. We had in hand each barcode which had been sequenced as a junction
401	with a unique genomic location in the Tn-seq step and had passed quality control there (see *Tn-*
402	*seq data analysis* above), and which was now detected in competition cultures. We interpreted
403	each such barcode as reporting a hemizygote clone bearing a transposon insertion at the
404	respective position of the respective species' allele (*S. cerevisiae* or *S. paradoxus*), with the
405	other species' allele retained as wild-type at that locus. In what follows, we refer to each such
406	barcode as reporting an inferred hemizygote clone, with respect to its growth behavior in
407	competition cultures. As in (Abrams et al. 2021), for a given biological replicate we normalized
408	the abundances attributed to each inferred hemizygote genotype to the total number of
409	sequencing reads in the respective sample, and we eliminated from further analysis insertions
410	which had been annotated as intergenic, or as corresponding to the plasmid used to generate
411	this library. For reciprocal hemizygosity tests, we excluded from consideration any gene with
412	fewer than three inferred hemizygote genotypes per allele. Of the retained genes, for each
413	inferred hemizygote genotype, we tabulated the quantity $a_{experimental,i}$, the sequencing-based
414	abundance measured after the competition culture in biological replicate i of growth at the
415	experimental temperature (36 or 37°C), and, separately, we calculated $a_{control,i}$, the analogous
416	quantity from growth at the control temperature (28°C), for $i = [1,12]$. We then took the mean of
417	the latter and used it to tabulate the temperature effect on the inferred hemizygote genotype in
418	replicate i, $t_i = \log_2(a_{experimental,i}/a_{control,mean})$. As in (Abrams et al. 2021), we eliminated an inferred
419	hemizygote genotype if the coefficient of variation of this quantity exceeded 2.0, or there were
420	fewer than 1.1 normalized reads. With the data for the remaining inferred hemizygote genotypes
421	(Tables S5 and S6), for a given gene, we compiled the vector of the t measurements across
422	replicates and all inferred hemizygote genotypes with each species' allele of the hybrid
423	disrupted in turn, and discarded genes where the coefficient of variation of the t measurements
424	across hemizygote inserts for one or both alleles exceeded 10. For the remainder, we used the
425	Mann-Whitney test to compare these two vectors, with Benjamini-Hochberg correction for
426	multiple testing (Tables S7 and S8). For a given gene, we calculated the effect size as the
427	difference between two values: the $\log_2$(abundance at the experimental temperature/abundance
428	at 28°C) of the average inferred hemizygote genotype representing a transposon insertion in the
429	*S. cerevisiae* allele, and the analogous quantity among inferred hemizygote genotypes
430	representing insertions in the *S. paradoxus* allele of the gene. Scripts for this modified RH-seq
431	analysis pipeline are available at https://github.com/melanieabrams-pub/RH-
432	seq_with_barcoding. We earmarked top candidate genes for factors contributing to the
433	thermotolerance of *S. cerevisiae* as those with corrected Mann-Whitney $p < 0.05$ in the
434	reciprocal hemizygosity test, and an effect size < -0.5, *i.e.* disrupting the *S. cerevisiae* allele was
435	associated with a strong defect in thermotolerance relative to disruption of the *S. paradoxus*
436	allele; we refer to this gene set as our top barcoded RH-seq hit gene list.

Analysis of inferred interactions between top hit genes from barcoded RH-seq

440	For the circos plot reporting inferred interactions between top hit genes from barcoded RH-seq,
441	we used the STRING database (Szklarczyk et al. 2021), accessed September 30, 2021, which
442	incorporates experimental/biochemical data from DIP, BioGRID, HPRD, IntAct, MINT, and PDB,
443	and curated data from Biocarta, BioCyc, Gene Ontology, KEGG, and Reactome. Widths of

444   edges between nodes in the circos plot represent STRING confidence scores, each the
445   probability of a true positive interaction between a given two genes (Szklarczyk et al. 2021).
446
To test the encoded proteins of top barcoded RH-seq hit genes for enrichment of physical
interactions with each other, we used curated known interactions from BioGRID (Oughtred et al.
2021) as housed in the Saccharomyces Gene Database, downloaded February 19, 2021. We
tabulated the number of physical interactions between the proteins encoded by RH-seq hit
genes, and we divided that by the total number of interactions involving one RH-seq hit gene
and any other gene in the genome; call this ratio r_{true}. Then, we drew a random sample of genes
from the genome, as described above for GO term resampling. We tabulated, in this random
gene set, the number of physical interactions between genes in that sample, and we divided
that by the total number of interactions involving one gene in the random sample and any other
gene in the genome, to yield $r_{resample}$. We repeated this procedure 10,000 times, and we used
the proportion of resampled groups where $r_{resample}$ was greater than or equal to r_{true} as a one-
sided p value assessing the significance of enrichment of interactions between our genes of
interest.
460
461   **Gene Ontology analyses of top hit genes from barcoded RH-seq**
462
To test top barcoded RH-seq hit genes for enrichment for overrepresentation of a particular
Gene Ontology (GO) term, we mapped each gene to its Gene Ontology groups based on data
from geneontology.com (Ashburner et al. 2000). We filtered out GO terms with fewer than five
or with more than 200 gene members. We also filtered out GO terms with identical membership
in the genome. We took the subset of the remaining GO terms with at least one member among
our top barcoded RH-seq hit genes. Then, we randomly sampled genes from the genome,
ensuring the same proportion of essential genes as in our set of top barcoded RH-seq hit genes
based on the essentiality annotations of (Winzeler 1999). We tabulated whether our random
sample had greater or fewer genes with that GO term than our candidate set. We repeated this
procedure 10,000 times and used the proportion of these resampled groups that had more
genes in the given GO term as the initial p value assessing the significance of the enrichment of
that GO term. Then, we applied Benjamini-Hochberg correction for multiple hypothesis testing to
generate final, adjusted p-values for the enrichment of the given GO term among top barcoded
RH-seq hit genes.
477
To test Biological Process ontologies for enrichment for large magnitudes of the effect of allelic
variation on thermotolerance, we used the latter as tabulated in **Reciprocal hemizygosity**
testing above. We filtered GO terms as above, and then excluded all genes absent in our
barcoded RH seq analysis, which would have no associated quantity for the effect of allelic
variation to resample. For each retained term in turn, we first tabulated the median absolute
value of the effect size of the gene members for which we had data, e_{true}. Then, we tabulated
the analogous quantity for a random sample of the same number of genes from the genome,
$e_{resample}$, ensuring the same proportion of essential genes as above. We repeated this procedure
100 times, and used the proportion of the resampled groups for which $e_{resample}$ was greater than
or equal to e_{true} as an initial p value assessing the enrichment of large effects of allelic variation
in the genes the term. For all GO terms with an initial p value < 0.1, we repeated this procedure
10,000 times to calculate a more precise p value. Then, we applied the Benjamini-Hochberg
correction for multiple hypothesis testing to generate final, adjusted p-values for the enrichment
of the given GO term for large effects of allelic variation on thermotolerance.
492
Molecular evolution analysis of RH-seq hit genes

Branch length PAML analysis with codeML was performed as in (Dubin et al. 2020). Hits were
manually inspected for the quality of the alignment, and one, *YAL026C*, was discarded for poor
alignment quality leading to an artifactually high branch length. We used the inferred branch
lengths as input into a resampling test as in **Gene Ontology analyses of top hit genes from**
barcoded RH-seq above, and we performed a one-sided significance test for long branch
lengths along the *S. cerevisiae* lineage. Branch-site PAML analysis with codeML was performed
as in (Abrams et al. 2021). Jalview version 2 was used to visualize the percent identity of amino
acid sequence alignments (Waterhouse et al. 2009). McDonald-Kreitman analysis statistics
were calculated as in (Abrams et al. 2021). Fisher's exact test was used to compute *p*-values for
individual loci, and these were adjusted using the Benjamini-Hochberg correction for multiple
hypothesis testing.
505
506 **Data availability statement**
507
508 Sequencing data are deposited in the Sequence Read Archive under the accession
509 PRJNA735401. Strains and plasmids are available upon request. Custom scripts for the
510 barcoded RH-seq analysis are available at https://github.com/melanieabrams-pub/RH-
511 seq_with_barcoding. The authors affirm that all data necessary for confirming the conclusions of
512 the article are present within the article, figures, and tables.
513
514 <div align="center">**RESULTS**</div>
515
516 **Dissecting thermotolerance divergence between species by barcoded transposon**
517 **mutagenesis**
518
With the goals of boosting RH-seq throughput and power, and achieving new insights into the
genetics and evolution of yeast thermotolerance, we set out to generate an RH-seq reagent for
yeast incorporating barcoded transposons (Wetmore et al. 2015). For this purpose, we first
generated a pool of plasmids, each encoding a barcoded copy of the piggyBac transposon and
its transposase (Figure S1A-C). To use these in RH-seq, we revisited our previously
characterized model system: a comparison between DVBPG1373, a thermotolerant Dutch soil
strain of *S. cerevisiae,* and Z1, an *S. paradoxus* isolate from the UK (Weiss et al. 2018; AlZaben
et al. 2021; Abrams et al. 2021). The F1 hybrid formed from the mating of these strains exhibits
a thermotolerance phenotype intermediate between those of the two species parents, and thus
is well-suited to mapping of allelic effects on the trait (Weiss et al. 2018). We transformed this
F1 hybrid with barcoded plasmids, yielding a pool of hemizygote mutants, which we expanded
and then banked (Figure S1D). Next, to catalog the genomic locations of transposon insertions,
we used the DNA from a culture of the pool in standard conditions as input into a first round of
sequencing library construction, whose primers recognized a common site on the transposon
and a common DNA adapter ligated to DNA fragment ends ("Tn-seq"; Figure 1A). Sequencing
and data analysis, with quality controls to eliminate barcodes that could not be uniquely
associated with a single transposon insertion location (see Methods), yielded a catalog of
548,129 barcoded hemizygotes in the pool whose genomic insertion locations were tabulated.
At this point we could harness the pool for highly replicated screens, each of which could
quantify hemizygote abundance in a condition of interest via relatively cheap and
straightforward barcode sequencing ("Bar-seq"; Figure 1B).
540
Thus, with our barcoded hemizygote pool, we implemented an RH-seq screen to search for
genes at which *S. cerevisiae* and *S. paradoxus* alleles drove differences in strain abundance at
high temperature. For this, we subjected the pool to growth assays with 12 biological replicate
cultures at 37°C, alongside controls at 28°C. We used DNA from each culture as input into

barcode sequencing (Figure 1B). The resulting data revealed a total of 301,349 cases where a
barcode, representing a hemizygote clone with a transposon insertion catalogued by Tn-seq
(Figure 1A), was detectable in our growth assays. Transposon insertion positions corresponding
to these informative barcodes were evenly split between *S. cerevisiae* and *S. paradoxus* alleles
of genes throughout the F1 hybrid genome (Figure S3). We took the normalized count of a
given barcode in a sequencing data set as a report of the fitness of the respective hemizygote,
i.e. its relative abundance after growth in the pool in the respective condition. We then used the
complete set of such counts as the input into reciprocal hemizygosity tests to compare, for a
given gene, the temperature-dependent abundance of strains harboring a disruption in the *S.*
cerevisiae allele, relative to that of strains with the *S. paradoxus* allele disrupted. A pipeline for
these tests, including filters for coverage and reproducibility and multiple testing correction (see
Methods), revealed 83 genes at a 5% false discovery rate (Figure 2 and Table S7). This
contrasted with the much smaller set of eight genes at which species' alleles drove differences
in high-temperature growth, in our original non-barcoded RH-seq approach (Weiss et al. 2018),
which had involved only three biological replicates. The 10-fold increase in the number of
significant hits in our barcoded RH-seq screen reflects the statistical power afforded by our
highly-replicated method to detect even quite small effects.
562
In our barcoded RH-seq screen hits, as a positive control we first examined the set of genes
known to contribute to thermotolerance divergence from our earlier study (*AFG2, APC1, CEP3,*
DYN1, ESP1, MYO1, SCC2, and *DYN1*) (Weiss et al. 2018). Several did not meet the
experiment-wide statistical thresholds of our barcoded RH-seq pipeline (Figure S4A),
suggesting an appreciable false negative rate of the latter overall. However, manual inspection
made clear that hemizygosity effects at all gold-standard thermotolerance loci were borne out: in
each case, in barcoded RH-seq data, strains with disruptions in the *S. cerevisiae* allele, and a
wild-type copy of the *S. paradoxus* allele, had worse thermotolerance than did strains with only
the *S. cerevisiae* allele intact (Figure S4A-B), as we had previously reported (Weiss et al. 2018).
Furthermore, the list of gene hits from barcoded RH-seq also included *HFA1* (Figure 2B and
Tables S7 and S9) which was reported and validated separately as a determinant of
thermotolerance differences between yeast species (Li et al. 2019). On the strength of these
controls, we considered our deep sampling of thermotolerance loci to serve as a useful proof of
concept for the barcoded RH-seq method.
577
578 **Functional-genomic analysis of thermotolerance genes**
579
We next aimed to pursue deeper analyses of the novel gene hits from barcoded RH-seq in our
yeast thermotolerance application. We considered that a focus on the strongest and most
evolutionarily relevant sources of mapping signal would likely yield the most informative results.
As such, in light of our interest in explaining the exceptional thermotolerance of purebred *S.*
cerevisiae, we earmarked the 44 genes from our larger candidate set at which the *S. cerevisiae*
allele boosted the trait most dramatically relative to that of *S. paradoxus* (Figure 2 and Table
S9). In what follows, we refer to these genes as our top RH-seq hits, and we analyze them as
our highest-confidence predictions for factors that nature would have used in evolving the *S.*
cerevisiae phenotype.
589
We sought to use our mapped loci to explore potential functional mechanisms underlying the
thermotolerance trait. We hypothesized that *S. cerevisiae* thermotolerance genes could
participate in an interacting network, jointly shoring up particular aspects of cell machinery that
were critical for growth at high temperature (AlZaben et al. 2021). Consistent with this notion,
the STRING database, which collates experimentally detected protein-protein interactions,
genetic interactions, and pathway membership (Szklarczyk et al. 2021), inferred multiple

interactions among our top genes from barcoded RH-seq, with salient signal involving cell cycle
factors (Figure 3). A more focused analysis revealed an enrichment, among our top barcoded
RH-seq hits, for protein-protein interactions with one another as tabulated in BioGRID (Oughtred
et al. 2021), to an extent beyond the null expectation (resampling p = 0.014). We also
implemented qualitative gene set enrichment tests, which revealed that chromosome
segregation and mitosis factors, although relatively few in number among our top barcoded RH-
seq hit loci, were significantly enriched relative to the genomic null (Table 1). And we developed
a complementary, quantitative test to screen Gene Ontology terms for large allelic effect size
(the impact on thermotolerance when the *S. cerevisiae* allele of a given gene was disrupted in
the hybrid, as a difference from the analogous quantity for the *S. paradoxus* allele; see
Methods). Top-scoring in this test was a mitosis gene group, encoding components of the septin
ring (GO:0000921; resampling p < 0.0001). Together, these results suggest that our top
thermotolerance gene hits share commonalities in function, most notably involving cell cycle
factors. This dovetails with previous phenotypic and genetic characterization of yeast
thermotolerance, including the breakdown of cell division in heat-treated *S. paradoxus* (Weiss et
al. 2018), and supports a model in which *S. cerevisiae* acquired thermotolerance in part by
resolving the latter cell cycle defect.

The genetics of yeast thermotolerance likely also involves mechanisms beside mitosis, given
the known role of mitochondrial genes (Baker et al. 2019; Li et al. 2019) and those operating
during stationary phase (AlZaben et al. 2021). Indeed, functional-genomic tests revealed
enrichment for secretion genes and for regulatory factors in our top RH-seq hits, although no
such group constituted a large proportion of the total hit list (Table 1). Annotations in
transcription and translation, mitochondrial function, and signaling were also apparent in our top
thermotolerance loci (Figure 2B). These trends are consistent with a scenario in which evolution
built the trait in *S. cerevisiae* by tweaking an array of housekeeping mechanisms, beside those
that involve cell cycle machinery.

Evolutionary analysis of thermotolerance genes

We anticipated that sequence analyses of the genes we had mapped to thermotolerance by
barcoded RH-seq could shed light on the evolutionary history of the trait. To explore this, we
used a phylogenetic approach in *Saccharomyces sensu stricto*. We first inferred species-
specific branch lengths in the phylogeny of each gene in turn, and focused on the lineage
leading to *S. cerevisiae*. The distribution of branch lengths along this lineage among top
thermotolerance gene hits was not detectably different from that of the genome as a whole, with
the exception of two rapidly evolving thermotolerance genes, *TAF2* and *BUL1*, encoding a
transcription initiation factor and ubiquitin ligase adapter respectively (Figure S5). Separately,
we quantified protein evolutionary rates in top hits from barcoded RH-seq. A branch-site
phylogenetic modeling approach (Yang 2007) detected striking evidence for positive selection
along the *S. cerevisiae* lineage in the amino acid permease *GNP1*, the kinetochore DNA binding
factor *CBF2*, and the sister chromatid cohesion factor *CTF18* (Figure 4). Interestingly, however,
McDonald-Kreitman tests (McDonald and Kreitman 1991) on population-genomic data did not
detect an overall excess of amino acid variation relative to synonymous changes, at these three
genes or any other barcoded RH-seq hit locus (Table S10). Thus, even at genes harboring
individual codons with likely signatures of selection, we could not detect evidence for a scenario
where *S. cerevisiae* stacked up a large number of unique amino acid changes, in the evolution
of thermotolerance. Together, however, our analyses do highlight thermotolerance genes with
marked signal for derived alleles in *S. cerevisiae* at single codons or in the overall DNA
sequence—cases where species divergence is likely to be of phenotypic and evolutionary
importance.

647
648 **DISCUSSION**
649
650 **RH-seq power and the interpretation of mapped loci**
651
652 In this work, we established the barcoded RH-seq method for genetic dissection of trait variation
653 between diverged lineages. RH-seq falls into a family of recently-developed methods that can
654 dissect natural trait variation across species barriers (Weiss and Brem 2019). A chief distinction
655 of RH-seq is its low cost and low overhead, and the barcoding feature we add here cuts down
656 labor and cost even further, enabling high replication.
657
Our application to yeast thermotolerance serves as an informative model for the performance of
barcoded RH-seq on highly genetically complex traits. We pinpointed dozens of candidate
genes at which species-level variation contributes to growth at high temperature. And yet we
also observed evidence for a sizeable false negative rate among our barcoded RH-seq results,
since some validated thermotolerance loci from our earlier screen did not appear among the hits
here. Likewise, a separate barcoded RH-seq mapping of yeast species' differences in growth
under milder heat stress revealed little signal above the noise (Table S6 and Table S8), likely
reflecting very weak genetic effects under this condition. We thus expect that, as would be true
for a classical linkage or association scan, the statistical power of a barcoded RH-seq
experiment is a function of signal to-noise, genetic complexity, and genetic effect size; and that
many thermotolerance loci remain to be identified even in our very deep set of screen results
from high-temperature growth.
670
By virtue of our focus on pro-thermotolerance alleles in *S. cerevisiae*, our work has left open the
functional and evolutionary genomics of loci at which the allele from *S. cerevisiae* instead
conferred worse thermotolerance than that of *S. paradoxus*, when each in turn was uncovered
in the hybrid. Our barcoded RH-seq identified a number of such genes at high statistical
significance. These loci may well reflect the accumulation of advantageous alleles in *S.*
paradoxus, or deleterious alleles in *S. cerevisiae*, by drift, even as the latter was under selection
to improve the trait in evolutionary history. Analogously, in linkage mapping results, the effect of
an allele in recombinant progeny from a cross often does not conform to that expected from the
respective parent's phenotype (Burke and Arnold 2001; Brem and Kruglyak 2005). It is also
possible that some such allelic effects are the product of epistatic interactions between a locus
of interest and the hybrid background, and would be phenotypically buffered (and thus
evolutionarily irrelevant) in the purebred species. Molecular validation will be necessary to
confirm the phenotypic impact of variation at our mapped loci, and its potential dependence on
genetic background.
685
That said, we consider genes with pro-thermotolerance *S. cerevisiae* alleles according to
barcoded RH-seq to be strong candidates for *bona fide* determinants of the trait from the wild in
this species. Indeed, earlier work has shown that for such genes mapped by RH-seq in the
hybrid, the advantage of *S. cerevisiae* alleles is borne out in tests in purebred backgrounds
(Weiss et al. 2018). Accordingly, we have shown here that as a cohort, barcoded RH-seq hits
with advantageous *S. cerevisiae* alleles exhibit functional and sequence-based attributes
consistent with a role in thermotolerance evolution in the wild.
693
694 **Cellular and molecular mechanisms of thermotolerance**
695
Our top RH-seq hits revealed strong evidence for chromosome segregation and other mitosis
functions as a linchpin of *S. cerevisiae* thermotolerance. As a complement to earlier

characterization of six such genes (*APC1, ESP1, DYN1, MYO1, CEP3,* and *SCC2*) (Weiss et al.
2018; Abrams et al. 2021), we now report seven new thermotolerance determinants that
function in cell division (*MEC1, MLH1, CTF13, CTF18, MCM21, CBF2,* and *MYO2*). The
emerging picture is one in which the ancestor of modern-day *S. cerevisiae*, faced with
dysfunction of a slew of mitotic factors at high temperature, acquired variants across the
genome to shore up their activity under these conditions. Under one model of *S. cerevisiae*
evolution, the particular niche to which this species specialized was one of avid fermentation,
producing (and resisting) heat and ethanol at levels that eliminated its microbial competitors
(Goddard 2008; Salvadó et al. 2011). In such a scenario, the maximum benefit could well
accrue to the organism if it were able to undergo rapid cell division under the challenging
conditions of its own making. Consistent with this notion, another budding yeast,
Hanseniaspora, which often dominates in early fermentation prior to takeover by *S. cerevisiae*
(Fleet 2003), underwent evolutionary loss of much of the cell-cycle checkpoint machinery,
consistent with a strategy of accelerated growth at any cost to outcompete other species at the
respective stage (Steenwyk et al. 2019).
713
However, since our current hit list includes many genes from other housekeeping pathways,
from transcription/translation to transport and lipid metabolism, mitosis does not appear to be
the whole mechanistic story for the thermotolerance trait in *S. cerevisiae*. Indeed, other
housekeeping factors also showed up in our previous screen (Weiss et al. 2018) and in an
elegant complementary study of mitochondrial determinants of thermotolerance divergence
between yeast species (Baker et al. 2019; Li et al. 2019). The panoply of functions detected
among our mapped loci conforms well to current models of the mechanisms of thermotolerance,
which invoke many essential genes and housekeeping processes (Leuenberger et al. 2017).
722
The latter idea emerged largely from a proteomic study which showed that thermotolerant
organisms had higher thermostability of essential proteins of many functions, across the tree of
life (Leuenberger et al. 2017). Were sequence changes that led to improved protein stability a
linchpin of thermotolerance evolution in *S. cerevisiae*? Our data are consistent with a
mechanistic role for properties of the protein sequences of many thermotolerance genes, in that
variation in coding regions has come to the fore in our sequence tests here and those of an
earlier small-scale analysis (Abrams et al. 2021). And interestingly, an experimental case study
of one of our mapped thermotolerance loci revealed no impact on the trait from variation in the
promoter, only from that in the coding region (Abrams et al. 2021). We cannot rule out
noncoding determinants in some cases, especially given that a few hundred genes exhibit
temperature-dependent *cis*-regulatory programs unique to *S. cerevisiae* (Tirosh et al. 2009; Li
and Fay 2017). But if coding regions do hold the exclusive key to the mechanism of *S.
cerevisiae* thermotolerance, they could well involve variants that improve protein function and
regulation alongside folding/structure at high temperature. Overall, then, we envision that nature
could have used a variety of molecular mechanisms in building the trait, given the apparent
complexity of the problem. Biochemical studies will be necessary to nail down exactly how *S.
cerevisiae* alleles advance thermotolerance.
740
In summary, our data reveal a newly detailed picture of the highly polygenic architecture for a
natural trait divergence between species. It is tempting to speculate that evolution may draw on
a vast number of variants across the genome to refine a trait over millions of generations,
making effects stronger, weaker, or less pleiotropic, adding regulatory control, and so on (Orr
1998). If so, these architectures may ultimately conform to the omnigenic model (Boyle et al.
2017)–which was originally applied to human disease genetics, but may also prove to be an apt
description of ancient adaptations.
748

ACKNOWLEDGEMENTS

This work was supported by NSF GRFP DGE 1752814 to M.A. and NIH R01 GM120430 to R.B.B. The authors thank Adam Arkin for his generosity with computational resources; Morgan Price, Lori Huberman, and members of the John Dueber lab for discussions about barcoded transposon mutagenesis; and sequencing; and Abel Duarte for cloning advice.

REFERENCES

Abrams MB, Dubin CA, AlZaben F, Bravo J, Joubert PM, Weiss CV, Brem RB. 2021. Population and comparative genetics of thermotolerance divergence between yeast species. G3 Genes|Genomes|Genetics 11(7):jkab139. doi: 10.1093/g3journal/jkab139.

AlZaben F, Chuong JN, Abrams MB, Brem RB. 2021. Joint effects of genes underlying a temperature specialization tradeoff in yeast. PLoS Genet. 17(9):e1009793. doi:10.1371/journal.pgen.1009793.

Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, Davis AP, Dolinski K, Dwight SS, Eppig JT, et al. 2000. Gene Ontology: tool for the unification of biology. Nat Genet. 25(1):25–29. doi:10.1038/75556.

Baker EP, Peris D, Moriarty RV, Li XC, Fay JC, Hittinger CT. 2019. Mitochondrial DNA and temperature tolerance in lager yeasts. Sci Adv. 5(1):eaav1869. doi:10.1126/sciadv.aav1869.

Boyle EA, Li YI, Pritchard JK. 2017. An Expanded View of Complex Traits: From Polygenic to Omnigenic. Cell. 169(7):1177–1186. doi:10.1016/j.cell.2017.05.038.

Brem RB, Kruglyak L. 2005. The landscape of genetic complexity across 5,700 gene expression traits in yeast. Proc Natl Acad Sci U S A. 102(5):1572–1577. doi:10.1073/pnas.0408709102.

Burke JM, Arnold ML. 2001. Genetics and the fitness of hybrids. Annu Rev Genet. 35:31–52. doi:10.1146/annurev.genet.35.102401.085719.

Coradetti ST, Pinel D, Geiselman GM, Ito M, Mondo SJ, Reilly MC, Cheng Y-F, Bauer S, Grigoriev IV, Gladden JM, et al. 2018. Functional genomics of lipid metabolism in the oleaginous yeast Rhodosporidium toruloides. Bohlmann J, editor. eLife. 7:e32110. doi:10.7554/eLife.32110.

Dubin CA, Roop JI, Brem RB. 2020. Divergence of Peroxisome Membrane Gene Sequence and Expression Between Yeast Species. G3: Genes, Genomes, Genetics. 10(6):2079–2085. doi:10.1534/g3.120.401304.

Fleet GH. 2003. Yeast interactions and wine flavour. International Journal of Food Microbiology. 86(1):11–22. doi:10.1016/S0168-1605(03)00245-9.

Goddard MR. 2008. Quantifying the complexities of Saccharomyces cerevisiae's ecosystem engineering via fermentation. Ecology. 89(8):2077–2082. doi:10.1890/07-2060.1.

Gonçalves P, Valério E, Correia C, Almeida JMGCF de, Sampaio JP. 2011. Evidence for Divergent Evolution of Growth Temperature Preference in Sympatric Saccharomyces Species. PLOS ONE. 6(6):e20739. doi:10.1371/journal.pone.0020739.

Hittinger CT. 2013. Saccharomyces diversity and evolution: a budding model genus. Trends Genet. 29(5):309–317. doi:10.1016/j.tig.2013.01.002.

Huh W-K, Falvo JV, Gerke LC, Carroll AS, Howson RW, Weissman JS, O'Shea EK. 2003. Global analysis of protein localization in budding yeast. Nature. 425(6959):686–691. doi:10.1038/nature02026.

Lee ME, DeLoache WC, Cervantes B, Dueber JE. 2015. A Highly Characterized Yeast Toolkit for Modular, Multipart Assembly. ACS Synth Biol. 4(9):975–986. doi:10.1021/sb500366v.

Leuenberger P, Ganscha S, Kahraman A, Cappelletti V, Boersema PJ, von Mering C, Claassen M, Picotti P. 2017. Cell-wide analysis of protein thermal unfolding reveals determinants of thermostability. Science. 355(6327). doi:10.1126/science.aai7825.

Li XC, Fay JC. 2017. Cis-Regulatory Divergence in Gene Expression between Two Thermally Divergent Yeast Species. Genome Biol Evol. 9(5):1120–1129. doi:10.1093/gbe/evx072.

Li XC, Peris D, Hittinger CT, Sia EA, Fay JC. 2019. Mitochondria-encoded genes contribute to evolution of heat and cold tolerance in yeast. Sci Adv. 5(1):eaav1848. doi:10.1126/sciadv.aav1848.

McDonald JH, Kreitman M. 1991. Adaptive protein evolution at the Adh locus in Drosophila. Nature. 351(6328):652–654. doi:10.1038/351652a0.

Morellet N, Li X, Wieninger SA, Taylor JL, Bischerour J, Moriau S, Lescop E, Bardiaux B, Mathy N, Assrir N, et al. 2018. Sequence-specific DNA binding activity of the cross-brace zinc finger motif of the piggyBac transposase. Nucleic Acids Research. 46(5):2660–2677. doi:10.1093/nar/gky044.

Orr HA. 1998. The Population Genetics of Adaptation: The Distribution of Factors Fixed During Adaptive Evolution. Evolution. 52(4):935–949. doi:10.1111/j.1558-5646.1998.tb01823.x.

Ott J, Wang J, Leal SM. 2015. Genetic linkage analysis in the age of whole-genome sequencing. Nat Rev Genet. 16(5):275–284. doi:10.1038/nrg3908.

Oughtred R, Rust J, Chang C, Breitkreutz B-J, Stark C, Willems A, Boucher L, Leung G, Kolas N, Zhang F, et al. 2021. The BioGRID database: A comprehensive biomedical resource of curated protein, genetic, and chemical interactions. Protein Sci. 30(1):187–200. doi:10.1002/pro.3978.

Peter J, Chiara MD, Friedrich A, Yue J-X, Pflieger D, Bergström A, Sigwalt A, Barre B, Freel K, Llored A, et al. 2018. Genome evolution across 1,011 Saccharomyces cerevisiae isolates. Nature. 556(7701):339–344. doi:10.1038/s41586-018-0030-5.

Pierleoni A, Martelli PL, Fariselli P, Casadio R. 2007. eSLDB: eukaryotic subcellular localization database. Nucleic Acids Res. 35(Database issue):D208–D212. doi:10.1093/nar/gkl775.

Salvadó Z, Arroyo-López FN, Guillamón JM, Salazar G, Querol A, Barrio E. 2011. Temperature adaptation markedly determines evolution within the genus Saccharomyces. Appl Environ Microbiol. 77(7):2292–2302. doi:10.1128/AEM.01861-10.

Steenwyk JL, Opulente DA, Kominek J, Shen X-X, Zhou X, Labella AL, Bradley NP, Eichman BF, Čadež N, Libkind D, et al. 2019. Extensive loss of cell-cycle and DNA repair genes in an ancient lineage of bipolar budding yeasts. PLOS Biology. 17(5):e3000255. doi:10.1371/journal.pbio.3000255.

Stern DL. 2014. Identification of loci that cause phenotypic variation in diverse species with the reciprocal hemizygosity test. Trends Genet. 30(12):547–554. doi:10.1016/j.tig.2014.09.006.

Sweeney JY, Kuehne HA, Sniegowski PD. 2004. Sympatric natural Saccharomyces cerevisiae and S. paradoxus populations have different thermal growth profiles. FEMS Yeast Res. 4(4–5):521–525. doi:10.1016/S1567-1356(03)00171-5.

Szklarczyk D, Gable AL, Nastou KC, Lyon D, Kirsch R, Pyysalo S, Doncheva NT, Legeay M, Fang T, Bork P, et al. 2021. The STRING database in 2021: customizable protein–protein networks, and functional characterization of user-uploaded gene/measurement sets. Nucleic Acids Research. 49(D1):D605–D612. doi:10.1093/nar/gkaa1074.

Tam V, Patel N, Turcotte M, Bossé Y, Paré G, Meyre D. 2019. Benefits and limitations of genome-wide association studies. Nat Rev Genet. 20(8):467–484. doi:10.1038/s41576-019-0127-1.

Tirosh I, Reikhav S, Levy AA, Barkai N. 2009. A Yeast Hybrid Provides Insight into the Evolution
of Gene Expression Regulation. Science. 324(5927):659–662. doi:10.1126/science.1169766.
Waterhouse AM, Procter JB, Martin DMA, Clamp M, Barton GJ. 2009. Jalview Version 2—a
multiple sequence alignment editor and analysis workbench. Bioinformatics. 25(9):1189–1191.
doi:10.1093/bioinformatics/btp033.
Weiss CV, Brem RB. 2019. Dissecting Trait Variation across Species Barriers. Trends in
Ecology & Evolution. 34(12):1131–1136. doi:10.1016/j.tree.2019.07.013.
Weiss CV, Roop JI, Hackley RK, Chuong JN, Grigoriev IV, Arkin AP, Skerker JM, Brem RB.
2018. Genetic dissection of interspecific differences in yeast thermotolerance. Nature Genetics.
50(11):1501. doi:10.1038/s41588-018-0243-4.
Wetmore KM, Price MN, Waters RJ, Lamson JS, He J, Hoover CA, Blow MJ, Bristow J, Butland
G, Arkin AP, et al. 2015. Rapid Quantification of Mutant Fitness in Diverse Bacteria by
Sequencing Randomly Bar-Coded Transposons. Moran MA, editor. mBio. 6(3):e00306-15.
doi:10.1128/mBio.00306-15.
Winzeler EA. 1999. Functional Characterization of the S. cerevisiae Genome by Gene
Deletion and Parallel Analysis. Science. 285(5429):901–906.
doi:10.1126/science.285.5429.901.
Yang Z. 2007. PAML 4: Phylogenetic Analysis by Maximum Likelihood. Molecular Biology and
Evolution. 24(8):1586–1591. doi:10.1093/molbev/msm088.
Yusa K, Zhou L, Li MA, Bradley A, Craig NL. 2011. A hyperactive piggyBac transposase for
mammalian applications. PNAS. 108(4):1531–1536. doi:10.1073/pnas.1008322108.
863

FIGURE CAPTIONS

Figure 1. Barcoded RH-seq mapping of yeast thermotolerance loci. (A) Barcoded RH-seq sequencing analysis steps. Left, in a pool of *S. cerevisiae* x *S. paradoxus* hybrid hemizygotes, each harboring a transposon (grey rectangle) marked with a unique 20-mer barcode (multicolored) flanked by universal primer sites (U1 and U2), each barcode is associated with its insertion location by transposon sequencing (Tn-seq). Genomic DNA from the pool is extracted, sheared, and ligated to universal adapters (pink ovals), followed by PCR amplification with a transposon-specific primer (forward black arrow) and an adapter-specific primer (reverse black arrow) and sequencing. Right, for barcode sequencing (Bar-seq) to quantify hemizygote strain abundance after pool growth in a condition of interest, genomic DNA is used as input to PCR with primers to universal primer sites for sequencing. (B) Thermotolerance RH-seq screen design. An aliquot of the hemizygote pool was thawed and cultured in large format, then split into small replicate cultures, each maintained in logarithmic growth phase at the temperature of interest by back-dilution, followed by quantification by Bar-seq.

Figure 2. Hits from barcoded RH-seq mapping of yeast thermotolerance. (A) Each panel reports barcoded RH-seq results for a gene at which the *S. cerevisiae* allele is associated with better thermotolerance than the *S. paradoxus* allele, when uncovered in the hybrid background. In a given panel, the *x*-axis reports the $\log_2$ of abundance, measured by RH-seq after selection at 37°C, of a clone harboring a barcoded transposon insertion in the indicated species' allele in a given replicate, as a difference from the analogous quantity for that clone after selection at 28°C on average across replicates. The *y*-axis reports the proportion of observations of all clones bearing insertions in the indicated allele that exhibited the abundance ratio on the *x*, as a kernel density estimate. Shown are the top six genes from among all barcoded RH-seq hit loci in terms of allelic effect size; see Table S7 for effect sizes of the complete set of hits. (B) Subcellular localization of RH-seq hit genes, where available from (Pierleoni et al. 2007) and (Huh et al. 2003). Genes at which effects of allelic variation on thermotolerance were reported previously (Weiss et al. 2018; Li et al. 2019) are denoted in bold type.

Figure 3. Interactions between thermotolerance loci. Each node represents a top hit gene from barcoded RH-seq mapping of thermotolerance. Each chord represents an inferred interaction, taking into account physical and genetic interactions as well as pathway membership, from the STRING database (Szklarczyk et al. 2021). Chords are weighted by the confidence of the inference of interactions; nodes are colored by the number of inferred interactions with other top hits, such that genes with higher numbers of interactions among the hits are represented by warmer colored nodes.

Figure 4. Codons under positive selection in thermotolerance loci. Each panel shows the amino acid sequence context of the codon(s) (red bar) inferred to be under positive selection along the *S. cerevisiae* lineage, in a hit gene from RH-seq thermotolerance mapping. Alignments are colored by percent identity, with darker purples indicating a higher percent identity. (A) *YDR508C/GNP1*. (B) *YGR140W/CBF2*. (C) *YMR078C/CTF18*.

TABLES

GO term	$n_{observed}$: $n_{expected}$	Adjusted p	Name	Total n
Cellular Component				
GO:0000775	5:1	0.0366	chromosome, centromeric region	75
GO:0000778	4:0	0.0256	kinetochore	40
Molecular Function				
GO:0000149	3:0	0.0701	SNARE binding	28
GO:0008081	2:0	0.0256	phosphoric diester hydrolase activity	11
GO:0004843	2:0	0.0998	thiol-dependent deubiquitinase	24
Biological Process				
GO:0007165	3:0	0.0923	signal transduction	59
GO:0001403	3:0	0.0923	invasive growth in response to glucose limitation	42
GO:0046580	2:0	0.0256	negative regulation of Ras protein signal transduction	6
GO:0001934	2:0	0.0256	positive regulation of protein phosphorylation	5
GO:0016042	2:0	0.0923	lipid catabolic process	26
GO:0034087	2:0	0.0923	establishment of mitotic sister chromatid cohesion	16

906
Table 1. Functional enrichment among thermotolerance loci. Each row with numerical data reports a
Gene Ontology (GO) term enriched for RH-seq hit genes. $n_{observed}$, the number of genes from among top
hits from thermotolerance RH-seq that were annotated with the term. $n_{expected}$, the number of genes
annotated with the term in the same number of randomly chosen genes from the genome, as a median
across samples. Adjusted p, resampling-based significance of the enrichment after Benjamini-Hochberg
correction. Total n, the total number of genes annotated in the GO term in *S. cerevisiae*.

Supplementary Figure 1. Making barcoded hemizygotes in a yeast hybrid background for RH-seq.
(A) A pool of random N20 barcodes (colors), each flanked by universal priming sites (U1 and U2), was
used as input into a PCR with primers containing recognition sites for the BbsI type IIS restriction enzyme.
(B) In a plasmid harboring an un-barcoded piggyBac transposon (gray rectangle) (the kanamycin
resistance cassette, kan^R, flanked by left and right transposon arms) and transposase (teal rectangle)
(Weiss et al. 2018), a 42 nucleotide stuffer sequence, consisting of two BbsI restriction enzyme sites
flanking a NotI restriction enzyme site and custom overhang sequences (Lee et al. 2015), replaced 42
nucleotides in the right arm of the transposon. (C) BbsI digestion of the barcodes and stuffer-containing
plasmid, followed by stuffer loss and ligation of a barcode into each plasmid, yielded a pool of barcoded
plasmids. (D) Transformation of the barcoded transposase plasmid into *S. cerevisiae* x *S. paradoxus*
hybrids, followed by transposition and plasmid loss, yielded a pool of marked transposon hemizygote
insertion genotypes in the hybrid background.

**Supplementary Figure 2. Modifying yeast piggyBac to test barcode insertion positions and
transposase optimization, and effects of barcode insertion positions and transposase sequence
on yeast piggyBac transposition efficiency.** (A) Left, a plasmid from (Weiss et al. 2018) containing the
unbarcoded piggyBac transposon (gray) and transposase (teal) was modified to eliminate three BbsI
restriction enzyme sites, and used as a backbone for further modifications. Right, test plasmids were
mutated at transposase sites designed to optimize codons and increase activity (Yusa et al. 2011).
Bottom, test plasmids were modified to incorporate into the transposon a single 20 nucleotide barcode
flanked by universal priming regions and custom two-nucleotide overhang sequences (blue squares),
either by insertion between the 3'-most end of the left arm and 5' end of the TEF promoter of the
kanamycin cassette (bottom left) or replacing endogenous nucleotides inside the right arm of the
transposon (bottom right). Pink rectangles indicate transposase binding sites from (Morellet et al. 2018).
(B) Each pair of boxes reports transposition test results from a plasmid schematized in (A) in the *S.
cerevisiae* x *S. paradoxus* F1 hybrid, with transformation at the indicated temperature. For a given box,
the thick black line reports the median; the box extent report quartiles; whiskers report outliers.

Supplementary Figure 3. Gene coverage and read depth in thermotolerance Bar-seq. (A) The *x*-axis
reports the number of inferred hemizygote clones in a given gene (corresponding to transposon insertion
mutants) whose abundance was detectable in Bar-seq (see Figure 1B), and each bar height reports the
number of genes with the number of detectable hemizygotes on the *x*, for the indicated species' allele in
the diploid hybrid. The dotted red line indicates the cutoff used in our quality control pipeline for tests of
allelic impact on thermotolerance, whereby only genes with greater than three inserts for an allele in the
Bar-seq counts were considered. (B) The *x*-axis reports the total number of Bar-seq reads, for a given
inferred hemizygote clone in the indicated species' allele, in competition cultures grown at 28°C; each bar
height reports the number of inferred hemizygote clones with the Bar-seq abundance on the *x*. (C) Data
are as in (B) except that competitions at 37°C were analyzed.

**Supplementary Figure 4. Impact on high-temperature growth of allelic variation, in barcoded RH-
seq, at genes from a previous thermotolerance screen.** (A) Each row reports the allelic effect, the
thermotolerance conferred by disruption of the *S. cerevisiae* allele, relative to the analogous quantity for
the *S. paradoxus* allele, as measured in barcoded RH-seq, of a gene at which allelic variation was
previously reported to impact thermotolerance (Weiss et al. 2018). Genes marked with asterisks were
significant at *p* < 0.05, after quality control for noise and number of inserts and multiple-hypothesis
correction. (B) The *x*-axis reports allelic effect for a given gene as in (A); the *y*-axis reports the proportion
of genes with the allelic effect on the *x*, with the blue trace showing the distribution across all genes with
barcoded RH-seq data, as a kernel density estimate. Red dotted vertical lines represent genes from (A).

Supplementary Figure 5. **Accelerated evolution of thermotolerance loci.** Shown are results of
analyses of branch length of top hit genes from barcoded RH-seq mapping of thermotolerance, as
inferred from gene trees and normalized for gene length. Each vertical bar reports inferred branch length,
along the *S. cerevisiae* lineage, of the indicated RH-seq hit gene. Horizontal lines report median branch
lengths across the indicated gene sets. A resampling test for long branches on the *S. cerevisiae* lineage

969   among top RH-seq hits revealed significant evidence for enrichment (p = 0.0465) but not when *TAF2* and
970   *BUL1* were eliminated (p = 0.1574), attesting to the particularly strong inference of accelerated evolution
971   in the latter two genes.

972 **SUPPLEMENTARY TABLES**
973
974 **Supplementary Table 1. Plasmids used in this study.**

Oligo name	Sequence (5' to 3')	Index Identifier	Index Sequence	Notes
Barcoding cloning				
Random Barcodes U1 — N20 — U2	GATGTCCACGAGGTCTCTNNNNNNNN NNNNNNNNNNNNNNCGTACGCTGCAG GTCGAC			Random barcodes flanked by universal primer sites
FW_BbsI_JC	TCACACAAGTTTGTACAAAAAAGCAG GCTGGAGCTCGgaagacATCCCTGATG TCCACGAGGTCTCT			Forward primer to amplify barcodes for cloning into vector
REV_BbsI_JC	CTCAACCACTTTGTACAAGAAAGCTG GGTGGATCCgaagacCGCGTTGTCGAC CTGCAGCGTACG			Reverse primer for to amplify barcodes for cloning into vector
Tn-Seq				
JC8 Tn-specific FW Primer	ATGATACGGCGACCACCGAGATCTAC ACTCTTTCCCTACACGACGCTCTTCC GATCTNNNNNNCCCTGATGTCCACGA GGTCTCT			Forward primer to sequence transposon insertions; homologous to transposon (Figure 1A, left)
P7_MOD_TS_index1	CAAGCAGAAGACGGCATACGAGATC GTGATGTGACTGGAGTTCAGACGTGT GCTCTTCCGATCT	Index1	ATCACG	Indexed reverse primer to sequence transposon insertions; homologous to adapter (Figure 1A, left)
P7_MOD_TS_index2	CAAGCAGAAGACGGCATACGAGATA CATCGGTGACTGGAGTTCAGACGTGT GCTCTTCCGATCT	Index2	CGATGT	Indexed reverse primer to sequence transposon insertions; homologous to adapter (Figure 1A, left)
P7_MOD_TS_index3	CAAGCAGAAGACGGCATACGAGATG CCTAAGTGACTGGAGTTCAGACGTGT GCTCTTCCGATCT	Index3	TTAGGC	Indexed reverse primer to sequence transposon insertions; homologous to adapter (Figure 1A, left)
P7_MOD_TS_index4	CAAGCAGAAGACGGCATACGAGATT GGTCAGTGACTGGAGTTCAGACGTG TGCTCTTCCGATCT	Index4	TGACCA	Indexed reverse primer to sequence transposon

				insertions; homologous to adapter (Figure 1A, left)
P7_MOD_TS_index5	CAAGCAGAAGACGGCATACGAGATC ACTGTGTGACTGGAGTTCAGACGTGT GCTCTTCCGATCT	Index5	ACAGTG	Indexed reverse primer to sequence transposon insertions; homologous to adapter (Figure 1A, left)
P7_MOD_TS_index6	CAAGCAGAAGACGGCATACGAGATAT TGGCGTGACTGGAGTTCAGACGTGT GCTCTTCCGATCT	Index6	GCCAAT	Indexed reverse primer to sequence transposon insertions; homologous to adapter (Figure 1A, left)
P7_MOD_TS_index7	CAAGCAGAAGACGGCATACGAGATG ATCTGGTGACTGGAGTTCAGACGTGT GCTCTTCCGATCT	Index7	CAGATC	Indexed reverse primer to sequence transposon insertions; homologous to adapter (Figure 1A, left)
P7_MOD_TS_index8	CAAGCAGAAGACGGCATACGAGATT CAAGTGTGACTGGAGTTCAGACGTGT GCTCTTCCGATCT	Index8	ACTTGA	Indexed reverse primer to sequence transposon insertions; homologous to adapter (Figure 1A, left)
Bar-seq				
P1_BS3_IT001	AATGATACGGCGACCACCGAGATCTA CACTCTTTCCCTACACGACGCTCTTC CGATCTNGCACTAGTCGACCTGCAG CGTACG	IT001	ATCACG	Indexed forward Bar-seq primer (Figure 1A, right)
P1_BS3_IT002	AATGATACGGCGACCACCGAGATCTA CACTCTTTCCCTACACGACGCTCTTC CGATCTNNTGTAGCGTCGACCTGCA GCGTACG	IT002	CGATGT	Indexed forward Bar-seq primer (Figure 1A, right)
P1_BS3_IT003	AATGATACGGCGACCACCGAGATCTA CACTCTTTCCCTACACGACGCTCTTC CGATCTNNNCGGATTGTCGACCTGCA GCGTACG	IT003	TTAGGC	Indexed forward Bar-seq primer (Figure 1A, right)
P1_BS3_IT004	AATGATACGGCGACCACCGAGATCTA CACTCTTTCCCTACACGACGCTCTTC	IT004	TGACCA	Indexed forward Bar-seq primer

	CGATCTNNNNACCAGTGTCGACCTG CAGCGTACG			(Figure 1A, right)
P1_BS3_IT005	AATGATACGGCGACCACCGAGATCTA CACTCTTTCCCTACACGACGCTCTTC CGATCTNGTGACAGTCGACCTGCAG CGTACG	IT005	ACAGTG	Indexed forward Bar-seq primer (Figure 1A, right)
P1_BS3_IT006	AATGATACGGCGACCACCGAGATCTA CACTCTTTCCCTACACGACGCTCTTC CGATCTNNTAACCGGTCGACCTGCA GCGTACG	IT006	GCCAAT	Indexed forward Bar-seq primer (Figure 1A, right)
P1_BS3_IT007	AATGATACGGCGACCACCGAGATCTA CACTCTTTCCCTACACGACGCTCTTC CGATCTNNNCTAGACGTCGACCTGCA GCGTACG	IT007	CAGATC	Indexed forward Bar-seq primer (Figure 1A, right)
P1_BS3_IT008	AATGATACGGCGACCACCGAGATCTA CACTCTTTCCCTACACGACGCTCTTC CGATCTNNNNAGTTCAGTCGACCTGC AGCGTACG	IT008	ACTTGA	Indexed forward Bar-seq primer (Figure 1A, right)
P1_BS3_IT009	AATGATACGGCGACCACCGAGATCTA CACTCTTTCCCTACACGACGCTCTTC CGATCTNGACTAGGTCGACCTGCAG CGTACG	IT009	GATCAG	Indexed forward Bar-seq primer (Figure 1A, right)
P1_BS3_IT010	AATGATACGGCGACCACCGAGATCTA CACTCTTTCCCTACACGACGCTCTTC CGATCTNNTTCGATGTCGACCTGCAG CGTACG	IT010	TAGCTT	Indexed forward Bar-seq primer (Figure 1A, right)
P1_BS3_IT011	AATGATACGGCGACCACCGAGATCTA CACTCTTTCCCTACACGACGCTCTTC CGATCTNNNCATCGGGTCGACCTGC AGCGTACG	IT011	GGCTAC	Indexed forward Bar-seq primer (Figure 1A, right)
P1_BS3_IT012	AATGATACGGCGACCACCGAGATCTA CACTCTTTCCCTACACGACGCTCTTC CGATCTNNNNATGTTCGTCGACCTGC AGCGTACG	IT012	CTTGTA	Indexed forward Bar-seq primer (Figure 1A, right)
P1_BS3_IT013	AATGATACGGCGACCACCGAGATCTA CACTCTTTCCCTACACGACGCTCTTC CGATCTNAACTGAGTCGACCTGCAGC GTACG	IT013	AGTCAA	Indexed forward Bar-seq primer (Figure 1A, right)
P1_BS3_IT014	AATGATACGGCGACCACCGAGATCTA CACTCTTTCCCTACACGACGCTCTTC CGATCTNNCCTTGAGTCGACCTGCAG CGTACG	IT014	AGTTCC	Indexed forward Bar-seq primer (Figure 1A, right)
P1_BS3_IT015	AATGATACGGCGACCACCGAGATCTA CACTCTTTCCCTACACGACGCTCTTC	IT015	ATGTCA	Indexed forward Bar-seq primer

	CGATCTNNNACTGTAGTCGACCTGCAGCGTACG			(Figure 1A, right)
P1_BS3_IT016	AATGATACGGCGACCACCGAGATCTACACTCTTTCCCTACACGACGCTCTTCCGATCTNNNNCCTGCCGTCGACCTGCAGCGTACG	IT016	CCGTCC	Indexed forward Bar-seq primer (Figure 1A, right)
P1_BS3_IT017	AATGATACGGCGACCACCGAGATCTACACTCTTTCCCTACACGACGCTCTTCCGATCTNGAGATGGTCGACCTGCAGCGTACG	IT017	GTAGAG	Indexed forward Bar-seq primer (Figure 1A, right)
P1_BS3_IT018	AATGATACGGCGACCACCGAGATCTACACTCTTTCCCTACACGACGCTCTTCCGATCTNNCGCCTGGTCGACCTGCAGCGTACG	IT018	GTCCGC	Indexed forward Bar-seq primer (Figure 1A, right)
P1_BS3_IT019	AATGATACGGCGACCACCGAGATCTACACTCTTTCCCTACACGACGCTCTTCCGATCTNNNAAAGTGGTCGACCTGCAGCGTACG	IT019	GTGAAA	Indexed forward Bar-seq primer (Figure 1A, right)
P1_BS3_IT020	AATGATACGGCGACCACCGAGATCTACACTCTTTCCCTACACGACGCTCTTCCGATCTNNNNCCGGTGGTCGACCTGCAGCGTACG	IT020	GTGGCC	Indexed forward Bar-seq primer (Figure 1A, right)
P1_BS3_IT021	AATGATACGGCGACCACCGAGATCTACACTCTTTCCCTACACGACGCTCTTCCGATCTNGCTTTGGTCGACCTGCAGCGTACG	IT021	GTTTCG	Indexed forward Bar-seq primer (Figure 1A, right)
P1_BS3_IT022	AATGATACGGCGACCACCGAGATCTACACTCTTTCCCTACACGACGCTCTTCCGATCTNNGCATGCGTCGACCTGCAGCGTACG	IT022	CGTACG	Indexed forward Bar-seq primer (Figure 1A, right)
P1_BS3_IT023	AATGATACGGCGACCACCGAGATCTACACTCTTTCCCTACACGACGCTCTTCCGATCTNNNGGTGAGGTCGACCTGCAGCGTACG	IT023	GAGTGG	Indexed forward Bar-seq primer (Figure 1A, right)
P1_BS3_IT024	AATGATACGGCGACCACCGAGATCTACACTCTTTCCCTACACGACGCTCTTCCGATCTNNNNCGATGGGTCGACCTGCAGCGTACG	IT024	GGTAGC	Indexed forward Bar-seq primer (Figure 1A, right)
P2_BS3_IT001	CAAGCAGAAGACGGCATACGAGATCGTGATGTGACTGGAGTTCAGACGTGTGCTCTTCCGATCTGATGTCCACGAGGTCTCT	IT001	ATCACG	Indexed reverse Bar-seq primer (Figure 1A, right)
P2_BS3_IT002	CAAGCAGAAGACGGCATACGAGATACATCGGTGACTGGAGTTCAGACGTGT	IT002	CGATGT	Indexed reverse Bar-seq primer

	GCTCTTCCGATCTGATGTCCACGAGG TCTCT			(Figure 1A, right)
P2_BS3_IT003	CAAGCAGAAGACGGCATACGAGATG CCTAAGTGACTGGAGTTCAGACGTGT GCTCTTCCGATCTGATGTCCACGAGG TCTCT	IT003	TTAGGC	Indexed reverse Bar-seq primer (Figure 1A, right)
P2_BS3_IT004	CAAGCAGAAGACGGCATACGAGATT GGTCAGTGACTGGAGTTCAGACGTG TGCTCTTCCGATCTGATGTCCACGAG GTCTCT	IT004	TGACCA	Indexed reverse Bar-seq primer (Figure 1A, right)
P2_BS3_IT005	CAAGCAGAAGACGGCATACGAGATC ACTGTGTGACTGGAGTTCAGACGTGT GCTCTTCCGATCTGATGTCCACGAGG TCTCT	IT005	ACAGTG	Indexed reverse Bar-seq primer (Figure 1A, right)
P2_BS3_IT006	CAAGCAGAAGACGGCATACGAGATAT TGGCGTGACTGGAGTTCAGACGTGT GCTCTTCCGATCTGATGTCCACGAGG TCTCT	IT006	GCCAAT	Indexed reverse Bar-seq primer (Figure 1A, right)
P2_BS3_IT007	CAAGCAGAAGACGGCATACGAGATG ATCTGGTGACTGGAGTTCAGACGTGT GCTCTTCCGATCTGATGTCCACGAGG TCTCT	IT007	CAGATC	Indexed reverse Bar-seq primer (Figure 1A, right)
P2_BS3_IT008	CAAGCAGAAGACGGCATACGAGATT CAAGTGTGACTGGAGTTCAGACGTGT GCTCTTCCGATCTGATGTCCACGAGG TCTCT	IT008	ACTTGA	Indexed reverse Bar-seq primer (Figure 1A, right)
P2_BS3_IT009	CAAGCAGAAGACGGCATACGAGATC TGATCGTGACTGGAGTTCAGACGTGT GCTCTTCCGATCTGATGTCCACGAGG TCTCT	IT009	GATCAG	Indexed reverse Bar-seq primer (Figure 1A, right)
P2_BS3_IT010	CAAGCAGAAGACGGCATACGAGATA AGCTAGTGACTGGAGTTCAGACGTGT GCTCTTCCGATCTGATGTCCACGAGG TCTCT	IT010	TAGCTT	Indexed reverse Bar-seq primer (Figure 1A, right)
P2_BS3_IT011	CAAGCAGAAGACGGCATACGAGATG TAGCCGTGACTGGAGTTCAGACGTGT GCTCTTCCGATCTGATGTCCACGAGG TCTCT	IT011	GGCTAC	Indexed reverse Bar-seq primer (Figure 1A, right)
P2_BS3_IT012	CAAGCAGAAGACGGCATACGAGATTA CAAGGTGACTGGAGTTCAGACGTGT GCTCTTCCGATCTGATGTCCACGAGG TCTCT	IT012	CTTGTA	Indexed reverse Bar-seq primer (Figure 1A, right)
P2_BS3_IT013	CAAGCAGAAGACGGCATACGAGATTT GACTGTGACTGGAGTTCAGACGTGT	IT013	AGTCAA	Indexed reverse Bar-seq primer

	GCTCTTCCGATCTGATGTCCACGAGG TCTCT			(Figure 1A, right)
P2_BS3_IT014	CAAGCAGAAGACGGCATACGAGATG GAACTGTGACTGGAGTTCAGACGTGT GCTCTTCCGATCTGATGTCCACGAGG TCTCT	IT014	AGTTCC	Indexed reverse Bar-seq primer (Figure 1A, right)
P2_BS3_IT015	CAAGCAGAAGACGGCATACGAGATT GACATGTGACTGGAGTTCAGACGTGT GCTCTTCCGATCTGATGTCCACGAGG TCTCT	IT015	ATGTCA	Indexed reverse Bar-seq primer (Figure 1A, right)
P2_BS3_IT016	CAAGCAGAAGACGGCATACGAGATG GACGGGTGACTGGAGTTCAGACGTG TGCTCTTCCGATCTGATGTCCACGAG GTCTCT	IT016	CCGTCC	Indexed reverse Bar-seq primer (Figure 1A, right)
P2_BS3_IT017	CAAGCAGAAGACGGCATACGAGATC TCTACGTGACTGGAGTTCAGACGTGT GCTCTTCCGATCTGATGTCCACGAGG TCTCT	IT017	GTAGAG	Indexed reverse Bar-seq primer (Figure 1A, right)
P2_BS3_IT018	CAAGCAGAAGACGGCATACGAGATG CGGACGTGACTGGAGTTCAGACGTG TGCTCTTCCGATCTGATGTCCACGAG GTCTCT	IT018	GTCCGC	Indexed reverse Bar-seq primer (Figure 1A, right)
P2_BS3_IT019	CAAGCAGAAGACGGCATACGAGATTT TCACGTGACTGGAGTTCAGACGTGTG CTCTTCCGATCTGATGTCCACGAGGT CTCT	IT019	GTGAAA	Indexed reverse Bar-seq primer (Figure 1A, right)
P2_BS3_IT020	CAAGCAGAAGACGGCATACGAGATG GCCACGTGACTGGAGTTCAGACGTG TGCTCTTCCGATCTGATGTCCACGAG GTCTCT	IT020	GTGGCC	Indexed reverse Bar-seq primer (Figure 1A, right)
P2_BS3_IT021	CAAGCAGAAGACGGCATACGAGATC GAAACGTGACTGGAGTTCAGACGTGT GCTCTTCCGATCTGATGTCCACGAGG TCTCT	IT021	GTTTCG	Indexed reverse Bar-seq primer (Figure 1A, right)
P2_BS3_IT022	CAAGCAGAAGACGGCATACGAGATC GTACGGTGACTGGAGTTCAGACGTG TGCTCTTCCGATCTGATGTCCACGAG GTCTCT	IT022	CGTACG	Indexed reverse Bar-seq primer (Figure 1A, right)
P2_BS3_IT023	CAAGCAGAAGACGGCATACGAGATC CACTCGTGACTGGAGTTCAGACGTGT GCTCTTCCGATCTGATGTCCACGAGG TCTCT	IT023	GAGTGG	Indexed reverse Bar-seq primer (Figure 1A, right)
P2_BS3_IT024	CAAGCAGAAGACGGCATACGAGATG CTACCGTGACTGGAGTTCAGACGTGT	IT024	GGTAGC	Indexed reverse Bar-seq primer

	GCTCTTCCGATCTGATGTCCACGAGG TCTCT			(Figure 1A, right)

977 **Supplementary Table 2. Oligonucleotides used in this study.**

Experiment	Library	Reads
	RBJC009_IT013	93,277,156
	RBJC009_IT014	73,914,499
	RBJC009_IT015	79,273,122
	RBJC009_IT016	81,404,878
	RBJC45_IT013	100,794,463
	RBJC45_IT014	92,410,111
	RBJC45_IT015	91,931,624
	RBJC45_I0T16	96,631,037
	RBJC009_IT013	93,277,156
	RBJC009_IT014	73,914,499
	RBJC009_IT015	79,273,122
E. coli vector pool Bar-seq	RBJC45_IT016	96,631,037
	RMBA038A_IT001	16,042,537
	RBMA038A_IT002	19,451,725
	RBMA038A_IT003	16,313,279
	RBMA038A_IT004	19,512,043
	RBMA038A_IT005	13,625,004
	RBMA038A_IT006	22,257,379
	RBMA038A_IT007	22,385,133
	RBMA038A_IT008	16,359,601
	RBMA038A_IT009	20,118,293
	RBMA038A_IT010	19,121,369
	RBMA038A_IT011	18,887,044
	RBMA038A_IT012	15,895,905
	RBMA038A_IT013	18,922,229
	RBMA038A_IT014	26,602
	RBMA038A_IT015	16,571,744
	RBMA038A_IT016	18,783,445
	RBMA038A_IT017	17,671,257
	RBMA038A_IT018	18,040,605
	RBMA038A_IT019	16,929,985
	RBMA038A_IT020	15,319,721
	RBMA038A_IT021	16,897,401
	RBMA038A_IT022	16,372,841
37°C vs. 28°C barcoded RH-seq in yeast	RBMA038A_IT023	16,679,583
	RBMA038A_IT024	13,875,240
	RMBA039A_IT001	28,004,341
	RBMA039A_IT002	19,283,097
	RBMA039A_IT003	12,427,040
	RBMA039A_IT004	13,934,236
	RBMA039A_IT005	12,068,707
	RBMA039A_IT006	60,072,532
	RBMA039A_IT008	12,500,340
	RBMA039A_IT009	25,029,516
36°C vs. 28°C barcoded RH-seq in yeast	RBMA039A_IT010	14,765,007
	RBMA039A_IT011	7,059,133

RBMA039A_IT012	19,837,690
RBMA039A_IT013	14,393,602
RBMA039A_IT014	19,572,649
RBMA039A_IT015	8,814,383
RBMA039A_IT016	14,587,181
RBMA039A_IT017	8,619,661
RBMA039A_IT018	13,216,843
RBMA039A_IT019	12,499,486
RBMA039A_IT020	11,963,866
RBMA039A_IT021	6,626,989
RBMA039A_IT022	11,649,542
RBMA039A_IT023	5,813,357
RBMA039A_IT024	9,349,172

980
981 **Supplementary Table 3. Bar-seq sequencing data sets.** Each row reports numbers of reads
982 sequenced for the indicated Bar-seq experiment. The first set of rows reports results from a check of
983 barcoded piggyBac transposon plasmids as in Figure S1C; the remaining rows report results from
984 quantification of yeast hemizygote insertion genotypes after competition in the indicated condition, as in
985 Figure 1B of the main text. Experiment identifiers are from BioProject PRJNA735401.

Pool	Library	Reads	Platform	Facility
67	RBJC37	38,713,102	Novaseq SP PE150	UC Berkeley
69	RBJC38	38,875,221		
69	RBJC39	43,194,450		
69	RBJC40	39,778,862		
69	RBJC41	38,836,065		
67	RBJC42	39,265,466		
67	RBJC43	47,124,575		
67	RBJC44	39,762,187		
67	RBJC48	91,531,071	HiSeq4000 PE150	Novogene
67	RBJC48_reseq	86,892,060		
70	RBCJ51	86,254,426		
70	RBCJ51_reseq	86,130,880		
70	RBJC52	52,108,306		
70	RBJC52_reseq	53,363,169		
71	RBJC54	88,154,532		
71	RBJC54_reseq	86,878,835		
71	RBJC55	90,265,981		
71	RBJC55_reseq	82,130,170		
69	RBJC57	84,296,399		
69	RBJC57_reseq	85,606,080		

Supplementary Table 4. Tn-seq sequencing data sets. Each row reports numbers of reads from the
indicated sequencing of insertion positions of barcoded transposons in the *S. cerevisiae* x *S. paradoxus*
hybrid, as in Figure 1A, left, of the main text. Experiment identifiers are from PRJNA735401; "reseq"
indicates the reads from a technical replicate performed to gather additional reads for the indicated
library.

Supplementary Table 5. Abundance of inferred hemizygote insertion genotypes from
thermotolerance RH-seq. Each row reports the results of sequencing from one inferred transposon
insertion in the *S. cerevisiae* x *S. paradoxus* diploid hybrid after selection of the barcoded transposon pool
after competitions comparing growth at 37°C and 28°C, reflecting the abundance in the pool of the
respective hemizygote clone harboring the insertion. Chromosome, strand, location, and gene report the
fine-scale position of the inferred insertion. Allele, the species parent's homolog in which the transposon
insertion lay. Abundance, read counts of the transposon insertion sequenced after selection of the
barcoded transposon pool at the indicated temperature, normalized for library size and averaged across
the biological replicate cultures. Transposon insertions not detected in any replicate of the indicated
selection were assigned an abundance of 1 prior to normalization by library size. CV, coefficient of
variation over biological replicates of normalized read counts after selection at the indicated temperature.
Barcode, the unique barcode identifier of the transposon insertion.

Supplementary Table 6. Abundance of hemizygote insertion genotypes from RH-seq at 36°C. Data
are as in Table S5, except that RH-seq was done using 36°C as the high-temperature condition.

Supplementary Table 7. Effects of allelic variation in thermotolerance RH-seq. Each row reports the
results of reciprocal hemizygote tests on thermotolerance at the indicated gene in the *S. cerevisiae* x *S.*
paradoxus diploid hybrid at 37°C. Columns B-G report analyses of abundance upon the aggregation at
the gene level of inferred hemizygote genotypes (Table S5) from all biological replicate experiments,
filtered for quality control (see Methods). Columns B-D report results of a two-tailed Mann-Whitney
statistical test for a difference in the abundance after growth at 37°C, relative to the abundance after
growth at 28°C, of hemizygotes harboring transposon insertions in the two species parents' homologs.
The Benjamini-Hochberg method was used to correct for multiple testing. Column E reports the
$\log_2$(abundance at 37°C/abundance at 28°C) of the average insert in the *S. cerevisiae* allele. Column F
reports the analogous quantity among inserts in the *S. paradoxus* allele of the gene. Column G reports
the allele-specific effect size, calculated as the difference between the measures of Columns E and F.

Supplementary Table 8. **Effects of allelic variation in RH-seq at 36°C.** Data are as in Table S7, except
that the high temperature growth condition was 36°C.

Hit	Description
YGR198W/YPP1	Cargo-transport protein involved in endocytosis; interacts with phosphatidylinositol-4-kinase Stt4; GFP-fusion protein localizes to the cytoplasm; YGR198W is an essential gene
YMR207C/HFA1	Mitochondrial acetyl-coenzyme A carboxylase, catalyzes the production of malonyl-CoA in mitochondrial fatty acid biosynthesis
YGL082W/MIY1	Putative protein of unknown function; predicted prenylation/proteolysis target of Afc1p and Rce1p; green fluorescent protein (GFP)-fusion protein localizes to the cytoplasm and nucleus; YGL082W is not an essential gene
YNL049C/SFB2	Component of the Sec23p-Sfb2p heterodimer of the COPII vesicle coat, required for cargo selection during vesicle formation in ER to Golgi transport; homologous to Sec24p and Sfb3p
YDL035C/GPR1	Plasma membrane G protein coupled receptor (GPCR) that interacts with the heterotrimeric G protein alpha subunit, Gpa2p, and with Plc1p; sensor that integrates nutritional signals with the modulation of cell fate via PKA and cAMP synthesis
YDR508C/GNP1	High-affinity glutamine permease, also transports Leu, Ser, Thr, Cys, Met and Asn; expression is fully dependent on Grr1p and modulated by the Ssy1p-Ptr3p-Ssy5p (SPS) sensor of extracellular amino acids
YBR136W/MEC1	Genome integrity checkpoint protein and PI kinase superfamily member; signal transducer required for cell cycle arrest and transcriptional responses prompted by damaged or unreplicated DNA; monitors and participates in meiotic recombination
YML099C/ARG81	Zinc-finger transcription factor of the Zn(2)-Cys(6) binuclear cluster domain type, involved in the regulation of arginine-responsive genes; acts with Arg80p and Arg82p
YPL254W/HFI1	Adaptor protein required for structural integrity of the SAGA complex, a histone acetyltransferase-coactivator complex that is involved in global regulation of gene expression through acetylation and transcription functions
YIL152W/VPR1	Putative protein of unknown function
YKL017C/HCS1	Hexameric DNA polymerase alpha-associated DNA helicase A involved in lagging strand DNA synthesis; contains single-stranded DNA stimulated ATPase and dATPase activities; replication protein A stimulates helicase and ATPase activities
YGR140W/CBF2	Essential kinetochore protein, component of the CBF3 multisubunit complex that binds to the CDEIII region of the centromere; Cbf2p also binds to the CDEII region possibly forming a different multimeric complex, ubiquitinated in vivo
YJR127C/RSF2	Zinc-finger protein involved in transcriptional control of both nuclear and mitochondrial genes, many of which specify products required for glycerol-based growth, respiration, and other functions
YDR375C/BCS1	Mitochondrial protein of the AAA ATPase family; has ATP-dependent chaperone activity; required for assembly of Rip1p and Qcr10p into cytochrome bc(1) complex; mutations in human homolog BCS1L are linked to neonatal mitochondrial diseases
YOR091W/TMA46	Protein of unknown function that associates with translating ribosomes; interacts with GTPase Rbg1p
YLR397C/AFG2	ATPase of the CDC48/PAS1/SEC18 (AAA) family, forms a hexameric complex; is essential for pre-60S maturation and release of several preribosome maturation factors; may be involved in degradation of aberrant mRNAs
YNL132W/KRE33	Essential protein, required for biogenesis of the small ribosomal subunit; heterozygous mutant shows haploinsufficiency in K1 killer toxin resistance
YMR078C/CTF18	Subunit of a complex with Ctf8p that shares some subunits with Replication Factor C and is required for sister chromatid cohesion; may have overlapping functions with Rad24p in the DNA damage replication checkpoint
YLR422W/DCK1	Protein of unknown function with similarity to human DOCK proteins (guanine nucleotide exchange factors); interacts with Ino4p; green fluorescent protein (GFP)-fusion protein localizes to the cytoplasm, YLR422W is not an essential protein

YMR125W/STO1	Large subunit of the nuclear mRNA cap-binding protein complex, interacts with Npl3p to carry nuclear poly(A)+ mRNA to cytoplasm; also involved in nuclear mRNA degradation and telomere maintenance; orthologous to mammalian CBP80
YOR371C/GPB1	Multistep regulator of cAMP-PKA signaling; inhibits PKA downstream of Gpa2p and Cyr1p, thereby increasing cAMP dependency; promotes ubiquitin-dependent proteolysis of Ira2p; regulated by G-alpha protein Gpa2p; homolog of Gpb2p
YMR094W/CTF13	Subunit of the CBF3 complex, which binds to the CDE III element of centromeres, bending the DNA upon binding, and may be involved in sister chromatid cohesion during mitosis
YMR167W/MLH1	Protein required for mismatch repair in mitosis and meiosis as well as crossing over during meiosis; forms a complex with Pms1p and Msh2p-Msh3p during mismatch repair; human homolog is associated with hereditary non-polyposis colon cancer
YDR103W/STE5	Pheromone-response scaffold protein that controls the mating decision; binds Ste11p, Ste7p, and Fus3p kinases, forming a MAPK cascade complex that interacts with the plasma membrane and Ste4p-Ste18p; allosteric activator of Fus3p
YDR318W/MCM21	Protein involved in minichromosome maintenance; component of the COMA complex (Ctf19p, Okp1p, Mcm21p, Ame1p) that bridges kinetochore subunits that are in contact with centromeric DNA and the subunits bound to microtubules
YAL026C/DRS2	Aminophospholipid translocase (flippase) that maintains membrane lipid asymmetry in post-Golgi secretory vesicles; contributes to clathrin-coated vesicle formation and endocytosis; mutations in human homolog ATP8B1 result in liver disease
YDR180W/SCC2	Subunit of cohesin loading factor (Scc2p-Scc4p), a complex required for loading of cohesin complexes onto chromosomes; involved in establishing sister chromatid cohesion during DSB repair via histone H2AX; evolutionarily-conserved adherin
YOR092W/ECM3	Non-essential protein of unknown function; involved in signal transduction and the genotoxic response; induced rapidly in response to treatment with 8-methoxypsoralen and UVA irradiation
YDR235W/PRP42	U1 snRNP protein involved in splicing, required for U1 snRNP biogenesis; contains multiple tetriatricopeptide repeats
YER151C/UBP3	Ubiquitin-specific protease that interacts with Bre5p to co-regulate anterograde and retrograde transport between the ER and Golgi; inhibitor of gene silencing; cleaves ubiquitin fusions but not polyubiquitin; also has mRNA binding activity
YMR275C/BUL1	Ubiquitin-binding component of the Rsp5p E3-ubiquitin ligase complex, functional homolog of Bul2p, disruption causes temperature-sensitive growth, overexpression causes missorting of amino acid permeases
YKL114C/APN1	Major apurinic/apyrimidinic endonuclease, 3'-repair diesterase involved in repair of DNA damage by oxidation and alkylating agents; also functions as a 3'-5' exonuclease to repair 7,8-dihydro-8-oxodeoxyguanosine
YOL081W/IRA2	GTPase-activating protein that negatively regulates RAS by converting it from the GTP- to the GDP-bound inactive form, required for reducing cAMP levels under nutrient limiting conditions, has similarity to Ira1p and human neurofibromin
YPR049C/ATG11	Adapter protein for pexophagy and the cytoplasm-to-vacuole targeting (Cvt) pathway; directs receptor-bound cargo to the phagophore assembly site (PAS) for packaging into vesicles; required for recruiting other proteins to the (PAS)
YGL095C/VPS45	Protein of the Sec1p/Munc-18 family, essential for vacuolar protein sorting; required for the function of Pep12p and the early endosome/late Golgi SNARE Tlg2p; essential for fusion of Golgi-derived vesicles with the prevacuolar compartment
YDR456W/NHX1	Na+/H+ and K+/H+ exchanger, required for intracellular sequestration of Na+ and K+; located in the vacuole and late endosome compartments; required for osmotolerance to acute hypertonic shock and for vacuolar fusion
YKL197C/PEX1	AAA-peroxin that heterodimerizes with AAA-peroxin Pex6p and participates in the recycling of peroxisomal signal receptor Pex5p from the peroxisomal membrane to the cystosol; induced by oleic acid and upregulated during anaerobiosis
YIL068C/SEC6	Essential 88kDa subunit of the exocyst complex, which mediates polarized targeting of secretory vesicles to active sites of exocytosis; dimeric form of Sec6p interacts with Sec9p in vitro and inhibits t-SNARE assembly

YOR326W/MYO2	One of two type V myosin motors (along with MYO4) involved in actin-based transport of cargos; required for the polarized delivery of secretory vesicles, the vacuole, late Golgi elements, peroxisomes, and the mitotic spindle
YNR045W/PET494	Mitochondrial translational activator specific for the COX3 mRNA, acts together with Pet54p and Pet122p; located in the mitochondrial inner membrane
YJR107W/LIH1	Putative protein of unknown function; has sequence or structural similarity to lipases
YPL268W/PLC1	Phospholipase C, hydrolyzes phosphatidylinositol 4,5-biphosphate (PIP2) to generate the signaling molecules inositol 1,4,5-triphosphate (IP3) and 1,2-diacylglycerol (DAG); involved in regulating many cellular processes
YJL062W/LAS21	Integral plasma membrane protein involved in the synthesis of the glycosylphosphatidylinositol (GPI) core structure; mutations affect cell wall integrity
YCR042C/TAF2	TFIID subunit (150 kDa), involved in RNA polymerase II transcription initiation

1022
1023 **Supplementary Table 9. Annotations of top hit loci from barcoded RH-seq of thermotolerance.**
1024 Shown are hits from thermotolerance mapping by barcoded RH-seq (Table S7) that met quality control
1025 thresholds and at which disruption of the *S. cerevisiae* allele compromised thermotolerance to a greater
1026 extent than did disruption of the *S. paradoxus* allele in the interspecific hybrid.

gene	D_s	D_n	P_s	P_n	NI	*p*	adjusted *p*
YBR136W/MEC1	293	38	770	363	3.63	5.71×10^{-15}	1.69×10^{-12}
YLR422W/DCK1	122	26	335	280	3.92	1.13×10^{-10}	8.70×10^{-9}
YOR326W/MYO2	35	0	1486	1273	inf	5.15×10^{-10}	2.77×10^{-3}
YAL026C/DRS2	32	1	1399	1595	36.48	5.81×10^{-10}	2.97×10^{-8}
YMR207C/HFA1	77	16	2215	2103	4.57	6.16×10^{-10}	3.08×10^{-8}
YOR371C/GPB1	79	18	347	323	4.09	1.42×10^{-8}	4.02×10^{-7}
YML099C/ARG81	96	16	351	247	4.22	1.61×10^{-8}	4.43×10^{-7}
YJL062W/LAS21	74	10	170	131	5.70	2.19×10^{-8}	5.77×10^{-7}
YPL268W/PLC1	112	16	245	146	4.17	4.57×10^{-8}	1.05×10^{-6}
YCR042C/TAF2	155	45	661	470	2.45	1.70×10^{-7}	2.90×10^{-6}
YPL254W/HFI1	76	10	150	106	5.37	1.66×10^{-7}	2.90×10^{-6}
YPR049C/ATG11	131	45	448	395	2.57	1.67×10^{-7}	2.90×10^{-6}
YNL049C/SFB2	32	1	773	587	24.30	3.22×10^{-7}	4.88×10^{-6}
YKL114C/APN1	44	3	120	87	10.63	9.80×10^{-7}	1.19×10^{-5}
YIL068C/SEC6	80	5	271	111	6.55	1.36×10^{-6}	1.52×10^{-5}
YGR198W/YPP1	79	25	295	282	3.02	1.97×10^{-6}	2.10×10^{-5}
YDR375C/BCS1	67	4	107	54	8.45	2.20×10^{-6}	2.28×10^{-5}
YKL017C/HCS1	78	13	219	143	3.92	3.56×10^{-6}	3.28×10^{-5}
YDR235W/PRP42	61	7	177	109	5.37	5.34×10^{-6}	4.57×10^{-5}
YDR180W/SCC2	173	72	523	431	1.98	6.47×10^{-6}	5.35×10^{-5}
YMR167W/MLH1	94	26	286	209	2.64	2.47×10^{-5}	1.54×10^{-4}
YKL197C/PEX1	132	58	415	351	1.92	1.55×10^{-4}	6.57×10^{-4}
YMR078C/CTF18	73	26	295	260	2.47	1.63×10^{-4}	6.82×10^{-4}
YGL095C/VPS45	67	14	202	122	2.89	3.72×10^{-4}	1.30×10^{-3}
YMR094W/CTF13	54	21	152	158	2.67	4.44×10^{-4}	1.50×10^{-3}
YNL132W/KRE33	75	4	187	52	5.21	5.13×10^{-4}	1.69×10^{-3}
YDR103W/STE5	93	56	344	373	1.80	1.57×10^{-3}	4.16×10^{-3}
YOR092W/ECM3	16	1	648	498	12.30	1.92×10^{-3}	4.86×10^{-3}
YNR045W/PET494	61	20	177	136	2.34	2.15×10^{-3}	5.32×10^{-3}
YJR107W/LIH1	19	1	102	61	11.36	2.40×10^{-3}	5.79×10^{-3}
YLR397C/AFG2	102	27	288	155	2.03	2.56×10^{-3}	6.10×10^{-3}
YGL082W/MIY1	41	8	132	84	3.26	2.60×10^{-3}	6.17×10^{-3}
YOR091W/TMA46	40	6	103	46	2.98	0.0212	0.0351
YDR456W/NHX1	87	13	187	59	2.11	0.0278	0.0443
YDR508C/GNP1	3	0	959	797	inf	0.2562	0.3009
YIL152W/VPR1	19	11	70	59	1.46	0.4184	0.4669

**1029 Supplementary Table 10. Whole-gene tests for evidence of non-neutral protein evolution at
thermotolerance loci.** Each row reports results from the McDonald-Kreitman test on sequences from
strains of European populations of *S. cerevisiae* and *S. paradoxus* of the indicated top hit from barcoded
RH-seq mapping of thermotolerance. D_s, number of sites of synonymous nucleotide divergence between
species; number of sites of D_n, nonsynonymous nucleotide divergence between species; P_s, number of

sites of synonymous nucleotide polymorphisms within species; P_n, number of sites of nonsynonymous
nucleotide polymorphisms within species. NI, neutrality index. The sixth column reports the p-value from a
Fisher's exact test on D_s, D_n, P_s, and P_n, and the seventh column reports the adjusted p-value after
applying the Benjamini-Hochberg correction for multiple hypothesis testing. All loci exhibited NI > 1,
corresponding to a dearth of divergent amino acid changes relative to synonymous changes and
polymorphisms.

A

Tn-seq: map barcode to genomic location

grow hemizygote pool in standard conditions

extract genomic DNA

Amplify with transposon-specific and adapter-specific primers and sequence

Bar-seq: quantify fitness of mutants by sequencing barcodes

grow hemizygote pool in condition of interest

extract genomic DNA

Amplify barcodes and sequence

B

2-3 generations

Frozen

Fresh (T0)

High temperature

Low temperature

x 12

x 12

backdilute for ~10 generations

Extract genomic DNA
Amplify barcodes

A

YGR198W/YPP1

YMR207C/HFA1

YGL082W/MIY1

YNL049C/SFB2

YDL035C/GPR1

YDR508C/GNP1

All plots: x-axis $\log_2(37°C/28°C)$, y-axis Density of Inserts

- Insert in *S. paradoxus*
- Insert in *S. cerevisiae*

B

Yeast cell diagram with compartments:

Golgi Body
DRS1

Bud
SEC6

Vacuole
VPS45

Nucleus
RSF2   HFI1
STE5   CBF2   STO1
APN1   CTF18   **SCC2**
MIY1   MCM21   **TAF2**
KRE33   MEC1   CTF13
ARG81   PRP42   MLH1

Bud Neck
MYO2

Cytoplasm
HSC1   GBP1
TMA46   UBP3
AFG2   BUL1
DCK1   IRA2

Punctae Composites
SFB2
PEX1
ATG11
YPP1

Endoplasmic Reticulum
ECM3   LAS21

Mitochondrion
HFA1   PET494
GNP1   NHX1
BCS1

Plasma Membrane
GPR1