1

Trehalose increases tomato drought tolerance, induces defenses, and increases resistance to bacterial wilt disease

April M. MacIntyre[1,2,6], Valerian Meline[3], Zachary Gorman[4], Steven P. Augustine[5], Carolyn J.
Dye[1,7], Corri D. Hamilton[1], Anjali S. Iyer-Pascuzzi[3], Michael V. Kolomiets[4], Katherine A.
McCulloh[5], and Caitilyn Allen[1*]

*Corresponding author: callen@wisc.edu   +1-608-556-3369

[1]University of Wisconsin-Madison Department of Plant Pathology
[2]University of Wisconsin-Madison Department of Bacteriology
[3]Purdue University, Department of Botany and Plant Pathology
[4]Texas A&M University, Department of Plant Pathology and Microbiology
[5] University of Wisconsin-Madison Department of Botany
[6] Current address: Valent Biosciences, Libertyville, IL
[7] Current address: Soil Carbon Co., Minneapolis-St. Paul, MN

Contributions: AMM and CA designed the experiments and wrote the text with contributions from VM, AIP, and ZG. AMM carried out most of the experiments and created most of the figures. VM performed the meta-transcriptomic analysis. ZG performed the xylem sap metabolite analysis. SPA advised and assisted with LI-COR experiments and created the trehalose stomatal conductance, photosynthesis, transpiration, and water efficiency graphs. CJD assisted with trehalose physiology, ABA treatment, and *Xanthomonas* experiments. CDH advised on xylem sap collection. AIP, MK, CDH, and KM provided feedback and edited the manuscript.

Abstract

Ralstonia solanacearum causes bacterial wilt disease, leading to severe crop losses. Xylem sap

from *R. solanacearum*-infected tomato is enriched in the disaccharide trehalose. Water-stressed plants

also accumulate trehalose, which increases drought tolerance via abscisic acid (ABA) signaling. Because

R. solanacearum-infected plants suffer reduced water flow, we hypothesized that bacterial wilt

physiologically mimics drought stress, which trehalose could mitigate. We found that *R. solanacearum*-

infected plants differentially expressed drought-associated genes, including those involved in ABA and

trehalose metabolism, and had more ABA in xylem sap. Consistent with this, treating tomato roots with

ABA reduced both stomatal conductance and stem colonization by *R. solanacearum*. Treating roots with

trehalose increased xylem sap ABA and reduced plant water use by lowering stomatal conductance and

temporarily improving water use efficiency. Trehalose treatment also upregulated expression of salicylic

acid (SA)-dependent tomato defense genes; increased xylem sap levels of SA and other antimicrobial

compounds; and increased bacterial wilt resistance of SA-insensitive *NahG* tomato plants. Additionally,

trehalose treatment increased xylem concentrations of jasmonic acid and related oxylipins. Finally,

trehalose-treated plants were substantially more resistant to bacterial wilt disease. Together, these data

show that exogenous trehalose reduced both water stress and bacterial wilt disease and triggered

systemic disease resistance, possibly through a Damage Associated Molecular Pattern (DAMP) response

pathway. This suite of responses revealed unexpected linkages between plant responses to biotic and

abiotic stress and suggested that *R. solanacearum*-infected plants increase trehalose to improve water

use efficiency and increase wilt disease resistance. The pathogen may degrade trehalose to counter

these efforts. Together, these results suggest that treating tomatoes with exogenous trehalose could be

a practical strategy for bacterial wilt management.

Keywords: *Ralstonia solanacearum,* bacterial wilt disease, drought tolerance, trehalose, plant

hormones, xylem water physiology

Introduction

Ralstonia solanacearum is a vascular plant pathogen that causes bacterial wilt disease in over

200 species of plants worldwide (Denny 2006). It is widely studied because it is economically damaging

to food, fiber, and feed production (Elphinstone 2005, Mansfield, Genin et al. 2012). *R. solanacearum* is

also notoriously difficult to eradicate because it persists long-term in soil, water, and latently infected

hosts (Álvarez, Bosca et al. 2010).

Most *R. solanacearum* strains infect plants through the soil (Elphinstone 2005). *R. solanacearum*

chemo- or aero-taxes towards roots, evades root border cell traps, enters roots at points of lateral root

emergence, moves to the root cortex, and eventually migrates to the xylem, all with the help of secreted

proteins such as hydrolytic enzymes and Type III secreted effectors (Vasse, Frey et al. 1995, Tans-

Kersten, Huang et al. 2001, Liu, Zhang et al. 2005, Yao and Allen 2006, Yao and Allen 2007, Digonnet,

Martinez et al. 2012, Deslandes and Genin 2014, Tran, MacIntyre et al. 2016). Once in xylem vessels, *R.*

solanacearum forms biofilm matrices composed of extracellular DNA and polysaccharide (EPS) and

proliferates to >10^9 CFU/g of stem tissue (Orgambide, Montrozier et al. 1991, Mori, Inoue et al. 2016,

Tran, MacIntyre et al. 2016). *R. solanacearum* also conditions the xylem environment to favor bacterial

growth (Lowe-Power, Hendrich et al. 2018). The large amounts of bacteria and EPS clog xylem vessels

and disrupt water flow (Ingel, Caldwell et al. 2021). If the plant cannot contain bacterial growth and

spread, it will wilt and eventually die.

Researchers long assumed that *R. solanacearum* and other vascular pathogens cause drought-

like physiological symptoms in host plants and that severity of the disease is affected by environmental

water availability (Abd El-Rahim, Fahmy et al. 1998, McElrone, Sherald et al. 2003, Daugherty, Lopes et

al. 2010). *R. solanacearum* dies more quickly in drought-stressed chickpea plants, and *R. solanacearum*

infection and water stress induce similar osmolyte biosynthesis and expression of genes related to the

plant signaling molecule abscisic acid (ABA), a well-characterized marker of water stress (Sinha, Gupta et

al. 2016, Sinha, Gupta et al. 2017). Further, stomatal conductance decreases in *R. solanacearum* infected

plants (Lowe-Power, Hendrich et al. 2018). These studies suggest but do not prove that plant response

to bacterial wilt mimics the drought stress response.

During water and osmotic stress, many plants, including tomatoes, accumulate trehalose and

trehalose-6-phosphate (T6P), which contribute to tolerance of these stresses (Romero, Bellés et al.

1997, Garg, Kim et al. 2002, Cortina and Culiáñez-Macià 2005, Lyu, Min et al. 2012, Khater, Dawood et

al. 2018). Xylem sap from *R. solanacearum*-infected tomato plants contains about 20-fold more

trehalose than healthy plants (Lowe-Power, Hendrich et al. 2018). This non-reducing disaccharide

composed of glucose and UDP-glucose is best known for its role as a compatible solute. Trehalose

synthesis helps *R. solanacearum* tolerate osmotic stress and contributes to pathogen fitness and

virulence (MacIntyre, Barth et al. 2019). However, the tomato host produces the trehalose in xylem sap

during infection; *R. solanacearum* does not export its trehalose in culture and it aggressively expresses

the secreted trehalase TreA *in planta* (Jacobs, Babujee et al. 2012).

The roles of trehalose and T6P as nutritional signaling metabolites in plants have been reviewed

(Lunn 2007, Paul 2007, Iordachescu and Imai 2008, Paul, Primavesi et al. 2008, Lunn, Delorge et al.

2014). In *Arabidopsis,* most trehalose metabolism-related proteins have predicted functions in

development or signaling (Vandesteene, Ramon et al. 2010, Vandesteene, Lopez-Galvis et al. 2012). The

current paradigm is that T6P, and by extension trehalose, are direct indicators of sucrose levels that help

plants maintain sucrose-to-starch ratios through SnRK1 signaling (Schluepmann, Pellny et al. 2003,

Kolbe, Tiessen et al. 2005, Lunn, Feil et al. 2006, Schluepmann, Berke et al. 2012). Trehalose also likely

improves drought tolerance in plants by two mechanisms: by regulating carbohydrate levels during

stress conditions, possibly via the SnRK1 kinase; and by altering ABA signaling and stomatal conductance

(Garg, Kim et al. 2002, Paul, Gonzalez-Uriarte et al. 2018, Yu, Zhao et al. 2019). Members of another

kinase family, SnRK2, also integrate with ABA signaling, and trehalose metabolism may connect to this

network via the TOR regulator (Signorelli, Tarkowski et al. 2019, Chen, Li et al. 2020).

Some studies suggest that trehalose may mediate host-microbe interactions and contribute to

plant defense. For example, treating *Arabidopsis* seedlings in liquid culture with 30 mM exogenous

trehalose triggered the expression of defense genes in ethylene and methyl-jasmonate pathways (Bae,

Herman et al. 2005). Infusing tobacco leaves with a 10-50 mM trehalose solution increased resistance to

tobacco mosaic virus and triggered a transcriptomic response suggestive of biotic and abiotic stress

tolerance (Shi, Sun et al. 2019). A *Xanthomonas citri* trehalose synthesis mutant had reduced fitness and

infusing citrus leaves with a trehalose solution induced defense gene expression, suggesting that

trehalose might simultaneously be a virulence factor and an indicator of pathogen attack (Piazza, Zimaro

et al. 2015). In another study, spraying tomato leaves with the trehalase inhibitor Validamycin A

increased resistance to bacterial wilt and *Fusarium* wilt and activated systemic acquired resistance

against a wide range of other pathogens in *Arabidopsis* (Ishikawa, Fujimori et al. 1996, Ishikawa,

Shirouzu et al. 2005, Bian, Duan et al. 2020). Additionally, silencing some tomato trehalose-6-phosphate

synthase genes decreased resistance to *Botrytis cinerea* and *P. syringae* pv. *tomato* DC3000 and affected

expression of JA, ET, and SA signaling responsive, defense related genes (Zhang, Hong et al. 2016).

We investigated the relationship between bacterial wilt and drought stress in tomatoes and

explored the role of trehalose in this interaction. We tested the following hypotheses: 1) *R.*

solanacearum infection mimics abiotic stress triggered by disrupted water flow; and 2) *Exogenous*

trehalose inhibits bacterial wilt development in tomatoes by enhancing plant water use efficiency and

*inducing defense response*s. To test H_o1, we used meta-transcriptomic analyses to show that tomato

genes involved in trehalose metabolism and water stress responses are differentially expressed during

R. solanacearum infection in susceptible tomato hosts. Using ABA treatment to close stomates, we

found that *R. solanacearum* depends on stomatal conductance and water flow for effective host

colonization. We also discovered that *R. solanacearum* infection increases xylem sap ABA levels.

Testing H_o2 revealed that treating plants with exogenous trehalose improved water use

efficiency in drought-stressed tomatoes, decreased short-term stomatal conductance and transpiration,

and increased ABA concentration in xylem sap. Trehalose-treated tomato plants were more resistant to

bacterial wilt. This protective effect appeared to be mediated by up-regulation of salicylic acid (SA)- and

jasmonic acid (JA)-related defense genes and by increased levels of jasmonates and other defense

compounds in xylem sap.

Results

Bacterial wilt disease altered expression of tomato trehalose and water stress genes

To investigate the relationship between bacterial wilt and drought stress, we measured

expression of all tomato genes annotated as related to trehalose metabolism, ABA, or water stress

responses. Candidate genes included those known or predicted to encode synthesis and degradation of

trehalose, including trehalose-6-phosphate phosphatase (TPP), trehalose-6-phosphate synthase (TPS),

and trehalase (Leyman, Van Dijck et al. 2001, Vandesteene, Ramon et al. 2010, Vandesteene, Lopez-

Galvis et al. 2012). Genes encoding water stress-associated proteins included: 1) those involved in

biosynthesis, regulation, and response to ABA, a known indicator of water stress (Song, Huang et al.

2016, Zhou, Yu et al. 2019); 2) late embryogenesis/dehydrins (LEAs), which accumulate in water-stressed

plants (Cao and Li 2015); and 3) aquaporins, channel proteins that allow living cells to move water and

small solutes more efficiently (Kapilan, Vaziri et al. 2018).

We used four datasets that profiled transcriptomes of two wilt-susceptible tomato plants

(WV700 and Bonny Best) responding to infection by *R. solanacearum* strain K60 or GMI1000 (French,

Kim et al. 2018). After normalizing the datasets to allow comparisons across these experiments, we

measured the expression of these candidate genes in healthy plants relative to their levels in *R.*

solanacearum-infected plants.

The datasets came from diverse experiments involving different plant genotypes, tissues, and *R.*

solanacearum strains. Despite considerable experiment-associated variation across datasets, this

analysis revealed that *R. solanacearum*-infected tomato plants significantly changed the expression of

several water stress physiology genes (Fig. 1). Nine genes meeting the annotation criteria were

significantly differentially expressed across all datasets. Notably, the trehalose-6-phosphate synthase

gene *SlTPS7* was upregulated in all of the susceptible tomatoes in response to *Rs,* suggesting this may be

the main T6P protein responsible for elevated trehalose levels in xylem sap during infection. All infected

plants upregulated an ABA hydroxylase and an ABA- and environmental stress-inducible dehydrin

protein TAS14 and downregulated many diverse aquaporins, consistent with transcript changes

observed in plants undergoing abiotic water stress. These results led us to directly measure

physiological and metabolic indicators of drought stress in *R. solanacearum*-infected plants.

Figure 1. *R. solanacearum* infection consistently changes expression of nine tomato genes involved in

trehalose metabolism and water stress physiology. Meta-analysis of transcriptomic data from four

different experiments measuring gene expression of tomato plants responding to infection by *R.*

solanacearum identified nine genes of interest that were commonly differentially expressed across all

data sets. The heatmaps show relative expression (as log$_2$ fold change) compared to mock-inoculated

control plants of all tomato genes that were both: 1) Annotated with the terms *trehalose*, *drought*, or

abscisic acid / ABA; and 2) Identified by the meta-transcriptomic analysis as significantly differentially

expressed across four different tomato RNAseq studies. The four studies were: **S1**, roots of wilt-

susceptible West Virginia 700 tomato plants (WV) sampled 24 h after infection with *R. solanacearum*

strain K60; **S2**, roots of WV sampled 48 h after infection with *R. solanacearum* strain K60; **S3**, seedling

roots of wilt-susceptible cv. Bonny Best (BB) sampled 24 h after infection with *R. solanacearum* strain

GMI1000; **S4**, mid-stems of BB sampled 72 h after inoculation with *R. solanacearum* strain GMI1000.

Tomato locus tags, gene names, and PANTHER families are from NCBI and the Sol Genomics Network,

accessed in 03/2020. Transcriptomic data were normalized across experiments to allow comparisons as

described in Methods.

Bacterial wilt disease increased ABA concentration in xylem sap

Plants can use ABA to mitigate drought stress. This plant hormone, synthesized locally in guard

cells or the vasculature, closes stomates to reduce water loss through transpiration (Raghavendra,

Gonugunta et al. 2010). To test the hypothesis that tomato plants produce ABA in response to bacterial

wilt, we used LCMS/MS to compare ABA levels in xylem sap of healthy plants and *R. solanacearum*-

inoculated plants showing early wilt symptoms. Xylem sap from the *R. solanacearum*-infected plants

contained 40% more ABA than sap from mock-inoculated plants (Fig. 2A).

Figure 2. ABA concentration increased in *R. solanacearum*-infected plants, and ABA treatment

reduced stomatal conductance, which correlated with reduced *R. solanacearum* stem colonization.

A) ABA concentration in xylem sap harvested from healthy and *R. solanacearum*-infected Bonny Best

tomato plants showing early wilt symptoms (1-25% of leaf area wilted) (T-test, *P=.03). Bars show

mean and standard deviation of xylem sap ABA levels, measured by GC-MS, from six plants per

treatment. **B)** Stomatal conductance in tomato leaves as measured with LI-COR instrument.

Conductance of plants watered with 50 or 100 uM of ABA differed from that of mock-inoculated

controls at all time points (2-way ANOVA Tukey's multiple comparisons, *P<.05). Data represent four

biological replicate experiments, each with three plants analyzed per treatment per time point. Bars

represent standard deviation. **C)** *R. solanacearum* population sizes as quantified in tomato plant stems

five days after soil-soak inoculation with *R. solanacearum* strain GMI1000. *R. solanacearum*

colonization was lower in ABA-treated plants at 5 dpi, which corresponds to 168 h of ABA treatment

(Kruskal Wallis, 50 uM ABA, ***P<.001; 100 uM ABA, **P<.01). The proportion of ABA-treated plants

containing detectable *R. solanacearum* cells (shown below X-axis labels) was also lower (Chi-square,

*P=.0016). Dots represent *R. solanacearum* population sizes in individual plants. Black bars represent

the geometric mean and standard deviation of the data. Plants with undetectable *R. solanacearum*

populations were given a value of 1 for statistical analysis. Data represent three biological replicate

experiments, each with 12 plants per treatment (36 total plants/treatment).

***R. solanacearum* depends on host stomatal conductance to establish infection**

We previously observed that *R. solanacearum*-infected tomatoes had reduced stomatal

conductance (Lowe-Power, Khokhani et al. 2018). To test the hypothesis that the increased ABA levels

and lowered stomatal conductance in infected plants also affect bacterial wilt disease development, we

measured the effects of exogenous ABA treatment on stomatal conductance, transpiration rates, and *R.*

solanacearum colonization of susceptible tomato plant stems. We watered tomato plants daily with 50

mL of either 100 µM ABA, 50 µM ABA, or water as control and measured stomatal conductance four to

six hours later. After 48 h of ABA treatment, plants were drench-inoculated by adding a suspension of *R.*

solanacearum strain GMI1000 to the ABA treatment solutions, which we determined were neither

nutritious nor cytotoxic to *R. solanacearum* (Supp. Fig. 1). Treatment with ABA reliably reduced stomatal

conductance in the plants over the seven days of the experiment; 100 µM and 50 µM ABA treatment

had similar effects (Fig. 2B).

ABA treatments continued for four days after plants were inoculated with *R. solanacearum*. At

five dpi we measured *R. solanacearum* population sizes in the stems. Stems of ABA-treated plants were

less frequently colonized by *R. solanacearum* than control plants (50 µM ABA = 17% plants colonized;

100 µM = 23% plants colonized; water control = 53% plants colonized) and they contained lower *R.*

solanacearum population sizes than untreated plants (Fig. 2C). The correlation between ABA treatment,

reduced stomatal conductance, and lower *R. solanacearum* colonization in tomato stems suggests that

plant water movement assists *R. solanacearum* infection either at the roots or in the stem and that

reducing host water flow reduces or slows *R. solanacearum* spread in the plant.

Trehalose treatment transiently improved water use efficiency in tomatoes

Tomato plants become more drought tolerant following exposure to exogenous trehalose via

ABA signaling and stomatal closure (Yu, Zhao et al. 2019). Because ABA and trehalose concentrations

increased in xylem sap from *R. solanacearum*-infected plants and pathogen spread depends on host

xylem water transport, we asked whether exogenous trehalose could also protect plants from bacterial

wilt.

We first confirmed that trehalose increases tomato drought tolerance under our experimental

conditions. Two-week-old wilt-susceptible Bonny Best tomato plants were treated with 50 mL of 30 mM

trehalose solution or water and then were not watered for the rest of the experiment. Trehalose-

treated plants remained turgid for three days after the water-treated control plants began wilting,

showing that trehalose treatment delayed wilting from lack of water under our conditions (Fig. 3A).

Quantitative mass spectrometry analysis of tomato xylem sap revealed that ABA content was 5-fold

higher in trehalose-treated plants at 24 h (Fig. 3B). However, at 6 and 48 h after treatment, ABA levels

were no different from those of water-treated control plants. Gravimetric assays quantifying water loss

in trehalose-treated plants compared to water controls confirmed that trehalose-treated plants lose

water at a slower rate due to transpiration than controls (data not shown).

Figure 3. Trehalose treatment delayed tomato wilting due to drought, likely by transiently

increasing ABA content in xylem sap and by decreasing stomatal conductance and transpiration. A)

Curves show how long it took plants to wilt following a single treatment with either water (blue) or

30 mM trehalose (green) and then no water for two weeks (*P*=.01, T-test of areas under the curve).

Data shown represent four biological replicates totaling 38 plants per treatment. **B)** ABA

concentration in trehalose treated plants (one-way ANOVA Fisher's LSD multiple comparisons to

water-treated control: 6 h tre *P*=.63; 24 h tre *P*<.0001; 48 h tre *P*=.76). Water controls were collected

concurrently with the treated samples and averaged. Data shown represent xylem sap samples from

six plants.   **C)** Transpiration and **D)** stomatal conductance were measured with LI-COR. Asterisks

indicate *P*>.05, mixed-model ANOVA, Tukey's HSD. Data shown represent 20 plants/treatment.

Measurements were taken 10 min after treatment with trehalose or water, then each day after that

in the morning two hours after growth chamber lights went on. Shaded gray vertical bars on the

graphs represent night periods. Bars on the data points represent the standard deviation. All data

were calculated from the same LICOR time points.

Next, we assessed stomatal conductance, transpiration, and photosynthesis of trehalose-treated plants.

We measured tomato leaf responses with a LI-COR portable photosynthesis machine at 10 min, 6, 24,

48, and 72 h after treatment with 30mM trehalose or water. Transpiration in trehalose-treated plants

was lower than in controls at 6, 24, and 48 h (Fig. 3B). Relative to untreated plants, trehalose treatment

detectably reduced stomatal conductance after just 10 min, an effect that persisted for at least 6 h. By

24 h after treatment, stomatal conductance of treated plants was indistinguishable from that of water

controls (Fig. 3C). Water use efficiency, the relationship between water loss and plant productivity

(photosynthesis/transpiration), was temporarily improved at 6 h by trehalose treatment (Supp. Fig. 2A).

Trehalose treatment did not negatively affect photosynthesis (Supp. Fig. 2B).

We found that tomato plant growth was not significantly altered by trehalose treatment and the

leaf relative water content was the same in trehalose-treated and water-treated plants at 48h (Supp.

Fig. 2C). Further, 1-3 days of watering with trehalose solution did not affect tomato seedling root growth

in vitro relative to water controls (Supp. Fig. 2D). Chlorophyll content, measured by SPAD meter, was

transiently lower in trehalose treated plants 2-10 days after treatment, although it eventually returned

to the same levels as in untreated plants (Supp. Fig. 2E). Together, these diverse physiological and

biochemical analyses showed that treating tomato plants with exogenous trehalose temporarily

increased their drought tolerance, likely because trehalose increased ABA levels and ABA signaling

closes stomates.

Trehalose is a potent inhibitor of bacterial wilt disease

After finding that bacterial wilt impacts host water stress physiology and that exogenous

trehalose reduces stomatal conductance in tomatoes, we wondered if the decreased water movement

in trehalose-treated plants would slow bacterial wilt disease development. We therefore tested the

hypothesis that trehalose makes plants more wilt resistant. Forty-eight hours after plants had been

treated with 30 mM trehalose solution or water, they were drench inoculated with a suspension of *R.*

solanacearum bacteria. Bacterial wilt incidence and severity was reduced by 25-75% across biological

replicates in the trehalose-treated plants, demonstrating that exogenous trehalose increases tomato

resistance to bacterial wilt (Fig. 4A).

Figure 4. Treating plants with trehalose at the root delayed bacterial wilt symptoms and reduced *R. solanacearum* stem colonization, but only following soil inoculation. A) Wilt disease progress of cv. Bonny Best tomato plants treated with trehalose, then after 48 h soil soak inoculated with *R. solanacearum* GMI1000. Disease symptoms were rated on a 0-4 disease index scale where 0=no wilt, and 4=76-100% wilted or dead. Trehalose treatment delayed appearance of wilt symptoms (*P*=.0082, t-test comparing areas under the curves or AUC). Data represent three replicate experiments, each with 15 plants/treatment. Bars represent standard error. **B)** Bacterial population sizes were smaller in stems of trehalose-treated plants at 5-7 days after inoculation (Mann-Whitney, *P*=.0009). Data represent three biological replicates with 15 plants/day sampled/treatment; each dot represents the *R. solanacearum* population size in one plant (including plants without detectable *R. solanacearum*). Black bars represent geometric mean and standard deviation of the data. The percentage of plants that contained any detectable *R. solanacearum* cells is shown below the X-axis; the limit of detection is ~10^2 cells. Relative to controls, fewer trehalose-treated plants contained detectable *R. solanacearum* populations (Chi-square, *P*=.0016). C**)** Disease progress of plants treated with trehalose (green) or water (blue), then 48 h later inoculated by introducing *R. solanacearum* directly into the xylem through a cut leaf petiole. Trehalose treatment did not affect wilt disease development in petiole-inoculated plants (*P*=.2067, T-test of AUC). Data represent three replicate experiments, each containing 15 plants/treatment. Bars represent standard error. **D)** Disease progress of plants treated with 30 mM trehalose or water, then 120 h later inoculated with *R. solanacearum* the soil drench

method (*P*=.0003, T-test of AUC) Data represent four replicate experiments, each containing 15
plants/treatment. Bars represent standard error of the mean.

Since *R. solanacearum* can infect plants without causing visible symptoms, the absence of wilt in
trehalose treated plants could indicate either the lack of disease or latent infection. To distinguish
between these possibilities, we measured *R. solanacearum* colonization of tomato midstems at 5, 6, and
7 dpi. Water-treated control plants were three times more likely to contain detectable *R. solanacearum*
cells than trehalose-treated plants (20.7% compared to 7.4%), and control plants contained significantly
higher bacterial population sizes (Fig. 4B). This showed that trehalose treatment did not increase the
frequency of latent infections but rather that *R. solanacearum* was less likely to colonize stems of
trehalose treated plants. The reduced colonization frequency at this relatively early time point is
consistent with the observed trehalose-associated delay in disease development.
We tested the protective limits of trehalose treatment by varying inoculation sites, treatment
methods, and pathogens. Interestingly, inoculating *R. solanacearum* cells directly into the xylem through
a cut petiole overwhelmed any protective benefit of trehalose treatment (Fig. 4C). This result suggests
that *R. solanacearum* must overcome a trehalose-enhanced checkpoint at or in tomato roots to colonize
plant stems. However, the protective effect of root drenching is enduring; trehalose treatment reduced
bacterial wilt incidence even when plants were inoculated five days later (Fig. 4D).
R. solanacearum produces a trehalase enzyme and can use trehalose as a carbon source, so we
wondered if treating tomato roots with trehalose nutritionally enriched the soil environment, thereby
de-incentivizing *R. solanacearum* root invasion. We tested this hypothesis by inoculating trehalose-
treated tomato plants with either wild-type *R. solanacearum* or a Δ*treA* trehalase mutant that cannot
grow on trehalose (MacIntyre, Barth et al. 2019). Trehalose treatment similarly reduced disease caused
by wild type and Δ*treA R. solanacearum*, so bacterial catabolism of trehalose cannot explain trehalose
treatment's protective effect (Suppl. Fig. 3A).
Spraying trehalose solution onto leaves and then inoculating *R. solanacearum* by soil-soak 48 h
later also did not affect bacterial wilt progress (Supp. Fig. 3B). To explore the effects of trehalose
treatment on another bacterial pathogen, we dip-inoculated tomato leaves with the tomato leaf spot
pathogen *Xanthomonas gardneri* 72 h after plant roots were treated with trehalose. Trehalose had no
significant effect on leafspot disease severity or *X. gardneri* colonization of leaves sampled 3, 6, and 9
dpi (Supp. Fig. 3C). These experiments demonstrated that trehalose needs to be applied at the roots to
protect against bacterial wilt and that trehalose might increase resistance only to *R. solanacearum.*

340   **Exogenous trehalose increases the expression of tomato defense genes, and SA signaling may mediate**

341   **its protective effect**

342   Although trehalose treatment of roots decreased plant transpiration and stomatal conductance

343   temporarily (6-48 h), this treatment protected plants from bacterial wilt for at least five days (Fig. 4D),

344   suggesting that trehalose induced plant defense responses. We tested this by quantifying the effect of

345   trehalose treatment on defense gene expression, bacterial wilt resistance of defense hormone-

346   insensitive tomato plants, and levels of hormones and defense compounds.

347   To measure the effect of trehalose on defense gene expression, we extracted RNA from tomato

348   stem sections 6 or 48 h after roots were treated with 30 mM trehalose solution or water. We measured

349   expression of two JA-responsive genes, *PIN2* and *LOXA* (Peña-Cortés, Fisahn et al. 1995, Zhao, Thilmony

350   et al. 2003). *PIN2* was significantly downregulated 6 h after trehalose treatment (Fig. 5A), suggesting JA

351   levels may have been affected by trehalose treatment, while *LOXA* expression did not change after

352   trehalose treatment. G*LUA*, *PR1A*, and *PR1B* are tomato pathogenesis-related (PR) genes induced by SA

353   and by pathogen challenge (Joosten and De Wit 1989, Tornero, Gadea et al. 1997, Block, Schmelz et al.

354   2005). Both *GLUA* and *PR1A* were expressed 10-fold more in trehalose-treated plants compared to

355   water controls 6 h after treatment (Fig. 5A) and their expression increased to over 100-fold at 48 h (Fig.

356   5B). *PR1B* and *OSM,* which respond to the defense hormone ethylene (ET), were also significantly

357   upregulated 48 h after trehalose treatment (Tornero, Gadea et al. 1997, Smart, Myers et al. 2003). The

358   ET synthesis gene *ACO5* was also 10-fold more highly expressed compared to water controls 6 and 48 h

359   after trehalose treatment (Fig. 5A+B) (Jacobs, Milling et al. 2013). Transcripts of the ABA-responsive

360   genes *RD22* and *DHN_TAS* were slightly but significantly upregulated 48 h after trehalose treatment,

361   possibly in response to the increased levels of ABA in xylem sap shown in Fig. 5B (Yamaguchi-Shinozaki

362   and Shinozaki 1993, Shinozaki, Yamaguchi-Shinozaki et al. 2003, Kissoudis, Seifi et al. 2016). Together,

363   these qRT-PCR experiments revealed that tomato plants respond to trehalose by increasing expression

364   of genes in JA, SA, ET, and ABA hormonal pathways involved in, among other things, biotic and abiotic

365   responses.

Figure 5. Exogenous trehalose treatment affects expression of hormone-responsive tomato defense genes, and trehalose-mediated disease resistance is partially regulated by salicylic acid signaling. Effects of trehalose treatment on expression of diverse defense genes in Bonny Best tomato stems, determined by qRT-PCR at **A)** 6 h and **B)** 48 h after trehalose treatment at the roots (one-sample t-test to a hypothetical mean of 1.0 where 1.0 = no difference in expression; asterisks: *, P=.01-.05; **, P=.001-.01; ***, P=.0001-.001; ****, P<.0001). Expression levels are shown as fold-change relative to those in mock (water)-treated plants. Data represent three biological replicates per time point, each containing five plants per treatment. Bars represent standard deviation. **C)** Wilt disease progress following *R. solanacearum* inoculation of water- or trehalose-treated *NeverRipe* ET-insensitive tomatoes and their parent cultivar 'Pearson' (one-way ANOVA of Area Under the Curve or AUC; 'Pearson' H_2O vs. *NeverRipe* H_2O, P=.91; 'Pearson' tre vs. *NeverRipe* tre, P=.98). **D)** Wilt disease progress following *R. solanacearum* inoculation of water- or trehalose-treated SA-insensitive *NahG* over-expressing plants and their parent cv. Moneymaker (one-way ANOVA of AUC, Moneymaker H_2O control vs. *NahG* H_2O control, P=.74; Moneymaker tre vs. *NahG* tre, P=.04). For **C)** and **D)** plants were treated at the roots with trehalose, then inoculated with *R. solanacearum* via soil drenching 48 h later. Data represent three biological replicates, each containing 15 plants/treatment (45 plants total). Bars represent standard error.

The finding that SA- and ET-responsive tomato genes were upregulated 6 and 48 h after

trehalose treatment suggested that ET perception and SA responses might be necessary for trehalose-

mediated increase in bacterial wilt resistance. We tested this hypothesis by treating with trehalose the

roots of ET-insensitive *NeverRipe* mutant tomatoes (cv. Pearson) and SA-degrading *NahG* transgenic

tomatoes (cv. Moneymaker). After 48 h, all plants were soil drench inoculated with *R. solanacearum*. As

observed for cv. Bonny Best, trehalose treatment increased bacterial wilt resistance of Pearson and

Moneymaker plants. Trehalose also increased bacterial wilt resistance of both the *NeverRipe* and *NahG*

plants compared to untreated controls (Fig. 5C and 5D). However, trehalose-mediated bacterial wilt

resistance does not require ET perception since bacterial wilt developed similarly on trehalose-treated

NeverRipe and the parental Pearson genotype (*P*<.98). In contrast, trehalose-treated *NahG* tomatoes

were significantly more resistant to bacterial wilt than trehalose-treated Moneymaker parental line

(*P*<.04). These results suggest that either a lack of downstream SA responses or overaccumulation of SA

degradation products contributes to the trehalose-mediated increase in bacterial wilt resistance.

Alternatively, trehalose may improve resistance downstream of SA responses.

More than half of the *NahG* tomato plants displayed a transient marginal leaf necrosis

phenotype following trehalose treatment (data not shown), which could have been due to increased

ROS production. To test this, we infused leaves from four-week-old wild type Bonny Best, Moneymaker,

or *NahG* tomato plants with water, *R. solanacearum* cells, or 30 mM trehalose solution and observed

ROS production with DAB staining. Infusing leaves with trehalose did not trigger excess ROS production

(Supp. Fig. 4), suggesting that another physiological effect caused the transient marginal necrosis in the

NahG plants.

Trehalose treatment promoted accumulation of hormones and defense compounds in xylem sap

To better understand the mechanisms of trehalose-mediated wilt resistance, we measured

levels of a suite of plant hormones, oxylipins, and defense-related compounds in xylem sap from

infected and trehalose-treated Bonny Best tomatoes. We harvested sap from tomatoes displaying a

disease index of 1 (DI=1, partially wilted, late-stage infection) and from uninoculated plants at 6, 24, and

48 h after trehalose treatment. Using targeted quantitative mass spectrometry, we measured

metabolites from the phenylpropanoid biosynthesis pathway, several oxylipins from the 9-lipoxygenase

and 13-lipoxygenase pathways, and other selected defense metabolites (Supplemental Table 2).

Sap from *R. solanacearum*-infected plants and trehalose-treated plants contained different

levels of targeted metabolites compared to sap from healthy tomatoes. Sap collected 24 h after

trehalose treatment had the highest number of enriched compounds. These were consistent with a

shock or wound response even though plants were not wounded (Vallad and Goodman 2004).

Surprisingly, trehalose treatment, but not *R. solanacearum* infection, increased SA concentration at 24 h

(Fig. 6A). This result could mean that trehalose plays a role in SA signaling and suggested that SA might

not have a significant role in xylem sap at late-stage disease. However, benzoic acid levels did increase in

xylem sap from both trehalose treated and *R. solanacearum* infected plants (Fig. 6B). Benzoic acid is a

potent antimicrobial compound produced through the phenylpropanoid pathway that can be a direct

precursor to SA (Chen, Zheng et al. 2009, Widhalm and Dudareva 2015). Interestingly, coumaric acid,

another phenylpropanoid defense compound that *R. solanacearum* can degrade, was significantly

increased in sap from *R. solanacearum*-infected plants but not in sap from trehalose treated plants (Fig.

6C) (Lowe, Ailloud et al. 2015). Levels of traumatic acid, which results from the oxidation of traumatin

produced by wounded plants, were higher in sap from both *R. solanacearum*-infected plants and

trehalose-treated plants at 24 h (Fig. 6D) (Zimmerman and Coudron 1979).

Figure 6. Trehalose treatment increased metabolites related to defense and wounding in tomato

xylem sap. LC-MS/MS determined concentrations of hormones, oxylipins, and other metabolites in

xylem sap collected from cv. Bonny Best tomato plants that were mock-inoculated (H_2O), inoculated

with *R. solanacearum* GMI1000 when wilt symptoms first appeared (*Rs* DI=1), or had been treated at

the roots with 30mM trehalose (Tre (6h), etc.) Differences from water-treated control plants were

determined by one-way ANOVA Fisher's LSD multiple comparison test: *, *P*=.01-.05; **, *P*=.001-.01;

, *P*=.0001-.001; *, *P*<.0001. Data shown represent the means of samples from 4-6 individual

plants. Bars indicate mean, vertical lines show standard deviation. **A)** SA=salicylic acid; **B)** BA=benzoic

acid; **C)** COUMA=coumaric acid; **D)** TA= Traumatic acid; **E)** 12-OPDA= 12-oxo-8(*Z*),13(*Z*)-phytodienoic

acid; **F)** JA=jasmonic acid; **G)** JA-Ile=jasmonic acid isoleucine; **H)** 9-HOT= 9(*S*)-hydroxy-

10(*E*),12(*Z*),15(*Z*)-octadecatrienoic acid; **I)** 9-HOD= 9(*S*)-hydroxy-10(*E*),12(*Z*)-octadecadienoic acid; **J)**

13-HOD= 9(*S*)-hydroxy-10(*E*),12(*Z*)-octadecadienoic acid; and **K)** 13-KOD= 13-oxo-9(*Z*),11(*E*)-

octadecadienoic acid.

The chemical signatures exclusively present in trehalose treated plants were the persistent

increases in JA, the JA precursor 12-OPDA, and JA-Ile, the biologically active form of JA (Fig. 6E-G). This

accumulation of jasmonates, together with the upregulation of JA and ET-pathway defense genes,

suggested that trehalose treatment induces a systemic defense response that contributes to its

protective effect (Vallad and Goodman 2004, Choudhary, Prakash et al. 2007).

Curiously, several defense-related oxylipins from the reductase and lipoxygenase branches of

the LOX pathway were enriched in sap from *R. solanacearum*-infected plants and 24 h post-trehalose

treatment plants (Fig. 6H-K). These oxylipins include 9(*S*)-hydroxy-10(*E*),12(*Z*),15(*Z*)-octadecatrienoic

acid (9-HOT, Fig. 6H); 9(*S*)-hydroxy-10(*E*),12(*Z*)-octadecadienoic acid (9-HOD, Fig. 6I); 13(S)-hydroxy-

9(Z),11(E)-octadecatrienoic acid (13-HOD, Fig. 6J); and 13-oxo-9(*Z*),11(*E*)-octadecadienoic acid (13-KOD,

Fig. 6K). All these oxylipins have antimicrobial and signaling activity in defense against pathogens (Prost,

Dhondt et al. 2005, Vicente, Cascón et al. 2012, Marcos, Izquierdo et al. 2015). Induction of these

oxylipins in this study suggests involvement in resistance to bacterial wilt. Altogether, trehalose

treatment had an unexpectedly large impact on xylem levels of defense metabolites, particularly

jasmonates and SA, which have often been considered antagonistic signals.

Discussion

To survive drought, plants must respond to and overcome multiple adverse physiological events

(Shinozaki, Yamaguchi-Shinozaki et al. 2003, Osakabe, Osakabe et al. 2014). Our stringent meta-analysis

of four different transcriptomic studies found that plants infected with *R. solanacearum* have a gene

expression signature like that of water-stressed plants, suggesting that bacterial wilt disease

physiologically mimics abiotic drought stress. The consistency of this response across multiple tomato

cultivars suggest that this is a robust, replicable phenomenon.

Infected tomatoes slightly upregulated TPS7, (Zhang, Hong et al. 2016), suggesting this gene

could be responsible for the observed increased trehalose synthesis during bacterial wilt. This gene was

also upregulated 36 h after infection with *P. syringae and* silencing this gene (annotated as *SlTPS3* in

that study) decreased tomato trehalose content in response to the fungal pathogen *Botrytis cinerea* and

also decreased plant resistance to *B. cinerea* (Zhang, Hong et al. 2016).

In response to *R. solanacearum* infection, tomatoes downregulated multiple aquaporins,

suggesting that plants may try to alter water flow during bacterial wilt. In particular, SlTIP2;2

(Solyc03g120470), an aquaporin that improved transpiration and yield when overexpressed in tomato

plants, was notably downregulated in our meta-transcriptomic analysis (Sade, Vinocur et al. 2009).

Accordingly, *Arabidopsis* PIP family aquaporin proteins were generally downregulated during drought

stress (Alexandersson, Fraysse et al. 2005). Tomato aquaporins are encoded by diverse families and

channel other solutes in addition to water, so it would not be expected that all aquaporins play the

same role during bacterial wilt (Reuscher, Akiyama et al. 2013, Patel and Mishra 2021).

An ABA-inducible dehydrin and an ABA hydroxylase gene were strongly upregulated in infected

plants. In *Arabidopsis* and tomato*,* CYP707A ABA hydroxylases, which catalyze the first step in ABA

degradation, are involved in dehydration and rehydration (Kushiro, Okamoto et al. 2004, Saito, Hirai et

al. 2004, Umezawa, Okamoto et al. 2006, Ji, Kai et al. 2014). This suggests both that tomatoes tightly

regulate ABA catabolism during bacterial wilt and that bacterial wilt causes water stress. Expression of

TAS14, a dehydrin inducible by environmental stress and ABA, was highly upregulated in infected plants,

further evidence that bacterial wilt causes drought stress (Godoy, Pardo et al. 1990). *Arabidopsis* ABA

synthesis, signaling, and responsive genes were similarly affected during late-stage bacterial wilt

infection (Hu, Barlet et al. 2008). Elevated ABA levels are usually associated with drought stress;

together with the observation that *R. solanacearum* infection lowers plant stomatal conductance, this

indicates that bacterial wilt disease significantly disrupts tomato water physiology. Further

characterization of ABA-responsive networks in tomatoes could identify targets for breeding more

hydraulically efficient tomatoes that are also more resistant to bacterial wilt.

Our data suggest that there might be an indirect defensive component to ABA concentration
and signaling. The levels of ABA in xylem sap increased in response to both *R. solanacearum* infection
and trehalose treatment, and our transcriptomic data suggest tomato plants degrade ABA during
disease via an ABA hydroxylase.   Although it would be interesting to test the effects of trehalose and *R.*
solanacearum on tomato ABA synthesis mutants, existing ABA mutants (*flacca, notabilis, sitiens)* share a
wilting phenotype due to ethylene overproduction and lack of control over stomatal closure. This
physiological wilting interferes with assessing pathogen-caused wilting symptoms (Tal 1966, Thompson,
Thorne et al. 2004, Kaiser, Morales et al. 2020). However, *Arabidopsis* studies also indicate that ABA is
important for tomato resistance to wilt disease. Deleting several cellulose synthase (*irx*) genes increased
Arabidopsis resistance to bacterial wilt in an ABA-dependent manner and ABA synthesis mutants were
more susceptible to *R. solanacearum* (Hernández-Blanco, Feng et al. 2007). Further, ABA perception
mutants are also hyper-susceptible to bacterial wilt (Zhao, Wang et al. 2019).

Treating tomato roots with ABA reduced both stomatal conductance and *R. solanacearum*
colonization of stems, suggesting that *R. solanacearum* spread depends, at least in part, on active
hydraulic conductance and a negative pressure gradient in host xylem. It has long been assumed that *R.*
solanacearum uses the plant's own water transport to spread within its host, but this is now
demonstrated experimentally. When this pathogen blocks host plant water flow, it decreases its own
spread and pathogenic fitness. Interestingly, some wilt-resistant tomato varieties thrive even when they
are latently infected. This supports the theory that resistant hosts can, to some extent, locally limit
vascular bacterial growth enough to prevent occlusion of xylem vessels (Beattie 2011). We speculate
that ancestors of *R. solanacearum* were once endophytes, but lost the ability to moderate their
population growth in susceptible hosts, possibly as a result of evolutionary selection pressures imposed
by agriculture, as has been proposed for *Xylella fastidiosa* (Beattie and Lindow 1995).

In our system, exogenous trehalose temporarily improved water conservation in uninfected
tomatoes and delayed abiotic wilting of drought-stressed plants. This was likely because trehalose
treatment temporarily reduced stomatal conductance and transpiration and increased water use
efficiency. Trehalose increased ABA content in xylem sap 24 h post-treatment, and disease symptoms
often occurred after ABA concentration peaked, consistent with a potential signaling role for ABA.
These results confirm and extend findings that trehalose, ABA signaling, and water use interact in
complex ways. Additional studies are needed to connect ABA signaling and concentration in xylem sap
to downstream effects on disease resistance.

Other *Arabidopsis* and tomato trehalose/water stress studies have confirmed some of the

phenotypes we observed. The *Arabidopsis TPS1* gene, which is responsible for maintaining T6P levels,

responds to ABA and helps regulate stomatal aperture (Avonce, Leyman et al. 2004, Gomez, Gilday et al.

2010). At least two *Arabidopsis* TPP proteins (AtTPPI and AtTPPE) that are controlled by ABA-responsive

transcription factors contribute to trehalose production and mitigate drought stress (Lin, Yang et al.

2019, Lin, Wang et al. 2020, Wang, Chen et al. 2020). Yu and co-workers found that applying trehalose

to tomato roots improved water conservation, decreased ABA biosynthesis gene expression, increased

expression of ABA-responsive genes, kept stomates closed five days after trehalose treatment without

water, and increased ABA levels in leaves after 24 h (Yu, Zhao et al. 2019). Our results contrast in some

ways with this paper. We found that ABA content temporarily increased in xylem sap 24 h after

trehalose treatment, while stomatal conductance was reduced for less than 24 h. These differences may

be explained by methodological variation: Yu and co-workers did not include non-drought stressed

tomato controls, they measured ABA in leaves rather than xylem, and they measured stomatal aperture

rather than conductance. We observed a lag between the stomatal conductance response in leaves and

ABA concentration in stem xylem sap, leading us to hypothesize that ABA concentration in leaves would

be a more biologically relevant marker for behavior of leaf stomates. It would be interesting to

determine if either exogenous trehalose or ABA treatment changes trehalose concentration in tomato

xylem sap.

In our system, trehalose protected plants against disease when applied via soil drenching, but

not when applied as a spray. Spraying wheat leaves with trehalose did increase plant growth, improve

water stress tolerance, and provide resistance to powdery mildew, a leaf disease (Reignault, Cogan et al.

2000, Renard-Merlier, Randoux et al. 2007, Smeekens 2017). Multiple applications of trehalose to leaves

via spray also increased tobacco growth under nitrogen limiting conditions (Lin, Zhang et al. 2017). We

found no studies demonstrating that spraying tomato leaves with trehalose promotes resistance to

pathogens, although leaf infiltration provides resistance to aphid predation (Singh and Shah 2012).

These findings suggest that all plants do not respond to exogenous trehalose in the leaves in the same

way. Additionally, the receptors for sugar-mediated DAMP response in tomatoes are unknown; grafting

experiments may offer insights into whether these are primarily expressed in the roots.

The plant hormone SA, which mediates resistance to *R. solanacearum* in many systems,

is also necessary for induction of plant innate immunity and activation of systemic acquired resistance

(Deslandes, Olivier et al. 2002, Hirsch, Deslandes et al. 2002, Chen, Lin et al. 2009, An and Mou 2011,

Boatwright and Pajerowska-Mukhtar 2013, Jacobs, Milling et al. 2013).. We found SA-responsive tomato

defense genes were highly upregulated 6 and 48 h after trehalose treatment. These same genes are

upregulated in diseased tomato plants following *R. solanacearum* soil soak infection, suggesting

trehalose treatment activates SA-mediated resistance pathways that could affect *R. solanacearum*

infection (Milling, Babujee et al. 2011). Expression of the SA-responsive marker gene PR1b is affected in

several tomato trehalose metabolism knockdown mutants (decreased in *SlTPS4,* increased in *SLTPS5* and

SlTPP2) after *P. syringae* pv. *tomato* challenge (Zhang, Hong et al. 2016), further suggesting interplay

between trehalose metabolism and resistance to bacterial pathogens.

Trehalose treatment increased wilt resistance of SA-degrading *NahG* tomato plants, but not of

the parental Moneymaker cultivar. There are several possible explanations for this result. First,

trehalose treatment could have increased SA levels so much that overproduction of catechol, the

breakdown product of SA, contributed to resistance (Retig and Chet 1974). Second, the excess ET and JA

in *NahG* plants could promote resistance to *R. solanacearum*. After infection with *P. syringae* DC3000

NahG tomatoes produce more ET and JA, both of which can contribute to disease resistance (Spoel,

Koornneef et al. 2003, López-Gresa, Lisón et al. 2016). ET is involved in *R. solanacearum*-host

interactions and ET-responsive defense genes were upregulated at 6 and 48 h post trehalose treatment.

Interestingly, ET and JA signaling-responsive defense-related genes were downregulated in tomatoes

with silenced *SlTPS3*, *4*, and *7* (Zhang, Hong et al. 2016)*.* However, ET-insensitive *NeverRipe* tomatoes

were neither more nor less resistant to *R. solanacearum* than wild type after trehalose treatment.

Treatment with a biocontrol strain of the oomycete *Pythium* enhanced resistance to *R. solanacearum* in

NahG tomatoes, but not in a JA mutant (Hase, Takahashi et al. 2008). Since the ET defense pathway

appeared less relevant in our study, it may be useful to determine if JA enhances *R. solanacearum*

resistance in tomatoes. Our experiments with tomato defense hormone mutants also demonstrated

that trehalose treatment significantly delays disease incidence relative to water-treated control plants in

multiple tomato genotypes.

Plants use hormones to fine-tune their growth, development, and defense, and xylem vessels

can act as signal highways for long-distance transport of auxin, ABA, JA, and SA (Jackson 1997, Park, Lee

et al. 2017). We observed increased SA concentration in xylem sap of trehalose-treated plants but not in

sap from *R. solanacearum*-infected plants. This was surprising because SA signaling and concentration

are integral to plant response to bacterial wilt (Jacobs, Milling et al. 2013). However, *R. solanacearum*

actively degrades SA, which is toxic to the pathogen (Lowe-Power, Jacobs et al. 2016), so SA

concentrations might be higher in root xylem than in stem xylem. For instance, an *Arabidopsis* mutant

impaired in secondary wall deposition had higher SA concentrations in the root and increased resistance

to *R. solanacearum* infection, possibly through vascular immunity (Denancé, Ranocha et al. 2013). Taken

together, the increased SA levels in xylem sap, the increased expression of SA-dependent defense genes,

and the trehalose-dependent *NahG* resistance phenotype suggest that the protective effect of trehalose

requires systemic resistance induced via SA.

Unexpectedly, jasmonate levels were elevated in trehalose-treated plants but not in *R.*

solanacearum-infected plants. As discussed earlier, there is some evidence that JA can play a role in *R.*

solanacearum defense. Genes related to ET, JA, and auxin signaling were upregulated in resistant LS-89

tomato seedlings one day after stem inoculation with *R. solanacearum* (Ishihara, Mitsuhara et al. 2012).

The higher concentration of jasmonates in trehalose-treated xylem sap may have contributed to

resistance. This hypothesis could be tested using JA-deficient tomato mutants.

SA- and JA-mediated plant defense responses are often antagonistic (Glazebrook 2005, Derksen,

Rampitsch et al. 2013). Our gene expression analysis followed this pattern; at 6 h post trehalose

treatment JA-responsive *pin2* was downregulated, while SA-responsive *PR1b* and *gluA* were

upregulated. Roots of wilt resistant Hawaii7996 tomatoes displayed a similar gene expression profile 24

h after *R. solanacearum* infection (French, Kim et al. 2018). Despite the antagonism suggested by the

gene expression data, the concentrations of SA, JA, and the JA precursor 12-OPDA all increased in

trehalose-treated plants at 24 h. In maize, levels of OPDA and JA are irrelevant for the classic SA/JA-Ile

antagonism and 12-OPDA and SA are co-regulated (Wang, Borrego et al. 2020). Infusing maize xylem

sap with 12-OPDA increased resistance to the fungal pathogen *Colletotrichum graminicola* while infusing

similar concentrations of JA increased susceptibility. Disease experiments with JA-insensitive tomato

mutants could determine if 12-OPDA or JA contribute to resistance and define their relationship to SA

signaling.

JA signaling also induces leaf senescence*,* and chlorosis is associated with increased JA

production (He, Fukushige et al. 2002). Consistent with this, trehalose treatment significantly reduced

chlorophyll content 2-10 days post-treatment, which correlated with the increased JA production in

trehalose-treated plants. Similarly, applying 5 mM trehalose to soybean roots did not affect JA

concentration, but it did decrease chlorophyll content at ten days (Asaf, Khan et al. 2017).

Trehalose treatment and *R. solanacearum* infection both increased xylem sap levels of several

antimicrobial compounds, notably phenolics. *R. solanacearum* requires at least two multi-drug efflux

pumps and a hydroxycinnamic acid degradation pathway for full virulence and fitness, showing that

antimicrobials, including phenolics, protect tomato plants from bacterial wilt (Brown, Swanson et al.

2007, Lowe, Ailloud et al. 2015). Plants have specialized cells that store and release phenolics into the

xylem as part of their defense response (Beckman 2000), and our data indicate that trehalose treatment

and *R. solanacearum* infection both induce this release.

Levels of the oxylipins 9-HOT, 9-HOD, 13-HOD, and 13-KOD increased in sap from both *R.*

solanacearum-infected plants and trehalose-treated plants 24 h after treatment. These metabolites,

which accumulate in wounded maize, bean, and *Arabidopsis*, can be potent inducers of defense gene

response (Alméras, Stolz et al. 2003, Walters, Cowley et al. 2006, He, Borrego et al. 2020). Further, these

oxylipins have direct antimicrobial activity against fungal and bacterial pathogens (Prost, Dhondt et al.

2005). The data generated by this study deserve further analysis since little is known about oxylipin

signatures during bacterial wilt or if oxylipins are toxic to *R. solanacearum*. However, 9-LOX-derived

oxylipins, including 9-HOT and 9-HOD, increase in tobacco leaves inoculated with *R. solanacearum*

GMI100, which triggers an intense defensive hypersensitive response (HR) on tobacco (Montillet, Agnel

et al. 2002). We speculate that the presence of related oxylipins in our study may indicate vascular HR. It

would be interesting to see if defense compound concentration changed in the xylem sap of resistant

tomatoes challenged with *R. solanacearum.*

We propose that tomato plants activated induced systemic resistance because they recognized

trehalose as an endogenous (self-derived) damage associated molecular pattern or DAMP (Tanaka and

Heil 2021). DAMPs are products released during pathogen invasion or wounding that alert the plant to

an attack (Ranf, Eschen-Lippold et al. 2011, De Lorenzo, Ferrari et al. 2018, Hou, Liu et al. 2019). A DAMP

can be almost anything that is usually intracellular like DNA, sugars, or NAD (indicating cellular leakage),

or products of lignin and cellulose degradation (indicating mechanical or enzymatic disruption of plant

cell walls). Plant systemic resistance response to DAMPs, mechanical wounding, or pathogen challenge

typically involves accumulation of SA, JA, and defense oxylipins, and increased expression of SA, JA, and

ET-responsive defense genes (Vallad and Goodman 2004, Choudhary, Prakash et al. 2007, Savatin,

Gramegna et al. 2014, Wang, Borrego et al. 2020). Consistent with this definition, we found that treating

tomato roots with trehalose increased SA, JA, and defense oxylipins, and upregulated SA, JA, and ET

defense genes. Our finding adds to the growing evidence that sugars, specifically trehalose, contribute

to plant innate immunity (Bolouri Moghaddam and Van den Ende 2012).

Could treating roots with compatible solutes like trehalose in the field provide temporary

resilience until disease or drought pressure have passed? Other chemical inducers of plant defense

responses can reduce bacterial wilt*.* Acibenzolar-S-methyl (Actigard) and DL-3-aminobutyric acid (BABA)

both lowered wilt disease incidence in greenhouse and field by promoting defense gene expression and

defense compound production, respectively (Jakab, Cottier et al. 2001, Pradhanang, Ji et al. 2005,

Hassan and Abo-Elyousr 2013). Additionally, growing tomatoes hydroponically with SA and JA increased

defense enzyme production and reduced bacterial wilt incidence (Mandal, Kar et al. 2013). Finally,

exogenous L-histidine induced resistance to bacterial wilt in an ET-dependent manner in *Arabidopsis* and

tomato (Seo, Nakaho et al. 2016). Further studies are needed to know if treating tomatoes with

exogenous trehalose could be a practical strategy for bacterial wilt management.

R. solanacearum GMI1000 produces and secretes abundant amounts of a trehalase (TreA)

during tomato infection. Trehalose is not an important nutrient for the pathogen but the ability to

degrade trehalose contributes to bacterial fitness and virulence by an unknown mechanism (Jacobs,

Babujee et al. 2012, Hamilton, Steidl et al. 2021). Interestingly, treating plants with the trehalase

inhibitor Validamycin A increased bacterial wilt resistance (Ishikawa, Fujimori et al. 1996). We previously

showed that the trehalose in xylem sap of *R. solanacearum*-infected tomato plants is produced by the

plant (MacIntyre, Barth et al. 2019). Could *R. solanacearum* secrete trehalase to counter the effects of

plant-produced trehalose?

Breaking down plant-produced trehalose could benefit *R. solanacearum* in several ways. First,

the pathogen may degrade xylem sap trehalose in order to reduce host plant defenses by removing this

potential DAMP. Second, as discussed above, trehalose reduces tomato transpiration, and transpiration

helps *R. solanacearum* spread and colonize host xylem. *R. solanacearum* may degrade trehalose to

maintain this disseminating sap flow by preventing trehalose-mediated stomatal closing. Third, *R.*

solanacearum could degrade trehalose to interfere with the trehalose-T6P signaling that plants use to

fine-tune their sucrose to starch ratios.   Trehalose degradation could thus indirectly increase the

amount of available sucrose, which is a key carbon source fueling *R. solanacearum* growth in xylem

(Jacobs, Babujee et al. 2012, Hamilton, Steidl et al. 2021). The fact that the *R. solanacearum* effector

RipTPS is a functional trehalose-6-phosphate synthase is consistent with the hypothesis that this

pathogen benefits from manipulating host plant trehalose metabolism (Poueymiro et al., 2014).

Additional experiments are needed to test these non-exclusive hypotheses about how trehalase benefits

R. solanacearum.

Conclusion

This study argues that trehalose is present in xylem sap of *R. solanacearum*-infected tomatoes

as a response to wilt disease-mediated water stress. Both trehalose levels and the expression of water

stress genes changed in response to *R. solanacearum* infection. Further, ABA content, an indicator of

drought status, increased in xylem sap of diseased plants. We confirmed the link between trehalose and

ABA signaling and drought stress in plants by demonstrating that trehalose treated plants temporarily

reduce stomatal conductance, contain more ABA in their xylem sap, and are more drought tolerant.

When applied to plant roots, trehalose reduced bacterial wilt incidence and reduced host xylem

conductance, which aids bacterial colonization. SA-responsive defense genes were upregulated in

trehalose-treated plants, SA non-responsive *NahG* plants were more resistant to *R. solanacearum* after

trehalose treatment, and SA concentration transiently spiked in xylem sap from treated plants. JA, other

oxylipins, and many antimicrobial compounds also increased in xylem sap of trehalose-treated plants.

Taken together, these results suggest that plants perceive trehalose as a DAMP and respond with

systemic disease resistance. In turn, *R. solanacearum* degrades trehalose as a counter-defense.

Methods

Total RNA extraction and RNA-seq sample preparation | RNA extraction and RNA-seq sample

preparation for the transcriptomic studies were performed as described (Khokhani, Lowe-Power et al.

2017, French, Kim et al. 2018).

RNA-seq data analysis | Illumina HiSeq2500 paired-end 100-bp; Illumina HiSeq2500 paired-end 150 bp

and Illumina HiSeq2000 paired-end 100-bp RNA sequencing were performed on the 18, 6, and 12

samples from three independent studies (two previously unpublished [Studies A+B], and [Study C]

French, Kim et al. 2018). RNA-sequencing data were analyzed through the same analysis pipeline. Read

quality was controlled using FastQC version 0.11.9 (Andrews). Reads were trimmed to remove low-

quality sequences and the Illumina sequencing adaptors (TruSeq3-PE-2) using Trimmomatic version 0.39

(Bolger, Lohse et al. 2014). After trimming and mapping, a total of 930,004,276, 135,532,392, and

103,831,164 paired-end reads were generated for Studies A, B, and C, respectively. Reads for each

sample were mapped to the ITAG4.0 *S. lycopersicum* reference genome using STAR version 2.7.4.a

(Dobin, Davis et al. 2012). Library type was set to strand-specific (reverse) for Study A and Study B, and

set to no-stranded in Study C. Gene expression was measured as the total number of reads for each

sample uniquely mapped to the reference using HTSeq version 0.11.1 and samtools version 1.10 (Li,

Handsaker et al. 2009, Anders, Pyl et al. 2014). Each sample averaged about 12, 9, and 4 million uniquely

aligned reads for Studies A, B, and C, respectively. "EdgeR" package version 3.28.0 was used to filter for

low counts such that at least three of the samples in each study had at least three counts per million,

resulting in a total of 19123, 18193, and 18917 genes remaining for Studies A, B, and C, respectively

(Robinson, McCarthy et al. 2009).

Differential gene expression within each study was conducted using the DESeq2 package

Bioconductor version 3.11 and a Benjamini-Hochberg false discovery rate multiple testing correction of

P=.05 (Benjamini and Hochberg 1995, Love, Huber et al. 2014). Meta-transcriptomic analysis was

performed using the metaRNASeq package version 1.0, and differential gene expression within the

meta-analysis was determined with the DESeq2 results, Fisher, and inverse normal techniques with a

P=.05, as described (Rau, Marot et al. 2014). Differentially expressed genes were filtered using a log2

fold change >|0.585| (corresponds to a fold change of |1.5|).

Gene Ontology analysis was performed using the ShinyGO analysis tool version 0.61, agriGO

version 2.0, and PANTHER GO analysis tool version 14 (Tian, Liu et al. 2017, Mi, Muruganujan et al. 2018,

Ge, Jung et al. 2019). *Arabidopsis thaliana* homologs of S. *lycopersicum* genes coding for transcription

factors were identified using Phytozome platform version 12.1 (Goodstein, Shu et al. 2011). Heatmaps

were visualized with R software version 3.6.1 package "pheatmap" version 1.0.12 (R , Kolde 2019).

Boxplots were visualized using package "ggplot2" (Wickham 2016).

Candidate TPS and TPP proteins were identified in the most recent annotations of the tomato

(cv. 'Heinz 1706') genome available on NCBI and the Sol Genomics Network (ITAG4.0,

https://solgenomics.net/tools/blast/, accessed 03/2020) using BLASTP with standard settings (Sato,

Tabata et al. 2012). BLASTP protein sequences were derived from the *E. coli* K-12 *otsA* and *otsB* proteins,

as well as *Arabidopsis* trehalose-related proteins (Leyman, Van Dijck et al. 2001, Vandesteene, Ramon et

al. 2010, Vandesteene, Lopez-Galvis et al. 2012). Tomato cv. Heinz 1706 genome contained ten TPS

genes, eight TPP genes, and one trehalase gene. These genes were previously identified as summarized

(Zhang, Hong et al. 2016). Tomato ABA-related, LEA, and aquaporin genes were identified by their

PANTHER annotation, and many were previously annotated (Ma, Szostkiewicz et al. 2009, Reuscher,

Akiyama et al. 2013).

The transcriptomic studies were performed and labeled as follows: S1, roots of wilt-susceptible

West Virginia 700 tomato plants (WV) sampled 24 h after infection with *R. solanacearum* strain K60; S2,

roots of WV sampled 48 h after infection with *R. solanacearum* strain K60; S3, seedling roots of wilt-

susceptible cv. Bonny Best (BB) sampled 24 h after infection with *R. solanacearum* strain GMI1000; S4,

mid-stems of BB sampled 72 h after inoculation with *R. solanacearum* strain GMI1000.

Hormone analysis leaf and sap collection | For the hormone analysis, xylem sap was harvested and

collected as described (Lowe-Power, Hendrich et al. 2018). Unwounded three-week-old wilt-susceptible

Bonny Best plants were inoculated with wild-type *R. solanacearum* or water via naturalistic soil soak in

which 50 mL of $1x10^8$ CFU/mL were poured into a pot containing one plant. Plants were harvested when

they reached DI=1 (1-25% of leaf area wilted). Trehalose-treated plants were treated with 50 mL of 30

mM trehalose solution or water, then harvested 6, 24, or 48 h later. Xylem sap was harvested by de-

topping plants 1 cm above the cotyledon and collecting the exuded sap over 30 minutes. The sap that

seeped within the first five minutes was discarded to reduce phloem contamination. Sap was kept on

ice. Bacteria were removed from sap via chilled centrifugation, and sap was subsequently flash frozen.

Stems were destructively sampled to quantify *R. solanacearum* colonization as CFU/g stem by serial

dilution plating. Xylem sap was normalized by sap volume (250-500 µL) and concentration of bacteria in

the stem. Each test group contained six technical replicates (individual plants).

For metabolite analysis of sap, 95 µL of xylem sap was mixed with 5 µL of 50 µM of internal

standard consisting of; d-ABA ($[^2H_6]$(+)-*cis, trans*-abscisic acid, Olchem, Olomouc, Czech Republic), d-CA

(d_7-CA, *trans*-cinnamic acid), d-IAA ($[^2H_5]$ indole-3-acetic acid, Olchem), d-JA (2,4,4-d_3; acetyl-2,2-d_2

jasmonic acid, CDN Isotopes, Pointe Claire, Quebec, Canada), and d-SA (d_6-SA; Sigma-Aldrich, St Louis,

MO, USA). Samples were transferred into autosampler vials for LC-MS/MS analysis. Authentic analytical

external standards for each metabolite were used to identify select ions and retention times for each

metabolite (Supplemental Table 3). The simultaneous detection of several phytohormones used the

following methods with modifications (Müller and Munné-Bosch 2011). We used an Ascentis Express C-

18 Column (3 cm × 2.1 mm, 2.7 µm) (Sigma-Aldrich, St. Louis, MO, USA) connected to an API 3200 LC-

MS/MS (Sciex, Framingham, MA, USA) using electrospray ionization in negative mode with multiple

reaction mentoring. The injection volume was 10 µL and had a 500 µL min^{-1} mobile phase consisting of

Solution A (0.2% acetic acid in water) and Solution B (0.2% acetic acid in acetonitrile) with a gradient

consisting of (time – %B): 0.5 – 10%, 1.0 – 20%, 21.0 – 70%, 24.6 – 100%, 24.8 – 10%, 29 – stop. After

integration with Analyst v1.6.3 (Sciex), the concentration of endogenous metabolites was determined by

comparing their peak areas to those of isotopically labeled internal standards using calculated response

factors.

Stomatal conductance, transpiration, and photosynthesis measurements with LI-COR |To assess the

effect of trehalose treatment on tomato stomatal conductance (g_s), transpiration (E), and

photosynthesis (A_{net}), we used a LI-COR portable photosynthesis system (model LI-6400XT, LI-COR

Biosciences, Lincoln, NE, USA) equipped with a CO_2-controller and a red-blue LED chamber (model LI-

6400-02b). The ambient CO_2 concentration was set to 400 ppm, flow rate to 300 μmol s^{-1}, and the

chamber light source set at an intensity of 1200 μmol m^{-2} s^{-1} for all measurements. Fifteen-day-old

Bonny Best plants were treated with 50 mL of trehalose solution or water, then watered every day after

for the duration of the experiment. LI-COR measurements were taken at 10min, 6 h, 24 h, 48 h, and 72 h

post trehalose treatment. The experiment was repeated twice with ten plants per treatment.

R. solanacearum **culture conditions |** *R. solanacearum* strains were cultured from water stocks or -80 °C

glycerol stocks on TZC plates containing casamino acids, peptone, glucose, and tetrazolium chloride, and

incubated at 28 °C for 48 h (Hendrick and Sequeira 1984). *Rs ΔtreA* cultures were supplemented with 20

mg L^{-1} spectinomycin. The kanamycin-resistant variant of wild-type GMI1000 was supplemented with 25

mg L^{-1} kanamycin on plates and in liquid culture. Overnight cultures of *R. solanacearum* were grown at

28 °C in CPG in a shaking incubator.

ABA treatments | We assessed the effect of ABA treatment on two-week-old tomato stomatal

conductance and subsequent bacterial wilt incidence. ABA was prepared at a stock concentration of

50mg/mL in DMSO. The water + DMSO control plants were watered daily with 50 mL of water

supplemented with the volume of DMSO used to make the 100 μM ABA solution. The ABA-treated

plants were watered daily with 50 mL of 50 or 100 μM ABA solution in water. After 48 h of ABA

treatment, plants were soil drenched with 50 mL of a 1×10^8 CFU/mL of *R. solanacearum* GMI1000 with

kanamycin resistance (to aid in recovery from the plant) (Lowe, Ailloud et al. 2015). Bacteria were

resuspended at the desired O.D. in water + DMSO or the ABA solutions, so plants were inoculated and

ABA treated concurrently. Tomatoes were destructively sampled five days post-infection (after 168 h of

ABA treatment) to assess bacterial colonization of stems. Stomatal conductance, transpiration, and

photosynthesis measurements were taken for the entirety of the experiment with the LI-COR as

described 4-6 h after watering/ABA treating in the morning, which was the length of time it took ABA

treatment to affect conductance.

We assessed ABA toxicity to *R. solanacearum* by supplementing CPG broth with 50-100 μM ABA

or the equivalent volume of DMSO, and growing *R. solanacearum* cells in this media for 48 h. Cells were

grown in a shaking 96 well plate at 28°C in a BioTek Synergy HT plate reader, which measured Abs_{600}

every half hour. The experiment was performed once with eight technical replicates per treatment.

Plant assays with *R. solanacearum* | Plant growth conditions, colonization, and disease curve assays

were performed as described (Khokhani, Tuan et al. 2018, MacIntyre, Barth et al. 2019). All plant assays

were grown on 12 h light/dark cycles at 28 °C with ~60% relative humidity unless stated otherwise. For

the disease progress assays with trehalose treatments, fifteen-day old Bonny Best tomatoes were

treated with either water (control) or 50 mL of 30 mM trehalose solution. Plants were not watered 24 h

post-treatment per standard procedure for soil soak inoculation assays. At 48 h post-treatment plants

were soil drenched with 50 mL of $1x10^8$ CFU/mL of wild type *R. solanacearum* GMI1000 (or Δ*treA* for the

trehalose treatment experiments with this mutant). For the petiole inoculations, 3-week-old tomatoes

were treated with trehalose or water as described above; after 48 h a true leaf was removed, and 2000

wild type GMI1000 cells in 10 μL of water were pipetted onto the cut stem. To assess the effect of

spraying trehalose on bacterial wilt incidence, fifteen-day-old Bonny Best plants were either treated

with 30 mM trehalose solution through the soil as described or leaves were sprayed with trehalose

solution (or water) to run-off. For all assays, plants were inoculated with *R. solanacearum* 48 h post

trehalose root or leaf treatment.

After treatments and inoculations, all plants in disease assays were watered on alternating days

post-inoculation with half-strength Hoagland's solution. Disease assay plants were rated daily on a 0-4

disease index scale where 0= no symptoms, 1=1-25% of leaf area wilted, and 4=76-100% of leaf area

wilted or dead. All disease curves represent four biological replicates, each containing 15

plants/treatment.

To assess *R. solanacearum* colonization of trehalose treated plants, fifteen-day-old Bonny Best

plants were treated with 50 mL 30 mM trehalose solution or water. 48 h post-treatment, plants were

inoculated with 50 mL of $1x10^8$ CFU/mL wild type *R. solanacearum* GMI100. At 5, 6, and 7 dpi plant

stems were destructively sampled 1 cm above the cotyledon, ground, and serially dilution plated.

NahG salicylic insensitive tomato seeds ('Moneymaker' background) and *NeverRipe* ET-

insensitive seeds ('Pearson' background) were propagated in our lab (Lanahan, Yen et al. 1994, Brading,

Hammond-Kosack et al. 2000). Disease assays with trehalose-treated hormone insensitive plants were

as described above.

Water conservation | Two-week-old Bonny Best plants grown as described above were treated with

trehalose or water. Water was subsequently withheld and plants were assessed for wilting symptoms,

using the same visual index scale as for disease progress assays. The experiment included four biological

replicates, each with ten plants per treatment.

QRT-PCR measurement of tomato trehalose and defense gene expression | JA, SA, and ET defense

gene primers were previously characterized in tomato *R. solanacearum* response (Supplemental Table 1)

(Milling, Babujee et al. 2011). The *rd22* and *dhn_tas* ABA-responsive primers were previously described

(Yamaguchi-Shinozaki and Shinozaki 1993, Kissoudis, Seifi et al. 2016). *ACTIN* and *DNAJ* primers served

as our constitutively expressed normalizing genes.

To measure defense gene expression, two-week-old Bonny Best tomatoes were treated with 50

mL of 30 mM trehalose or water. 100 mg of stem tissue was harvested 6 or 48 h post-treatment, then

flash frozen. For all samples, stems were homogenized for a minute in a Powerlyzer (Qiagen) at room

temperature in 450 µL of Buffer RLT from the RNAeasy Plant Mini Kit (Qiagen). RNA extraction

proceeded as specified in the kit protocol with off-column DNase I treatment using a DNA-free kit

(Thermo-Fisher). RNA quality and quantity were assessed with a NanoDrop (Thermo-Fisher) and reverse

transcribed into cDNA using Superscript III First-Strand Synthesis SuperMix (Invitrogen). Lack of genomic

DNA contamination was confirmed with PCR of no reverse transcriptase controls. Quantitative real-time

PCR was performed in 10 µL reactions with EvaGreen Mastermix (Biotium) using kit-specified reaction

parameters in a Quant Studio 5 Real-Time PCR machine (ThermoFisher Scientific). Averaged actin and

dnaJ values were used as the stably expressed normalization genes. Fold change values were calculated

using the $\Delta\Delta C_T$ method as described (Milling, Babujee et al. 2011). The data are presented as fold change

in gene expression compared to mock-inoculated controls and represent 16 plants per treatment over

four biological replicates.

Relative water content | To assess the water status and osmotic adjustment of tomato plants treated

with trehalose, we measured relative water content (RWC) in leaves. This directly measures cellular

hydration. Three-week-old Bonny Best tomato plants were treated with 50 mL of water or 30 mM

trehalose solution. 48 h post-treatment, leaves were excised at the petiole and weighed for fresh

weight. Leaves were then rehydrated by floating on distilled water in closed petri dishes for 4 h, at

which time they were blotted, and their turgid weight was assessed. Leaves were then thoroughly dried

in a food dehydrator oven for 24 h, and dry weight was evaluated. RWC was calculated as follows: %

RWC= [(Leaf fresh weight – dry leaf weight)/(turgid leaf weight-leaf dry weight)] x 100 (Barrs and

Weatherley 1962). The data represent two biological replicates with five technical reps (leaf from an

individual plant) per treatment.

Root length assays | Root length assays to quantify trehalose treatment effect on root length were

performed as described with modification (Tran, MacIntyre et al. 2016). Tomato seeds were sterilized in

10% bleach for 10 min, then 95% ethanol for 10 min, then washed five times with sterile water and

germinated in the dark on 1% water agar plates at room temperature. Four-day-old tomato seedlings

were transferred to germination pouches (Mega International, Newport, MN) then watered with either

10 mL of water or 30 mM trehalose solution and incubated in a 28°C growth chamber. Following the

initial trehalose treatment, pouches were kept moist as needed with sterile water and imaged ten days

post-treatment. This experiment was performed three times with ten tomato

seedlings/pouch/treatment. New root growth was calculated as % new root growth= (new growth/total

root length) x 100.

Chlorophyll determination | To measure the amount of chlorophyll in tomatoes, we used a SPAD-502

meter (Konica Minolta, Inc.) (Ling, Huang et al. 2011). Readings were performed on the first true leaves

of two-week-old Bonny Best plants treated with 50 mL of a 30 mM trehalose solution or water. The data

represent two biological replicates with ten plants per treatment per replicate.

Bacterial leaf spot of tomato inoculations | *Xanthomonas gardneri* culture conditions, infection, and

colonization analysis were as described (Potnis, Soto-Arias et al. 2014). *X. gardneri* strains were cultured

on nutrient agar plates (Millipore-Sigma) supplemented with nalidixic acid (50 µg/mL) at 28°C. Inoculum

was grown for 48 h in a shaking incubator at 28°C in nutrient broth supplemented with nalidixic acid.

The inoculum was adjusted to OD_{600}=0.3 and diluted 1:100 in 400 mL sterile water with 75 µL Silwet

(Fisher Scientific). Four-week-old Bonny Best tomato plants that were treated with water or 50 mL of 30

mM trehalose solution 72 h previously were dip-inoculated for 30 s in the inoculum. Plants were

incubated in high humidity conditions (85-90%) in plastic containers in the growth chamber for 48 h.

After 48 h, plants were moved to low humidity conditions during the day (taken out of containers, 45-

60%) and high humidity conditions at night on a 12 h cycle. Bacterial spot symptoms appeared 5-6 days

post-inoculation. Leaf tissue from individual plants was sampled at 3, 6, and 9 days post-inoculation.

Two 2 cm^2 portions of leaf were sampled from each plant, ground in a Powerlyzer (Qiagen) in water, and

serially diluted and enumerated on nutrient agar + nalidixic acid. The data represent three biological

replicates with 12 technical reps/treatment/timepoint.

DAB staining tomato leaves for ROS production | Tomato leaves were stained with 3,3'-

Diaminobenzidine (DAB, Sigma-Aldrich) as described (Flores-Cruz and Allen 2009). Leaves from 4-5-

week-old Bonny Best tomato plants were infused with water, 30 mM of trehalose solution, or $1x10^9$

CFU/mL *R. solanacearum* cells by pricking with a needle then flushing the apoplast with the solution.

Plants were returned to the growth chamber overnight. At 24 h, leaves were cut and placed in 0.1

mg/mL DAB solution prepared at pH 3.8 at room temperature (Thordal-Christensen, Zhang et al. 1997).

Containers holding the stain and leaves were wrapped in aluminum foil to exclude light and incubated

for 8 h at room temperature with gentle shaking. At 8 h, leaves were bleached by boiling in 95% ethanol

for 10 min, then further cleared with overnight incubation in 70% ethanol before imaging. These

experiments were performed three times with leaves from four individual tomato plants per treatment.

Statistical Analyses | Statistical analyses and graphs were generated with GraphPad Prism 8.

Acknowledgments: Thanks to Dr. Raka Mitra (Carleton College) for manuscript review and valuable

discussions. Thank you to Dr. Eric Kruger for the use of equipment and Dr. Kim Cowles for *X. gardneri*

cultures and technical advice. This research was supported by a USDA-NIFA Predoctoral Fellowship to

AMM and by the University of Wisconsin-Madison College of Agricultural and Life Sciences.

Captions for Supplemental Materials

Supplemental Table 1. Strains and primers used in this study

Supplemental Table 2. Concentration of hormones, oxylipins, and phenylpropanoids in tomato xylem

sap following various treatments

Supplemental Table 3. Metabolite key for hormone analysis

Supplemental Figure 1. ABA and DMSO are not toxic to *Rs*. Growth of Rs cultured in rich CPG broth

supplemented with ABA in DMSO, or water with DMSO because ABA stock solutions were dissolved in

DMSO (100 µM treatments contained 0.1 µL of either DMSO or ABA stock solution, and 50 µM

treatments contained 0.05 µL DMSO or ABA stock solution) (ANOVA Area Under Curve/AUC, Fisher's LSD

multiple comparisons to CPG control, 100 µM DMSO, P=.55; 50 µM DMSO, P=.53; 100 µM ABA, P=.67;

50 µM ABA, P=.45). Growth was measured spectrophotometrically using a Bio-Tek plate reader; the data

represent eight technical reps/treatment. The bars represent the standard error of the mean.

Supplemental Figure 2. Trehalose treatment temporarily improves water efficiency without affecting

photosynthesis; trehalose treatment does not affect root growth and relative water content of leaves

but lowers chlorophyll concentration. A) Water efficiency, measured with a LI-COR as photosynthetic

rate/transpiration, in trehalose treated plants compared to water only controls (t-test, $P<0.5$). **B)**

Photosynthesis, measured as A_{net}, in trehalose treated plants (Mixed model ANOVA, Tukey's HSD,

$P>.05$). The water efficiency and photosynthetic data represent 20 plants/treatment. LI-COR

measurements were taken ~10 min after treatment and then in the morning every time point after.

Shaded bars on the graphs represent night periods. All data were calculated from the same LICOR time

points. The barson the data points represent the standard deviation. **C)** Relative water content of leaves

was calculated from the fresh weight, hydrated weight, and dry weight of leaves from water control or

trehalose treated BB tomatoes (t-test, $P=.14$). The data represent two biological replicates with five

technical replicates (leaves)/treatment). The bars represent the standard deviation. **D)** 'Bonny Best'

tomato seedlings were placed in root growth pouches, watered with 10mL of either water or trehalose

for one day, then watered as needed for ten days. Root length was measured after ten days (t-test,

$P=.31$). The data represent three biological replicates with ten roots/treatment. Percent of new root

growth was calculated as %new root growth= (new growth/total root length)*100. The bars represent

the standard deviation. **E)** SPAD meter measurements of chlorophyll content in leaves following

trehalose treatment (2-way ANOVA Fishers LSD multiple comparison, * = $P<0.5$ or lower). The bars

represent the standard deviation.

Supplemental Figure 3. Trehalose treatment does not nutritionally enrich the soil environment for *R.*

***solanacearum*, and its protection is limited to root application. A)** Disease progress curve of plants

treated with trehalose or water, then infected with either wild-type *Rs* or a Δ*treA* mutant unable to

catabolize trehalose (one-way ANOVA of areas under the curve, WT H_2O vs. WT tre, $P=.0008$, Δ*treA* H_2O

vs. Δ*treA* tre, $P=.017$). The data represent three bioreps each containing 13-15 plants per treatment. The

bars represent the standard error. **B)** Disease development in plants sprayed once with 30 mM trehalose

or water, and then soil-soak inoculated with *Rs* 48 h later (ANOVA of AUC, Fisher's LSD multiple

comparisons to H_2O soil, H_2O spray, $P=.67$; tre spray, $P=.63$; tre soil, $P=.63$). The data represent three

biological replicates each containing fifteen plants per treatment. The bars represent the standard error.

C) Colonization of tomato leaves by the bacterial leaf spot pathogen *Xanthomonas gardneri*; plants were

dip-inoculated with *X. gardneri* 72 h after root treatment with water or 30 mM trehalose (Mann-

Whitney, day 3, *P*=.06; day 6, *P*=.61; day 9, *P*=08)*.* The data represent three biological replicates with

twelve plants/treatment/timepoint. The graph displays the geometric means; bars indicate the

geometric standard deviation of the data.

Supplemental Figure 4. Infusing leaves with trehalose did not trigger ROS production in 'Bonny Best'

tomato leaves. DAB staining of 'Bonny best' tomato leaves infused with water, 10^9 CFU/mL *Rs*, or 30

mM trehalose solution to assess the effect of trehalose treatment on ROS production. The data

represent three biological replicates, with four plants per biological replicate per treatment. Photos are

representative samples and images were uniformly sharpened 25% to increase contrast.

References

Abd El-Rahim, M. F., G. M. Fahmy and Z. M. Fahmy (1998). "Alterations in transpiration and stem
vascular tissues of two maize cultivars under conditions of water stress and late wilt disease." Plant
Pathology **47**(2): 216-223.
Alexandersson, E., L. Fraysse, S. Sjövall-Larsen, S. Gustavsson, M. Fellert, M. Karlsson, U. Johanson and P.
Kjellbom (2005). "Whole gene family expression and drought stress regulation of aquaporins." Plant Mol
Biol **59**(3): 469-484.
Alméras, E., S. Stolz, S. Vollenweider, P. Reymond, L. Mène-Saffrané and E. E. Farmer (2003). "Reactive
electrophile species activate defense gene expression in *Arabidopsis*." The Plant Journal **34**(2): 205-216.
Álvarez, B., E. G. Bosca and M. M. López (2010). "On the life of *Ralstonia solanacearum*, a destructive
bacterial plant pathogen." Technology and Education Topics in Applied Mcrobiology and Microbial
Biotechnology: 267–279.
An, C. and Z. Mou (2011). "Salicylic acid and its function in plant immunity." Journal of Integrative Plant
Biology **53**(6): 412-428.
Anders, S., P. T. Pyl and W. Huber (2014). "HTSeq—a Python framework to work with high-throughput
sequencing data." Bioinformatics **31**(2): 166-169.
Andrews, S. FastQC A Quality Control Tool for High Throughput Sequence Data, Babraham
Bioinformatics.
Asaf, S., A. L. Khan, M. A. Khan, Q. M. Imran, B.-W. Yun and I.-J. Lee (2017). "Osmoprotective functions
conferred to soybean plants via inoculation with *Sphingomonas* sp. *LK11* and exogenous trehalose."
Microbiological Research **205**: 135-145.
Avonce, N., B. Leyman, J. O. Mascorro-Gallardo, P. Van Dijck, J. M. Thevelein and G. Iturriaga (2004).
"The *Arabidopsis* trehalose-6-P synthase AtTPS1 gene is a regulator of glucose, abscisic acid, and stress
signaling." Plant Physiology **136**(3): 3649-3659.
Bae, H., E. Herman, B. Bailey, H.-J. Bae and R. Sicher (2005). "Exogenous trehalose alters *Arabidopsis*
transcripts involved in cell wall modification, abiotic stress, nitrogen metabolism, and plant defense."
Physiologia Plantarum **125**(1): 114-126.
Barrs, H. D. and P. E. Weatherley (1962). "A re-examination of the relative turgidity technique for
estimating water deficits in leaves." Australian Journal of Biological Sciences **15**(3): 413-428.
Beattie, G. A. (2011). "Water relations in the interaction of foliar bacterial pathogens with plants."
Annual Review of Phytopathology **49**(1): 533-555.
Beattie, G. A. and S. E. Lindow (1995). "The secret life of foliar bacterial pathogens on leaves." Annual
Review of Phytopathology **33**: 145-172.
Beckman, C. H. (2000). "Phenolic-storing cells: keys to programmed cell death and periderm formation
in wilt disease resistance and in general defence responses in plants?" Physiological and Molecular Plant
Pathology **57**(3): 101-110.
Benjamini, Y. and Y. Hochberg (1995). "Controlling the false discovery rate: A practical and powerful
approach to multiple testing." Journal of the Royal Statistical Society. Series B (Methodological) **57**(1):
289-300.
Bian, C., Y. Duan, J. Wang, Q. Xiu, J. Wang, Y. Hou, X. Song and M. Zhou (2020). "Validamycin A induces
broad-spectrum resistance involving salicylic acid and jasmonic acid/ethylene signaling pathways."
Molecular Plant-Microbe Interactions **33**(12): 1424-1437.
Block, A., E. Schmelz, P. J. O'Donnell, J. B. Jones and H. J. Klee (2005). "Systemic acquired tolerance to
virulent bacterial pathogens in tomato." Plant Physiology **138**(3): 1481-1490.
Boatwright, J. L. and K. Pajerowska-Mukhtar (2013). "Salicylic acid: an old hormone up to new tricks."
Molecular Plant Pathology **14**(6): 623-634.

Bolger, A. M., M. Lohse and B. Usadel (2014). "Trimmomatic: a flexible trimmer for Illumina sequence
data." Bioinformatics **30**(15): 2114-2120.
Bolouri Moghaddam, M. R. and W. Van den Ende (2012). "Sugars and plant innate immunity." Journal of
Experimental Botany **63**(11): 3989-3998.
Brading, P. A., K. E. Hammond-Kosack, A. Parr and J. D. G. Jones (2000). "Salicylic acid is not required for
Cf-2- and *Cf-9*-dependent resistance of tomato to *Cladosporium fulvum*." The Plant Journal **23**(3): 305-
318.
Brown, D. G., J. K. Swanson and C. Allen (2007). "Two host-induced *Ralstonia solanacearum* genes, *acrA*
and *dinF*, encode multidrug efflux pumps and contribute to bacterial wilt virulence." Applied and
Environmental Microbiology **73**(9): 2777-2786.
Cao, J. and X. Li (2015). "Identification and phylogenetic analysis of late embryogenesis abundant
proteins family in tomato (*Solanum lycopersicum*)." Planta **241**(3): 757-772.
Chen, K., G.-J. Li, R. A. Bressan, C.-P. Song, J.-K. Zhu and Y. Zhao (2020). "Abscisic acid dynamics,
signaling, and functions in plants." Journal of Integrative Plant Biology **62**(1): 25-54.
Chen, Y. Y., Y. M. Lin, T. C. Chao, J. F. Wang, A. C. Liu, F. I. Ho and C. P. Cheng (2009). "Virus-induced gene
silencing reveals the involvement of ethylene-, salicylic acid- and mitogen-activated protein kinase-
related defense pathways in the resistance of tomato to bacterial wilt." Physiologia Plantarum **136**(3):
324-335.
Chen, Z., Z. Zheng, J. Huang, Z. Lai and B. Fan (2009). "Biosynthesis of salicylic acid in plants." Plant
signaling & behavior **4**(6): 493-496.
Choudhary, D. K., A. Prakash and B. N. Johri (2007). "Induced systemic resistance (ISR) in plants:
mechanism of action." Indian Journal of Microbiology **47**(4): 289-297.
Cortina, C. and F. A. Culiáñez-Macià (2005). "Tomato abiotic stress enhanced tolerance by trehalose
biosynthesis." Plant Science **169**(1): 75-82.
Daugherty, M. P., J. R. S. Lopes and R. P. P. Almeida (2010). "Strain-specific alfalfa water stress induced
by *Xylella fastidiosa*." European Journal of Plant Pathology **127**(3): 333-340.
De Lorenzo, G., S. Ferrari, F. Cervone and E. Okun (2018). "Extracellular DAMPs in plants and mammals:
Immunity, tissue damage and repair." Trends in Immunology **39**(11): 937-950.
Denancé, N., P. Ranocha, N. Oria, X. Barlet, M.-P. Rivière, K. A. Yadeta, L. Hoffmann, F. Perreau, G.
Clément, A. Maia-Grondard, G. C. M. van den Berg, B. Savelli, S. Fournier, Y. Aubert, S. Pelletier, B. P. H.
J. Thomma, A. Molina, L. Jouanin, Y. Marco and D. Goffner (2013). "*Arabidopsis wat1* (*walls are thin1*)-
mediated resistance to the bacterial vascular pathogen, *Ralstonia solanacearum*, is accompanied by
cross-regulation of salicylic acid and tryptophan metabolism." The Plant Journal **73**(2): 225-239.
Denny, T. P. (2006). Plant pathogenic *Ralstonia* species. Plant-Associated Bacteria. S. S. Gnanamanickam,
Springer**:** 573–644.
Derksen, H., C. Rampitsch and F. Daayf (2013). "Signaling cross-talk in plant disease resistance." Plant
Science **207**: 79-87.
Deslandes, L. and S. Genin (2014). "Opening the *Ralstonia solanacearum* type III effector tool box:
insights into host cell subversion mechanisms." Current Opinion in Plant Biology **20**: 110-117.
Deslandes, L., J. Olivier, Theuli, xe, F. res, xe, xe, ric, J. Hirsch, D. X. Feng, P. Bittner-Eddy, J. Beynon and Y.
Marco (2002). "Resistance to *Ralstonia solanacearum* in *Arabidopsis thaliana* is conferred by the
recessive RRS1-R gene, a member of a novel family of resistance genes." Proceedings of the National
Academy of Sciences **99**(4): 2404-2409.
Digonnet, C., Y. Martinez, N. Denance, M. Chasseray, P. Dabos, P. Ranocha, Y. Marco, A. Jauneau and D.
Goffner (2012). "Deciphering the route of *Ralstonia solanacearum* colonization in *Arabidopsis thaliana*
roots during a compatible interaction: focus at the plant cell wall." Planta **236**(5): 1419-1431.
Dobin, A., C. A. Davis, F. Schlesinger, J. Drenkow, C. Zaleski, S. Jha, P. Batut, M. Chaisson and T. R.
Gingeras (2012). "STAR: ultrafast universal RNA-seq aligner." Bioinformatics **29**(1): 15-21.

Elphinstone, J. G. (2005). The Current Bacterial Wilt Situation: A Global Overview. <u>Bacterial Wilt Disease</u>
<u>and the *Ralstonia solanacearum* Species Complex</u>. C. Allen, P. Prior and A. C. Hayward. St. Paul, MN, APS
Press.
Flores-Cruz, Z. and C. Allen (2009). "*Ralstonia solanacearum* encounters an oxidative environment
during tomato infection." <u>Molecular Plant-Microbe Interactions</u> **22**(7): 773–782.
French, E., B. S. Kim, K. Rivera-Zuluaga and A. S. Iyer-Pascuzzi (2018). "Whole root transcriptomic
analysis suggests a role for auxin pathways in resistance to *Ralstonia solanacearum* in tomato."
<u>Molecular Plant-Microbe Interactions</u> **31**(4): 432-444.
Garg, A. K., J.-K. Kim, T. G. Owens, A. P. Ranwala, Y. D. Choi, L. V. Kochian and R. J. Wu (2002). "Trehalose
accumulation in rice plants confers high tolerance levels to different abiotic stresses." <u>Proceedings of</u>
<u>the National Academy of Sciences</u> **99**(25).
Ge, S. X., D. Jung and R. Yao (2019). "ShinyGO: a graphical gene-set enrichment tool for animals and
plants." <u>Bioinformatics</u> **36**(8): 2628-2629.
Glazebrook, J. (2005). "Contrasting mechanisms of defense against biotrophic and necrotophic
pathogens." <u>Annual Review of Phytopathology</u> **43**(1): 205-227.
Godoy, J. A., J. M. Pardo and J. A. Pintor-Toro (1990). "A tomato cDNA inducible by salt stress and
abscisic acid: nucleotide sequence and expression pattern." <u>Plant Molecular Biology</u> **15**(5): 695-705.
Gomez, L. D., A. Gilday, R. Feil, J. E. Lunn and I. A. Graham (2010). "AtTPS1-mediated trehalose 6-
phosphate synthesis is essential for embryogenic and vegetative growth and responsiveness to ABA in
germinating seeds and stomatal guard cells." <u>The Plant Journal</u> **64**(1): 1-13.
Goodstein, D. M., S. Shu, R. Howson, R. Neupane, R. D. Hayes, J. Fazo, T. Mitros, W. Dirks, U. Hellsten, N.
Putnam and D. S. Rokhsar (2011). "Phytozome: a comparative platform for green plant genomics."
<u>Nucleic Acids Research</u> **40**(D1): D1178-D1186.
Hamilton, C. D., O. R. Steidl, A. M. MacIntyre, C. G. Hendrich and C. Allen (2021). "*Ralstonia*
solanacearum depends on catabolism of myo-inositol, sucrose, and trehalose for virulence in an
infection stage-dependent manner." <u>Molecular Plant-Microbe Interactions</u>.
Hase, S., S. Takahashi, S. Takenaka, K. Nakaho, T. Arie, S. Seo, Y. Ohashi and H. Takahashi (2008).
"Involvement of jasmonic acid signalling in bacterial wilt disease resistance induced by biocontrol agent
Pythium oligandrum in tomato." <u>Plant Pathology</u> **57**(5): 870-876.
Hassan, M. A. E. and K. A. M. Abo-Elyousr (2013). "Activation of tomato plant defence responses against
bacterial wilt caused by *Ralstonia solanacearum* using DL-3-aminobutyric acid (BABA)." <u>European</u>
<u>Journal of Plant Pathology</u> **136**(1): 145-157.
He, Y., E. J. Borrego, Z. Gorman, P.-C. Huang and M. V. Kolomiets (2020). "Relative contribution of
LOX10, green leaf volatiles and JA to wound-induced local and systemic oxylipin and hormone signature
in *Zea mays* (maize)." <u>Phytochemistry</u> **174**: 112334.
He, Y., H. Fukushige, D. F. Hildebrand and S. Gan (2002). "Evidence supporting a role of jasmonic acid in
Arabidopsis leaf senescence." <u>Plant Physiology</u> **128**(3): 876.
Hendrick, C. A. and L. Sequeira (1984). "Lipopolysaccharide defective mutants of the wilt pathogen
Pseudomonas solanacearum." <u>Applied and Environmental Microbiology</u> **48**(1): 94-101.
Hernández-Blanco, C., D. X. Feng, J. Hu, A. Sánchez-Vallet, L. Deslandes, F. Llorente, M. Berrocal-Lobo, H.
Keller, X. Barlet, C. Sánchez-Rodríguez, L. K. Anderson, S. Somerville, Y. Marco and A. Molina (2007).
"Impairment of cellulose synthases required for *Arabidopsis* secondary cell wall formation enhances
disease resistance." <u>The Plant Cell</u> **19**(3): 890.
Hirsch, J., L. Deslandes, D. X. Feng, C. Balagué and Y. Marco (2002). "Delayed symptom development in
ein2-1, an *Arabidopsis* ethylene-insensitive mutant, in response to bacterial wilt caused by *Ralstonia*
solanacearum." <u>Phytopathology</u> **92**(10): 1142-1148.
Hou, S., Z. Liu, H. Shen and D. Wu (2019). "Damage-Associated Molecular Pattern-triggered immunity in
plants." <u>Frontiers in Plant Science</u> **10**: 646.

Hu, J., X. Barlet, L. Deslandes, J. Hirsch, D. X. Feng, I. Somssich and Y. Marco (2008). "Transcriptional
responses of *Arabidopsis thaliana* during wilt disease caused by the soil-borne phytopathogenic
bacterium, *Ralstonia solanacearum*." PLOS ONE **3**(7): e2589-e2589.
Ingel, B., D. Caldwell, F. Duong, D. Y. Parkinson, K. A. McCulloh, A. S. Iyer-Pascuzzi, A. J. McElrone and T.
M. Lowe-Power (2021). "Revisiting the source of wilt symptoms: X-ray microcomputed tomography
provides direct evidence that *Ralstonia* biomass clogs xylem vessels." bioRxiv: 2021.2003.2019.436187.
Iordachescu, M. and R. Imai (2008). "Trehalose biosynthesis in response to abiotic stresses." Journal of
Integrative Plant Biology **50**(10): 1223-1229.
Ishihara, T., I. Mitsuhara, H. Takahashi and K. Nakaho (2012). "Transcriptome analysis of quantitative
resistance-specific response upon *Ralstonia solanacearum* infection in tomato." PLOS ONE **7**(10):
e46763.
Ishikawa, R., K. Fujimori and K. Matsuura (1996). "Antibacterial activity of Validamycin A against
Pseudomonas solanacearum and its efficacy against tomato bacterial wilt." Annals of the
Phytopathological Society of Japan **62**(5): 478-482.
Ishikawa, R., K. Shirouzu, H. Nakashita, H.-Y. Lee, T. Motoyama, I. Yamaguchi, T. Teraoka and T. Arie
(2005). "Foliar spray of Validamycin A or Validoxylamine A controls tomato *Fusarium* wilt."
Phytopathology **95**(10): 1209-1216.
Jackson, M. (1997). "Hormones from roots as signals for the shoots of stressed plants." Trends in Plant
Science **2**(1): 22-28.
Jacobs, J. M., L. Babujee, F. Meng, A. Milling and C. Allen (2012). "The *in planta* transcriptome of
Ralstonia solanacearum: conserved physiological and virulence strategies during bacterial wilt of
tomato." MBio **3**(4).
Jacobs, J. M., A. Milling, R. M. Mitra, C. S. Hogan, F. Ailloud, P. Prior and C. Allen (2013). "*Ralstonia
solanacearum* requires PopS, an ancient AvrE-family effector, for virulence and to overcome salicylic
acid-mediated defenses during tomato pathogenesis." MBio **4**(6): e00875-00813.
Jakab, G., V. Cottier, V. Toquin, G. Rigoli, L. Zimmerli, J.-P. Métraux and B. Mauch-Mani (2001). "β-
Aminobutyric acid-induced resistance in plants." European Journal of Plant Pathology **107**(1): 29-37.
Ji, K., W. Kai, B. Zhao, Y. Sun, B. Yuan, S. Dai, Q. Li, P. Chen, Y. Wang, Y. Pei, H. Wang, Y. Guo and P. Leng
(2014). "SlNCED1 and SlCYP707A2: key genes involved in ABA metabolism during tomato fruit ripening."
Journal of Experimental Botany **65**(18): 5243-5255.
Joosten, M. H. A. J. and P. J. G. M. De Wit (1989). "Identification of several Pathogenesis-Related
proteins in tomato leaves inoculated with *Cladosporium fulvum* (syn. *Fulvia fulva*) as 1,3-β-glucanases
and chitinases." Plant Physiology **89**(3): 945.
Kaiser, E., A. Morales, J. Harbinson, E. Heuvelink and L. F. M. Marcelis (2020). "High stomatal
conductance in the tomato *flacca* mutant allows for faster photosynthetic induction." Frontiers in Plant
Science **11**(1317).
Kapilan, R., M. Vaziri and J. J. Zwiazek (2018). "Regulation of aquaporins in plants under stress."
Biological Research **51**(1): 4.
Khater, M. A., M. G. Dawood, M. S. Sadak, M. A. F. Shalaby, M. E. El-Awadi and K. Gamal El-Din (2018).
"Enhancement of the performance of cowpea plants grown under drought conditions via trehalose
application." Middle East Journal of Agriculture Research **7**(3): 782-800.
Khokhani, D., T. M. Lowe-Power, T. M. Tran and C. Allen (2017). "A single regulator mediates strategic
switching between attachment/spread and growth/virulence in the plant pathogen *Ralstonia
solanacearum*." MBio **8**(5).
Khokhani, D., T. Tuan, T. Lowe-Power and C. Allen (2018). "Plant assays for quantifying *Ralstonia
solanacearum* virulence." Bio-Protocol **8**(18).

Kissoudis, C., A. Seifi, Z. Yan, A. T. Islam, H. van der Schoot, C. C. van de Wiel, R. G. Visser, C. G. van der
Linden and Y. Bai (2016). "Ethylene and abscisic acid signaling pathways differentially influence tomato
resistance to combined powdery mildew and salt stress." Frontiers in Plant Science **7**: 2009.
Kolbe, A., A. Tiessen, H. Schluepmann, M. Paul, S. Ulrich and P. Geigenberger (2005). "Trehalose 6-
phosphate regulates starch synthesis via posttranslational redox activation of ADP-glucose
pyrophosphorylase." Proceedings of the National Academy of Sciences **102**(31): 11118–11123.
Kolde, R. (2019). "pheatmap: Pretty Heatmaps."   Retrieved June 7, 2020, from https://cran.r-
project.org/web/packages/pheatmap/index.html.
Kushiro, T., M. Okamoto, K. Nakabayashi, K. Yamagishi, S. Kitamura, T. Asami, N. Hirai, T. Koshiba, Y.
Kamiya and E. Nambara (2004). "The Arabidopsis cytochrome P450 CYP707A encodes ABA 8'-
hydroxylases: key enzymes in ABA catabolism." The EMBO Journal **23**(7): 1647-1656.
Lanahan, M. B., H.-C. Yen, J. J. Giovannoni and H. J. Klee (1994). "The *Never Ripe* mutation blocks
ethylene perception in tomato." The Plant Cell **6**: 521-530.
Leyman, B., P. Van Dijck and J. M. Thevelein (2001). "An unexpected plethora of trehalose biosynthesis
genes in *Arabidopsis thaliana*." Trends in Plant Science **6**(11): 510-513.
Li, H., B. Handsaker, A. Wysoker, T. Fennell, J. Ruan, N. Homer, G. Marth, G. Abecasis, R. Durbin and S.
Genome Project Data Processing (2009). "The Sequence Alignment/Map format and SAMtools."
Bioinformatics **25**(16): 2078-2079.
Lin, Q., S. Wang, Y. Dao, J. Wang and K. Wang (2020). "The *Arabidopsis thaliana* trehalose-6-phosphate
phosphatase gene *AtTPPI* enhances drought tolerance by regulating stomatal apertures." Journal of
Experimental Botany.
Lin, Q., J. Yang, Q. Wang, H. Zhu, Z. Chen, Y. Dao and K. Wang (2019). "Overexpression of the trehalose-
6-phosphate phosphatase family gene *AtTPPF* improves the drought tolerance of *Arabidopsis thaliana*."
BMC Plant Biology **19**(1): 381.
Lin, Y., J. Zhang, W. Gao, Y. Chen, H. Li, D. W. Lawlor, M. J. Paul and W. Pan (2017). "Exogenous trehalose
improves growth under limiting nitrogen through upregulation of nitrogen metabolism." BMC Plant Biol
17(1): 247.
Ling, Q., W. Huang and P. Jarvis (2011). "Use of a SPAD-502 meter to measure leaf chlorophyll
concentration in *Arabidopsis thaliana*." Photosynth Res **107**(2): 209-214.
Liu, H., S. Zhang, M. A. Schell and T. P. Denny (2005). "Pyramiding unmarked deletions in *Ralstonia
solanacearum* shows that secreted proteins in addition to plant cell-wall-degrading enzymes contribute
to virulence." Molecular Plant-Microbe Interactions **18**(12): 1296-1305.
López-Gresa, M. P., P. Lisón, L. Yenush, V. Conejero, I. Rodrigo and J. M. Bellés (2016). "Salicylic acid is
involved in the basal resistance of tomato plants to Citrus Excortis Viroid and Tomato Spotted Wilt
Virus." PLOS ONE **11**(11): e0166938.
Love, M. I., W. Huber and S. Anders (2014). "Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2." Genome Biology **15**(12): 550.
Lowe, T. M., F. Ailloud and C. Allen (2015). "Hydroxycinnamic acid degradation, a broadly conserved
trait, protects *Ralstonia solanacearum* from chemical plant defenses and contributes to root
colonization and virulence." Molecular Plant-Microbe Interactions **28**(3): 286-297.
Lowe-Power, T. M., C. G. Hendrich, E. von Roepenack-Lahaye, B. Li, D. Wu, R. Mitra, B. L. Dalsing, P.
Ricca, J. Naidoo, D. Cook, A. Jancewicz, P. Masson, B. Thomma, T. Lahaye, A. J. Michael and C. Allen
(2018). "Metabolomics of tomato xylem sap during bacterial wilt reveals *Ralstonia solanacearum
produces abundant putrescine, a metabolite that accelerates wilt disease." Environmental Microbiology
20(4): 1330-1349.
Lowe-Power, T. M., J. M. Jacobs, F. Ailloud, B. Fochs, P. Prior and C. Allen (2016). "Degradation of the
plant defense signal salicylic acid protects *Ralstonia solanacearum* from toxicity and enhances virulence
on tobacco." MBio **7**(3).

Lowe-Power, T. M., D. Khokhani and C. Allen (2018). "How *Ralstonia solanacearum* exploits and thrives
in the flowing plant xylem environment." Trends Microbiol **26**(11): 929-942.
Lunn, J. E. (2007). "Gene families and evolution of trehalose metabolism in plants." Functional Plant
Biology **34**(6).
Lunn, J. E., I. Delorge, C. M. Figueroa, P. Van Dijck and M. Stitt (2014). "Trehalose metabolism in plants."
The Plant Journal **79**(4): 544-567.
Lunn, J. E., R. Feil, J. H. Hendriks, Y. Gibon, R. Morcuende, D. Osuna, W. R. Scheible, P. Carillo, M. R.
Hajirezaei and M. Stitt (2006). "Sugar-induced increases in trehalose 6-phosphate are correlated with
redox activation of ADP-glucose pyrophosphorylase and higher rates of starch synthesis in *Arabidopsis*
thaliana." Biochem J **397**(1): 139-148.
Lyu, J. I., S. R. Min, J. H. Lee, Y. H. Lim, J.-K. Kim, C.-H. Bae and J. R. Liu (2012). "Overexpression of a
trehalose-6-phosphate synthase/phosphatase fusion gene enhances tolerance and photosynthesis
during drought and salt stress without growth aberrations in tomato." Plant Cell, Tissue and Organ
Culture **112**(2): 257-262.
Ma, Y., I. Szostkiewicz, A. Korte, D. Moes, Y. Yang, A. Christmann and E. Grill (2009). "Regulators of PP2C
Phosphatase Activity Function as Abscisic Acid Sensors." Science **324**(5930): 1064.
MacIntyre, A. M., J. X. Barth, M. C. Pellitteri Hahn, C. O. Scarlett, S. Genin and C. Allen (2019). "Trehalose
synthesis contributes to osmotic stress tolerance and virulence of the bacterial wilt pathogen *Ralstonia*
solanacearum." Molecular Plant-Microbe Interactions: MPMI-08-19-0218-R.
Mandal, S., I. Kar, A. K. Mukherjee and P. Acharya (2013). "Elicitor-induced defense responses in
Solanum lycopersicum against *Ralstonia solanacearum*." The Scientific World Journal **2013**: 561056.
Mansfield, J., S. Genin, S. Magori, V. Citovsky, M. Sriariyanum, P. Ronald, M. Dow, V. Verdier, S. V. Beer,
M. A. Machado, I. Toth, G. Salmond and G. D. Foster (2012). "Top 10 plant pathogenic bacteria in
molecular plant pathology." Molecular Plant Pathology **13**(6): 614-629.
Marcos, R., Y. Izquierdo, T. Vellosillo, S. Kulasekaran, T. Cascón, M. Hamberg and C. Castresana (2015).
"9-Lipoxygenase-derived oxylipins activate brassinosteroid signaling to promote cell wall-based defense
and limit pathogen infection." Plant Physiology **169**(3): 2324-2334.
McElrone, A. J., J. L. Sherald and I. N. Forseth (2003). "Interactive effects of water stress and xylem-
limited bacterial infection on the water relations of a host vine." Journal of Experimental Botany
54(381): 419-430.
Mi, H., A. Muruganujan, D. Ebert, X. Huang and P. D. Thomas (2018). "PANTHER version 14: more
genomes, a new PANTHER GO-slim and improvements in enrichment analysis tools." Nucleic Acids
Research **47**(D1): D419-D426.
Milling, A., L. Babujee and C. Allen (2011). "*Ralstonia solanacearum* extracellular polysaccharide is a
specific elicitor of defense responses in wilt-resistant tomato plants." PLOS ONE **6**(1): e15853.
Montillet, J.-L., J.-P. Agnel, M. Ponchet, F. Vailleau, D. Roby and C. Triantaphylidès (2002).
"Lipoxygenase-mediated production of fatty acid hydroperoxides is a specific signature of the
hypersensitive reaction in plants." Plant Physiology and Biochemistry **40**(6): 633-639.
Mori, Y., K. Inoue, K. Ikeda, H. Nakayashiki, C. Higashimoto, K. Ohnishi, A. Kiba and Y. Hikichi (2016). "The
vascular plant-pathogenic bacterium *Ralstonia solanacearum* produces biofilms required for its
virulence on the surfaces of tomato cells adjacent to intercellular spaces." Molecular Plant Pathology
17(6): 890-902.
Müller, M. and S. Munné-Bosch (2011). "Rapid and sensitive hormonal profiling of complex plant
samples by liquid chromatography coupled to electrospray ionization tandem mass spectrometry." Plant
Methods **7**(1): 37.
Orgambide, G., H. Montrozier, P. Servin, J. Roussel, D. Trigalet-Demery and A. Trigaley (1991). "High
heterogeneity of the exopolysaccharides of *Pseudomonas solanacearum* strain GMI1000 and the
complete structure of the major polysaccharide." Journal of Biological Chemistry **266**(13): 8312-8321.

Osakabe, Y., K. Osakabe, K. Shinozaki and L.-S. Tran (2014). "Response of plants to water stress." Frontiers in Plant Science **5**: 86.

Park, J., Y. Lee, E. Martinoia and M. Geisler (2017). "Plant hormone transporters: what we know and what we would like to know." BMC biology **15**(1): 93-93.

Patel, J. and A. Mishra (2021). "Plant aquaporins alleviate drought tolerance in plants by modulating cellular biochemistry, root-architecture, and photosynthesis." Physiologia Plantarum **172**(2): 1030-1044.

Paul, M. (2007). "Trehalose 6-phosphate." Current Opinion in Plant Biology **10**(3): 303-309.

Paul, M. J., A. Gonzalez-Uriarte, C. A. Griffiths and K. Hassani-Pak (2018). "The role of trehalose 6-phosphate in crop yield and resilience." Plant Physiology **177**(1): 12-23.

Paul, M. J., L. F. Primavesi, D. Jhurreea and Y. Zhang (2008). "Trehalose metabolism and signaling." Annu Rev Plant Biol **59**: 417-441.

Peña-Cortés, H., J. Fisahn and L. Willmitzer (1995). "Signals involved in wound-induced proteinase inhibitor II gene expression in tomato and potato plants." Proceedings of the National Academy of Sciences **92**: 4106-4113.

Piazza, A., T. Zimaro, B. S. Garavaglia, F. A. Ficarra, L. Thomas, C. Marondedze, R. Feil, J. E. Lunn, C. Gehring, J. Ottado and N. Gottig (2015). "The dual nature of trehalose in citrus canker disease: a virulence factor for *Xanthomonas citri* subsp. *citri* and a trigger for plant defence responses." Journal of Experimental Botany **66**(9): 2795-2811.

Potnis, N., J. P. Soto-Arias, K. N. Cowles, A. H. C. van Bruggen, J. B. Jones and J. D. Barak (2014). "*Xanthomonas perforans* colonization influences *Salmonella enterica* in the tomato phyllosphere." Applied and environmental microbiology **80**(10): 3173-3180.

Pradhanang, P. M., P. Ji, M. T. Momol, S. M. Olson, J. L. Mayfield and J. B. Jones (2005). "Application of Acibenzolar-S-Methyl enhances host resistance in tomato against *Ralstonia solanacearum*." Plant Disease **89**(9): 989-993.

Prost, I., S. Dhondt, G. Rothe, J. Vicente, M. J. Rodriguez, N. Kift, F. Carbonne, G. Griffiths, M.-T. Esquerré-Tugayé, S. Rosahl, C. Castresana, M. Hamberg and J. Fournier (2005). "Evaluation of the antimicrobial activities of plant oxylipins supports their involvement in defense against pathogens." Plant Physiology **139**(4): 1902.

R. "R: The R Project for Statistical Computing." Retrieved June 7, 2020, from https://www.r-project.org/.

Raghavendra, A. S., V. K. Gonugunta, A. Christmann and E. Grill (2010). "ABA perception and signalling." Trends in Plant Science **15**(7): 395-401.

Ranf, S., L. Eschen-Lippold, P. Pecher, J. Lee and D. Scheel (2011). "Interplay between calcium signalling and early signalling elements during defence responses to microbe- or damage-associated molecular patterns." The Plant Journal **68**(1): 100-113.

Rau, A., G. Marot and F. Jaffrézic (2014). "Differential meta-analysis of RNA-seq data from multiple studies." BMC Bioinformatics **15**(1): 91.

Reignault, P., A. Cogan, J. Muchembled, A. Lounes-Hadj Sahraoui, R. Durand and M. Sancholle (2000). "Trehalose induces resistance to powdery mildew in wheat." New Phytologist **149**(3): 519-529.

Renard-Merlier, D., B. Randoux, E. Nowak, F. Farcy, R. Durand and P. Reignault (2007). "Iodus 40, salicylic acid, heptanoyl salicylic acid and trehalose exhibit different efficacies and defence targets during a wheat/powdery mildew interaction." Phytochemistry **68**(8): 1156-1164.

Retig, N. and I. Chet (1974). "Catechol-induced resistance of tomato plants to *Fusarium* wilt." Physiological Plant Pathology **4**(4): 469-475.

Reuscher, S., M. Akiyama, C. Mori, K. Aoki, D. Shibata and K. Shiratake (2013). "Genome-wide identification and expression analysis of aquaporins in tomato." PLOS ONE **8**(11): e79052.

Robinson, M. D., D. J. McCarthy and G. K. Smyth (2009). "edgeR: a Bioconductor package for differential expression analysis of digital gene expression data." Bioinformatics **26**(1): 139-140.

Romero, C., J. M. Bellés, J. L. Vayá, R. Serrano and A. Francisco (1997). "Expression of the yeast
trehalose-6-phosphate synthase gene in transgenic tobacco plants: Pleiotropic phenotypes include
drought tolerance." Planta **201**(3): 293-297.
Sade, N., B. J. Vinocur, A. Diber, A. Shatil, G. Ronen, H. Nissan, R. Wallach, H. Karchi and M. Moshelion
(2009). "Improving plant stress tolerance and yield production: is the tonoplast aquaporin SlTIP2;2 a key
to isohydric to anisohydric conversion?" New Phytologist **181**(3): 651-661.
Saito, S., N. Hirai, C. Matsumoto, H. Ohigashi, D. Ohta, K. Sakata and M. Mizutani (2004). "Arabidopsis
CYP707As encode (+)-abscisic acid 8'-hydroxylase, a key enzyme in the oxidative catabolism of abscisic
acid." Plant Physiology **134**(4): 1439-1449.
Sato, S., S. Tabata, H. Hirakawa and e. al (2012). "The tomato genome sequence provides insights into
fleshy fruit evolution." Nature **485**(7400): 635-641.
Savatin, D. V., G. Gramegna, V. Modesti and F. Cervone (2014). "Wounding in the plant tissue: the
defense of a dangerous passage." Frontiers in Plant Science **5**: 470.
Schluepmann, H., L. Berke and G. F. Sanchez-Perez (2012). "Metabolism control over growth: a case for
trehalose-6-phosphate in plants." Journal of Experimental Botany **63**(9): 3379-3390.
Schluepmann, H., T. Pellny, A. van Dijken, S. Smeekens and M. Paul (2003). "Trehalose 6-phosphate is
indispensable for carbohydrate utilization and growth in *Arabidopsis thaliana*." Proceedings of the
National Academy of Sciences **100**(11): 6849.
Seo, S., K. Nakaho, S. W. Hong, H. Takahashi, H. Shigemori and I. Mitsuhara (2016). "L-Histidine induces
resistance in plants to the bacterial pathogen *Ralstonia solanacearum* partially through the activation of
ethylene signaling." Plant Cell Physiol **57**(9): 1932-1942.
Shi, Y., H. Sun, X. Wang, W. Jin, Q. Chen, Z. Yuan and H. Yu (2019). "Physiological and transcriptomic
analyses reveal the molecular networks of responses induced by exogenous trehalose in plant." PLOS
ONE **14**(5): e0217204.
Shinozaki, K., K. Yamaguchi-Shinozaki and M. Seki (2003). "Regulatory network of gene expression in the
drought and cold stress responses." Current Opinion in Plant Biology **6**(5): 410-417.
Signorelli, S., L. P. Tarkowski, W. Van den Ende and D. C. Bassham (2019). "Linking Autophagy to Abiotic
and Biotic Stress Responses." Trends in Plant Science **24**(5): 413-430.
Singh, V. and J. Shah (2012). "Tomato responds to green peach aphid infestation with the activation of
trehalose metabolism and starch accumulation." Plant Signal Behav **7**(6): 605-607.
Sinha, R., A. Gupta and M. Senthil-Kumar (2016). "Understanding the impact of drought on foliar and
xylem invading bacterial pathogen stress in chickpea." Frontiers in Plant Science **7**: 902.
Sinha, R., A. Gupta and M. Senthil-Kumar (2017). "Concurrent drought stress and vascular pathogen
infection induce common and distinct transcriptomic responses in chickpea." Frontiers in Plant Science
8(333).
Smart, C. D., K. L. Myers, S. Restrepo, G. B. Martin and W. E. Fry (2003). "Partial resistance of tomato to
Phytophthora infestans is not dependent upon ethylene, jasmonic acid, or salicylic acid signaling
pathways." Molecular Plant-Microbe Interactions **2**: 141-148.
Smeekens, S. (2017). "Drought resistance: Spraying for yield." Nature Plants **3**(3).
Song, L., S.-s. C. Huang, A. Wise, R. Castanon, J. R. Nery, H. Chen, M. Watanabe, J. Thomas, Z. Bar-Joseph
and J. R. Ecker (2016). "A transcription factor hierarchy defines an environmental stress response
network." Science **354**(6312): aag1550.
Spoel, S. H., A. Koornneef, S. M. C. Claessens, J. P. Korzelius, J. A. Van Pelt, M. J. Mueller, A. J. Buchala, J.-
P. Métraux, R. Brown, K. Kazan, L. C. Van Loon, X. Dong and C. M. J. Pieterse (2003). "NPR1 modulates
cross-talk between salicylate- and jasmonate-dependent defense pathways through a novel function in
the cytosol." The Plant Cell **15**(3): 760.
Tal, M. (1966). "Abnormal stomatal behavior in wilty mutants of tomato." Plant Physiology **41**(8): 1387-
1391.

Tanaka, K. and M. Heil (2021). "Damage-Associated Molecular Patterns (DAMPs) in plant innate
immunity: applying the danger model and evolutionary perspectives." <u>Annual Review of Phytopathology</u>
59(1): 53-75.
Tans-Kersten, J., H. Huang and C. Allen (2001). "*Ralstonia solanacearum* needs motility for invasive
virulence on tomato." <u>J Bacteriol</u> **183**(12): 3597-3605.
Thompson, A. J., E. T. Thorne, A. Burbidge, A. C. Jackson, R. E. Sharp and I. B. Taylor (2004).
"Complementation of *notabilis*, an abscisic acid-deficient mutant of tomato: importance of sequence
context and utility of partial complementation." <u>Plant, Cell & Environment</u> **27**(4): 459-471.
Thordal-Christensen, H., Z. Zhang, Y. Wei and D. B. Collinge (1997). "Subcellular localization of H_2O_2 in
plants. H_2O_2 accumulation in papillae and hypersensitive response during the barley—powdery mildew
interaction." <u>The Plant Journal</u> **11**(6): 1187-1194.
Tian, T., Y. Liu, H. Yan, Q. You, X. Yi, Z. Du, W. Xu and Z. Su (2017). "AgriGO v2.0: a GO analysis toolkit for
the agricultural community, 2017 update." <u>Nucleic Acids Research</u> **45**(W1): W122-W129.
Tornero, P., J. Gadea, V. Conejero and P. Vera (1997). "Two PR-1 genes from tomato are differentially
regulated and reveal a novel mode of expression for a pathogenesis-related gene during the
hypersensitive response and development." <u>Molecular Plant-Microbe Interactions</u> **5**: 624-634.
Tran, T., A. MacIntyre, M. C. Hawes and C. Allen (2016). "Escaping underground Nets: Extracellular
DNases degrade plant extracellular traps and contribute to virulence of the plant pathogenic bacterium
Ralstonia solanacearum." <u>PLOS Pathogens</u> **12**(6): e1005686.
Tran, T., A. MacIntyre, D. Khokhani, M. C. Hawes and C. Allen (2016). "Extracellular DNases of *Ralstonia*
solanacearum modulate biofilms and facilitate bacterial wilt virulence." <u>Environmental Microbiology</u>
18(11): 4103-4117.
Umezawa, T., M. Okamoto, T. Kushiro, E. Nambara, Y. Oono, M. Seki, M. Kobayashi, T. Koshiba, Y.
Kamiya and K. Shinozaki (2006). "CYP707A3, a major ABA 8'-hydroxylase involved in dehydration and
rehydration response in Arabidopsis thaliana." <u>The Plant Journal</u> **46**(2): 171-182.
Vallad, G. E. and R. M. Goodman (2004). "Systemic acquired resistance and induced systemic resistance
in conventional agriculture." <u>Crop Science</u> **44**(6): 1920-1934.
Vandesteene, L., L. Lopez-Galvis, K. Vanneste, R. Feil, S. Maere, W. Lammens, F. Rolland, J. E. Lunn, N.
Avonce, T. Beeckman and P. Van Dijck (2012). "Expansive evolution of the trehalose-6-phosphate
phosphatase gene family in *Arabidopsis*." <u>Plant Physiology</u> **160**(2): 884-896.
Vandesteene, L., M. Ramon, K. Le Roy, P. Van Dijck and F. Rolland (2010). "A single active trehalose-6-P
synthase (TPS) and a family of putative regulatory TPS-like proteins in *Arabidopsis*." <u>Molecular Plant</u>
3(2): 406-419.
Vasse, J., P. Frey and A. Trigalet (1995). "Microscopic studies of intercellular infection and protoxylem
invasion of tomato roots by *Pseudomonas solanacearum*." <u>Molecular Plant-Microbe Interactions</u> **8**(2):
241-251.
Vicente, J., T. Cascón, B. Vicedo, P. García-Agustín, M. Hamberg and C. Castresana (2012). "Role of 9-
lipoxygenase and α-dioxygenase oxylipin pathways as modulators of local and systemic defense."
<u>Molecular Plant</u> **5**(4): 914-928.
Walters, D. R., T. Cowley and H. Weber (2006). "Rapid accumulation of trihydroxy oxylipins and
resistance to the bean rust pathogen *Uromyces fabae* following wounding in *Vicia faba*." <u>Annals of</u>
<u>Botany</u> **97**(5): 779-784.
Wang, K.-D., E. J. Borrego, C. M. Kenerley and M. V. Kolomiets (2020). "Oxylipins other than jasmonic
acid are xylem resident signals regulating systemic resistance induced by *Trichoderma virens* in maize."
<u>The Plant Cell</u> **32**(1): 166.
Wang, W., Q. Chen, S. Xu, W.-C. Liu, X. Zhu and C.-P. Song (2020). "Trehalose-6-phosphate phosphatase
e modulates ABA-controlled root growth and stomatal movement in *Arabidopsis*." <u>Journal of Integrative</u>
<u>Plant Biology</u>.

Wickham, H. (2016). "ggplot2: Elegant graphics for data analysis." from https://ggplot2-book.org/.
Widhalm, Joshua R. and N. Dudareva (2015). "A familiar ring to it: biosynthesis of plant benzoic acids."
Molecular Plant **8**(1): 83-97.
Yamaguchi-Shinozaki, K. and K. Shinozaki (1993). "The plant hormone abscisic acid mediates the
drought-induced expression but not the seed-specific expression of *rd22*, a gene responsive to
dehydration stress in *Arabidopsis thaliana*." Molecular and General Genetics MGG **238**(1): 17-25.
Yao, J. and C. Allen (2006). "Chemotaxis is required for virulence and competitive fitness of the bacterial
wilt pathogen *Ralstonia solanacearum*." J Bacteriol **188**(10): 3697-3708.
Yao, J. and C. Allen (2007). "The plant pathogen *Ralstonia solanacearum* needs aerotaxis for normal
biofilm formation and interactions with its tomato host." J Bacteriol **189**(17): 6415-6424.
Yu, W., R. Zhao, L. Wang, S. Zhang, R. Li, J. Sheng and L. Shen (2019). "ABA signaling rather than ABA
metabolism is involved in trehalose-induced drought tolerance in tomato plants." Planta **250**(2): 643-
655.
Zhang, H., Y. Hong, L. Huang, S. Liu, L. Tian, Y. Dai, Z. Cao, L. Huang, D. Li and F. Song (2016). "Virus-
Induced Gene Silencing-based functional analyses revealed the involvement of several putative
trehalose-6-phosphate synthase/phosphatase genes in disease resistance against *Botrytis cinerea* and
Pseudomonas syringae pv. *tomato* DC3000 in tomato." Frontiers in Plant Science **7**(1176).
Zhao, C., H. Wang, Y. Lu, J. Hu, L. Qu, Z. Li, D. Wang, Y. He, M. Valls, N. S. Coll, Q. Chen and H. Lu (2019).
"Deep sequencing reveals early reprogramming of *Arabidopsis* root transcriptomes upon *Ralstonia*
solanacearum infection." Molecular Plant-Microbe Interactions® **32**(7): 813-827.
Zhao, Y., R. Thilmony, C. L. Bender, A. Schaller, S. Y. He and G. A. Howe (2003). "Virulence systems of
Pseudomonas syringae pv. *tomato* promote bacterial speck disease in tomato by targeting the
jasmonate signaling pathway." The Plant Journal **36**(4): 485-499.
Zhou, R., X. Yu, T. Zhao, C.-O. Ottosen, E. Rosenqvist and Z. Wu (2019). "Physiological analysis and
transcriptome sequencing reveal the effects of combined cold and drought on tomato leaf." BMC Plant
Biology **19**(1): 377.
Zimmerman, D. C. and C. A. Coudron (1979). "Identification of traumatin, a wound hormone, as 12-oxo-
trans-10-dodecenoic acid." Plant Physiology **63**(3): 536.

1412