1

Trehalose induces tomato defenses and increases drought tolerance and bacterial wilt disease resistance

April M. MacIntyre[1,2], Valerian Meline[3], Zachary Gorman[4], Steven P. Augustine[5], Carolyn J. Dye[6],
Corri D. Hamilton[1], Anjali S. Iyer-Pascuzzi[3], Michael V. Kolomiets[4], Katherine A. McCulloh[5], and
Caitilyn Allen[1*]

[1]University of Wisconsin-Madison Department of Plant Pathology
[2]University of Wisconsin-Madison Department of Bacteriology
[3]Purdue University, Department of Botany and Plant Pathology
[4]Texas A&M University, Department of Plant Pathology and Microbiology
[5] University of Wisconsin-Madison Department of Botany
[6] Soil Carbon Co., St. Paul, MN

21
22
23
*Corresponding author: callen@wisc.edu   +1-608-556-3369

Abstract

Ralstonia solanacearum causes plant bacterial wilt disease, leading to severe crop losses. Xylem

sap from *R. solanacearum*-infected tomato is enriched in host produced trehalose. Water stressed

plants accumulate the disaccharide trehalose, which increases drought tolerance via abscisic acid (ABA)

signaling networks. Because infected plants have reduced water flow, we hypothesized that bacterial

wilt physiologically mimics drought stress, which trehalose could mitigate. Transcriptomic responses of

susceptible vs. resistant tomato plants to *R. solanacearum* infection revealed differential expression of

drought-associated genes, including those involved in ABA and trehalose metabolism. ABA was enriched

in xylem sap from *R. solanacearum*-infected plants. Treating roots with ABA lowered stomatal

conductance and reduced *R. solanacearum* stem colonization. Treating roots with trehalose increased

ABA in xylem sap and reduced plant water use by reducing stomatal conductance and temporarily

improving water use efficiency. Further, trehalose-treated plants were more resistant to bacterial wilt

disease. Trehalose treatment also upregulated expression of salicylic acid (SA)-dependent defense

genes, increased xylem sap levels of SA and other antimicrobial compounds, and increased wilt

resistance of SA-insensitive *NahG* tomato plants. Additionally, trehalose treatment increased xylem

concentrations of jasmonic acid and related oxylipins. Together, these data show that exogenous

trehalose reduced both water stress and bacterial wilt disease and triggered systemic resistance. This

suite of responses revealed unexpected linkages between plant responses to biotic and abiotic stress

and suggests that that *R. solanacearum*-infected tomato plants produce more trehalose to improve

water use efficiency and increase wilt disease resistance. In turn, *R. solanacearum* degrades trehalose as

a counter-defense.

Keywords: *Ralstonia solanacearum,* bacterial wilt disease, drought tolerance, trehalose, trehalase, plant

hormones, xylem water physiology

Introduction

*Ralstonia solanacearum*is a vascular plant pathogen that causes bacterial wilt disease (BW) in

over 200 species of plants worldwide (Denny 2006). It is widely studied because it is economically

damaging to food, fiber, and feed production (Elphinstone 2005, Mansfield, Genin et al. 2012). *R.*

solanacearum is also notoriously difficult to eradicate because it persists long-term in soil, water, and

latently infected hosts (Álvarez, Bosca et al. 2010).

Most *R. solanacearum* strains infect plants through the soil (Elphinstone 2005). *R. solanacearum*

chemo- or aero-taxes towards roots, evades root border cell traps, enters roots at points of lateral root

emergence, moves to the root cortex, and eventually migrates to the xylem, all with the help of secreted

proteins such as hydrolytic enzymes and Type III secreted effectors (Vasse, Frey et al. 1995, Tans-

Kersten, Huang et al. 2001, Liu, Zhang et al. 2005, Yao and Allen 2006, Yao and Allen 2007, Digonnet,

Martinez et al. 2012, Deslandes and Genin 2014, Tran, MacIntyre et al. 2016). Once in xylem vessels, *R.*

solanacearum forms biofilm matrices composed of extracellular DNA and polysaccharide (EPS) and

proliferates to >10^9 CFU/g of stem tissue (Orgambide, Montrozier et al. 1991, Mori, Inoue et al. 2016,

Tran, MacIntyre et al. 2016). *R. solanacearum* also conditions the xylem environment to favor bacterial

growth (Lowe-Power, Hendrich et al. 2018). The large amounts of bacteria and EPS clog xylem vessels

and disrupt water flow (Ingel, Caldwell et al. 2021). If the plant cannot contain bacterial growth and

spread, it will wilt and eventually die.

Researchers long assumed that *R. solanacearum* and other vascular pathogens cause drought-

like physiological symptoms in host plants and that severity of the disease is affected by environmental

water availability (Abd El-Rahim, Fahmy et al. 1998, McElrone, Sherald et al. 2003, Daugherty, Lopes et

al. 2010). *R. solanacearum* dies more quickly in drought-stressed chickpea plants, and *R. solanacearum*

infection and water stress induce similar osmolyte biosynthesis and expression of genes related to the

plant signaling molecule abscisic acid (ABA), a well-characterized marker of water stress (Sinha, Gupta et

al. 2016, Sinha, Gupta et al. 2017). Further, stomatal conductance decreases in *R. solanacearum* infected

plants (Lowe-Power, Hendrich et al. 2018). These studies suggest but do not prove that plant response

to BW mimics the drought stress response.

During water and osmotic stress, many plants, including tomato, accumulate trehalose and

trehalose-6-phosphate (T6P), which contribute to tolerance of these stresses (Romero, Bellés et al.

1997, Garg, Kim et al. 2002, Cortina and Culiáñez-Macià 2005, Lyu, Min et al. 2012, Khater, Dawood et

al. 2018). Xylem sap from *R. solanacearum*-infected tomato plants contains about 20-fold more

trehalose than healthy plants (Lowe-Power, Hendrich et al. 2018). This non-reducing disaccharide

composed of glucose and UDP-glucose is best known for its role as a compatible solute. Trehalose

synthesis helps *R. solanacearum* tolerate osmotic stress and contributes to pathogen fitness and

virulence (MacIntyre, Barth et al. 2019). However, the tomato host produces the trehalose in xylem sap

during infection; *R. solanacearum* does not export its trehalose in culture and it aggressively expresses

the secreted TreA trehalase *in planta* (Jacobs, Babujee et al. 2012).

Some studies suggest that trehalose may mediate host-microbe interactions and contribute to

plant defense. For example, treating *Arabidopsis* seedlings in liquid culture with 30 mM exogenous

trehalose triggered expression of defense genes in ethylene and methyl-jasmonate pathways (Bae,

Herman et al. 2005). Infusing tobacco leaves with a 10-50 mM trehalose solution increased resistance to

tobacco mosaic virus and triggered a transcriptomic response suggestive of biotic and abiotic stress

tolerance (Shi, Sun et al. 2019). A *Xanthomonas citri* trehalose synthesis mutant has reduced fitness and

infusing citrus leaves with a trehalose solution induced defense gene expression, suggesting that

trehalose might simultaneously be a virulence factor and an indicator of pathogen attack (Piazza, Zimaro

et al. 2015). In another study, spraying leaves with the trehalase inhibitor Validamycin A increased

tomato resistance to BW and *Fusarium* wilt and activated systemic acquired resistance against a wide

range of other pathogens in *Arabidopsis* (Ishikawa, Fujimori et al. 1996, Ishikawa, Shirouzu et al. 2005,

Bian, Duan et al. 2020).

We investigated the relationship between BW and drought stress in tomatoes and explored the

role of trehalose in this interaction. We tested the following hypotheses: 1) *R. solanacearum infection*

mimics abiotic stress triggered by disrupted water flow; and 2) *exogenous trehalose inhibits BW*

*development in tomato by sustaining plant water use efficiency and inducing defense response*s. To test

H_o1, we used meta-transcriptomic analyses to show that tomato genes involved in trehalose metabolism

and water stress responses are differentially expressed during *R. solanacearum* infection and between

susceptible and resistant tomato hosts. Using ABA treatment to close stomates, we found that *R.*

solanacearum depends on stomatal conductance and water flow for effective host colonization. We also

discovered that *R. solanacearum* infection increases xylem sap ABA levels.

Testing H_o2 revealed that treating plants with exogenous trehalose improved water use

efficiency in drought-stressed tomatoes, decreased short-term stomatal conductance and transpiration,

and increased ABA concentration in xylem sap. Trehalose-treated tomato plants were more resistant to

BW. This protective effect appeared to be mediated by up-regulation of salicylic acid (SA)- and jasmonic

acid (JA)-related defense genes and by increased levels of jasmonates and other defense compounds in

xylem sap.

Results

Bacterial wilt disease altered expression of tomato trehalose and water stress genes

118   To investigate the relationship between BW and drought stress, we measured tomato gene

expression related to trehalose metabolism and water stress responses. Candidate genes included those

known or predicted to encode synthesis and degradation of trehalose, including trehalose-6-phosphate

phosphatase (TPP), trehalose-6-phosphate synthase (TPS), and trehalase (Leyman, Van Dijck et al. 2001,

Vandesteene, Ramon et al. 2010, Vandesteene, Lopez-Galvis et al. 2012). We also focused on genes

encoding water stress-associated proteins such as: 1) those involved in biosynthesis, regulation, and

response to ABA, a known indicator of water stress (Song, Huang et al. 2016, Zhou, Yu et al. 2019); 2)

late embryogenesis/dehydrins (LEAs), which accumulate in water-stressed plants (Cao and Li 2015); and

3) aquaporins, channel proteins that allow living cells to move water and small solutes more efficiently

(Kapilan, Vaziri et al. 2018).

128   We used seven datasets that profiled transcriptomes of a wilt-resistant tomato breeding line

129   (Hawaii 7996) and two wilt-susceptible tomato plants (WV700 and Bonny Best) responding to infection

130   by various *R. solanacearum* strains (K60 and GMI1000) (French, Kim et al. 2018). After normalizing the

131   datasets to allow comparisons across these experiments, we measured the expression of these

132   candidate genes in healthy plants relative to their levels in *R. solanacearum*-infected plants.

133   Although the datasets came from diverse plant genotypes, tissues, and *R. solanacearum* strains,

this analysis revealed that *R. solanacearum*-infected tomato plants significantly changed expression of

multiple genes related to trehalose metabolism and water stress physiology (Fig. 1). Expression trends

varied across infected tissue types, cultivar disease resistance, and within gene families. Some genes in

these categories were differentially expressed in both resistant and susceptible infected tomato plants.

For example, resistant and susceptible plants upregulated an ABA hydroxylase and an ABA and

environmental stress-inducible protein TAS14 (Fig. 1C) and downregulated many diverse aquaporins

(Fig. 1D). In contrast, with the exception of TPS-7, many tomato trehalose synthase and phosphatase

genes (Fig. 1A) and LEA genes (Fig. 1B) were downregulated in resistant tomatoes, suggesting that

trehalose and LEA genes could be involved in resistance against *R. solanacearum.* These results led us to

directly measure physiological and metabolic indicators of drought stress in *R. solanacearum*-infected

plants.

Bacterial wilt disease increases ABA concentration in xylem sap

Plants can use ABA to mitigate drought stress. This plant hormone, synthesized locally in guard

cells or in the vasculature, closes stomates to reduce water loss through transpiration (Raghavendra,

Gonugunta et al. 2010). To test the hypothesis that tomato plants produce ABA in response to BW, we

used LCMS/MS to compare ABA levels in xylem sap of healthy plants and *R. solanacearum*-inoculated

plants showing early wilt symptoms. Xylem sap from the *R. solanacearum*-infected plants contained 40%

more ABA than sap from mock-inoculated plants (Fig. 2A).

***R. solanacearum* depends on normal host stomatal conductance to establish infection**

We previously observed that *R. solanacearum*-infected tomatoes have reduced stomatal

conductance (Lowe-Power, Khokhani et al. 2018). To test the hypothesis that the increased ABA levels

and lowered stomatal conductance in infected plants also affects BW development, we measured the

effects of exogenous ABA treatment on stomatal conductance, transpiration rates, and *R. solanacearum*

colonization of susceptible tomato plant stems. We watered tomato plants daily with 50 mL of either

100 µM ABA, 50 µM ABA, or water + DMSO as control and measured stomatal conductance four to six

hours later. After 48 h of ABA treatment, plants were drench-inoculated by adding a suspension of *R.*

solanacearum strain GMI1000 to the ABA treatment solutions, which we determined were not

nutritious or cytotoxic to *R. solanacearum* (Supp. Fig. 1). Treatment with ABA reliably reduced stomatal

conductance in the plants over the seven days of the experiment; 100 µM and 50 µM ABA treatment

had similar effects (Fig. 2B).

ABA treatments continued for four days after plants were inoculated with *R. solanacearum*. At

five dpi we measured *R. solanacearum* population sizes in the stems. Stems of ABA-treated plants were

less frequently colonized by *R. solanacearum* than control plants (50 µM ABA = 17% plants colonized;

100 µM = 23% plants colonized; water control = 53% plants colonized) and they contained lower *R.*

solanacearum population sizes than untreated plants (Fig. 2C). The correlation between ABA treatment,

reduced stomatal conductance, and lower *R. solanacearum* colonization in tomato stems suggests that

plant water movement assists *R. solanacearum* infection either at the roots or in the stem and that

reducing host water flow reduces or slows *R. solanacearum* spread in the plant.

Trehalose treatment transiently improves water use efficiency in tomatoes

Tomato plants become more drought tolerant following exposure to exogenous trehalose, via

the mechanisms of ABA signaling and stomatal closure (Yu, Zhao et al. 2019). Because ABA and trehalose

concentrations increase in xylem sap from *R. solanacearum*-infected plants and pathogen spread

depends on host xylem water transport, we asked whether exogenous trehalose could also protect

plants from bacterial wilt.

We first confirmed that trehalose increases tomato drought tolerance under our experimental

conditions. Two-week-old wilt-susceptible Bonny Best tomato plants were treated with 50 mL of 30 mM

trehalose solution or water and then did not water them again. Trehalose-treated plants remained

turgid for three days after control plants began wilting, showing that trehalose treatment delayed

wilting from lack of water under our conditions (Fig. 3A).

Next, we assessed stomatal conductance, transpiration, and photosynthesis of trehalose-treated

plants. After watering with 30 mM trehalose as above, we measured tomato leaf responses with a LI-

COR portable photosynthesis machine at 10 min, 6, 24, 48, and 72 h after treatment. Relative to

untreated plants, trehalose treatment detectably reduced stomatal conductance in 10 min, an effect

that persisted for at least 6 h. By 24 h after treatment, stomatal conductance of treated plants was

indistinguishable from that of water controls (Fig. 3B). However, transpiration in trehalose-treated

plants was lower than controls at 6, 24, and 48 h (Fig. 3C). Water use efficiency, the relationship

between water loss and plant productivity (photosynthesis/transpiration), was temporarily improved at

6 h by trehalose treatment (Supp. Fig. 2A). Quantitative mass spectrometry analysis of tomato xylem sap

revealed that ABA content was 5-fold higher in trehalose-treated plants at 24 h (Fig. 3D). However, at 6

and 48 h after treatment ABA levels were no different from those of water-treated control plants.

Trehalose treatment did not negatively affect photosynthesis (Supp. Fig. 2B), plant growth

determined by measuring plant mass (data not shown), relative water content (RWC), or seedling

growth. Tomato leaf RWC was the same in trehalose-treated and water-treated plants at 48h (Supp. Fig.

2C). Further, 1-3 days of watering with trehalose solution did not affect tomato seedling root growth *in*

vitro relative to water controls (Supp. Fig. 2D). Chlorophyll content, measured by SPAD meter, was

transiently lower in trehalose treated plants 2-10 days after treatment, although chlorophyll content

eventually returned to the same levels as in untreated plants (Supp. Fig. 2E). Together, these diverse

physiological and biochemical analyses showed that treating tomato plants with exogenous trehalose

temporarily increased their drought tolerance, likely because trehalose increased ABA levels and ABA

signaling closes stomates.

Trehalose is a potent inhibitor of bacterial wilt disease

Since we found that BW impacts host water stress physiology and that exogenous trehalose

reduces stomatal conductance in tomatoes, we wondered if this decreased water movement would

affect BW development. We tested the hypothesis that treating plants with trehalose would make them

more wilt resistant by watering tomato plants with 30 mM trehalose solution or water as above. After

48 h, treated plants were drench inoculated with a suspension of *R. solanacearum* bacteria. Bacterial

wilt incidence and severity was reduced by 25-75% across biological replicates in the trehalose-treated

plants, demonstrating that exogenous trehalose increases tomato resistance to BW (Fig. 4A). This

protective effect was enduring; trehalose treatment reduced BW incidence even when plants were

inoculated five days later (data not shown).

Since *R. solanacearum* can infect plants without causing visible symptoms, the absence of wilt in

trehalose treated plants could indicate either the lack of disease or latent infection. To distinguish

between these possibilities we measured *R. solanacearum* colonization of tomato midstems at 5, 6, and

7 dpi. Water-treated control plants were three times more likely to contain detectable *R. solanacearum*

cells than trehalose-treated plants (20.7% compared to 7.4%), and control plants contained significantly

higher bacterial population sizes (Fig. 4B). This showed that trehalose treatment did not increase the

frequency of latent infections but rather that *R. solanacearum* was less likely to colonize stems of

trehalose treated plants. The reduced colonization frequency at this relatively early time point is

consistent with the observed trehalose-associated delay in disease development.

We tested the protective limits of trehalose treatment by varying inoculation sites, treatment

methods, and pathogens. Interestingly, inoculating *R. solanacearum* cells directly into the xylem through

a cut petiole overwhelmed any protective benefit of trehalose treatment (Fig. 4C). This result suggests

that *R. solanacearum* must overcome a trehalose-enhanced checkpoint at or in tomato roots in order to

colonize plant stems.

R. solanacearum produces a trehalase enzyme and can use trehalose as a carbon source, so we

wondered if treating tomato roots with trehalose nutritionally enriched the soil environment*,* thereby

de-incentivizing *R. solanacearum* root invasion. We tested this hypothesis by inoculating trehalose-

treated tomato plants with either wild-type *R. solanacearum* or a Δ*treA* trehalase mutant that cannot

grow on trehalose (MacIntyre, Barth et al. 2019). Trehalose treatment similarly reduced disease caused

by wild type and Δ*treA R. solanacearum*, so bacterial catabolism of trehalose cannot explain trehalose

treatment's protective effect (Suppl. Fig. 3A).

Spraying trehalose solution onto leaves and then inoculating *R. solanacearum* by soil-soak 48 h

later also did not affect BW progress (Supp. Fig. 3B). To explore the effects of trehalose treatment on

another bacterial pathogen, we dip-inoculated tomato leaves with the tomato leaf spot pathogen

Xanthomonas gardneri 72 h after plant roots were treated with trehalose. Trehalose had no effect on

leafspot disease severity or *X. gardneri* colonization of leaves sampled 3, 6, and 9 dpi (Supp. Fig. 3C).

These experiments demonstrated that trehalose needs to be applied at the roots to protect against BW

and that trehalose might increase resistance only to *R. solanacearum.*

Exogenous trehalose increases the expression of tomato defense genes, and SA signaling may mediate

its protective effect

Although trehalose treatment of roots decreased plant transpiration and stomatal conductance

temporarily (6-48 h), this treatment protected plants from BW for at least five days, suggesting that

trehalose might also induced plant defense responses. We tested this by quantifying the effect of

trehalose treatment on defense gene expression, BW resistance of defense hormone-insensitive tomato

plants, and levels of hormones and defense compounds.

To measure the effect of trehalose on defense gene expression, we extracted RNA from tomato

stem sections 6 or 48 h after roots were treated with 30 mM trehalose solution or water. We measured

expression of two JA-responsive genes, *pin2* and *loxA* (Peña-Cortés, Fisahn et al. 1995, Zhao, Thilmony et

al. 2003). *Pin2* was significantly downregulated 6 h after trehalose treatment (Fig. 5A), suggesting JA

levels may have been affected by trehalose treatment, while *loxA* expression did not change after

trehalose treatment. G*luA*, *PR1a*, and *PR1b* are tomato pathogenesis-related (PR) genes induced by SA

and by pathogen challenge (Joosten and De Wit 1989, Tornero, Gadea et al. 1997, Block, Schmelz et al.

2005). Both *GluA* and *PR1a* were expressed 10-fold more in trehalose-treated plants compared to water

controls 6 h after treatment (Fig. 5A) and their expression increased to over 100-fold at 48 h (Fig. 5B).

PR1b and *osm,* which respond to the defense hormone ethylene (ET), were also significantly

upregulated 48 h after trehalose treatment (Tornero, Gadea et al. 1997, Smart, Myers et al. 2003). The

ET synthesis gene *ACO5* was also 10-fold more highly expressed compared to water controls 6 and 48 h

after trehalose treatment (Fig. 5A+B) (Jacobs, Milling et al. 2013). Transcripts of the ABA-responsive

genes *rd22* and *dhn-tas* were slightly but significantly upregulated 48 h after trehalose treatment,

possibly in response to the increased levels of ABA in xylem sap shown in Fig. 5B (Yamaguchi-Shinozaki

and Shinozaki 1993, Shinozaki, Yamaguchi-Shinozaki et al. 2003, Kissoudis, Seifi et al. 2016). Together,

these qRT-PCR experiments revealed that tomato plants respond to trehalose by increasing expression

of genes in multiple biotic and abiotic stress signal pathways, including those responding to JA, SA, ET,

and ABA.

The finding that SA- and ET-responsive tomato genes were upregulated 6 and 48 h after

trehalose treatment suggested that ET perception and SA responses might be necessary for trehalose-

mediated increase in BW resistance. We tested this hypothesis by treating with trehalose the roots of

ET-insensitive *NeverRipe* mutant tomatoes (cv. Pearson) and SA-degrading *NahG* transgenic tomatoes

(cv. Moneymaker). After 48 h, all plants were soil drench inoculated with *R. solanacearum*. As observed

for cv. Bonny Best, trehalose treatment increased BW resistance of Pearson and Moneymaker plants.

Trehalose also increased BW resistance of both the *NeverRipe* and *NahG* plants compared to untreated

controls (Fig. 5C and 5D). However, trehalose-mediated BW resistance does not require ET perception

since BW developed similarly on trehalose-treated *NeverRipe* and the parental 'Pearson' genotype

(*P*<.98). In contrast, trehalose-treated *NahG* tomatoes were significantly more resistant to BW than

trehalose-treated Money Maker parental line (*P*<.04). These results suggest that either a lack of

downstream SA responses or overaccumulation of SA degradation products contributes to the

trehalose-mediated increase in BW resistance. Alternatively, trehalose may improve resistance

downstream of SA responses.

More than half of the *NahG* tomato plants displayed a transient marginal leaf necrosis

phenotype following trehalose treatment (data not shown), which could have been due to increased

ROS production. To test this, we infused leaves from four-week-old wild type Bonny Best, Moneymaker,

or *NahG* tomato plants with water, *R. solanacearum* cells, or 30 mM trehalose solution and observed

ROS production with DAB staining. Infusing leaves with trehalose did not trigger excess ROS production

(Supp. Fig. 4), suggesting that another physiological effect caused the transient marginal necrosis in the

NahG plants.

Trehalose treatment promotes accumulation of hormones and defense compounds in xylem sap

To better understand the mechanisms of trehalose-mediated wilt resistance, we measured

levels of a suite of plant hormones, oxylipins, and defense-related compounds in xylem sap from

infected and trehalose-treated Bonny Best tomatoes. We harvested sap from tomatoes displaying a

disease index of 1 (DI=1, partially wilted, late-stage infection) and from uninoculated tomatoes at 6, 24,

and 48 h after trehalose treatment. Using targeted quantitative mass spectrometry, we measured

metabolites from the phenylpropanoid biosynthesis pathway, several oxylipins from the 9-lipoxygenase

and 13-lipoxygenase pathways, and other selected defense metabolites (Supp. Table 3).

Sap from *R. solanacearum*-infected plants and trehalose-treated plants had different levels of

targeted metabolites compared to sap from healthy tomatoes. Sap collected 24 h after trehalose

treatment had the highest number of enriched compounds. These were consistent with a shock or

wound response even though plants were not wounded (Vallad and Goodman 2004). Surprisingly,

trehalose treatment, but not *R. solanacearum* infection, increased SA concentration at 24 h (Fig. 6A).

This result was consistent with the possibility that trehalose plays a role in SA signaling and suggested

that SA concentration might not have a significant role in xylem sap at late-stage disease. Benzoic acid is

a potent antimicrobial compound produced through the phenylpropanoid pathway that can serve as a

direct precursor to SA (Chen, Zheng et al. 2009, Widhalm and Dudareva 2015). Benzoic acid levels

increased in xylem sap from both trehalose treated and *R. solanacearum* infected plants (Fig. 6B).

Interestingly, coumaric acid, another phenylpropanoid defense compound that *R. solanacearum* can

degrade, was significantly increased in sap from *R. solanacearum*-infected plants but not in sap from

trehalose treated plants (Fig. 6C) (Lowe, Ailloud et al. 2015). Levels of traumatic acid, which results from

the oxidation of traumatin produced by wounded plants, were higher in sap from both *R. solanacearum*-

infected plants and trehalose-treated plants at 24 h (Fig. 6D) (Zimmerman and Coudron 1979).

The chemical signatures exclusively present in trehalose treated plants were the persistent

increases in JA, the JA precursor, 12-OPDA, and JA-Ile, the biologically active form of JA (Fig. 6E-G). This

accumulation of jasmonates, together with the upregulation of JA and ET-pathway defense genes,

suggested that trehalose treatment induces a systemic defense response that contributes to its

protective effect (Vallad and Goodman 2004, Choudhary, Prakash et al. 2007).

Curiously, several defense-related oxylipins from the reductase and lipoxygenase branches of

the LOX pathway were enriched in sap from *R. solanacearum*-infected plants and 24 h trehalose treated

plants (Fig. 6H-K). These oxylipins include 9(*S*)-hydroxy-10(*E*),12(*Z*),15(*Z*)-octadecatrienoic acid (9-HOT,

Fig. 6H), 9(*S*)-hydroxy-10(*E*),12(*Z*)-octadecadienoic acid (9-HOD, Fig. 6I), 13(S)-hydroxy-9(Z),11(E)-

octadecatrienoic acid (13-HOD, Fig. 6J), and 13-oxo-9(*Z*),11(*E*)-octadecadienoic acid (13-KOD, Fig. 6K). All

these oxylipins have antimicrobial and signaling activity in defense against pathogens (Prost, Dhondt et

al. 2005, Vicente, Cascón et al. 2012, Marcos, Izquierdo et al. 2015). Induction of these oxylipins in this

study indicates their potential involvement in resistance to BW. Altogether, trehalose treatment had an

unexpectedly large impact on xylem levels of defense metabolites, particularly jasmonates and SA,

which have often been considered antagonistic signals.

Discussion

To survive drought, plants must respond to and overcome multiple adverse physiological and

molecular events (Shinozaki, Yamaguchi-Shinozaki et al. 2003, Osakabe, Osakabe et al. 2014). Our

experiments indicated that *R. solanacearum* affects tomato water physiology; the transcriptomic

signatures of infected plants had striking similarities to the gene expression profiles associated with

water stress.

The roles of trehalose and T6P as nutritional signaling metabolites in plants have been reviewed

(Lunn 2007, Paul 2007, Iordachescu and Imai 2008, Paul, Primavesi et al. 2008, Lunn, Delorge et al.

2014). The current paradigm is that T6P, and by extension trehalose, are direct indicators of sucrose

levels that help plants maintain sucrose-to-starch ratios through SnRK1 signaling (Schluepmann, Pellny

et al. 2003, Kolbe, Tiessen et al. 2005, Lunn, Feil et al. 2006, Schluepmann, Berke et al. 2012). However,

little is known about the roles of plant trehalose-metabolizing enzymes beyond nutritional signaling.

Most organisms, including plants and *R. solanacearum*, synthesize trehalose via a two-step reaction.

Trehalose-6-phosphate synthase (TPS) makes trehalose-6-phosphate (T6P) from glucose and UDP

glucose. T6P phosphatase (TPP) then dephosphorylates T6P to make trehalose. In *Arabidopsis,* most of

these proteins have predicted functions in development or signaling (Vandesteene, Ramon et al. 2010,

Vandesteene, Lopez-Galvis et al. 2012). Infected susceptible tomatoes slightly upregulated TPS-7 (0.5 log

fold change), suggesting this gene could be responsible for trehalose synthesis during BW. Resistant

plants did not upregulate TPS-7. Instead, they downregulated two TPS and three TPP genes, implying

that tightly regulated trehalose concentration during disease might contribute to resistance and

defense. Experimentally manipulating expression of these TPS and TPP proteins could test this

hypothesis.

We focused on tomato LEA and aquaporin genes because these proteins accumulate in

response to dehydration. However, resistant tomatoes downregulated genes annotated as LEA

hydroxyproline-rich glycoprotein (HRGP) proteins (Fig. 1B) (Cao and Li 2015). LEA HRGPs are plant cell

wall components, and their downregulation in our data suggests that resistant plants may slow their

growth as part of a defense trade-off (Johnson, Cassin et al. 2017). Since tomato aquaporins are

encoded by diverse families and channel other solutes in addition to water, it is not easy to generalize

about the behavior of aquaporins during BW (Reuscher, Akiyama et al. 2013). However, in response to

R. solanacearum infection, both susceptible and resistant tomatoes downregulated far more aquaporins

than they upregulated, suggesting that plants may try to alter water flow during BW. SlTIP2;2

(Solyc03g120470), an aquaporin that improved transpiration and yield when overexpressed in tomato

plants (Sade, Vinocur et al. 2009), was downregulated in both resistant and susceptible plants (Fig. 1D).

The meta-transcriptome analysis revealed that one ABA hydroxylase gene was strongly

upregulated in resistant and susceptible plants, while another was downregulated only in resistant

plants (Fig. 1C). In *Arabidopsis* and tomato*,* CYP707A ABA hydroxylases, which catalyze the first step in

ABA degradation, are involved in dehydration and rehydration at various developmental stages (Kushiro,

Okamoto et al. 2004, Saito, Hirai et al. 2004, Umezawa, Okamoto et al. 2006, Ji, Kai et al. 2014). This

strongly suggests both that tomatoes tightly regulate ABA catabolism during BW and that BW causes

water stress. Expression of TAS14, a dehydrin inducible by environmental stress and ABA (Godoy, Pardo

et al. 1990), was highly upregulated in susceptible and resistant plants, further evidence that BW causes

abiotic stress. In *Arabidopsis,* ABA synthesis, signaling, and responsive genes were similarly affected

during late-stage BW infection (Hu, Barlet et al. 2008). Additional identification of ABA-responsive

networks in tomatoes could identify targets for breeding more hydraulically efficient tomatoes that are

also more resistant to BW. It would be interesting to see if the ABA transcription factors identified in our

transcriptomic analysis regulate expression of tomato TPP genes; if so, manipulating these genes could

contribute to BW resistance.

ABA concentration in xylem sap of susceptible tomatoes increased in response to both *R.*

solanacearum infection and trehalose treatment. Elevated ABA levels are usually associated with

drought stress; together with the observation that *R. solanacearum* infection lowers plant stomatal

conductance, this indicates that BW significantly disrupts tomato water physiology. However, there

might also be a defensive component to ABA concentration and signaling. Deleting several cellulose

synthase (*irx*) genes increased *Arabidopsis* resistance to BW in an ABA-dependent manner and ABA

synthesis mutants were more susceptible to *R. solanacearum* (Hernández-Blanco, Feng et al. 2007). This

suggests that tomatoes that over-produce ABA could have increased resistance to wilt diseases.

Treating tomato roots with ABA reduced both stomatal conductance and *R. solanacearum*

colonization in stems, suggesting that *R. solanacearum* colonization depends, at least in part, on active

hydraulic conductance and a negative pressure gradient in host xylem. *R. solanacearum* is known to use

swimming and type IV motility and cell wall degrading enzymes to move towards roots and migrate

through the xylem (Tans-Kersten, Huang et al. 2001, Liu, Zhang et al. 2005). However, our results show

that *R. solanacearum* also uses plant water transport caused by transpiration to spread within the plant.

This suggests that *R. solanacearum* decreases its own spread and pathogenic fitness when it blocks host

plant water flow. This idea has interesting implications for physiological differences between susceptible

and resistant tomato plants since wilt-resistant varieties thrive even though they are often latently

infected. It is possible that ancestors of *R. solanacearum* were once endophytes but lost the ability to

check their population growth in the vasculature of susceptible hosts, possibly as a result of evolutionary

selection pressures imposed by agriculture, as has been proposed for *Xylella fastidiosa* (Beattie and

Lindow 1995).

Trehalose likely improves drought tolerance in plants by two mechanisms: by regulating

carbohydrate levels during stress conditions, possibly via the SnRK1 kinase; and by altering ABA signaling

and stomatal conductance (Garg, Kim et al. 2002, Paul, Gonzalez-Uriarte et al. 2018, Yu, Zhao et al.

2019). Members of another kinase family, SnRK2, also integrate with ABA signaling, and trehalose

metabolism may also connect to this network via the TOR regulator (Signorelli, Tarkowski et al. 2019,

Chen, Li et al. 2020). In our system, exogenous trehalose temporarily improved water conservation in

uninfected tomatoes and delayed wilting of drought-stressed plants. This was most likely because

trehalose treatment temporarily reduced stomatal conductance and transpiration and increased water

use efficiency. Trehalose also increased ABA content in xylem sap 24 h post-treatment. These results

confirm and extend findings that trehalose, ABA signaling, and water use interact in complex ways. The

Arabidopsis TPS1 gene responsible for maintaining T6P levels responds to ABA and contributes to

regulating stomatal aperture (Avonce, Leyman et al. 2004, Gomez, Gilday et al. 2010). At least two

Arabidopsis TPP proteins (AtTPPI and AtTPPE) that are controlled by ABA-responsive transcription

factors contribute to trehalose production and mitigate drought stress (Lin, Yang et al. 2019, Lin, Wang

et al. 2020, Wang, Chen et al. 2020). Another study found applying trehalose to tomato roots improved

water conservation, decreased ABA biosynthesis gene expression, increased expression of ABA-

responsive genes, kept stomates closed five days after trehalose treatment without water, and

increased ABA levels in leaves after 24 h (Yu, Zhao et al. 2019). Our results contrast in some ways with

this paper. We found that ABA content temporarily increased in xylem sap 24 h after after trehalose

treatment while stomatal conductance was reduced for less than 24 h. These differences may be

explained by methodological variation: Yu and co-workers did not include non-drought stressed tomato

controls, they measured ABA in leaves rather than xylem, and they measured stomatal aperture rather

than conductance. There is a lag between stomatal conductance response in leaves and ABA

concentration in the xylem sap in our data, leading us to hypothesize that ABA concentration in leaves

might be a more biologically relevant marker for stomatal conductance. It would be interesting to

determine if exogenous trehalose or ABA treatment changes trehalose concentration in tomato xylem

sap.

The plant hormone SA, which mediates resistance to *R. solanacearum* in many systems, is

necessary for induction of plant innate immunity and activation of systemic acquired resistance

(Deslandes, Olivier et al. 2002, Hirsch, Deslandes et al. 2002, Chen, Lin et al. 2009, An and Mou 2011,

Boatwright and Pajerowska-Mukhtar 2013, Jacobs, Milling et al. 2013). SA-responsive tomato defense

genes were highly upregulated 6 and 48 h after trehalose treatment. These same genes are upregulated

in diseased tomato plants following *R. solanacearum* soil soak infection, suggesting trehalose treatment

activates SA-mediated resistance pathways that could affect *R. solanacearum* infection (Milling, Babujee

et al. 2011).

Trehalose treatment increased wilt resistance of SA-degrading *NahG* tomato plants, but not the

parental Moneymaker cultivar. There are several possible explanations for this result. First, trehalose

treatment could have increased SA levels so much that overproduction of catechol, the breakdown

product of SA, contributed to resistance (Retig and Chet 1974). Second, excess ET and JA in *NahG* plants

could promote resistance to *R. solanacearum*. After infection with *P. syringae* DC3000 *NahG* tomatoes

produce more ET and JA, both of which can contribute to disease resistance (Spoel, Koornneef et al.

2003, López-Gresa, Lisón et al. 2016). ET is involved in *R. solanacearum*-host interactions and ET-

responsive defense genes were upregulated at 6 and 48 h post trehalose treatment. However, ET-

insensitive *NeverRipe* tomatoes were neither more nor less resistant to *R. solanacearum* than wild type

after trehalose treatment. In another study, treatment with a biocontrol strain of the oomycete *Pythium*

enhanced resistance to *R. solanacearum* in *NahG* tomatoes, but not in a JA mutant (Hase, Takahashi et

al. 2008). Since the ET defense pathway appeared less relevant in our study, it may be useful to

determine if JA enhances *R. solanacearum* resistance in tomatoes.

Plants use hormones to fine-tune many of their growth, development, and defense functions,

and xylem vessels can act as signal transduction highways for long-distance transport of auxin, ABA, JA,

and SA (Jackson 1997, Park, Lee et al. 2017). We observed increased SA concentration in xylem sap of

trehalose treated plants but not in the xylem sap of symptomatic *R. solanacearum*-infected plants. This

was surprising because SA signaling and concentration are integral to plant response to BW (Jacobs,

Milling et al. 2013). However, *R. solanacearum* can actively degrade SA, which is toxic to the pathogen

(Lowe-Power, Jacobs et al. 2016). It is also possible that SA concentrations might be higher in root

xylem than in stem xylem. For instance, an *Arabidopsis* mutant impaired in secondary wall deposition

had higher SA concentrations in the root and increased resistance to *R. solanacearum* infection, possibly

through vascular immunity (Denancé, Ranocha et al. 2013). Taken together, the increased SA levels in

xylem sap, the increased expression of SA-dependent defense genes, and the trehalose-dependent

NahG resistance phenotype suggest that the protective effect of trehalose requires systemic resistance

induced via SA.

Unexpectedly, jasmonate levels were elevated in trehalose-treated plants but not in *R.*

solanacearum-infected plants. As discussed earlier, there is some evidence that JA can play a role in *R.*

solanacearum defense. Genes related to ET, JA, and auxin signaling were upregulated in resistant LS-89

tomato seedlings one day after stem inoculation with *R. solanacearum* (Ishihara, Mitsuhara et al. 2012).

The higher concentration of jasmonates in trehalose-treated xylem sap may have contributed to

resistance. This hypothesis could be tested using JA-deficient tomato mutants.

SA- and JA-mediated plant defense responses have often been seen as antagonistic (Glazebrook

2005, Derksen, Rampitsch et al. 2013). Our gene expression analysis displayed this antagonism; at six h

post trehalose treatment JA-responsive *pin2* was downregulated, while SA-responsive *PR1b* and *gluA*

were upregulated. Roots of wilt-resistant Hawaii7996 tomatoes displayed a similar gene expression

profile 24h after *R. solanacearum* infection (French, Kim et al. 2018). Despite the antagonism observed

in the gene expression data, the concentrations of SA, JA, and the JA precursor 12-OPDA all increased in

trehalose-treated plants at 24 h. Recently, it was demonstrated in maize that levels of OPDA and JA are

irrelevant for the classic SA/JA-Ile antagonism and that 12-OPDA and SA are co-regulated (Wang,

Borrego et al. 2020). Infusing maize xylem sap with 12-OPDA increased resistance to *Colletotrichum*

graminicola while infusing similar concentrations of JA increased susceptibility. Disease experiments

with JA-insensitive tomato mutants could determine if 12-OPDA or JA contribute to resistance and

define their relationship to SA signaling.

JA signaling also induces leaf senescence, and chlorosis is associated with increased JA

production (He, Fukushige et al. 2002). We found that trehalose treatment significantly decreased

chlorophyll content 2-10 days post-treatment, which correlates with the increased JA production in

trehalose-treated plants. This is consistent with the finding that applying 5 mM trehalose to soybean

roots did not affect JA concentration, but it did decrease chlorophyll content at ten days (Asaf, Khan et

al. 2017).

Trehalose treatment and *R. solanacearum* infection both increased xylem sap levels of several

antimicrobial compounds, including phenolics. *R. solanacearum* requires at least two multi-drug efflux

pumps and a hydroxycinnamic acid degradation pathway for full virulence and fitness, showing that

antimicrobials, including phenolics, protect tomato plants from BW (Brown, Swanson et al. 2007, Lowe,

Ailloud et al. 2015). Plants have specialized cells that store and release phenolics into the xylem as part

of their defense response, and trehalose treatment and *R. solanacearum* infection both induce this

release (Beckman 2000).

Levels of the oxylipins 9-HOT, 9-HOD, 13-HOD, and 13-KOD increased in sap from *R.*

solanacearum-infected plants and trehalose-treated plants 24 h after treatment. These metabolites,

which accumulate in wounded maize, bean, and *Arabidopsis*, can be potent inducers of defense gene

response (Alméras, Stolz et al. 2003, Walters, Cowley et al. 2006, He, Borrego et al. 2020). Further, these

oxylipins have direct antimicrobial activity against fungal and bacterial pathogens (Prost, Dhondt et al.

2005). The data generated by this study deserve further analysis since little is known about oxylipin

signatures during BW or oxylipin toxicity to *R. solanacearum*. However, 9-LOX-derived oxylipins,

including 9-HOT and 9-HOD, increase in tobacco leaves inoculated with *R. solanacearum* GMI100, which

is avirulent on tobacco (Montillet, Agnel et al. 2002). This increase in 9-LOX derivatives correlated with

the hypersensitive response (HR) in leaves. We speculate that the presence of related oxylipins in our

study may indicate vascular HR. It would be interesting to see if defense compound concentration

changed in the xylem sap of resistant tomatoes challenged with *R. solanacearum.*

Plant systemic resistance typically involves accumulation of SA, JA, and defense oxylipins, and

increased expression of SA, JA, and ET-responsive defense genes in response to mechanical wounding or

pathogen challenge (Vallad and Goodman 2004, Choudhary, Prakash et al. 2007, Savatin, Gramegna et

al. 2014, Wang, Borrego et al. 2020). We found that treating tomato roots with trehalose similarly

increased SA, JA, and defense oxylipins, and upregulated SA, JA, and ET defense genes. We propose that

tomato plants activated induced systemic resistance because they recognized trehalose as a damage

associated molecular pattern (DAMP). DAMPs are waste products produced by pathogen invasion or

wounding that alert the plant that it is under attack (Ranf, Eschen-Lippold et al. 2011, De Lorenzo,

Ferrari et al. 2018, Hou, Liu et al. 2019). A DAMP can be almost anything that is usually intracellular or

results from lignin and cellulose polymer degradation. Our finding adds to the growing evidence that

sugars contribute to plant innate immunity (Bolouri Moghaddam and Van den Ende 2012).

Could treating roots with compatible solutes like trehalose in the field provide temporary

resilience until disease and drought pressure have passed? Other chemical inducers of plant defense

responses can reduce BW. Acibenzolar-S-methyl (Actigard) and DL-3-aminobutyric acid (BABA) both

lowered wilt disease incidence in greenhouse and field by promoting defense gene expression and

defense compound production, respectively (Jakab, Cottier et al. 2001, Pradhanang, Ji et al. 2005,

Hassan and Abo-Elyousr 2013). Additionally, growing tomatoes hydroponically with SA and JA increased

defense enzyme production and reduced BW incidence (Mandal, Kar et al. 2013). Finally, exogenous L-

histidine induced resistance to BW in an ET-dependent manner in *Arabidopsis* and tomato (Seo, Nakaho

et al. 2016). Further experiments are needed to determine if treating tomatoes with exogenous

trehalose could be a practical BW management strategy.

R. solanacearum GMI1000 produces and secretes abundant amounts of the TreA trehalase

during tomato infection, and ability to degrade trehalose contributes to bacterial fitness and virulence

even though trehalose is not an important nutritional source for the pathogen (Jacobs, Babujee et al.

2012, Hamilton, Steidl et al. 2021). Interestingly, treating plants with the trehalase inhibitor Validamycin

A increased BW resistance (Ishikawa, Fujimori et al. 1996). We previously showed that the trehalose in

xylem sap of *R. solanacearum*-infected tomato plants is produced by the plant (MacIntyre, Barth et al.

2019). Could *R. solanacearum* secrete trehalase to counter the effects of plant-produced trehalose?

Our results suggest several ways *R. solanacearum* could benefit from breaking down plant-

produced trehalose. First, the pathogen may degrade xylem sap trehalose in order to reduce host plant

defenses by removing this potential DAMP. Second, as discussed above, trehalose reduces tomato

transpiration, which helps *R. solanacearum* spread and colonize host xylem. *R. solanacearum* may

degrade trehalose to maintain this helpful sap flow by preventing trehalose-mediated stomatal closing.

Third, *R. solanacearum* could degrade trehalose to interfere with the trehalose and T6P signaling that

plants use to fine tune their sucrose to starch ratios.   Trehalose degradation could thus indirectly

increase the amount of available sucrose, which is a key carbon source fueling *R. solanacearum* growth

in xylem (Jacobs, Babujee et al. 2012, Hamilton, Steidl et al. 2021). The fact that the *R. solanacearum*

effector RipTPS is a functional trehalose-6-phosphate synthase is consistent with the hypothesis that this

pathogen benefits from manipulating host plant trehalose metabolism (Poueymiro et al., 2014).

Additional experiments are needed to test these non-exclusive hypotheses about how trehalase benefits

R. solanacearum.

Conclusion

This study argues that trehalose is present in xylem sap of *R. solanacearum*-infected tomatoes

as a response to wilt disease-mediated water stress. Both trehalose levels and the expression of water

stress genes change in response to *R. solanacearum* infection. Further, ABA content, an indicator of

drought status, increases in xylem sap of diseased plants. We confirmed the link between trehalose and

ABA signaling and drought stress in plants by demonstrating that trehalose treated plants temporarily

reduce stomatal conductance, contain more ABA in their xylem sap, and are more drought tolerant.

When applied to plant roots, trehalose reduced BW incidence and reduced host xylem

conductance, which aids bacterial colonization. SA-responsive defense genes were upregulated in

trehalose treated plants, SA non-responsive *NahG* plants were more resistant to *R. solanacearum* after

trehalose treatment, and SA concentration transiently spiked in xylem sap from treated plants. JA, other

oxylipins, and antimicrobial compounds also all increased in xylem sap of trehalose-treated plants.

Taken together, these results suggest that plants perceive trehalose as a DAMP and respond with

systemic disease resistance. In turn, *R. solanacearum* degrades trehalose as a counter-defense.

Methods

Total RNA extraction and RNA-seq sample preparation | RNA extraction and RNA-seq sample

preparation for the transcriptomic studies were performed as described (Khokhani, Lowe-Power et al.

2017, French, Kim et al. 2018).

RNA-seq data analysis | Illumina HiSeq2500 paired-end 100-bp; Illumina HiSeq2500 paired-end 150 bp

and Illumina HiSeq2000 paired-end 100-bp RNA sequencing were performed on the 18, 6, and 12

samples from three independent studies (two previously unpublished [Studies A+B], and [Study C]

French, Kim et al. 2018). RNA-sequencing data were analyzed through the same analysis pipeline. Read

quality was controlled using FastQC version 0.11.9 (Andrews). Reads were trimmed to remove low-

quality sequences and the Illumina sequencing adaptors (TruSeq3-PE-2) using Trimmomatic version 0.39

(Bolger, Lohse et al. 2014). After trimming and mapping, a total of 930,004,276, 135,532,392, and

103,831,164 paired-end reads were generated for Studies A, B, and C, respectively. Reads for each

sample were mapped to the ITAG4.0 *S. lycopersicum* reference genome using STAR version 2.7.4.a

(Dobin, Davis et al. 2012). Library type was set to strand-specific (reverse) for Study A and Study B, and

set to no-stranded in Study C. Gene expression was measured as the total number of reads for each

sample uniquely mapped to the reference using HTSeq version 0.11.1 and samtools version 1.10 (Li,

Handsaker et al. 2009, Anders, Pyl et al. 2014). Each sample averaged about 12, 9, and 4 million uniquely

aligned reads for Studies A, B, and C, respectively. "EdgeR" package version 3.28.0 was used to filter for

low counts such that at least three of the samples in each study had at least three counts per million,

resulting in a total of 19123, 18193, and 18917 genes remaining for Studies A, B, and C, respectively

(Robinson, McCarthy et al. 2009).

Differential gene expression within each study was conducted using the DESeq2 package

Bioconductor version 3.11 and a Benjamini-Hochberg false discovery rate multiple testing correction of

P=.05 (Benjamini and Hochberg 1995, Love, Huber et al. 2014). Meta-transcriptomic analysis was

performed using the metaRNASeq package version 1.0, and differential gene expression within the

meta-analysis was determined with the DESeq2 results, Fisher, and inverse normal techniques with a

P=.05, as described (Rau, Marot et al. 2014). Differentially expressed genes were filtered using a log2

fold change >|0.585| (corresponds to a fold change of |1.5|).

Gene Ontology analysis was performed using the ShinyGO analysis tool version 0.61, agriGO

version 2.0, and PANTHER GO analysis tool version 14 (Tian, Liu et al. 2017, Mi, Muruganujan et al. 2018,

Ge, Jung et al. 2019). *Arabidopsis thaliana* homologs of S. *lycopersicum* genes coding for transcription

factors were identified using Phytozome platform version 12.1 (Goodstein, Shu et al. 2011). Heatmaps

were visualized with R software version 3.6.1 package "pheatmap" version 1.0.12 (R , Kolde 2019).

Boxplots were visualized using package "ggplot2" (Wickham 2016).

Candidate TPS and TPP proteins were identified in the most recent annotations of the tomato

(cv. 'Heinz 1706') genome available on NCBI and the Sol Genomics Network (ITAG4.0,

https://solgenomics.net/tools/blast/, accessed 03/2020) using BLASTP with standard settings (Sato,

Tabata et al. 2012). BLASTP protein sequences were derived from the *E. coli* K-12 *otsA* and *otsB* proteins,

as well as *Arabidopsis* trehalose-related proteins (Leyman, Van Dijck et al. 2001, Vandesteene, Ramon et

al. 2010, Vandesteene, Lopez-Galvis et al. 2012). Tomato cv. Heinz 1706 contained ten TPS genes, eight

TPP genes, and one trehalase gene (Supplemental Table 1). Tomato ABA-related, LEA, and aquaporin

were identified by their PANTHER annotation, and many were previously annotated (Ma, Szostkiewicz et

al. 2009, Reuscher, Akiyama et al. 2013).

The transcriptomic studies were performed and labeled as follows: R1, roots of resistant tomato

breeding line Hawaii7966 (HW) sampled 24 h after infection with *R. solanacearum* strain K60; R2, roots

of HW sampled 48 h after infection with *R. solanacearum* strain K60; R3, seedling roots of HW sampled

24 h after infection with *R. solanacearum* strain GMI1000; S1, roots of wilt-susceptible West Virginia 700

tomato plants (WV) sampled 24 h after infection with *R. solanacearum* strain K60; S2, roots of WV

sampled 48 h after infection with *R. solanacearum* strain K60; S3, seedling roots of wilt-susceptible cv.

Bonny Best (BB) sampled 24 h after infection with *R. solanacearum* strain GMI1000; S4, mid-stems of BB

sampled 72 h after inoculation with *R. solanacearum* strain GMI1000.

Hormone analysis leaf and sap collection | For the hormone analysis, xylem sap was harvested and

collected as described (Lowe-Power, Hendrich et al. 2018). Unwounded three-week-old wilt-susceptible

Bonny Best plants were inoculated with wild-type *R. solanacearum* or water via naturalistic soil soak in

which 50 mL of $1x10^8$ CFU/mL were poured into a pot containing one plant. Plants were harvested when

they reached DI=1 (1-25% of leaf area wilted). Trehalose-treated plants were treated with 50 mL of 30

mM trehalose solution or water, then harvested 6, 24, or 48 h later. Xylem sap was harvested by de-

topping plants 1 cm above the cotyledon and collecting the exuded sap over 30 minutes. The sap that

seeped within the first five minutes was discarded to reduce phloem contamination. Sap was kept on

ice. Bacteria were removed from sap via chilled centrifugation, and sap was subsequently flash frozen.

Stems were destructively sampled to quantify *R. solanacearum* colonization as CFU/g stem by serial

dilution plating. Xylem sap was normalized by sap volume (250-500 µL) and concentration of bacteria in

the stem. Each test group contained six technical replicates (individual plants).

For metabolite analysis of sap, 95 µL of xylem sap was mixed with 5 µL of 50 µM of internal

standard consisting of; d-ABA ($[^2H_6]$(+)-*cis*, *trans*-abscisic acid, Olchem, Olomouc, Czech Republic), d-CA

(d_7-CA, *trans*-cinnamic acid), d-IAA ($[^2H_5]$ indole-3-acetic acid, Olchem), d-JA (2,4,4-d_3; acetyl-2,2-d_2

jasmonic acid, CDN Isotopes, Pointe Claire, Quebec, Canada), and d-SA (d_6-SA; Sigma-Aldrich, St Louis,

MO, USA). Samples were transferred into autosampler vials for LC-MS/MS analysis. Authentic analytical

external standards for each metabolite were used to identify select ions and retention times for each

metabolite (Supplemental Table 4). The simultaneous detection of several phytohormones used the

following methods with modifications (Müller and Munné-Bosch 2011). We used an Ascentis Express C-

18 Column (3 cm × 2.1 mm, 2.7 μm) (Sigma-Aldrich, St. Louis, MO, USA) connected to an API 3200 LC-

MS/MS (Sciex, Framingham, MA, USA) using electrospray ionization in negative mode with multiple

reaction mentoring. The injection volume was 10 μL and had a 500 μL min^{-1} mobile phase consisting of

Solution A (0.2% acetic acid in water) and Solution B (0.2% acetic acid in acetonitrile) with a gradient

consisting of (time – %B): 0.5 – 10%, 1.0 – 20%, 21.0 – 70%, 24.6 – 100%, 24.8 – 10%, 29 – stop. After

integration with Analyst v1.6.3 (Sciex), the concentration of endogenous metabolites was determined by

comparing their peak areas to those of isotopically labeled internal standards using calculated response

factors.

Stomatal conductance, transpiration, and photosynthesis measurements with LI-COR |To assess the

effect of trehalose treatment on tomato stomatal conductance (g_s), transpiration (E), and

photosynthesis (A_{net}), we used a LI-COR portable photosynthesis system (model LI-6400XT, LI-COR

Biosciences, Lincoln, NE, USA) equipped with a CO_2-controller and a red-blue LED chamber (model LI-

6400-02b). The ambient CO_2 concentration was set to 400 ppm, flow rate to 300 μmol s^{-1}, and the

chamber light source set at an intensity of 1200 μmol m^{-2} s^{-1} for all measurements. Fifteen-day-old

Bonny Best plants were treated with 50 mL of trehalose solution or water, then watered every day after

for the duration of the experiment. LI-COR measurements were taken at 10min, 6 h, 24 h, 48 h, and 72 h

post trehalose treatment. The experiment was repeated twice with ten plants per treatment.

R. solanacearum **culture conditions |** *R. solanacearum* strains were cultured from water stocks or -80 °C

glycerol stocks on TZC plates containing casamino acids, peptone, glucose, and tetrazolium chloride, and

incubated at 28 °C for 48 h (Hendrick and Sequeira 1984). Δ*treA* cultures were supplemented with 20 mg

L^{-1} spectinomycin. The kanamycin-resistant variant of wild-type GMI1000 was supplemented with 25 mg

L^{-1} kanamycin on plates and in liquid culture. Overnight cultures of *R. solanacearum* were grown at 28 °C

in CPG in a shaking incubator.

ABA treatments | We assessed the effect of ABA treatment on two-week-old tomato stomatal

conductance and subsequent bacterial wilt incidence. ABA was prepared at a stock concentration of

50mg/mL in DMSO. The water + DMSO control plants were watered daily with 50 mL of water

supplemented with the volume of DMSO used to make the 100 µM ABA solution. The ABA-treated

groups were watered daily with 50 mL of 50 or 100 µM ABA solution in water. After 48 h of ABA

treatment, plants were soil drenched with 50 mL of a $1x10^8$ CFU/mL of *R. solanacearum* GMI1000 with

kanamycin resistance (to aid in recovery from the plant) (Lowe, Ailloud et al. 2015). Bacteria were

resuspended at the desired O.D. in water + DMSO or the ABA solutions, so plants were inoculated and

ABA treated concurrently. Tomatoes were destructively sampled five days post-infection (after 168 h of

ABA treatment) to assess bacterial colonization of stems. Stomatal conductance, transpiration, and

photosynthesis measurements were taken for the entirety of the experiment with the LI-COR as

described 4-6 h after watering/ABA treating in the morning, which was the length of time it took ABA

treatment to affect conductance.

We assessed ABA toxicity to *R. solanacearum* by supplementing CPG broth with 50-100 µM ABA

or the equivalent volume of DMSO, and growing *R. solanacearum* cells in this media for 48 h. Cells were

grown in a shaking 96 well plate at 28°C in a BioTek Synergy HT plate reader, which measured Abs_{600}

every half hour. The experiment was performed once with eight technical replicates per treatment.

Plant assays with *R. solanacearum* | Plant growth conditions, colonization, and disease curve assays

were performed as described (Khokhani, Tuan et al. 2018, MacIntyre, Barth et al. 2019). All plant assays

were grown on 12 h light/dark cycles at 28 °C with ~60% relative humidity unless stated otherwise. For

the disease curve assays with trehalose treatments, fifteen-day old Bonny Best tomatoes were treated

with either water (control) or 50 mL of 30 mM trehalose solution. Plants were not watered 24 h post-

treatment per standard procedure for soil soak assays. At 48 h post-treatment plants were soil drenched

with 50 mL of $1x10^8$ CFU/mL of wild type *R. solanacearum* GMI1000 (or Δ*treA* for the later trehalose

treatment experiments with this mutant). For the petiole inoculations, 3-week-old tomatoes were

treated with trehalose or water as described above; after 48 h a true leaf was removed, and 2000 wild

type GMI1000 cells in 10 µL of water were pipetted onto the cut stem. To assess the effectiveness of

spraying trehalose to delay BW incidence, fifteen-day-old Bonny Best plants were either treated with 30

mM trehalose solution through the soil as described or sprayed with trehalose solution (or water) to

run-off. For all assays, plants were inoculated with *R. solanacearum* 48 h post trehalose root or leaf

treatment.

After treatments and inoculations, all plants in disease assays were watered on alternating days

post-inoculation with half-strength Hoagland's solution. Disease assay plants were rated daily on a 0-4

disease index scale where 0= no symptoms, 1=1-25% of leaf area wilted, and 4=76-100% of leaf area

wilted or dead. All disease curves represent four biological replicates, each containing 15

plants/treatment.

To assess *R. solanacearum* colonization of trehalose treated plants, fifteen-day-old Bonny Best

plants were treated with 50 mL 30 mM trehalose solution or water. 48 h post-treatment, plants were

infected with 50 mL of $1x10^8$ CFU/mL wild type *R. solanacearum* GMI100. Plant stems were destructively

sampled 1 cm above the cotyledon at 5, 6, and 7 dpi, ground, and serially dilution plated.

NahG salicylic insensitive tomato seeds ('Moneymaker' background) and *NeverRipe* ET-

insensitive seeds ('Pearson' background) were propagated in our lab (Lanahan, Yen et al. 1994, Brading,

Hammond-Kosack et al. 2000). Disease assays with trehalose-treated hormone insensitive plants were

as described above.

Water conservation | Two-week-old Bonny Best plants were treated with trehalose or water. Water

was subsequently withheld and plants were assessed daily to see how long it took them to wilt. Wilting

was rated using the same scale as disease curve assays. The experiment included four biological

replicates, each with ten plants per treatment.

QRT-PCR measurement of tomato trehalose and defense gene expression | JA, SA, and ET defense

gene primers were previously characterized in tomato *R. solanacearum* response (Supplemental Table 2)

(Milling, Babujee et al. 2011). The *rd22* and *dhn_tas* ABA-responsive primers were previously described

(Yamaguchi-Shinozaki and Shinozaki 1993, Kissoudis, Seifi et al. 2016). Actin and *dnaJ* primers served as

our constitutively expressed normalizing genes.

To measure defense gene expression, two-week-old Bonny Best tomatoes were treated with 50

mL of 30 mM trehalose or water. 100 mg of stem tissue was harvested 6 or 48 h post-treatment, then

flash frozen. For all samples, stems were homogenized for a minute in a Powerlyzer (Qiagen) at room

temperature in 450 µL of Buffer RLT from the RNAeasy Plant Mini Kit (Qiagen). RNA extraction

proceeded as specified in the kit protocol with off-column DNase I treatment using a DNA-free kit

(Thermo-Fisher). RNA quality and quantity were assessed with a NanoDrop (Thermo-Fisher) and reverse

transcribed into cDNA using Superscript III First-Strand Synthesis SuperMix (Invitrogen). Lack of genomic

DNA contamination was confirmed with PCR of no reverse transcriptase controls. Quantitative real-time

PCR was performed in 10 µL reactions with EvaGreen Mastermix (Biotium) using kit-specified reaction

parameters in a Quant Studio 5 Real-Time PCR machine (ThermoFisher Scientific). Averaged actin and

dnaJ values were used as the stably expressed normalization genes. Fold change values were calculated

using the $\Delta\Delta C_T$ method as described (Milling, Babujee et al. 2011). The data are presented as fold change

in gene expression compared to mock-inoculated controls and represent 16 plants per treatment over

four biological replicates.

Relative water content | To assess the water status and osmotic adjustment of tomato plants treated

with trehalose, we measured relative water content (RWC) in leaves. This directly measures cellular

hydration. Three-week-old Bonny Best tomato plants were treated with 50 mL of water or 30 mM

trehalose solution. 48 h post-treatment, leaves were excised at the petiole and weighed for fresh

weight. Leaves were then rehydrated by floating on distilled water in closed petri dishes for 4 h, at

which time they were blotted, and their turgid weight was assessed. Leaves were then thoroughly dried

in a food dehydrator oven for 24 h, and dry weight was evaluated. RWC was calculated as follows: %

RWC= [(Leaf fresh weight – dry leaf weight)/(turgid leaf weight-leaf dry weight)] x 100 (Barrs and

Weatherley 1962). The data represent two biological replicates with five technical reps (leaf from an

individual plant) per treatment.

Root length assays | Root length assays to quantify trehalose treatment effect on root length were

performed as described with modification (Tran, MacIntyre et al. 2016). Tomato seeds were sterilized in

10% bleach for 10 min, then 95% ethanol for 10 min, then washed five times with sterile water and

germinated in the dark on 1% water agar plates at room temperature. Four-day-old tomato seedlings

were transferred to germination pouches (Mega International, Newport, MN) then watered with either

10 mL of water or 30 mM trehalose solution and incubated in a 28°C growth chamber. Following the

initial trehalose treatment, pouches were kept moist as needed with sterile water and imaged ten days

post-treatment. This experiment was performed three times with ten tomato

seedlings/pouch/treatment. New root growth was calculated as % new root growth= (new growth/total

root length) x 100.

Chlorophyll determination | To measure the amount of chlorophyll in tomatoes, we used a SPAD-502

meter (Konica Minolta, Inc.) (Ling, Huang et al. 2011). Readings were performed on the first true leaves

of two-week-old Bonny Best plants treated with 50 mL of a 30 mM trehalose solution or water. The data

represent two biological replicates with ten plants per treatment per replicate.

Bacterial leaf spot of tomato inoculations | *Xanthomonas gardneri* culture conditions, infection, and

colonization analysis were as described (Potnis, Soto-Arias et al. 2014). *X. gardneri* strains were cultured

on nutrient agar plates (Millipore-Sigma) supplemented with nalidixic acid (50 μg/mL) at 28°C. Inoculum

was grown for 48 h in a shaking incubator at 28°C in nutrient broth supplemented with nalidixic acid.

The inoculum was adjusted to OD_{600}=0.3 and diluted 1:100 in 400 mL sterile water with 75 μL Silwet

(Fisher Scientific). Four-week-old Bonny Best tomato plants that were treated with water or 50 mL of 30

mM trehalose solution 72 h previously were dip-inoculated for 30 s in the inoculum. Plants were

incubated in high humidity conditions (85-90%) in plastic containers in the growth chamber for 48 h.

After 48 h, plants were moved to low humidity conditions during the day (taken out of containers, 45-

60%) and high humidity conditions at night on a 12 h cycle. Bacterial spot symptoms appeared 5-6 days

post-inoculation. Leaf tissue from individual plants was sampled at 3, 6, and 9 days post-inoculation.

Two 2 cm^2 portions of leaf were sampled from each plant, ground in a Powerlyzer (Qiagen) in water, and

serially diluted and enumerated on nutrient agar + nalidixic acid. The data represent three biological

replicates with 12 technical reps/treatment/timepoint.

DAB staining tomato leaves for ROS production | Tomato leaves were stained with 3,3'-

Diaminobenzidine (DAB, Sigma-Aldrich) as described (Flores-Cruz and Allen 2009). Leaves from 4-5-

week-old Bonny Best tomato plants were infused with water, 30 mM of trehalose solution, or $1x10^9$

CFU/mL *R. solanacearum* cells by pricking with a needle then flushing the apoplast with the solution.

Plants were returned to the growth chamber overnight. At 24 h, leaves were cut and placed in 0.1

mg/mL DAB solution prepared at pH 3.8 at room temperature (Thordal-Christensen, Zhang et al. 1997).

Containers holding the stain and leaves were wrapped in aluminum foil to exclude light and incubated

for 8 h at room temperature with gentle shaking. At 8 h, leaves were bleached by boiling in 95% ethanol

for 10 min, then further cleared with overnight incubation in 70% ethanol before imaging. These

experiments were performed three times with leaves from four individual tomato plants per treatment.

Statistical Analyses | Statistical analyses and graphs were generated with GraphPad Prism 8.

Acknowledgments and contributions: Thanks to Raka Mitra (Carleton College) for manuscript review

and valuable discussions. Thank you to Eric Kruger for the equipment loan and Kim Cowles for *X.*

gardneri cultures and technical advice. April MacIntyre and Caitilyn Allen designed the experiments and

wrote the text with contributions from Valerian Meline, Anjali Iyer-Pascuzzi, and Zachary Gorman. April

MacIntyre carried out most of the experiments and created most of the figures. Valerian Meline

performed the meta-transcriptomic analysis and created Figure 1. Zachary Gorman performed the xylem

sap metabolite analysis. Steven Augustine advised and assisted with all LI-COR experiments and created

the trehalose stomatal conductance, photosynthesis, transpiration, and water efficiency graphs. Carolyn

Dye assisted with the trehalose physiology, ABA treatment, and *Xanthomonas* experiments. Corri

Hamilton advised on xylem sap collection. Anjali Iyer-Pascuzzi, Michael Kolomiets, and Kate McCulloh

provided feedback and edited the manuscript.

Figures

Figure 1. *R. solanacearum* **infection changes expression of tomato genes involved in trehalose**
metabolism and water stress physiology. Comparative analysis of transcriptomic responses to *R.*
solanacearum from seven experiments measuring commonly differentially expressed genes across all
data sets in wilt-resistant and susceptible tomato plants. Heatmaps show relative expression (as $\log_2$
fold change) compared to mock-inoculated control plants of: **A)** Genes encoding trehalose metabolism;
B) Late embryogenesis/dehydrin (LEA) genes; **C)** Genes encoding ABA synthesis and signaling; and **D)**
Aquaporin genes. Gray boxes (NA) in heatmaps indicate that a gene's expression was not different from
that of mock-inoculated control plants. Transcriptomic datasets are: R1, roots of wilt-resistant tomato
breeding line Hawaii7966 (HW) sampled 24 h after infection with *R. solanacearum* strain K60; R2, roots
of HW sampled 48 h after infection with *R. solanacearum* strain K60; R3, seedling roots of HW sampled
24 h after infection with *R. solanacearum* strain GMI1000; S1, roots of wilt-susceptible West Virginia 700
tomato plants (WV) sampled 24 h after infection with *R. solanacearum* strain K60; S2, roots of WV
sampled 48 h after infection with *R. solanacearum* strain K60; S3, seedling roots of wilt-susceptible cv.
Bonny Best (BB) sampled 24 h after infection with *R. solanacearum* strain GMI1000; S4, mid-stems of BB
sampled 72 h after inoculation with *R. solanacearum* strain GMI1000. Tomato gene names and locus
tags are from NCBI and the Sol Genomics Network, accessed in 03/2020. Transcriptomic data were
normalized across experiments to allow comparisons as described in Methods.

Figure 2. ABA concentration increased in *R. solanacearum*-infected plants, and ABA treatment reduced stomatal conductance, which correlated with reduced *R. solanacearum* stem colonization. A) ABA concentration in xylem sap harvested from healthy and *R. solanacearum*-infected tomato plants showing early wilt symptoms (1-25% of leaf area wilted) (T-test, *P=.03). Bars show means of xylem sap ABA levels, measured by GC-MS, from four to six plants per treatment. **B)** Stomatal conductance in tomato leaves as measured with LI-COR instrument. Conductance of plants watered with 50 or 100 uM of ABA differed from that of mock-inoculated controls at all time points (2-way ANOVA Tukey's multiple comparisons, *P*<.05). Data represent four biological replicate experiments, with three plants analyzed per treatment per time point. **C)** *R. solanacearum* population sizes were quantified in tomato plant stems five days after soil-soak inoculation with *R. solanacearum* strain GMI1000. *R. solanacearum* colonization was lower in ABA-treated plants at 5 dpi, which corresponds to 168 h of ABA treatment (Kruskall Wallis, 50 uM ABA, ***P<.001; 100 uM ABA, **P<.01). The proportion of plants containing detectable *R. solanacearum* cells (shown below X-axis labels) was also lower (Chi-square, *P*=.0016). Dots represent *R. solanacearum* population sizes in individual plants. The horizontal bars represent the geometric mean of the data and vertical bars indicate standard deviation. Plants with undetectable *R. solanacearum* populations were given a value of 1 in the statistical analysis. Data represent three biological replicate experiments, each with 12 plants per treatment.

Figure 3. Trehalose treatment delayed tomato wilting due to drought, likely by transiently decreasing stomatal conductance and transpiration and increasing ABA content in xylem sap. A) Curves showing how long it took plants to wilt following a single treatment with either water (blue) or 30 mM trehalose (green) and then no water for two weeks (P=.01, T-test of areas under the curve). Data shown represent four biological replicates with ten plants/treatment/replicate. **B)** Stomatal conductance and **C)** transpiration were measured with LI-COR (P>0.05, repeated measures ANOVA). Data shown represent 20 plants/treatment. Measurements were taken 10 min after treatment with trehalose or water, then each day after that in the morning two hours after growth chamber lights turned on. The shaded bars on graphs represent night periods. All data were calculated from the same LICOR time points. **D)** ABA concentration in trehalose treated plants (One-way ANOVA Fisher's LSD multiple comparisons to water-treated control: 6 h tre P=.85, 24 h tre P=.016, 48 h tre P=.95). Data shown represent measurements of xylem sap samples from four to six plants.

Figure 4. Treating plants with trehalose at the root delayed BW symptoms and reduced *R.*
***solanacearum* stem colonization, but only following soil inoculation. A)** Wilt disease progress of
tomato plants treated with trehalose, then after 48 h soil soak inoculated with *R. solanacearum*
GMI1000. Trehalose treatment delayed appearance of wilt symptoms (*P*=.0082, T-test comparing areas
under the curves). Data represent three replicate experiments, each with 15 plants/treatment. Disease
symptoms were rated on a 0-4 disease index scale where 0=no wilt, 1=1-25% of the leaves wilted,
4=plant is 76-100% wilted or dead. **B)** Bacterial population sizes were smaller in stems of trehalose-
treated plants at 5-7 days after inoculation (Mann-Whitney, *P*=.0009). Data represent three biological
replicates with 15 plants/day sampled/treatment; each dot represents the *R. solanacearum* population
size in one plant (including plants without detectable *R. solanacearum*). Black bars represent the
geometric mean and standard deviation of the data. The percentage of plants that contained any
detectable *R. solanacearum* cells is shown below the X-axis; the limit of detection is ~10^2 cells. Relative
to controls, fewer trehalose-treated plants contained detectable *R. solanacearum* populations (chi-
square, *P*=.0016). **C)** Disease progress of plants treated with trehalose (green) or water (blue), then 48 h
later inoculated by introducing *R. solanacearum* directly into the xylem through a cut leaf petiole.
Trehalose treatment did not affect wilt disease development in petiole-inoculated plants (*P*=.2067, T-
test of areas under curves). Data represent three replicate experiments, each containing 15
plants/treatment.

Figure 5. Exogenous trehalose treatment affects the expression of hormone-responsive tomato defense genes, and trehalose-mediated disease resistance is partially regulated by salicylic acid signaling. Effects of trehalose treatment on expression of diverse defense genes in Bonny Best tomato stems, determined by qRT-PCR at **A)** 6 h and **B)** 48 h after trehalose treatment at the roots (one-sample t- test to a hypothetical mean of 1.0 where 1.0 = no difference in expression; asterisks: *, P=.01-.05; **, P=.001-.01; ***, P=.0001-.001; ****, P<.0001). Expression levels are shown as fold-change relative to those in water-treated plants. Data represent three biological replicates per time point, each containing five plants per treatment. **C)** Wilt disease progress following *R. solanacearum* inoculation of water- or trehalose-treated *NeverRipe* ET-insensitive tomatoes and their 'Pearson' parent cultivar (One-way ANOVA of AUC; 'Pearson' H_2O vs. *NeverRipe* H_2O, P=.91; 'Pearson' tre vs. *NeverRipe* tre, P=.98). **D)** Disease progress following *R. solanacearum* inoculation of water- or trehalose-treated SA-insensitive *NahG* over-expressing plants and their parent cv. Moneymaker (One-way ANOVA of AUC, Moneymaker H_2O control vs. *NahG* H_2O control, P=.74; Moneymaker tre vs. *NahG* tre, P=.04). For **C)** and **D)**, plants were treated at the roots with trehalose, then inoculated with *R. solanacearum* via soil drenching 48 h later. Data represent three biological replicates, each containing 15 plants/treatment (45 plants total).

Figure 6. Trehalose treatment increased metabolites related to defense and wounding in tomato xylem sap. LC-MS/MS was used to measure concentrations of hormones, oxylipins, and other metabolites in xylem sap collected from cv. Bonny Best tomato plants that were either mock-inoculated (H_2O), inoculated with *R. solanacearum* GMI1000 (*R. solanacearum* DI=1), or had been treated at the roots with 30mM trehalose (Tre treat (6h), etc.) Differences from water-treated control plants were determined by one-way ANOVA Fisher's LSD multiple comparison test, *, P=.01-.05; **, P=.001-.01; ***, P=.0001-.001; ****, P<.0001). Data shown represent the means of samples from 4-6 individual plants. Bars indicate the mean with standard deviation. **A)** SA=salicylic acid; **B)** BA=benzoic acid; **C)** COUMA=coumaric acid; **D)** TA= Traumatic acid; **E)** 12-OPDA= 12-oxo-8(*Z*),13(*Z*)-phytodienoic acid; **F)** JA=jasmonic acid; **G)** JA-Ile=jasmonic acid isoleucine; **H)** 9-HOT= 9(*S*)-hydroxy-10(*E*),12(*Z*),15(*Z*)-octadecatrienoic acid; **I)** 9-HOD= 9(*S*)-hydroxy-10(*E*),12(*Z*)-octadecadienoic acid; **J)** 13-HOD= 9(*S*)-hydroxy-10(*E*),12(*Z*)-octadecadienoic acid; and **K)** 13-KOD= 13-oxo-9(*Z*),11(*E*)-octadecadienoic acid.

883   **Supplemental Tables and Figures**

884   MacIntyre et al *Trehalose induces tomato defenses and increases drought tolerance and bacterial wilt*
885   *disease resistance*

887   **Supplemental Table 1. Predicted TPS/TPP and trehalase genes in tomato cv. 'Heinz1706'**

Protein Name[a]	Locus tag (Solgenomics)[b]	Protein Accession (NCBI)[c]	Renamed[d]	Protein Length	Notes
trehalose-phosphate synthase [UDP-forming] 1-like isoform X2	Solyc02g071590.3.1	XP_010316884.1	SlTPSX2	943	
trehalose-phosphate synthase [UDP-forming] 6	Solyc01g005210.4.1	XP_004228746.1	SlTPS6	857	
trehalose-phosphate synthase [UDP-forming] 9	Solyc04g025940.3.1	XP_004237260.1	SlTPS9a	857	
trehalose-phosphate synthase [UDP-forming] 9	Solyc05g005750.3.1	XP_004238680.1	SlTPS9b	851	
trehalose-phosphate synthase 5	Solyc08g076650.3.1	XP_004245918.1	SlTPS5	862	
trehalose-phosphate synthase 7	Solyc07g055300.3.1	XP_004243268.1	SlTPS7a	858	
trehalose-phosphate synthase 7	Solyc02g072150.3.1	XP_004233035.1	SlTPS7b	851	
trehalose-phosphate synthase 7	Solyc10g007950.4.1	XP_004248198.1	SlTPS7c	857	
trehalose-phosphate synthase isoform X1	Solyc07g006500.3.1	XP_010323144.1	SlTPS11	852	Homologous to *Arabdopsis SlTPS1* (Singh and Shah 2012)
trehalose-phosphate synthase 1	Solyc07g062140.3.1	NP_001234879.1	SlTPS1	926	
trehalose-phosphate phosphatase H isoform X1	Solyc05g051880.4.1	XP_019069376.1	SlTPP_HX1	348	May have two additional splice variants

Protein Name[a]	Locus tag (Solgenomics)[b]	Protein Accession (NCBI)[c]	Renamed[d]	Protein Length	Notes
trehalose-phosphate phosphatase 2	Solyc03g007290.4.1	XP_004234173.1	SlTPP_2	348	
trehalose-phosphate phosphatase J	Solyc04g072920.4.1	XP_004237894.1	SlTPP_Ja	365	
trehalose-phosphate phosphatase J	Solyc04g054930.3.1	XP_004237406.1	SlTPP_Jb	388	
trehalose-phosphate phosphatase F	Solyc08g079060.4.1	XP_004245739.1	SlTPP_Fa	386	
trehalose-phosphate phosphatase F	Solyc06g060600.3.1	XP_004242008.1	SlTPP_Fb	339	
trehalose-phosphate phosphatase D	Solyc04g082550.2.1	XP_025886308.1	SlTPP_D	296	
trehalose-phosphate phosphatase A	Solyc03g083960.3.1	XP_025885725.1	SlTPP_A	384	
trehalase	Solyc08g082860.4.1	XP_025888255	SltreA	532	

[a] Tomato TPS/TPP and trehalase proteins were identified via a BLASTP search through NCBI (https://blast.ncbi.nlm.nih.gov/Blast.cgi?PAGE=Proteins) and the SolGenomics network (https://solgenomics.net/tools/blast/)(accessed: 3/11/2020) as described in the Methods section using the most up to date tomato reference genome (SL4.0 and ITAG4.0)
[b] Locus tags assigned by the SolGenomics network (https://solgenomics.net/)
[c] Protein accession tags through NCBI (https://www.ncbi.nlm.nih.gov/protein/)
[d] Proteins were renamed/abbreviated by the authors for clarity

893 **Supplemental Table 2. Strains and primers used in this study**

Strain	Notes	Reference
Wild type GMI1000	*Ralstonia solanacearum* phylotype one sequevar 18, isolated from tomato in French Guyana	(Boucher, Barberis et al. 1985)
GMI1000 Δ*treA*	GMI1000 trehalase mutant cannot degrade trehalose; Spectinomycin/streptomycin-resistant	(MacIntyre, Barth et al. 2019)
Primer Name	**Sequence**	**Reference**
loxA F	TGGTAGACCACCAACACGAA	(Milling, Babujee et al. 2011)
loxA R	GACCAAAACGCTCGTCTCTC	"
osmF	TGTACCACGTTTGGAGGACA	"
osmR	ACCAGGGCAAGTAAATGTGC	"
PR1b B F	TTGGTGACTGCGGGATGA	"
PR1b B R	GGCGGCGGCTAGGTT T	"
PR1a B F	GAGGGCAGCCGTGCAA	"
PR1a B R	CACATTTTTCCACCAACACATTG	"
GluA F	TCA GCA GGG TTG CAA AAT CA	"
GluA R	CTCTAGGTGGGTAGGTGTTGGTTAA	"
Pin2 F	TGATGCCAAGGCTTGTACTAGAGA	"
Pin2 R	AGCGGACTTCCTTCTGAACGT	"
ACO5 F	AGATGGGCATTGGGTGAACA	"
ACO5 R	TTCAGCCATCACTCGGTGTC	"
DnaJ F	ATGAAGCGCCAGATACCATC	"
DnaJ R	TCAAGGCTCAATGTGTGCTC	"
Actin F	TCAGCAACTGGGATGATATG	"
Actin R	TTAGGGTTGAGAGGTGCTTC	"
rd22_F	ACGTGGCGTTATTTTTCCTG	(Yamaguchi-Shinozaki and Shinozaki 1993)
rd22_R	ATCTCCGGCATCTTCTCTGA	"
dhn_tas_F	CACCATGAGGGGCAACAGCA	(Kissoudis, Seifi et al. 2016)
dhn_tas_R	TCACCTTCATGTTGTCCAGGCATC	"

895 **Supplemental Table 3. Concentration of hormones, oxylipins, and phenylpropanoids in tomato xylem**
896 **sap following various treatments**

Compound (nM)[a]	Healthy Avg[b]	WT avg[c]	WT FC[d]	WT p-val[e]	6 hr tre avg[f]	6 hr tre FC	6 hr tre p-val	24 hr tre avg[g]	24 hr tre FC	24 hr tre p-val	48 hr tre avg[h]	48 hr tre FC	48 hr tre p-val
12,13-diHOM	0.83	1.30	2	0.04	1.26	2	0.27	2.69	3	0.00	1.50	2	0.02
12,13-Ep-9KOM	5.38	6.86	1	0.24	7.93	1	0.02	4.82	1	0.64	7.62	1	0.04
12OH-JA	1.43	1.70	1	0.63	5.21	4	0.30	1.38	1	0.84	1.17	1	0.06
12OH-JA-Ile	12.37	15.23	1	0.81	20.67	2	0.29	22.97	2	0.14	57.67	5	0.00
12-OPDA	35.86	47.89	1	0.44	50.75	1	0.43	138.72	4	0.00	91.84	3	0.01
13-HOD	4.66	8.52	2	0.00	5.09	1	0.66	9.02	2	0.01	5.49	1	0.14
13-KOD	4.50	10.78	2	0.01	6.27	1	0.31	9.72	2	0.00	5.81	1	0.12
13,12-KODA	0.27	0.75	3	0.34	0.11	0	0.37	4.89	18	0.00	0.22	1	0.80
13,12-KOMA	1.78	3.18	2	0.23	1.23	1	0.47	9.67	5	0.00	2.23	1	0.58
2-HOD	0.24	0.35	1	0.44	0.36	1	0.18	0.30	1	0.55	0.29	1	0.66
9,10,11-THOD	88.94	191.22	2	0.02	134.71	2	0.09	214.49	2	0.00	153.30	2	0.02
9,10,11-THOM	237.57	811.16	3	0.00	430.01	2	0.02	572.57	2	0.00	379.05	2	0.00
9,10,13-THOD	3.47	4.47	1	0.28	9.91	3	0.00	5.78	2	0.00	6.06	2	0.02
9,10,13-THOM	62.08	100.56	2	0.03	147.39	2	0.00	128.47	2	0.00	96.87	2	0.01
9,10-diHOM	0.95	1.25	1	0.43	1.48	2	0.52	3.64	4	0.00	1.02	1	0.86
9,12,13-THOD	39.77	50.62	1	0.18	43.68	1	0.56	84.76	2	0.00	60.09	2	0.02
9,12,13-THOM	127.69	231.37	2	0.01	153.78	1	0.36	305.89	2	0.00	179.49	1	0.04
9-HOD	6.15	21.29	3	0.01	10.27	2	0.02	14.60	2	0.00	8.64	1	0.17
9-HOT	3.04	9.78	3	0.03	5.02	2	0.16	9.75	3	0.00	4.87	2	0.12
9-KOD	4.35	8.90	2	0.15	6.60	2	0.34	19.32	4	0.00	12.16	3	0.00
9-KOT	1.06	3.68	3	0.17	5.51	5	0.05	7.61	7	0.00	3.90	4	0.05
9OH-10KOM	8.81	5.36	1	0.07	10.81	1	0.15	9.05	1	0.88	8.97	1	0.92
9OH-12KOD	0.20	0.13	1	0.51	0.27	1	0.53	0.69	4	0.00	0.34	2	0.02
9OH-12KOM	1.88	2.48	1	0.03	2.96	2	0.24	3.97	2	0.00	2.11	1	0.36
9-oxo-NA	13.94	17.47	1	0.37	16.53	1	0.56	16.06	1	0.51	17.25	1	0.32
ABA	8.20	49.59	6	0.03	5.71	1	0.60	40.37	5	0.00	9.72	1	0.73

Compound (nM)[a]	Healthy Avg[b]	WT avg[c]	WT FC[d]	WT p-val[e]	6 hr tre avg[f]	6 hr tre FC	6 hr tre p-val	24 hr tre avg[g]	24 hr tre FC	24 hr tre p-val	48 hr tre avg[h]	48 hr tre FC	48 hr tre p-val
AZA	14.08	19.39	1	0.11	19.21	1	0.00	20.44	1	0.00	18.03	1	0.01
BA	7.18	12.37	2	0.13	11.34	2	0.19	13.29	2	0.00	9.74	1	0.02
CA	7.49	5.19	1	0.52	5.30	1	0.56	22.69	3	0.00	14.19	2	0.16
COUMA	2.04	149.10	73	0.05	2.42	1	0.83	5.22	3	0.11	0.00	0	0.12
dnOPDA	0.12	0.21	2	0.25	0.26	2	0.02	0.53	4	0.00	0.44	4	0.00
IAA	3.26	36.76	11	0.03	2.84	1	0.59	4.01	1	0.29	2.20	1	0.09
JA	16.71	9.42	1	0.22	26.64	2	0.33	49.41	3	0.01	71.36	4	0.00
JA-Ile	26.59	3.67	0	0.09	35.48	1	0.62	60.08	2	0.04	103.68	4	0.00
OPC-4	0.31	0.53	2	0.09	0.52	2	0.13	1.09	4	0.00	1.05	3	0.00
p12,13-DiHOD*	0.16	0.91	6	0.00	0.85	5	0.05	0.82	5	0.00	0.18	1	0.69
p9,10-DiHOD*	0.63	1.75	3	0.02	1.48	2	0.04	1.73	3	0.01	0.91	1	0.42
p9OH-TAN*	3.62	3.35	1	0.62	3.36	1	0.71	8.86	2	0.00	4.07	1	0.46
pOTD*	0.47	0.42	1	0.72	0.52	1	0.71	1.19	3	0.00	0.36	1	0.33
SA	67.53	31.73	0	0.21	55.50	1	0.69	173.44	3	0.01	55.09	1	0.65
TA	6.03	10.56	2	0.04	7.94	1	0.19	17.41	3	0.00	7.35	1	0.23
TAN	0.54	0.77	1	0.32	0.53	1	0.91	0.74	1	0.01	0.60	1	0.51

[a] Compound color key: Blue fill= 13-LOX pathway derived compound; Red fill= 9-LOX pathway derived compound; Purple fill= phenylpropanoid pathway derived compound; Gray fill= other; metabolites were measured with LC/MS/MS

[b] Xylem sap harvested from mock-inoculated/treated healthy BB plants

[c] Sap harvested from DI=1 *Rs* infected plants

[d] Fold-change values ≥ 2 are shaded in light red with dark red text.

[e] P-value derived from t-test of treatment compared to healthy; *P*<0.5 was shaded yellow with darker yellow text.

[f-g] Xylem sap harvested from uninoculated, trehalose treated plants 6, 24, and 48 h post-treatment

898   **Supplemental Table 4. Metabolite key for hormone analysis**

Q1[a]	Q3[b]	RT[c]	Metabolite	Formal Name
313.3	183.3	10.16	12,13-DiHOM	threo-12,13-dihydroxy-9(Z)-octadecenoic acid
309.4	208.9	12.46	(E)-KODA	12,13-epoxy-9-oxo-10(E)-octadecenoic acid
225	59	2.44	12OH-JA	12-hydroxy-jasmonic acid
338.3	130.1	3.19	12OH-JA-Ile	12-hydroxy-jasmonic acid isoluecine
291.3	165	11.13	12-OPDA	12-oxo-10(Z),15(Z)-phytodienoic acid
295.3	195.2	13.46	13-HOD	13(S)-hydroxy-9(Z),11(E)-octadecatrienoic acid
293.2	113.2	14.16	13-KOD	13-oxo-9(Z),11(E)-octadecadienoic acid
309.4	180.8	10.32	13,12-KODA	13-hydroxy-12-oxo-9(Z),15(Z)-octadecadienoic acid
311.3	194.9	11.42	13,12-KOMA	13-hydroxy-12-oxo-9(Z)-octadecenoic acid
293.2	191.1	15.15	2-HOD	2-hydroxy-9(Z),15(Z)-octadecadienoic acid
327.1	137	5.57	9,10,11-THOD	9(S),10(S),11(R)-trihydroxy-12(Z),15(Z)-octadecadienoic acid
329.5	201	6.69	9,10,11-THOM	9(S),10(S),11(R)-trihydroxy-12(Z)-octadecenoic acid
327.1	137	4.94	9,10,13-THOD	9(S),10(S),13(S)-trihydroxy-11(E),15(Z)-octadecadienoic acid
329.5	139	5.8	9,10,13-THOM	9(S),10(S),13(S)-trihydroxy-11(E)-octadecenoic acid
313.2	201.2	10.46	9,10-DiHOM	threo-9,10-dihydroxy-12(Z)-octadecenoic acid
326.9	229.1	5	9,12,13-THOD	9(S),12(S),13(S)-trihydroxy-10(E),15(Z)-octadecadienoic acid
329.5	210.9	5.74	9,12,13-THOM	9(S),12(S),13(S)-trihydroxy-10(E)-octadecenoic acid
295.3	171.1	13.53	9-HOD	9(S)-hydroxy-10(E),12(Z)-octadecadienoic acid
293.2	171.2	12.04	9-HOT	9(S)-hydroxy-10(E),12(Z),15(Z)-octadecatrienoic acid
293.2	195.1	12.25	9-KOD	9(S)-oxo-10(E),12(Z)-octadecatrienoic acid
291.3	184.7	13.04	9-KOT	9(S)-oxo-10(E),12(Z),15(Z)-octadecatrienoic acid
311.3	293.2	11.76	9,10-KOMA	9-hydroxy-10-oxo-12(Z)-octadecadienoic acid
309.1	291.3	9.02	9,12-KODA	9-hydroxy-12-oxo-10(E),15(Z)-octadecadienoic acid
311.3	293.1	10.29	9,12-KOMA	9-hydroxy-12-oxo-10(E)-octadecenoic acid
263.4	153.1	3.33	ABA	Abscisic acid
186.9	124.9	2.83	AZA	Azelaic Acid
121.1	76.9	2.54	BA	Benzoic acid
147	102.9	3.6	CA	Cinnamic acid
162.9	118.9	2.05	COUMA	Coumaric acid
263.1	165	8.29	dn-12-OPDA	dinor-12-oxo-8(Z),13(Z)-phytodienoic acid
174	129.9	2.77	IAA	Indole-3-acetic acid
209.1	59	4.24	JA	(+)-7-iso-jasmonic acid
322.2	130	6.82	JA-Ile	(+)-7-Jasmonoyl-L-isoleucine
237	165.1	6.38	OPC-4	8-[3-oxo-2-cis-[(Z)-2-pentenylcyclopentyl]butanoic acid
311.2	183.2	10.3	12,13-DiHOD*	threo-12,13-dihydroxy-9(Z),15(Z)-octadecenoic acid
311.2	201.2	10.35	9,10-DiHOD*	threo-9,10-dihydroxy-12(Z),15(Z)-octadecenoic acid
226.9	208.7	3.2	9OH-TAN*	9-hydroxy-12-oxo-10(E)-dodecenoic acid
223	195	7.3	OTD*	13-oxo-9(Z),11(E)-tridecadienoic acid
137	92.9	2.45	SA	Salicylic acid
226.9	182.7	5.36	TA	Traumatic acid
210.9	182.7	6.56	TAN	Traumatin

[A-B] Select m/z ions for compound identification (see Methods section)
[C] Compound retention time

Supplemental Figures

Supplemental Figure 1. ABA and DMSO are not toxic to *Rs*. Growth of *Rs* cultured in rich CPG broth

supplemented with ABA in DMSO, or with DMSO alone since ABA stock solutions were dissolved in

DMSO (100 μM treatments contained 0.1 μL of DMSO or ABA stock solution, and 50 μM treatments

contained 0.05 μL DMSO or ABA stock solution) (ANOVA AUC, Fisher's LSD multiple comparisons to CPG

control, 100 μM DMSO, P=.55; 50 μM DMSO, P=.53; 100 μM ABA, P=.67; 50 μM ABA, P=.45). Growth

was measured spectrophotometrically using a Bio-Tek plate reader; the data represent eight technical

reps/treatment.

Supplemental Figure 2. Trehalose treatment temporarily improves water efficiency without affecting

photosynthesis; treatment does not affect root growth and relative water content of leaves but

lowers chlorophyll concentration. A) Water efficiency, measured with a LI-COR as photosynthetic

rate/transpiration, in trehalose treated plants compared to water only controls (t-test, *P*<0.5). **B)**

Photosynthesis, measured as A$_{net}$, in trehalose treated plants (repeated measures ANOVA, *P*>0.05). The

water efficiency and photosynthetic data represent 20 plants/treatment. LI-COR measurements were

taken ~10min after treatment and then in the morning every time point after. Shaded bars on the

graphs represent night periods. All data were calculated from the same LICOR time points. **C)** Relative

water content of leaves was calculated from the fresh weight, hydrated weight, and dry weight of leaves

from water control or trehalose treated BB tomatoes (t-test, *P*=.14). The data represent two biological

replicates with five technical replicates (leaves)/treatment. **D)** 'Bonny Best' tomato seedlings were

placed in root growth pouches, watered with 10mL of either water or trehalose for one day, then

watered as needed for ten days. Root length was measured after ten days (t-test, *P*=.31). The data

represent three biological replicates with ten roots/treatment. %New root growth was calculated as

%new root growth= (new growth/total root length)*100. **E)** SPAD meter measurements of chlorophyll

content in leaves following trehalose treatment (2-way ANOVA Fishers LSD multiple comparison, * =

P<0.5 or lower).

Supplemental Figure 3 Trehalose treatment does not enrich the soil environment, and its protection is

limited to roots. A) Disease progress curve of plants treated with trehalose or water, then infected with

either wild-type *Rs* or a Δ*treA* mutant unable to catabolize trehalose (One-way ANOVA of areas under

the curve, WT H_2O vs. WT tre, *P*=.0008, Δ*treA* H_2O vs. Δ*treA* tre, *P*=.017). The data represent three

bioreps with 13-15 plants per treatment. **B)** Disease development in plants sprayed once with 30 mM

trehalose or water, and then soil-soak inoculated with *Rs* 48 h later (ANOVA of AUC, Fisher's LSD

multiple comparisons to H_2O soil, H_2O spray, *P*=.67; tre spray, *P*=.63; tre soil, *P*=.63). The data represent

three biological replicates with fifteen plants per treatment. **C)** Colonization of tomato leaves by the

bacterial leaf spot pathogen *Xanthomonas gardneri*; plants were dip-inoculated with *X. gardneri* 72 h

after root treatment with water or 30 mM trehalose (Mann-Whitney, day 3, *P*=.06; day 6, *P*=.61; day 9,

P=08)*. The data represent three biological replicates with twelve plants/treatment/timepoint. The graph

displays the geometric means; bars indicate the geometric standard deviation of the data.

Supplemental Figure 4 Infusing leaves with trehalose did not trigger ROS production in 'Bonny Best'

tomato leaves. DAB staining of 'Bonny best' tomato leaves infused with water, 10^9 CFU/mL *Rs*, or 30

mM trehalose solution to assess the effect of trehalose treatment on ROS production. The data

represent three biological replicates, with four plants per biological replicate per treatment. Photos are

representative samples and sharpened 25% for contrast.

References

Abd El-Rahim, M. F., G. M. Fahmy and Z. M. Fahmy (1998). "Alterations in transpiration and stem
vascular tissues of two maize cultivars under conditions of water stress and late wilt disease." Plant
Pathology **47**(2): 216-223.
Alméras, E., S. Stolz, S. Vollenweider, P. Reymond, L. Mène-Saffrané and E. E. Farmer (2003). "Reactive
electrophile species activate defense gene expression in *Arabidopsis*." The Plant Journal **34**(2): 205-
216.
Álvarez, B., E. G. Bosca and M. M. López (2010). "On the life of *Ralstonia solanacearum*, a destructive
bacterial plant pathogen." Technology and Education Topics in Applied Mcrobiology and Microbial
Biotechnology: 267–279.
An, C. and Z. Mou (2011). "Salicylic acid and its function in plant immunity." Journal of Integrative Plant
Biology **53**(6): 412-428.
Anders, S., P. T. Pyl and W. Huber (2014). "HTSeq—a Python framework to work with high-throughput
sequencing data." Bioinformatics **31**(2): 166-169.
Andrews, S. FastQC A Quality Control Tool for High Throughput Sequence Data, Babraham
Bioinformatics.
Asaf, S., A. L. Khan, M. A. Khan, Q. M. Imran, B.-W. Yun and I.-J. Lee (2017). "Osmoprotective functions
conferred to soybean plants via inoculation with *Sphingomonas* sp. *LK11* and exogenous trehalose."
Microbiological Research **205**: 135-145.
Avonce, N., B. Leyman, J. O. Mascorro-Gallardo, P. Van Dijck, J. M. Thevelein and G. Iturriaga (2004).
"The *Arabidopsis* trehalose-6-P synthase AtTPS1 gene is a regulator of glucose, abscisic acid, and
stress signaling." Plant Physiology **136**(3): 3649-3659.
Bae, H., E. Herman, B. Bailey, H.-J. Bae and R. Sicher (2005). "Exogenous trehalose alters *Arabidopsis*
transcripts involved in cell wall modification, abiotic stress, nitrogen metabolism, and plant
defense." Physiologia Plantarum **125**(1): 114-126.
Barrs, H. D. and P. E. Weatherley (1962). "A re-examination of the relative turgidity technique for
estimating water deficits in leaves." Australian Journal of Biological Sciences **15**(3): 413-428.
Beattie, G. A. and S. E. Lindow (1995). "The secret life of foliar bacterial pathogens on leaves." Annu Rev
Phytopathol **33**: 145-172.
Beckman, C. H. (2000). "Phenolic-storing cells: keys to programmed cell death and periderm formation
in wilt disease resistance and in general defence responses in plants?" Physiological and Molecular
Plant Pathology **57**(3): 101-110.
Benjamini, Y. and Y. Hochberg (1995). "Controlling the false discovery rate: A practical and powerful
approach to multiple testing." Journal of the Royal Statistical Society. Series B (Methodological)
57(1): 289-300.
Bian, C., Y. Duan, J. Wang, Q. Xiu, J. Wang, Y. Hou, X. Song and M. Zhou (2020). "Validamycin A induces
broad-spectrum resistance involving salicylic acid and jasmonic acid/ethylene signaling pathways."
Molecular Plant-Microbe Interactions **33**(12): 1424-1437.
Block, A., E. Schmelz, P. J. O'Donnell, J. B. Jones and H. J. Klee (2005). "Systemic acquired tolerance to
virulent bacterial pathogens in tomato." Plant Physiology **138**(3): 1481-1490.

Boatwright, J. L. and K. Pajerowska-Mukhtar (2013). "Salicylic acid: an old hormone up to new tricks."
Molecular Plant Pathology **14**(6): 623-634.

Bolger, A. M., M. Lohse and B. Usadel (2014). "Trimmomatic: a flexible trimmer for Illumina sequence
data." Bioinformatics **30**(15): 2114-2120.

Bolouri Moghaddam, M. R. and W. Van den Ende (2012). "Sugars and plant innate immunity." Journal of
Experimental Botany **63**(11): 3989-3998.

Brading, P. A., K. E. Hammond-Kosack, A. Parr and J. D. G. Jones (2000). "Salicylic acid is not required for
Cf-2- and *Cf-9*-dependent resistance of tomato to *Cladosporium fulvum*." The Plant Journal **23**(3):
305-318.

Brown, D. G., J. K. Swanson and C. Allen (2007). "Two host-induced *Ralstonia solanacearum* genes, *acrA*
and *dinF*, encode multidrug efflux pumps and contribute to bacterial wilt virulence." Applied and
Environmental Microbiology **73**(9): 2777-2786.

Cao, J. and X. Li (2015). "Identification and phylogenetic analysis of late embryogenesis abundant
proteins family in tomato (*Solanum lycopersicum*)." Planta **241**(3): 757-772.

Chen, K., G.-J. Li, R. A. Bressan, C.-P. Song, J.-K. Zhu and Y. Zhao (2020). "Abscisic acid dynamics,
signaling, and functions in plants." Journal of Integrative Plant Biology **62**(1): 25-54.

Chen, Y. Y., Y. M. Lin, T. C. Chao, J. F. Wang, A. C. Liu, F. I. Ho and C. P. Cheng (2009). "Virus-induced gene
silencing reveals the involvement of ethylene-, salicylic acid- and mitogen-activated protein kinase-
related defense pathways in the resistance of tomato to bacterial wilt." Physiologia Plantarum
136(3): 324-335.

Chen, Z., Z. Zheng, J. Huang, Z. Lai and B. Fan (2009). "Biosynthesis of salicylic acid in plants." Plant
signaling & behavior **4**(6): 493-496.

Choudhary, D. K., A. Prakash and B. N. Johri (2007). "Induced systemic resistance (ISR) in plants:
mechanism of action." Indian Journal of Microbiology **47**(4): 289-297.

Cortina, C. and F. A. Culiáñez-Macià (2005). "Tomato abiotic stress enhanced tolerance by trehalose
biosynthesis." Plant Science **169**(1): 75-82.

Daugherty, M. P., J. R. S. Lopes and R. P. P. Almeida (2010). "Strain-specific alfalfa water stress induced
by *Xylella fastidiosa*." European Journal of Plant Pathology **127**(3): 333-340.

De Lorenzo, G., S. Ferrari, F. Cervone and E. Okun (2018). "Extracellular DAMPs in plants and mammals:
Immunity, tissue damage and repair." Trends in Immunology **39**(11): 937-950.

Denancé, N., P. Ranocha, N. Oria, X. Barlet, M.-P. Rivière, K. A. Yadeta, L. Hoffmann, F. Perreau, G.
Clément, A. Maia-Grondard, G. C. M. van den Berg, B. Savelli, S. Fournier, Y. Aubert, S. Pelletier, B. P.
H. J. Thomma, A. Molina, L. Jouanin, Y. Marco and D. Goffner (2013). "*Arabidopsis wat1* (*walls are*
thin1)-mediated resistance to the bacterial vascular pathogen, *Ralstonia solanacearum*, is
accompanied by cross-regulation of salicylic acid and tryptophan metabolism." The Plant Journal
73(2): 225-239.

Denny, T. P. (2006). Plant pathogenic *Ralstonia* species. Plant-Associated Bacteria. S. S. Gnanamanickam,
Springer**:** 573–644.

Derksen, H., C. Rampitsch and F. Daayf (2013). "Signaling cross-talk in plant disease resistance." Plant
Science **207**: 79-87.

Deslandes, L. and S. Genin (2014). "Opening the *Ralstonia solanacearum* type III effector tool box:
insights into host cell subversion mechanisms." Current Opinion in Plant Biology **20**: 110-117.

Deslandes, L., J. Olivier, Theuli, xe, F. res, xe, xe, ric, J. Hirsch, D. X. Feng, P. Bittner-Eddy, J. Beynon and Y.
Marco (2002). "Resistance to *Ralstonia solanacearum* in *Arabidopsis thaliana* is conferred by the
recessive RRS1-R gene, a member of a novel family of resistance genes." Proceedings of the National
Academy of Sciences **99**(4): 2404-2409.
Digonnet, C., Y. Martinez, N. Denance, M. Chasseray, P. Dabos, P. Ranocha, Y. Marco, A. Jauneau and D.
Goffner (2012). "Deciphering the route of *Ralstonia solanacearum* colonization in *Arabidopsis*
thaliana roots during a compatible interaction: focus at the plant cell wall." Planta **236**(5): 1419-
1431.
Dobin, A., C. A. Davis, F. Schlesinger, J. Drenkow, C. Zaleski, S. Jha, P. Batut, M. Chaisson and T. R.
Gingeras (2012). "STAR: ultrafast universal RNA-seq aligner." Bioinformatics **29**(1): 15-21.
Elphinstone, J. G. (2005). The Current Bacterial Wilt Situation: A Global Overview. Bacterial Wilt Disease
and the *Ralstonia solanacearum* Species Complex. C. Allen, P. Prior and A. C. Hayward. St. Paul, MN,
APS Press.
Flores-Cruz, Z. and C. Allen (2009). "*Ralstonia solanacearum* encounters an oxidative environment
during tomato infection." Molecular Plant-Microbe Interactions **22**(7): 773–782.
French, E., B. S. Kim, K. Rivera-Zuluaga and A. S. Iyer-Pascuzzi (2018). "Whole root transcriptomic
analysis suggests a role for auxin pathways in resistance to *Ralstonia solanacearum* in tomato."
Molecular Plant-Microbe Interactions **31**(4): 432-444.
Garg, A. K., J.-K. Kim, T. G. Owens, A. P. Ranwala, Y. D. Choi, L. V. Kochian and R. J. Wu (2002). "Trehalose
accumulation in rice plants confers high tolerance levels to different abiotic stresses." Proceedings
of the National Academy of Sciences **99**(25).
Ge, S. X., D. Jung and R. Yao (2019). "ShinyGO: a graphical gene-set enrichment tool for animals and
plants." Bioinformatics **36**(8): 2628-2629.
Glazebrook, J. (2005). "Contrasting mechanisms of defense against biotrophic and necrotophic
pathogens." Annual Review of Phytopathology **43**(1): 205-227.
Godoy, J. A., J. M. Pardo and J. A. Pintor-Toro (1990). "A tomato cDNA inducible by salt stress and
abscisic acid: nucleotide sequence and expression pattern." Plant Molecular Biology **15**(5): 695-705.
Gomez, L. D., A. Gilday, R. Feil, J. E. Lunn and I. A. Graham (2010). "AtTPS1-mediated trehalose 6-
phosphate synthesis is essential for embryogenic and vegetative growth and responsiveness to ABA
in germinating seeds and stomatal guard cells." The Plant Journal **64**(1): 1-13.
Goodstein, D. M., S. Shu, R. Howson, R. Neupane, R. D. Hayes, J. Fazo, T. Mitros, W. Dirks, U. Hellsten, N.
Putnam and D. S. Rokhsar (2011). "Phytozome: a comparative platform for green plant genomics."
Nucleic Acids Research **40**(D1): D1178-D1186.
Hamilton, C. D., O. R. Steidl, A. M. MacIntyre, C. G. Hendrich and C. Allen (2021). "*Ralstonia*
solanacearum depends on catabolism of myo-inositol, sucrose, and trehalose for virulence in an
infection stage-dependent manner." Molecular Plant-Microbe Interactions.
Hase, S., S. Takahashi, S. Takenaka, K. Nakaho, T. Arie, S. Seo, Y. Ohashi and H. Takahashi (2008).
"Involvement of jasmonic acid signalling in bacterial wilt disease resistance induced by biocontrol
agent *Pythium oligandrum* in tomato." Plant Pathology **57**(5): 870-876.
Hassan, M. A. E. and K. A. M. Abo-Elyousr (2013). "Activation of tomato plant defence responses against
bacterial wilt caused by *Ralstonia solanacearum* using DL-3-aminobutyric acid (BABA)." European
Journal of Plant Pathology **136**(1): 145-157.

He, Y., E. J. Borrego, Z. Gorman, P.-C. Huang and M. V. Kolomiets (2020). "Relative contribution of LOX10, green leaf volatiles and JA to wound-induced local and systemic oxylipin and hormone signature in *Zea mays* (maize)." Phytochemistry **174**: 112334.

He, Y., H. Fukushige, D. F. Hildebrand and S. Gan (2002). "Evidence supporting a role of jasmonic acid in *Arabidopsis* leaf senescence." Plant Physiology **128**(3): 876.

Hendrick, C. A. and L. Sequeira (1984). "Lipopolysaccharide defective mutants of the wilt pathogen *Pseudomonas solanacearum*." Applied and Environmental Microbiology **48**(1): 94-101.

Hernández-Blanco, C., D. X. Feng, J. Hu, A. Sánchez-Vallet, L. Deslandes, F. Llorente, M. Berrocal-Lobo, H. Keller, X. Barlet, C. Sánchez-Rodríguez, L. K. Anderson, S. Somerville, Y. Marco and A. Molina (2007). "Impairment of cellulose synthases required for *Arabidopsis* secondary cell wall formation enhances disease resistance." The Plant Cell **19**(3): 890.

Hirsch, J., L. Deslandes, D. X. Feng, C. Balagué and Y. Marco (2002). "Delayed symptom development in *ein2-1*, an *Arabidopsis* ethylene-insensitive mutant, in response to bacterial wilt caused by *Ralstonia solanacearum*." Phytopathology **92**(10): 1142-1148.

Hou, S., Z. Liu, H. Shen and D. Wu (2019). "Damage-Associated Molecular Pattern-triggered immunity in plants." Frontiers in Plant Science **10**: 646.

Hu, J., X. Barlet, L. Deslandes, J. Hirsch, D. X. Feng, I. Somssich and Y. Marco (2008). "Transcriptional responses of *Arabidopsis thaliana* during wilt disease caused by the soil-borne phytopathogenic bacterium, *Ralstonia solanacearum*." PLOS ONE **3**(7): e2589-e2589.

Ingel, B., D. Caldwell, F. Duong, D. Y. Parkinson, K. A. McCulloh, A. S. Iyer-Pascuzzi, A. J. McElrone and T. M. Lowe-Power (2021). "Revisiting the source of wilt symptoms: X-ray microcomputed tomography provides direct evidence that *Ralstonia* biomass clogs xylem vessels." bioRxiv: 2021.2003.2019.436187.

Iordachescu, M. and R. Imai (2008). "Trehalose biosynthesis in response to abiotic stresses." Journal of Integrative Plant Biology **50**(10): 1223-1229.

Ishihara, T., I. Mitsuhara, H. Takahashi and K. Nakaho (2012). "Transcriptome analysis of quantitative resistance-specific response upon Ralstonia solanacearum infection in tomato." PLOS ONE **7**(10): e46763.

Ishikawa, R., K. Fujimori and K. Matsuura (1996). "Antibacterial activity of Validamycin A against *Pseudomonas solanacearum* and its efficacy against tomato bacterial wilt." Annals of the Phytopathological Society of Japan **62**(5): 478-482.

Ishikawa, R., K. Shirouzu, H. Nakashita, H.-Y. Lee, T. Motoyama, I. Yamaguchi, T. Teraoka and T. Arie (2005). "Foliar spray of Validamycin A or Validoxylamine A controls tomato *Fusarium* wilt." Phytopathology **95**(10): 1209-1216.

Jackson, M. (1997). "Hormones from roots as signals for the shoots of stressed plants." Trends in Plant Science **2**(1): 22-28.

Jacobs, J. M., L. Babujee, F. Meng, A. Milling and C. Allen (2012). "The *in planta* transcriptome of *Ralstonia solanacearum*: conserved physiological and virulence strategies during bacterial wilt of tomato." MBio **3**(4).

Jacobs, J. M., A. Milling, R. M. Mitra, C. S. Hogan, F. Ailloud, P. Prior and C. Allen (2013). "*Ralstonia solanacearum* requires PopS, an ancient AvrE-family effector, for virulence and to overcome salicylic acid-mediated defenses during tomato pathogenesis." MBio **4**(6): e00875-00813.

Jakab, G., V. Cottier, V. Toquin, G. Rigoli, L. Zimmerli, J.-P. Métraux and B. Mauch-Mani (2001). "β-
Aminobutyric acid-induced resistance in plants." European Journal of Plant Pathology **107**(1): 29-37.
Ji, K., W. Kai, B. Zhao, Y. Sun, B. Yuan, S. Dai, Q. Li, P. Chen, Y. Wang, Y. Pei, H. Wang, Y. Guo and P. Leng
(2014). "SlNCED1 and SlCYP707A2: key genes involved in ABA metabolism during tomato fruit
ripening." Journal of Experimental Botany **65**(18): 5243-5255.
Johnson, K. L., A. M. Cassin, A. Lonsdale, G. K.-S. Wong, D. E. Soltis, N. W. Miles, M. Melkonian, B.
Melkonian, M. K. Deyholos, J. Leebens-Mack, C. J. Rothfels, D. W. Stevenson, S. W. Graham, X.
Wang, S. Wu, J. C. Pires, P. P. Edger, E. J. Carpenter, A. Bacic, M. S. Doblin and C. J. Schultz (2017).
"Insights into the evolution of Hydroxyproline-Rich Glycoproteins from 1000 plant transcriptomes."
Plant Physiology **174**(2): 904.
Joosten, M. H. A. J. and P. J. G. M. De Wit (1989). "Identification of several Pathogenesis-Related
proteins in tomato leaves inoculated with *Cladosporium fulvum* (syn. *Fulvia fulva*) as 1,3-β-
glucanases and chitinases." Plant Physiology **89**(3): 945.
Kapilan, R., M. Vaziri and J. J. Zwiazek (2018). "Regulation of aquaporins in plants under stress."
Biological Research **51**(1): 4.
Khater, M. A., M. G. Dawood, M. S. Sadak, M. A. F. Shalaby, M. E. El-Awadi and K. Gamal El-Din (2018).
"Enhancement of the performance of cowpea plants grown under drought conditions via trehalose
application." Middle East Journal of Agriculture Research **7**(3): 782-800.
Khokhani, D., T. M. Lowe-Power, T. M. Tran and C. Allen (2017). "A single regulator mediates strategic
switching between attachment/spread and growth/virulence in the plant pathogen *Ralstonia*
solanacearum." MBio **8**(5).
Khokhani, D., T. Tuan, T. Lowe-Power and C. Allen (2018). "Plant assays for quantifying *Ralstonia*
solanacearum virulence." Bio-Protocol **8**(18).
Kissoudis, C., A. Seifi, Z. Yan, A. T. Islam, H. van der Schoot, C. C. van de Wiel, R. G. Visser, C. G. van der
Linden and Y. Bai (2016). "Ethylene and abscisic acid signaling pathways differentially influence
tomato resistance to combined powdery mildew and salt stress." Frontiers in Plant Science **7**: 2009.
Kolbe, A., A. Tiessen, H. Schluepmann, M. Paul, S. Ulrich and P. Geigenberger (2005). "Trehalose 6-
phosphate regulates starch synthesis via posttranslational redox activation of ADP-glucose
pyrophosphorylase." Proceedings of the National Academy of Sciences **102**(31): 11118–11123.
Kolde, R. (2019). "pheatmap: Pretty Heatmaps."   Retrieved June 7, 2020, from https://cran.r-
project.org/web/packages/pheatmap/index.html.
Kushiro, T., M. Okamoto, K. Nakabayashi, K. Yamagishi, S. Kitamura, T. Asami, N. Hirai, T. Koshiba, Y.
Kamiya and E. Nambara (2004). "The Arabidopsis cytochrome P450 CYP707A encodes ABA 8'-
hydroxylases: key enzymes in ABA catabolism." The EMBO Journal **23**(7): 1647-1656.
Lanahan, M. B., H.-C. Yen, J. J. Giovannoni and H. J. Klee (1994). "The *Never Ripe* mutation blocks
ethylene perception in tomato." The Plant Cell **6**: 521-530.
Leyman, B., P. Van Dijck and J. M. Thevelein (2001). "An unexpected plethora of trehalose biosynthesis
genes in *Arabidopsis thaliana*." Trends in Plant Science **6**(11): 510-513.
Li, H., B. Handsaker, A. Wysoker, T. Fennell, J. Ruan, N. Homer, G. Marth, G. Abecasis, R. Durbin and S.
Genome Project Data Processing (2009). "The Sequence Alignment/Map format and SAMtools."
Bioinformatics **25**(16): 2078-2079.

Lin, Q., S. Wang, Y. Dao, J. Wang and K. Wang (2020). "The *Arabidopsis thaliana* trehalose-6-phosphate
phosphatase gene *AtTPPI* enhances drought tolerance by regulating stomatal apertures." <u>Journal of</u>
<u>Experimental Botany</u>.
Lin, Q., J. Yang, Q. Wang, H. Zhu, Z. Chen, Y. Dao and K. Wang (2019). "Overexpression of the trehalose-
6-phosphate phosphatase family gene *AtTPPF* improves the drought tolerance of *Arabidopsis*
thaliana." <u>BMC Plant Biology</u> **19**(1): 381.
Ling, Q., W. Huang and P. Jarvis (2011). "Use of a SPAD-502 meter to measure leaf chlorophyll
concentration in *Arabidopsis thaliana*." <u>Photosynth Res</u> **107**(2): 209-214.
Liu, H., S. Zhang, M. A. Schell and T. P. Denny (2005). "Pyramiding unmarked deletions in *Ralstonia*
solanacearum shows that secreted proteins in addition to plant cell-wall-degrading enzymes
contribute to virulence." <u>Molecular Plant-Microbe Interactions</u> **18**(12): 1296-1305.
Liu, H., S. Zhang, M. A. Schell and T. P. Denny (2005). "Pyramiding unmarked deletions in *Ralstonia*
solanacearum shows that secreted proteins in addition to plant cell-wall-degrading enzymes
contribute to virulence." <u>Molecular Plant-Microbe Interactions</u>® **18**(12): 1296-1305.
López-Gresa, M. P., P. Lisón, L. Yenush, V. Conejero, I. Rodrigo and J. M. Bellés (2016). "Salicylic acid is
involved in the basal resistance of tomato plants to Citrus Excortis Viroid and Tomato Spotted Wilt
Virus." <u>PLOS ONE</u> **11**(11): e0166938.
Love, M. I., W. Huber and S. Anders (2014). "Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2." <u>Genome Biology</u> **15**(12): 550.
Lowe, T. M., F. Ailloud and C. Allen (2015). "Hydroxycinnamic acid degradation, a broadly conserved
trait, protects *Ralstonia solanacearum* from chemical plant defenses and contributes to root
colonization and virulence." <u>Molecular Plant-Microbe Interactions</u> **28**(3): 286-297.
Lowe-Power, T. M., C. G. Hendrich, E. von Roepenack-Lahaye, B. Li, D. Wu, R. Mitra, B. L. Dalsing, P.
Ricca, J. Naidoo, D. Cook, A. Jancewicz, P. Masson, B. Thomma, T. Lahaye, A. J. Michael and C. Allen
(2018). "Metabolomics of tomato xylem sap during bacterial wilt reveals *Ralstonia solanacearum*
produces abundant putrescine, a metabolite that accelerates wilt disease." <u>Environmental</u>
<u>Microbiology</u> **20**(4): 1330-1349.
Lowe-Power, T. M., J. M. Jacobs, F. Ailloud, B. Fochs, P. Prior and C. Allen (2016). "Degradation of the
plant defense signal salicylic acid protects *Ralstonia solanacearum* from toxicity and enhances
virulence on tobacco." <u>MBio</u> **7**(3).
Lowe-Power, T. M., D. Khokhani and C. Allen (2018). "How *Ralstonia solanacearum* exploits and thrives
in the flowing plant xylem environment." <u>Trends Microbiol</u> **26**(11): 929-942.
Lunn, J. E. (2007). "Gene families and evolution of trehalose metabolism in plants." <u>Functional Plant</u>
<u>Biology</u> **34**(6).
Lunn, J. E., I. Delorge, C. M. Figueroa, P. Van Dijck and M. Stitt (2014). "Trehalose metabolism in plants."
<u>The Plant Journal</u> **79**(4): 544-567.
Lunn, J. E., R. Feil, J. H. Hendriks, Y. Gibon, R. Morcuende, D. Osuna, W. R. Scheible, P. Carillo, M. R.
Hajirezaei and M. Stitt (2006). "Sugar-induced increases in trehalose 6-phosphate are correlated
with redox activation of ADP-glucose pyrophosphorylase and higher rates of starch synthesis in
Arabidopsis thaliana." <u>Biochem J</u> **397**(1): 139-148.
Lyu, J. I., S. R. Min, J. H. Lee, Y. H. Lim, J.-K. Kim, C.-H. Bae and J. R. Liu (2012). "Overexpression of a
trehalose-6-phosphate synthase/phosphatase fusion gene enhances tolerance and photosynthesis

during drought and salt stress without growth aberrations in tomato." <u>Plant Cell, Tissue and Organ</u>
<u>Culture</u> **112**(2): 257-262.

Ma, Y., I. Szostkiewicz, A. Korte, D. Moes, Y. Yang, A. Christmann and E. Grill (2009). "Regulators of PP2C
Phosphatase Activity Function as Abscisic Acid Sensors." <u>Science</u> **324**(5930): 1064.

MacIntyre, A. M., J. X. Barth, M. C. Pellitteri Hahn, C. O. Scarlett, S. Genin and C. Allen (2019). "Trehalose
synthesis contributes to osmotic stress tolerance and virulence of the bacterial wilt pathogen
Ralstonia solanacearum." <u>Molecular Plant-Microbe Interactions</u>: MPMI-08-19-0218-R.

Mandal, S., I. Kar, A. K. Mukherjee and P. Acharya (2013). "Elicitor-induced defense responses in
Solanum lycopersicum against *Ralstonia solanacearum*." <u>The Scientific World Journal</u> **2013**: 561056.

Mansfield, J., S. Genin, S. Magori, V. Citovsky, M. Sriariyanum, P. Ronald, M. Dow, V. Verdier, S. V. Beer,
M. A. Machado, I. Toth, G. Salmond and G. D. Foster (2012). "Top 10 plant pathogenic bacteria in
molecular plant pathology." <u>Molecular Plant Pathology</u> **13**(6): 614-629.

Marcos, R., Y. Izquierdo, T. Vellosillo, S. Kulasekaran, T. Cascón, M. Hamberg and C. Castresana (2015).
"9-Lipoxygenase-derived oxylipins activate brassinosteroid signaling to promote cell wall-based
defense and limit pathogen infection." <u>Plant Physiology</u> **169**(3): 2324-2334.

McElrone, A. J., J. L. Sherald and I. N. Forseth (2003). "Interactive effects of water stress and xylem-
limited bacterial infection on the water relations of a host vine." <u>Journal of Experimental Botany</u>
54(381): 419-430.

Mi, H., A. Muruganujan, D. Ebert, X. Huang and P. D. Thomas (2018). "PANTHER version 14: more
genomes, a new PANTHER GO-slim and improvements in enrichment analysis tools." <u>Nucleic Acids</u>
<u>Research</u> **47**(D1): D419-D426.

Milling, A., L. Babujee and C. Allen (2011). "*Ralstonia solanacearum* extracellular polysaccharide is a
specific elicitor of defense responses in wilt-resistant tomato plants." <u>PLOS ONE</u> **6**(1): e15853.

Montillet, J.-L., J.-P. Agnel, M. Ponchet, F. Vailleau, D. Roby and C. Triantaphylidès (2002).
"Lipoxygenase-mediated production of fatty acid hydroperoxides is a specific signature of the
hypersensitive reaction in plants." <u>Plant Physiology and Biochemistry</u> **40**(6): 633-639.

Mori, Y., K. Inoue, K. Ikeda, H. Nakayashiki, C. Higashimoto, K. Ohnishi, A. Kiba and Y. Hikichi (2016). "The
vascular plant-pathogenic bacterium *Ralstonia solanacearum* produces biofilms required for its
virulence on the surfaces of tomato cells adjacent to intercellular spaces." <u>Molecular Plant</u>
<u>Pathology</u> **17**(6): 890-902.

Müller, M. and S. Munné-Bosch (2011). "Rapid and sensitive hormonal profiling of complex plant
samples by liquid chromatography coupled to electrospray ionization tandem mass spectrometry."
<u>Plant Methods</u> **7**(1): 37.

Orgambide, G., H. Montrozier, P. Servin, J. Roussel, D. Trigalet-Demery and A. Trigaley (1991). "High
heterogeneity of the exopolysaccharides of *Pseudomonas solanacearum* strain GMI1000 and the
complete structure of the major polysaccharide." <u>Journal of Biological Chemistry</u> **266**(13): 8312-
8321.

Osakabe, Y., K. Osakabe, K. Shinozaki and L.-S. Tran (2014). "Response of plants to water stress."
<u>Frontiers in Plant Science</u> **5**: 86.

Park, J., Y. Lee, E. Martinoia and M. Geisler (2017). "Plant hormone transporters: what we know and
what we would like to know." <u>BMC biology</u> **15**(1): 93-93.

Paul, M. (2007). "Trehalose 6-phosphate." <u>Current Opinion in Plant Biology</u> **10**(3): 303-309.

Paul, M. J., A. Gonzalez-Uriarte, C. A. Griffiths and K. Hassani-Pak (2018). "The role of trehalose 6-
phosphate in crop yield and resilience." Plant Physiology **177**(1): 12-23.
Paul, M. J., L. F. Primavesi, D. Jhurreea and Y. Zhang (2008). "Trehalose metabolism and signaling." Annu
Rev Plant Biol **59**: 417-441.
Peña-Cortés, H., J. Fisahn and L. Willmitzer (1995). "Signals involved in wound-induced proteinase
inhibitor II gene expression in tomato and potato plants." Proceedings of the National Academy of
Sciences **92**: 4106-4113.
Piazza, A., T. Zimaro, B. S. Garavaglia, F. A. Ficarra, L. Thomas, C. Marondedze, R. Feil, J. E. Lunn, C.
Gehring, J. Ottado and N. Gottig (2015). "The dual nature of trehalose in citrus canker disease: a
virulence factor for *Xanthomonas citri* subsp. *citri* and a trigger for plant defence responses." Journal
of Experimental Botany **66**(9): 2795-2811.
Poueymiro, M, A C Cazalé, J M François, J L Parrou, N Peeters, and S Genin. 2014. "A *Ralstonia
solanacearum* Type III Effector Directs the Production of the Plant Signal Metabolite Trehalose-6-
Phosphate." *MBio* 5 (6): 1–9. https://doi.org/10.1128/mBio.02065-14.

Potnis, N., J. P. Soto-Arias, K. N. Cowles, A. H. C. van Bruggen, J. B. Jones and J. D. Barak (2014).
"*Xanthomonas perforans* colonization influences *Salmonella enterica* in the tomato phyllosphere."
Applied and environmental microbiology **80**(10): 3173-3180.
Pradhanang, P. M., P. Ji, M. T. Momol, S. M. Olson, J. L. Mayfield and J. B. Jones (2005). "Application of
Acibenzolar-S-Methyl enhances host resistance in tomato against *Ralstonia solanacearum*." Plant
Disease **89**(9): 989-993.
Prost, I., S. Dhondt, G. Rothe, J. Vicente, M. J. Rodriguez, N. Kift, F. Carbonne, G. Griffiths, M.-T.
Esquerré-Tugayé, S. Rosahl, C. Castresana, M. Hamberg and J. Fournier (2005). "Evaluation of the
antimicrobial activities of plant oxylipins supports their involvement in defense against pathogens."
Plant Physiology **139**(4): 1902.
R. "R: The R Project for Statistical Computing." Retrieved June 7, 2020, from https://www.r-
project.org/.
Raghavendra, A. S., V. K. Gonugunta, A. Christmann and E. Grill (2010). "ABA perception and signalling."
Trends in Plant Science **15**(7): 395-401.
Ranf, S., L. Eschen-Lippold, P. Pecher, J. Lee and D. Scheel (2011). "Interplay between calcium signalling
and early signalling elements during defence responses to microbe- or damage-associated molecular
patterns." The Plant Journal **68**(1): 100-113.
Rau, A., G. Marot and F. Jaffrézic (2014). "Differential meta-analysis of RNA-seq data from multiple
studies." BMC Bioinformatics **15**(1): 91.
Retig, N. and I. Chet (1974). "Catechol-induced resistance of tomato plants to *Fusarium* wilt."
Physiological Plant Pathology **4**(4): 469-475.
Reuscher, S., M. Akiyama, C. Mori, K. Aoki, D. Shibata and K. Shiratake (2013). "Genome-wide
identification and expression analysis of aquaporins in tomato." PLOS ONE **8**(11): e79052.
Robinson, M. D., D. J. McCarthy and G. K. Smyth (2009). "edgeR: a Bioconductor package for differential
expression analysis of digital gene expression data." Bioinformatics **26**(1): 139-140.

Romero, C., J. M. Bellés, J. L. Vayá, R. Serrano and A. Francisco (1997). "Expression of the yeast
trehalose-6-phosphate synthase gene in transgenic tobacco plants: Pleiotropic phenotypes include
drought tolerance." Planta **201**(3): 293-297.

Sade, N., B. J. Vinocur, A. Diber, A. Shatil, G. Ronen, H. Nissan, R. Wallach, H. Karchi and M. Moshelion
(2009). "Improving plant stress tolerance and yield production: is the tonoplast aquaporin SlTIP2;2 a
key to isohydric to anisohydric conversion?" New Phytologist **181**(3): 651-661.

Saito, S., N. Hirai, C. Matsumoto, H. Ohigashi, D. Ohta, K. Sakata and M. Mizutani (2004). "Arabidopsis
CYP707As encode (+)-abscisic acid 8'-hydroxylase, a key enzyme in the oxidative catabolism of
abscisic acid." Plant Physiology **134**(4): 1439-1449.

Sato, S., S. Tabata, H. Hirakawa and e. al (2012). "The tomato genome sequence provides insights into
fleshy fruit evolution." Nature **485**(7400): 635-641.

Savatin, D. V., G. Gramegna, V. Modesti and F. Cervone (2014). "Wounding in the plant tissue: the
defense of a dangerous passage." Frontiers in Plant Science **5**: 470.

Schluepmann, H., L. Berke and G. F. Sanchez-Perez (2012). "Metabolism control over growth: a case for
trehalose-6-phosphate in plants." Journal of Experimental Botany **63**(9): 3379-3390.

Schluepmann, H., T. Pellny, A. van Dijken, S. Smeekens and M. Paul (2003). "Trehalose 6-phosphate is
indispensable for carbohydrate utilization and growth in *Arabidopsis thaliana*." Proceedings of the
National Academy of Sciences **100**(11): 6849.

Seo, S., K. Nakaho, S. W. Hong, H. Takahashi, H. Shigemori and I. Mitsuhara (2016). "L-Histidine induces
resistance in plants to the bacterial pathogen *Ralstonia solanacearum* partially through the
activation of ethylene signaling." Plant Cell Physiol **57**(9): 1932-1942.

Shi, Y., H. Sun, X. Wang, W. Jin, Q. Chen, Z. Yuan and H. Yu (2019). "Physiological and transcriptomic
analyses reveal the molecular networks of responses induced by exogenous trehalose in plant."
PLOS ONE **14**(5): e0217204.

Shinozaki, K., K. Yamaguchi-Shinozaki and M. Seki (2003). "Regulatory network of gene expression in the
drought and cold stress responses." Current Opinion in Plant Biology **6**(5): 410-417.

Signorelli, S., L. P. Tarkowski, W. Van den Ende and D. C. Bassham (2019). "Linking Autophagy to Abiotic
and Biotic Stress Responses." Trends in Plant Science **24**(5): 413-430.

Sinha, R., A. Gupta and M. Senthil-Kumar (2016). "Understanding the impact of drought on foliar and
xylem invading bacterial pathogen stress in chickpea." Frontiers in Plant Science **7**: 902.

Sinha, R., A. Gupta and M. Senthil-Kumar (2017). "Concurrent drought stress and vascular pathogen
infection induce common and distinct transcriptomic responses in chickpea." Frontiers in Plant
Science **8**(333).

Smart, C. D., K. L. Myers, S. Restrepo, G. B. Martin and W. E. Fry (2003). "Partial resistance of tomato to
Phytophthora infestans is not dependent upon ethylene, jasmonic acid, or salicylic acid signaling
pathways." Molecular Plant-Microbe Interactions **2**: 141-148.

Song, L., S.-s. C. Huang, A. Wise, R. Castanon, J. R. Nery, H. Chen, M. Watanabe, J. Thomas, Z. Bar-Joseph
and J. R. Ecker (2016). "A transcription factor hierarchy defines an environmental stress response
network." Science **354**(6312): aag1550.

Spoel, S. H., A. Koornneef, S. M. C. Claessens, J. P. Korzelius, J. A. Van Pelt, M. J. Mueller, A. J. Buchala, J.-
P. Métraux, R. Brown, K. Kazan, L. C. Van Loon, X. Dong and C. M. J. Pieterse (2003). "NPR1

modulates cross-talk between salicylate- and jasmonate-dependent defense pathways through a
novel function in the cytosol." The Plant Cell **15**(3): 760.

Tans-Kersten, J., H. Huang and C. Allen (2001). "*Ralstonia solanacearum* needs motility for invasive
virulence on tomato." J Bacteriol **183**(12): 3597-3605.

Thordal-Christensen, H., Z. Zhang, Y. Wei and D. B. Collinge (1997). "Subcellular localization of H_2O_2 in
plants. H_2O_2 accumulation in papillae and hypersensitive response during the barley—powdery
mildew interaction." The Plant Journal **11**(6): 1187-1194.

Tian, T., Y. Liu, H. Yan, Q. You, X. Yi, Z. Du, W. Xu and Z. Su (2017). "AgriGO v2.0: a GO analysis toolkit for
the agricultural community, 2017 update." Nucleic Acids Research **45**(W1): W122-W129.

Tornero, P., J. Gadea, V. Conejero and P. Vera (1997). "Two PR-1 genes from tomato are differentially
regulated and reveal a novel mode of expression for a pathogenesis-related gene during the
hypersensitive response and development." Molecular Plant-Microbe Interactions **5**: 624-634.

Tran, T., A. MacIntyre, M. C. Hawes and C. Allen (2016). "Escaping underground Nets: Extracellular
DNases degrade plant extracellular traps and contribute to virulence of the plant pathogenic
bacterium *Ralstonia solanacearum*." PLOS Pathogens **12**(6): e1005686.

Tran, T., A. MacIntyre, D. Khokhani, M. C. Hawes and C. Allen (2016). "Extracellular DNases of *Ralstonia*
solanacearum modulate biofilms and facilitate bacterial wilt virulence." Environmental Microbiology
18(11): 4103-4117.

Umezawa, T., M. Okamoto, T. Kushiro, E. Nambara, Y. Oono, M. Seki, M. Kobayashi, T. Koshiba, Y.
Kamiya and K. Shinozaki (2006). "CYP707A3, a major ABA 8'-hydroxylase involved in dehydration and
rehydration response in Arabidopsis thaliana." The Plant Journal **46**(2): 171-182.

Vallad, G. E. and R. M. Goodman (2004). "Systemic acquired resistance and induced systemic resistance
in conventional agriculture." Crop Science **44**(6): 1920-1934.

Vandesteene, L., L. Lopez-Galvis, K. Vanneste, R. Feil, S. Maere, W. Lammens, F. Rolland, J. E. Lunn, N.
Avonce, T. Beeckman and P. Van Dijck (2012). "Expansive evolution of the trehalose-6-phosphate
phosphatase gene family in *Arabidopsis*." Plant Physiology **160**(2): 884-896.

Vandesteene, L., M. Ramon, K. Le Roy, P. Van Dijck and F. Rolland (2010). "A single active trehalose-6-P
synthase (TPS) and a family of putative regulatory TPS-like proteins in *Arabidopsis*." Molecular Plant
3(2): 406-419.

Vasse, J., P. Frey and A. Trigalet (1995). "Microscopic studies of intercellular infection and protoxylem
invasion of tomato roots by *Pseudomonas solanacearum*." Molecular Plant-Microbe Interactions
8(2): 241-251.

Vicente, J., T. Cascón, B. Vicedo, P. García-Agustín, M. Hamberg and C. Castresana (2012). "Role of 9-
lipoxygenase and α-dioxygenase oxylipin pathways as modulators of local and systemic defense."
Molecular Plant **5**(4): 914-928.

Walters, D. R., T. Cowley and H. Weber (2006). "Rapid accumulation of trihydroxy oxylipins and
resistance to the bean rust pathogen *Uromyces fabae* following wounding in *Vicia faba*." Annals of
Botany **97**(5): 779-784.

Wang, K.-D., E. J. Borrego, C. M. Kenerley and M. V. Kolomiets (2020). "Oxylipins other than jasmonic
acid are xylem resident signals regulating systemic resistance induced by *Trichoderma virens* in
maize." The Plant Cell **32**(1): 166.

Wang, W., Q. Chen, S. Xu, W.-C. Liu, X. Zhu and C.-P. Song (2020). "Trehalose-6-phosphate phosphatase
e modulates ABA-controlled root growth and stomatal movement in *Arabidopsis*." Journal of
Integrative Plant Biology.
Wickham, H. (2016). "ggplot2: Elegant graphics for data analysis." from https://ggplot2-book.org/.
Widhalm, Joshua R. and N. Dudareva (2015). "A familiar ring to it: biosynthesis of plant benzoic acids."
Molecular Plant **8**(1): 83-97.
Yamaguchi-Shinozaki, K. and K. Shinozaki (1993). "The plant hormone abscisic acid mediates the
drought-induced expression but not the seed-specific expression of *rd22*, a gene responsive to
dehydration stress in *Arabidopsis thaliana*." Molecular and General Genetics MGG **238**(1): 17-25.
Yao, J. and C. Allen (2006). "Chemotaxis is required for virulence and competitive fitness of the bacterial
wilt pathogen *Ralstonia solanacearum*." J Bacteriol **188**(10): 3697-3708.
Yao, J. and C. Allen (2007). "The plant pathogen *Ralstonia solanacearum* needs aerotaxis for normal
biofilm formation and interactions with its tomato host." J Bacteriol **189**(17): 6415-6424.
Yu, W., R. Zhao, L. Wang, S. Zhang, R. Li, J. Sheng and L. Shen (2019). "ABA signaling rather than ABA
metabolism is involved in trehalose-induced drought tolerance in tomato plants." Planta **250**(2):
643-655.
Zhao, Y., R. Thilmony, C. L. Bender, A. Schaller, S. Y. He and G. A. Howe (2003). "Virulence systems of
Pseudomonas syringae pv. *tomato* promote bacterial speck disease in tomato by targeting the
jasmonate signaling pathway." The Plant Journal **36**(4): 485-499.
Zhou, R., X. Yu, T. Zhao, C.-O. Ottosen, E. Rosenqvist and Z. Wu (2019). "Physiological analysis and
transcriptome sequencing reveal the effects of combined cold and drought on tomato leaf." BMC
Plant Biology **19**(1): 377.
Zimmerman, D. C. and C. A. Coudron (1979). "Identification of traumatin, a wound hormone, as 12-oxo-
trans-10-dodecenoic acid." Plant Physiology **63**(3): 536.

1386