Synaptic Mechanisms of Top-Down Control by The Auditory Cortico-Collicular Pathway

Hannah M. Oberle[1,2], Alex Ford[1], Jordyn Czarny[1], & Pierre F. Apostolides[1,3]*
1) Kresge Hearing Research Institute & Department of Otolaryngology, University of Michigan
Medical School
2) Neuroscience Graduate Program, University of Michigan
3) Molecular and Integrative Physiology, University of Michigan Medical School
*corresponding author: piaposto@med.umich.edu

Abstract

Abstract
Corticofugal projections to evolutionarily ancient, sub-cortical structures are ubiquitous across
mammalian sensory systems. These "descending" pathways enable the neocortex to control
ascending sensory representations in a predictive or feedback manner, but the underlying cellular
mechanisms are poorly understood. Here we combine optogenetic approaches with *in vivo* and *in*
vitro patch-clamp electrophysiology to study the projection from auditory cortex to the inferior
colliculus (IC), a major descending auditory pathway that controls IC neuron feature selectivity,
plasticity and auditory perceptual learning. Although individual auditory cortico-collicular synapses
were generally weak, IC neurons often integrated inputs from multiple corticofugal axons that
generated reliable, tonic depolarizations even during prolonged presynaptic activity. Latency
measurements *in vivo* showed that descending signals reach the IC within 30 ms of sound onset,
which in IC neurons corresponded to the peak of synaptic depolarizations evoked by short sounds.
Activating ascending and descending pathways at latencies expected *in vivo* caused a NMDA
receptor dependent, supra-linear EPSP summation, indicating that descending signals can non-
linearly amplify IC neurons' moment-to-moment acoustic responses. Our results shed light upon the
synaptic bases of descending sensory control, and imply that heterosynaptic cooperativity
contributes to the auditory cortico-collicular pathway's role in plasticity and perceptual learning.

Introduction

The auditory system is organized as a network of feedback loops, such that most central auditory
nuclei receive descending projections from higher levels of the processing hierarchy (Diamond et
al., 1969; Saldaña et al., 1996; Winer et al., 1998, 2001; Doucet et al., 2003; Coomes and
Schofield, 2004; Schofield et al., 2006; Suthakar and Ryugo, 2017). The auditory cortex is a major
source of excitatory (glutamatergic) descending projections, with the density of descending fibers
often rivaling that of ascending fiber tracts (Winer et al., 2001; Winer, 2006; Stebbings et al., 2014).
These corticofugal projections likely play a major role in hearing by providing an anatomical
substrate for "top-down" information to control early acoustic processing. Indeed, stimulating or
silencing the auditory cortex *in vivo* changes spontaneous and sound-evoked activity throughout
the central auditory system (Massopust and Ordy, 1962; Ryugo and Weinberger, 1976; Yan and
Suga, 1998, 1999; Xiao and Suga, 2002; Yu et al., 2004; Nakamoto et al., 2008, 2010; Anderson
and Malmierca, 2013; Kong et al., 2014; Vila et al., 2019; Blackwell et al., 2020; Qi et al., 2020),
indicating that high-level activity regulates the moment-to-moment function of sub-cortical auditory
circuits. However, little is known regarding the biophysical properties of auditory corticofugal
synapses, nor do we understand how descending signals are integrated with ascending
information. Given that synaptic dynamics and pathway integration are fundamental building blocks
of neural circuit computations (Abbott et al., 1997; Zucker and Regehr, 2002; Stuart and Spruston,
2015), addressing these knowledge gaps is necessary to understand how the auditory cortex
exerts control over early auditory processing.
48
Of particular interest is the descending projection from auditory cortex to the inferior colliculus (IC),
a midbrain hub important for sound localization, speech perception, and an early site of divergence
for primary and higher-order auditory pathways (Masterton et al., 1968; Krishna and Semple, 2000;
Champoux et al., 2007; Sinex and Li, 2007; Joswig et al., 2015). The IC is generally sub-divided
into a "lemniscal" central core and "non-lemniscal" dorsal and lateral shell regions whose neurons
have distinct afferent and efferent connections (Faye-Lund and Osen, 1985; Loftus et al., 2008;
Ayala et al., 2015; Chen et al., 2018). Whereas central IC neurons project mainly to the primary
auditory thalamus (ventral medial geniculate nucleus; Mellott et al., 2014; Oliver, 1984), shell IC
neurons preferentially project to secondary, higher-order auditory thalamic nuclei that subsequently
funnel acoustic information to the amygdala and striatum (Oliver and Hall, 1978; LeDoux et al.,
1990; Bordi and LeDoux, 1994; Mellott et al., 2014; Cai et al., 2019; Ponvert and Jaramillo, 2019).
Auditory cortico-collicular axons terminate primarily in the shell IC, with comparatively fewer fibers
in the central IC (Bajo et al., 2019; Chen et al., 2018; Lesicko et al., 2016; Song et al., 2018; Winer
et al., 1998; Xiong et al., 2015; but see Saldaña et al., 1996). Thus, auditory cortico-collicular
synapses seem uniquely positioned to modulate acoustic signals destined for limbic circuits
supporting learned valence and habit formation; this prediction is further supported by the fact that
chemical ablation of auditory cortico-collicular neurons selectively impairs certain forms of auditory
perceptual learning while sparing the performance of previously learned task associations (Bajo et
al., 2010). Nevertheless, little is known regarding how auditory cortico-collicular synapses control
activity in single IC neurons. Intriguingly, auditory cortex inactivation typically does not abolish IC
neuron sound responses, but rather causes divisive, non-monotonic changes in receptive field
properties and feature selectivity (Yan and Suga, 1999; Nakamoto et al., 2008, 2010; Anderson and

Malmierca, 2013). Thus, descending transmission might operate in part via heterosynaptic
interactions, perhaps by controlling how IC neurons respond to ascending acoustic inputs.
73
Here we employ electrophysiology and optogenetic approaches to identify how auditory cortico-
collicular synapses transmit descending signals, and to understand how descending synapses
control IC neuron responses to ascending inputs. We find that the majority of shell IC neurons
receive monosynaptic inputs from auditory cortex and often integrate information from multiple
distinct corticofugal axons. Synaptic latency measurements *in vivo* show that descending excitation
reaches IC neurons ~5-7 ms after spike initiation in auditory cortex, such that cortical feedback will
rapidly follow the onset of acoustically driven excitation. Somewhat surprisingly, NMDA receptors
only modestly contribute to descending transmission. By contrast, excitatory intra-collicular
synapses from the central IC, which are probably a major source of ascending acoustic signals to
shell IC neurons, had a much larger NMDA component. Consequently, appropriately timed activity
in ascending and descending pathways integrates supra-linearly owing to the cooperative activation
of NMDA receptors. Our data reveal a key role for heterosynaptic non-linearities in the descending
modulation of early acoustic processing. In addition, the results place important biophysical
constraints on the synaptic learning rules that might support the auditory cortico-collicular
pathway's role in experience-dependent plasticity and perceptual learning.
89
90 **Results**
91
Auditory cortico-collicular synapses robustly target superficial IC neurons
Corticofugal axons are predominantly restricted to the shell IC (Winer et al., 1998; Song et al.,
2018), but little is known about the extent of functional synaptic connectivity between auditory
cortex and IC neurons. We tested how auditory cortico-collicular synapses impact IC neurons by
transducing the optogenetic activator Chronos in the auditory cortex of mice via intracranial AAV
injections (Figure 1A), and 2-4 weeks later, performing *in vivo* whole-cell recordings from the
ipsilateral IC of urethane anesthetized mice. GFP-tagged, Chronos expressing auditory cortico-
collicular axons were primarily restricted to the ipsilateral dorsal-medial and lateral IC (Figure 1B),
in agreement with previous studies. Single flashes of blue light from an optic fiber positioned over
auditory cortex (1-5 ms duration; Figure 1C) reliably triggered excitatory postsynaptic potentials
(EPSPs) in n = 21/38 IC neurons recorded from N = 8 mice (Figure 1D). EPSPs were primarily
observed in superficial IC neurons, with the majority of unresponsive neurons located more
ventrally, presumably in the central IC (Figure 1E; mean depth of cortico recipient and non-recipient
neurons: 220 ± 23 vs. 354 ± 38 μm from surface, p=0.0036, Kolmogorov-Smirnov test). EPSPs had
an onset latency of 10.3 ± 0.7 ms following light stimulation (Figure 1F), a peak amplitude of 3.5 ±
0.7 mV (Figure 1G), and a full-width at half-maximum of 18.4 ± 1.5 ms (Figure 1H). Interestingly,
EPSP amplitudes varied over 2 orders of magnitude across different cells and were occasionally
large enough to drive IC neurons beyond spike threshold (Supplemental Figure 1A). These data
indicate that descending excitation can be quite potent, such that even brief and presumably sparse
activity of auditory cortico-collicular neurons could in principle drive IC efferent signals
independently of ascending inputs.
113

Figure 1: *Biophysical properties of auditory cortico-collicular synapses* **A)** Cartoon of experiment. **B)** Confocal stack of GFP-Chronos labeled axons mainly restricted to the shell IC. **C)** *In vivo* whole-cell recordings are obtained from IC neurons 2-4 weeks following Chronos injections; an optic fiber is positioned above the auditory cortex **D)** Example EPSPs following *in vivo* optogenetic stimulation. Gray traces are single trials; magenta is average. Inset is the EPSP rising phase at a faster timebase. **E)** Dorsal-ventral locations (relative to dura) for IC neurons where auditory cortical stimulation did (magenta) and did not (gray) evoke an EPSP. **F-H)** Summary of EPSP onset (F) amplitude (G), and half-width (H). **I)** whole-cell recordings obtained from dorso-medial shell IC neurons *in vitro*. **J)** EPSPs evoked by *in vitro* optogenetic stimulation (2 ms light flash). Inset: EPSP rising phase. **K-M)** Histograms of EPSP onset (K), amplitudes (L), and half-widths (M) *in vitro*.

Because the auditory cortex projects to many sub-cortical targets besides the IC, *in vivo* stimulation
could drive polysynaptic excitation onto IC neurons that would complicate estimates of
monosynaptic connectivity. We thus prepared acute IC brain slices from mice injected with Chronos
in auditory cortex to quantify the functional properties of descending synapses in a more controlled
setting. We targeted whole-cell current-clamp recordings specifically to neurons in the dorso-medial
shell IC, as this region shows the highest density of corticofugal axons (Song et al., 2018; our
Figure 1B). Stimulating auditory cortico-collicular axons via single blue light flashes delivered
through the microscope objective (1-10 ms duration) drove EPSPs in n = 78 neurons from N = 40
mice (Figure 1I,J). EPSPs had short-latency onsets following photo-stimulation (Figure 1K; 3.0 ±
0.1 ms), indicating a monosynaptic rather than polysynaptic origin. EPSPs *in vitro* had a similar
range and mean peak amplitude as those recorded *in vivo* (Figure 1L; 2.97 ± 0.35 mV, p=0.3,
Mann-Whitney test), and similarly could drive spikes in a subset of recordings (Supplemental Figure
1B). Although the EPSP half-width was significantly slower *in vitro* compared to *in vivo* (Figure 1M;
39.6±2.3 ms, p<0.001, rank-sum test), this result is not surprising: The constant barrage of synaptic
inputs *in vivo* is expected to generate a "high conductance state" that accelerates the membrane
time constant (Destexhe et al., 2003).
130
In a separate set of experiments (n = 11 neurons from N = 6 mice), we quantified the kinetics of
auditory cortical excitatory postsynaptic currents (EPSCs) as they appear at the soma using
voltage-clamp recordings with a Cs^+ based internal solution (Supplemental Figure 2. Peak
amplitude = 60.1 ± 14.4 pA; weighted decay time constant = 4.5 ± 0.5 ms; 10-90% rise time of 1.2 ±
0.2 ms). Altogether, our results show that auditory cortico-collicular synapses substantially
depolarize shell IC neurons independent of network-level, polysynaptic activity. In addition, EPSP
kinetics are such that descending excitation will undergo significant temporal summation at firing
rates observed during sound-evoked activity of auditory cortico-collicular neurons (20-50 Hz;
Williamson and Polley, 2019).
140
Synaptic strength reflects presynaptic convergence rather than unitary EPSP amplitudes
EPSP amplitudes spanned two orders of magnitude under our conditions (Figure 1G,L). Does this
variability reflect a differential potency of individual synapses, or alternatively, differences in the
number of presynaptic auditory cortical axons impinging onto individual shell IC neurons? We first
estimated unitary EPSP amplitudes using a minimal stimulation paradigm designed to activate one
(or very few) auditory cortico-collicular fibers. In these experiments, the LED intensity was titrated to
the minimum power required for optogenetic responses to fluctuate between successful EPSPs and
failures on a trial-by-trial basis (Figure 2A,B; mean failure rate across experiments: 44 ± 3%, n = 18
cells from N = 14 mice). The mean amplitude of successful EPSPs was generally small (0.84 ± 0.09
mV; Figure 2C) and similar to previous reports of unitary synapses between layer 5 pyramidal
neurons in sensory cortex (Brown and Hestrin, 2009; Lefort et al., 2009). By contrast, progressively
stronger LED flashes increased the amplitude of successful EPSPs compared to the minimal
stimulation condition in many cells tested (n=15 cells from N = 11 mice; Figure 2D,E), indicating
that multiple corticofugal axons can converge onto individual shell IC neurons. The ratio of
maximum to minimum EPSP showing varied >10-fold across different experiments (median: 1.67,
range: 0.91-24.04, Figure 2F), indicating that the EPSP amplitude variability across individual
neurons likely reflects the number of presynaptic auditory cortical fibers recruited during stimulation

rather than differences in unitary strength. Importantly, the EPSP half-width was constant across
the range of stimulus intensities (Figure 2D, inset. EPSP halfwidth ratios at maximal and minimal
LED intensities: 1.04 ±0.07). This result indicates that increased LED intensities recruit more axons
rather than prolonging Chronos activation and temporally dispersing vesicle release from single
presynaptic boutons. We conclude that although the strength of individual auditory cortico-collicular
synapses is weak, the convergence of multiple presynaptic fibers ensures that descending signals
will substantially increase shell IC neuron excitability.

Figure 2: *Multiple auditory cortico-collicular fibers converge onto single shell IC neurons*. **A)** Example threshold optogenetic stimulation experiment. Thin black and gray traces are successes failures, respectively. Thick traces are averages. **B)** Diary plot from the experiment in panel A. black and gray symbols are successes and failures, respectively. **C)** Summary of putative unitary EPSP amplitudes **D)** Magenta: putative unitary EPSP (average of successes recorded at threshold stimulation). Black and Gray traces: averages of successes at increasing LED intensity. Data are from a different neuron as in A. Inset: Peak scaled and onset aligned EPSPs recorded at threshold and maximal intensity, revealing an identical time-course. **E)** EPSP peak amplitude is plotted as a function of LED intensity for the recording in (D). Magenta line is Hill fit. **F)** EPSP amplitude ratios at maximal and threshold LED intensities. Most values are >1, indicating convergence of at least 2 fibers. Of note, data are on a log scale.

Auditory cortico-collicular synapses contact shell IC neurons with diverse biophysical properties
IC neurons have a variety of firing patterns and membrane properties that potentially correspond to
distinct neuronal subtypes (Smith, 1992; Li et al., 1998; Peruzzi et al., 2000; Sivaramakrishnan and
Oliver, 2001; Ahuja and Wu, 2007; Sun and Wu, 2008; Moore and Trussell, 2017; Goyer et al.,
2019; Naumov et al., 2019; Silveira et al., 2020). However, whether auditory cortico-collicular
synapses contact biophysically homogenous or diverse neurons is unknown. We injected 300 ms
positive and negative current steps to quantify spiking patterns and passive membrane properties
in a subset of our *in vitro* experiments described above (n = 64 cells). 62/64 cortico-recipient shell
IC neurons could be classified into one of four general categories. Over half of neurons (38/64) had
significant spike rate adaptation during positive current injections (Figure 3A,B) and qualitatively
similar firing patterns as the shell IC neurons recorded in rat slices by Smith (1992). However, 18 of
these "adapting" neurons responded to negative current with a sustained hyperpolarization (Figure
3A), whereas 20 displayed a prominent I_h-like "sag" likely mediated by HCN channels (Figure 3B,

middle trace). These data suggest a minimum of two shell IC neuron sub-types with adapting firing
patterns, which can be differentiated based on the extent of I_h sag. By contrast, other neurons had
delayed first spikes (9/64, Figure 3C) or showed non-adapting discharge patterns (15/64, Figure
3D). Finally, 2/64 cells displayed a strikingly distinct phenotype, with a burst of spikes riding atop a
"hump"-like depolarization similar to neurons expressing T-type Ca^{2+} channels (Figure 3E). EPSP
amplitude and half-width of were similar across the four major categories (Peak amplitude: p=0.45;
half-width: p=0.5, Kruskal-Wallis tests) despite marked differences in electrical properties. Thus, the
auditory cortex broadcasts similarly strong signals to multiple, putatively distinct IC neurons,
arguing that biophysical properties alone do not predict the strength of descending synapses.

Figure 3: *Shell IC neurons receiving descending signals have diverse biophysical properties.* **A-E)** Examples of different shell IC neurons. Top and Middle traces are spike output and membrane hyperpolarization following positive and negative current steps, respectively. Lower traces: average auditory cortical EPSPs from the same neuron. **F, G)** EPSP peak amplitude (F) and half-width (G) as a function of biophysical category. Bursting neurons are omitted due to low n.

Temporal integration of auditory cortical inputs is moderately sub-linear
Auditory cortico-collicular neurons in awake mice respond to acoustic stimuli with ~10-50 Hz spike
trains (Williamson and Polley, 2019). Given the kinetics of auditory cortico-collicular EPSPs (Figure
1), these spike rates are expected to result in significant temporal summation of descending
signals. However, the use-dependent dynamics neurotransmitter release, as well as postsynaptic
ion channels, can enforce sub- or supra-linear summation that effectively dictates the temporal
integration of EPSPs (Markram et al., 1998; Pouille and Scanziani, 2004; Polsky et al., 2009). How
do IC neurons integrate sustained cortical activity? We addressed this question by repetitively
stimulating auditory cortico-collicular axons (10 light pulses at 20 or 50 Hz, 2 ms pulse width; Figure
4A,B, black traces; n = 12 neurons from N = 7 mice). We quantified temporal integration by
comparing the peak EPSP amplitudes observed after each light flash in the train to the amplitudes
expected from the linear summation of a single auditory cortico-collicular EPSP recorded in the
same neuron (Figure 4A,B; magenta traces). The observed peak amplitude of the 10th EPSP in the
train reached 92 ± 13 % and 77 ± 16% of that expected from linear summation at 20 and 50 Hz,
respectively (Figure 4C,D). These results argue that at the population-level, auditory cortical firing
rates are read out as moderately sub-linear shifts of the membrane potential towards threshold.
203
Auditory cortico-collicular neurons display elevated firing rates at ~20 Hz for the entirety of long-
duration, complex sounds (e.g., 4 s long dynamic chord stimuli; Williamson and Polley, 2019). We
thus wondered if descending synapses maintain transmission during sustained acoustic
processing, or if synaptic depression instead limits descending signals to sound onset. Stimulating
auditory cortical axons with 4 s trains of light pulses at 20 Hz (Figure 4E) drove fast EPSPs riding
atop a steady-state, tonic depolarization (Figure 4F; mean amplitude of DC component during the
final 1 s of stimulation: 1.18 ± 0.01 mV, p = 0.013, one-sample t-test compared to a hypothetical
value of 0. n = 10 cells from n = 7 mice). Importantly, these effects were not an artifact of directly
stimulating auditory cortico-collicular nerve terminals in brain slices: A similar tonic depolarization
was observed in superficial IC neurons recorded *in vivo* using an optic fiber positioned over
auditory cortex (Figure 4G,H; mean amplitude of DC component: 1.5 ± 0.7 mV, n=19 cells from n =
6 mice, p=0.53 compared to *in vitro* data, rank-sum test).
216
In addition, cessation of cortical stimulation *in vivo* also caused a long-lasting after-
hyperpolarization (AHP) in 14/19 cells: The membrane potential rapidly fell below baseline after the
last stimulus (τ = 76.5 ± 1.5 ms) and recovered over several seconds (τ = 2.6 ± 0.6 s; Supplemental
Figure 3). Interestingly, this AHP was independent of postsynaptic spiking and thus may reflect
buildup of feed-forward inhibition from local and long-range sources, or alternatively, a transient
cessation of tonic descending excitation. Altogether, these experiments show that auditory cortico-
collicular synapses can sustain transmission on seconds time scales via tonic and phasic
excitation. In addition, the profound AHP following *in vivo* stimulation suggests that IC neuron
excitability is bi-directionally yoked to auditory cortical firing patterns. Thus, increases as well as
pauses in auditory cortico-collicular neuron activity may be of comparable significance to IC
neurons.

Figure 4: *Repetitive synaptic activity tonically depolarizes shell IC neurons despite sub-linear temporal summation of EPSPs.* **A,B)** Black traces: Average EPSPs evoked by 20 (A) or 50 Hz (B) trains of 10 light flashes. Gray: Average EPSPs evoked by a single 2 ms light flash recorded in the same neuron. Magenta: The expected response assuming linear summation of single EPSPs at each frequency. Of note is that the recorded EPSP amplitudes in the train are smaller than the expected linear sum. Data in A and B are from the same neuron. **C,D)** Summary of observed over expected amplitudes for EPSPs during 20 (C) or 50 Hz (D) trains. Gray dotted lines are individual neurons; black + shading are mean ± SEM **E)** Sustained auditory cortical transmission generates phasic EPSPs (black) riding atop a tonic depolarization (magenta) *in vitro*. **F)** Group data quantifying membrane potential changes at steady-state (final 1 s of stimulation). **G,H)** Same as E and F but during *in vivo* recordings. Of note is the large AHP.

NMDA receptors contribute to temporal integration of descending signals
Central excitatory transmission is predominantly mediated by AMPA and NMDA type glutamate
receptors, with NMDA receptors being particularly crucial for dendritic integration (Larkum et al.,
2009; Branco et al., 2010), associative plasticity (Kerchner and Nicoll, 2008), and learning (Morris
et al., 1986). The auditory cortico-collicular pathway is involved in perceptual learning following
monaural hearing loss (Bajo et al., 2010), and NMDA receptors in the avian IC shell homologue
preferentially contribute to receptive fields generated by experience-dependent plasticity (Feldman
et al., 1996). We thus asked to what extent auditory cortico-collicular synapses activate NMDA
receptors in shell IC neurons. Interestingly, bath application of the AMPA/kainate receptor
antagonist NBQX (10 µM) completely abolished EPSPs in all neurons tested (Figure 5A,B. Peak
amplitude control: 1.93 ± 0.02 mV; NBQX: 0.09 ± 0.03 mV, p=0.004, n=9 cells from N = 7 mice,
Wilcoxon sign-rank test), indicating that AMPA receptors mediate the overwhelming majority of
synaptic charge at descending synapses.
241
By contrast, although the NMDA receptor antagonist R-CPP (5 µM) had minimal effect on the peak
amplitude of auditory cortico-collicular EPSPs evoked with single light flashes (Figure 5C,D; n = 9
cells from N = 6 mice. Control: 3.3 ± 0.9 mV, R-CPP: 3.4 ± 0.9 mV, p = 0.67, paired t-test), R-CPP
caused a modest but significant reduction in the EPSP half-width (Figure 5C,E; control: 35.7 ± 4.4
ms, R-CPP: 30.4 ± 4.7 ms. 17.6% reduction, p=0.016, paired t-test). In addition, depolarizing
neurons to +30 to +40 mV in voltage-clamp (using a Cs^+-rich internal solution) slowed the weighted
decay time constant of optogenetically evoked EPSCs by 8.2 fold compared to negative holding
potentials (Supplemental Figure 4; -60 to -70 mV: 6.1 ± 0.9 ms, +30 to +40 mV: 50.0 ±0.8 ms, n = 5
cells from N = 3 mice), consistent with the biophysical properties of NMDA receptors. We conclude
that although AMPA receptors mediate the bulk of descending transmission at resting membrane
potentials, glutamate released from descending synapses nevertheless reaches synaptic NMDA
receptors to shape EPSP kinetics. Thus, although NMDA receptors contribute little to the peak
amplitude of EPSPs during sparse stimulation, they may nevertheless control the integration of
repetitive activity across time.
256
Accordingly, R-CPP significantly reduced summation of auditory cortico-collicular EPSPs evoked by
a train of 10 stimuli at 50 Hz (Figure 5F, G; n=10 cells from N = 8 mice. Main effect of drug
condition in a two-way repeated measures ANOVA; $F(1,9)=5.91$, p=0.038), resulting in a 26%
reduction in the cumulative integral of the synaptic depolarization (Figure 5F,H; control: 1.42 ± 0.35
mV * s, R-CPP: 1.03 ± 0.26 mV * s; p=0.012, paired t-test). These data show that NMDA receptors
prolong the integration time window for descending signals, thereby boosting postsynaptic
depolarizations during sustained cortical activity.

Figure 5. *Pharmacology of descending transmission.* **A)** EPSPs (mean ± SEM) from a single neuron before (black) and after (magenta) bath application of NBQX (10 µM). Of note is the absence of residual synaptic depolarization after NBQX. **B)** Summary data. For the effect of NBQX on descending EPSPs. **C)** EPSPs before and after R-CPP (black and magenta traces, respectively). **D, E)** Group data for the effect of R-CPP on EPSP peak amplitude (D) and half-width (E). Asterisks denote statistical significance. **F)** Upper panel: EPSPs (average ± SEM) evoked by 10 light flashes at 50 Hz. Black and magenta are in control and R-CPP. Lower panel is the cumulative integral of the waveforms. **G)** Group data showing amplitude of each EPSP in the train (mean ± SEM) before and after R-CPP (black and magenta, respectively). **H)** Group data for the effect of R-CPP on voltage integral as in F. Asterisks denote significance.

Differential contribution of NMDA receptors at ascending and descending synapses
The modest contribution of NMDA receptors at descending synapses is surprising, as previous
studies suggest a rather prominent NMDA component at excitatory synapses in the IC (Smith,
1992; Wu, 2004; Kitagawa and Sakaba, 2019). We thus wondered if our results reflected a relative
paucity of NMDA receptors at all excitatory synapses onto shell IC neurons, or rather a unique
feature of auditory cortico-collicular synapses. Shell IC neurons receive a prominent intra-collicular
projection from the central IC which likely transmits a significant amount of ascending acoustic
information (Saldaña and Merchán, 1992; Saldaña et al., 1996; Sun and Wu, 2009). We tested if
the NMDA component differed between ascending and descending EPSPs in the same neuron
using a dual pathway stimulation approach *in vitro*: A bipolar stimulating electrode was positioned in
the central IC and Chronos was expressed in auditory cortex to activate ascending and descending
synapses, respectively (Figure 6A). 2.5-5 µM gabazine was present in all experiments to isolate
excitatory transmission. Repetitive stimulation of either pathway (5x, 50 Hz) led to summating
EPSPs which were differentially sensitive to NMDA receptor blockade (Figure 6B-C): 5 µM R-CPP
reduced the cumulative integral of central IC and auditory cortical EPSPs to 44±4% and 78±4% of
baseline, respectively (Figure 6D, n=9 cells from n = 8 mice, p=0.0004, paired t-test), indicating that
NMDA receptors contributed less at descending compared to ascending synapses. Thus, the
extent of NMDA receptor contribution to descending transmission reflects a synapse-specific
property of auditory cortico-collicular inputs rather than the global distribution of NMDARs in shell
IC neurons.

Figure 6. *Differential contribution of NMDA receptors at ascending and descending synapses.* **A)**
Cartoon of experiment: Blue light flashes delivered through the microscope objective activate
descending auditory cortico-collicular axons; a bipolar electrode in the central IC stimulates
ascending central ->shell IC axons. **B,C)** EPSPs evoked by 5x 50 Hz stimulation of central IC (B) or
auditory cortex (C) axons, in absence or presence of R-CPP (black and magenta, respectively). Data
in panels B and C are from the same neuron. **D)** Group data quantifying the fractional EPSP
remaining in R-CPP (as measured by the cumulative integral of the EPSP waveform) for central IC
and auditory cortico-collicular synapses.

Auditory cortical inputs are predicted to arrive at the peak of EPSPs evoked by transient sounds
Layer 5 auditory cortico-collicular neurons in awake mice respond to sound with a mean first-spike
latency of ~21 ms (Williamson and Polley, 2019). This value is surprisingly shorter than reported
first spike latencies in shell IC neurons (~35 ms; Lumani and Zhang, 2010). Whether auditory
cortical excitation arrives before, during, or after the onset of sound-evoked EPSPs in shell IC
neurons effectively determines how ascending and descending signals integrate at the single cell

level, but the relative timing of distinct inputs onto IC neurons is unknown. We thus quantified the
relative timing of ascending sound-evoked and descending cortical EPSPs using *in vivo* whole-cell
recordings from superficial IC neurons of anesthetized mice. We first determined the onset latency
of descending EPSPs in IC neurons using electrical stimulation of the auditory cortex (Figure 7A).
We employed electrical, rather than optogenetic stimulation for these experiments because spike
onset following optogenetic stimulation is limited by the cell's membrane time constant and effective
spike threshold, whereas electrical stimulation bypasses somato-dendritic depolarization by directly
triggering axonal spikes. Single shocks delivered to the auditory cortex evoked EPSPs with an
onset latency of 5.4 ± 0.6 ms (n = 7 cells from n = 4 mice; mean depth of recorded neurons 188 ±
14 μm, Figure 7B,C), indicating that descending information reaches IC neurons within a few ms of
AP initiation in auditory cortex. Since the mean first-spike latency of auditory cortico-collicular
neurons is ~21 ms (Williamson & Polley, 2019) and the synaptic latency is 5-8 ms (Figure 7B,C),
these data collectively argue that that cortical feedback begins to excite IC neurons <30 ms after
sound onset. Furthermore, assuming an axon path length of ~8 mm from auditory cortex to shell IC
(Llano et al., 2014) and a synaptic release delay of ~2 ms, the data suggest a minimum conduction
velocity of ~2.35 m/s. These values are similar to conduction velocity estimates for the myelinated
axons of layer 5 pyramidal neurons in rodents (2.9 m/s; Kole et al., 2007), suggesting that layer 5,
and not unmyelinated layer 6 neurons, are the dominant source of descending signals to the IC.

Figure 7: *Auditory Cortical feedback is predicted to collide with the peak of ascending excitation.* **A**) Cartoon of experiment. **B)** Example recording showing short latency EPSPs following a single shock to the auditory cortex. Dotted line is onset of stimulation. Gray traces are single trials; black is average. The stimulation artifact, as well as APs riding atop the cortical EPSP, were blanked for clarity. **C)** Group data for EPSP onset latency. **D)** EPSPs evoked by a 200 μs click. Arrow and speaker show sound onset. Data are from a different neuron than in B. Black and magenta bars above the traces show the reported first spike latency of auditory cortico-collicular neurons from previous work and the synaptic latency calculated from panels B and C, respectively. **E)** Group data for onset and peak latency of click-evoked EPSPs.

We next calculated the timing of sound-evoked excitation onto superficial IC neurons using
broadband "click" transients (Figure 7D, n = 10 cells from N = 7 mice. Mean depth: 204±23 μm).
Although sound-evoked EPSPs were typically sub-threshold under our conditions, click stimuli
occasionally drove spikes in 5/10 experiments (33.8 ± 16.6% of trials), with mean first spike
latencies of 29.4±7.6 ms, similar to previous reports (Lumani and Zhang, 2010; Geis et al., 2011).
Sound evoked EPSPs had mean onset and peak latencies of 15.7 ± 1.7 and 26.5 ± 3.9 ms,
respectively (Figure 7E), indicating that cortical feedback lags the rising phase of sound-evoked
excitation by a mere 10-15 ms. Importantly, the data further suggest that cortical excitation will
arrive prior to, or in very close succession with, the peak of EPSPs evoked by transient sounds,
such that cortical feedback could in principle dictate the first spike latencies of shell IC neurons.
318
NMDA receptor dependent, supra-linear pathway integration in shell IC neurons
Our latency measurements (Figure 7) suggest that ascending information is rapidly followed by
descending cortical excitation. This temporal overlap is intriguing because ascending and
descending synapses express NMDA receptors (Figure 6), which in other cell-types, enable
cooperative interactions between co-active pathways onto the same neuron (Takahashi and
Magee, 2009; Harnett et al., 2012). We thus hypothesized that appropriately timed cortical
feedback might integrate non-linearly with ascending inputs from the central IC, thereby generating
a synaptic depolarization larger than expected from the sum of either pathway active in isolation.
We tested this idea *in vitro* using our dual pathway stimulation approach (Figure 6) while recording
from shell IC neurons. We first recorded the synaptic depolarization following stimulation of
ascending and descending pathways in isolation (5 stimuli at 50 Hz; Figure 8A, upper traces).

Figure 8: *NMDA-receptor dependent, supralinear pathway integration.* **A)** Upper and lower traces are during baseline conditions and after bath applying R-CPP. Left and middle panels: EPSPs during electrical or optogenetic stimulation of central IC or auditory cortical axons, respectively (5x, 50 Hz stimuli). Right panel ("Combined"): Black traces are the observed depolarization during simultaneous stimulation of both pathways. Magenta is the depolarization expected from the arithmetic sum of the waveforms following stimulation of either pathway alone (e.g., left and middle traces). Of note, the observed depolarization under control conditions is larger than expected from linear summation; blocking NMDA receptors linearizes pathway integration (compare black and magenta traces in R-CPP). **B)** Group data plotting the ratio of observed and expected peak amplitude for each of 5 EPSPs in a 50 Hz train during synchronous activation of central IC and auditory cortical synapses. Asterisks denote statistical significance of Bonferroni post-hoc test for the fourth and fifth stimuli following a main effect of drug condition (p=0.038, F(1,9), Two-way repeated measures ANOVA). **C)** Group data, observed over expected ratio of the cumulative integral during combined pathway activation in control conditions and in the presence of R-CPP. Asterisks denote statistical significance (sign-rank test).

We next simultaneously activated the two pathways such that the onset of cortical EPSPs collided
with the peak of ascending EPSPs, as predicted from our *in vivo* latency measurements (Figure 7).
The observed depolarization during synchronous pathway activation was on average significantly
larger than expected from the arithmetic sum of each pathway stimulated alone (Figure 8A; n=12
cells from N=9 mice. Cumulative integral observed: 1.90 ± 0.35 mV*ms, expected: 1.45±0.22
mV*ms, p=0.007, paired t-test test), indicating that coincident activity of ascending and descending
pathways summates supra-linearly. Importantly, R-CPP mostly abolished this supra-linear
summation, such that the depolarization during coincident activation in R-CPP now equaled the
expected sum of each pathway activated alone (Figure 8B,C, lower traces; n=7 cells from N = 7
mice. Observed/Expected control: 1.39±0.05; in R-CPP: 1.02±0.04, p=0.0156, sign-rank test).
Thus, synaptic NMDA receptors provide a supra-linear boost when descending excitation follows
ascending activity, thereby promoting the non-linear mixing of distinct pathways in single IC
neurons.
344
345
346 **Discussion**
347
We have shown that the majority of neurons in the superficial (shell) IC layers receive reasonably
strong excitation from auditory cortex that is predicted to arrive ~30 ms following sound onset.
These synaptic properties could enable a context-dependent modulation of IC neurons across
multiple timescales. Indeed, the rapid onset of descending EPSPs following cortical spikes (~5-8
ms) is more than twice as fast as sensory-evoked cortical gamma rhythms (30-50 Hz). Thus,
descending signals could effectively synchronize neural ensembles across the ascending auditory
hierarchy either to the temporal envelope of sound (Weible et al., 2020) or to internally generated
rhythms. Furthermore, descending synapses sustained transmission and drove tonic
depolarizations even during seconds-long activity patterns, such that IC neurons may also integrate
slower cortical state fluctuations. Intriguingly, auditory cortical neurons in behaving animals show
enhanced firing rates during the delay period of auditory working memory tasks (Gottlieb et al.,
1989) which apparently, precedes similar activity patterns in prefrontal cortex (Huang et al., 2016).
If these working memory related neuronal ensembles include auditory cortico-collicular neurons,
sustained transmission from descending synapses could cause seconds-long increases in IC
neuron excitability based on working memory content. Accordingly, persistent delay period activity
is observed in ~10% of IC neurons when rats engaged in an auditory working memory task
(Sakurai, 1990), although future studies are necessary to determine the extent to which this activity
is inherited from descending auditory cortical pathways.
366
Finally, although our data suggest that shell IC neurons are the major target of descending fibers,
the non-uniform thickness of the IC shell across medial-lateral axis (Barnstedt et al., 2015)
indicates we cannot rule out that some of our *in vivo* data are from neurons in the most dorsal
region of the central IC. Indeed, *in vivo* intracellular recordings from neurons in the deep IC layers
report EPSPs following auditory cortex stimulation (Mitani et al., 1983; Qi et al., 2020), implying that
some functional cortical synapses may in fact target central IC neurons. However, *in vitro* circuit
mapping experiments imply that monosynaptic connections between auditory cortex and central IC

are likely very rare (Xiong et al., 2015; Song et al., 2018), such that shell IC neurons likely process
the bulk of descending signals.
376
Synapse-specific contribution of NMDA receptors in shell IC neurons
The minor contribution of NMDA receptors to auditory cortico-collicular transmission is somewhat
surprising, as synaptic NMDA receptors are activated even at hyperpolarized membrane potentials
in central IC neurons (Ma et al., 2002; Wu, 2004; Goyer et al., 2019; Kitagawa and Sakaba, 2019).
However, our results do not simply reflect a global paucity of NMDA receptors at excitatory
synapses onto dorso-medial shell IC neurons, but rather can be explained by a pathway specific
contribution of specific glutamate receptor sub-types to the synaptic depolarization: NMDA receptor
blockade reduced EPSPs to a greater extent at central IC -> shell IC compared to auditory cortico-
collicular synapses. Although a simple explanation is that the total number of synaptic NMDA
receptors in shell IC neurons differs in a pathway specific manner, we cannot exclude differences in
NMDA receptor subunit composition (Schwartz et al., 2012), glutamate diffusion (Arnth-Jensen et
al., 2002), and synapse location (Branco et al., 2010; Song et al., 2018) as contributing factors.
389
Non-linear integration of ascending and descending signals
Intra-collicular synapses originating from the central IC likely provide a significant amount of
ascending acoustic input to shell IC neurons. Indeed, reported first spikes latencies of central IC
neurons typically lead the onset of sound evoked EPSPs in shell IC neurons (Syka et al., 2000;
Hurley and Pollak, 2005), and central IC neurons send tonotopically organized axonal projections to
the IC shell that conspicuously mirror the tonotopic distribution of best frequencies in the shell IC
(Saldaña and Merchán, 1992; Wong and Borst, 2019). By contrast, the onset of sound-evoked
EPSPs in shell IC neurons typically begin prior to the reported first spikes of auditory cortico-
collicular neurons, thereby ruling out the possibility that acoustic responses are solely inherited
from auditory cortex. Instead, axonal conduction velocities along ascending and descending
pathways impose an obligatory delay such that cortical feedback excitation arrives ~25-30 ms
following the onset of sound-evoked EPSPs. Notably, while glutamate released from ascending
terminals will have unbound and diffused away from low-affinity synaptic AMPA receptors prior to
the onset of cortical feedback excitation (Clements et al., 1992), high-affinity NMDA receptors are
expected to remain bound with glutamate during this time (Lester and Jahr, 1992). Extracellular
Mg^{2+} imparts a voltage-dependence to the NMDA receptor channel; the additional depolarization
provided by cortical feedback would thus be expected to cooperatively enhance current flowing
through NMDA receptors at ascending synapses active immediately prior to the onset of cortical
feedback. This prediction is supported by our data showing a NMDA receptor-dependent, supra-
linear summation of ascending and descending inputs that are activated similar to their expected
timing *in vivo*. These single-cell, biophysical operations can potentially explain the non-linear
changes in IC neuron receptive fields during cortical inactivation (Yan and Suga, 1999; Nakamoto
et al., 2008, 2010).
413
NMDA receptor dependent non-linearities are typically thought of as unique features of cortical
pyramidal neurons that support to the computational power of these high-level microcircuits.
Nevertheless, multiplicative integration of sound localization cues has been observed in single
neurons of barn owl IC (Peña and Konishi, 2001; 2002), and the NMDA receptor dependent non-

linearity we observe is comparable in magnitude to that reported during clustered activation of
neighboring synapses in CA1 neurons (Harnett et al., 2012). Together, these data suggest that
cooperative interactions between temporally correlated inputs may be a common neuronal
operation throughout the central nervous system. An important distinction however is that most, if
not all excitatory synapses in pyramidal neurons reside on dendritic spines; this
compartmentalization greatly limits any cooperative interactions to neighboring synaptic inputs on
the same branch (Gasparini and Magee, 2006; Losonczy and Magee, 2006). By contrast, auditory
cortico-collicular axons often form large (~5 μm^3) synapses on the soma of dorsal IC neurons
(Song et al., 2018). Depending on the impedance mismatch between the somatic and dendritic
compartments, synaptic depolarizations at the soma could propagate passively throughout the
neuron's multiple dendrites, thereby enabling cortical signals to non-linearly control ascending
information irrespective of the spatial relationship of co-active inputs. However, further studies are
necessary to identify the precise anatomical relationship between ascending and descending
synapses in single IC neurons.
432
Implications for experience-dependent plasticity and perceptual learning
Several studies now show that layer 5 corticofugal pyramidal neurons are necessary for perceptual
learning in multiple different sensory tasks. Optogenetic inhibition of layer 5 pyramidal neurons in
somatosensory cortex prevents behavioral adaptation following cue-related changes in a tactile
detection task, although the same manipulation had no effect on touch perception (Ranganathan et
al., 2018). Similarly, lesioning visual corticostriatal neurons prevents acquisition, but not
performance of a visual detection task (Ruediger and Scanziani, 2020). In the auditory system,
chemical lesions of auditory cortico-collicular neurons prevent the experience-dependent recovery
of sound localization following monaural hearing loss (Bajo et al., 2010), although auditory cortex
becomes dispensable for sound-localization once animals have learned to localize sounds using
monaural cues (Bajo et al., 2019). Thus, although necessary for perceptual learning, corticofugal
synapses may not be the primary locus of experience-dependent plasticity. Indeed, classic studies
in barn owls suggest that ascending central IC -> external (shell) IC synapses are the first site of
experience-dependent, spatial map plasticity in the auditory system (Brainard and Knudsen, 1993).
In tandem with our current study, these results suggest that auditory cortico-collicular synapses'
contributions to perceptual learning may not lie in their explicit ability to undergo classical Hebbian
associative plasticity, but rather as permissive forces of heterosynaptic plasticity at ascending
synapses.
451
452
453   **Methods**
454
Surgery for viral injections: All experiments were approved by the University of Michigan's IACUC
and performed in accordance with NIH's Guide for the care and use of laboratory animals. All
surgical procedures were performed under aseptic conditions. Surgeries were performed on 4-7
week old male or female C57BL6/J mice purchased from Jackson Labs or offspring of CBA x
C57BL6/J matings bred in house. Mice were deeply anesthetized with 4-5% isoflurane vaporized in
O_2 and mounted in a rotating stereotaxic frame (model 1430, David Kopf Instruments). Isoflurane
was subsequently lowered to 1-2% to maintain a deep anesthetic plane, as assessed by the

absence of paw withdrawal reflex and stable respiration (1-1.5 breaths/s). Body temperature was
maintained near 37-38° C using a feedback controlled, homeothermic heating blanket (Harvard
Apparatus). Mice were administered 5 mg/kg carprofen after induction as a pre-surgical analgesic.
The scalp was clear of hair, swabbed with betadine, and a small incision was made in the skin
overlying the left hemisphere. Topical 2% lidocaine was then applied to the wound margins. The
stereotaxic frame was rotated ~50 degrees, allowing a vertical approach perpendicular to the layers
of auditory cortex. A 200-400 µm craniotomy was carefully opened over the left auditory cortex (-
2.75 mm from Bregma, centered on the lateral ridge) using a 0.5 mm diameter dental burr (19007-
05, Fine Science Tools) and Foredom microdrill. The skull was frequently irrigated with chilled
phosphate buffered saline (PBS) to prevent overheating during drilling. Following the craniotomy, a
glass pipette (0.1-0.2 mm diameter at the tip) containing the pAAV-Syn-Chronos-GFP virus
(Addgene #59170-AAV1) penetrated the auditory cortex at a rate of <10 µm/s using a motorized
micromanipulator. A total of 100-200 nL virus was injected at 2-4 sites 810 and 710 µm below the
pial surface (25-50 nL per site). Following injections, the pipette was maintained in place for an
additional 5 min before slowly retracting at a rate of <10 µm/s. At the end of the surgery, the
craniotomy was filled with bone wax, the skin was sutured, and the mouse was removed from the
stereotax. Immediately following surgery, mice were given an analgesic injection of buprenorphine
(0.03 mg/kg, s.c.) and allowed to recover on a heating pad before returning to their home cage. An
additional post-operative dose of carprofen was administered 24 hours following surgery.
481
In vivo electrophysiology: 2-4 weeks following viral injections, mice were deeply anesthetized with
isoflurane and mounted in a stereotax as described above. The skin overlying the skull was
removed, the left temporal muscle was retracted, the stereotax was rotated ~50 degrees, and a 2-
2.5 mm craniotomy was carefully opened over the left auditory cortex. For optogenetic stimulation
in Figure 1, the dura overlying the auditory cortex was left intact and a cranial window was
implanted over the exposed brain with cyanoacrylate glue and dental cement. For the electrical
stimulation experiments in Figure 7, a small slit was carefully made in the dura and the craniotomy
was subsequently sealed with silicone elastomer. The stereotaxic frame was returned to the
horizontal position and a custom titanium headbar was affixed to the skull with dental cement. A
300-500 µm craniotomy was opened over the left IC and filled with a silicone elastomer plug. The
mouse was then removed from the stereotax, anesthetized with urethane (1.5 g/kg, i.p.), and head-
fixed in a custom-made sound attenuation chamber. Body temperature during the experiment was
maintained at 37-38° C with a custom designed, feedback-controlled heating blanket. For
optogenetic stimulation, a 0.5 NA, 400 µm core optic fiber (Thorlabs M45L02) coupled to a 470 nm
LED (Thorlabs M470F3) was mounted on a micromanipulator and positioned <1 mm away from the
auditory cortex cranial window. For electrical stimulation experiments, the silicone plug over
auditory cortex was removed and a bipolar platinum-iridium electrode (FHC 30210) was carefully
inserted ~800 µm into auditory cortex at an angle roughly perpendicular to the cortical layers.
Electrical stimuli were delivered via a custom stimulus isolator designed in house. Sound clicks (0.2
ms duration) were presented at ~91 dB peak equivalent SPL via a free-field speaker (Peerless
XT25SC90-04) positioned ~10 cm from the mouse's right ear. For whole-cell recordings, the
silicone plug over the IC was removed and patch-clamp recordings were obtained from IC neurons
via the "blind patch" approach using pipettes filled with K^+-rich internal solution containing (in mM):
115 K-Gluconate, 4 KCl, 0.1 EGTA, 10 HEPES, 14 Tris-Phosphocreatine, 4 Mg-ATP, 0.5 Tris-GTP,

4 NaCl, pH 7.2-7.3, 290 mOsm (open tip resistance: 5-10 MΩ). Data were acquired using an AM
Systems model 2400 patch-clamp amplifier, online filtered at 2-10 kHz, and digitized at 50 kHz with
a National Instruments PCI-6343 card + BNC2090A interface controlled by Matlab based
acquisition software (Wavesurfer). Data were recorded with the amplifier's pipette capacitance
neutralization circuitry activated. Series resistance was typically between 20-60 MΩ.
511
In vitro electrophysiology: 2-4 weeks following viral injections, mice were deeply anesthetized with
isoflurane, swiftly decapitated, and the brains carefully removed in warm (~34° C), oxygenated
ACSF containing (in mM): 119 NaCl, 25 $NaHCO_3$, 3 KCl, 1.25 NaH_2PO_4, 15 glucose, 1 $MgCl_2$, 1.3
$CaCl_2$, 1 ascorbate, 3 pyruvate. 200-300 μm thick coronal slices of the IC were prepared with a
vibratome (Campden Instruments). On each slice, a small cut was made in the lateral portion of the
right cerebellum or right IC to aid with visual identification of the un-injected hemisphere. Slices
were then incubated at 34° C in a holding chamber filled ACSF for 25-30 min and subsequently
stored at room temperature. Experiments were generally performed within 3-4 hours following slice
preparation. Following incubation, a slice was transferred to a recording chamber and held in place
with single strands of unwaxed dental floss tightly strung around a platinum "harp". The slice was
continuously perfused with oxygenated ACSF heated to 32-34° C (2-4 mL/min; chamber volume: ~
1 mL). 2-5 μM SR95531 was added to the ACSF to block $GABA_A$ receptors in experiments of
Figure 5C-H, Figure 6, Figure 8, Supplemental Figures 2 and 4, and some experiments of Figure 1.
Neurons in the dorso-medial shell IC were visualized via DIC or Dodt contrast optics using a 40x or
63x objective (Zeiss Axioskop 2 FS Plus or Olympus BXW51 microscope). Neurons were targeted
for whole-cell current-clamp recordings with pipettes filled with the same K^+-rich internal solution
used for *in vivo* recordings (open tip resistance: 3-6 MΩ). For whole-cell voltage clamp
experiments, the internal solution contained (in mM): 110 Cesium Methanesulfonate, 10 QX-314-
Bromide, 0.1 EGTA, 10 HEPES, 0.5 Tris-GTP, 4.5 MgATP, 5 TEA-Cl, 10 Tris-phosphocreatine. In
some experiments, 30 μM Alexa 594 or 0.1% biocytin were added to the internal solution to
visualize neuronal morphology via online fluorescence or post-hoc histological reconstruction.
533
Data were acquired with a Multiclamp 700B or AM System model 2400 amplifier, online filtered at
2-10 kHz, and digitized at 50 kHz with a National Instruments PCI-6343 card + BNC2090A interface
controlled by Wavesurfer. In current clamp, pipette capacitance neutralization was employed and
bridge balance was maintained (series resistance typically 10-30 MΩ). In a few instances, a small
amount of negative bias current (-5 to -50 pA) was injected to hyperpolarize neurons and prevent
spike initiation during optogenetic activation of cortico-collicular synapses. Series resistance
compensation was employed in voltage clamp experiments (60-80%, bandwidth: 3 kHz). For dual
pathway experiments, the central IC was electrically stimulated using an AM systems model 2100
stimulus isolator delivering biphasic shocks to a theta glass bipolar electrode placed ~500 μm from
the recorded neuron. Drugs were obtained from Tocris or HelloBio, aliquoted as stock solutions in
distilled water, and stored at -20 C until the day of the experiment.
545
Data analysis: Electrophysiology data were analyzed using custom Matlab scripts. EPSP analyses
were performed on averages of multiple trials (typically >10 trials per condition) after baseline
membrane potential subtraction and lowpass filtering at 1 kHz unless explicitly noted in the text.
Peak amplitudes of single EPSPs were calculated by averaging data points ±0.1 ms around the

local maximum following optogenetic stimulation Halfwidths are calculated as the full width at half-
maximum of the peak. EPSP onset latency was defined as the time following optogenetic stimulus
onset when the membrane potential reaches 20% of peak. The tonic EPSP amplitude during 20 Hz
trains was calculated as follows: We first linearly interpolated the membrane potential data between
each light flash to remove the phasic EPSP component. The trace was then smoothed using a 50
ms sliding window. Datapoints during the final 1s period of the stimulus train were then averaged to
estimate the amplitude of tonic membrane potential change. In certain experiments of Figures 4, 7,
and 8, train stimuli and click sounds occasionally triggered APs in IC neurons both *in vivo* and *in*
vitro. In these cases, APs were digitally removed prior to averaging the traces by linearly
interpolating 0.1-0.2 ms of datapoints after the membrane potential crossed spike threshold (~20
mV/ms). Shock artifacts during electrical stimulation experiments were similarly removed via linear
interpolation. In summary plots, black symbols are individual cells, magenta is mean ± SEM, and
lines connect data from the same recording unless otherwise stated.
563
564 The expected linear waveforms for temporal summation experiments were calculated as follows.
565 The average waveform of a single optogenetically evoked EPSP was peak normalized to the first
566 EPSP in the recorded 20 or 50 Hz train from the same cell. We subsequently convolved the single
567 EPSP waveform with a 20 or 50 Hz binary pulse train using the Matlab function convr(). We then
568 calculated the peak amplitude ratios for each EPSP in observed and expected trains.
569
570 For the free-field sound presentation experiments of Figure 7D-E, we limited our analyses to
571 superficial IC neurons that showed onset EPSPs in response to clicks. Other IC neurons
572 encountered during these experiments showed either sound-evoked IPSPs (n=7 cells from N=5
573 mice) or IPSPs followed by rebound depolarizations (n=6 cells from N=5 mice); analyses of these
574 data will be presented in a separate report.
575
In dual pathway experiments of Figure 8, the onset latency of ascending and descending EPSPs
varied across cells. Thus, the relative timing of electrical and optogenetic stimulation during
combined pathway activation was calculated online and on a cell-by-cell basis, such that the onset
of descending auditory cortical EPSPs collided with the peak of ascending EPSPs from central IC
as predicted from our *in vivo* latency measurements (Figure 7; range of Δt between stimulation of
descending and ascending synapses: -1.3 to 16.4 ms). These stimulation parameters were held
constant across control and R-CPP conditions for each cell. The expected linear summation was
calculated by digitally summing the average synaptic waveforms following stimulation of either
pathway alone, accounting for the temporal offset employed during synchronous pathway
activation.
586
Histology and Confocal Imaging: Mice were deeply anesthetized in a glass induction chamber
circulated with 4.2 mL isoflurane and transcardially perfused with ~80-100 mL of PBS followed by
~80-100 mL of 10% buffered formalin (Fisher Scientific catalog # 23-245684). Brains were carefully
removed, stored in 10% formalin and protected from light for 24 hours. Subsequently, brains were
stored in PBS for up to 72 hours and 100 μm thick coronal slices were cut using a ceramic blade
(Cadence Endurium) and a Leica VT1000s vibratome, mounted onto slides and coverslipped using
Fluoromount, then protected from light and allowed to dry at room temperature for ~12-24 hours.

Slides were then stored at 4 C until ready for use. Images were collected using a Leica TCS SP8
laser scanning confocal microscope equipped with a 10x objective.
596
Statistics: Although not explicitly pre-determined prior to data collection, sample sizes reflect
commonly accepted standards in the field. Data were tested for normality using a Lilliefors test prior
to statistical comparisons. Parametric, two-tailed t-tests are employed for normally distributed data.
Non-parametric rank-sum or sign-rank tests are used when one or more of the distributions deviate
from normal. Alpha is corrected for multiple comparisons in post-hoc significance tests following
ANOVA. Statistics were run in Matlab or Graphpad Prism 9.
603

604 Author Contributions: HMO conducted and analyzed most electrophysiology experiments in Figure
605 7, with contributions from PFA. HMO, AF, and JC performed viral injections. AF performed
606 histology and confocal microscopy experiments in Figure 1. PFA conducted and analyzed most
607 electrophysiology experiments in Figures 1-6 and Figure 8, with contributions from HMO. PFA
608 designed the experiments, interpreted the results, and wrote the paper.
609
Acknowledgments: Funding was generously provided by the Whitehall Foundation, Hearing Health
Foundation, and NIH/NIDCD R01DC019090 to PFA, as well as T32 DC005356 and T32 DC000011
pre-doctoral awards to HMO. We thank Mr. Deepak Dileepkumar for outstanding technical support,
Drs. Michael Roberts, Marina Silveira, Gunnar Quass and Meike Rogalla for critical comments on
the manuscript.

References
617

Abbott LF, Varela JA, Sen K, Nelson SB (1997) Synaptic depression and cortical gain control. Science 275:220–224.

Ahuja TK, Wu SH (2007) Intrinsic membrane properties and synaptic response characteristics of neurons in the rat's
external cortex of the inferior colliculus. Neuroscience 145:851–865.

Anderson LA, Malmierca MS (2013) The effect of auditory cortex deactivation on stimulus-specific adaptation in the
inferior colliculus of the rat. Eur J Neurosci 37:52–62.

Arnth-Jensen N, Jabaudon D, Scanziani M (2002) Cooperation between independent hippocampal synapses is
controlled by glutamate uptake. Nat Neurosci 5:325–331.

Ayala YA, Udeh A, Dutta K, Bishop D, Malmierca MS, Oliver DL (2015) Differences in the strength of cortical and
brainstem inputs to SSA and non-SSA neurons in the inferior colliculus. Sci Rep 5:10383.

Bajo VM, Nodal FR, Korn C, Constantinescu AO, Mann EO, Boyden ES, King AJ (2019) Silencing cortical activity
during sound-localization training impairs auditory perceptual learning. Nat Commun 10:3075.

Bajo VM, Nodal FR, Moore DR, King AJ (2010) The descending corticocollicular pathway mediates learning-induced
auditory plasticity. Nat Neurosci 13:253–260.

Barnstedt O, Keating P, Weissenberger Y, King AJ, Dahmen JC (2015) Functional Microarchitecture of the Mouse
Dorsal Inferior Colliculus Revealed through In Vivo Two-Photon Calcium Imaging. J Neurosci 35:10927–10939.

Blackwell JM, Lesicko AM, Rao W, De Biasi M, Geffen MN (2020) Auditory cortex shapes sound responses in the
inferior colliculus. eLife 9:e51890.

Bordi F, LeDoux JE (1994) Response properties of single units in areas of rat auditory thalamus that project to the
amygdala: II. Cells receiving convergent auditory and somatosensory inputs and cells antidromically activated
by amygdala stimulation. Exp Brain Res 98:275–286.

Brainard MS, Knudsen EI (1993) Experience-dependent plasticity in the inferior colliculus: a site for visual calibration of
the neural representation of auditory space in the barn owl. J Neurosci Off J Soc Neurosci 13:4589–4608.

Branco T, Clark BA, Häusser M (2010) Dendritic discrimination of temporal input sequences in cortical neurons.
Science 329:1671–1675.

Brown SP, Hestrin S (2009) Intracortical circuits of pyramidal neurons reflect their long-range axonal targets. Nature
457:1133–1136.

Cai D, Yue Y, Su X, Liu M, Wang Y, You L, Xie F, Deng F, Chen F, Luo M, Yuan K (2019) Distinct Anatomical
Connectivity Patterns Differentiate Subdivisions of the Nonlemniscal Auditory Thalamus in Mice. Cereb Cortex
29:2437–2454.

Champoux F, Paiement P, Mercier C, Lepore F, Lassonde M, Gagné J-P (2007) Auditory processing in a patient with a
unilateral lesion of the inferior colliculus: Role of the inferior colliculus in humans. Eur J Neurosci 25:291–297.

Chen C, Cheng M, Ito T, Song S (2018) Neuronal Organization in the Inferior Colliculus Revisited with Cell-Type-
Dependent Monosynaptic Tracing. J Neurosci 38:3318–3332.

Clements JD, Lester RA, Tong G, Jahr CE, Westbrook GL (1992) The time course of glutamate in the synaptic cleft.
Science 258:1498–1501.

Coomes DL, Schofield BR (2004) Projections from the auditory cortex to the superior olivary complex in guinea pigs.
Eur J Neurosci 19:2188–2200.

Destexhe A, Rudolph M, Paré D (2003) The high-conductance state of neocortical neurons in vivo. Nat Rev Neurosci
4:739–751.

Diamond IT, Jones EG, Powell TPS (1969) The projection of the auditory cortex upon the diencephalon and brain stem
in the cat. Brain Res 15:305–340.

Doucet JR, Molavi DL, Ryugo DK (2003) The source of corticocollicular and corticobulbar projections in area Te1 of the
rat. Exp Brain Res 153:461–466.

Faye-Lund H, Osen KK (1985) Anatomy of the inferior colliculus in rat. Anat Embryol (Berl) 171:1–20.

Feldman DE, Brainard MS, Knudsen EI (1996) Newly learned auditory responses mediated by NMDA receptors in the
owl inferior colliculus. Science 271:525–528.

Gasparini S, Magee JC (2006) State-dependent dendritic computation in hippocampal CA1 pyramidal neurons. J
Neurosci Off J Soc Neurosci 26:2088–2100.

Geis H-RAP, van der Heijden M, Borst JGG (2011) Subcortical input heterogeneity in the mouse inferior colliculus. J
Physiol 589:3955–3967.

Gottlieb Y, Vaadia E, Abeles M (1989) Single unit activity in the auditory cortex of a monkey performing a short term
memory task. Exp Brain Res 74:139–148.

Goyer D, Silveira MA, George AP, Beebe NL, Edelbrock RM, Malinski PT, Schofield BR, Roberts MT (2019) A novel
class of inferior colliculus principal neurons labeled in vasoactive intestinal peptide-Cre mice. eLife 8:e43770.

Harnett MT, Makara JK, Spruston N, Kath WL, Magee JC (2012) Synaptic amplification by dendritic spines enhances
input cooperativity. Nature 491:599–602.

Huang Y, Matysiak A, Heil P, König R, Brosch M (2016) Persistent neural activity in auditory cortex is related to
auditory working memory in humans and nonhuman primates. eLife 5.

Hurley LM, Pollak GD (2005) Serotonin shifts first-spike latencies of inferior colliculus neurons. J Neurosci Off J Soc
Neurosci 25:7876–7886.

Joswig H, Schönenberger U, Brügge D, Richter H, Surbeck W (2015) Reversible pure word deafness due to inferior
colliculi compression by a pineal germinoma in a young adult. Clin Neurol Neurosurg 139:62–65.

Kerchner GA, Nicoll RA (2008) Silent synapses and the emergence of a postsynaptic mechanism for LTP. Nat Rev
Neurosci 9:813–825.

Kitagawa M, Sakaba T (2019) Developmental changes in the excitatory short-term plasticity at input synapses in the rat
inferior colliculus. Eur J Neurosci 50:2830–2846.

Kole MHP, Letzkus JJ, Stuart GJ (2007) Axon initial segment Kv1 channels control axonal action potential waveform
and synaptic efficacy. Neuron 55:633–647.

Kong L, Xiong C, Li L, Yan J (2014) Frequency-specific corticofugal modulation of the dorsal cochlear nucleus in mice.
Front Syst Neurosci 8 Available at: http://journal.frontiersin.org/article/10.3389/fnsys.2014.00125/abstract
[Accessed November 16, 2020].

Krishna BS, Semple MN (2000) Auditory Temporal Processing: Responses to Sinusoidally Amplitude-Modulated Tones
in the Inferior Colliculus. J Neurophysiol 84:255–273.

Larkum ME, Nevian T, Sandler M, Polsky A, Schiller J (2009) Synaptic Integration in Tuft Dendrites of Layer 5
Pyramidal Neurons: A New Unifying Principle. Science 325:756–760.

LeDoux J, Farb C, Ruggiero D (1990) Topographic organization of neurons in the acoustic thalamus that project to the
amygdala. J Neurosci 10:1043–1054.

Lefort S, Tomm C, Floyd Sarria J-C, Petersen CCH (2009) The excitatory neuronal network of the C2 barrel column in
mouse primary somatosensory cortex. Neuron 61:301–316.

Lesicko AMH, Hristova TS, Maigler KC, Llano DA (2016) Connectional Modularity of Top-Down and Bottom-Up
Multimodal Inputs to the Lateral Cortex of the Mouse Inferior Colliculus. J Neurosci 36:11037–11050.

Lester RA, Jahr CE (1992) NMDA channel behavior depends on agonist affinity. J Neurosci Off J Soc Neurosci
12:635–643.

Li Y, Evans MS, Faingold CL (1998) In vitro electrophysiology of neurons in subnuclei of rat inferior colliculus. Hear Res
121:1–10.

Llano DA, Slater BJ, Lesicko AMH, Stebbings KA (2014) An auditory colliculothalamocortical brain slice preparation in
mouse. J Neurophysiol 111:197–207.

Loftus WC, Malmierca MS, Bishop DC, Oliver DL (2008) The cytoarchitecture of the inferior colliculus revisited: A
common organization of the lateral cortex in rat and cat. Neuroscience 154:196–205.

Losonczy A, Magee JC (2006) Integrative properties of radial oblique dendrites in hippocampal CA1 pyramidal
neurons. Neuron 50:291–307.

Lumani A, Zhang H (2010) Responses of neurons in the rat's dorsal cortex of the inferior colliculus to monaural tone
bursts. Brain Res 1351:115–129.

Ma CL, Kelly JB, Wu SH (2002) AMPA and NMDA receptors mediate synaptic excitation in the rat's inferior colliculus.
Hear Res 168:25–34.

Markram H, Wang Y, Tsodyks M (1998) Differential signaling via the same axon of neocortical pyramidal neurons. Proc
Natl Acad Sci U S A 95:5323–5328.

Massopust LC, Ordy JM (1962) Auditory organization of the inferior colliculi in the cat. Exp Neurol 6:465–477.

Masterton RB, Jane JA, Diamond IT (1968) Role of brain-stem auditory structures in sound localization. II. Inferior
717   colliculus and its brachium. J Neurophysiol 31:96–108.

718   Mellott JG, Foster NL, Ohl AP, Schofield BR (2014) Excitatory and inhibitory projections in parallel pathways from the
719   inferior colliculus to the auditory thalamus. Front Neuroanat 8 Available at:
720   http://journal.frontiersin.org/article/10.3389/fnana.2014.00124/abstract [Accessed July 24, 2020].

721   Mitani A, Shimokouchi M, Nomura S (1983) Effects of stimulation of the primary auditory cortex upon
722   colliculogeniculate neurons in the inferior colliculus of the cat. Neurosci Lett 42:185–189.

723   Moore LA, Trussell LO (2017) Corelease of Inhibitory Neurotransmitters in the Mouse Auditory Midbrain. J Neurosci Off
724   J Soc Neurosci 37:9453–9464.

725   Morris RG, Anderson E, Lynch GS, Baudry M (1986) Selective impairment of learning and blockade of long-term
726   potentiation by an N-methyl-D-aspartate receptor antagonist, AP5. Nature 319:774–776.

727   Nakamoto KT, Jones SJ, Palmer AR (2008) Descending projections from auditory cortex modulate sensitivity in the
728   midbrain to cues for spatial position. J Neurophysiol 99:2347–2356.

729   Nakamoto KT, Shackleton TM, Palmer AR (2010) Responses in the inferior colliculus of the guinea pig to concurrent
730   harmonic series and the effect of inactivation of descending controls. J Neurophysiol 103:2050–2061.

731   Naumov V, Heyd J, de Arnal F, Koch U (2019) Analysis of excitatory and inhibitory neuron types in the inferior
732   colliculus based on I_h properties. J Neurophysiol 121:2126–2139.

733   Oliver DL (1984) Neuron types in the central nucleus of the inferior colliculus that project to the medial geniculate body.
734   Neuroscience 11:409–424.

735   Oliver DL, Hall WC (1978) The medial geniculate body of the tree shrew,Tupaia glis I. Cytoarchitecture and midbrain
736   connections. J Comp Neurol 182:423–458.

737   Peña JL, Konishi M (2001) Auditory spatial receptive fields created by multiplication. Science 292:249–252.

738   Pena JL, Konishi M (2002) From postsynaptic potentials to spikes in the genesis of auditory spatial receptive fields. J
739   Neurosci Off J Soc Neurosci 22:5652–5658.

740   Peruzzi D, Sivaramakrishnan S, Oliver DL (2000) Identification of cell types in brain slices of the inferior colliculus.
741   Neuroscience 101:403–416.

742   Polsky A, Mel B, Schiller J (2009) Encoding and decoding bursts by NMDA spikes in basal dendrites of layer 5
743   pyramidal neurons. J Neurosci Off J Soc Neurosci 29:11891–11903.

744   Ponvert ND, Jaramillo S (2019) Auditory Thalamostriatal and Corticostriatal Pathways Convey Complementary
745   Information about Sound Features. J Neurosci 39:271–280.

746   Pouille F, Scanziani M (2004) Routing of spike series by dynamic circuits in the hippocampus. Nature 429:717–723.

747   Qi J, Zhang Z, He N, Liu X, Zhang C, Yan J (2020) Cortical Stimulation Induces Excitatory Postsynaptic Potentials of
748   Inferior Colliculus Neurons in a Frequency-Specific Manner. Front Neural Circuits 14:591986.

749   Ranganathan GN, Apostolides PF, Harnett MT, Xu N-L, Druckmann S, Magee JC (2018) Active dendritic integration
750   and mixed neocortical network representations during an adaptive sensing behavior. Nat Neurosci 21:1583–
751   1590.

752   Ruediger S, Scanziani M (2020) Learning speed and detection sensitivity controlled by distinct cortico-fugal neurons in
753   visual cortex. eLife 9.

Ryugo DK, Weinberger NM (1976) Corticofugal modulation of the medial geniculate body. Exp Neurol 51:377–391.

Sakurai Y (1990) Cells in the rat auditory system have sensory-delay correlates during the performance of an auditory
working memory task. Behav Neurosci 104:856–868.

Saldaña E, Feliciano M, Mugnaini E (1996) Distribution of descending projections from primary auditory neocortex to
inferior colliculus mimics the topography of intracollicular projections. J Comp Neurol 371:15–40.

Saldaña E, Merchán MA (1992) Intrinsic and commissural connections of the rat inferior colliculus. J Comp Neurol
319:417–437.

Schofield BR, Coomes DL, Schofield RM (2006) Cells in Auditory Cortex that Project to the Cochlear Nucleus in
Guinea Pigs. J Assoc Res Otolaryngol 7:95–109.

Schwartz EJ, Rothman JS, Dugué GP, Diana M, Rousseau C, Silver RA, Dieudonné S (2012) NMDA receptors with
incomplete Mg^{2+} block enable low-frequency transmission through the cerebellar cortex. J Neurosci Off J Soc
Neurosci 32:6878–6893.

Silveira MA, Anair JD, Beebe NL, Mirjalili P, Schofield BR, Roberts MT (2020) Neuropeptide Y Expression Defines a
Novel Class of GABAergic Projection Neuron in the Inferior Colliculus. J Neurosci 40:4685–4699.

Sinex DG, Li H (2007) Responses of Inferior Colliculus Neurons to Double Harmonic Tones. J Neurophysiol 98:3171–
3184.

Sivaramakrishnan S, Oliver DL (2001) Distinct K currents result in physiologically distinct cell types in the inferior
colliculus of the rat. J Neurosci Off J Soc Neurosci 21:2861–2877.

Smith PH (1992) Anatomy and physiology of multipolar cells in the rat inferior collicular cortex using the in vitro brain
slice technique. J Neurosci Off J Soc Neurosci 12:3700–3715.

Song JH, Lucaci D, Calangiu I, Brown MTC, Park JS, Kim J, Brickley SG, Chadderton P (2018) Combining mGRASP
and Optogenetics Enables High-Resolution Functional Mapping of Descending Cortical Projections. Cell Rep
24:1071–1080.

Stebbings KA, Lesicko AMH, Llano DA (2014) The auditory corticocollicular system: Molecular and circuit-level
considerations. Hear Res 314:51–59.

Stuart GJ, Spruston N (2015) Dendritic integration: 60 years of progress. Nat Neurosci 18:1713–1721.

Sun H, Wu SH (2008) Modification of membrane excitability of neurons in the rat's dorsal cortex of the inferior colliculus
by preceding hyperpolarization. Neuroscience 154:257–272.

Sun H, Wu SH (2009) The physiological role of pre- and postsynaptic GABA(B) receptors in membrane excitability and
synaptic transmission of neurons in the rat's dorsal cortex of the inferior colliculus. Neuroscience 160:198–211.

Suthakar K, Ryugo DK (2017) Descending projections from the inferior colliculus to medial olivocochlear efferents: Mice
with normal hearing, early onset hearing loss, and congenital deafness. Hear Res 343:34–49.

Syka J, Popelár J, Kvasñák E, Astl J (2000) Response properties of neurons in the central nucleus and external and
dorsal cortices of the inferior colliculus in guinea pig. Exp Brain Res 133:254–266.

Takahashi H, Magee JC (2009) Pathway interactions and synaptic plasticity in the dendritic tuft regions of CA1
pyramidal neurons. Neuron 62:102–111.

Vila C-H, Williamson RS, Hancock KE, Polley DB (2019) Optimizing optogenetic stimulation protocols in auditory
corticofugal neurons based on closed-loop spike feedback. J Neural Eng 16:066023.

Weible AP, Yavorska I, Wehr M (2020) A Cortico-Collicular Amplification Mechanism for Gap Detection. Cereb Cortex
N Y N 1991 30:3590–3607.

Williamson RS, Polley DB (2019) Parallel pathways for sound processing and functional connectivity among layer 5
and 6 auditory corticofugal neurons. eLife 8:e42974.

Winer JA (2006) Decoding the auditory corticofugal systems. Hear Res 212:1–8.

Winer JA, Diehl JJ, Larue DT (2001) Projections of auditory cortex to the medial geniculate body of the cat. J Comp
Neurol 430:27–55.

Winer JA, Larue DT, Diehl JJ, Hefti BJ (1998) Auditory cortical projections to the cat inferior colliculus. J Comp Neurol
400:147–174.

Wong AB, Borst JGG (2019) Tonotopic and non-auditory organization of the mouse dorsal inferior colliculus revealed
by two-photon imaging. eLife 8:e49091.

Wu SH (2004) Contribution of AMPA, NMDA, and GABAA Receptors to Temporal Pattern of Postsynaptic Responses
in the Inferior Colliculus of the Rat. J Neurosci 24:4625–4634.

Xiao Z, Suga N (2002) Modulation of cochlear hair cells by the auditory cortex in the mustached bat. Nat Neurosci
5:57–63.

Xiong XR, Liang F, Zingg B, Ji X, Ibrahim LA, Tao HW, Zhang LI (2015) Auditory cortex controls sound-driven innate
defense behaviour through corticofugal projections to inferior colliculus. Nat Commun 6:7224.

Yan J, Suga N (1999) Corticofugal Amplification of Facilitative Auditory Responses of Subcortical Combination-
Sensitive Neurons in the Mustached Bat. J Neurophysiol 81:817–824.

Yan W, Suga N (1998) Corticofugal modulation of the midbrain frequency map in the bat auditory system. Nat Neurosci
1:54–58.

Yu Y-Q, Xiong Y, Chan Y-S, He J (2004) Corticofugal gating of auditory information in the thalamus: an in vivo
intracellular recording study. J Neurosci Off J Soc Neurosci 24:3060–3069.

Zucker RS, Regehr WG (2002) Short-term synaptic plasticity. Annu Rev Physiol 64:355–405.

A In Vivo

10 mV
10 ms

B In Vitro

20 mV
5 ms

Supplemental Figure 1: *Auditory cortico-collicular EPSPs can trigger spikes in IC neurons in vivo*
and in vitro.
A) Example overlaid trials from an *in vivo* whole-cell recording of a superficial IC neuron.
Optogenetic stimulation of auditory cortex denoted by arrowhead.
B) Example trials from an *in vitro* recording.

Supplemental Figure 2: *Auditory Cortico-collicular transmission in voltage-clamp.*
A) Example EPSCs recorded in voltage clamp. Gray and black traces are individual trials and
average, respectively. **B-D)** Summary data for EPSC peak amplitudes (B), decay time constant (C),
and 10-90% rise time (D) in n=11 cells.

Supplemental Figure 3: *Large AHPs follow the cessation of auditory cortico-collicular activity in vivo.*

A) Membrane potential (average of multiple trials) during prolonged auditory cortical activity *in vivo*. Magenta are mono-exponential fits to the membrane potential fall and rise upon cessation of optogenetic stimulation.

B) Peak AHP amplitude (relative to baseline Vm) in n=14 cells.

C) Time constants for the fall and raise of the membrane potential for n = 14 cells. The data are on a log scale due to the order of magnitude difference in exponent values for fall and rise.

834   Supplemental Figure 4: *Auditory cortico-collicular EPSCs recorded at positive potentials are slower*
835   *than at negative potentials, consistent with the presence of NMDA receptors.*
836   **A)** Example average EPSCs recorded at negative and positive potentials. Magenta are double
837   exponential fits to the EPSC decay.
838   **B)** Decay time constant for n=5 cells recorded at positive and negative voltages.