1   **The Alzheimer's gene *SORL1* is a key regulator of endosomal recycling in human**

2   **neurons**

3   Swati Mishra[1,2*], Allison Knupp[1,2*], Marcell Szabo[1,2], Chizuru Kinoshita[1,2], Dale W. Hailey[1,2],

4   Yuliang Wang[2,3], and Jessica E. Young[1,2,#]

[1]Department of Laboratory Medicine and Pathology

[2]Institute for Stem Cell and Regenerative Medicine

[3]School of Computer Science and Engineering

University of Washington Seattle WA, 98195, USA

[#]Correspondence: jeyoung@uw.edu

[*]Equal Contribution

12   **ABSTRACT**

Background Loss of the Sortilin-related receptor 1 (*SORL1*) gene seems to act as a causal event

for Alzheimer's disease (AD). Recent studies have established that loss of *SORL1*, as well as

mutations in autosomal dominant AD genes *APP* and *PSEN1/2*, pathogenically converge by

swelling early endosomes, AD's cytopathological hallmark. Acting together with the retromer

trafficking complex, *SORL1* has been shown to regulate the recycling of the amyloid precursor

protein (APP) out of the endosome, contributing to endosomal swelling and to APP

misprocessing. We hypothesized that *SORL1* plays a broader role in neuronal endosomal

recycling and used human induced pluripotent stem cell derived neurons (hiPSC-Ns) to test this

hypothesis. We examined endosomal recycling of three transmembrane proteins linked to AD

pathophysiology: APP, the BDNF receptor Tropomyosin-related kinase B (TRKB), and the

glutamate receptor subunit AMPA1 (GLUA1).

Methods We used isogenic hiPSCs engineered to have *SORL1* depleted or to have enhanced

SORL1 expression. We differentiated neurons from these cell lines and mapped the trafficking of

APP, TRKB and GLUA1 within the endosomal network using confocal microscopy. We also

performed cell surface recycling and lysosomal degradation assays to assess the functionality of

the endosomal network. Finally, we analyzed alterations in gene expression in *SORL1* depleted

neurons using RNA-sequencing.

Results We find that as with APP, endosomal trafficking of GLUA1 and TRKB is impaired by loss

of *SORL1*. Conversely, increased *SORL1* expression enhances endosomal recycling for APP and

GLUA1. Our unbiased transcriptomic data further support *SORL1*'s role in endosomal recycling.

We observe altered expression networks that regulate cell surface trafficking and neurotrophic

signaling

Conclusion Collectively, and together with other recent observations, these findings suggest that

SORL1 is a key and broad regulator of retromer-dependent endosomal recycling in neurons, a

conclusion that has both pathogenic and therapeutic implications for Alzheimer's disease.

43   **BACKGROUND**

Alzheimer's disease (AD) is a progressive neurodegenerative disorder and the most common

cause of dementia. The underlying contributors to AD pathology encompass several biological

pathways, including endosomal function, amyloid precursor protein (APP) processing, immune

function, synaptic function, and lipid metabolism(Karch and Goate, 2015). Among these,

endosomal dysfunction in neurons is emerging as a potential causal mechanism(Small and

Petsko, 2020). Mutations in the amyloid precursor protein (*APP*) and the two presenilins (*PSEN1*

and *PSEN2*) lead to early-onset autosomal dominant AD. When these mutations are modelled in

human neurons and other systems they cause endosomal swelling, indicative of traffic jams, a

phenotype that is a cytopathological hallmark of AD(Cataldo et al., 2000; Choi et al., 2013; Kwart

et al., 2019). Recent genetic studies have identified a fourth gene, the trafficking receptor 'sortilin

related receptor 1' (*SORL1*), which, when harboring frame-shift mutations leading to premature

stop codons, is described as causal for AD(Holstege et al., 2017; Raghavan et al., 2018;

Scheltens et al., 2021). Interestingly, *SORL1* is also linked to the more common, late-onset form

of AD(Lambert et al., 2013; Rogaeva et al., 2007b) and its expression is lost in sporadic AD brains

(Dodson et al., 2006; Thonberg et al., 2017). When modelled in human neurons, *SORL1* depletion

phenocopies *APP* and *PSEN* mutations by causing endosomal swelling(Hung et al., 2021; Knupp

et al., 2020).

The *SORL1* gene codes for the protein SORLA, which functions as part of the retromer trafficking

complex (Fjorback et al., 2012; Rogaeva et al., 2007a; Small and Gandy, 2006). Retromer

recycles cargo out of the early endosome, either from the endosome to the trans-Golgi network

or, with greater importance for neurons, back to the cell surface(Fjorback et al., 2012; Seaman,

2012). To date, the best evidence for *SORL1*'s role in retromer-dependent endosomal recycling

comes from studies investigating APP trafficking(Fjorback et al., 2012; Schmidt et al., 2007;

Willnow and Andersen, 2013). Our previous work demonstrated that *SORL1* depletion retains

APP in early endosomes, which may contribute to endosomal swelling by blocking

recycling(Knupp et al., 2020).

Retromer-dependent trafficking in neurons, however, also recycles cargo other than APP. For

example, retromer is required for the normal recycling of glutamate receptors, a trafficking event

that mediates synaptic plasticity and synaptic health, and this dependency occurs independent of

retromer's role in APP recycling(Park et al., 2004; Temkin et al., 2017). Neurotrophin receptors

are also trafficked through the endosomal system, in a retromer-dependent manner, and are

important for synaptic health(Klinger et al., 2015; Patapoutian and Reichardt, 2001; Rohe et al.,

2013).

Here we used human induced pluripotent stem cell derived-neurons (hiPSC-Ns) to test the

hypothesis that *SORL1* plays a broader role in neuronal endosomal recycling. We use our

previously described *SORL1*-depleted hiPSC lines to generate hiPSC-Ns, which model the loss

of *SORL1* expression that occurs in AD(Knupp et al., 2020). Furthermore, we used previously

established cell lines engineered to overexpress *SORL1* 2-3-fold over wild-type levels(Young et

al., 2015) to test the effects of enhanced *SORL1* expression in hiPSC-Ns on these trafficking

pathways. Importantly, all cell lines are isogenic. We map the trafficking effects these

manipulations have on three specific receptors, APP, the GLUA1 subunit of the AMPA receptor,

and neurotrophin receptor TRKB, all of which are implicated in AD (Devi and Ohno, 2015; Dewar

et al., 1991; Ginsberg et al., 2019; Martin-Belmonte et al., 2020; Wakabayashi et al., 1999;

Yasuda et al., 1995).

Finally, we performed RNA-sequencing on the *SORL1* depleted cell lines to explore an unbiased

transcriptomics analysis induced by *SORL1* depletion. The results generally confirmed our

hypothesis, showing that *SORL1* is a key and broad regulator of endosomal recycling in neurons,

a conclusion that has both pathogenic and therapeutic implications.

RESULTS

***SORL1* depletion increases neuronal cargo localization in early endosomes.**

Using CRISPR/Cas9 genome editing techniques, we previously generated hiPSC-derived

neurons (hiPSC-Ns) deficient in *SORL1* expression due to indels introduced in exon 6. We

demonstrated that loss of *SORL1* expression in these neurons leads to enlarged early endosomes

and an increased colocalization of APP within early endosomes, indicative of endosomal traffic

jams(Knupp et al., 2020). We utilized these same cell lines (hereafter referred to as *SORL1*KO

and their guide-matched isogenic wild-type clones referred to as WT) to examine localization of

the BDNF receptor TRKB and the GLUA1 subunit of the neuronal AMPA receptor. TRKB has

been shown to bind to SORLA and this interaction mediates trafficking of TRKB to synaptic

plasma membranes(Rohe et al., 2013). GLUA1 is trafficked via the retromer complex, of which

SORLA is an adaptor protein(Fjorback et al., 2012; Temkin et al., 2017) and both of these cargo

are important in maintaining healthy neuronal function. Because we previously observed an

increase in APP localization in early endosomes, resulting in a decrease in localization in

downstream vesicles such as Ras-related protein (Rab)7+ late endosomes with *SORL1*

depletion(Knupp et al., 2020), we performed an immunocytochemical analysis of both TRKB and

GLUA1 localization with the early endosome marker EEA1. Similar to our previous observations

for APP, we documented significantly increased localization of both TRKB (Figure 1A) and GLUA1

(Figure 1B) in early endosomes in *SORL1*KOneurons as compared to isogenic WT control

neurons. Accumulation of neuronal cargo in early endosomes is indicative of endosomal traffic

jams, which are thought to impact the transit of cellular cargo through other arms of the endo-

lysosomal network.

SORL1 **depletion mis-traffics cargo throughout the endo-lysosomal system**

The early endosome serves as a hub in which internalized cargo can be retrogradely transported

to the trans-Golgi, recycled back to the cell surface or degraded as endosomes mature into late

endosomes and lysosomes(Mayle et al., 2012). We have previously observed that APP

localization within the trans-Golgi network was decreased in *SORL1*KO neurons(Knupp et al.,

2020). Here we tested whether trafficking to the degradative arm of the endo-lysosomal network

was affected in our *SORL1*KO neurons. Trafficking of substrates out of the early endosome to

late endosomes and, subsequently, lysosomes is important for protein degradation and SORLA

has been previously implicated in promoting Aβ degradation via lysosomes(Caglayan et al.,

2014). We treated *SORL1* deficient neurons with DQ Red BSA, a proteolysis sensitive fluorogenic

substrate that generates fluorescence only when enzymatically cleaved in intracellular lysosomal

compartments. Since substrate degradation primarily occurs in lysosomes, altered fluorescence

intensity of this reagent is a readout of altered lysosomal degradation(Marwaha and Sharma,

2017). We analyzed DQ Red BSA treated neurons for fluorescence intensity using confocal

microscopy. Consistent with loss of *SORL1* leading to endosomal traffic jams, we observed a

significant reduction of DQ Red BSA fluorescence intensity in *SORL1*KO neurons compared to

isogenic WT controls (Figure 2A). We next performed immunocytochemical staining to quantify

the colocalization of our selected neuronal cargo with Rab7, a marker of late endosomes, and

LAMP1 (Lysosomal Associated Membrane Protein 1), a lysosome marker. We show a significant

decrease in co-localization of TRKB (Figure 2B) and GLUA1 (Figure 2C) with Rab7. This result is

consistent with our previous observation for APP (Knupp et al., 2020). We analyzed colocalization

of these cargo with LAMP1 and we observed a significant decrease with APP (Figure 2D) and

TRKB (Figure 2E) and a trend of a decrease with GLUA1 (Figure 2F). These data indicate some

fluidity in the network but suggest that trafficking of APP, TRKB and GLUA1 to late

endosomes/lysosomes is all decreased by *SORL1*KO, although GLUA1 may be more likely to be

trafficked to cell surface pathways or utilizes other adaptor proteins for late endosome to

lysosomal trafficking.

Lysosome size can influence lysosome function and is altered in AD(de Araujo et al., 2020; Hwang

et al., 2019). Similarly, location and number of lysosomes within neurons can alter degradative

activity (Cheng et al., 2018; Farfel-Becker et al., 2019; Gowrishankar et al., 2015; Yap et al.,

2018) and in some cases, altered lysosomal distribution may represent an early neuropathological

defect(Zigdon et al., 2017).Recently, loss of *SORL1* in hiPSC neurons was shown to contribute

to lysosome dysfunction as indicated by both increased lysosome size and number as well as

decreased cathepsin-D activity(Hung et al., 2021). Therefore, we first analyzed LAMP1-

immunopositive puncta and also documented a significant increase in lysosome size and number

in our *SORL1*KO neurons (Supplemental Figure 1A).

Next, we analyzed co-localization of Cathepsin-D and LAMP1 to determine if loss of *SORL1*

expression leads to altered Cathepsin-D trafficking in neurons. Retromer trafficking is required to

deliver one of the most abundant lysosomal proteases, pro-cathepsin D, to lysosomes via the

mannose-6-phosphate receptor (M6PR)(Qureshi et al., 2018; Seaman, 2004). The SORLA

protein has GGA domains similar to that of M6PR(Spoelgen et al., 2006), and mis-trafficking of

Cathepsin-D to lysosomes could affect the maturation and degradative capacity of these

organelles. Therefore, we analyzed co-localization of Cathepsin-D and LAMP1 to determine if

loss of *SORL1* expression leads to altered Cathepsin-D trafficking in neurons. However, we did

not observe a change in Cathepsin-D colocalization between WT and *SORL*1KO (Supplemental

Figure 1B).   Taken together, our data suggest that *SORL1* loss in neurons reduces trafficking of

cargo out of the early endosome to the late endosome and lysosome, contributing to lysosome

stress as evidenced by an increase in size and number in these conditions while the decreased

cathepsin-D activity observed upon *SORL1* loss(Hung et al., 2021) may not be due to impairment

of lysosomal trafficking of the enzyme.

***SORL1* depletion targets the endosomal recycling pathway**

Another route out of the early endosome is via the endocytic recycling complex (ERC) which can

send cargo either to the cell surface or to the trans-Golgi network(Grant and Donaldson, 2009;

Mallard et al., 1998; Marsh et al., 1995; Maxfield and McGraw, 2004). To directly examine if

SORL1 expression alters recycling function, we performed a transferrin recycling assay using

confocal microscopy. Transferrin can be recycled via a fast pathway within approximately 5-10

minutes after being internalized or via a slower pathway involving the ERC over longer periods of

time(Ouellette and Carabeo, 2010; Sonnichsen et al., 2000). We examined the fluorescence

intensity of Alexa Fluor 647-conjugated transferrin over a 40-minute time course in WT and

*SORL1*KO neurons and observed that a higher percentage of intracellular fluorescent transferrin

persisted in *SORL1*KO neurons at both early and later time points as compared to WT neurons,

indicating reduced recycling pathway function (Figure 3A). Cargo destined for the cell surface can

transit to the ERC via Rab11+ recycling endosomes (Ren et al., 1998). Altered size of recycling

endosomes can be indicative of dysfunctional recycling of cargo through these compartments.

We tested whether loss of *SORL1* expression affected the size of Rab11+ recycling endosomes.

Interestingly, we observed a significant increase in the size of Rab11+ recycling endosomes in

the *SORL1*KO neurons (Figure 3B), suggesting that this endosomal compartment is also under

stress. To test if increased size is due to abnormal cargo trafficking through recycling endosomes,

we assessed colocalization of APP, TRKB and GLUA1 with Rab11 and observed increased co-

localization of all three cargo with Rab11+ structures in *SORL1*KO neurons compared to WT

neurons (Figure 3C-E). Together, these data demonstrate that loss of *SORL1* impacts neuronal

recycling endosome pathways by causing traffic jams in the recycling endosomes, similar to the

effect that *SORL1* loss has on early endosomes.

SORL1 depletion reduces cell surface levels of cargo.

Together, our data indicate that *SORL1*KO neurons have impaired cargo recycling with increased

retention of cargo in recycling endosomes. These observations led us to test whether this cargo

was indeed trafficked to the cell surface. A portion of APP has been shown to return to the cell

surface via recycling endosomes(Das et al., 2016) and SORLA can interact with the sorting nexin

SNX27 to return APP to the cell surface(Das et al., 2016; Huang et al., 2016), although in that

study the exact compartment was not described. Furthermore, recycling endosomes are the

source for AMPA receptors during long-term potentiation(Park et al., 2004). We therefore

examined cell surface levels of APP and GLUA1 using immunofluorescence and confocal

microscopy. We documented a significant decrease in cell surface staining of both APP (Figure

4A) and GLUA1 (Figure 4B) in *SORL1*KO neurons as compared to WT, consistent with our

hypothesis that SORLA is involved in regulating traffic from recycling endosomes.

SORL1 overexpression enhances endosomal recycling

Defects in cell surface recycling have severe consequences in neurons, especially as these

processes are necessary for healthy neuronal function. Synapse loss is a main component of AD

neurodegeneration and enhancing pathways that support healthy synaptic function may be

beneficial. We utilized previously generated cell lines that overexpress _SORL1_ cDNA using the

piggybac transposon system(Young et al., 2015) to test whether increased _SORL1_ expression

may enhance the trafficking pathways that are impaired in the _SORL1_KO neurons. Importantly,

the _SORL1_ overexpressing (_SORL1_OE) cell line and control were generated in the same genetic

background as our _SORL1_KO and isogenic WT cell lines. While we found no effect of enhanced

SORL1 expression on DQ Red BSA signal in _SORL1_OE neurons (Supplemental Figure 2A), we

did observe increased localization of the three tested cargo (APP, TRKB, and GLUA1) in late

endosomes and lysosomes in these cells (Supplemental Figure 2B-D and 2E-G), suggesting that

increased _SORL1_ expression may enhance cargo trafficking to the degradative pathway.

We analyzed recycling function using the transferrin recycling assay and observed that _SORL1_OE

neurons showed significantly faster transferrin recycling (Figure 5A). We next tested whether

colocalization of cargo with recycling endosomes and cell surface recycling was altered between

_SORL1_OE and WT neurons. Interestingly, the size of Rab11+ recycling endosomes was

significantly smaller in _SORL1_OE neurons (Figure 5B) possibly indicating that increased _SORL1_

expression is clearing cargo more rapidly from this compartment. We observed a significant

increase in localization of cargo with Rab11+ recycling endosomes (Figure 5C-E). While this result

was initially surprising, as we also saw increased colocalization with Rab11+recycling endosomes

in our _SORL1_KO neurons (Figure 3C-E), we further documented a significant increase of APP

and GLUA1 on the cell surface compared to WT neurons with only endogenous _SORL1_

expression (Figure 5F and 5G), as opposed to decreased APP and GLUA1 localization on the

cell surface in _SORL1_KO neurons (Figure 4A and 4B). These results suggest that cell surface

trafficking via a Rab11 pathway is enhanced by increased *SORL1* expression and that a crucial

action of SORLA is the trafficking out of recycling endosomes. Thus, our data support a critical

role for SORLA for trafficking cargo from recycling endosomes to the cell surface. In addition, we

show for the first time that SORLA levels may regulate cell surface recycling of AMPA receptor

subunits in human neurons.

***SORL1* depletion's effect on gene expression**.

To determine a more global effect of chronic *SORL1* loss in human neurons, we performed bulk

RNA sequencing of *SORL1*KO neurons compared to WT neurons. Interestingly, we observed that

there were significantly more down-regulated genes in *SORL1*KO neurons than upregulated ones

(Supplemental Figure 2B). While none of the cargo we explicitly studied in this work was

differentially expressed, GO analysis showed that the top downregulated molecular function

pathways in the *SORL1*KO cells were related to receptor-ligand activity and extracellular matrix

organization (Figure 6A). The top upregulated molecular function pathways were related to ion

channel activity (Figure 6B). To understand these data in the context of an integrated network,

we used an analysis method that infers ligand receptor interactions from bulk RNA-seq

data(Ramilowski et al., 2015; Wang et al., 2020). We observed several nodes of altered receptor-

ligand interactions that indicate altered cell surface recycling and neurotrophic activity (Figure

6C). These include alterations in β-integrin signaling, which is consistent with previous work

showing reduced β-integrin on the cell surface in *SORL1*KO cancer cells(Pietila et al., 2019), and

altered interactions in ephrins/ephrin receptors, also corroborating previous work implicating

SORL1 expression in ephrin signaling and synapse regulation(Huang et al., 2017). Our analysis

also showed nodes with alterations in nerve growth factor/nerve growth factor receptor

(NGF/NGFR) and fibroblast growth factor/fibroblast growth factor receptor (FGF/FGFR) signaling,

indicating alterations in neurotrophin and growth factor signaling and suggesting that the presence

of endosomal traffic jams in *SORL1*KO neurons may ultimately impact multiple pathways

important for neuronal health and development.

262   **DISCUSSION**

Trafficking through the endo-lysosomal network regulates intra-cellular location of proteins,

dictating their homeostasis, function and influence on cellular physiology. Functional studies by

our group and others document endosomal abnormalities in hiPSC-derived neuronal models of

AD(Hung et al., 2021; Knupp et al., 2020; Kwart et al., 2019). Emerging from this evidence is the

role of *SORL1* as an endosomal gene that plays essential roles in mediating cargo trafficking.

Recent work has implicated *SORL1* as AD's fourth causal gene(Olav M. Andersen, 2021;

Scheltens et al., 2021), and of these genes it is the only one linked to the common late-onset form

of the disease. Understanding *SORL1*'s function is paramount for understanding AD's pathogenic

mechanisms and for potential therapeutic interventions.

Acting as a receptor of the retromer trafficking complex, SORLA has already been pathogenically

linked to AD by its role in recycling APP out of endosomes(Andersen et al., 2005; Herskowitz et

al., 2012; Offe et al., 2006). This current work and our previous study(Knupp et al., 2020) shows

that *SORL1* depletion leads to increased APP localization in early and recycling endosomes. By

lengthening the residence time of APP in these endosomal compartments, accelerated

amyloidogenic cleavage of APP occurs due to the close proximity of APP and BACE1(Sun and

Roy, 2018). Indeed, loss of *SORL1* leads to the accumulation of Aβ peptides, an antecedent of

'amyloid pathology' (Andersen et al., 2005; Knupp et al., 2020; Rogaeva et al., 2007b). We

hypothesized that loss of *SORL1* in neurons would impact other cargo important for healthy

neuronal function. To test this hypothesis, in addition to APP, we examined localization of the

neurotrophin receptor TRKB and the GLUA1 subunit of the AMPA receptor. These cargo link to

another key pathology of AD: neurodegeneration, a slowly progressive process that begins with

synaptic dysfunction characterized by glutamate receptor loss, which then progresses to synaptic

loss before ultimately, over years, leading to widespread neuronal cell death (Selkoe, 2002).

Our experiments point to *SORL1*'s key role in cell surface recycling. By using a prototypical cargo,

transferrin, we demonstrate a reciprocal role between loss and enhancement of *SORL1*

expression in cell surface recycling. Specifically, by showing that *SORL1* plays a role in recycling

glutamate receptors, a trafficking event critical for preventing synaptic dysfunction and synaptic

loss, our results link *SORL1* to AD's early-stage neurodegenerative process. Since retromer-

dependent glutamate receptor recycling has been shown to occur independent of APP(Temkin et

al., 2017), our results suggest that *SORL1* mutations can, at least in principle, drive two key AD

pathologies, amyloid pathology and synaptic pathology, through parallel mechanisms(Small and

Petsko, 2020).

Interestingly, AD's defining cytopathology, swollen endosomes, also seems to occur independent

of Aβ peptides. In the case of early-onset causal genes, studies have shown that APP and PSEN

mutations lead to endosomal swelling even when the production of Aβ peptides are arrested, but

is driven instead by an APP intermediate, the β−C-terminal fragment produced by BACE1

cleavage(Kwart et al., 2019). We previously demonstrated that endosomal swelling occurs in

*SORL1*KO neurons even when BACE1 is inhibited(Knupp et al., 2020). However, a recent study

showed that there is a reduction in endosomal swelling when total APP expression is

reduced(Hung et al., 2021). One interpretation that reconciles these findings is that reduction of

cargo in general, especially one as abundant as APP, may relieve some of the stress on neuronal

endosomes induced by *SORL1* deficiency. Collectively, however, our data suggest that AD's core

histopathology, amyloid plaques, and cytopathology, swollen endosomes, occur through parallel

events.

To test the hypothesis that *SORL1* functions more broadly in neuronal endosomal recycling, we

measured recycling endosome size and showed that, in addition to APP and GLUA1, the

neurotrophin receptor TRKB, is dependent on *SORL1* for normal endosomal trafficking. TRKB is

part of a different family of proteins that depend on retromer-dependent endosomal recycling for

its normal trafficking, and one that has been linked to synaptic health(Klinger et al., 2015; Rohe

et al., 2013) (Patapoutian and Reichardt, 2001).

Our unbiased transcriptomic screen further supported that neurotrophic signaling and cell surface

recycling pathways are impacted by *SORL1* deficiency. Our goal for this analysis was to

determine the global effect of *SORL1* loss on neuronal networks. Indeed, our analysis does not

show that the specific cargo proteins described here are differentially expressed. However, the

analysis does indicate that loss of *SORL1* in human neurons impacts cell surface networks,

including receptor ligand interactions in neurotrophic and growth factor pathways, β-integrin

signaling, and ephrin signaling. Our analysis corroborates previous work and the altered networks

we observe impact neuronal health, axonal guidance, and synapse formation(Huang et al., 2017;

Huang et al., 2006; Pietila et al., 2019).

The early endosome is considered the central station in the sorting and trafficking of cargo

throughout the many stations of the endo-lysosomal system. While the early endosome is the

station that is affected first and foremost in AD, it is not surprising that a primary dysfunction in

this central station will secondarily influence trafficking throughout the system. Our work, along

with other recent work(Hung et al., 2021) also supports a role for *SORL1* in lysosomal trafficking

in neurons. We observed an increase in lysosome size and altered localization of APP, TRKB,

and GLUA1 in late endosomes and lysosomes in *SORL1*KO neurons. Hung et al., also reported

decreased Cathepsin-D activity in *SORL1* deficient neurons, suggesting that *SORL1* loss directly

impacts lysosome function. We interpret our functional and colocalization data to suggest that

loss of *SORL1* expression mainly affects trafficking of cargo to lysosomes, but our data does not

rule out a role of SORLA in neuronal lysosome function. While we did not observe a difference

in Cathepsin-D localization to lysosomes, it is important to note that that LAMP1 only partially

colocalizes with Cathepsin-D in neurons (Cheng et al., 2018). Furthermore, the loss of proteolytic

activity evidenced by decreased intensity of DQ Red BSA in *SORL1*KO neurons may not be

completely due to reduced trafficking to lysosomes but could be a result of abnormal lysosomal

function as DQ Red BSA is internalized by a process called macropinocytosis wherein

macropinosomes can be directly trafficked to lysosomes (Hamasaki et al., 2004; Lorenzen et al.,

2010; Racoosin and Swanson, 1993). Thus, the loss of proteolytic activity evidenced by

decreased intensity of DQ Red BSA in *SORL1*KO neurons may not be completely due to reduced

trafficking to lysosomes but could also be a result of abnormal lysosomal function.

347   **CONCLUSION**

While the secondary downstream effects induced by *SORL1* depletion in the endo-lysosomal

system are interesting and likely relevant to AD's ultimate pathogenesis, from a therapeutic

perspective it is best to target *SORL1*'s primary defect, which seems to localize to the endosomal

recycling pathway. Interestingly, recent biomarker studies suggest that defects in retromer-

dependent endosomal recycling occur in a majority of patients with 'sporadic' AD(Simoes et al.,

2020), suggesting that the observed *SORL1*-induced defects may generalize across early and

late onset forms of the disorder. Collectively, our results support the conclusion that *SORL1*, and

the retromer-dependent pathway in which it functions, is a valid therapeutic target and

interventions directed at this pathway may ameliorate endosomal recycling defects that seem to

act as, at least, one primary driver of AD.

359   **METHODS**

Cell lines

Cell lines generated by CRIPSR/Cas9 gene editing technology

The generation of the cell lines used in this paper is described in our previously published work

(Knupp et al., 2020) and consists of four clones: Two wild-type clones, designated clone A6 and

clone A7, and two *SORL1*KO clones, designated clone E1 and clone E4. Cell lines were

generated from our previously published and characterized CV background human induced

pluripotent stem cell line(Young et al., 2015). This cell line is male and has a APOE ε3/ε4

genotype(Levy et al., 2007). All four clones were shown to have normal karyotypes and are

routinely tested for mycoplasma (MycoAlert).

CRISPR/Cas9 gRNA, ssODN, and Primer Sequences gRNA: ATTGAACGACATGAACCCTC

ssODN:

GGGAATTGATCCCTATGACAAACCAAATACCATCTACATTGAACGACATGAACCCTCTGGC

TACTCCACGTCTTCCGA AGTACAGATTTCTTCCAGTCCCGGGAAAACCAGGAAG Forward

primer: ctctatcctgagtcaaggagtaac Reverse primer: ccttccaattcctgtgtatgc PCR amplifies 458 bp

sequence. These sequences have been previously published in (Knupp et al., 2020)

SORL1 overexpression cell lines Isogenic cell lines with overexpression of SORL1 were

generated as previously described(Young et al., 2015). These lines are generated from the CV

parental line, the same parental line as the *SORL1*KO cell lines were made from. Briefly, stable

integration of SORL1 cDNA into the genome was achieved by using piggybac transposon system

(Systems Biosciences). Vector alone (WT) or vector with SORL1 cDNA (*SORL1*OE) constructs

were introduced into iPSCs by electroporation and stable cell lines were selected with puromycin

(2ug/ml) treatment. For all overexpression experiments *SORL1*OE cells were compared to the

vector alone controls.

hiPSC Neuronal Differentiation

hiPSCs were differentiated to neurons using dual-SMAD inhibition(Chambers et al., 2009; Shi et

al., 2012). Briefly, hiPSCs were plated on Matrigel coated 6-well plates at a density of 3.5 million

cells per well and fed with Basal Neural Maintenance Media (1:1 DMEM/F12 + glutamine

media/neurobasal media, 0.5% N2 supplement, 1% B27 supplement, 0.5% GlutaMax, 0.5%

insulin-transferrin-selenium, 0.5% NEAA, 0.2% β-mercaptoethanol; Gibco, Waltham, MA) +

10mM SB-431542 + 0.5mM LDN-193189 (Biogems, Westlake Village, CA). Cells were fed daily

for seven days. On day eight, cells were incubated with Versene, gently dissociated using cell

scrapers, and passaged at a ratio of 1:3. On day nine, media was switched to Basal Neural

Maintenance Media and fed daily. On day 13, media was switched to Basal Neural Maintenance

Media with 20 ng/mL FGF (R&D Systems, Minneapolis, MN) and fed daily. On day sixteen, cells

were passaged again at a ratio of 1:3. Cells were fed until approximately day twenty-three. At this

time, cells were FACS sorted to obtain the CD184/CD24 positive, CD44/CD271 negative neural

precursor cell (NPC) population. Following sorting, NPCs were expanded for neural

differentiation. For cortical neuronal differentiation, NPCs were plated out in 10cm cell culture

dishes at a density of 6 million cells/10cm plate. After 24 hours, cells were switched to Neural

Differentiation media (DMEM-F12 + glutamine, 0.5% N2 supplement, 1% B27 supplement, 0.5%

GlutaMax) + 0.02ug/mL brain-derived neurotrophic factor (PeproTech, Rocky Hill, NJ) +

0.02ug/mL glial-cell-derived neurotrophic factor (PeproTech) + 0.5mM dbcAMP (Sigma Aldrich,

St Louis, MO). Media was refreshed twice a week for three weeks. After three weeks, neurons

were selected for CD184/CD44/CD271 negative population by MACS sorting and plated for

experiments.

Purification of Neurons

Following three weeks of differentiation, neurons were dissociated with accutase and

resuspended in Magnet Activated Cell Sorting (MACS) buffer (PBS + 0.5% bovine serum albumin

[Sigma Aldrich, St Louis, MO] + 2mM ethylenediaminetetraacetic acid [Thermo Fisher Scientific,

Waltham, MA]). Following a modification of (Yuan et al., 2011), cells were incubated with PE-

conjugated mouse anti-Human CD44 and mouse anti-Human CD184 antibodies (BD

Biosciences, San Jose, CA) at a concentration of 5µl/10 million cells. Following antibody

incubation, cells were washed with MACS buffer and incubated with anti-PE magnetic beads (BD

Biosciences, San Jose, CA) at a concentration of 25µl/10 million cells. Bead-antibody complexes

were pulled down using a rare earth magnet, supernatants were selected, washed, and plated at

an appropriate density.

DQ Red BSA assay for visualization of lysosomal degradation

Lysosomal proteolytic degradation was evaluated using DQ Red BSA (#D-12051; Thermo Fisher

Scientific), a fluorogenic substrate for lysosomal proteases, that generates fluorescence only

when enzymatically cleaved in intracellular lysosomal compartments. hiPSC-derived neurons

were seeded at a density of 400,000 cells/well of a matrigel coated 48 well plate. After 24 hours,

cells were washed once with DPBS, treated with complete media containing either 10µg/ml DQ

Red BSA or vehicle (PBS) and incubated for 5 hours at 37°C in a 5% CO_2 incubator as described

in(Marwaha and Sharma, 2017). At the end of 5 hours, cells were washed with PBS, fixed with

4% PFA and immunocytochemistry was performed as described in methods. Cells were imaged

using a Leica SP8 confocal microscope and all image processing was completed with ImageJ

software. Cell bodies were identified by MAP2 labeling, and fluorescence intensity of DQ Red

BSA was measured in regions of the images containing the MAP2 label.

Immunocytochemistry

For immunocytochemistry, cells were fixed with 4% PFA for 20 minutes. Fixed cells were washed

three times with PBST (PBS with 0.05% tween 20), permeabilized with Triton X-100 in PBS for

15 minutes, washed twice again with PBST, blocked with 5% BSA in PBS at room temperature

for 1h and incubated with appropriate primary antibodies overnight at 4°C. The next day, cells

were incubated with appropriate secondary antibodies and 1µg/ml DAPI for 1 hour at RT, washed

three times with PBST and mounted on glass slides with Prolong Gold Antifade mountant

(#P36930; Thermo Fisher Scientific).

Colocalization analysis

To investigate colocalization with endo-lysosomal compartments, hiPSC-derived neurons were

labeled with markers specific for each intra-cellular compartment (EEA1 for early endosomes,

Rab7 for late endosomes, LAMP1 for lysosomes and Rab11 for recycling endosomes) using

immunocytochemistry. A minimum of 10 fields of confocal z-stack images were captured under

blinded conditions using a Yokogawa W1 spinning disk confocal microscope (Nikon) and a 100X

plan apochromat oil immersion objective. Median filtering was used to remove noise from images

and Otsu thresholding was applied to all images. Colocalization was quantified using the JACOP

plugin(Bolte and Cordelieres, 2006) in Image J software(Schindelin et al., 2012) and presented

as Mander's correlation coefficient.(Dunn et al., 2011; Manders et al., 1993)

Cell Surface Staining: Cell surface expression of GLUA1 and APP was determined using

immunocytochemistry and confocal microscopy. To label proteins at the cell surface, cells were

fixed with 4% PFA, washed and treated with primary and secondary antibodies as described in

the 'immunocytochemistry' section of methods. Permeabilization with 0.1% Triton X-100 was not

performed for this experiment. To label total protein levels, cells were fixed with 4% PFA, washed,

permeabilized with 0.1% Triton X-100 and treated with primary and secondary antibodies as

described in the 'immunocytochemistry' section of methods. Analysis of fluorescence intensity

was done using Image J software. Cell surface expression was represented as ratio of

fluorescence intensity measured under non-permeabilized conditions and fluorescence intensity

measured under permeabilized conditions.

 Antibodies

The following primary antibodies were used: Early endosome antigen 1 (EEA1) at 1:500

(#610456; BD Biosciences); amyloid precursor protein (APP) at 1:500 (#ab32136; Abcam);

microtubule-associated protein 2 (MAP2) at 1:1000 (ab92434; Abcam); Ras-related protein Rab-

7a (Rab7) at 1:1000 (ab50533; Abcam); Ras-related protein Rab-11 (Rab11) at 1:250 (#610656;

BD Biosciences); Lysosome associated membrane protein-1 (LAMP1) at 1:250 (#sc 2011; Santa

Cruz); Tropomyosin receptor kinase B (TRKB; # ab18987; abcam) and GLUA1(# MAB2263;

Millipore sigma) at 1:500(DAPI at 1 mg/mL final (Alfa Aesar).

Transferrin recycling assay

To measure recycling pathway function, we utilized transferrin recycling assay as previously

described(Rapaport et al., 2010). Purified neurons were seeded at 400,000 cells/well of a 24 well

plate containing matrigel coated 12 mm glass coverslip/well. After 5 DIV, cells were washed once

with DMEM-F12 medium and incubated with starving medium (DMEM-F12 medium+25mM

HEPES+ 0.5% BSA) for 30 minutes at 37°C in a 5% CO_2 incubator to remove any residual

transferrin. Thereafter, cells were pulsed with either 100µg/ml transferrin from human serum

conjugated with Alexa Fluor™ 647(#T23366; Thermo Fisher Scientific) or vehicle (PBS) in

'starving medium'. At the end of 10 mins, cells were washed twice with ice-cold PBS to remove

any external transferrin and stop internalization of transferrin and washed once with acid stripping

buffer (25mM citric acid+24.5mM sodium citrate+280mM sucrose+0.01mM Deferoxamine) to

remove any membrane bound transferrin. Next, cells were either fixed in 4% PFA or 'Chase

medium' (DMEM-F12+50µM Deferoxamine+20mM HEPES+500µg/ml Holo-transferrin) was

added for different time points. Immunocytochemistry was done using MAP2 antibody to label

neurons, confocal images were captured using Leica SP8 confocal microscope under blinded

conditions. Fluorescence intensity of transferrin was measured using ImageJ software. Recycling

function was presented as transferrin fluorescence intensity as a percentage of the fluorescence

intensity measured at time zero.

Measurement of lysosome and recycling endosome size

Immunocytochemistry using antibodies for LAMP1 and MAP2 or RAB11 and MAP2 was

performed as described above. Using a Leica SP8 confocal microscope with an apochromatic

63X oil immersion lens, z-stack images were obtained under blinded conditions. For the LAMP1

analysis, 17-34 fields were analyzed for a total of **45 – 76** cells analyzed. For the RAB11 analysis,

15-30 fields were analyzed for a total of **59 – 124** cells analyzed. Vesicle size measurements were

performed using Cell Profiler software as previously described (Knupp et al., 2020) (McQuin et

al., 2018). Briefly, the vesicle channel was masked using the MAP2 channel and automated

segmentation algorithms were used to identify individual puncta. The pixel area of each puncta

was measured and is presented as mean area of all puncta per field.

Statistical Analysis

For all the experiments, at least two independent clones of each genotype were used. For all

imaging experiments the data was analyzed in a blinded manner. Experimental data was tested

for normal distributions using the ShapiroWilk normality test. Normally distributed data was

analyzed using parametric two-tailed unpaired t tests. Non-normally distributed data was analyzed

by non-parametric Mann-Whitney test. Significance was defined as a value of $p > 0.05$. All

statistical analysis was completed using GraphPad Prism software. Statistical details of individual

experiments, including biological and technical replicate information, can be found in the figure

legends.

RNA- sequencing analysis

RNA Extraction

RNA was collected from 3 separate differentiations including a combination of two WT clones and

two *SORL1*KO clones. Each sample includes 2-3 technical replicates. RNA was collected from 2

million purified neurons for each sample. Purification of total RNA was completed using the

PureLink RNA Mini Kit (Thermo Fisher 12183018A). Assessment of purified RNA was completed

using a NanoDrop. Final RNA quantification was completed using the Quant-iT RNA assay

(Invitrogen) and RNA integrity analysis was completed using a fragment analyzer (Advanced

Analytical).

Library Prep and Sequencing

Library preparation was completed using the TruSeq Stranded mRNA kit (Illumina RS-122-2103)

per manufacturer instructions. Sequencing was performed on a NovaSeq 6000 instrument.

Data analysis

Raw reads were aligned to GRCh38 with reference transcriptome GENCODE release 29 using

STAR v2.6.1d (Dobin, Alexander et al. "STAR: ultrafast universal RNA-seq

aligner." *Bioinformatics (Oxford, England)* vol. 29,1 (2013): 15-21.

doi:10.1093/bioinformatics/bts635). Gene-level expression quantification is generated by RSEM

v1.3.1(Li and Dewey, 2011). Genes with fewer than 20 normalized reads across all samples were

omitted from further analysis. We did observe variation in the transcriptome based on

differentiation (Supplemental Figure 3A), however this was corrected for using the sva

package(Leek et al., 2012).

To identify differentially expressed genes (DEGs), we used DESeq (Simon Anders and Wolfgang

Huber (2010): Briefly, we fit two models: a null model where gene expression only depends on

batch effects (i.e., differentiation), and an alternative model where gene expression depends on

both genotype (*SORL1*KO vs. WT) and batch effects. Chi-squared tests were performed to

compare both fits, and we declare a gene as differentially expressed only when the alternative

model fits the expression data better. DEGs are defined as genes with false discovery rate less

than 0.05 and fold change greater than 1.5. The top gene ontology (GO) package (Alexa A,

Rahnenfuhrer J (2021). *topGO: Enrichment Analysis for Gene Ontology*. R package version

2.44.0) was used to identify GO terms that were enriched. GO terms were tested according to the

Fisher's exact test. Finally, we mapped DEGs onto receptor-ligand interaction diagrams

generated by Ramilowski et al(Ramilowski et al., 2015) using the igraph plugin (Csardi G, Nepusz

T (2006). "The igraph software package for complex network research." *InterJournal*, **Complex**

Systems, 1695. https://igraph.org). To compare the amount of down-regulated vs. up-regulated

genes we used a 2-sample test for equality of proportions with continuity correction in R.

Declarations

Availability of Data and Material

All raw and processed RNA-seq data has been deposited at the NCBI Gene Expression Omnibus

(GSE180793). The data sets used and/or analyzed during the current study are available from

the corresponding author on reasonable request.

The authors declare no competing interests

Acknowledgements:

We thank Dr. Harald Frankowski, Dr. Yoshito Kinoshita, Ms. Shannon Rose, Ms. Eden

Cruickshank and all members of the Young Laboratory for critical discussions and feedback

during the preparation of this manuscript. We also thank Dr. Scott A. Small and Dr. Gregory A.

Petsko for critical comments, discussions and feedback on this work. We would like to

acknowledge the UW SLU Cell Analysis Facility and the Garvey Imaging Core at the UW Institute

for Stem Cell and Regenerative Medicine. This work was supported by a NIH grant

(R01AG062148) and a BrightFocus Foundation grant (A2018656S) to J.E.Y., a Biogen

Sponsored Research Agreement to J.E.Y., a Retromer Therapeutic Sponsored Research

Agreement to J.E.Y. and NIH traning grant (T32AG052354) to A.K. and a generous gift from the

Ellison Foundation (to UW).

Author Contributions:

Conceptualization: J.E.Y., S.M, A.K. Microscopy analysis: S.M, A.K., and D.W.H. RNA-seq

analysis and bioinformatics: Y.W and A.K. Methodology: S.M., A.K., M.S., C.K. Writing-Original

Draft: S.M., A.K. and J.E.Y. Writing-Reviewing and editing: J.E.Y., S.M., A.K. Funding Acquisition:

J.E.Y. and A.K. Supervision: J.E.Y.

References

Andersen, O.M., Reiche, J., Schmidt, V., Gotthardt, M., Spoelgen, R., Behlke, J., von Arnim, C.A.,
Breiderhoff, T., Jansen, P., Wu, X.*, et al.* (2005). Neuronal sorting protein-related receptor
sorLA/LR11 regulates processing of the amyloid precursor protein. Proc Natl Acad Sci U S A *102*,
13461-13466.
Bolte, S., and Cordelieres, F.P. (2006). A guided tour into subcellular colocalization analysis in light
microscopy. J Microsc *224*, 213-232.
Caglayan, S., Takagi-Niidome, S., Liao, F., Carlo, A.S., Schmidt, V., Burgert, T., Kitago, Y.,
Fuchtbauer, E.M., Fuchtbauer, A., Holtzman, D.M.*, et al.* (2014). Lysosomal sorting of amyloid-
beta by the SORLA receptor is impaired by a familial Alzheimer's disease mutation. Science
translational medicine *6*, 223ra220.
Cataldo, A.M., Peterhoff, C.M., Troncoso, J.C., Gomez-Isla, T., Hyman, B.T., and Nixon, R.A. (2000).
Endocytic pathway abnormalities precede amyloid beta deposition in sporadic Alzheimer's
disease and Down syndrome: differential effects of APOE genotype and presenilin mutations. Am
J Pathol *157*, 277-286.
Chambers, S.M., Fasano, C.A., Papapetrou, E.P., Tomishima, M., Sadelain, M., and Studer, L.
(2009). Highly efficient neural conversion of human ES and iPS cells by dual inhibition of SMAD
signaling. Nat Biotechnol *27*, 275-280.
Cheng, X.-T., Xie, Y.-X., Zhou, B., Huang, N., Farfel-Becker, T., and Sheng, Z.-H. (2018).
Characterization of LAMP1-labeled nondegradative lysosomal and endocytic compartments in
neurons. J Cell Biol *217*, 3127-3139.

Choi, J.H., Kaur, G., Mazzella, M.J., Morales-Corraliza, J., Levy, E., and Mathews, P.M. (2013). Early endosomal abnormalities and cholinergic neuron degeneration in amyloid-beta protein precursor transgenic mice. J Alzheimers Dis *34*, 691-700.

Das, U., Wang, L., Ganguly, A., Saikia, J.M., Wagner, S.L., Koo, E.H., and Roy, S. (2016). Visualizing APP and BACE-1 approximation in neurons yields insight into the amyloidogenic pathway. Nat Neurosci *19*, 55-64.

de Araujo, M.E.G., Liebscher, G., Hess, M.W., and Huber, L.A. (2020). Lysosomal size matters. Traffic *21*, 60-75.

Devi, L., and Ohno, M. (2015). TrkB reduction exacerbates Alzheimer's disease-like signaling aberrations and memory deficits without affecting β-amyloidosis in 5XFAD mice. Translational psychiatry *5*, e562-e562.

Dewar, D., Chalmers, D.T., Graham, D.I., and McCulloch, J. (1991). Glutamate metabotropic and AMPA binding sites are reduced in Alzheimer's disease: an autoradiographic study of the hippocampus. Brain Res *553*, 58-64.

Dodson, S.E., Gearing, M., Lippa, C.F., Montine, T.J., Levey, A.I., and Lah, J.J. (2006). LR11/SorLA expression is reduced in sporadic Alzheimer disease but not in familial Alzheimer disease. J Neuropathol Exp Neurol *65*, 866-872.

Dunn, K.W., Kamocka, M.M., and McDonald, J.H. (2011). A practical guide to evaluating colocalization in biological microscopy. Am J Physiol Cell Physiol *300*, C723-C742.

Farfel-Becker, T., Roney, J.C., Cheng, X.T., Li, S., Cuddy, S.R., and Sheng, Z.H. (2019). Neuronal Soma-Derived Degradative Lysosomes Are Continuously Delivered to Distal Axons to Maintain Local Degradation Capacity. Cell Rep *28*, 51-64.e54.

Fjorback, A.W., Seaman, M., Gustafsen, C., Mehmedbasic, A., Gokool, S., Wu, C., Militz, D., Schmidt, V., Madsen, P., Nyengaard, J.R.*, et al.* (2012). Retromer binds the FANSHY sorting motif in SorLA to regulate amyloid precursor protein sorting and processing. The Journal of neuroscience : the official journal of the Society for Neuroscience *32*, 1467-1480.

Ginsberg, S.D., Malek-Ahmadi, M.H., Alldred, M.J., Chen, Y., Chen, K., Chao, M.V., Counts, S.E., and Mufson, E.J. (2019). Brain-derived neurotrophic factor (BDNF) and TrkB hippocampal gene expression are putative predictors of neuritic plaque and neurofibrillary tangle pathology. Neurobiology of Disease *132*, 104540.

Gowrishankar, S., Yuan, P., Wu, Y., Schrag, M., Paradise, S., Grutzendler, J., De Camilli, P., and Ferguson, S.M. (2015). Massive accumulation of luminal protease-deficient axonal lysosomes at Alzheimer's disease amyloid plaques. Proceedings of the National Academy of Sciences *112*, E3699-E3708.

Grant, B.D., and Donaldson, J.G. (2009). Pathways and mechanisms of endocytic recycling. Nat Rev Mol Cell Biol *10*, 597-608.

Hamasaki, M., Araki, N., and Hatae, T. (2004). Association of early endosomal autoantigen 1 with macropinocytosis in EGF-stimulated A431 cells. Anat Rec A Discov Mol Cell Evol Biol *277*, 298-306.

Herskowitz, J.H., Offe, K., Deshpande, A., Kahn, R.A., Levey, A.I., and Lah, J.J. (2012). GGA1-mediated endocytic traffic of LR11/SorLA alters APP intracellular distribution and amyloid-beta production. Mol Biol Cell *23*, 2645-2657.

Holstege, H., van der Lee, S.J., Hulsman, M., Wong, T.H., van Rooij, J.G., Weiss, M., Louwersheimer, E., Wolters, F.J., Amin, N., Uitterlinden, A.G.*, et al.* (2017). Characterization of

pathogenic SORL1 genetic variants for association with Alzheimer's disease: a clinical
interpretation strategy. Eur J Hum Genet *25*, 973-981.

Huang, T.Y., Zhao, Y., Jiang, L.L., Li, X., Liu, Y., Sun, Y., Pina-Crespo, J.C., Zhu, B., Masliah, E.,
Willnow, T.E.*, et al.* (2017). SORLA attenuates EphA4 signaling and amyloid beta-induced
neurodegeneration. J Exp Med *214*, 3669-3685.

Huang, T.Y., Zhao, Y., Li, X., Wang, X., Tseng, I.C., Thompson, R., Tu, S., Willnow, T.E., Zhang, Y.W.,
and Xu, H. (2016). SNX27 and SORLA Interact to Reduce Amyloidogenic Subcellular Distribution
and Processing of Amyloid Precursor Protein. The Journal of neuroscience : the official journal of
the Society for Neuroscience *36*, 7996-8011.

Huang, Z., Shimazu, K., Woo, N.H., Zang, K., Muller, U., Lu, B., and Reichardt, L.F. (2006). Distinct
roles of the beta 1-class integrins at the developing and the mature hippocampal excitatory
synapse. The Journal of neuroscience : the official journal of the Society for Neuroscience *26*,
11208-11219.

Hung, C., Tuck, E., Stubbs, V., van der Lee, S.J., Aalfs, C., van Spaendonk, R., Scheltens, P., Hardy,
J., Holstege, H., and Livesey, F.J. (2021). SORL1 deficiency in human excitatory neurons causes
APP-dependent defects in the endolysosome-autophagy network. Cell Rep *35*, 109259.

Hwang, J., Estick, C.M., Ikonne, U.S., Butler, D., Pait, M.C., Elliott, L.H., Ruiz, S., Smith, K.,
Rentschler, K.M., Mundell, C.*, et al.* (2019). The Role of Lysosomes in a Broad Disease-Modifying
Approach Evaluated across Transgenic Mouse Models of Alzheimer's Disease and Parkinson's
Disease and Models of Mild Cognitive Impairment. Int J Mol Sci *20*.

Karch, C.M., and Goate, A.M. (2015). Alzheimer's disease risk genes and mechanisms of disease
pathogenesis. Biol Psychiatry *77*, 43-51.

Klinger, S.C., Siupka, P., and Nielsen, M.S. (2015). Retromer-Mediated Trafficking of
Transmembrane Receptors and Transporters. Membranes (Basel) *5*, 288-306.

Knupp, A., Mishra, S., Martinez, R., Braggin, J.E., Szabo, M., Kinoshita, C., Hailey, D.W., Small, S.A.,
Jayadev, S., and Young, J.E. (2020). Depletion of the AD Risk Gene SORL1 Selectively Impairs
Neuronal Endosomal Traffic Independent of Amyloidogenic APP Processing. Cell Rep *31*, 107719.

Kwart, D., Gregg, A., Scheckel, C., Murphy, E.A., Paquet, D., Duffield, M., Fak, J., Olsen, O., Darnell,
R.B., and Tessier-Lavigne, M. (2019). A Large Panel of Isogenic APP and PSEN1 Mutant Human
iPSC Neurons Reveals Shared Endosomal Abnormalities Mediated by APP beta-CTFs, Not Abeta.
Neuron *104*, 256-270 e255.

Lambert, J.C., Ibrahim-Verbaas, C.A., Harold, D., Naj, A.C., Sims, R., Bellenguez, C., Jun, G.,
Destefano, A.L., Bis, J.C., Beecham, G.W.*, et al.* (2013). Meta-analysis of 74,046 individuals
identifies 11 new susceptibility loci for Alzheimer's disease. Nat Genet *45*, 1452-1458.

Leek, J.T., Johnson, W.E., Parker, H.S., Jaffe, A.E., and Storey, J.D. (2012). The sva package for
removing batch effects and other unwanted variation in high-throughput experiments.
Bioinformatics *28*, 882-883.

Levy, S., Sutton, G., Ng, P.C., Feuk, L., Halpern, A.L., Walenz, B.P., Axelrod, N., Huang, J., Kirkness,
E.F., Denisov, G.*, et al.* (2007). The diploid genome sequence of an individual human. PLoS Biol *5*,
e254.

Li, B., and Dewey, C.N. (2011). RSEM: accurate transcript quantification from RNA-Seq data with
or without a reference genome. BMC Bioinformatics *12*, 323.

Lorenzen, A., Samosh, J., Vandewark, K., Anborgh, P.H., Seah, C., Magalhaes, A.C., Cregan, S.P., Ferguson, S.S.G., and Pasternak, S.H. (2010). Rapid and Direct Transport of Cell Surface APP to the Lysosome defines a novel selective pathway. Molecular Brain *3*, 11.

Mallard, F., Antony, C., Tenza, D., Salamero, J., Goud, B., and Johannes, L. (1998). Direct pathway from early/recycling endosomes to the Golgi apparatus revealed through the study of shiga toxin B-fragment transport. J Cell Biol *143*, 973-990.

Manders, E., Verbeek, F., and Aten, J. (1993). Measurement of co-localization of objects in dual-colour confocal images. Journal of microscopy *169*, 375-382.

Marsh, E.W., Leopold, P.L., Jones, N.L., and Maxfield, F.R. (1995). Oligomerized transferrin receptors are selectively retained by a lumenal sorting signal in a long-lived endocytic recycling compartment. J Cell Biol *129*, 1509-1522.

Martin-Belmonte, A., Aguado, C., Alfaro-Ruiz, R., Itakura, M., Moreno-Martinez, A.E., de la Ossa, L., Molnar, E., Fukazawa, Y., and Lujan, R. (2020). Age-Dependent Shift of AMPA Receptors From Synapses to Intracellular Compartments in Alzheimer's Disease: Immunocytochemical Analysis of the CA1 Hippocampal Region in APP/PS1 Transgenic Mouse Model. Front Aging Neurosci *12*, 577996.

Marwaha, R., and Sharma, M. (2017). DQ-Red BSA Trafficking Assay in Cultured Cells to Assess Cargo Delivery to Lysosomes. Bio Protoc *7*.

Maxfield, F.R., and McGraw, T.E. (2004). Endocytic recycling. Nature reviews Molecular cell biology *5*, 121-132.

Mayle, K.M., Le, A.M., and Kamei, D.T. (2012). The intracellular trafficking pathway of transferrin. Biochim Biophys Acta *1820*, 264-281.

McQuin, C., Goodman, A., Chernyshev, V., Kamentsky, L., Cimini, B.A., Karhohs, K.W., Doan, M., Ding, L., Rafelski, S.M., Thirstrup, D.*, et al.* (2018). CellProfiler 3.0: Next-generation image processing for biology. PLOS Biology *16*, e2005970.

Offe, K., Dodson, S.E., Shoemaker, J.T., Fritz, J.J., Gearing, M., Levey, A.I., and Lah, J.J. (2006). The lipoprotein receptor LR11 regulates amyloid beta production and amyloid precursor protein traffic in endosomal compartments. The Journal of neuroscience : the official journal of the Society for Neuroscience *26*, 1596-1603.

Olav M. Andersen, N.B., Anne M. Landau, Gro Grunnet Pløen, Anne Mette G. Jensen, Giulia Monti, Benedicte Parm Ulhøi, Jens Randel Nyengaard, Kirsten Rosenmay Jacobsen, Margarita Melnikova Jørgensen, Ida E. Holm, Marianne L. Kristensen, Esben Søvsø Szocska Hansen, Charlotte E. Teunissen, Laura Breidenbach, Mathias Droescher, Ying Liu, Hanne Skovsgaard Pedersen, Henrik Callesen, Yonglun Luo, Lars Bolund, David J. Brooks, Christoffer Laustsen, Scott A. Small, Lars F. Mikkelsen, Charlotte B. Sørensen (2021). In vivo evidence that SORL1, encoding the endosomal recycling receptor SORLA, can function as a causal gene in Alzheimer's Disease. bioRxiv *doi: https://doi.org/10.1101/2021.07.13.452149*.

Ouellette, S.P., and Carabeo, R.A. (2010). A Functional Slow Recycling Pathway of Transferrin is Required for Growth of Chlamydia. Front Microbiol *1*, 112.

Park, M., Penick, E.C., Edwards, J.G., Kauer, J.A., and Ehlers, M.D. (2004). Recycling endosomes supply AMPA receptors for LTP. Science *305*, 1972-1975.

Patapoutian, A., and Reichardt, L.F. (2001). Trk receptors: mediators of neurotrophin action. Curr Opin Neurobiol *11*, 272-280.

Pietila, M., Sahgal, P., Peuhu, E., Jantti, N.Z., Paatero, I., Narva, E., Al-Akhrass, H., Lilja, J.,
Georgiadou, M., Andersen, O.M.*, et al.* (2019). SORLA regulates endosomal trafficking and
oncogenic fitness of HER2. Nat Commun *10*, 2340.
Qureshi, Y.H., Patel, V.M., Berman, D.E., Kothiya, M.J., Neufeld, J.L., Vardarajan, B., Tang, M.,
Reyes-Dumeyer, D., Lantigua, R., Medrano, M.*, et al.* (2018). An Alzheimer's Disease-Linked Loss-
of-Function CLN5 Variant Impairs Cathepsin D Maturation, Consistent with a Retromer Trafficking
Defect. Mol Cell Biol *38*.
Racoosin, E.L., and Swanson, J.A. (1993). Macropinosome maturation and fusion with tubular
lysosomes in macrophages. J Cell Biol *121*, 1011-1020.
Raghavan, N.S., Brickman, A.M., Andrews, H., Manly, J.J., Schupf, N., Lantigua, R., Wolock, C.J.,
Kamalakaran, S., Petrovski, S., Tosto, G.*, et al.* (2018). Whole-exome sequencing in 20,197
persons for rare variants in Alzheimer's disease. Ann Clin Transl Neurol *5*, 832-842.
Ramilowski, J.A., Goldberg, T., Harshbarger, J., Kloppmann, E., Lizio, M., Satagopam, V.P., Itoh,
M., Kawaji, H., Carninci, P., Rost, B.*, et al.* (2015). A draft network of ligand-receptor-mediated
multicellular signalling in human. Nat Commun *6*, 7866.
Rapaport, D., Lugassy, Y., Sprecher, E., and Horowitz, M. (2010). Loss of SNAP29 Impairs Endocytic
Recycling and Cell Motility. PLOS ONE *5*, e9759.
Ren, M., Xu, G., Zeng, J., De Lemos-Chiarandini, C., Adesnik, M., and Sabatini, D.D. (1998).
Hydrolysis of GTP on rab11 is required for the direct delivery of transferrin from the pericentriolar
recycling compartment to the cell surface but not from sorting endosomes. Proc Natl Acad Sci U
S A *95*, 6187-6192.
Rogaeva, E., Meng, Y., Lee, J.H., Gu, Y., Kawarai, T., Zou, F., Katayama, T., Baldwin, C.T., Cheng,
R., Hasegawa, H.*, et al.* (2007a). The neuronal sortilin-related receptor SORL1 is genetically
associated with Alzheimer disease. Nat Genet.
Rogaeva, E., Meng, Y., Lee, J.H., Gu, Y., Kawarai, T., Zou, F., Katayama, T., Baldwin, C.T., Cheng,
R., Hasegawa, H.*, et al.* (2007b). The neuronal sortilin-related receptor SORL1 is genetically
associated with Alzheimer disease. Nat Genet *39*, 168-177.
Rohe, M., Hartl, D., Fjorback, A.N., Klose, J., and Willnow, T.E. (2013). SORLA-mediated trafficking
of TrkB enhances the response of neurons to BDNF. PLoS One *8*, e72164.
Scheltens, P., De Strooper, B., Kivipelto, M., Holstege, H., Chetelat, G., Teunissen, C.E., Cummings,
J., and van der Flier, W.M. (2021). Alzheimer's disease. Lancet *397*, 1577-1590.
Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S.,
Rueden, C., Saalfeld, S., Schmid, B.*, et al.* (2012). Fiji: an open-source platform for biological-
image analysis. Nat Methods *9*, 676-682.
Schmidt, V., Sporbert, A., Rohe, M., Reimer, T., Rehm, A., Andersen, O.M., and Willnow, T.E.
(2007). SorLA/LR11 regulates processing of amyloid precursor protein via interaction with
adaptors GGA and PACS-1. J Biol Chem *282*, 32956-32964.
Seaman, M.N. (2004). Cargo-selective endosomal sorting for retrieval to the Golgi requires
retromer. J Cell Biol *165*, 111-122.
Seaman, M.N. (2012). The retromer complex - endosomal protein recycling and beyond. J Cell Sci
125, 4693-4702.
Selkoe, D.J. (2002). Alzheimer's disease is a synaptic failure. Science *298*, 789-791.

Shi, Y., Kirwan, P., Smith, J., Robinson, H.P., and Livesey, F.J. (2012). Human cerebral cortex
development from pluripotent stem cells to functional excitatory synapses. Nat Neurosci *15*, 477-
486, S471.
Simoes, S., Neufeld, J.L., Triana-Baltzer, G., Moughadam, S., Chen, E.I., Kothiya, M., Qureshi, Y.H.,
Patel, V., Honig, L.S., Kolb, H.*, et al.* (2020). Tau and other proteins found in Alzheimer's disease
spinal fluid are linked to retromer-mediated endosomal traffic in mice and humans. Science
translational medicine *12*.
Small, S.A., and Gandy, S. (2006). Sorting through the cell biology of Alzheimer's disease:
intracellular pathways to pathogenesis. Neuron *52*, 15-31.
Small, S.A., and Petsko, G.A. (2020). Endosomal recycling reconciles the Alzheimer's disease
paradox. Science translational medicine *12*.
Sonnichsen, B., De Renzis, S., Nielsen, E., Rietdorf, J., and Zerial, M. (2000). Distinct membrane
domains on endosomes in the recycling pathway visualized by multicolor imaging of Rab4, Rab5,
and Rab11. J Cell Biol *149*, 901-914.
Spoelgen, R., von Arnim, C.A., Thomas, A.V., Peltan, I.D., Koker, M., Deng, A., Irizarry, M.C.,
Andersen, O.M., Willnow, T.E., and Hyman, B.T. (2006). Interaction of the cytosolic domains of
sorLA/LR11 with the amyloid precursor protein (APP) and beta-secretase beta-site APP-cleaving
enzyme. The Journal of neuroscience : the official journal of the Society for Neuroscience *26*, 418-
428.
Sun, J., and Roy, S. (2018). The physical approximation of APP and BACE-1: A key event in
alzheimer's disease pathogenesis. Dev Neurobiol *78*, 340-347.
Temkin, P., Morishita, W., Goswami, D., Arendt, K., Chen, L., and Malenka, R. (2017). The
Retromer Supports AMPA Receptor Trafficking During LTP. Neuron *94*, 74-82 e75.
Thonberg, H., Chiang, H.H., Lilius, L., Forsell, C., Lindstrom, A.K., Johansson, C., Bjorkstrom, J.,
Thordardottir, S., Sleegers, K., Van Broeckhoven, C.*, et al.* (2017). Identification and description
of three families with familial Alzheimer disease that segregate variants in the SORL1 gene. Acta
Neuropathol Commun *5*, 43.
Wakabayashi, K., Narisawa-Saito, M., Iwakura, Y., Arai, T., Ikeda, K., Takahashi, H., and Nawa, H.
(1999). Phenotypic down-regulation of glutamate receptor subunit GluR1 in Alzheimer's disease.
Neurobiol Aging *20*, 287-295.
Wang, Y., Eng, D.G., Kaverina, N.V., Loretz, C.J., Koirala, A., Akilesh, S., Pippin, J.W., and
Shankland, S.J. (2020). Global transcriptomic changes occur in aged mouse podocytes. Kidney Int
98, 1160-1173.
Willnow, T.E., and Andersen, O.M. (2013). Sorting receptor SORLA--a trafficking path to avoid
Alzheimer disease. J Cell Sci *126*, 2751-2760.
Yap, C.C., Digilio, L., McMahon, L.P., Garcia, A.D.R., and Winckler, B. (2018). Degradation of
dendritic cargos requires Rab7-dependent transport to somatic lysosomes. Journal of Cell Biology
217, 3141-3159.
Yasuda, R.P., Ikonomovic, M.D., Sheffield, R., Rubin, R.T., Wolfe, B.B., and Armstrong, D.M.
(1995). Reduction of AMPA-selective glutamate receptor subunits in the entorhinal cortex of
patients with Alzheimer's disease pathology: a biochemical study. Brain Res *678*, 161-167.
Young, J.E., Boulanger-Weill, J., Williams, D.A., Woodruff, G., Buen, F., Revilla, A.C., Herrera, C.,
Israel, M.A., Yuan, S.H., Edland, S.D.*, et al.* (2015). Elucidating Molecular Phenotypes Caused by

the SORL1 Alzheimer's Disease Genetic Risk Factor Using Human Induced Pluripotent Stem Cells.
Cell stem cell *16*, 373-385.
Yuan, S.H., Martin, J., Elia, J., Flippin, J., Paramban, R.I., Hefferan, M.P., Vidal, J.G., Mu, Y., Killian,
R.L., Israel, M.A*., et al.* (2011). Cell-surface marker signatures for the isolation of neural stem
cells, glia and neurons derived from human pluripotent stem cells. PLoS One *6*, e17540.
Zigdon, H., Meshcheriakova, A., Farfel-Becker, T., Volpert, G., Sabanay, H., and Futerman, A.H.
(2017). Altered lysosome distribution is an early neuropathological event in neurological forms
of Gaucher disease. FEBS Lett *591*, 774-783.
823

Figure legends

Figure 1. Loss of *SORL1* expression leads to increased TRKB and GLUA1 localization in early endosomes. Representative immunofluorescent images of WT and *SORL1*KO neurons showing increased colocalization of **(a)** TRKB (green) and **(b)** GLUA1 (green) with EEA1 (red). All neurons were immunolabeled with MAP2 (far-red) and counterstained with DAPI (blue). Scale bar: 10μm. In all cases, quantification of colocalization was represented as Mander's correlation co-efficient (MCC). 10-20 images were analyzed per genotype. Two isogenic clones of each genotype were used in all experiments. Data represented as mean ± SD. Normally distributed data was analyzed using parametric two-tailed unpaired t test and non-normally distributed data was analyzed by non-parametric Mann Whitney test. Significance was defined as a value of *p < 0.05, **p < 0.01, ***p < 0.001, and ****p < 0.0001.

Figure 2. Loss of *SORL1* expression impairs trafficking to late endosomes and lysosomes.

a) *SORL1*KO neurons show reduced lysosomal proteolytic activity as measured by DQ Red BSA. Representative immunofluorescent images of WT and *SORL1*KO neurons showing double immunolabeling for MAP2 (green) and DQ Red BSA (red). Scale bar: 10μm Quantification of fluorescence intensity of DQ Red BSA using ImageJ software. **(b-f)** *SORL1*KO neurons show reduced colocalization of cargo with late endosomes and lysosomes. Representative immunofluorescent images of WT and *SORL1*KO neurons showing reduced colocalization of **(b)** TRKB (green) and **(c)** GLUA1 (green) with Rab7 positive late endosomes (red) in *SORL1*KO neurons. Representative immunofluorescent images of WT and *SORL1*KO neurons showing reduced colocalization of **(d)** APP (green) and **(e)** TRKB (green) and **(f)** GLUA1 (green) with LAMP1 positive lysosomes (red) in *SORL1*KO neurons. Scale bar: 10μm. In all cases, quantification of colocalization was represented as Mander's correlation co-efficient (MCC). 10-20 images were analyzed per genotype. Two isogenic clones of each genotype were used in all experiments. Data represented as mean ± SD. Normally distributed data was analyzed using parametric

two-tailed unpaired t test and non-normally distributed data was analyzed by non-parametric Mann Whitney test. Significance was defined as a value of *p < 0.05, **p < 0.01, ***p < 0.001, and ****p < 0.0001.

Figure 3. Loss of *SORL1* impacts the cell surface recycling pathway

a) *SORL1*KO neurons show slower rate of transferrin recycling. Quantification of fluorescence intensity of intracellular transferrin at different time points after treating cells with Alexa Fluor 647-conjugated transferrin for 15 mins, using ImageJ software. Data represented as percent of time 0 fluorescence intensity. **(b)** *SORL1*KO neurons show larger recycling endosomes. Representative immunofluorescent images of WT and *SORL1*KO neurons labeled with antibodies for MAP2 (red) and Rab11 (green). Nuclei were counterstained with DAPI (blue). Scale bar: 5μm. Quantification of size of Rab11 labeled recycling endosomes using CellProfiler software. Representative immunofluorescent images of WT and *SORL1*KO neurons showing increased colocalization of **(c)** APP (green), **(d)** TRKB (green) and **(e)** GLUA1 (green) with Rab11 positive recycling endosomes (red) in *SORL1*KO neurons. Scale bar:10mm In all cases, quantification of colocalization was represented as Mander's correlation co-efficient (MCC). 10-20 images were analyzed per genotype. Two isogenic clones of each genotype were used in all experiments. Data represented as mean ± SD. Normally distributed data was analyzed using parametric two-tailed unpaired t test and non-normally distributed data was analyzed by non-parametric Mann Whitney test. Significance was defined as a value of *p < 0.05, **p < 0.01, ***p < 0.001, and ****p < 0.0001.

Figure 4. Loss of *SORL1* expression impairs recycling to the cell surface.

a-b) *SORL1*KO neurons show reduced cell surface expression of APP **(a)** and GLUA1 **(b)**. Representative immunofluorescent images of WT and *SORL1*KO neurons labeled with antibodies for APP **(a)** (red) and GLUA1 **(b)** (red). Scale bar: 5μm. Intensity of APP and GLUA1 measured using ImageJ software. Data is presented as a ratio of surface intensity to total intensity. Two isogenic clones of each genotype were used in all experiments.

Data represented as mean ± SD. Normally distributed data was analyzed using parametric two-tailed unpaired t test. Significance was defined as a value of *p < 0.05, **p < 0.01, ***p < 0.001, and ****p < 0.0001.

Figure 5. Overexpression of *SORL1* enhances endosomal recycling.

a) *SORL1*OE neurons show faster rate of transferrin recycling. Quantification of fluorescence intensity of intracellular transferrin at different time points after treating cells with Alexa Fluor 647-conjugated transferrin for 15 mins, using ImageJ software. Data represented as percent of time 0 fluorescence intensity. **(b)** *SORL1*OE neurons show reduced size of recycling endosomes. Representative immunofluorescent images of WT and *SORL1*OE neurons labeled with Rab11 (green) and MAP2 (red) showing smaller Rab11 positive recycling endosomes in *SORL1*OE neurons. Nuclei counterstained with DAPI (blue). Quantification of Rab11+ recycling endosome size performed using Cell Profiler software and represented as area of Rab11+ vesicles. Scale bar: 5μm. Representative immunofluorescent images of WT and *SORL1*OE neurons showing increased colocalization of **(c)** APP (green), **(d)** TRKB (green) and **(e)** GLUA1 (green) with Rab11 (red) positive recycling endosomes. *SORL1*OE neurons and controls have endogenous GFP expression due to the piggybac vector system. GFP fluorescence is pseudo-colored (Far-red) and was used to outline cell bodies. Quantification of colocalization with Rab11 represented as Mander's Correlation Co-efficient (MCC). Scale bar: 10μm. Representative immunofluorescent images of WT and *SORL1*OE neurons showing increased cell surface expression of **(f)** APP (red) and **(g)** GLUA1 (red) in *SORL1*OE neurons. Scale bar: 5μm Fluorescence intensity of APP and GLUA1 measured using ImageJ software. Data is presented as a ratio of surface intensity to total intensity. Nuclei counterstained with DAPI. Two isogenic clones of each genotype were used in all experiments. Data represented as mean ± SD. Normally distributed data was analyzed using parametric two-tailed unpaired t test. Significance defined as a value of *p < 0.05, **p < 0.01, ***p < 0.001, and ****p < 0.0001.

Figure 6. Analysis of bulk RNA-sequencing data indicates alterations in cell surface and extracellular trafficking, receptor-ligand and channel activity.

Gene ontology analysis of DEGs in WT and *SORL1*KO neurons. Shown here are the top upregulated **(a)** and downregulated **(b)** molecular function terms in *SORL1*KO neurons. GO annotation terms are listed on the y-axis, adjusted p-value is shown on the x-axis. **(c)** Ligand-receptor network changes in *SORL1*KO neurons, identified if genes are more than 1.5-fold increased or decreased in *SORL1*KO neurons with an adjusted p-value less than 0.05. Circles denote ligands, squares denote receptors, blue indicates genes expressed significantly lower in *SORL1*KO neurons, red indicates genes expressed significantly higher in *SORL1*KO neurons. Arrows point from ligand to receptor, denoting receptor-ligand interactions. Black arrows denote consistent expression changes between ligand and receptor, indicating that both genes in the pair are either upregulated or downregulated. Gray arrows denote inconsistent changes between ligand and receptor. Clusters impacted by cell surface recycling (β-integrins, ephrins) or neurotrophic signaling (FGF/FGFR, NGF/NGFR) are indicated by green dotted arrows.

Supplementary figures

Supplementary figure 1

a) *SORL1*KO neurons show larger lysosome size and increased lysosome number. Representative immunofluorescent images of WT and *SORL1*KO neurons labeled with LAMP1 (green) and MAP2 (red) showing increased LAMP1 positive vesicle size in *SORL1*KO neurons. Quantification of LAMP1 size and number was performed using Cell Profiler software. LAMP1 size is represented as area of LAMP1 positive vesicles, and LAMP1 number is represented as number of LAMP1 positive vesicles per square micron of cell area. Scale bar: 5μm **(b)** *SORL1*KO neurons show no change in colocalization of lysosomes with the lysosomal enzyme Cathepsin D. Representative immunofluorescent images of WT and *SORL1*KO neurons labeled with antibodies for LAMP1 (green), Cathepsin D (red) and MAP2 (Far-red) showing no alteration in colocalization of Cathepsin-D with LAMP1 in *SORL1*KO neurons. Nuclei counterstained with DAPI (blue). Scale bar: 10μm Quantification of colocalization of LAMP1 with Cathepsin-D represented as Mander's correlation coefficient (MCC). 10-20 images

were analyzed per genotype. Two isogenic clones of each genotype were used in all experiments. Data represented as mean ± SD. Significance was determined using two-tailed unpaired Student's t-test and defined as a value of $*p < 0.05$, $**p < 0.01$, $***p < 0.001$, and $****p < 0.0001$.

Supplemental figure 2

a) *SORL1*OE neurons show no change in lysosomal proteolytic activity as measured using DQ Red BSA. Representative immunofluorescent images of WT and *SORL1*OE neurons showing double immunolabeling for MAP2 (green) and DQ Red BSA (red). Scale bar: 10μm. Quantification of fluorescence intensity of DQ Red BSA using ImageJ software. Representative immunofluorescent images of WT and *SORL1*OE neurons showing increased colocalization of APP, TRKB and GLUA1 (green) with Rab7 (red) **(b-d)** and LAMP1 (red) **(e-g)** in *SORL1*OE neurons. *SORL1*OE neurons and controls have endogenous GFP expression due to the piggybac vector system. GFP fluorescence is pseudo-colored (Far-red) and was used to outline cell bodies. Scale bar: 10μm. Nuclei are counterstained with DAPI (blue). In all cases, quantification of colocalization was represented as Mander's correlation co-efficient (MCC). 10-20 images were analyzed per genotype. Two isogenic clones of each genotype were used in all experiments. Data represented as mean ± SD. Significance was determined using two-tailed unpaired Student's t-test and was defined as a value of $*p < 0.05$, $**p < 0.01$, $***p < 0.001$, and $****p < 0.0001$.

Supplemental figure 3

a) Principle component analysis (PCA) plot of all RNAseq samples using all expressed genes. Samples are color coded by differentiation batch. Triangles represent WT samples, circles represent *SORL1*KO. Genotype accounts for the highest variance (37%, PC1, x-axis). **b)** Volcano plot. Log2 fold change between *SORL1*KO and WT is shown along the x-axis. Statistical significance is shown along the y-axis, and is measured by adjusted p-value. Genes upregulated in *SORL1*KO neurons are shown by red circles, genes downregulated in *SORL1*KO neurons are shown by blue circles. We observed 6643 DEGs, with 2819 upregulated and 3824 downregulated.

There are significantly more down regulated genes than upregulated genes (p<2.2e-16). Grey circles represent genes that are not significantly differentially expressed.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

A

B

C

Ligand upregulated in *SORL1* KO
Ligand downregulated in *SORL1* KO
Receptor upregulated in *SORL1* KO
Receptor downregulated in *SORL1* KO
Both ligand and receptor up/down in *SORL1* KO
Ligand and receptor inconsistent expression change