**The Alzheimer's gene *SORL1* is a regulator of endosomal traffic and recycling in human**

**neurons**

Swati Mishra[1,2*], Allison Knupp[1,2*], Marcell P. Szabo[1,2], Charles A. Williams[1,2], Chizuru

Kinoshita[1,2], Dale W. Hailey[1,2], Yuliang Wang[2,3], and Jessica E. Young[1,2,#]

[1]Department of Laboratory Medicine and Pathology

[2]Institute for Stem Cell and Regenerative Medicine

[3]School of Computer Science and Engineering

University of Washington Seattle WA, 98195, USA

[#]Correspondence: jeyoung@uw.edu

[*]Equal Contribution

Author emails: swatim2@uw.edu, knuppa@uw.edu, szabom@uw.edu, cwil@uw.edu

ckino@uw.edu, dhailey@uw.edu, yuliangw@uw.edu, jeyoung@uw.edu

**ABSTRACT**

**Background** Loss of the Sortilin-related receptor 1 (*SORL1*) gene seems to act as a causal event

for Alzheimer's disease (AD). Recent studies have established that loss of *SORL1*, as well as

mutations in autosomal dominant AD genes *APP* and *PSEN1/2*, pathogenically converge by

swelling early endosomes, AD's cytopathological hallmark. Acting together with the retromer

trafficking complex, *SORL1* has been shown to regulate the recycling of the amyloid precursor

protein (APP) out of the endosome, contributing to endosomal swelling and to APP

misprocessing. We hypothesized that *SORL1* plays a broader role in neuronal endosomal

recycling and used human induced pluripotent stem cell derived neurons (hiPSC-Ns) to test this

hypothesis. We examined endosomal recycling of three transmembrane proteins linked to AD

pathophysiology: APP, the BDNF receptor Tropomyosin-related kinase B (TRKB), and the

glutamate receptor subunit AMPA1 (GLUA1).

**Methods** We used isogenic hiPSCs engineered to have *SORL1* depleted or to have enhanced

*SORL1* expression. We differentiated neurons from these cell lines and mapped the trafficking of

APP, TRKB and GLUA1 within the endosomal network using confocal microscopy. We also

performed cell surface recycling and lysosomal degradation assays to assess the functionality of

the endosomal network in both *SORL1* depleted and overexpressing neurons. Finally, we

analyzed alterations in gene expression in *SORL1* depleted neurons using RNA-sequencing.

**Results** We find that as with APP, endosomal trafficking of GLUA1 and TRKB is impaired by loss

of *SORL1*. We show that trafficking of all three cargo to late endosomes and lysosomes is affected

by manipulating SORL1 expression. We also show that depletion of *SORL1* significantly impacts

the endosomal recycling pathway for APP and GLUA1 at the level of the recycling endosome and

trafficking to the cell surface. This has a functional effect on neuronal activity as shown by multi-

electrode array (MEA). Conversely, increased *SORL1* expression enhances endosomal recycling

for APP and GLUA1. Our unbiased transcriptomic data further support *SORL1*'s role in

endosomal recycling.  We observe altered expression networks that regulate cell surface

trafficking and neurotrophic signaling in SORL1 depleted neurons.

**Conclusion** Collectively, and together with other recent observations, these findings suggest that

*SORL1* is a broad regulator of retromer-dependent endosomal recycling in neurons, a conclusion

that has both pathogenic and therapeutic implications for Alzheimer's disease.

48                                                    **KEY WORDS**

49      Human induced pluripotent stem cells, neurons, Alzheimer's disease, endosomal network,

50      *SORL1*.

51                                                    **BACKGROUND**

Alzheimer's disease (AD) is a progressive neurodegenerative disorder and the most common

cause of dementia. The underlying contributors to AD pathology encompass several biological

pathways, including endosomal function, amyloid precursor protein (APP) processing, immune

function, synaptic function, and lipid metabolism(Karch and Goate, 2015). Among these,

endosomal dysfunction in neurons is emerging as a potential causal mechanism(Small and

Petsko, 2020). Mutations in the amyloid precursor protein (*APP*) and the two presenilins (*PSEN1*

and *PSEN2*) lead to early-onset autosomal dominant AD. When these mutations are modelled in

human neurons and other systems they cause endosomal swelling, indicative of traffic jams, a

phenotype that is a cytopathological hallmark of AD(Cataldo et al., 2000; Choi et al., 2013; Kwart

et al., 2019). Recent genetic studies have identified a fourth gene, the trafficking receptor 'sortilin

related receptor 1' (*SORL1*), which, when harboring frame-shift mutations leading to premature

stop codons, is described as causal for AD(Holstege et al., 2017; Raghavan et al., 2018;

Scheltens et al., 2021). Interestingly, *SORL1* is also linked to the more common, late-onset form

of AD(Lambert et al., 2013; Rogaeva et al., 2007b) and its expression is lost in sporadic AD brains

(Dodson et al., 2006; Thonberg et al., 2017). When modelled in human neurons, *SORL1* depletion

phenocopies *APP* and *PSEN* mutations by causing endosomal swelling(Hung et al., 2021; Knupp

et al., 2020).

The *SORL1* gene codes for the protein SORLA, an endosomal sorting protein that is also an

adaptor molecule for the retromer trafficking complex (Fjorback et al., 2012; Rogaeva et al.,

2007a; Small and Gandy, 2006). Retromer recycles cargo out of the early endosome, either from

the endosome to the trans-Golgi network or, with greater importance for neurons, back to the cell

surface(Fjorback et al., 2012; Seaman, 2012). To date, the best evidence for *SORL1*'s role in

retromer-dependent    endosomal    recycling    comes    from    studies    investigating    APP

trafficking(Fjorback et al., 2012; Schmidt et al., 2007; Willnow and Andersen, 2013). Our previous

work demonstrated that *SORL1* depletion retains APP in early endosomes, which may contribute

to endosomal swelling by blocking recycling(Knupp et al., 2020).

Retromer-dependent trafficking in neurons, however, also recycles cargo other than APP.  For

example, retromer is required for the normal recycling of glutamate receptors, a trafficking event

that mediates synaptic plasticity and synaptic health, and this dependency occurs independent of

retromer's role in APP recycling(Park et al., 2004; Temkin et al., 2017). Neurotrophin receptors

are also trafficked through the endosomal system, in a retromer-dependent manner, and are

important for synaptic health(Klinger et al., 2015; Patapoutian and Reichardt, 2001; Rohe et al.,

2013).

Here we used human induced pluripotent stem cell derived-neurons (hiPSC-Ns) to test the

hypothesis that *SORL1* plays a broader role in neuronal endosomal recycling. We use our

previously described *SORL1*-depleted hiPSC lines to generate hiPSC-Ns, which model the loss

of *SORL1* expression that occurs in AD(Knupp et al., 2020). Furthermore, we used previously

established cell lines engineered to overexpress *SORL1* 2-3-fold over wild-type levels(Young et

al., 2015) to test the effects of enhanced *SORL1* expression in hiPSC-Ns on these trafficking

pathways.  Importantly,  all  cell  lines  are  isogenic.  We  map  the  trafficking  effects  these

manipulations have on three specific receptors, APP, the GLUA1 subunit of the AMPA receptor,

and neurotrophin receptor TRKB, all of which are implicated in AD (Devi and Ohno, 2015; Dewar

et al., 1991; Ginsberg et al., 2019; Martin-Belmonte et al., 2020; Wakabayashi et al., 1999;

Yasuda et al., 1995).

Finally, we performed RNA-sequencing on the *SORL1* depleted cell lines to explore an unbiased

transcriptomics  analysis  induced  by  *SORL1*  depletion.   The  results  generally  confirmed  our

hypothesis, showing that *SORL1* is a broad regulator of endosomal recycling in neurons, a

conclusion that has both pathogenic and therapeutic implications.

105                                                      **METHODS**

**Cell lines**

***Cell lines generated by CRIPSR/Cas9 gene editing technology***

The generation of the cell lines used in this paper is described in our previously published work

(Knupp et al., 2020) and consists of four clones: Two wild-type clones, designated clone A6 and

clone A7, and two *SORL1*KO clones, designated clone E1 and clone E4. Cell lines were

generated from our previously published and characterized CV background human induced

pluripotent stem cell line(Young et al., 2015). This cell line is male and has a APOE ε3/ε4

genotype(Levy et al., 2007). All four clones were shown to have normal karyotypes and are

routinely tested for mycoplasma (MycoAlert). The clones used in the experiments in this work are

listed in the figure legends.

**CRISPR/Cas9 gRNA, ssODN, and Primer Sequences** gRNA: ATTGAACGACATGAACCCTC

ssODN:

GGGAATTGATCCCTATGACAAACCAAATACCATCTACATTGAACGACATGAACCCTCTGGC

TACTCCACGTCTTCCGA  AGTACAGATTTCTTCCAGTCCCGGGAAAACCAGGAAG  Forward

primer: ctctatcctgagtcaaggagtaac Reverse primer: ccttccaattcctgtgtatgc PCR amplifies 458 bp

sequence. These sequences have been previously published in (Knupp et al., 2020)

***SORL1 overexpression cell lines*** Isogenic cell lines with overexpression of SORL1 were

generated as previously described(Young et al., 2015). These lines are generated from the CV

parental line, the same parental line as the *SORL1*KO cell lines were made from. Briefly, stable

integration of SORL1 cDNA into the genome was achieved by using piggybac transposon system

(Systems Biosciences). Vector alone (WT) or vector with SORL1 cDNA (*SORL1*OE) constructs

were introduced into iPSCs by electroporation and stable cell lines were selected with puromycin

(2ug/ml) treatment. We obtained one SORL1OE stable cell line and one vector alone stable cell

line. For all overexpression experiments *SORL1*OE cells were compared to the vector alone

controls.

**hiPSC Neuronal Differentiation**

hiPSCs were differentiated to neurons using dual-SMAD inhibition(Chambers et al., 2009; Shi et

al., 2012). Briefly, hiPSCs were plated on Matrigel coated 6-well plates at a density of 3.5 million

cells per well and fed with Basal Neural Maintenance Media (1:1 DMEM/F12 + glutamine

media/neurobasal media, 0.5% N2 supplement, 1% B27 supplement, 0.5% GlutaMax, 0.5%

insulin-transferrin-selenium, 0.5% NEAA, 0.2% β-mercaptoethanol; Gibco, Waltham, MA) +

10mM SB-431542 + 0.5mM LDN-193189 (Biogems, Westlake Village, CA). Cells were fed daily

for seven days. On day eight, cells were incubated with Versene, gently dissociated using cell

scrapers, and passaged at a ratio of 1:3. On day nine, media was switched to Basal Neural

Maintenance Media and fed daily. On day 13, media was switched to Basal Neural Maintenance

Media with 20 ng/mL FGF (R&D Systems, Minneapolis, MN) and fed daily. On day sixteen, cells

were passaged again at a ratio of 1:3. Cells were fed until approximately day twenty-three. At this

time, cells were FACS sorted to obtain the CD184/CD24 positive, CD44/CD271 negative neural

precursor cell (NPC) population. Following sorting, NPCs were expanded for neural

differentiation. For cortical neuronal differentiation, NPCs were plated out in 10cm cell culture

dishes at a density of 6 million cells/10cm plate. After 24 hours, cells were switched to Neural

Differentiation media (DMEM-F12 + glutamine, 0.5% N2 supplement, 1% B27 supplement, 0.5%

GlutaMax) + 0.02ug/mL brain-derived neurotrophic factor (PeproTech, Rocky Hill, NJ) +

0.02ug/mL glial-cell-derived neurotrophic factor (PeproTech) + 0.5mM dbcAMP (Sigma Aldrich,

St Louis, MO). Media was refreshed twice a week for three weeks. After three weeks, neurons

were selected for CD184/CD44/CD271 negative population by MACS sorting and plated for

experiments. The data presented in this study represent 2-3 neuronal differentiations.

**Purification of Neurons**

Following three weeks of differentiation, neurons were dissociated with accutase and

resuspended in Magnet Activated Cell Sorting (MACS) buffer (PBS + 0.5% bovine serum albumin

[Sigma Aldrich, St Louis, MO] + 2mM ethylenediaminetetraacetic acid [Thermo Fisher Scientific,

Waltham, MA]). Following a modification of (Yuan et al., 2011), cells were incubated with PE-

conjugated mouse anti-Human CD44 and mouse anti-Human CD184 antibodies (BD

Biosciences, San Jose, CA) at a concentration of 5µl/10 million cells. Following antibody

incubation, cells were washed with MACS buffer and incubated with anti-PE magnetic beads (BD

Biosciences, San Jose, CA) at a concentration of 25µl/10 million cells. Bead-antibody complexes

were pulled down using a rare earth magnet, supernatants were selected, washed, and plated at

an appropriate density.

**DQ Red BSA assay for visualization of lysosomal degradation**

Lysosomal proteolytic degradation was evaluated using DQ Red BSA (#D-12051; Thermo Fisher

Scientific), a fluorogenic substrate for lysosomal proteases, that generates fluorescence only

when enzymatically cleaved in intracellular lysosomal compartments. hiPSC-derived neurons

were seeded at a density of 400,000 cells/well of a matrigel coated 48 well plate. After 24 hours,

cells were washed once with DPBS, treated with complete media containing either 10µg/ml DQ

Red BSA or vehicle (PBS) and incubated for 6 hours or 24 hours (Davis et al., 2021; Romano et

al., 2021; Tian et al., 2015)at 37°C in a 5% $CO_2$ incubator as described in(Marwaha and Sharma,

2017). At the end of 6 or 24 hours, cells were washed with PBS, fixed with 4% PFA and

immunocytochemistry was performed as described in methods. Cells were imaged using a Leica

SP8 confocal microscope and all image processing was completed with ImageJ software. Cell

bodies were identified by MAP2 labeling, and fluorescence intensity of DQ Red BSA was

measured in regions of the images containing the MAP2 label.

**Immunocytochemistry**

For immunocytochemistry, cells were fixed with 4% PFA for 20 minutes. Fixed cells were washed

three times with PBST (PBS with 0.05% tween 20), permeabilized with Triton X-100 in PBS for

15 minutes, washed twice again with PBST, blocked with 5% BSA in PBS at room temperature

for 1h and incubated with appropriate primary antibodies overnight at 4°C. The next day, cells

were incubated with appropriate secondary antibodies and 1µg/ml DAPI for 1 hour at RT, washed

three times with PBST and mounted on glass slides with Prolong Gold Antifade mountant

(#P36930; Thermo Fisher Scientific).

**Colocalization analysis**

To investigate colocalization with endo-lysosomal compartments, hiPSC-derived neurons were

labeled with markers specific for each intra-cellular compartment (EEA1 for early endosomes,

Rab7 for late endosomes, LAMP1 for lysosomes and Rab11 for recycling endosomes) using

immunocytochemistry. A minimum of 10 fields of confocal z-stack images were captured under

blinded conditions using a Yokogawa W1 spinning disk confocal microscope (Nikon) and a 100X

plan apochromat oil immersion objective. Median filtering was used to remove noise from images

and Otsu thresholding was applied to all images. Colocalization was quantified using the JACOP

plugin(Bolte and Cordelieres, 2006) in Image J software(Schindelin et al., 2012) and presented

as Mander's correlation coefficient.(Dunn et al., 2011; Manders et al., 1993). For all imaging

experiments the data was analyzed in a blinded manner.

**Cell Surface Staining:** Cell surface expression of GLUA1 and APP was determined using

immunocytochemistry and confocal microscopy. To label proteins at the cell surface, cells were

fixed with 4% PFA, washed and treated with primary and secondary antibodies as described in

the 'immunocytochemistry' section of methods. Permeabilization with 0.1% Triton X-100 was not

performed for this experiment. To label total protein levels, cells were fixed with 4% PFA, washed,

permeabilized with 0.1% Triton X-100 and treated with primary and secondary antibodies as

described in the 'immunocytochemistry' section of methods. Analysis of fluorescence intensity

was done using Image J software. Cell surface expression was represented as ratio of

fluorescence intensity measured under non-permeabilized conditions and fluorescence intensity

measured under permeabilized conditions. For all imaging experiments the data was analyzed in

a blinded manner.

**Multielectrode (MEA) assay**

hiPSC-derived neural progenitor cells were differentiated into neurons and neurons were purified

as previously described in methods. Purified neurons were mixed with unpurified neurons in a

ratio of 5:1 and this mixture was plated onto a matrigel coated 48 well MEA plate (Axion

Biosystems; # M768-tMEA-48W) at a cell density of 8000 cells/ul (total number of cells/well =

50,000). MEA-plated neurons were initially cultured in neural differentiation media. Media was

gradually switched to BrainPhys media (Stem cell technologies; # 05790) by replacing half of a

well's media twice a week. BrainPhys media was supplemented with B27, N2, BDNF, GDNF, and

db-cAMP.

**Multielectrode (MEA) analysis**

Electrical signals from neurons in the MEA plates were recorded twice a week using Axion

Biosystems Maestro Pro system. Signals were recorded at a sampling frequency of 12.5 kHz with

a 3 kHz Kaiser Window low pass filter and 200 Hz high pass filter. Spikes were detected using

Axion Axis Navigator recording software with the adaptive threshold method. Recordings were

analyzed using the Axion Neural Metric Tool. Firing rate data were limited to active electrodes

that detected a minimum of five spikes a minute. Firing rate data from all active electrodes in a

well were averaged for plotting and statistical testing.

**Antibodies**

The following primary antibodies were used: Early endosome antigen 1 (EEA1) at 1:500 (#610456; BD Biosciences); amyloid precursor protein (APP) at 1:500 (#ab32136; Abcam); microtubule-associated protein 2 (MAP2) at 1:1000 (ab92434; Abcam); Ras-related protein Rab-7a (Rab7) at 1:1000 (ab50533; Abcam); Ras-related protein Rab-11 (Rab11) at 1:250 (#610656; BD Biosciences); Lysosome associated membrane protein-1 (LAMP1) at 1:250 (#sc 2011; Santa Cruz); Tropomyosin receptor kinase B (TRKB; # ab18987; abcam) at 1:1000 ,GLUA1(# MAB2263; Millipore sigma) at 1:500 andVPS35(Abcam; #ab97545) at 1:500. DAPI was used at a final concentration of 0.1µg/ml (Alfa Aesar).

**Transferrin recycling assay**

To measure recycling pathway function, we utilized transferrin recycling assay as previously described(Rapaport et al., 2010). Purified neurons were seeded at 400,000 cells/well of a 24 well plate containing matrigel coated 12 mm glass coverslip/well. After 5 DIV, cells were washed once with DMEM-F12 medium and incubated with starving medium (DMEM-F12 medium+25mM HEPES+ 0.5% BSA) for 30 minutes at 37°C in a 5% $CO_2$ incubator to remove any residual transferrin. Thereafter, cells were pulsed with either 100µg/ml transferrin from human serum conjugated with Alexa Fluor™ 647(#T23366; Thermo Fisher Scientific) or vehicle (PBS) in 'starving medium'. At the end of 10 mins, cells were washed twice with ice-cold PBS to remove any external transferrin and stop internalization of transferrin and washed once with acid stripping buffer (25mM citric acid+24.5mM sodium citrate+280mM sucrose+0.01mM Deferoxamine) to remove any membrane bound transferrin. Next, cells were either fixed in 4% PFA or 'Chase medium' (DMEM-F12+50µM Deferoxamine+20mM HEPES+500µg/ml Holo-transferrin) was added for different time points. Immunocytochemistry was done using MAP2 antibody to label neurons, confocal images were captured using Leica SP8 confocal microscope under blinded conditions. Fluorescence intensity of transferrin was measured using ImageJ software. Recycling

function was presented as transferrin fluorescence intensity as a percentage of the fluorescence

intensity measured at time zero.

**Measurement of lysosome and recycling endosome size**

Immunocytochemistry using antibodies for LAMP1 and MAP2 or RAB11 and MAP2 was

performed as described above. Using a Leica SP8 confocal microscope with an apochromatic

63X oil immersion lens, z-stack images were obtained under blinded conditions. For the LAMP1

analysis, 17-34 fields were analyzed for a total of **45 – 76** cells analyzed. For the RAB11 analysis,

15-30 fields were analyzed for a total of **59 – 124** cells analyzed. Vesicle size measurements were

performed using Cell Profiler software as previously described (Knupp et al., 2020) (McQuin et

al., 2018). Briefly, the vesicle channel was masked using the MAP2 channel and automated

segmentation algorithms were used to identify individual puncta. The pixel area of each puncta

was measured and is presented as mean area of all puncta per field.

**Statistical Analysis**

For all imaging experiments, data was collected and analyzed in a blinded manner. Data was

assessed for significance using parametric two-tailed unpaired Student's t-tests or two-way

ANOVA tests. Data is represented as mean $\pm$ standard deviation to show the spread of the data.

Significance was defined as a value of $p \leq 0.05$. All statistical analysis was completed using

GraphPad Prism software. Statistical details of individual experiments, including biological and

technical replicate information, can be found in the figure legends. All of the raw statistical data

for the experiments in this paper, including means, difference between the means $\pm$ SEM, and

95% confidence intervals are presented in Supplemental Table 1.

**RNA- sequencing analysis**

_RNA Extraction_

RNA was collected from 3 separate differentiations including a combination of two WT clones and

two *SORL1*KO clones. Each sample includes 2-3 technical replicates. RNA was collected from 2

million purified neurons for each sample. Purification of total RNA was completed using the

PureLink RNA Mini Kit (Thermo Fisher 12183018A). Assessment of purified RNA was completed

using a NanoDrop. Final RNA quantification was completed using the Quant-iT RNA assay

(Invitrogen) and RNA integrity analysis was completed using a fragment analyzer (Advanced

Analytical).

*Library Prep and Sequencing*

Library preparation was completed using the TruSeq Stranded mRNA kit (Illumina RS-122-2103)

per manufacturer instructions. Sequencing was performed on a NovaSeq 6000 instrument.

*Data analysis*

Raw reads were aligned to GRCh38 with reference transcriptome GENCODE release 29 using

STAR v2.6.1d(Dobin et al., 2013). Gene-level expression quantification is generated by RSEM

v1.3.1(Li and Dewey, 2011). Genes with fewer than 20 normalized reads across all samples were

omitted from further analysis. We did observe variation in the transcriptome based on

differentiation (Supplemental Figure 3A), however this was corrected for using the sva

package(Leek et al., 2012).

To identify differentially expressed genes (DEGs), we used DESeq(Anders and Huber, 2010).

Briefly, we fit two models: a null model where gene expression only depends on batch effects

(i.e., differentiation), and an alternative model where gene expression depends on both genotype

(*SORL1*KO vs. WT) and batch effects. Chi-squared tests were performed to compare both fits,

and we declare a gene as differentially expressed only when the alternative model fits the

expression data better. DEGs are defined as genes with false discovery rate less than 0.05 and

fold change greater than 1.5. The top gene ontology package(Alexa et al., 2006) and the SynGO

synaptic gene ontology annotations(Koopmans et al., 2019) were used to identify GO terms that

were enriched. GO terms were tested according to the Fisher's exact test. Finally, we mapped

DEGs onto receptor-ligand interaction diagrams generated by Ramilowski et al(Ramilowski et al.,

2015) using the igraph plugin(Csardi G, 2006). To compare the amount of down-regulated vs. up-

regulated genes we used a 2-sample test for equality of proportions with continuity correction in

R.

318                                    **RESULTS**

*SORL1* **depletion increases neuronal cargo localization in early endosomes.**

Using CRISPR/Cas9 genome editing techniques, we previously generated hiPSC-derived

neurons (hiPSC-Ns) deficient in *SORL1* expression due to indels introduced in exon 6. We

demonstrated that loss of *SORL1* expression in these neurons leads to enlarged early endosomes

and an increased colocalization of APP within early endosomes, indicative of endosomal traffic

jams(Knupp et al., 2020). We utilized these same cell lines (hereafter referred to as *SORL1*KO

and their guide-matched isogenic wild-type clones referred to as WT) to examine localization of

the BDNF receptor TRKB and the GLUA1 subunit of the neuronal AMPA receptor. TRKB has

been shown to bind to SORLA and this interaction mediates trafficking of TRKB to synaptic

plasma membranes(Rohe et al., 2013). GLUA1 is trafficked via the retromer complex, of which

SORLA is an adaptor protein(Fjorback et al., 2012; Temkin et al., 2017) and both of these cargo

are important in maintaining healthy neuronal function. Because we previously observed an

increase in APP localization in early endosomes, resulting in a decrease in localization in

downstream vesicles such as Ras-related protein (Rab)7+ late endosomes with *SORL1*

depletion(Knupp et al., 2020), we performed an immunocytochemical analysis of both TRKB and

GLUA1 localization with the early endosome marker EEA1. Similar to our previous observations

for APP, we documented significantly increased localization of both TRKB (Figure 1A) and GLUA1

(Figure 1B) in early endosomes in *SORL1*KOneurons as compared to isogenic WT control

neurons. Accumulation of neuronal cargo in early endosomes is indicative of endosomal traffic

jams, which are thought to impact the transit of cellular cargo through other arms of the endo-

lysosomal network. Due to SORLA's role as an adaptor protein for the retromer complex, we also

examined whether *SORL1* depletion led to changes in retromer subunit localization. We observed

that VPS35, a core subunit of the retromer cargo recognition complex is also mis-localized to

early endosomes, similar to what we observed for APP, TRKB, and GLUR1 (Supplemental Figure

1).

**Modulating *SORL1* expression impacts cargo trafficking throughout the endo-lysosomal**

**system**

The early endosome serves as a hub in which internalized cargo can be retrogradely transported

to the trans-Golgi, recycled back to the cell surface or degraded as endosomes mature into late

endosomes and lysosomes(Mayle et al., 2012). We have previously observed that APP

localization within the trans-Golgi network was decreased in *SORL1*KO neurons(Knupp et al.,

2020). Here we tested whether trafficking to the degradative arm of the endo-lysosomal network

was affected in our *SORL1*KO neurons. Trafficking of substrates out of the early endosome to

late endosomes and, subsequently, lysosomes is important for protein degradation and SORLA

has been previously implicated in promoting Aβ degradation via lysosomes(Caglayan et al.,

2014). We treated *SORL1* deficient neurons with DQ Red BSA, a proteolysis sensitive fluorogenic

substrate that generates fluorescence only when enzymatically cleaved in intracellular lysosomal

compartments. Since substrate degradation primarily occurs in lysosomes, altered fluorescence

intensity of this reagent is a readout of altered lysosomal degradation(Marwaha and Sharma,

2017). We treated neurons with DQ Red BSA for 6 and 24 hours and analyzed fluorescence

intensity using confocal microscopy. Consistent with loss of *SORL1* leading to endosomal traffic

jams, we observed a significant reduction of DQ Red BSA fluorescence intensity at both time

points in *SORL1*KO neurons compared to isogenic WT controls (Figure 2A). We next performed

immunocytochemical staining to quantify the colocalization of our selected neuronal cargo with

Rab7, a marker of late endosomes, and LAMP1 (Lysosomal Associated Membrane Protein 1), a

lysosome marker. We show a significant decrease in co-localization of TRKB (Figure 2B) and

GLUA1 (Figure 2C) with Rab7. This result is consistent with our previous observation for APP

(Knupp et al., 2020). We analyzed colocalization of these cargo with LAMP1 and we observed a

significant decrease with APP (Figure 2D) and TRKB (Figure 2E) and a trend of a decrease with

GLUA1 (Figure 2F). These data indicate some fluidity in the network but suggest that trafficking

of APP, TRKB and GLUA1 to late endosomes/lysosomes is all decreased by *SORL1*KO, although

GLUA1 may be more likely to be trafficked to cell surface pathways or utilizes other adaptor

proteins for late endosome to lysosomal trafficking. These changes in localization are not due to

changes in expression of cargo. We have previously shown that APP levels do not change in

*SORL1*KO neurons(Knupp et al., 2020) and also show here that protein expression of TRKB and

GLUA1 are not different (Supplemental Figure 3).

We next utilized previously generated cell lines that overexpress *SORL1* cDNA using the

piggybac transposon system(Young et al., 2015) to test whether increased *SORL1* expression

may enhance the trafficking pathways that are impaired in the *SORL1*KO neurons. Importantly,

the *SORL1* overexpressing (*SORL1*OE) cell line and control were generated in the same genetic

background as our *SORL1*KO and isogenic WT cell lines.   Interestingly, while there was no effect

of *SORL1* overexpression on DQ Red BSA fluorescence at the earlier time point (6 hours), we

did see a significant enhancement of DQ RED BSA trafficking at the 24-hour time point (Figure

3A). In accordance, we observed significantly increased localization of our studied cargo with late

endosomal and lysosomal markers in the *SORL1*OE neurons (Figure 3 B-G).

Lysosome size can influence lysosome function and is altered in AD(de Araujo et al., 2020; Hwang

et al., 2019). Similarly, location and number of lysosomes within neurons can alter degradative

activity  (Cheng et al., 2018; Farfel-Becker et al., 2019; Gowrishankar et al., 2015; Yap et al.,

2018) and in some cases, altered lysosomal distribution may represent an early neuropathological

defect(Zigdon et al., 2017).Recently, loss of *SORL1* in hiPSC neurons was shown to contribute

to lysosome dysfunction as indicated by both increased lysosome size and number as well as

decreased cathepsin-D activity(Hung et al., 2021). Therefore, we first analyzed LAMP1-

immunopositive puncta and also documented a significant increase in lysosome size and number

in our *SORL1*KO neurons (Supplemental Figure 2A). Interestingly, although the number of

lysosomes marked by LAMP1 puncta is increased in *SORL1KO* neurons, we did not observe a

significant change in LAMP1 protein expression (Supplemental Figure 3). This may be partially

explained by differences in autophagy in SORL1 KO neurons which we did not examine in this

study but that has been previously reported(Hung et al., 2021) and further underscores the

dynamic complexity of the endo-lysosomal network.

Next, we analyzed co-localization of Cathepsin-D and LAMP1 to determine if loss of *SORL1*

expression leads to altered Cathepsin-D trafficking in neurons. Retromer trafficking is required to

deliver one of the most abundant lysosomal proteases, pro-cathepsin D, to lysosomes via the

mannose-6-phosphate receptor (M6PR)(Qureshi et al., 2018; Seaman, 2004). The SORLA

protein has GGA domains similar to that of M6PR(Spoelgen et al., 2006), and mis-trafficking of

Cathepsin-D to lysosomes could affect the maturation and degradative capacity of these

organelles. Therefore, we analyzed co-localization of Cathepsin-D and LAMP1 to determine if

loss of *SORL1* expression leads to altered Cathepsin-D trafficking in neurons. However, we did

not observe a change in Cathepsin-D colocalization between WT and *SORL*1KO (Supplemental

Figure 2B).

Taken together, our data suggest that *SORL1* loss in neurons reduces trafficking of cargo out of

the early endosome to the late endosome and lysosome, contributing to lysosome stress as

evidenced by an increase in size and number in these conditions while the decreased cathepsin-

D activity observed upon *SORL1* loss(Hung et al., 2021) may not be due to impairment of

lysosomal trafficking of the enzyme.

**Loss of *SORL1* impacts the endosomal recycling pathway**

Another route out of the early endosome is via the endocytic recycling complex (ERC) which can

send cargo either to the cell surface or to the trans-Golgi network(Grant and Donaldson, 2009;

Mallard et al., 1998; Marsh et al., 1995; Maxfield and McGraw, 2004). To directly examine if

*SORL1* expression alters recycling function, we performed a transferrin recycling assay using

confocal microscopy. Transferrin can be recycled via a fast pathway within approximately 5-10

minutes after being internalized or via a slower pathway involving the ERC over longer periods of

time(Ouellette and Carabeo, 2010; Sonnichsen et al., 2000). We examined the fluorescence

intensity of Alexa Fluor 647-conjugated transferrin over a 40-minute time course in WT and

*SORL1*KO neurons and observed that a higher percentage of intracellular fluorescent transferrin

persisted in *SORL1*KO neurons at both early and later time points as compared to WT neurons,

indicating reduced recycling pathway function (Figure 4A). Cargo destined for the cell surface can

transit the ERC via Rab11+ recycling endosomes (Ren et al., 1998). Altered size of recycling

endosomes can be indicative of dysfunctional recycling of cargo through these compartments.

We tested whether loss of *SORL1* expression affected the size of Rab11+ recycling endosomes.

Interestingly, we observed a significant increase in the size of Rab11+ recycling endosomes,

although there was no change in Rab11 protein expression in the *SORL1*KO neurons (Figure 4B,

Supplemental Figure 3), suggesting that this endosomal compartment is also under stress. To

test if increased size is due to abnormal cargo trafficking through recycling endosomes, we

assessed colocalization of APP, TRKB and GLUA1 with Rab11 and observed increased co-

localization of all three cargo with Rab11+ structures in *SORL1*KO neurons compared to WT

neurons (Figure 4C-E). Together, these data demonstrate that loss of *SORL1* impacts neuronal

recycling endosome pathways by causing traffic jams in the recycling endosomes, similar to the

effect that *SORL1* loss has on early endosomes.

**SORL1 depletion reduces cell surface levels of cargo.**

Together, our data indicate that *SORL1*KO neurons have impaired cargo recycling with increased

retention of cargo in recycling endosomes. These observations led us to test whether this cargo

was indeed trafficked to the cell surface. A portion of APP has been shown to return to the cell

surface via recycling endosomes(Das et al., 2016) and SORLA can interact with the sorting nexin

SNX27 to return APP to the cell surface(Das et al., 2016; Huang et al., 2016), although in that

study the exact compartment was not described. Furthermore, recycling endosomes are the

source for AMPA receptors during long-term potentiation(Park et al., 2004). We therefore

examined cell surface levels of APP and GLUA1 using immunofluorescence and confocal

microscopy. We documented a significant decrease in cell surface staining of both APP (Figure

5A) and GLUA1 (Figure 5B) in *SORL1*KO neurons as compared to WT, consistent with our

hypothesis that SORLA is involved in regulating traffic from recycling endosomes. Due to the

importance of GLUA1 in the formation of functional excitatory synapses, we next analyzed

neuronal activity by culturing *SORL1*KO and isogenic WT neurons on multi-electrode array (MEA)

plates. Interestingly, we observed an early increase in the mean firing rate of *SORL1*KO neurons

at an early time point (27 days post-plating) however neuronal firing in *SORL1*KO neurons was

significantly reduced at a later time point (66 days post-plating) (Figure 5C) suggesting that

synaptic activity may be partially impaired in these cells.

**SORL1 overexpression enhances endosomal recycling**

Defects in cell surface recycling have severe consequences in neurons, especially as these

processes are necessary for healthy neuronal function and enhancing pathways that promote

endosomal recycling in neurons may be beneficial. Using our *SORL1*OE neurons we analyzed

recycling function using the transferrin recycling assay and observed that *SORL1*OE neurons

showed significantly faster transferrin recycling (Figure 6A). We next tested whether colocalization

of cargo with recycling endosomes and cell surface recycling was altered between *SORL1*OE

and WT neurons.  Interestingly, the size of Rab11+ recycling endosomes was significantly smaller

in *SORL1*OE neurons (Figure 6B) possibly indicating that increased *SORL1* expression is clearing

cargo more rapidly from this compartment. We observed a significant increase in localization of

cargo with Rab11+ recycling endosomes (Figure 6C-E). While this result was initially surprising,

as we also saw increased colocalization with Rab11+recycling endosomes in our *SORL1*KO

neurons (Figure 4C-E), we further documented a significant increase of APP and GLUA1 on the

cell surface compared to WT neurons with only endogenous *SORL1* expression (Figure 6F and

6G), as opposed to decreased APP and GLUA1 localization on the cell surface in *SORL1*KO

neurons (Figure 5A and 5B).  These results suggest that cell surface trafficking via a Rab11

pathway is enhanced by increased *SORL1* expression and that a crucial action of SORLA is the

trafficking out of recycling endosomes. Thus, our data support a critical role for SORLA for

trafficking cargo from recycling endosomes to the cell surface. In addition, we show for the first

time that SORLA levels may regulate cell surface recycling of AMPA receptor subunits in human

neurons.

**SORL1 depletion affects gene expression**.

To determine a more global effect of chronic *SORL1* loss in human neurons, we performed bulk

RNA sequencing of *SORL1*KO neurons compared to WT neurons. Interestingly, we observed that

there were significantly more down-regulated genes in *SORL1*KO neurons than upregulated ones

(Supplemental Figure 4B). While none of the cargo we explicitly studied in this work was

differentially expressed, GO analysis showed that the top downregulated molecular function

pathways in the *SORL1*KO cells were related to receptor-ligand activity and extracellular matrix

organization (Figure 7A). The top upregulated molecular function pathways were related to ion

channel activity (Figure 7B). To understand these data in the context of an integrated network,

we used an analysis method that infers ligand receptor interactions from bulk RNA-seq

data(Ramilowski et al., 2015; Wang et al., 2020). We observed several nodes of altered receptor-

ligand interactions that indicate altered cell surface recycling and neurotrophic activity (Figure

7C). These include alterations in β-integrin signaling, which is consistent with previous work

showing reduced β-integrin on the cell surface in *SORL1*KO cancer cells(Pietila et al., 2019), and

altered interactions in ephrins/ephrin receptors, also corroborating previous work implicating

*SORL1* expression in ephrin signaling and synapse regulation(Huang et al., 2017). Our analysis

also showed nodes with alterations in nerve growth factor/nerve growth factor receptor

(NGF/NGFR) and fibroblast growth factor/fibroblast growth factor receptor (FGF/FGFR) signaling,

indicating alterations in neurotrophin and growth factor signaling and suggesting that the presence

of endosomal traffic jams in *SORL1*KO neurons may ultimately impact multiple pathways

important for neuronal health and development. Because we documented a decrease of GLUA1

puncta on the cell surface and observed altered neuronal activity on MEAs, we further mined our

RNA-seq data for genes involved in synaptic function. Interestingly when we performed

enrichment analysis, we observed that pathways associated with synaptic function were

upregulated in *SORL1*KO neurons (Supplemental Figure 5).

510                                    **DISCUSSION**

Trafficking through the endo-lysosomal network regulates intra-cellular location of proteins,

dictating their homeostasis, function and influence on cellular physiology. Functional studies by

our group and others document endosomal abnormalities in hiPSC-derived neuronal models of

AD(Hung et al., 2021; Knupp et al., 2020; Kwart et al., 2019). Emerging from this evidence is the

role of *SORL1* as an endosomal gene that plays essential roles in mediating cargo trafficking.

Recent work has implicated *SORL1* as AD's fourth causal gene(Olav M. Andersen, 2021;

Scheltens et al., 2021), and of these genes it is the only one linked to the common late-onset form

of the disease. Understanding *SORL1*'s function is paramount for understanding AD's pathogenic

mechanisms and for potential therapeutic interventions.

Acting as an adaptor molecule of the retromer trafficking complex, SORLA has already been

pathogenically linked to AD by its role in recycling APP out of endosomes(Andersen et al., 2005;

Herskowitz et al., 2012; Offe et al., 2006). This current work and our previous study(Knupp et al.,

2020) shows that *SORL1* depletion leads to increased APP localization in early and recycling

endosomes. By lengthening the residence time of APP in these endosomal compartments,

accelerated amyloidogenic cleavage of APP occurs due to the close proximity of APP and

BACE1(Sun and Roy, 2018). Indeed, loss of *SORL1* leads to the accumulation of Aβ peptides,

an antecedent of 'amyloid pathology' (Andersen et al., 2005; Knupp et al., 2020; Rogaeva et al.,

2007b). We hypothesized that loss of *SORL1* in neurons would impact other cargo important for

healthy neuronal function. To test this hypothesis, in addition to APP, we examined localization

of the neurotrophin receptor TRKB and the GLUA1 subunit of the AMPA receptor and also

observed that these proteins are increased in early endosomes (Figure. 1). These cargo link to

another key pathology of AD: neurodegeneration, a slowly progressive process that begins with

synaptic dysfunction characterized by glutamate receptor loss, which then progresses to synaptic

loss before ultimately, over years, leading to widespread neuronal cell death (Selkoe, 2002).

The early endosome is considered the central station in the sorting and trafficking of cargo

throughout the many stations of the endo-lysosomal system.  While the early endosome is the

station that is affected first and foremost in AD, it is not surprising that a primary dysfunction in

this central station will secondarily influence trafficking throughout the system including the

recycling and degradative pathways. Indeed, SORLA was shown to traffic Aβ to lysosomes in

neuroblastoma cells, a function that is impaired by an AD-associated variant(Caglayan et al.,

2014). Our work presented in this study, along with other recent work(Hung et al., 2021) also

supports a role for *SORL1* in lysosomal trafficking in neurons.  We observe a decrease in the pH-

sensitive fluorogenic substrate DQ RED BSA in our *SORL1*KO neurons and decreased

localization of APP, TRKB, and GLUA1 in late endosomes and lysosomes in *SORL1*KO neurons

and that this is reversed in *SORL1*OE neurons (Figure. 2, Figure. 3). We interpret our functional

and colocalization data to suggest that loss of *SORL1* expression mainly affects trafficking of

cargo to lysosomes, but our data does not rule out a role of SORLA in neuronal lysosome function.

Hung et al., reported decreased Cathepsin-D activity in *SORL1* deficient neurons, suggesting that

*SORL1* loss directly impacts lysosome function(Hung et al., 2021). While we did not observe a

difference in Cathepsin-D localization to lysosomes, it is important to note that that LAMP1 only

partially colocalizes with Cathepsin-D in neurons (Cheng et al., 2018). Furthermore, the loss of

proteolytic activity evidenced by decreased intensity of DQ Red BSA in *SORL1*KO neurons may

not be completely due to reduced trafficking to lysosomes but could be a result of abnormal

lysosomal function as DQ Red BSA is internalized by a process called macropinocytosis wherein

macropinosomes can be directly trafficked to lysosomes (Hamasaki et al., 2004; Lorenzen et al.,

2010; Racoosin and Swanson, 1993). Interestingly, while we did not observe an effect on DQ Red

BSA at a shorter time point in *SORL1*OE neurons, we did see increased fluorescence of this

reagent at a 24-hour time-point (Figure. 3), suggesting that overtime, increased SORL1

expression impacts lysosome trafficking and/or function in cortical neurons.

Our experiments also point to *SORL1*'s role in cell surface recycling. By using a prototypical cargo,

transferrin, we demonstrate a reciprocal role between loss and enhancement of *SORL1*

expression in cell surface recycling. Specifically, we show that *SORL1*KO neurons have defects

in transferrin recycling at both early (10-minute) and late (40-minute) time points while *SORL1*OE

neurons have faster recycling at these time points (Figure 4, Figure 6). This suggests that SORLA

functions in both fast and slow endosomal recycling. Our data further implicates the recycling

pathway by showing that modulation of SORLA expression affects recycling endosome size and

the amount of cargo (APP, TRKB and GLUA1) localized to recycling endosomes (Figure. 4,

Figure. 6).

Endocytic recycling comprises returning cargo, primarily membrane proteins, to the cell

surface(Cullen and Steinberg, 2018). We studied a canonical SORLA cargo, APP,  and show that

loss of *SORL1* expression results in reduced cell surface APP while enhanced expression

increases cell surface APP (Figure 5). These results corroborate previous work showing that

*SORL1* and SNX27 work to return APP to the cell surface(Das et al., 2016; Huang et al., 2016)

in a human model.

We also show that *SORL1* plays a role in recycling glutamate receptors (Figure. 5) This finding is

critically important as recent work indicates that in mouse cortical neurons, SORLA interacts with

a neuronal specific retromer subunit, VPS26b, to promote recycling of glutamate

receptors(Simoes S., 2021). In our cortical neurons when *SORL1* is depleted, there is a reduction

of GLUA1 subunits on the cell surface and this may result in synaptic impairment. MEA data

comparing *SORL1*KO and isogenic WT neurons shows alterations in weighted mean firing rate

as neurons mature (Figure 5). Interestingly, we observed an increase in neuronal firing at an early

time point and a significant decrease in firing at a later time point. *SORL1*KO mice live to

adulthood but have been described to have some deficits in learning and memory that may also

be age dependent(Glerup et al., 2013; Hojland et al., 2018). Some of these alterations could be

explained by compensatory expression changes ion channels or synaptic genes induced by

chronic loss of *SORL1* during the course of neuronal differentiation from pluripotent stem cells.

Our RNA-seq data does show up-regulation of ion channels and channel activity (Figure. 7).

Interestingly, when we further interrogated our RNA-seq data for pathways enriched in synaptic

function using the SynGo database (Koopmans 2019), we observed an upregulation of

differentially expressed genes in synaptic pathways (Supplemental Fig. 5). This data suggests

that *SORL1*KO neurons may attempt to compensate for altered trafficking of synaptic receptors

by upregulating gene expression.

Our unbiased transcriptomic screen further supported that neurotrophic signaling and cell

surface recycling pathways are impacted by *SORL1* deficiency (Figure 7). The SORLA

cytoplasmic tail has been shown to translocate to the nucleus and activate transcription in a

reporter gene assay(Bohm et al., 2006). Despite this, distinct genes regulated by SORLA are

not known. Rather than looking for a direct effect on gene regulation, our goal for the analysis

was to determine the global effect of *SORL1* loss or overexpression on neuronal networks.

Indeed, our data does not show that the specific cargo proteins described here are differentially

expressed. However, the analysis does indicate that loss of *SORL1* in human neurons impacts

cell surface networks, including receptor ligand interactions in neurotrophic and growth factor

pathways, β-integrin signaling, and ephrin signaling. This corroborates previous work and the

altered networks we observe impact neuronal health, axonal guidance, and synapse

formation(Huang et al., 2017; Huang et al., 2006; Pietila et al., 2019).

Importantly, enhancing SORL1 expression improves cell surface trafficking of GLUA1 (Figure 6).

Trafficking of glutamate receptors is an event that is critical for preventing synaptic dysfunction

and synaptic loss, thus our results link *SORL1* to AD's early-stage neurodegenerative process.

Since retromer-dependent glutamate receptor recycling has been shown to occur independent of

APP(Temkin et al., 2017), our previous and current results suggest that *SORL1* mutations can,

at least in principle, drive two key AD pathologies, amyloid pathology and synaptic pathology,

through parallel mechanisms(Small and Petsko, 2020).

We summarize our findings in Figure. 8. However, our study has certain limitations. For example,

our results encompass only one human genome. Future studies will benefit from looking at

*SORL1* deficiency or overexpression in multiple human genetic backgrounds. Furthermore, in this

work we are describing purely neuronal phenotypes although *SORL1* is expressed in other CNS

cells. Future work looking at cell- autonomous and non-cell autonomous mechanisms of *SORL1*

depletion or overexpression in human glial or brain organoid models will be informative.

**CONCLUSIONS**

In this work, we report that *SORL1* depletion affects endosomal trafficking by retaining cargo in

early and recycling endosomes and impacts cell surface recycling and lysosomal trafficking of

neuronal cargo. In particular, we demonstrate that *SORL1* expression in neurons affects cell

surface localization of GLUA1, a phenotype that may ultimately impact synaptic dysfunction and

neurodegeneration in AD. Interestingly, increasing *SORL1* expression enhances endosomal

recycling and increases cell surface GLUA1. While the secondary downstream effects induced

by *SORL1* depletion in the endo-lysosomal system are interesting and likely relevant to AD's

ultimate pathogenesis, from a therapeutic perspective it is best to target *SORL1*'s primary defect,

which seems to localize to the endosomal recycling pathway. Interestingly, recent biomarker

studies suggest that defects in retromer-dependent endosomal recycling occur in a majority of

patients with 'sporadic' AD(Simoes et al., 2020), suggesting that the observed *SORL1*-induced

defects may generalize across early and late onset forms of the disorder. Collectively, our results

support the conclusion that *SORL1*, and the retromer-dependent pathway in which it functions, is

a valid therapeutic target and interventions directed at this pathway may ameliorate endosomal

recycling defects that seem to act as, at least, one primary driver of AD.

**ABBREVIATIONS**

AD: Alzheimer's Disease; SORL1: Sortilin-related receptor 1; SORLA: Sortilin-related receptor

with A-type repeats;  APP: Amyloid Precursor Protein;  TRKB: Tropomyosin Related Kinase B;

GLUA1: Glutamate receptor subunit AMPA1; AMPA: α-amino-3-hydroxy-5-methyl-4-

isoxazolepropionic acid; OE: Overexpression;  KO: Knock-out; WT: Wild-type; LAMP1:

Lysosome-associated membrane glycoprotein 1; M6PR: mannose-6-phosphate receptor; GGA:

Golgi-Localized γ-Ear-Containing ARF-Binding; Rab: Ras-related protein.

646                                                         **DECLARATIONS**

**Ethics Approval and Consent to Participate**

Not applicable

**Consent for Publication**

Not applicable

**Availability of Data and Material**

All raw and processed RNA-seq data has been deposited at the NCBI Gene Expression Omnibus

(GSE180793). The data sets used and/or analyzed during the current study are available from

the corresponding author on reasonable request.

**Availability of Supporting Material**

All supporting data and material in the current study are available from the corresponding author

on reasonable request.

**Competing Interests**

The authors declare no competing interests

**Acknowledgements**

We thank Dr. Harald Frankowski, Dr. Yoshito Kinoshita, Ms. Shannon Rose, Ms. Eden

Cruickshank and all members of the Young Laboratory for critical discussions and feedback

during the preparation of this manuscript. We also thank Dr. Scott A. Small and Dr. Gregory A.

Petsko for critical comments, discussions and feedback on this work. We would like to

acknowledge the UW SLU Cell Analysis Facility and the Garvey Imaging Core at the UW Institute

for Stem Cell and Regenerative Medicine. This work was supported by a NIH grant

(R01AG062148) and a BrightFocus Foundation grant (A2018656S) to J.E.Y., a Biogen

Sponsored Research Agreement to J.E.Y., a Retromer Therapeutic Sponsored Research

Agreement to J.E.Y. and NIH traning grant (T32AG052354) to A.K. and a generous gift from the

Ellison Foundation (to UW).

**Author Contributions:**

Conceptualization: J.E.Y., S.M, A.K. Microscopy analysis: S.M, A.K., and D.W.H. RNA-seq

analysis and bioinformatics: Y.W and A.K. Methodology: S.M., A.K., M.S., C.A.W., C.K. Writing-

Original Draft: S.M., A.K. and J.E.Y. Writing-Reviewing and editing: J.E.Y., S.M., A.K. Funding

Acquisition: J.E.Y. and A.K. Supervision: J.E.Y. All authors read and approved the final

manuscript.

678                     **REFERENCES**

Alexa, A., Rahnenfuhrer, J., and Lengauer, T. (2006). Improved scoring of functional groups from
gene expression data by decorrelating GO graph structure. Bioinformatics *22*, 1600-1607.
Anders, S., and Huber, W. (2010). Differential expression analysis for sequence count data.
Genome Biol *11*, R106.
Andersen, O.M., Reiche, J., Schmidt, V., Gotthardt, M., Spoelgen, R., Behlke, J., von Arnim, C.A.,
Breiderhoff, T., Jansen, P., Wu, X.*, et al.* (2005). Neuronal sorting protein-related receptor
sorLA/LR11 regulates processing of the amyloid precursor protein. Proc Natl Acad Sci U S A *102*,
13461-13466.
Bohm, C., Seibel, N.M., Henkel, B., Steiner, H., Haass, C., and Hampe, W. (2006). SorLA signaling
by regulated intramembrane proteolysis. J Biol Chem *281*, 14547-14553.
Bolte, S., and Cordelieres, F.P. (2006). A guided tour into subcellular colocalization analysis in light
microscopy. J Microsc *224*, 213-232.
Caglayan, S., Takagi-Niidome, S., Liao, F., Carlo, A.S., Schmidt, V., Burgert, T., Kitago, Y.,
Fuchtbauer, E.M., Fuchtbauer, A., Holtzman, D.M.*, et al.* (2014). Lysosomal sorting of amyloid-
beta by the SORLA receptor is impaired by a familial Alzheimer's disease mutation. Science
translational medicine *6*, 223ra220.
Cataldo, A.M., Peterhoff, C.M., Troncoso, J.C., Gomez-Isla, T., Hyman, B.T., and Nixon, R.A. (2000).
Endocytic pathway abnormalities precede amyloid beta deposition in sporadic Alzheimer's
disease and Down syndrome: differential effects of APOE genotype and presenilin mutations. Am
J Pathol *157*, 277-286.
Chambers, S.M., Fasano, C.A., Papapetrou, E.P., Tomishima, M., Sadelain, M., and Studer, L.
(2009). Highly efficient neural conversion of human ES and iPS cells by dual inhibition of SMAD
signaling. Nat Biotechnol *27*, 275-280.

Cheng, X.-T., Xie, Y.-X., Zhou, B., Huang, N., Farfel-Becker, T., and Sheng, Z.-H. (2018). Characterization of LAMP1-labeled nondegradative lysosomal and endocytic compartments in neurons. J Cell Biol *217*, 3127-3139.

Choi, J.H., Kaur, G., Mazzella, M.J., Morales-Corraliza, J., Levy, E., and Mathews, P.M. (2013). Early endosomal abnormalities and cholinergic neuron degeneration in amyloid-beta protein precursor transgenic mice. J Alzheimers Dis *34*, 691-700.

Csardi G, N.T. (2006). The igraph software package for complex network research. InterJournal, Complex systems *1695*.

Cullen, P.J., and Steinberg, F. (2018). To degrade or not to degrade: mechanisms and significance of endocytic recycling. Nat Rev Mol Cell Biol *19*, 679-696.

Das, U., Wang, L., Ganguly, A., Saikia, J.M., Wagner, S.L., Koo, E.H., and Roy, S. (2016). Visualizing APP and BACE-1 approximation in neurons yields insight into the amyloidogenic pathway. Nat Neurosci *19*, 55-64.

Davis, S.E., Roth, J.R., Aljabi, Q., Hakim, A.R., Savell, K.E., Day, J.J., and Arrant, A.E. (2021). Delivering progranulin to neuronal lysosomes protects against excitotoxicity. Journal of Biological Chemistry *297*.

de Araujo, M.E.G., Liebscher, G., Hess, M.W., and Huber, L.A. (2020). Lysosomal size matters. Traffic *21*, 60-75.

Devi, L., and Ohno, M. (2015). TrkB reduction exacerbates Alzheimer's disease-like signaling aberrations and memory deficits without affecting β-amyloidosis in 5XFAD mice. Translational psychiatry *5*, e562-e562.

Dewar, D., Chalmers, D.T., Graham, D.I., and McCulloch, J. (1991). Glutamate metabotropic and AMPA binding sites are reduced in Alzheimer's disease: an autoradiographic study of the hippocampus. Brain Res *553*, 58-64.

Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson, M., and Gingeras, T.R. (2013). STAR: ultrafast universal RNA-seq aligner. Bioinformatics *29*, 15-21.

Dodson, S.E., Gearing, M., Lippa, C.F., Montine, T.J., Levey, A.I., and Lah, J.J. (2006). LR11/SorLA expression is reduced in sporadic Alzheimer disease but not in familial Alzheimer disease. J Neuropathol Exp Neurol *65*, 866-872.

Dunn, K.W., Kamocka, M.M., and McDonald, J.H. (2011). A practical guide to evaluating colocalization in biological microscopy. Am J Physiol Cell Physiol *300*, C723-C742.

Farfel-Becker, T., Roney, J.C., Cheng, X.T., Li, S., Cuddy, S.R., and Sheng, Z.H. (2019). Neuronal Soma-Derived Degradative Lysosomes Are Continuously Delivered to Distal Axons to Maintain Local Degradation Capacity. Cell Rep *28*, 51-64.e54.

Fjorback, A.W., Seaman, M., Gustafsen, C., Mehmedbasic, A., Gokool, S., Wu, C., Militz, D., Schmidt, V., Madsen, P., Nyengaard, J.R.*, et al.* (2012). Retromer binds the FANSHY sorting motif in SorLA to regulate amyloid precursor protein sorting and processing. The Journal of neuroscience : the official journal of the Society for Neuroscience *32*, 1467-1480.

Ginsberg, S.D., Malek-Ahmadi, M.H., Alldred, M.J., Chen, Y., Chen, K., Chao, M.V., Counts, S.E., and Mufson, E.J. (2019). Brain-derived neurotrophic factor (BDNF) and TrkB hippocampal gene expression are putative predictors of neuritic plaque and neurofibrillary tangle pathology. Neurobiology of Disease *132*, 104540.

Glerup, S., Lume, M., Olsen, D., Nyengaard, J.R., Vaegter, C.B., Gustafsen, C., Christensen, E.I.,
Kjolby, M., Hay-Schmidt, A., Bender, D*., et al.* (2013). SorLA controls neurotrophic activity by
sorting of GDNF and its receptors GFRalpha1 and RET. Cell Rep *3*, 186-199.

Gowrishankar, S., Yuan, P., Wu, Y., Schrag, M., Paradise, S., Grutzendler, J., De Camilli, P., and
Ferguson, S.M. (2015). Massive accumulation of luminal protease-deficient axonal lysosomes at
Alzheimer's disease amyloid plaques. Proceedings of the National Academy of Sciences *112*,
E3699-E3708.

Grant, B.D., and Donaldson, J.G. (2009). Pathways and mechanisms of endocytic recycling. Nat
Rev Mol Cell Biol *10*, 597-608.

Hamasaki, M., Araki, N., and Hatae, T. (2004). Association of early endosomal autoantigen 1 with
macropinocytosis in EGF-stimulated A431 cells. Anat Rec A Discov Mol Cell Evol Biol *277*, 298-
306.

Herskowitz, J.H., Offe, K., Deshpande, A., Kahn, R.A., Levey, A.I., and Lah, J.J. (2012). GGA1-
mediated endocytic traffic of LR11/SorLA alters APP intracellular distribution and amyloid-beta
production. Mol Biol Cell *23*, 2645-2657.

Hojland, A., Richner, M., Molgaard, S., Dieu, R.S., Eskelund, A., Nykjaer, A., Nyengaard, J.R.,
Lykkesfeldt, J., Glerup, S., and Nielsen, M.S. (2018). Biochemical and cognitive effects of
docosahexaenoic acid differ in a developmental and SorLA dependent manner. Behav Brain Res
*348*, 90-100.

Holstege, H., van der Lee, S.J., Hulsman, M., Wong, T.H., van Rooij, J.G., Weiss, M.,
Louwersheimer, E., Wolters, F.J., Amin, N., Uitterlinden, A.G*., et al.* (2017). Characterization of
pathogenic SORL1 genetic variants for association with Alzheimer's disease: a clinical
interpretation strategy. Eur J Hum Genet *25*, 973-981.

Huang, T.Y., Zhao, Y., Jiang, L.L., Li, X., Liu, Y., Sun, Y., Pina-Crespo, J.C., Zhu, B., Masliah, E.,
Willnow, T.E*., et al.* (2017). SORLA attenuates EphA4 signaling and amyloid beta-induced
neurodegeneration. J Exp Med *214*, 3669-3685.

Huang, T.Y., Zhao, Y., Li, X., Wang, X., Tseng, I.C., Thompson, R., Tu, S., Willnow, T.E., Zhang, Y.W.,
and Xu, H. (2016). SNX27 and SORLA Interact to Reduce Amyloidogenic Subcellular Distribution
and Processing of Amyloid Precursor Protein. The Journal of neuroscience : the official journal of
the Society for Neuroscience *36*, 7996-8011.

Huang, Z., Shimazu, K., Woo, N.H., Zang, K., Muller, U., Lu, B., and Reichardt, L.F. (2006). Distinct
roles of the beta 1-class integrins at the developing and the mature hippocampal excitatory
synapse. The Journal of neuroscience : the official journal of the Society for Neuroscience *26*,
11208-11219.

Hung, C., Tuck, E., Stubbs, V., van der Lee, S.J., Aalfs, C., van Spaendonk, R., Scheltens, P., Hardy,
J., Holstege, H., and Livesey, F.J. (2021). SORL1 deficiency in human excitatory neurons causes
APP-dependent defects in the endolysosome-autophagy network. Cell Rep *35*, 109259.

Hwang, J., Estick, C.M., Ikonne, U.S., Butler, D., Pait, M.C., Elliott, L.H., Ruiz, S., Smith, K.,
Rentschler, K.M., Mundell, C*., et al.* (2019). The Role of Lysosomes in a Broad Disease-Modifying
Approach Evaluated across Transgenic Mouse Models of Alzheimer's Disease and Parkinson's
Disease and Models of Mild Cognitive Impairment. Int J Mol Sci *20*.

Karch, C.M., and Goate, A.M. (2015). Alzheimer's disease risk genes and mechanisms of disease
pathogenesis. Biol Psychiatry *77*, 43-51.

Klinger, S.C., Siupka, P., and Nielsen, M.S. (2015). Retromer-Mediated Trafficking of
Transmembrane Receptors and Transporters. Membranes (Basel) *5*, 288-306.
Knupp, A., Mishra, S., Martinez, R., Braggin, J.E., Szabo, M., Kinoshita, C., Hailey, D.W., Small, S.A.,
Jayadev, S., and Young, J.E. (2020). Depletion of the AD Risk Gene SORL1 Selectively Impairs
Neuronal Endosomal Traffic Independent of Amyloidogenic APP Processing. Cell Rep *31*, 107719.
Koopmans, F., van Nierop, P., Andres-Alonso, M., Byrnes, A., Cijsouw, T., Coba, M.P., Cornelisse,
L.N., Farrell, R.J., Goldschmidt, H.L., Howrigan, D.P.*, et al.* (2019). SynGO: An Evidence-Based,
Expert-Curated Knowledge Base for the Synapse. Neuron *103*, 217-234 e214.
Kwart, D., Gregg, A., Scheckel, C., Murphy, E.A., Paquet, D., Duffield, M., Fak, J., Olsen, O., Darnell,
R.B., and Tessier-Lavigne, M. (2019). A Large Panel of Isogenic APP and PSEN1 Mutant Human
iPSC Neurons Reveals Shared Endosomal Abnormalities Mediated by APP beta-CTFs, Not Abeta.
Neuron *104*, 256-270 e255.
Lambert, J.C., Ibrahim-Verbaas, C.A., Harold, D., Naj, A.C., Sims, R., Bellenguez, C., Jun, G.,
Destefano, A.L., Bis, J.C., Beecham, G.W.*, et al.* (2013). Meta-analysis of 74,046 individuals
identifies 11 new susceptibility loci for Alzheimer's disease. Nat Genet *45*, 1452-1458.
Leek, J.T., Johnson, W.E., Parker, H.S., Jaffe, A.E., and Storey, J.D. (2012). The sva package for
removing batch effects and other unwanted variation in high-throughput experiments.
Bioinformatics *28*, 882-883.
Levy, S., Sutton, G., Ng, P.C., Feuk, L., Halpern, A.L., Walenz, B.P., Axelrod, N., Huang, J., Kirkness,
E.F., Denisov, G.*, et al.* (2007). The diploid genome sequence of an individual human. PLoS Biol *5*,
e254.
Li, B., and Dewey, C.N. (2011). RSEM: accurate transcript quantification from RNA-Seq data with
or without a reference genome. BMC Bioinformatics *12*, 323.
Lorenzen, A., Samosh, J., Vandewark, K., Anborgh, P.H., Seah, C., Magalhaes, A.C., Cregan, S.P.,
Ferguson, S.S.G., and Pasternak, S.H. (2010). Rapid and Direct Transport of Cell Surface APP to
the Lysosome defines a novel selective pathway. Molecular Brain *3*, 11.
Mallard, F., Antony, C., Tenza, D., Salamero, J., Goud, B., and Johannes, L. (1998). Direct pathway
from early/recycling endosomes to the Golgi apparatus revealed through the study of shiga toxin
B-fragment transport. J Cell Biol *143*, 973-990.
Manders, E., Verbeek, F., and Aten, J. (1993). Measurement of co-localization of objects in dual-
colour confocal images. Journal of microscopy *169*, 375-382.
Marsh, E.W., Leopold, P.L., Jones, N.L., and Maxfield, F.R. (1995). Oligomerized transferrin
receptors are selectively retained by a lumenal sorting signal in a long-lived endocytic recycling
compartment. J Cell Biol *129*, 1509-1522.
Martin-Belmonte, A., Aguado, C., Alfaro-Ruiz, R., Itakura, M., Moreno-Martinez, A.E., de la Ossa,
L., Molnar, E., Fukazawa, Y., and Lujan, R. (2020). Age-Dependent Shift of AMPA Receptors From
Synapses to Intracellular Compartments in Alzheimer's Disease: Immunocytochemical Analysis of
the CA1 Hippocampal Region in APP/PS1 Transgenic Mouse Model. Front Aging Neurosci *12*,
577996.
Marwaha, R., and Sharma, M. (2017). DQ-Red BSA Trafficking Assay in Cultured Cells to Assess
Cargo Delivery to Lysosomes. Bio Protoc *7*.
Maxfield, F.R., and McGraw, T.E. (2004). Endocytic recycling. Nature reviews Molecular cell
biology *5*, 121-132.

Mayle, K.M., Le, A.M., and Kamei, D.T. (2012). The intracellular trafficking pathway of transferrin. Biochim Biophys Acta *1820*, 264-281.

McQuin, C., Goodman, A., Chernyshev, V., Kamentsky, L., Cimini, B.A., Karhohs, K.W., Doan, M., Ding, L., Rafelski, S.M., Thirstrup, D.*, et al.* (2018). CellProfiler 3.0: Next-generation image processing for biology. PLOS Biology *16*, e2005970.

Offe, K., Dodson, S.E., Shoemaker, J.T., Fritz, J.J., Gearing, M., Levey, A.I., and Lah, J.J. (2006). The lipoprotein receptor LR11 regulates amyloid beta production and amyloid precursor protein traffic in endosomal compartments. The Journal of neuroscience : the official journal of the Society for Neuroscience *26*, 1596-1603.

Olav M. Andersen, N.B., Anne M. Landau, Gro Grunnet Pløen, Anne Mette G. Jensen, Giulia Monti, Benedicte Parm Ulhøi, Jens Randel Nyengaard, Kirsten Rosenmay Jacobsen, Margarita Melnikova Jørgensen, Ida E. Holm, Marianne L. Kristensen, Esben Søvsø Szocska Hansen, Charlotte E. Teunissen, Laura Breidenbach, Mathias Droescher, Ying Liu, Hanne Skovsgaard Pedersen, Henrik Callesen, Yonglun Luo, Lars Bolund, David J. Brooks, Christoffer Laustsen, Scott A. Small, Lars F. Mikkelsen, Charlotte B. Sørensen (2021). In vivo evidence that SORL1, encoding the endosomal recycling receptor SORLA, can function as a causal gene in Alzheimer's Disease. bioRxiv *doi: https://doi.org/10.1101/2021.07.13.452149*.

Ouellette, S.P., and Carabeo, R.A. (2010). A Functional Slow Recycling Pathway of Transferrin is Required for Growth of Chlamydia. Front Microbiol *1*, 112.

Park, M., Penick, E.C., Edwards, J.G., Kauer, J.A., and Ehlers, M.D. (2004). Recycling endosomes supply AMPA receptors for LTP. Science *305*, 1972-1975.

Patapoutian, A., and Reichardt, L.F. (2001). Trk receptors: mediators of neurotrophin action. Curr Opin Neurobiol *11*, 272-280.

Pietila, M., Sahgal, P., Peuhu, E., Jantti, N.Z., Paatero, I., Narva, E., Al-Akhrass, H., Lilja, J., Georgiadou, M., Andersen, O.M.*, et al.* (2019). SORLA regulates endosomal trafficking and oncogenic fitness of HER2. Nat Commun *10*, 2340.

Qureshi, Y.H., Patel, V.M., Berman, D.E., Kothiya, M.J., Neufeld, J.L., Vardarajan, B., Tang, M., Reyes-Dumeyer, D., Lantigua, R., Medrano, M.*, et al.* (2018). An Alzheimer's Disease-Linked Loss-of-Function CLN5 Variant Impairs Cathepsin D Maturation, Consistent with a Retromer Trafficking Defect. Mol Cell Biol *38*.

Racoosin, E.L., and Swanson, J.A. (1993). Macropinosome maturation and fusion with tubular lysosomes in macrophages. J Cell Biol *121*, 1011-1020.

Raghavan, N.S., Brickman, A.M., Andrews, H., Manly, J.J., Schupf, N., Lantigua, R., Wolock, C.J., Kamalakaran, S., Petrovski, S., Tosto, G.*, et al.* (2018). Whole-exome sequencing in 20,197 persons for rare variants in Alzheimer's disease. Ann Clin Transl Neurol *5*, 832-842.

Ramilowski, J.A., Goldberg, T., Harshbarger, J., Kloppmann, E., Lizio, M., Satagopam, V.P., Itoh, M., Kawaji, H., Carninci, P., Rost, B.*, et al.* (2015). A draft network of ligand-receptor-mediated multicellular signalling in human. Nat Commun *6*, 7866.

Rapaport, D., Lugassy, Y., Sprecher, E., and Horowitz, M. (2010). Loss of SNAP29 Impairs Endocytic Recycling and Cell Motility. PLOS ONE *5*, e9759.

Ren, M., Xu, G., Zeng, J., De Lemos-Chiarandini, C., Adesnik, M., and Sabatini, D.D. (1998). Hydrolysis of GTP on rab11 is required for the direct delivery of transferrin from the pericentriolar recycling compartment to the cell surface but not from sorting endosomes. Proc Natl Acad Sci U S A *95*, 6187-6192.

Rogaeva, E., Meng, Y., Lee, J.H., Gu, Y., Kawarai, T., Zou, F., Katayama, T., Baldwin, C.T., Cheng,
R., Hasegawa, H.*, et al.* (2007a). The neuronal sortilin-related receptor SORL1 is genetically
associated with Alzheimer disease. Nat Genet.
Rogaeva, E., Meng, Y., Lee, J.H., Gu, Y., Kawarai, T., Zou, F., Katayama, T., Baldwin, C.T., Cheng,
R., Hasegawa, H.*, et al.* (2007b). The neuronal sortilin-related receptor SORL1 is genetically
associated with Alzheimer disease. Nat Genet *39*, 168-177.
Rohe, M., Hartl, D., Fjorback, A.N., Klose, J., and Willnow, T.E. (2013). SORLA-mediated trafficking
of TrkB enhances the response of neurons to BDNF. PLoS One *8*, e72164.
Romano, R., Rivellini, C., De Luca, M., Tonlorenzi, R., Beli, R., Manganelli, F., Nolano, M., Santoro,
L., Eskelinen, E.-L., Previtali, S.C.*, et al.* (2021). Alteration of the late endocytic pathway in
Charcot–Marie–Tooth type 2B disease. Cellular and Molecular Life Sciences *78*, 351-372.
Scheltens, P., De Strooper, B., Kivipelto, M., Holstege, H., Chetelat, G., Teunissen, C.E., Cummings,
J., and van der Flier, W.M. (2021). Alzheimer's disease. Lancet *397*, 1577-1590.
Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S.,
Rueden, C., Saalfeld, S., Schmid, B.*, et al.* (2012). Fiji: an open-source platform for biological-
image analysis. Nat Methods *9*, 676-682.
Schmidt, V., Sporbert, A., Rohe, M., Reimer, T., Rehm, A., Andersen, O.M., and Willnow, T.E.
(2007). SorLA/LR11 regulates processing of amyloid precursor protein via interaction with
adaptors GGA and PACS-1. J Biol Chem *282*, 32956-32964.
Seaman, M.N. (2004). Cargo-selective endosomal sorting for retrieval to the Golgi requires
retromer. J Cell Biol *165*, 111-122.
Seaman, M.N. (2012). The retromer complex - endosomal protein recycling and beyond. J Cell Sci
*125*, 4693-4702.
Selkoe, D.J. (2002). Alzheimer's disease is a synaptic failure. Science *298*, 789-791.
Shi, Y., Kirwan, P., Smith, J., Robinson, H.P., and Livesey, F.J. (2012). Human cerebral cortex
development from pluripotent stem cells to functional excitatory synapses. Nat Neurosci *15*, 477-
486, S471.
Simoes, S., Neufeld, J.L., Triana-Baltzer, G., Moughadam, S., Chen, E.I., Kothiya, M., Qureshi, Y.H.,
Patel, V., Honig, L.S., Kolb, H.*, et al.* (2020). Tau and other proteins found in Alzheimer's disease
spinal fluid are linked to retromer-mediated endosomal traffic in mice and humans. Science
translational medicine *12*.
Simoes S., G.J., Buitrago L., Qureshi Y.H., Feng X., Kothiya M., Cortes E., Patel V., Kannan S., Kim
Y.-H., Chang K.-T., ADNI, Hussaini A., Moreno H., Di Paolo G., Andersen O.M., Small S.A (2021).
Alzheimer's vulnerable brain region relies on a distinct retromer core dedicated to endosomal
recycling. . Cell Reports *Accepted*.
Small, S.A., and Gandy, S. (2006). Sorting through the cell biology of Alzheimer's disease:
intracellular pathways to pathogenesis. Neuron *52*, 15-31.
Small, S.A., and Petsko, G.A. (2020). Endosomal recycling reconciles the Alzheimer's disease
paradox. Science translational medicine *12*.
Sonnichsen, B., De Renzis, S., Nielsen, E., Rietdorf, J., and Zerial, M. (2000). Distinct membrane
domains on endosomes in the recycling pathway visualized by multicolor imaging of Rab4, Rab5,
and Rab11. J Cell Biol *149*, 901-914.
Spoelgen, R., von Arnim, C.A., Thomas, A.V., Peltan, I.D., Koker, M., Deng, A., Irizarry, M.C.,
Andersen, O.M., Willnow, T.E., and Hyman, B.T. (2006). Interaction of the cytosolic domains of

sorLA/LR11 with the amyloid precursor protein (APP) and beta-secretase beta-site APP-cleaving
enzyme. The Journal of neuroscience : the official journal of the Society for Neuroscience *26*, 418-
428.

Sun, J., and Roy, S. (2018). The physical approximation of APP and BACE-1: A key event in
alzheimer's disease pathogenesis. Dev Neurobiol *78*, 340-347.

Temkin, P., Morishita, W., Goswami, D., Arendt, K., Chen, L., and Malenka, R. (2017). The
Retromer Supports AMPA Receptor Trafficking During LTP. Neuron *94*, 74-82 e75.

Thonberg, H., Chiang, H.H., Lilius, L., Forsell, C., Lindstrom, A.K., Johansson, C., Bjorkstrom, J.,
Thordardottir, S., Sleegers, K., Van Broeckhoven, C.*, et al.* (2017). Identification and description
of three families with familial Alzheimer disease that segregate variants in the SORL1 gene. Acta
Neuropathol Commun *5*, 43.

Tian, X., Gala, U., Zhang, Y., Shang, W., Nagarkar Jaiswal, S., di Ronza, A., Jaiswal, M., Yamamoto,
S., Sandoval, H., Duraine, L.*, et al.* (2015). A Voltage-Gated Calcium Channel Regulates Lysosomal
Fusion with Endosomes and Autophagosomes and Is Required for Neuronal Homeostasis. PLOS
Biology *13*, e1002103.

Wakabayashi, K., Narisawa-Saito, M., Iwakura, Y., Arai, T., Ikeda, K., Takahashi, H., and Nawa, H.
(1999). Phenotypic down-regulation of glutamate receptor subunit GluR1 in Alzheimer's disease.
Neurobiol Aging *20*, 287-295.

Wang, Y., Eng, D.G., Kaverina, N.V., Loretz, C.J., Koirala, A., Akilesh, S., Pippin, J.W., and
Shankland, S.J. (2020). Global transcriptomic changes occur in aged mouse podocytes. Kidney Int
*98*, 1160-1173.

Willnow, T.E., and Andersen, O.M. (2013). Sorting receptor SORLA--a trafficking path to avoid
Alzheimer disease. J Cell Sci *126*, 2751-2760.

Yap, C.C., Digilio, L., McMahon, L.P., Garcia, A.D.R., and Winckler, B. (2018). Degradation of
dendritic cargos requires Rab7-dependent transport to somatic lysosomes. Journal of Cell Biology
*217*, 3141-3159.

Yasuda, R.P., Ikonomovic, M.D., Sheffield, R., Rubin, R.T., Wolfe, B.B., and Armstrong, D.M.
(1995). Reduction of AMPA-selective glutamate receptor subunits in the entorhinal cortex of
patients with Alzheimer's disease pathology: a biochemical study. Brain Res *678*, 161-167.

Young, J.E., Boulanger-Weill, J., Williams, D.A., Woodruff, G., Buen, F., Revilla, A.C., Herrera, C.,
Israel, M.A., Yuan, S.H., Edland, S.D.*, et al.* (2015). Elucidating Molecular Phenotypes Caused by
the SORL1 Alzheimer's Disease Genetic Risk Factor Using Human Induced Pluripotent Stem Cells.
Cell stem cell *16*, 373-385.

Yuan, S.H., Martin, J., Elia, J., Flippin, J., Paramban, R.I., Hefferan, M.P., Vidal, J.G., Mu, Y., Killian,
R.L., Israel, M.A.*, et al.* (2011). Cell-surface marker signatures for the isolation of neural stem
cells, glia and neurons derived from human pluripotent stem cells. PLoS One *6*, e17540.

Zigdon, H., Meshcheriakova, A., Farfel-Becker, T., Volpert, G., Sabanay, H., and Futerman, A.H.
(2017). Altered lysosome distribution is an early neuropathological event in neurological forms
of Gaucher disease. FEBS Lett *591*, 774-783.

**Figure legends**

**Figure 1.** Loss of *SORL1* expression leads to increased TRKB and GLUA1 localization in early endosomes.

Representative immunofluorescent images of WT and *SORL1*KO neurons showing increased colocalization of **(a)** TRKB (green) and **(b)** GLUA1 (green) with EEA1 (red). All neurons were immunolabeled with MAP2 (far-red) and counterstained with DAPI (blue). Scale bar: 10μm. In all cases, quantification of colocalization was represented as Mander's correlation co-efficient (MCC). 1 WT and 2 *SORL1*KO isogenic clones were used for these experiments and 10 images per clone per genotype were analyzed. Data represented as mean ± SD. Data was analyzed using parametric two-tailed unpaired t test. Significance was defined as a value of *p < 0.05, **p < 0.01, ***p < 0.001, and ****p < 0.0001.

**Figure 2.**  Loss of *SORL1* expression impairs trafficking to late endosomes and lysosomes.

**a)** *SORL1*KO neurons show reduced lysosomal proteolytic activity as measured by DQ Red BSA. Representative immunofluorescent images of WT and *SORL1*KO neurons showing double immunolabeling for MAP2 (green) and DQ Red BSA (red). Scale bar: 10μm Quantification of fluorescence intensity of DQ Red BSA using ImageJ software. **(b-f)** *SORL1*KO neurons show reduced colocalization of cargo with late endosomes and lysosomes. Representative immunofluorescent images of WT and *SORL1*KO neurons showing reduced colocalization of **(b)** TRKB (green) and **(c)** GLUA1 (green) with Rab7 positive late endosomes (red) in *SORL1*KO neurons. Representative immunofluorescent images of WT and *SORL1*KO neurons showing reduced colocalization of **(d)** APP (green) and **(e)** TRKB (green) and **(f)** GLUA1 (green) with LAMP1 positive lysosomes (red) in *SORL1*KO neurons. Scale bar: 10μm. In all cases, quantification of colocalization was represented as Mander's correlation co-efficient (MCC). 1 WT and 2 *SORL1*KO isogenic clones were used for these experiments and 10 images per clone per genotype were analyzed. Data represented as mean ± SD. Data was analyzed using parametric two-tailed unpaired t test. Significance was defined as a value of *p < 0.05, **p < 0.01, ***p < 0.001, and ****p < 0.0001.

**Figure 3:** Enhancing *SORL1* Expression improves trafficking to late endosomes and lysosomes

**a)** *SORL1*OE neurons show no change in lysosomal proteolytic activity as measured using DQ Red BSA after a 6 hour treatment but do show an enhancement of trafficking at 24 hours. Representative immunofluorescent images of WT and *SORL1*OE neurons showing double immunolabeling for MAP2 (green) and DQ Red BSA (red). Scale bar: 10 μm. Quantification of fluorescence intensity of DQ Red BSA using ImageJ software. 1 cell line of each genotype (WT vs. SORL1 OE) were used for these experiments. 10 images per genotype were analyzed. Representative immunofluorescent images of WT and *SORL1*OE neurons showing increased colocalization of APP, TRKB and GLUA1 (green) with Rab7 (red) **(b-d)** and LAMP1 (red) **(e-g)** in *SORL1*OE neurons. *SORL1*OE neurons and controls have endogenous GFP expression due to the piggybac vector system. GFP fluorescence is pseudo-colored (Far-red) and was used to outline cell bodies. Scale bar: 10μm. Nuclei are counterstained with DAPI (blue). In all cases, quantification of colocalization was represented as Mander's correlation co-efficient (MCC). 1 cell line of each genotype (WT vs. SORL1 OE) were used for these experiments. 10 images per genotype were analyzed. Data represented as mean ± SD. Data was analyzed using parametric two-tailed unpaired t test. Significance was defined as a value of $*p < 0.05$, $**p < 0.01$, $***p < 0.001$, and $****p < 0.0001$.

**Figure 4.** Loss of *SORL1* impacts the cell surface recycling pathway

**a)** *SORL1*KO neurons show slower rate of transferrin recycling. Quantification of fluorescence intensity of intracellular transferrin at different time points after treating cells with Alexa Fluor 647-conjugated transferrin for 15 mins, using ImageJ software. Data represented as percent of time 0 fluorescence intensity. 2 WT and 2 *SORL1*KO isogenic clones were used for these experiments. 12 images per clone per genotype were analyzed.

**(b)** *SORL1*KO neurons show larger recycling endosomes. Representative immunofluorescent images of WT and *SORL1*KO neurons labeled with antibodies for MAP2 (red) and Rab11 (green). Nuclei were counterstained with DAPI (blue). Scale bar: 5μm. Quantification of size of Rab11 labeled recycling endosomes using CellProfiler software. 1 WT and 2 *SORL1*KO isogenic clones were used for these experiments. 15 images per clone per

genotype were analyzed. Representative immunofluorescent images of WT and *SORL1*KO neurons showing increased colocalization of **(c)** APP (green), **(d)** TRKB (green) and **(e)** GLUA1 (green) with Rab11 positive recycling endosomes (red) in *SORL1*KO neurons. Scale bar:10μm In all cases, quantification of colocalization was represented as Mander's correlation co-efficient (MCC). 1 WT and 2 *SORL1*KO isogenic clones were used for these experiments. 10 images per clone per genotype were analyzed. Data represented as mean ± SD. Data was analyzed using parametric two-tailed unpaired t test and two-was ANOVA. Significance was defined as a value of $*p < 0.05$, $**p < 0.01$, $***p < 0.001$, and $****p < 0.0001$.

**Figure 5.** Loss of *SORL1* expression impairs recycling to the cell surface.

**a-b)** *SORL1*KO neurons show reduced cell surface expression of APP **(a)** and GLUA1 **(b)**. Representative immunofluorescent images of WT and *SORL1*KO neurons labeled with antibodies for APP **(a)** (red) and GLUA1 **(b)** (red). Scale bar: 5μm. Intensity of APP and GLUA1 measured using ImageJ software. Data is presented as a ratio of surface intensity to total intensity. 2 WT and 2 *SORL1*KO clones were used in these experiments. 16 images per clone per genotype were analyzed. Data represented as mean ± SD. Normally distributed data was analyzed using parametric two-tailed unpaired t test. Significance was defined as a value of $*p < 0.05$, $**p < 0.01$, $***p < 0.001$, and $****p < 0.0001$. **(c)** Multielectrode array (MEA) analysis of WT and *SORL1*KO neurons at early (d27) and late (d66) time points of differentiation. 1 WT and 1 *SORL1*KO clone was used for these experiments. Data represented as mean ± SD. Data was analyzed using parametric two-tailed unpaired t test. Significance was defined as a value of $*p < 0.05$, $**p < 0.01$, $***p < 0.001$, and $****p < 0.0001$.

**Figure 6.** Overexpression of *SORL1* enhances endosomal recycling.

**a)** *SORL1*OE neurons show faster rate of transferrin recycling. Quantification of fluorescence intensity of intracellular transferrin at different time points after treating cells with Alexa Fluor 647-conjugated transferrin for 15 mins, using ImageJ software. Data represented as percent of time 0 fluorescence intensity. 1 cell line of each

genotype (WT vs. SORL1 OE) were used for these experiments. 10 images per genotype were analyzed. **(b)** *SORL1*OE neurons show reduced size of recycling endosomes. Representative immunofluorescent images of WT and *SORL1*OE neurons labeled with Rab11 (green) and MAP2 (red) showing smaller Rab11 positive recycling endosomes in *SORL1*OE neurons. Nuclei counterstained with DAPI (blue). Quantification of Rab11+ recycling endosome size performed using Cell Profiler software and represented as area of Rab11+ vesicles. Scale bar: 5μm. 1 cell line of each genotype (WT vs. SORL1 OE) were used for these experiments. 26 images per genotype were analyzed. Representative immunofluorescent images of WT and *SORL1*OE neurons showing increased colocalization of **(c)** APP (green), **(d)** TRKB (green) and **(e)** GLUA1 (green) with Rab11 (red) positive recycling endosomes. *SORL1*OE neurons and controls have endogenous GFP expression due to the piggybac vector system. GFP fluorescence is pseudo-colored (Far-red) and was used to outline cell bodies. Quantification of colocalization with Rab11 represented as Mander's Correlation Co-efficient (MCC). Scale bar: 10μm. 1 cell line of each genotype (WT vs. SORL1 OE) were used for these experiments. 10 images per genotype were analyzed. Representative immunofluorescent images of WT and *SORL1*OE neurons showing increased cell surface expression of **(f)** APP (red) and **(g)** GLUA1 (red) in *SORL1*OE neurons. Scale bar: 5μm Fluorescence intensity of APP and GLUA1 measured using ImageJ software. Data is presented as a ratio of surface intensity to total intensity. Nuclei counterstained with DAPI. 1 cell line of each genotype (WT vs. SORL1 OE) were used for these experiments. 12-14 images per genotype were analyzed. Data represented as mean ± SD. Data was analyzed using parametric two-tailed unpaired t test and two-way ANOVA. Significance defined as a value of *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, and ****$p < 0.0001$.

**Figure 7.** Analysis of bulk RNA-sequencing data indicates alterations in cell surface and extracellular trafficking, receptor-ligand and channel activity. Gene ontology analysis of DEGs in WT and *SORL1*KO neurons. Shown here are the top upregulated **(a)** and downregulated **(b)** molecular function terms in *SORL1*KO neurons. GO annotation terms are listed on the y-axis, adjusted p-value is shown on the x-axis. **(c)** Ligand-receptor network changes in *SORL1*KO neurons, identified if genes are more than 1.5-fold increased or decreased in *SORL1*KO

neurons with an adjusted p-value less than 0.05. Circles denote ligands, squares denote receptors, blue indicates genes expressed significantly lower in *SORL1*KO neurons, red indicates genes expressed significantly higher in *SORL1*KO neurons. Arrows point from ligand to receptor, denoting receptor-ligand interactions. Black arrows denote consistent expression changes between ligand and receptor, indicating that both genes in the pair are either upregulated or downregulated. Gray arrows denote inconsistent changes between ligand and receptor. Clusters impacted by cell surface recycling (β-integrins, ephrins) or neurotrophic signaling (FGF/FGFR, NGF/NGFR) are indicated by green dotted arrows. RNA was collected from 3 separate differentiations including a combination of two WT clones and two *SORL1*KO clones. Each sample includes 2-3 technical replicates.

**Figure 8. Model of modulation of SORL1 expression in hiPSC-derived cortical neurons.**

Modulation of SORLA expression regulates endosomal recycling and degradative pathways in hiPSC-derived neurons: The early endosome is a sorting hub where various cargo can be trafficked to degradative or cell surface recycling pathways in neurons (A). In this study, we mapped trafficking of three important neuronal cargo under conditions of depletion (B) or enhancement (C) of the AD risk gene SORL1/SORLA. As depicted in panel B, our data suggest that modulation of SORL1 expression significantly impacts the neuronal recycling pathway as we observe increased size and accumulation of cargo in recycling endosomes in SORL1 KO neurons compared to isogenic WT cells. Specifically, loss of SORL1 impacts recycling to the cell surface because, while there is increased cargo in recycling endosomes, there is a reduction of this cargo on the neuronal surface. Loss of SORL1 also impacts the degradative pathway out of the early endosome as there is a reduction of cargo in late endosome and lysosomes. As depicted in panel C, enhancement of SORL1 expression reciprocally impacts these pathways. SORL1 OE neurons also have an increase in cargo in recycling endosomes, but unlike KO cells, this results in more cell surface trafficking of neuronal cargo, leading to reduced stress (small size) of recycling endosomes. Additionally, SORL1 OE enhances trafficking out of the early endosome towards the degradative pathway as well. Taken together, our data solidifies the AD risk gene SORL1 as a key modulator of neuronal endosomal trafficking. Created with Biorender.com.

**Supplementary figures**

**Supplementary figure 1**.

Loss of *SORL1* expression leads to increased VPS35 localization in early endosomes. **(a)** Representative immunofluorescent images of WT and *SORL1*KO neurons showing increased colocalization of VPS35 (green) with EEA1 (red). All neurons were immunolabeled with MAP2 (far-red) and counterstained with DAPI (blue). Scale bar: 10μm. In all cases, quantification of colocalization was represented as Mander's correlation co-efficient (MCC). 1 WT and 2 SORL1KO isogenic clones were used for these experiments and 10 images per clone per genotype were analyzed. Data represented as mean ± SD. Significance was determined using parametric two-tailed unpaired t test and was defined as a value of $*p < 0.05$, $**p < 0.01$, $***p < 0.001$, and $****p < 0.0001$.

**Supplementary figure 2**

**a)** *SORL1*KO neurons show larger lysosome size and increased lysosome number. Representative immunofluorescent images of WT and *SORL1*KO neurons labeled with LAMP1 (green) and MAP2 (red) showing increased LAMP1 positive vesicle size in *SORL1*KO neurons. Quantification of LAMP1 size and number was performed using Cell Profiler software. LAMP1 size is represented as area of LAMP1 positive vesicles, and LAMP1 number is represented as number of LAMP1 positive vesicles per square micron of cell area. Scale bar: 5μm **(b)** *SORL1*KO neurons show no change in colocalization of lysosomes with the lysosomal enzyme Cathepsin D. Representative immunofluorescent images of WT and *SORL1*KO neurons labeled with antibodies for LAMP1 (green), Cathepsin D (red) and MAP2 (Far-red) showing no alteration in colocalization of Cathepsin-D with LAMP1 in *SORL1*KO neurons. Nuclei counterstained with DAPI (blue). Scale bar: 10μm Quantification of colocalization of LAMP1 with Cathepsin-D represented as Mander's correlation coefficient (MCC). 10-20 images were analyzed per genotype. Two isogenic clones of each genotype were used in all experiments. Data

represented as mean ± SD. Normally distributed data was analyzed using parametric two-tailed unpaired t test. Significance was defined as a value of *p < 0.05, **p < 0.01, ***p < 0.001, and ****p < 0.0001.

**Supplemental figure 3.**

Loss of SORL1 does not change protein expression of the compartments or cargo analyzed in this study as analyzed by Western blot. Representative blots in (a), quantification in (b).

**Supplemental figure 4**

**a)** Principal component analysis (PCA) plot of all RNAseq samples using all expressed genes. Samples are color coded by differentiation batch. Triangles represent WT samples, circles represent *SORL1*KO. Genotype accounts for the highest variance (37%, PC1, x-axis). **b)** Volcano plot. Log2 fold change between *SORL1*KO and WT is shown along the x-axis. Statistical significance is shown along the y-axis, and is measured by adjusted p-value. Genes upregulated in *SORL1*KO neurons are shown by red circles, genes downregulated in *SORL1*KO neurons are shown by blue circles. We observed 6643 DEGs, with 2819 upregulated and 3824 downregulated. There are significantly more down regulated genes than upregulated genes (p<2.2e-16). Grey circles represent genes that are not significantly differentially expressed.

**Supplemental figure 5 .**

Loss of *SORL1* expression alters synaptic pathways. Analysis of bulk RNA-sequencing data indicates alterations in synaptic pathway functioning. We conducted gene ontology analysis of DEGs in WT and *SORL1*KO neurons using the SynGO synaptic annotation database. Shown here are the top upregulated and biological process and cellular component terms in *SORL1*KO neurons. GO annotation terms are listed on the y-axis, adjusted p-value is shown on the x-axis. No downregulated pathways in the SynGO database were shown to be enriched in *SORL1*KO neurons.

**Supplemental Table 1.**

Summary of statistical analyses for data presented in this manuscript. In this table we present the statistical data that correspond to the experiments presented in the figures. This includes the group means, the difference between the means $\pm$ SEM, and the 95% confidence interval.

Figure 1

# Figure 2

# Figure 3

# Figure 4

# Figure 5

# Figure 6

# Figure 7

# Figure 8

A   **WT neuron**

B   ***SORL1* KO neuron**

C   ***SORL1* OE neuron**

EE   Early endosome
RE   Recycling endosome
LE   Late endosome
LY   Lysosome

TrkB
APP
GLUA1
DQ-BSA

Neuronal soma

# Supplemental Figure 2

# Supplemental Figure 3

# Supplemental Figure 4

A

B

Significantly more downregulated
than upregulated genes
p<2.2e-16

Supplemental Figure 5

A      Biological Process Synaptic Terms
Upregulated in *SORL1* KO neurons

B      Cellular Component Synaptic Terms
Upregulated in *SORL1* KO neurons

| Figure Number | Experimental Measure | Group A | Group B | Group A Mean | Group B Mean | Difference between means (B - A) ± SEM | 95% Confidence Interval |
|---|---|---|---|---|---|---|---|
| 1A | TRKB colocalization with EEA1 | WT Neurons | SORL1KO Neurons | 0.0799 | 0.1228 | 0.04290 ± 0.01042 | 0.02155 to 0.06425 |
| 1B | GLUA1 colocalization with EEA1 | WT Neurons | SORL1KO Neurons | 0.0496 | 0.1727 | 0.1231 ± 0.02257 | 0.07682 to 0.1693 |
| 2A | 6hr DQ Red BSA Intensity | WT Neurons | SORL1KO Neurons | 52.1 | 29.48 | -22.62 ± 2.684 | -28.09 to -17.15 |
| 2A | 24hr DQ Red BSA Intensity | WT Neurons | SORL1KO Neurons | 62.54 | 43.35 | -19.19 ± 3.608 | -26.58 to -11.80 |
| 2B | TRKB colocalization with RAB7 | WT Neurons | SORL1KO Neurons | 0.3102 | 0.2149 | -0.09530 ± 0.01234 | -0.1212 to -0.06937 |
| 2C | GLUA1 colocalization with RAB7 | WT Neurons | SORL1KO Neurons | 0.3828 | 0.286 | -0.09685 ± 0.02779 | -0.1538 to -0.03992 |
| 2D | APP colocalization with LAMP1 | WT Neurons | SORL1KO Neurons | 0.2983 | 0.1799 | -0.1184 ± 0.01951 | -0.1584 to -0.07844 |
| 2E | TRKB colocalization with LAMP1 | WT Neurons | SORL1KO Neurons | 0.071 | 0.04461 | -0.02639 ± 0.004876 | -0.03643 to -0.01635 |
| 2F | GLUA1 colocalization with LAMP1 | WT Neurons | SORL1KO Neurons | 0.1967 | 0.1588 | -0.03790 ± 0.02085 | -0.08060 to 0.004801 |
| 3A | 6hr DQ Red BSA Intensity | WT Neurons | SORL1OE Neurons | 30.31 | 27.84 | -2.473 ± 2.059 | -6.799 to 1.854 |
| 3A | 24hr DQ Red BSA Intensity | WT Neurons | SORL1OE Neurons | 32.61 | 39.73 | 7.123 ± 1.887 | 3.157 to 11.09 |
| 3B | APP colocalization with RAB7 | WT Neurons | SORL1OE Neurons | 0.7041 | 0.8008 | 0.09670 ± 0.02520 | 0.04375 to 0.1496 |
| 3C | TRKB colocalization with RAB7 | WT Neurons | SORL1OE Neurons | 0.44 | 0.5261 | 0.08610 ± 0.02417 | 0.03532 to 0.1369 |
| 3D | GLUA1 colocalization with RAB7 | WT Neurons | SORL1OE Neurons | 0.4287 | 0.5512 | 0.1225 ± 0.02485 | 0.07029 to 0.1747 |

| | | | | | | | |
|---|---|---|---|---|---|---|---|
| 3E | APP colocalization with LAMP1 | WT Neurons | SORL1OE Neurons | 0.1397 | 0.2949 | 0.1552 ± 0.02066 | 0.1118 to 0.1986 |
| 3F | TRKB colocalization with LAMP1 | WT Neurons | SORL1OE Neurons | 0.1539 | 0.2674 | 0.1135 ± 0.01501 | 0.08196 to 0.1450 |
| 3G | GLUA1 colocalization with LAMP1 | WT Neurons | SORL1OE Neurons | 0.1353 | 0.3206 | 0.1853 ± 0.02279 | 0.1374 to 0.2332 |
| 4A - T0 | Transferrin Intensity T0 | WT Neurons - T0 | SORL1KO Neurons - T0 | 100 | 100 | -4.167E-08 | -4.634 to 4.634 |
| 4A - T10 | Transferrin Intensity T10 | WT Neurons - T10 | SORL1KO Neurons - T10 | 64.14 | 82.82 | -18.68 | -24.35 to -13.00 |
| 4A - T20 | Transferrin Intensity T20 | WT Neurons - T20 | SORL1KO Neurons - T20 | 58.44 | 67.48 | -9.038 | -13.67 to -4.404 |
| 4A - T40 | Transferrin Intensity T40 | WT Neurons - T40 | SORL1KO Neurons - T40 | 56.41 | 68.96 | -12.55 | -17.19 to -7.920 |
| 4B | Rab11 Area | WT Neurons | SORL1KO Neurons | 0.06625 | 0.073 | 0.006747 ± 0.002882 | 0.0009338 to 0.01256 |
| 4C | APP colocalization with RAB11 | WT Neurons | SORL1KO Neurons | 0.2689 | 0.3147 | 0.04580 ± 0.01442 | 0.01626 to 0.07534 |
| 4D | TRKB colocalization with RAB11 | WT Neurons | SORL1KO Neurons | 0.1071 | 0.2277 | 0.1206 ± 0.02924 | 0.06071 to 0.1805 |
| 4E | GLUA1 colocalization with RAB11 | WT Neurons | SORL1KO Neurons | 0.2605 | 0.3065 | 0.04595 ± 0.01599 | 0.01320 to 0.07870 |
| 5A | Surface:Total APP Intensity | WT Neurons | SORL1KO Neurons | 1.128 | 0.5124 | -0.6157 ± 0.05825 | -0.7331 to -0.4982 |
| 5B | Surface:Total GLUA1 Intensity | WT Neurons | SORL1KO Neurons | 1.022 | 0.5769 | -0.4451 ± 0.04588 | -0.5368 to -0.3534 |
| 5C | D27 Weighted Mean Firing | WT Neurons | SORL1KO Neurons | 3.768 | 7.696 | 3.928 ± 1.266 | 1.169 to 6.687 |
| 5C | D66 Weighted Mean Firing | WT Neurons | SORL1KO Neurons | 7.074 | 3.784 | -3.289 ± 1.101 | -5.688 to -0.8903 |

| | | | | | | | |
|---|---|---|---|---|---|---|---|
| 6A - T0 | Transferrin Intensity T0 | WT Neurons - T0 | SORL1OE Neurons - T0 | 100 | 100 | -0.0000002 | -11.47 to 11.47 |
| 6A - T10 | Transferrin Intensity T10 | WT Neurons - T10 | SORL1OE Neurons - T10 | 82.52 | 69.57 | 12.95 | 1.483 to 24.42 |
| 6A - T20 | Transferrin Intensity T20 | WT Neurons - T20 | SORL1OE Neurons - T20 | 74.47 | 59.35 | 15.12 | 3.653 to 26.59 |
| 6A - T40 | Transferrin Intensity T40 | WT Neurons - T40 | SORL1OE Neurons - T40 | 68.31 | 55.43 | 12.89 | 1.419 to 24.35 |
| 6B | Rab11 Area | WT Neurons | SORL1OE Neurons | 0.101 | 0.09005 | -0.01099 ± 0.003199 | -0.01742 to -0.004558 |
| 6C | APP colocalization with RAB11 | WT Neurons | SORL1OE Neurons | 0.3903 | 0.5878 | 0.1975 ± 0.02717 | 0.1404 to 0.2546 |
| 6D | TRKB colocalization with RAB11 | WT Neurons | SORL1OE Neurons | 0.1846 | 0.2969 | 0.1123 ± 0.02416 | 0.06155 to 0.1631 |
| 6E | GLUA1 colocalization with RAB11 | WT Neurons | SORL1OE Neurons | 0.2509 | 0.2965 | 0.04560 ± 0.01822 | 0.007315 to 0.08389 |
| 6F | Surface:Total APP Intensity | WT Neurons | SORL1OE Neurons | 0.9734 | 1.189 | 0.2155 ± 0.09083 | 0.02710 to 0.4038 |
| 6G | Surface:Total GLUA1 Intensity | WT Neurons | SORL1OE Neurons | 0.5727 | 0.9505 | 0.3778 ± 0.06831 | 0.2374 to 0.5182 |
| Supplementary 1A | VPS35 colocalization with EEA1 | WT Neurons | SORL1KO Neurons | 0.457 | 0.5838 | 0.1268 ± 0.02249 | 0.08129 to 0.1724 |
| Supplementary 2A | LAMP1 area | WT Neurons | SORL1KO Neurons | 0.1008 | 0.1112 | 0.01032 ± 0.004276 | 0.001730 to 0.01892 |
| Supplementary 2B | Cathepsin-D colocalization with LAMP1 | WT Neurons | SORL1KO Neurons | 0.3621 | 0.3499 | -0.01219 ± 0.02558 | -0.06488 to 0.04051 |
| Supplementary 2C | Number of LAMP1 puncta per $\mu m^2$ | WT Neurons | SORL1KO Neurons | 1.452 | 3.339 | 1.887 ± 0.5339 | 0.8145 to 2.960 |
| Supplementary 2C | Number of APP puncta per $\mu m^2$ | WT Neurons | SORL1KO Neurons | 0.6442 | 0.5906 | -0.05352 ± 0.03725 | -0.1288 to 0.02175 |

| | | | | | | | |
|---|---|---|---|---|---|---|---|
| Supplementary 2C | Number of TRKB puncta per $\mu m^2$ | WT Neurons | SORL1KO Neurons | 1.318 | 1.208 | -0.1100 ± 0.06986 | -0.2509 to 0.03084 |
| Supplementary 2C | Number of GLUA1 puncta per $\mu m^2$ | WT Neurons | SORL1KO Neurons | 0.5202 | 0.4951 | -0.02509 ± 0.01880 | -0.06303 to 0.01285 |
| Supplementary 2C | Number of EEA1 puncta per $\mu m^2$ | WT Neurons | SORL1KO Neurons | 0.5488 | 0.5986 | 0.04975 ± 0.03228 | -0.01560 to 0.1151 |
| Supplementary 3B | SORLA/Actin | WT Neurons | SORL1KO Neurons | 1 | 0.02221 | -0.9778 ± 0.06426 | -1.135 to -0.8205 |
| Supplementary 3B | TRKB/Actin | WT Neurons | SORL1KO Neurons | 1 | 1.043 | 0.04315 ± 0.3219 | -0.7444 to 0.8307 |
| Supplementary 3B | LAMP1/Actin | WT Neurons | SORL1KO Neurons | 1 | 1.141 | 0.1413 ± 0.1599 | -0.2499 to 0.5324 |
| Supplementary 3B | VPS35/Actin | WT Neurons | SORL1KO Neurons | 1 | 1.146 | 0.1459 ± 0.1848 | -0.3064 to 0.5982 |
| Supplementary 3B | GLUA1/Actin | WT Neurons | SORL1KO Neurons | 1 | 0.9428 | -0.05717 ± 0.07946 | -0.2516 to 0.1373 |

Supplemental Table 1.